Brain aromatase and its relationship with parental experience and

behavior in male mice

Paula Duarte-Guterman[1]*, Dimitri A. Skandalis[1,2] †, Ariane Merkl †, Diana B. Geissler,

Günter Ehret*

Institute of Neurobiology, University of Ulm, D-89069 Ulm, Germany.

[1] Present address: Department of Psychology and Centre for Neuroscience, Brock University,

ON, Canada.

[2] Present address: Psychological & Brain Sciences and Johns Hopkins Kavli Neuroscience

Discovery Institute, Johns Hopkins University, MD, USA.

† These authors contributed equally to this work

* Correspondence:

Paula Duarte-Guterman; email: pduarteg@brocku.ca

Günter Ehret; email: guenter.ehret@uni-ulm.de

Keywords

brain plasticity, estrogen, hemisphere lateralization, infanticide, limbic system, paternal

care

Abbreviations: BNST, bed nucleus of the stria terminalis; CAM, cortical amygdala; LS,

lateral septum; LSD, lateral septum dorsal; LDI, lateral septum intermediate; LSV, lateral

septum ventral; MAM, medial amygdala; MPOA, medial preoptic area; NAC, nucleus

accumbens; PIR, piriform cortex; VMH, ventromedial nucleus of the hypothalamus.

Abstract

In most mammals, paternal care is not mandatory for raising offspring. In house mice, experience with pups governs the extent and quality of paternal care. First-time fathers undergo a dramatic transition from ignoring or killing pups to caring for pups. The behavioral shift occurs together with changes in brain estrogen signaling as indicated by changes in estrogen receptor presence and distribution in multiple areas regulating olfaction, emotion, and motivation. Here, we report changes in the expression of aromatase, the enzyme converting testosterone into estrogen, as an indirect measure of estrogen synthesis. The amount of paternal experience (5 or 27 days) was associated with increased numbers of immunocytochemically-identified aromatase expressing cells in the medial and cortical amygdala, posterior piriform cortex, and ventromedial hypothalamus. Functionally, these changes can be related to the disappearance of aggression or neglect towards pups when first-time fathers or, even more, well-experienced fathers are handling their own pups. In the lateral septum, the anterior piriform cortex and to some extent in the medial preoptic area, parental experience increased the number of aromatase-positive cells only in fathers with 27 days of experience, and only in the right hemisphere. This is a new case of brain-functional lateralization due to experience that has activated certain instinctive behavior. Nuclei/areas associated with maternal care (medial preoptic area, bed nucleus of stria terminalis, nucleus accumbens) exhibited a left-hemisphere advantage in aromatase expressing cells, both in pup-naïve and pup-experienced males. This newly found lateralization may contribute to the left-hemisphere dominant processing and perception of pup calls to release parental behavior. In general, the experience-dependent changes in aromatase expression we observed in most brain areas did not mirror the previously reported changes in estrogen receptors (ERα) when pup-naïve males became pup-caring fathers. Hence, paternal behavior may depend in a brain area-specific way on the differential action of estrogen through its receptors and/or direct local modulation of neural processing.

Introduction

Mammalian paternal care is facultative, and the behavioral and neural dynamics that result in becoming a caring father are less well understood compared to those of obligatory maternal care (e.g., Bales and Saltzman, 2016; Brunton and Russell, 2008; Dulac et al., 2014; Guoynes and Marler, 2020; Horrell et al., 2021; Lonstein and De Vries, 2000; Numan, 2020; Numan and Insel, 2003; Rilling and Mascaro, 2017; Woodroffe and Vincent, 1994)). In rodents and primates, it is hypothesized that maternal care is induced by hormonal changes during pregnancy and parturition (onset phase) whereas the continuation of pup-caring (maintenance phase) is governed by sensory stimuli from the pups (Brunton and Russell, 2008; Fahrbach and Pfaff, 1982; Rosenblatt and Siegel, 1981). In house mice, pup-naïve females (i.e., having no sexual or pup caring experience), males, and even previous fathers do not experience the onset phase. Rather, they require various amounts of experience with pups (from hours up to several days of contact) for the activation of caring behavior (Akther et al., 2021; Alsina-Llanes et al., 2015; Ehret and Buckenmaier, 1994; Ehret and Koch, 1989; Ehret et al., 1987,1993; Elwood, 1985; Koch and Ehret, 1989; Martín-Sánchez et al., 2015; Stolzenberg and Rissman, 2011; Tachikawa et al., 2013). Nonetheless, pup-naïve females engage in consummatory (pup retrieval) and appetitive (preference for pup ultrasounds) behaviors (Koch and Ehret, 1991; Numan, 2006) after fewer days of experience with pups (and sometimes spontaneously; Alsina-Llanes et al., 2015; Martín-Sánchez et al., 2015) compared to either pup-naïve males or previous fathers (Ehret and Buckenmaier, 1994; Ehret et al., 1987,1993). Pup naïve females thus appear to be more easily sensitized or motivated by pup cues than males and even fathers to provide parental care. This observation is suggestive of sex differences in the neural mechanisms underlying the activation of maternal versus paternal instincts.

Numerous studies have characterized the role of the limbic regions in the activation

and regulation of parental behaviors. These regions include the medial preoptic area (MPOA),

bed nucleus of the stria terminalis (BNST), lateral septum (LS), nucleus accumbens (NAC),

nuclei of the amygdala, and ventromedial hypothalamus (VMH). Olfaction is thought to

provide the main modulatory sensory input, through pathways including the vomeronasal

organ to the BNST and amygdala, the olfactory bulb to the amygdala, via the VMH to the LS,

and via the piriform cortex (PIR) and hippocampus to the LS (e.g. Bridges, 2015; Brunton and

Russell, 2008; Dulac et al., 2014; Ehret et al., 1991; Kohl and Dulac, 2018; Kohl et al., 2017;

Numan, 2006, 2020). Estradiol treatment can activate both maternal and paternal instincts

(Ehret and Koch 1989; Koch and Ehret, 1989; Poindron et al., 1988; Romero-Morales et al.,

2018; Rosenblatt and Ceus, 1998; Rosenblatt et al., 1994; Siegel and Rosenblatt, 1975) and

changes in brain estrogen signaling (through estrogen receptors) may be part of the underlying

mechanism. In house mice, expression of parental behaviors coincides with changes in the

distribution of estrogen receptor (ERα) positive cells in limbic brain areas of pup-naïve and

pup-experienced females (Ehret and Buckenmaier, 1994; Koch, 1990) and males (Ehret et al.,

1993). In pup-naïve male and female voles, high levels of spontaneous pup care are correlated

with higher number of ERα positive cells in the MPOA (Li et al., 2015), potentially indicating

convergent circuit dynamics.

Low levels of circulating estrogens in males (Nilsson et al., 2015) suggest that brain-

active estrogens are primarily synthesized locally in the brain from testosterone by the

enzyme aromatase (e.g., Cornil et al., 2006,2013; Garcia-Segura, 2008). Locally synthesized

estrogens have been implicated in the onset of paternal behaviors in California mice (Trainor

and Marler, 2002; Trainor et al., 2003). Aromatase is expressed in the limbic system of male

mice (LS, BNST, MPOA, nuclei of the amygdala, VMH, and PIR) (Foidart et al., 1995;

Schleicher et al., 1986; Stanić et al., 2014; Wozniak et al., 1992). In the present study, we

sought to examine changes in aromatase expression during the onset and maintenance of

paternal behavior of male mice (pup-naïve and fathers with increasing experience). We

focused on LS, BNST, MPOA, nuclei of the amygdala, VMH, and PIR, which are areas in

which ERα positive cells have also been located in pup-experienced fathers (Ehret et al.,

1993).

Materials and methods

Animals

Adult male and female laboratory mice (outbred strain NMRI, 2-3 months old) with no

previous sexual or parental experience were housed at the University of Ulm. As previously

(Ehret et al., 1993), mice were randomly assigned to one of three treatment groups consisting

of (1) naïve males without any experience with pups (N); (2) males co-caring with their

female mate for their first litter for five days (5D); or (3) males co-caring with their female

mate for their first litter until weaning and their second litter for five days. The total co-caring

time of the latter males (first and second litter) was 26-27 days (27D) since after delivery of

the first litter, females in post-partum estrus paired immediately and, while nursing the first

litter up to weaning at 21 days, delivered the second litter after 21-22 days of gestation. Mice

were housed at 22ºC and a 12 h light-dark cycle (starting at 7.00 h) in plastic cages (26.5 × 20

× 14 cm) either with other naïve males (group N; 4 brothers per cage) or with one female

(groups 5D and 27D). Water and food were available *ad libitum*. The experiments were

carried out in accordance with the European Communities Council Directive (86/609/EEC)

and were approved by the appropriate authority (Regierungspräsidium Tübingen, Germany).

Pup retrieval tests

Testing of pup retrieval behavior was performed as previously described (Ehret and Koch,

1989; Ehret et al., 1993). Male mice (alone or with litter mate and pups) were placed

overnight on a running board (length 110 cm, width 8 cm) with a central nest depression (Fig.

1A) suspended in a sound-proof and anechoic room. The running board was supplied with

mouse chow and nest material from the cage and covered with a plexiglass hood having an

opening for the tip of a suspended water bottle. The next day, retrieving tests were done under

dim red light (<1 lx) ensuring that males relied on the sense of smell, audition, and touch to

respond and retrieve pups. Before the start of testing, the hood, and the female with the pups

(groups 5D and 27D) were removed and placed back to their home cage outside the

soundproof room. Seven pups were kept in the room to serve as targets for retrieving. The

males were habituated for at least 30 min to the new situation after the hood (group N) or the

hood and female with pups (groups 5D and 27D) were removed. Prior to testing males of the

N group, five-day old foster pups were rubbed with nest material of the respective males to

ensure a familiar scent. During the test, the seven pups (always 1-3 pups at a time) were

randomly placed on the running board, at least 30 cm away from the nest. A male was scored

as 'retriever' if he retrieved all seven pups to the nest within 10 min. A male was scored as

'non-retriever' if he had not retrieved any pups after 10 mins. The distinction between

retriever and non-retriever was clear-cut, because once a male had retrieved the first pup, he

retrieved all. Retrieving tests were immediately stopped and the pups removed if a male

attacked any of the pups ('aggressor').

Immunocytochemistry and microscopy

Immediately following the pup retrieval test, males were euthanized by cervical dislocation

and their brains quickly dissected and frozen over liquid nitrogen. Brains were mounted using

O.C.T compound and frontal sections were serially cut (30 μm) using a cryostat (Thermo

Scientific Microm HM560), and immediately mounted on slides to keep track of the left and

right sides of the brain. Sections were fixed in 4% paraformaldehyde in 0.1 M phosphate

buffer (PB; pH = 7.4) for 60 min. Sections were rinsed three times for 5 min with 0.1 M PB in

between all subsequent processing steps. Slides were incubated with 0.2% Triton X-100

(Sigma) in 0.1 M PB for 45 min to permeabilize cells and with 1% H_2O_2 (Merck) in 0.1 M PB

for 20 min to block endogenous peroxidase. In a humidified chamber, the sections were

treated with 2% normal goat serum in 0.1 M PB for 60 min to block non-specific staining,

then with the polyclonal rabbit anti-aromatase antibody (Abcam Cat# ab35604,

RRID:AB_867729; 0.002 mg/ml in 2% normal goat serum) at 4ºC for 72h. Sections were

then incubated with the secondary antibody from goat against rabbit horseradish peroxidase

(Dako, Cat# PO448, 0.00125 mg/ml in 0.1 M phosphate buffer) at room temperature for 60

min. Immunoreactivity was revealed by incubating sections in 0.015% diaminobenzidine

tetrachloride (DAB; Sigma), 0.023% NiCl, and 0.013% H_2O_2 in 0.1 M PB for 7 min. Sections

were rinsed in PB, dehydrated in a series of alcohols and xylene, and coverslipped with

Entellan (Merck). At the time of purchase, the primary antibody was recommended for

aromatase immunohistochemistry on mouse tissue and the supplier provided positive quality

controls with mouse ovarian tissue. Dutta et al. (2014) also performed a Western blot with

mouse tissue that confirmed specificity of the antibody. Previous studies have used this

antibody to detect aromatase immunoreactivity in mouse ovary (Laws et al., 2014), brain stem

(Ji et al., 2017) and auditory cortex (Charitidi and Canlon, 2010). We tested the specificity of

the secondary antibody labeling by incubating sections with PB instead of the primary

antibody. In all these control sections, immunoreactivity was completely absent (Fig. 2A).

Sections were independently analyzed by two blinded experimenters. In view of

varying non-specific background labeling, the experimenters started the analysis by checking

many sections from all animal groups in order to agree which kind of labeling would be

considered aromatase-specific. Brain areas were identified with 10x objective magnification,

aromatase-positive cells were identified by focusing with a 40x objective magnification. Cells

were considered as expressing aromatase if they were (1) more intensely stained with shades

of gray than the background, and (2) contained black granulae often grouped around the cell

nucleus (Fig. 2B). In cases in which we observed a single or many black granulae without an

associated cellular structure in shades of gray, the structure was not counted as an aromatase-

positive cell (Fig. 2C). This procedure led to (a) a high concordance of counting results of the

two experimenters, (b) counts of only definite aromatase-positive cells, and (c) relatively low

numbers of aromatase-positive cells per evaluated brain area in a given section.

Sections were analyzed with a light microscope (Axiophot, Zeiss Germany). The brain

areas in which aromatase-positive cells were counted are listed in Table 1. Areas were

matched to the mouse brain atlas plates and associated Bregma coordinates (Paxinos and

Franklin, 2001) and anatomically matched across all experimental animals. These areas

largely overlap with those showing ERα positive cell dynamics during the acquisition of pup-

caring experience in male (Ehret et al., 1993) and female (Ehret and Buckenmaier, 1994)

mice.. Aromatase-positive cells were counted separately in the right and left hemispheres of

the brain. Counts per section for a given animal, brain area and hemisphere were averaged

between the maximum number of consecutive sections that could be matched across all

animals, i.e., two sections for LS, BNST, VMH, and nucleus accumbens (NAC), or three

sections for PIR, MPOA, cortical amygdala (CAM), and medial amygdala (MAM). These

average counts from the individual animals entered further statistical analyses concerning the

experimental groups of animals (N, 5D, 27D) and comparisons among the groups.

Statistical analyses

All statistical analyses were carried out using R v4.2.0 (R Core Team, 2022).

Behavior: Male behavior in response to pup presentation was classified as ignoring,

attacking, or retrieving a pup. The outcomes of behavioral trials were analysed as the

multinomial probability of exhibiting a given response to pup presentation (aggression,

ignore, retrieval) as a function of paternal experience (experience as ordered factor, N < 5D <

27D). To perform pairwise comparisons between the groups for each behavior, we calculated

the probability that the difference in the posterior predictive intervals between groups was

significantly directional. We then corrected for multiple comparisons using the false

discovery rate.

Aromatase immunolabeling: Aromatase cell counts were conducted in two data sets,

which were combined to increase statistical power for comparisons. The data sets were

obtained with the same immunocytochemical protocols and the same chemicals, however, the

experiments were performed as two separate cohorts. We observed some count variation

between the data of the cohorts (e.g., for the PIR -1.34 data; see Supplemental Fig. 2), and so

each data set was standardized by the mean and variance of the paternal groups (5D and 27D;

the groups for which the datasets were overlapping). Sections for some brain areas were

missing in some mice. The pattern of missingness would have led to omitting eight of 56

individuals (14%). In order not to totally exclude these animals from the data analysis, we

elected to impute the missing data (16 cases, 5.7% of 280 total region counts) using predictive

mean matching (van Buuren and Groothuis-Oudshoorn, 2011; R package mice). We

generated five such candidate data sets and integrated over the resulting uncertainty.

Our statistical model of aromatase variation was designed to address non-

independence arising from correlated variation within individuals among areas. Such non-

independence could be an especially important factor when comparing identifiable subregions

of an area, such as in the piriform cortex and each part of the lateral septum. We therefore

examined variation in normalized cell counts among all brain areas simultaneously

(multivariate ANOVA), treating individual mouse identification (ID) as a correlated group

(random) effect to account for hemisphere differences within individual. Responses were

modelled as t-distributed to increase robustness against outlier counts. The interaction of

Experience x Hemisphere was examined together.

All models were fit using the R package *brms* (Bürkner, 2017) using four chains of

2500 iterations each, and a thinning rate of 5. We assumed uniform priors throughout. Model

convergence was assessed by $\hat{R} = 1$ (Vehtari et al., 2021); the $\hat{R}$ statistic was computed within

each imputed data set. We calculated the posterior means of each data set based on the fitted

statistical model (*emmeans* package; Lenth, 2022), and used these to examine the posterior

distributions of contrasts between hemispheres (left *minus* right for each brain area and

experimental group of animals) and between the experimental groups for each brain area and

hemisphere, i.e. 5D minus N (5D-N), 27D minus N (27D-N), 27D minus 5D (27D-5D). These

data are plotted in Figure 3. Significance of each contrast was judged as a Bayesian

probability of direction P > 0.975 (Makowski et al., 2019), which is an index of effect

existence, i.e., our level of certainty in whether a reported change is different from zero, either

positive or negative. This threshold closely corresponds to the two-tailed p-value of 0.05, and

so for simplicity, we report the p-value. Plots were generated using the *ggplot2* package

(Wickham, 2016).

Residual correlations among areas represent relationships that are not explained by the

other population- and group-effects, including correlated expression and unexpected technical

covariation. Inspection of residual correlations can therefore inform biological hypotheses.

However, direct interpretation of the residual correlation matrix is challenging because the

correlation between two brain areas could be due to mutual covariation with another area. We

dichotomized relationships among brain areas by studying the matrix of *partial* correlations

(de la Fuente et al., 2004; Shipley, 2000). We constructed an initial graph in which edges

were created between all nodes with significant (p < 0.05) pairwise correlations. Edges were

iteratively pruned by testing whether the association remained statistically significant after

testing it against all other paths between two variables, up to third order. The independence

test was determined from the posterior distribution of residual correlation matrices, and

implemented in the R package pcalg (Kalisch et al., 2012). The result is an undirected, acyclic

graph representing a network of associations among brain areas. The graph is shown in Figure

4. The large number of tests may be conservative and may reject true associations, so we also

report tentative links observed when setting a less stringent threshold of $p < 0.10$.

Pup ignoring and aromatase immunolabeling: Pup retrieval and aggression

behaviors are completely colinear with the treatment groups (see below), so we could not

examine behavioral differences simultaneously with experience. Pup ignoring, however, was

observed in all experimental groups of mice. Therefore, we sought to find factors predicting

why some males, regardless of experience, ignore pups (or why some males react to pups).

Behavior toward pups was coded as reacting (aggressiveness or retrieval) or ignoring, and we

then conducted a similar analysis to that described above for aromatase cell counts. All males

were examined together without considering differences in experiences or hemispherical

variation in aromatase cell counts.

Results

Male mice retrieve pups after acquiring parental experience

Only males with paternal experience retrieved pups (Fig. 1A). Figure 1B shows that

naïve males with no previous sexual experience and pup contact mostly ignored the pups (n =

5/8) or were aggressive (n = 3/8). Most fathers with 5 days (5D) or 26-27 days (27D) of pup-

experience retrieved pups (n = 10/12 and 11/12, respectively) and only few ignored pups (n =

2/12 and 1/12, respectively). No father was aggressive.

The probability of observing aggressive behavior toward pups was significantly higher

in naïve males compared to fathers regardless of experience (5D and 27D, Fig. 1B). The

probability of other non-paternal behavior like ignoring pups was higher in naïve males

compared to 27D fathers. Parental behavior measured as pup retrieval was significantly higher

in fathers (5D and 27D) compared to naïve males. There were no significant differences in the

probability of pup-retrieving between the groups of fathers, 5D and 27D (Fig. 1B).

Aromatase expression in the male mouse brain, general picture

We found aromatase-positive cells in each examined area and with each level of

experience. In absolute terms, largest numbers of aromatase positive cells per section were

seen in PIR, LSI, MAM, and CAM (see unstandardized data in Supplement, Fig. 2).

Examples of aromatase labeling in the LSV, PIR, VMH and MAM of experimental groups N

and 27D showed qualitative differences between the groups and between left and right sides

of the brain (Fig. 2). Standardized cell counts together with the group means are shown in Fig.

3A along with our estimates of the posterior differences in aromatase counts by hemisphere

(Left – Right, Fig. 3B), and between levels of experience (Fig. 3C).

Lateralized occurrence of aromatase positive cells

We found evidence for significantly greater numbers of aromatase positive cells in the

left hemisphere of the MPOA and BNST in all three experimental groups (Left – Right > 0,

Fig. 3B). The NAC exhibited a similar trend to MPOA and BNST, but with a just

insignificant contrast in the 27D group. We did not detect significant lateralization of

aromatase counts in any of the experimental groups in the amygdala (CAM and MAM),

VMH, or posterior PIR (Bregma -1.34). In the other brain areas, the appearance of

lateralization strongly depended on the level of pup-experience. Long-term experience (27D)

led to greater numbers of aromatase positive cells in the right hemispheres of anterior PIR

(Bregma 0.98), the LSV, and the anterior LSI (Bregma 0.14). Posterior LSI (Bregma 0.98)

exhibited a similar trend that was just non-significant. In the LSD, pup-experience had a

short-term effect (5D) during which counts in the left hemisphere were greater. This effect

was not present in the 27D group (Fig. 3B).

Experience-dependent expression of aromatase

In all cases in which we found significant changes of aromatase counts with pup-experience, the number of aromatase positive cells increased (Fig. 3C). This experience-dependent increase was pervasive for the amygdala (CAM and MAM), the VMH, and the posterior PIR (Bregma -1.34). For these areas, the increase in aromatase positive cells was present in both hemispheres and with increasing experience, i.e., experience contrasts had significant positive values for both brain hemispheres and all experimental groups. In the LSD, LSI, and LSV (both Bregma coordinates), and the anterior PIR (Bregma 0.98), aromatase positive cells increased significantly only in the right hemisphere of 27D males. Experience-dependent effects were absent in the NAC, the BNST, and the MPOA, except for a significant effect in the right MPOA of 27D males, i.e., 27D males had significantly more aromatase positive cells than 5D males (Fig. 3C).

Graphical analysis of covariation among brain areas

We examined the unexplained (residual) variation in aromatase cell numbers among brain areas as graphically shown in Figure 4. We found positive residual correlations between all brain areas, with the sole exception of CAM (-1.34) and LSD (0.98), which were negatively correlated. Three subgroups of covariation are obvious. First, the LS is highly connected and each subregion (D, I, V) is correlated across Bregma coordinates and to at least one other subregion. The LS appears to form the backbone of variation, to which other nodes are connected. This includes subgroups comprised of MPOA (connected to LSV), of NAC and BNST (connected to LSI), and of CAM and VMH (connected to LSD). Second, the subgroup of MAM and anterior PIR appears to be weakly connected to other parts of the graph (MPOA and VMH), because the edges are significant only with $p < 0.10$. Third, posterior PIR did not appear to be even weakly correlated with any other analyzed brain area

despite the increase in aromatase expression with experience that was similar to CAM, MAM,

and VMH (see Fig. 3).

Variation in aromatase expression in males that react or ignore pups

The loss of aggressiveness towards pups and the gain of pup retrieval behavior starkly

marks the transition to fatherhood (Fig. 1). Nonetheless, some males appear to ignore pups,

and even one second-time father failed to retrieve pups during the pup retrieval test. We

therefore examined if variation in aromatase cell counts in any area might predict whether a

male reacted to or did not react to pups (contrast of reacting – ignoring mean cell counts, Fig.

5). Reacting males had significantly higher numbers of aromatase cells in the MAM

(p=0.011), and a weak tendency to higher numbers in the CAM (p=0.06).

Discussion

In the present study, we found that aromatase, which catalyses the conversion of

testosterone to estrogen, is expressed throughout areas of the limbic system of male mice,

regardless of their experience with pups. However, we identified important factors

contributing to hierarchical variation in expression levels among individuals. First, the

dramatic changes in behavior associated with paternal experience were associated with

changes in aromatase expression in areas/nuclei of the limbic system. However, not only were

these changes area-specific, but also hemisphere-specific (summarised in Fig. 6). Finally,

layered on these widespread patterns, we detected correlated aromatase levels among brain

areas within individual mice. We briefly review the role of aromatase and the limbic areas to

discuss how such expression patterns may influence, and be regulated by, paternal behaviors

and experience.

Aromatase and the expression of paternal care

Aromatase expression in the male mouse brain has been shown via immunostaining with several protocols using different antibodies (Akther et al., 2015; Foidart et al., 1995), partly with enhancement through transgenic techniques (Stanić et al., 2014). In agreement with the published data, we found aromatase expression in all studied brain areas of our three experimental groups of male mice (Fig. 3). Differences in the levels of aromatase expression between studies relate to differences in the staining methods, data analysis, and the social environment of the animals prior to and during the behavioral tests. For example, in our study, dynamics of paternal behavior and associated changes in aromatase levels have been examined while the fathers were continually housed with their female mate except during the pup-retrieval tests. This is different from the study of Akther et al. (2015) in which the relationship between pup-retrieving behavior in males and the dynamics of aromatase levels was tested throughout different ethological contexts (presence, absence and reintroduction of their female mates and pups). In that previous study, brain aromatase expression was highest when the fathers were housed with their female partner, but without their pups, followed by fathers housed with the whole family (Akther et al., 2015). Therefore, the observed changes in aromatase levels in that study can be related to male-female interaction. Nevertheless, the strong reduction of pup-retrieval in fathers after injection with an aromatase inhibitor (letrozole; Akther et al., 2015) shows, along with the results of our present study, that proper levels of aromatase expression are integral to the display of paternal behavior in mice.

Interestingly, circulating testosterone is negatively correlated with paternal behavior in various rodent species (reviewed in Bales and Saltzman, 2016; Numan and Insel, 2003) and testosterone can inhibit paternal care in mice (Okabe et al., 2013). Our study did not examine the source of testosterone, the substrate of aromatase in the brain. However, previous studies have shown that changes in circulating testosterone may not be reflected locally in the brain.

Neurosteroids can be locally produced, and levels in the brain may be independent of

circulation or gonadal status (Giatti et al., 2019; Hojo et al., 2008, 2004).

In the following, we discuss the observed presence and changes of aromatase levels in

the studied brain areas of the limbic system in order to see how these areas may contribute to

changing male behavior towards pups from aggressive or ignoring in naïve males to caring in

experienced fathers (Fig. 1). And further, the comparison of dynamics of aromatase-positive

cells with the dynamics of ERα-positive cells in the given brain areas will show in how far

they are in parallel and support the hypothesis about aromatase influencing indirectly (via

estrogen synthesis and ER binding) paternal behavior.

Amygdala and ventromedial hypothalamus

Three nuclei, the MAM, CAM, and VMH were most affected by pup experience with

increases in the number of aromatase positive cells from N to 5D to 27D animals (Fig. 3 A

and C; Supplement Fig. 2). These areas are associated with the regulation of aggression

compared to sexual and parental behavior (e.g., Chen et al., 2019; Dwyer et al., 2022;

Hashikawa et al., 2016; Lee et al., 2014; Sheehan et al., 2001; Unger et al., 2015; Yang et al.,

2017). Important changes in aggression in the course of becoming a caring father are

mediated by olfactory signals from the vomeronasal organ via the accessory olfactory bulb to

the CAM and MAM (e.g., Hashikawa et al., 2016) and concern the switch from aggressive to

caring behavior towards pups (Tashikawa et al., 2013). Concomitantly, aggressive motivation

in fathers increases against male intruders in order to protect their own pups from potential

pup killers as shown in house mice (Shabalova et al., 2020), Californian mice (Trainor et al.,

2008) and prairie voles (Getz et al., 1981). This increased aggression towards intruders was

not mediated by changes in ERα or ERß expression in limbic centers, but reflected in

increased neuronal activation (FOS-expression) in the MPOA, MAM and, potentially the

VMH in California mice (Trainor et al., 2008). In house mice fathers, aggressive behavior

toward intruders is regulated by oxytocin and activation of neurons in areas of the

hypothalamus (periventricular and supraoptic nuclei; Shabalova et al., 2020). Future work

could examine whether changes in aggression in naïve vs fathers are also related to aromatase

levels in the amygdala and VMH.

In contrast to aromatase expression (present study), the number of estrogen receptors

(ERα) in MAM, CAM, and VMH did not change with parental experience in male mice (5D

or 27D; (Ehret et al., 1993). Together, this indicates that the increased presence of local

estrogen synthesis does not facilitate the expression of ERα, as has been shown in the VMH

(Devidze et al., 2005; Malikov and Madeira, 2013), especially in combination with parental

experience (Koch, 1990). Very limited work has been done on how other estrogen receptors

are involved in paternal care in rodents (Hyer et al., 2017; Torres et al., 2024). It is possible

that other subtypes (ERβ and membrane receptor GPER) are involved in the expression of

parental behavior. Our work indicates that parental experience is associated with changes in

aromatase expression, and this can potentially affect the balance of androgen to estrogen

levels in the brain. Future work should also examine whether androgen receptors, expressed in

the same brain areas as aromatase (Dart et al., 2024), are implicated in parental behavior (e.g.,

see work in Mongolian gerbils; Martinez et al., 2019). We suggest that the increase in

aromatase expression in the MAM, CAM, and VMH in new (5D) and highly experienced

(27D) fathers (Fig. 6) supports the control of instinctive behavior (Honda et al., 2011)

including inhibition of infanticide, via increased local synthesis of estrogens on neurons (e.g.,

Cornil et al., 2006, 2013).

Piriform cortex

In house mouse mothers, the performance of pup-retrieving does not depend on

olfactory signals from the main olfactory bulb and its projections to the PIR (Koch and Ehret,

1991). Pup-naïve females, however, needed this olfactory information for learning to become

as maternal as the mothers (Ehret and Buckenmaier, 1994; Koch and Ehret, 1991). While

learning pup cues for becoming maternal, virgin females with maternal experience have an

increase in ERα positive cells in the posterior PIR compared to females without experience

(also in the entorhinal cortex; Koch, 1990). Similarly, as males became caring fathers, ERα

positive cells become visible in the posterior PIR (also in the entorhinal cortex; Ehret et al.,

1993). In view of the significant increase of aromatase expression in the posterior PIR with

increasing paternal experience (Fig. 3C), it is possible that in PIR, unlike MAM, CAM, and

VMH, estrogen may have induced its own receptors (tested for ERα) and enhanced olfactory

learning of pup cues for increased paternal performance.

The significant increase of aromatase positive cells only in the right hemisphere in the

anterior PIR of 27D fathers does not directly correlate with any evidence known to us. In

studies on ERα expression in the limbic areas of female and male mice under various

conditions of the reproductive cycle, experience with pups, and intact or lesioned olfaction, no

hemisphere differences have been reported in any area (Ehret et al., 1993; Ehret and

Buckenmaier, 1994; Koch, 1990; Koch and Ehret, 1991). The splitting of PIR in regions with

experience-dependent lateralized (anterior PIR) and non-lateralized (posterior PIR) aromatase

change shows compartmentalization of the PIR. Functional relationships need to be elucidated

in further studies.

Lateral septum

The main effects of pup-experience on aromatase expression in the LS were changes

in the left-right balance (Fig. 3B) and significant increase of aromatase-positive cells only in

the right LS of highly experienced (27D) fathers (Fig. 3C). As in the PIR, ERα positive cells

in the LS become visible in caring fathers compared to pup-naïve males (Ehret et al., 1993).

Since hemisphere differences in ERα positive cells have not been found, we cannot directly

relate increased right-hemisphere aromatase expression with non-lateralized increase of ERα

expression in the LS of both hemispheres. Lateralized input to the rostral parts of the LSD and

LSI comes from the hippocampal CA1 and subiculum region (work in mice and rats; Rizzi-

Wise and Wang, 2021; Risold and Swanson, 1997). In rodents, the CA1 shows lateralization

in synapse morphology and receptor distribution (Shinohara et al., 2008; Xiao and Jordan,

2002) and left-right dissociation of memory processes (Shipton et al., 2014) with right-

hemisphere dominance of spatial memory in mice (Shinohara et al., 2012). Also, androgen

receptor levels are higher in the left hippocampus in male rats (Xiao and Jordan, 2002). It is

unclear, however, whether and how lateralized hippocampal input to the LS could have

caused lateralized aromatase expression in the LS of 27D fathers (Fig. 3C). Right hemisphere

lateralization is associated with the regulation of mood, affect, and stress, and with sustained

and global arousal and attention (e.g., Ehret, 2006; Palomero-Gallagher and Amunts, 2022).

Speculatively, our observed right-hemisphere LS bias in 27D fathers could be related to the

stress of continuous arousal and steady involvement in a routine of pup-care over several

weeks.

Medial preoptic area, bed nucleus stria terminalis, and nucleus accumbens

The MPOA together with the BNST and their output via the NAC are essential for

controlling parental behavior in rodents (e.g., Numan, 2006; Bridges, 2015; Wu et al., 2014).

Significant increases have been observed in the MPOA both of the number of and area

occupied by ERα positive cells in 5D and 27D pup-experienced father mice (Ehret et al.,

1993) and, although absent in pup-naïve males, ERα-positive cells are present in the BNST of

27D fathers (Ehret et al., 1993; Koch and Ehret, 1989). In females, the number of ERα

positive cells are also higher in the MPOA and BNST of one-day pup-experienced virgin

females (Ehret and Buckenmaier, 1994). In the current study, we did not observe changes in

the number of aromatase positive cells in the BNST and NAC and found only a small increase

in the MPOA in just the 27D males relative to 5D. This suggests that changes in sensory

processing and behavior with experience in MPOA, BNST, and NAC may be more dependent

on regulation of estrogen signaling through binding to ERα than on local estrogen action after

synthesis via aromatase. However further work on other estrogen receptors is needed. Brain

areas that regulate social behavior, including parental care, are conserved in vertebrates

(O'Connell and Hofmann, 2012). However, neuroendocrine ligand spatial profiles are more

variable than receptor distributions within the social decision-making network among

vertebrates, although there are regional differences (O'Connell and Hofmann, 2012). Our

work supports experience-dependent regional differences and specifically that influences of

MPOA, BNST, and NAC on parental behavior may depend on changes in ERα distribution,

whereas the influences of VMH and amygdala may depend on changes in aromatase

expression. Overall, our results point to an area-specific regulation of receptor and ligand

synthesis with parental experience.

The brain areas MPOA, BNST, and NAC exhibited substantial lateralization of

presence of aromatase-positive cells, biased toward the left hemisphere (Fig. 3B). Left-

hemisphere dominance has been noted in a number of contexts, including maternal perception

and processing of pup-ultrasounds, in the latter case, through cFos expression (Ehret, 1987;

Geissler et al., 2016). Maternal pup retrieval behavior likewise is associated with left- but not

right-hemisphere oxytocin receptor expression and action (Marlin et al., 2015). To what

extent can these sources of left-hemisphere bias in females (primarily auditory cortex) inform

our findings? The left-hemisphere bias in females has primarily been observed in the auditory

cortex (Geissler et al., 2016; Marlin et al., 2015). Lateralization of cFos expression has not

been observed in the neural responses to pup presence in the limbic system of female mice,

MPOA and BNST included (Geissler et al., 2013). However, neural activation characterized

by cFos labeling concerns the input side of neurons (e.g. Sheng et al., 1990, 1993). Provided

such absence of hemisphere differences in activation at the input side of the MPOA and

BNST also existed in males/fathers, we can hypothesize that the greater left hemisphere

aromatase expression in the MPOA, BNST, may lead to a left-hemisphere bias of activation at

the output of the MPOA and BNST after local estrogen action (enabled via aromatase) had

taken place. Whether such a functional bias actually exists and how it may contribute to a

preferred processing of pup ultrasounds in order to activate appetitive pup-searching behavior

and possible left-hemisphere dominant ultrasound perception in pup-experienced male mice

similar to mothers remains to be shown.

Correlated individual variation in aromatase-positive cells between brain areas

The brain's connectivity suggests that counts of active cells (e.g., c-Fos as indicative

of brain activation) within individuals may be correlated across regions. For example,

aromatase could be uniformly lower or higher in some individuals compared to others, or

could vary highly non-uniformly across brain areas. We expect that such variation will

contribute to individual differences in behavior, such as some fathers continuing to ignore

pups (Fig. 1B). Our study found significant residual correlated variation among brain areas

(Fig. 4), generally not depicted in the data shown in Fig. 3. Our analysis points to a central

role of the LS in the variation of aromatase expression in the mouse brain. The LS has a

central functional role in the brain integrating cognitive and emotional information in the

regulation of social relationships and instincts (Besnard and Leroy, 2022; Menon et al., 2022;

Rizzi-Wise and Wang, 2021) including parental behavior (Fleischer and Slotnick, 1978;

Slotnick and Nigrosh, 1975). Interestingly, these partial correlations (shown in Fig. 4) reflect

main neural connections of the subregions LSD and LSV. For example, there is input from the

amygdala to the LS supporting the LSD-CAM partial correlation and there is an output from

the LS to the MPOA supporting the LSV-MPOA partial correlation (Besnard and Leroy,

2022; Menon et al., 2022; Rizzi-Wise and Wang, 2021; Risold and Swanson, 1997). Thus,

residual variation in aromatase expression (Fig. 4) appears to mirror functional connections

and relationships of the LS. It is noteworthy that the posterior PIR and, to a lesser degree, the

anterior PIR and the MAM remained outside this connectivity. Indeed, Supplementary Fig. 2

shows that posterior PIR (Bregma -1.34) was the sole case in which we observed substantial

divergence in overall cell counts between the two cohorts of animals. Together, these

observations on correlated individual variations indicate both coherent relationships between

brain areas in a functional network and the presence of possibly unknown additional factors

intrinsic to given experiments. In our present case, such factors modulated the numbers of

aromatase-positive cells in PIR and MAM. Therefore, our study shows the usefulness of

correlative modelling to assess and estimate the relationships between contributions of the

members of a neuronal network to its overall functional properties. This is another example

that correlative modelling supports the adequate interpretation of absolute data.

Amygdala aromatase expression is related to reacting to pups

Our study design cannot fully link changes in individual behavior to changes in

aromatase in the brain because the switch in the expression of both is so stark. We therefore

sought to examine why some fathers *do not* appear to react to their own pups (i.e., they ignore

their pups) and, like most naïve males, continue to ignore them. Although tentative, our

results (Fig. 5) point to the possibility that the amygdala (especially the MAM) mediates

strong emotional responses to pups and the particular expression of is dictated by the network

of brain areas. It would be interesting to study the longitudinal responses of males (naïve and

fathers) to pups, to address the individual repeatability of their responses and determine

whether ignoring is an intermediate state (between aggression and retrieval) or a distinct

behavioral state. Such a study will require a much larger sample size than was possible here,

to opportunistically sample the natural variation in individual behavior.

Conclusion

Aromatase expression in the male limbic brain is regulated with parental experience in

an area-, hemisphere-, and experience-specific manner. Acquiring paternal pup-caring

experience enhanced aromatase expression (number of aromatase positive cells) in brain areas

processing olfactory information (PIR, CAM, MAM) and regulating aggression, sexual

behavior and other social relations (MAM, VMH, LS). In centers essential for controlling

parental behavior (MPOA, BNST, NAC), aromatase expression was instead present with left-

hemisphere dominance and largely independent of experience with pups. In general, the

experience-dependent changes in aromatase expression we observed in most brain areas did

not mirror the previously reported changes in estrogen receptors (ERα) when pup-naïve males

became pup-caring fathers.

Acknowledgements

We thank H. Sabine Schmidt for assistance with animal care, immunocytochemistry,

and microscopy. This study was supported by the Deutsche Forschungsgemeinschaft,

EH53/19 to GE. PDG was funded by a postdoctoral fellowship from the Fonds Quebecois

Recherche Nature et Technologie. PDG is currently supported by the Canada Research Chair

Program. DAS was partially funded by a postdoctoral fellowship from the Kavli Neuroscience

Discovery Institute.

References

Akther S, Huang Z, Liang M, Zhong J, Fakhrul AAKM, Yuhi T, et al. (2015). Paternal
retrieval behavior regulated by brain estrogen synthetase (aromatase) in mouse sires that
engage in communicative interactions with pairmates. Front. Neurosci. 9, 450. doi:
10.3389/fnins.2015.00450

Alsina-Llanes, M., De Brun, V., Olazábal, D.E. (2015). Development and expression of
maternal behavior in naïve female C57BL/6 mice. Dev. Psychobiol. 57, 189–200.
https://doi.org/10.1002/dev.21276

Bales KL, Saltzman W. (2016). Fathering in rodents: Neurobiological substrates and
consequences for offspring. Hormones and Behavior, 77, 249–259.

Besnard A, Leroy F. (2022). Top-down regulation of motivated behaviors via lateral
septum sub-circuits. Mol. Psychiatry, 27(8), 3119–3128.

Bridges RS. (2015). Neuroendocrine regulation of maternal behavior. Front.
Neuroendocrinol. 36, 178–196. doi: 10.1016/j.yfrne.2014.11.007

Brunton PJ, Russell JA. (2008). The expectant brain: adapting for motherhood. Nat. Rev.
Neurosci. 9(1), 11–25.

Bürkner PC. (2017). brms: An R package for Bayesian multilevel models using Stan.
Journal of Statistical Software, 80, 1–28.

Charitidi K, Canlon B. (2010). Estrogen receptors in the central auditory system of male
and female mice. Neuroscience, 165(3), 923–933.

Chen PB, Hu RK, Wu YE, Pan L, Huang S, Micevych PE, et al. (2019). Sexually
dimorphic control of parenting behavior by the medial amygdala. Cell, 176(5), 1206-1221.

Cornil CA, Ball GF, Balthazart J. (2006). Functional significance of the rapid regulation of
brain estrogens: Where do the estrogens come from? Brain Res. 1126(1), 2–26.

Cornil CA, Seredynski AL, de Bournonville C, Dickens MJ, Charlier TD, Ball GF, et al.
(2013). Rapid control of reproductive behaviour by locally synthesised oestrogens: focus on
aromatase. J. Neuroendocrinol. 25(11), 1070–1078.

de la Fuente A, Bing N, Hoeschele I, Mendes P. (2004). Discovery of meaningful
associations in genomic data using partial correlation coefficients. Bioinformatics, 20(18),
3565–3574.

Devidze N, Mong JA, Jasnow AM, Kow LM, Pfaff DW. (2005). Sex and estrogenic effects
on coexpression of mRNAs in single ventromedial hypothalamic neurons. Proc. Natl. Acad.
Sci. USA, 102(40), 14446–14451.

Dulac C, O'Connell LA, Wu Z. (2014). Neural control of maternal and paternal behaviors.
Science, 345(6198), 765–770.

Dutta S, Mark-Kappeler CJ, Hoyer PB, Pepling ME. (2014). The steroid hormone
environment during primordial follicle formation in perinatal mouse ovaries. Biol. Reprod.
91(3):68. doi: 10.1095/biolreprod.114.119214

Dwyer J, Kelly DA, Bergan J. (2022). Brain-wide synaptic inputs to aromatase-expressing
neurons in the medial amygdala suggest complex circuitry for modulating social behavior.
eNeuro, 9(2). doi: 10.1523/ENEURO.0329-21.2021

Ehret G. (1987). Left hemisphere advantage in the mouse brain for recognizing ultrasonic
communication calls. Nature, 325(6101), 249–51.

Ehret G. (2006). "Hemisphere dominance of brain function - which functions are
lateralized and why?" in 23 Problems in Systems Neuroscience, eds. L van Hemmen, TJ
Sejnowski (New York, NY: Oxford University Press) 44–61.

Ehret G, Buckenmaier J. (1994). Estrogen-receptor occurrence in the female mouse brain:
Effects of maternal experience, ovariectomy, estrogen and anosmia. J. Physiol. Paris, 88(5),
315–329.

Ehret G, Jürgens A, Koch M. (1993). Oestrogen receptor occurrence in the male mouse
brain: modulation by paternal experience. NeuroReport, 4(11), 1247–1250.

Ehret G, Koch M. (1989). Ultrasound-induced parental behaviour in house mice is
controlled by female sex hormones and parental experience. Ethology, 80, 81–93.

Ehret G, Koch M, Haack B, Markl H. (1987). Sex and parental experience determine the
onset of an instinctive behavior in mice. Naturwissenschaften, 74(1), 47–47.

Ehret G, Koch M, Jirikowski GF. (1991). Hormone, ZNS und Fortpflanzungsverhalten.
Allergologie, 14, 383–94.

Elwood RW. (1985). Inhibition of infanticide and onset of paternal care in male mice (Mus
musculus). J. Comp. Psychol. 99, 457–467.

Fahrbach SE, Pfaff DW. (1982). "Hormonal and neural mechanisms underlying maternal
behavior in the rat," in The Physiological Mechanisms of Motivation, ed. DW Pfaff (New
York, NY: Springer) 253–285. doi.org/10.1007/978-1-4612-5692-2_9

Fleischer S, Slotnick BM. (1978). Disruption of maternal behavior in rats with lesions of
the septal area. Physiol. Behav. 21(2), 189–200.

Foidart A, Harada N, Balthazart J. (1995). Aromatase-immunoreactive cells are present in
mouse brain areas that are known to express high levels of aromatase activity. Cell Tissue
Res. 280(3), 561–574.

Garcia-Segura LM. (2008). Aromatase in the brain: not just for reproduction anymore. J.
Neuroendocrinol. 20(6), 705–712.

Geissler DB, Sabine Schmidt H, Ehret G. (2013). Limbic brain activation for maternal
acoustic perception and responding is different in mothers and virgin female mice. J. Physiol.
Paris, 107, 62–71.

Geissler DB, Schmidt HS, Ehret G. (2016). Knowledge about sounds—context-specific
meaning differently activates cortical hemispheres, auditory cortical fields, and layers in
house mice. Front. Neurosci. 10:98. doi: 10.3389/fnins.2016.00098

Getz LL, Carter CS, Gavish L. (1981). The mating system of the prairie vole, Microtus
ochrogaster: Field and laboratory evidence for pair-bonding. Behav. Ecol. Sociobiol. 8(3),
189–194.

Giatti, S., Diviccaro, S., Garcia-Segura, L.M., Melcangi, R.C. (2019). Sex differences in
the brain expression of steroidogenic molecules under basal conditions and after
gonadectomy. J Neuroendocrinol 31, e12736. https://doi.org/10.1111/jne.12736

Guoynes C, Marler C. (2020). "Paternal behavior from a neuroendocrine perspective." in
Oxford Research Encyclopedia of Neuroscience.
https://oxfordre.com/neuroscience/display/10.1093/acrefore/9780190264086.001.0001/acrefo
re-9780190264086-e-9

Hashikawa K, Hashikawa Y, Falkner A, Lin D. (2016). The neural circuits of mating and fighting in male mice. Curr. Opin. Neurobiol. 38, 27–37.

Hojo, Y., Hattori, T.-A., Enami, T., Furukawa, A., Suzuki, K., Ishii, H.-T., Mukai, H., Morrison, J.H., Janssen, W.G.M., Kominami, S., Harada, N., Kimoto, T., Kawato, S. (2004). Adult male rat hippocampus synthesizes estradiol from pregnenolone by cytochromes P45017alpha and P450 aromatase localized in neurons. Proc Natl Acad Sci U S A 101, 865–870. https://doi.org/10.1073/pnas.2630225100

Hojo, Y., Murakami, G., Mukai, H., Higo, S., Hatanaka, Y., Ogiue-Ikeda, M., Ishii, H., Kimoto, T., Kawato, S. (2008). Estrogen synthesis in the brain--role in synaptic plasticity and memory. Mol Cell Endocrinol 290, 31–43. https://doi.org/10.1016/j.mce.2008.04.017

Honda SI, Wakatsuki T, Harada N. (2011). Behavioral analysis of genetically modified mice indicates essential roles of neurosteroidal estrogen. Front. Endocrinol. 2:40. doi: 10.3389/fendo.2011.00040

Horrell ND, Acosta MC, Saltzman W. (2021). Plasticity of the paternal brain: Effects of fatherhood on neural structure and function. Dev. Psychobiol. 63(5), 1499-1520. doi: 10.1002/dev.22097

Hyer, M.M., Khantsis, S., Venezia, A.C., Madison, F.N., Hallgarth, L., Adekola, E., Glasper, E.R. (2017). Estrogen-dependent modifications to hippocampal plasticity in paternal California mice (Peromyscus californicus). Horm. Behav. 96, 147–155. https://doi.org/10.1016/j.yhbeh.2017.09.015

Ji Y xiao, Zhao M, Liu Y ling, Chen L sha, Hao P li, Sun C. Expression of aromatase and estrogen receptors in lumbar motoneurons of mice. Neurosci. Lett. 653, 7–11.

Kalisch M, Mächler M, Colombo D, Maathuis MH, Bühlmann P. (2012). Causal inference using graphical models with the R package pcalg. Journal of Statistical Software. 47, 1–26.

Koch M. (1990). Effects of treatment with estradiol and parental experience on the number and distribution of estrogen-binding neurons in the ovariectomized mouse brain. Neuroendocrinol. 51(5), 505–514.

Koch M, Ehret G. (1989a). Estradiol and parental experience, but not prolactin are necessary for ultrasound recognition and pup-retrieving in the mouse. Physiol. Behav. 45(4), 771–776.

Koch M, Ehret G. (1989b). Immunocytochemical localization and quantitation of estrogen-binding cells in the male and female (virgin, pregnant, lactating) mouse brain. Brain Res. 489(1), 101–112.

Koch M, Ehret G. (1991). Parental behavior in the mouse: effects of lesions in the entorhinal/piriform cortex. Behav. Brain Res. 42(1), 99–105.

Kohl J, Dulac C. (2018). Neural control of parental behaviors. Curr. Opin. Neurobiol. 49, 116–122.

Kohl J, Autry AE, Dulac C. (2017). The neurobiology of parenting: A neural circuit perspective. Bioessays, 39(1), 1–11.

Laws MJ, Kannan A, Pawar S, Haschek WM, Bagchi MK, Bagchi IC. (2014).
Dysregulated estrogen receptor signaling in the hypothalamic-pituitary-ovarian axis leads to
ovarian epithelial tumorigenesis in mice. PLoS Genetics, 10(3):e1004230. doi:
10.1371/journal.pgen.1004230

Lee H, Kim DW, Remedios R, Anthony TE, Chang A, Madisen L, et al. (2014). Scalable
control of mounting and attack by ESR1+ neurons in the ventromedial hypothalamus. Nature,
509(7502), 627–632.

Lenth R. (2022). emmeans: estimated marginal means, aka least-squares means. R package
version 1.8.0. https://CRAN.R-project.org/package=emmeans

Li Y, Lian Z, Wang B, Tai F, Wu R, Hao P, et al. (2015). Natural variation in paternal
behavior is associated with central estrogen receptor alpha and oxytocin levels. J. Comp.
Physiol. A, 201(3), 285–293.

Lonstein JS, De Vries GJ. (2000). Sex differences in the parental behavior of rodents.
Neurosci. Biobehav. Revs. 24(6), 669–686.

Makowski D, Ben-Shachar MS, Lüdecke D. (2019). bayestestR: Describing effects and
their uncertainty, existence and significance within the Bayesian framework. Journal of Open
Source Software, 4(40), 1541. https://doi.org/10.21105/joss.01541

Malikov V, Madeira MD. (2013). Regulation of ERα protein expression by 17β-estradiol in
cultured neurons of hypothalamic ventromedial nucleus. Neurochem. Res. 38(1), 82–89.

Marlin BJ, Mitre M, D'amour JA, Chao MV, Froemke RC. (2015) Oxytocin enables
maternal behaviour by balancing cortical inhibition. Nature, 520(7548), 499–504.

Martín-Sánchez, A., Valera-Marín, G., Hernández-Martínez, A., Lanuza, E., Martínez-
García, F., Agustín-Pavón, C. (2015). Wired for motherhood: induction of maternal care but
not maternal aggression in virgin female CD1 mice. Front. Behav. Neurosci. 9.
https://doi.org/10.3389/fnbeh.2015.00197

Menon R, Süß T, Oliveira VE de M, Neumann ID, Bludau A. (2022). Neurobiology of the
lateral septum: regulation of social behavior. Trends Neurosci. 45(1), 27–40.

Nilsson ME, Vandenput L, Tivesten Å, Norlén AK, Lagerquist MK, Windahl SH, et al.
(2015). Measurement of a comprehensive sex steroid profile in rodent serum by high-sensitive
gas chromatography-tandem mass spectrometry. Endocrinology, 156(7), 2492–2502.

Numan M. (2006). Hypothalamic neural circuits regulating maternal responsiveness
toward infants. Behav. Cogn. Neurosci. Rev. 5(4), 163–190.

Numan M. (2020). The Parental Brain: Mechanisms, Development, and Evolution. New
York: Oxford University Press.

Numan M, Insel TR. (2003). The Neurobiology of Parental Behavior. New York: Springer.

O'Connell LA, Hofmann HA. (2012). Evolution of a vertebrate social decision-making
network. Science, 336(6085), 1154–1157.

Okabe, S., Kitano, K., Nagasawa, M., Mogi, K., Kikusui, T. (2013). Testosterone inhibits
facilitating effects of parenting experience on parental behavior and the oxytocin neural
system in mice. Phys. Behav. 118, 159–164. https://doi.org/10.1016/j.physbeh.2013.05.017

Palomero-Gallagher N, Amunts K. (2022). A short review on emotion processing: a
lateralized network of neuronal networks. Brain Struct. Funct. 227(2), 673–684.

Paxinos G, Franklin K. (2001). The mouse brain in stereotaxic coordinates. 2nd ed. San
Diego: Academic Press.

Poindron P, Lévy F, Krehbiel D. (1988). Genital, olfactory, and endocrine interactions in
the development of maternal behaviour in the parturient ewe. Psychoneuroendocrinology, 13,
99–125.

R Core Team. (2022). R: A language and environment for statistical computing. R
Foundation for Statistical Computing, Vienna, Austria. https://www.R-project.org/

Rilling JK, Mascaro JS. (2017). The neurobiology of fatherhood. Curr. Opin. Psychol. 15,
26–32.

Risold PY, Swanson LW. (1997). Connections of the rat lateral septal complex. Brain Res.
Brain Res. Revs. 24, 115–195.

Rizzi-Wise CA, Wang DV. (2021). Putting together pieces of the lateral septum:
multifaceted functions and its neural pathways. eNeuro, 8(6):ENEURO.0315-21.2021. doi:
10.1523/ENEURO.0315-21.2021

Romero-Morales L, Martínez-Torres M, Cárdenas M, Álvarez C, Carmona A, Cedillo B, et
al. (2018). An increase in estradiol facilitates the onset of paternal behavior in the dwarf
hamster (Phodopus campbelli). Horm. Behav. 99, 35–40.

Rosenblatt JS, Ceus K. (1998). Estrogen implants in the medial preoptic area stimulate
maternal behavior in male rats. Horm. Behav. 33(1), 23–30.

Rosenblatt JS, Siegel HI. (1981). "Factors governing the onset and maintenance of
maternal behavior among nonprimate mammals." in Parental Care in Mammals, eds. DJ
Gubernick, PH Klopfer. (Boston, MA: Springer) 13–76. https://doi.org/10.1007/978-1-4613-
3150-6_2

Rosenblatt JS, Wagner CK, Morrell JI. (1994). Hormonal priming and triggering of
maternal behavior in the rat with special reference to the relations between estrogen receptor
binding and ER mRNA in specific brain regions. Psychoneuroendocrinology, 19, 543–552.

Schleicher G, Stumpf WE, Morin JK, Drews U. (1986). Sites of aromatization of
[3H]testosterone in forebrain of male, female and androgen receptor-deficient Tfm mice: an
autoradiographic study. Brain Res. 397(2), 290–296.

Shabalova, A.A., Liang, M., Zhong, J., Huang, Z., Tsuji, C., Shnayder, N.A., Lopatina, O.,
Salmina, A.B., Okamoto, H., Yamamoto, Y., Zhong, Z.-G., Yokoyama, S., Higashida, H.
(2020). Oxytocin and CD38 in the paraventricular nucleus play a critical role in paternal
aggression in mice. Horm. Behav. 120, 104695. https://doi.org/10.1016/j.yhbeh.2020.104695

Sheehan T, Paul M, Amaral E, Numan MJ, Numan M. (2001). Evidence that the medial amygdala projects to the anterior/ventromedial hypothalamic nuclei to inhibit maternal behavior in rats. Neuroscience, 106(2), 341–356.

Sheng HZ, Fields RD, Nelson PG. (1993). Specific regulation of immediate early genes by patterned neuronal activity. J. Neurosci. Res. 35(5), 459–467.

Sheng M, McFadden G, Greenberg ME. (1990). Membrane depolarization and calcium induce c-fos transcription via phosphorylation of transcription factor CREB. Neuron, 4(4), 571–82.

Shinohara Y, Hirase H, Watanabe M, Itakura M, Takahashi M, Shigemoto R. (2008). Left-right asymmetry of the hippocampal synapses with differential subunit allocation of glutamate receptors. Proc. Natl. Acad. Sci. USA, 105(49), 19498–19503.

Shinohara Y, Hosoya A, Yamasaki N, Ahmed H, Hattori S, Eguchi M, et al. Right-hemispheric dominance of spatial memory in split-brain mice. Hippocampus, 22(2), 117–121.

Shipley B. (2000). Cause and Correlation in Biology: a User's Guide to Path Analysis, Structural Equations, and Causal Inference in R. Cambridge, UK: Cambridge University Press.

Shipton OA, El-Gaby M, Apergis-Schoute J, Deisseroth K, Bannerman DM, Paulsen O, et al. (2014). Left–right dissociation of hippocampal memory processes in mice. Proc. Natl. Acad. Sci. USA, 111(42), 15238–15243.

Siegel HI, Rosenblatt JS. (1975). Estrogen-induced maternal behavior in hysterectomized-ovariectomized virgin rats. Physiol. Behav. 14(4), 465–471.

Slotnick BM, Nigrosh BJ. (1975). Maternal behavior of mice with cingulate cortical, amygdala, or septal lesions. J. Comp. Physiol. Psychol. 88(1), 118.

Stanić D, Dubois S, Chua HK, Tonge B, Rinehart N, Horne MK, et al. (2014). Characterization of aromatase expression in the adult male and female mouse brain. I. Coexistence with oestrogen receptors α and β, and androgen receptors. PLoS ONE, 9(3):e90451. doi: 10.1371/journal.pone.0090451

Stolzenberg DS, Rissman EF. (2011). Oestrogen-independent, experience-induced maternal behaviour in female mice. J. Neuroendocrinol. 23(4), 345–354.

Tachikawa KS, Yoshihara Y, Kuroda KO. (2013). Behavioral transition from attack to parenting in male mice: a crucial role of the vomeronasal system. J. Neurosci. 33(12), 5120–5126.

Torres, T., Adam, N., Mhaouty-Kodja, S., Naulé, L. (2024). Reproductive function and behaviors: an update on the role of neural estrogen receptors alpha and beta. Front. Endocrinol. 15. https://doi.org/10.3389/fendo.2024.1408677

Trainor BC, Marler CA. (2002). Testosterone promotes paternal behaviour in a monogamous mammal via conversion to oestrogen. Proc. R. Soc. Lond. B, 269(1493), 823–829.

Trainor BC, Bird IM, Alday NA, Schlinger BA, Marler CA. (2003). Variation in aromatase
activity in the medial preoptic area and plasma progesterone is associated with the onset of
paternal behavior. Neuroendocrinol. 78(1), 36–44.

Trainor BC, Finy MS, Nelson RJ. (2008). Paternal aggression in a biparental mouse:
parallels with maternal aggression. Horm. Behav. 53(1), 200–207.

Unger EK, Burke KJ, Yang CF, Bender KJ, Fuller PM, Shah NM. (2015). Medial
amygdalar aromatase neurons regulate aggression in both sexes. Cell Reports, 10(4), 453–
462.

van Buuren S, Groothuis-Oudshoorn K. (2011). mice: multivariate imputation by chained
equations in R. Journal of Statistical Software, 45, 1–67.

Vehtari A, Gelman A, Simpson D, Carpenter B, Bürkner PC. (2021). Rank-normalization,
folding, and localization: an improved Rˆ for assessing convergence of MCMC (with
discussion). Bayesian Analysis, 16(2), 667–718.

Wickham H. (2016). ggplot2: elegant graphics for data analysis. New York: Springer-
Verlag.

Woodroffe R, Vincent A. (1994). Mother's little helpers: Patterns of male care in mammals.
Trends Ecol. Evol. 9(8), 294–297.

Wozniak A, Hutchison RE, Hutchison JB. (1992). Localisation of aromatase activity in
androgen target areas of the mouse brain. Neurosci. Lett. 146(2), 191–194.

Wu Z, Autry AE, Bergan JF, Watabe-Uchida M, Dulac CG. (2014). Galanin neurons in the
medial preoptic area govern parental behaviour. Nature, 509(7500):325–330.

Xiao L, Jordan CL. (2002). Sex Differences, Laterality, and Hormonal Regulation of
Androgen Receptor Immunoreactivity in Rat Hippocampus. Horm. Behav, 42(3), 327–336.

Yang T, Yang CF, Chizari MD, Maheswaranathan N, Burke KJ, Borius M, et al. (2007).
Social control of hypothalamus-mediated male aggression. Neuron, 95(4), 955-970. doi:
10.1016/j.neuron.2017.06.046

Table 1. Brain areas analyzed for aromatase expression, their Bregma coordinates and plate

numbers from the mouse brain atlas (Paxinos and Franklin, 2001), and sample sizes per group

per brain area.

Brain area		Bregma Coordinates	Plate number	Sample Size
Lateral septum dorsal (LSD)	Anterior and posterior	0.98 mm and 0.14 mm	23 and 30	n=7 (N) n=10 (5D) n=11 (27D)
Lateral septum intermediate (LSI)	Anterior and posterior	0.98 mm and 0.14 mm	23 and 30	n=7 (N) n=10 (5D) n=11 (27D)
Lateral septum ventral (LSV)	Anterior and posterior	0.98 mm and 0.14 mm	23 and 30	n=7 (N) n=10 (5D) n=11 (27D)
Piriform cortex (PIR)	Anterior and posterior	0.98 mm and -1.34 mm	23 and 42	n=7 (N) n=8 (5D) n=9 (27D)
Nucleus accumbens (NAC)		0.98 mm	23	n=7 (N) n=10 (5D) n=11 (27D)
Bed nucleus stria terminalis (BNST)		0.14 mm	30	n=7 (N) n=10 (5D) n=11 (27D)
Medial preoptic area (MPOA)		0.14 mm	30	n=7 (N) n=10 (5D) n=11 (27D)
Cortical amygdala (CAM)		-1.34 mm	42	n=7 (N) n=9 (5D) n=8 (27D)
Medial amygdala (MAM)		-1.34 mm	42	n=7 (N) n=9 (5D) n=8 (27D)
Ventromedial hypothalamus (VMH)		-1.34 mm	42	n=7 (N) n=9 (5D) n=8 (27D)

    Legends to the Figures

Figure 1. Behavioral responses towards pups. A. Experimental setting for the pup-

retrieving tests on a running board with a central nest depression. Responses were scored as

aggression (attacking pups), ignoring pups, or retrieval of pups (returning pups to the nest). **B.**

Probabilities of behavioral expression in naïve (N) males and in fathers with five (5D) or 27

days of experience (27D). Aggression declined immediately from N to 5D fathers. Although

fathers occasionally still ignored pups, the probability of retrieval was very high. Different

letters indicate statistical differences between the groups of males. Sample size was n=8 for

N, n=12 for 5D, and n=12 for 27D.

Figure 2. Aromatase immunolabeling in the paternal brain. Cross sections (**A**) of the

brain areas analyzed here (Bregma coordinates: 0.98 mm; 0.14 mm; and -1.34 mm). Boxes

indicate approximate areas that were counted. Representative images from areas highlighted

in (**A**), showing aromatase immunostaining in naïve and experienced fathers (5D and 27D) in

the left (L) and right (R) hemisphere and lack of immunostaining in the no primary antibody

controls. High magnification (**B**) images of examples of aromatase expressing cells in the

MPOA, VMH, and MAM showing cytoplasmic expression and black granulae. In cases in

which the granulae were located perinuclear to an identifiable cell nucleus (marked with *)

the cells were counted as aromatase-positive. In cases in which the granulae could not be

associated with a cell nucleus (marked with §) the structure was not counted as an aromatase-

positive cell.

Figure 3. Aromatase expression differentially depends on paternal experience and brain

hemisphere. **A.** Standardized cell counts (light grey circles) and means with ± 95%

confidence interval (white circles) for each investigated brain area separately for the left and

right sides of the brain and levels of pup-experience (N, 5D, 27D). **B.** Hemisphere contrast

(left – right, with 95% confidence intervals) of the standardized counts of aromatase positive

cells for each brain area and experimental group. Contrasts with intervals significantly

different ($p < 0.05$) from zero are highlighted in purple (see text for description). We found

evidence for left-hemisphere dominant lateralization (left minus right difference was positive)

of aromatase cell counts both independent of (BNST, MPOA, and most likely NAC) and

dependent on pup-experience (dorsal LS), and an experience-dependent right-hemisphere

dominant lateralization in the intermediate and ventral LS and anterior PIR (Bregma 0.98). **C.**

Experience contrast (5D–N; 27D–N, 27D–5D, with 95% confidence intervals) for the

standardized counts of aromatase positive cells, separately for each hemisphere and brain

nucleus/area. We found that the occurrence of aromatase positive cells in posterior (Bregma -

1.34) PIR, CAM, MAM, and VMH in both brain hemispheres significantly increased with the

experience level. Expression of aromatase in the LS and anterior PIR significantly increased

only in the right hemisphere of the 27D group. No changes were observed in the NAC and

BNST. Contrasts with intervals significantly different ($p < 0.05$) from zero are highlighted in

purple (see text for description). Sample sizes are given in Table 1.

Figure 4. Residual correlations among brain nuclei/areas viewed as a network. Each

nucleus/area at each Bregma coordinate (see Table 1) is represented by a node, with edges

representing significant partial correlations (solid for $p < 0.05$, dashed for $p < 0.10$). Residual

correlations were positive between all brain areas, with the sole exception of CAM (-1.34)

and LSD (0.98), which were negatively correlated. PIR (-1.34) was not correlated with any

other brain area.

Figure 5. Aromatase expression in males that react to pups compared to those ignoring

pups. Contrasts (reacting – ignoring, with 95% confidence intervals) of the standardized

counts of aromatase positive cells for each brain area. Contrasts with intervals significantly

different (p < 0.05) from zero are highlighted in purple (see text for description). We found

that reacting males had significantly higher numbers of aromatase cells in the MAM

(p=0.011), and a weak tendency to higher numbers in the CAM (p=0.06).

Figure 6. Effects of paternal experience on aromatase expression in areas of the limbic

system. Sagittal view of the brain areas analyzed here (A) and summary of the changes in

aromatase expression in 5D **(B)** and 27D **(C)** males in the left and right hemispheres of the

brain nuclei/areas analyzed in the current study. **A.** Positions and extensions of nuclei/areas

are illustrative only and do not indicate anatomical relationships. The connectivity denoted in

(A) shown by the arrows are exemplars for main information pathways (Numan, 2020; Kohl

et al., 2017). **B.** Pup experience in first-litter fathers (5D group) increased the numbers of

aromatase positive cells (indicated by light purple) relative to naïve males in the amygdala

(CAM and MAM), hypothalamus (VMH), and in posterior piriform cortex (PIR, half shading

indicates these partial changes). **C.** Second-litter fathers (27D group) had further increased

aromatase expression in the same areas as 5D males (indicated as darker purple), i.e., the

amygdala (CAM and MAM), hypothalamus (VMH), and posterior piriform cortex (PIR) in

both sides of the brain. In the right hemisphere, the number of aromatase positive cells

increased (indicated by blue) in the anterior PIR, lateral septum (LS) relative to naïve and 5D

males, and potentially in the MPOA relative to 5D males.

 FIGURE 1

933 **FIGURE 2**

FIGURE 3

939 **FIGURE 4**
940

 FIGURE 5

	Region	Bregma	Reacting – Ignoring
Lateral septum (dorsal)	LSD	0.14	
Lateral septum (dorsal)	LSD	0.98	
Lateral septum (intermediate)	LSI	0.14	
Lateral septum (intermediate)	LSI	0.98	
Lateral septum (ventral)	LSV	0.14	
Lateral septum (ventral)	LSV	0.98	
Piriform cortex	PIR	0.98	
Piriform cortex	PIR	-1.34	
Nucleus accumbens	NAC	0.98	
Bed nucleus of the stria terminalis	BNST	0.14	
Medial preoptic area	MPOA	0.14	
Cortical amygdala	CAM	-1.34	
Medial amygdala	MAM	-1.34	
Ventromedial hypothalamus	VMH	-1.34	

Response contrast

945 **FIGURE 6**
946

A

B

C