Brain aromatase dynamics reflect parental experience and behaviour in male mice

Paula Duarte-Guterman[1], Dimitri A. Skandalis[2]*, Ariane Merkl*, Diana B. Geissler, Günter

Ehret

Institute of Neurobiology, University of Ulm, D-89069 Ulm, Germany

* Contributed equally to this project

[1] Present address: Department of Psychology and Centre for Neuroscience, Brock University,

ON, Canada.

[2] Present address: Psychological & Brain Sciences and Johns Hopkins Kavli Neuroscience

Discovery Institute, Johns Hopkins University, MD, USA

Corresponding authors:

Paula Duarte-Guterman; email: pduarteg@brocku.ca

Günter Ehret; email: guenter.ehret@uni-ulm.de

Keywords: brain plasticity, estrogen, hemisphere lateralization, infanticide, limbic system,

paternal care

Abbreviations: BNST, bed nucleus of the stria terminalis; CAM, cortical amygdala; LS,

lateral septum; LSD, lateral septum dorsal; LDI, lateral septum intermediate; LSV, lateral

septum ventral; MAM, medial amygdala; MPOA, medial preoptic area; NAC, nucleus

accumbens; PIR, piriform cortex; VMH, ventromedial nucleus of the hypothalamus.

Abstract

In most mammals, providing paternal care is not automatic. In house mice, experience with pups governs the extent and quality of paternal care. First-time fathers undergo a dramatic transition from ignoring or killing pups to caring for pups. The behavioural shift occurs together with changes in brain estrogen signalling as indicated by changes in estrogen receptor presence and distribution in multiple areas regulating olfaction, emotion, and motivation. Here, we report estrogen dynamics by altered local estrogen synthesis via changes in aromatase, a key enzyme catalysing the conversion of testosterone to estrogen. The amount of paternal experience (5 or 27 days) was associated with increased numbers of immunocytochemically-identified aromatase expressing cells in the medial and cortical amygdala, piriform cortex, and ventromedial hypothalamus. In the lateral septum, and to some extent in the medial preoptic area, parental experience increased aromatase only in fathers with 27 days of experience, and only in the right brain hemisphere, a new case of brain-functional lateralisation with experience. Nuclei/areas associated with maternal care (medial preoptic area, bed nucleus of stria terminalis, nucleus accumbens) exhibited a left-hemisphere advantage in aromatase expressing cells. This newly found lateralisation may contribute to the left-hemisphere dominant processing and perception of pup calls to release parental behaviour.

Introduction

Mammalian paternal care is facultative, but the behavioural and neural dynamics that result in becoming a caring father are less well understood compared to those of obligatory maternal care (e.g., (1–10)). In rodents and primates, it is hypothesised that maternal care is induced by hormonal changes during pregnancy and parturition (onset phase) whereas the continuation of pup-caring (maintenance phase) is governed by sensory stimuli from the pups (2,11,12). In house mice, pup-naïve females, males, and even previous fathers do not experience the onset phase but require various amounts of experience with pups (hours up to several days of contact) for the activation of caring behaviour (13–21). Nonetheless, pup-naïve females engage in consummatory (pup retrieval) and appetitive (preference for pup ultrasounds) behaviours (22,23) after fewer days of experience with pups than either pup-naïve males or fathers (14–16). Pup naïve females thus appear to be more easily sensitised or motivated by pup cues than males and even fathers to provide parental care, suggesting sex differences in neural mechanisms underlying the activation of maternal versus paternal instincts.

Numerous studies have characterised the interactions between limbic system nuclei and areas for the activation and regulation of parental behaviours. Principal nuclei and areas involved in parental care include the medial preoptic area (MPOA), the bed nucleus of the stria terminalis (BNST), the lateral septum (LS), the nucleus accumbens (NAC), nuclei of the amygdala, and the ventromedial hypothalamus (VMH). The main modulatory sensory inputs are thought to be olfaction via the vomeronasal organ to the BNST and amygdala, and via the olfactory bulb either to the amygdala and via the VMH to the LS or via the piriform cortex (PIR) and hippocampus to the LS (e.g., (2,3,7,23–26)). In females, neurons in these regions express estrogen receptors and so may be affected by levels of circulating estrogens during the course of the estrous cycle and during pregnancy and after parturition (12). However, estradiol treatment can activate both maternal and paternal instincts (17,19,27–31). In house

mice, expression of parental behaviours coincides with changes in the distribution of estrogen

receptor (ERα) positive cells in limbic brain regions of pup-naïve and pup-experienced

females (16,32) and males (14). For instance, in the MPOA of females and fathers, the

number of ERα positive cells increased with pup contact and retrieving experience (14,16,32).

In pup-naïve male and female voles, animals with spontaneously high levels of pup care

showed more ERα positive cells in the MPOA than animals with low levels (33).

Qualitatively similar distributions of ERα positive cells were observed in the limbic system

between pup-naïve males after five days of experience (14) and pup-naïve females (34),

potentially suggesting some levels of convergence in terms of the underlying circuit

dynamics.

Unlike in females, low circulating levels of estrogens in males (35) imply that

estrogens are primarily synthesised locally in the brain from testosterone by the enzyme

aromatase (e.g., (36–38)). Locally synthesised estrogens have been implicated in the onset of

paternal behaviours in California mice (39,40) and aromatase activity has been documented in

the brains of adult male mice (41–43). Aromatase is expressed in the male mouse limbic

system (LS, BNST, MPOA, nuclei of the amygdala, VMH, and PIR), in regions where ERα

positive cells have been located in pup-experienced fathers (14). This suggests that estrogen

signalling via ERα could be an important regulator of paternal behaviour via an aromatase-

dependent estrogen supply. Our objective was therefore to examine changes in aromatase

distributions during the onset and maintenance of paternal behaviour of male mice (pup-naïve

and fathers) in brain regions known to exhibit changes in ERα positive cells (14).

Materials and methods

Animals

Adult male and female laboratory mice (outbred strain NMRI, 2-3 months old) with no

previous sexual or parental experience were housed at the University of Ulm. As previously

(14), mice were randomly assigned to one of three treatment groups consisting of (1) naïve

males without any experience with pups (N); (2) males co-caring with their female mate for

their first litter for five days (5D); or (3) males co-caring with their female mate for their first

litter until weaning and their second litter for five days. The total co-caring time of the latter

males (first and second litter) was 26-27 days (27D) since after delivery of the first litter,

females in post-partum estrus paired immediately and, while nursing the first litter up to

weaning at 21 days, delivered the second litter after 21-22 days of gestation. Mice were

housed at 22ºC and a 12 h light-dark cycle (starting at 7.00 h) in plastic cages ($26.5 \times 20 \times 14$

cm) either with other naïve males (group N; 4 brothers per cage) or with one female (groups

5D and 27D). Water and food were available *ad libitum*. The experiments were carried out in

accordance with the European Communities Council Directive (86/609/EEC) and were

approved by the appropriate authority (Regierungspräsidium Tübingen, Germany).

Pup retrieval test

Testing of pup retrieval behaviours was done as previously described (14,17). Male

mice (alone or with litter mate and pups) were placed overnight on a running board (length

110 cm, width 8 cm) with a central nest depression (Fig. 1A) suspended in a sound-proof and

anechoic room. The running board was supplied with mouse chow and nest material from the

cage and covered with a plexiglass hood having an opening for the tip of a suspended water

bottle. Retrieving tests were done under dim red light. Before the start of testing, the hood,

and the female with the pups (groups 5D and 27D) were removed and placed back to their

home cage outside the soundproof room. Seven pups were kept in the room to serve as targets

for retrieving. The males were habituated for at least 30 min to the new situation after the

hood (group N) or the hood and female with pups (groups 5D and 27D) were removed. Prior

to testing males of the N group, five-day old foster pups were rubbed with nest material of the

respective males to ensure a familiar scent. During the test, the seven pups (always 1-3 pups

at a time) were randomly placed on the running board, at least 30 cm away from the nest. A

male was scored as 'retriever' if he retrieved all seven pups to the nest within 10 min. A male

was scored as 'non-retriever' if he had not retrieved any pups after 10 mins. The distinction

between retriever and non-retriever was clear-cut, because once a male had retrieved the first

pup, he retrieved all. Retrieving tests were immediately stopped and the pups removed if a

male attacked any of the pups ('aggressor').

Immunocytochemistry and microscopy

Immediately following the pup retrieval test, males were euthanised by cervical

dislocation and their brains quickly dissected and frozen over liquid nitrogen. Brain serial

frontal sections (30 μm) were cut on a freezing microtome (Thermo Scientific Microm

HM560), mounted on slides, and immediately fixed in 4% paraformaldehyde in 0.1 M

phosphate buffer (PB; pH=7.4) for 60 min. Sections were rinsed three times for 5 min with

0.1 M PB in between all subsequent processing steps. Slides were incubated with 0.2% Triton

X-100 (Sigma) in 0.1 M PB for 45 min to permeabilise cells and with 1% H_2O_2 (Merck) in

0.1 M PB for 20 min to block endogenous peroxidase. In a humidified chamber, the sections

were treated with 2% normal goat serum in 0.1 M PB for 60 min to block non-specific

staining, then with the polyclonal rabbit anti-aromatase antibody (Abcam Cat# ab35604,

RRID:AB_867729; 0.002 mg/ml in 2% normal goat serum) at 4ºC for 72h. Sections were

then incubated with the secondary antibody from goat against rabbit horseradish peroxidase

(Dako, Cat# PO448, 0.00125 mg/ml in 0.1 M phosphate buffer) at room temperature for 60

min, followed by staining with 0.015% diaminobenzidine tetrachloride (DAB; Sigma),

0.023% NiCl, and 0.013% H_2O_2 in 0.1 M PB for 7 min. Sections were rinsed in PB,

dehydrated in a series of alcohols and xylene, and coverslipped with Entellan (Merck). The

specificity of the aromatase antibody has previously been tested and confirmed for the mouse

brain (44). We further controlled for non-specific labeling by incubating sections with PB

instead of the primary antibody. In all these control sections only non-specific background

labelling was present. Aromatase-specific labelling was absent.

Sections were independently analysed by two experimenters blind to the source of the

sections. In view of varying non-specific background labelling, the experimenters started the

analysis by checking many sections from all animal groups in order to agree which kind of

labelling would be considered aromatase-specific. Cells were considered as expressing

aromatase if they were more intensely stained with shades of gray than the background and

contained black granulae often grouped around the cell nucleus (Supplement Fig. 1). In cases

in which a single or several black granulae without an associated cellular structure in shades

of gray was seen, the structure was not counted as an aromatase-positive cell. This procedure

led to (a) a high concordance of counting results of the two experimenters, (b) counts of only

definite aromatase-positive cells, and (c) relatively low numbers of aromatase-positive cells

per evaluated brain area.

Sections were analysed with a light microscope (Axiophot, Zeiss Germany). Brain

areas were identified with 10x objective magnification, aromatase-positive cells were

identified by focussing with a 40x objective magnification. The brain areas in which

aromatase-positive cells were counted are listed in Table 1. These areas largely overlap with

those showing ERα positive cell dynamics during the acquisition of pup-caring experience in

male (14) and female (16) mice. The Bregma coordinates of the brain areas were from the

mouse brain atlas (45) and anatomically matched across all experimental animals. Aromatase-

positive cells were counted separately on the right and left hemispheres of the brain. Counts

were averaged between two (LSD, LSI, LSV, BNST, NAC, VMH) or three (PIR, MPOA,

CAM, MAM) sections for each animal (Supplemental Fig. 2).

Statistical Analyses

All statistical analyses were carried out using R v4.2.0 (46). Male behaviour in

response to pup presentation was classified as ignoring, attacking, or retrieving a pup. The

outcomes of behavioural trials were analysed as the multinomial probability of exhibiting a

given response to pup presentation (aggression, ignore, retrieval) as a function of paternal

experience (experience as ordered factor, N < 5D < 27D). To perform pairwise comparisons

between the groups for each behaviour, we calculated the probability that the difference in the

posterior predictive intervals between groups was significantly directional. We then corrected

for multiple comparisons using the false discovery rate.

We combined two partially overlapping datasets of aromatase cell counts from two

cohorts of animals to increase our statistical power to identify variation among cohorts and

brain regions (Supplemental Fig 2). We observed significant count variation attributable to

cohort (e.g., count variation between cohorts for the PIR -1.34 data; Supplemental Fig 2). To

address this issue, each dataset was standardised by the mean and variance of paternal groups

(5D and 27D; the groups for which the datasets were overlapping). Counts for some brain

regions were missing in some mice. Rather than rejecting these mice altogether in our

statistical model (described below), we imputed the missing observations based on values

inferred from the non-missing entries (R package mice, 47). We generated five such candidate

data sets and integrated over the resulting uncertainty.

Our statistical model of aromatase variation was designed to address non-

independence arising from correlated variation within individuals among regions, which could

be an especially important factor when comparing identifiable subregions of an area, such as

in the piriform cortex and each part of the lateral septum. We therefore examined variation in

normalised cell counts among all brain regions simultaneously (multivariate ANOVA),

treating mouse ID as a correlated group (random) effect. Responses were modelled as *t*-

distributed to increase robustness against outlier counts. We examined the interaction of

Experience x Hemisphere together with the residual correlations among regions, which

represent relationships among brain regions which are not explained by the other population-

and group-effects.

All models were fit using the R package *brms* (48), using four chains of 2500

iterations each, and a thinning rate of 5. We assumed uniform priors throughout. Typically,

model convergence would be assessed by $\hat{R}=1$ (49) but because each imputed data set will

generate slightly different estimates at convergence, $\hat{R}$ may be misleading. We therefore

assured model convergence in a separate analysis of a single data set generated by averaging

over the imputed data sets. We calculated the posterior means of each group based on the

fitted statistical model (*emmeans* package (50)), and used these to examine the posterior

distributions of contrasts between hemispheres (left-right) and between levels of experience

(5D-N, 27D-N, 27D-5D) within each hemisphere. Significance of each contrast was judged as

a Bayesian probability of direction $P>0.975$ (51), which is an index of effect existence, i.e.,

our level of certainty in whether a reported change is positive or negative. This threshold

closely corresponds to the frequentist two-sided *p*-value of 0.05, and so for simplicity, we

report the *p*-value. Plots were generated using the *ggplot2* package (52).

Residual correlations can point to latent (hidden) factors that determine correlated

expression across the brain and/or unexpected technical covariation, so inspection of residual

correlations can inform biological hypotheses. However, direct interpretation of the residual

correlation matrix is challenging because the correlation between two brain regions could be

due to mutual covariation with another region. We therefore dichotomised relationships

among brain regions by studying the matrix of *partial* correlations (53,54). We constructed an

initial graph in which edges were created between all nodes with significant ($p<0.05$) pairwise

correlations. Each edge was iteratively pruned by testing whether the association remained

statistically significant after testing it against all other paths between two variables, up to third

order. The independence test was determined by the posterior distribution of residual

correlation matrices implemented in the R package pcalg (55). The result is an undirected,

acyclic graph representing a network of associations among brain regions. The large number

of tests may be conservative and may reject true associations, and thus we also report

tentative links observed when setting a less stringent threshold of $p<0.10$.

Results

Male mice retrieve pups after acquiring parental experience

Only males with paternal experience retrieved pups (Fig. 1). Naïve males with no

previous sexual experience and pup contact mostly ignored the pups (n = 5/8), although some

were aggressive (n = 3/8). Most fathers with 5 days (5D) and 26-27 days (27D) of experience

retrieved pups (n = 10/12 and 11/12, respectively). A small number of fathers ignored pups in

the 5D (n = 2/12) and 27D groups (n = 1/12) but none were aggressive.

The probability of observing aggressive behaviour toward pups was significantly

higher in naïve males compared to fathers regardless of experience (5D and 27D, Fig. 1B).

The probability of other non-paternal behaviour like ignoring pups was higher in naïve males

compared to 27D fathers. Parental behaviour measured as pup retrieval was significantly

higher in fathers (5D and 27D) compared to naïve males. There were no significant

differences in the probability of pup-retrieving between the groups of fathers, 5D and 27D

(Fig. 1B).

Aromatase expression in the male mouse brain, general picture

We found aromatase-positive cells in each examined region and with each level of

experience. In absolute terms, largest numbers of aromatase positive cells per section were

seen in PIR, LSI, MAM, and CAM (see unstandardised data in Supplement, Fig. 2). Examples

of aromatase labeling in the LSV, PIR, VMH and MAM of experimental groups N and 27D

show qualitative differences between the groups and between left and right sides of the brain

(Fig. 2). Standardised cell counts together with the group means are shown in Fig 3A along

with our estimates of the posterior differences in aromatase counts by hemisphere (Left –

Right, Fig. 3B), and between levels of experience (Fig. 3C).

Lateralised expression of aromatase positive cells

We found evidence for significantly greater numbers of aromatase positive cells in the

left hemisphere of the MPOA and BNST in all three experimental groups (Left – Right > 0,

Fig. 3B). The NAC exhibited a similar trend to MPOA and BNST, but with a weakly

insignificant contrast in the 27D group. We did not detect significant lateralisation of

aromatase counts in any of the experimental groups in the amygdala (CAM and MAM),

VMH, or posterior PIR (Bregma -1.34) in any groups. In the other regions, the appearance of

lateralisation strongly depended on the level of experience. Long-term pup-experience (27D)

led to greater numbers of aromatase positive cells in the right hemispheres of anterior PIR

(Bregma 0.98), the LSV, and the anterior LSI (Bregma 0.14). Posterior LSI (Bregma 0.98)

exhibited a similar trend that was weakly non-significant. In the LSD, pup-experience had a

short-term effect (5D) during which counts in the left hemisphere were greater, which was not

present in the 27D group (Fig. 3B).

Experience-dependent expression of aromatase

In all cases in which we found significant changes of aromatase counts with

experience, the number of aromatase positive cells increased (Fig. 3C). This experience-

dependent increase was pervasive for the amygdala (CAM and MAM), the VMH, and the

posterior PIR (Bregma -1.34). For these regions, the increase in aromatase positive cells was

present in both hemispheres and with increasing experience (i.e., experience contrasts had

significant positive values for both brain hemispheres and all experimental groups). In the

LSD, LSI, and LSV (both Bregma coordinates), and the anterior PIR (Bregma 0.98),

aromatase positive cells increased significantly only in the right hemisphere of 27D males.

Experience-dependent effects were absent in the NAC, the BNST, and the MPOA, with the

caveat of a significant effect in the right MPOA of 27D males (27D males had significantly

more aromatase positive cells than 5D males). In fact, both BNST and MPOA exhibit trends

toward increasing aromatase with experience, but this variation was not statistically

significant (Fig. 3C).

Graphical analysis of covariation among brain regions

We examined the unexplained (residual) variation among all brain regions in the study

(represented graphically in Fig. 4). Three subgroups of covariation are obvious. First, LS is

highly connected (although not in order from dorsal to ventral), and each subregion (D, I, V)

is correlated across Bregma coordinates and to at least one other subregion. LS appears to

form the backbone of variation, to which other nodes are connected. This includes subgroups

comprised of MPOA (connected to LSV), of NAC and BNST (connected to LSI), and of

CAM and VMH (connected to LSD). Second, the subgroup of MAM and anterior PIR

appears to be weakly connected to other parts of the graph (MPOA and VMH), because the

edges are significant only with $p<0.10$. Third, posterior PIR did not appear to be even weakly

correlated with any other analysed part of the brain despite an increase in aromatase

expression with experience that is similar to CAM, MAM, and VMH (see Fig. 3).

Discussion

Aromatase is expressed throughout the male brain at considerable levels (41,43) , even

in pup-naïve male mice (Fig. 3A). Together with this widespread distribution, levels of

aromatase vary in an experience-dependent manner (Fig. 3). In mice, fathers that exhibit pup

retrieval, have higher levels of aromatase immunoreactivity in certain limbic regions (VMH,

amygdala, MPOA, NAC, LSD) after social contact with the female partner following retrieval

tests (13). In mice, changes in aromatase expression have been observed in fathers in response

to pup and social experience, e.g. contact with the the mating partner after separation (13). In

addition, inhibition of aromatase in fathers from day 1 to day 5 after birth of their first litter

significantly inhibited the occurrence of pup-retrieval (13), which needs more than 1 day of

pup contact to be securely established in untreated fathers (15). We therefore consider it likely

that the presence and increase in brain aromatase expression observed here is integral to the

adoption of parental behaviours in house mice. However, we find that the change in

aromatase levels varies non-uniformly across limbic brain regions, broadly summarised in

Fig. 5. Moreover, the individual expression of parental behaviours will likely depend not only

on overall levels but the correlated changes among brain regions. We found significant

residual correlations in aromatase counts among brain regions, pointing to intrinsic sources of

variation in aromatase among individuals. Such correlations may ultimately contribute to

understanding why some fathers ignored pups even though none attacked them. Here, we

briefly review the contributions of limbic areas to paternal behaviours and experience.

Amygdala and ventromedial hypothalamus

Three nuclei, the MAM, CAM, and VMH were most affected by pup experience with

large increases in the number of aromatase positive cells from N to 5D to 27D animals (Fig. 3

A and C; Supplement Fig. 2). These regions are associated with the regulation of aggression

compared to sexual and parental behaviour (e.g., (56–62)). Important changes in aggression in

the course of becoming a caring father are mediated by olfactory signals from the

vomeronasal organ via the accessory olfactory bulb to the CAM and MAM (e.g., (58)) and

concern the switch from aggressive to caring behaviour towards pups (21). Concomitantly,

aggressive motivation in fathers increases against male intruders in order to protect their own

pups from potential pup killers as shown in Californian mice (63) and prairie voles (64). This

increased aggression towards intruders was not mediated by changes in ERα or ERß

expression in limbic centers, but reflected in increased neuronal activation (FOS-expression)

in the MPOA, MAM and, potentially the VMH (63). We are not aware of respective tests in

male house mice, or whether the decrease in aggression towards pups likewise alters

aggression toward potentially dangerous intruders. Such tests combined with aromatase

measurements could clarify whether the differentiation of aggression against kin or non-kin

depends on aromatase levels in the amygdala and VMH.

Notwithstanding the increased numbers of aromatase-positive cells, a long-term

increase in the number of estrogen receptors (ERα) in these nuclei has not been observed (14).

This might have been expected because the presence of estrogen can induce the expression of

its own receptors, for example in the VMH (65,66), especially in combination with parental

experience (32). We suggest that the increase in aromatase activity in the MAM, CAM, and

VMH in new (5D) and highly experienced (27D) fathers (Fig. 5) supports the control of

instinctive behaviour (67) via increased local estrogen action (e.g., (36,37)).

Piriform cortex

In house mouse mothers, the performance of pup-retrieving does not depend on

olfactory signals from the main olfactory bulb and its projections to the PIR (22). Pup-naïve

females, however, needed this olfactory information for learning to become as maternal as the

mothers (16,22). While learning pup cues for becoming maternal, virgin females with

maternal experience have an increase in ERα positive cells in the posterior PIR compared to

females without experience (also in the entorhinal cortex; (32)). Similarly, as males became

caring fathers, ERα positive cells become visible in the posterior PIR (also in the entorhinal

cortex; (14)). In view of the significant increase of aromatase expression in the posterior PIR

with increasing paternal experience (Fig. 3C), it is possible that in PIR, unlike MAM, CAM,

and VMH, may have induced its own receptors (tested for ERα) and enhanced olfactory

learning of pup cues for increased paternal performance. The significant increase of aromatase

positive cells only in the right hemisphere in the anterior PIR of 27D fathers does not directly

correlate with any evidence known to us. In studies on ERα expression in the limbic centres

of female and male mice under various conditions of the reproductive cycle, experience with

pups, and intact or lesioned olfaction, no hemisphere differences have been reported in any

centre (14,16,22,32). The splitting of PIR in regions with experience-dependent lateralised

(anterior PIR) and non-lateralised (posterior PIR) aromatase change shows

compartmentalization of the PIR. Functional relationships need to be elucidated in further

studies.

Lateral septum

The LS has a central functional role in the brain integrating cognitive and emotional

information in the regulation of social relationships and instincts (68–70) including parental

behaviour (71,72). Our analysis of residual correlations likewise (Fig. 4) points to a central

role of the LS in the variation of aromatase expression in the mouse brain. Interestingly, these

partial correlations reflect main neural connections of the subregions LSD and LSV. For

example, there is input from the amygdala to the LS supporting the LSD-CAM partial

correlation and there is an output from the LS to the MPOA supporting the LSV-MPOA

partial correlation (68–70,73). Thus, at least some of the residual variation in aromatase

expression (Fig. 4) appears to mirror functional connections and relationships of the LS.

The main effects of pup-experience on aromatase expression in the LS were changes

in the left-right balance (Fig. 3B) and significant increase of aromatase-positive cells only in

the right LS of highly experienced (27D) fathers (Fig. 3C). As in the PIR, ERα positive cells

in the LS become visible in caring fathers compared to pup-naïve males (14). Since

hemisphere differences in ERα positive cells have not been found, we cannot directly relate

increased right-hemisphere aromatase expression with non-lateralised increase of ERα

expression in the LS of both hemispheres. Lateralised input to the rostral parts of the LSD and

LSI comes from the hippocampal CA1 and subiculum region (70,73). In rodents, the CA1

shows lateralisation in synapse morphology and receptor distribution (74,75) and left-right

dissociation of memory processes (76) with right-hemisphere dominance of spatial memory in

mice (77). Also, androgen receptor levels are higher in the left hippocampus in male rats (75).

It is unclear, however, whether and how lateralised hippocampal input to the LS could have

caused lateralised aromatase expression in the LS of 27D fathers (Fig. 3C). Right hemisphere

lateralisation is associated with the regulation of mood, affect, and stress, and with sustained

and global arousal and attention (e.g., (78,79)). Speculatively, our observed right-hemisphere

LS bias in 27D fathers could be related to the stress of continuous arousal and involvement in

a routine of pup-care.

Medial preoptic area, bed nucleus stria terminalis, and nucleus accumbens

The MPOA together with the BNST and their output via the NAC are essential for

controlling parental behaviour in rodents (e.g., (23,24,80)). Significant increases have been

observed in the MPOA both of the number of and area occupied by ERα positive cells in one-

day pup-experienced virgin female mice (16) and in 5D and 27D pup-experienced father mice

(14). The number of ERα positive cells increase in the BNST of one-day pup-experienced

virgin females (16) and, although absent in pup-naïve males, ERα-positive cells are present in

the BNST of 27D fathers (14,34). We did not observe a statistically significant change in the

number of aromatase positive cells in the BNST and NAC and found only a small increase in

the MPOA in just the 27D males relative to 5D. This suggests that changes in sensory

processing and behavioural expression with experience in MPOA, BNST, and NAC may be

more dependent on regulation of estrogen signalling through receptors rather than synthesis.

Brain regions that regulate social behaviours, including parental care, are conserved in

vertebrates (81). However, neuroendocrine ligand spatial profiles are more variable than

receptor distributions within the social decision-making network among vertebrates, although

there are regional differences (81). Our work supports experience-dependent regional

differences, and specifically that MPOA, BNST, and NAC may depend on changes in ERα

distribution, whereas VMH and amygdala may depend on changes in the synthesis of the

ligand. Overall, our results point to a region-specific regulation of receptor and ligand with

parental experience.

The brain regions MPOA, BNST, and NAC in N and 5D mice exhibited substantial

lateralization of aromatase, biased toward the left hemisphere (Fig. 3B). There is a left-

hemisphere dominance for pup-ultrasound perception in mothers (82), a left-hemisphere

dominant activation for pup-ultrasound processing in the posterior-dorsal field of the auditory

cortex of mothers and pup-experienced females (83), and a left-hemisphere dominant

expression of oxytocin receptors and higher rate and precision of neural responding in the

primary auditory cortex of mothers and pup-experienced females compared to naïve animals

(84). Pup-retrieving behavior was also accelerated by oxytocin action in the left auditory

cortex (84). That is, ultrasounds of pups motivate caretaking adults to search for the caller

(appetitive behaviour of the parental instinct) in order to be able to perform the consummatory

act of pup-retrieving (17). This sound-dependent motivational activation is left-hemisphere

dominant suggesting that the acoustic input signals to the MPOA, BNST and NAC are also

more relevant when processed via the left hemisphere. Therefore, the greater left hemisphere

aromatase expression in the MPOA, BNST, and for 5D and 27D fathers in the NAC, may

prepare these centers for a preferred processing of pup ultrasounds in order to activate

instinctive parental behaviour. This hypothesis may be tested via unilateral enhancement

and/or suppression of neural processing in these neural centres.

Correlated individual variation in aromatase-positive cells between brain regions

The brain's connectivity suggests that cell counts within individuals may be correlated

across regions. For example, aromatase could be uniformly lower or higher in some

individuals compared to others, or could vary highly non-uniformly across brain regions. We

expect that such variation will contribute to individual differences in behaviour, such as some

fathers continuing to ignore pups (Fig. 1). Our study finds significant residual correlated

variation among brain regions (Fig. 4) that are not generally depicted in the data shown in Fig.

3. It is noteworthy that the posterior PIR and, to a lesser degree, the anterior PIR and the

MAM remained outside this connectivity. Indeed, Supplementary Fig. 2 shows that posterior

PIR (Bregma -1.34) is the sole case in which we observed substantial divergence in overall

cell counts. Together, these observations suggest to us that there are additional sensory and

genetic factors that modulate the numbers of aromatase-positive cells in PIR and MAM,

which may integrate other information as brain pathways reorganise to accommodate paternal

care, such as stress and arousal status. Our work therefore suggests that going forward, it will

be fruitful to consider both the overall regulation of the entire network, along with relative

and absolute differences in specific brain regions.

Conclusion

Aromatase expression in the male limbic brain is regulated with parental experience in

a region-, hemisphere-, and experience-specific manner. Acquiring paternal pup-caring

experience enhanced aromatase expression (number of aromatase positive cells) in brain

centres processing olfactory information (PIR, CAM, MAM) and regulating aggression,

sexual behaviour and other social relations (MAM, VMH, LS). In centers essential for

controlling parental behaviour (MPOA, BNST, NAC), aromatase expression was instead

present with left-hemisphere dominance. In general, the aromatase expression dynamics we

observed did not mirror the previously reported patterns in estrogen receptors (ERα). This

suggests that levels of estrogen receptors are not simply dependent on levels of locally

produced aromatase-dependent estrogen through an autoregulatory feedback loop. Rather,

paternal behaviour may depend on differential regulation of estrogen signalling through

receptor or synthesis in a region-dependent way.

Acknowledgements

We thank H. Sabine Schmidt with assistance with animal care, immunocytochemistry,

and microscopy. This study was supported by the Deutsche Forschungsgemeinschaft,

EH53/19 to GE. PDG was funded by a postdoctoral fellowship from the Fonds Quebecois

Recherche Nature et Technologie. PDG is currently supported by the Canada Research Chair

Program. DAS was partially funded by a postdoctoral fellowship from the Kavli Neuroscience

Discovery Institute.

Table 1. Brain regions analysed for aromatase expression and their Bregma coordinates from

the mouse brain atlas (45).

Brain region		Bregma Coordinates
Lateral septum dorsal (LSD)	Anterior and posterior	0.98 mm and 0.14 mm
Lateral septum intermediate (LSI)	Anterior and posterior	0.98 mm and 0.14 mm
Lateral septum ventral (LSV)	Anterior and posterior	0.98 mm and 0.14 mm
Piriform cortex (PIR)	Anterior and posterior	0.98 mm and -1.34 mm
Nucleus accumbens (NAC)		0.98 mm
Bed nucleus stria terminalis (BNST)		0.14 mm
Medial preoptic area (MPOA)		0.14 mm
Cortical amygdala (CAM)		-1.34 mm
Medial amygdala (MAM)		-1.34 mm
Ventromedial hypothalamus (VMH)		-1.34 mm

Figure 1. Pup-caring behaviour (retrieving) is determined by paternal experience.

A. Experimental setting for the pup-retrieving tests on a running board with a central nest depression. Responses were scored as aggression (attacking pups), ignoring, or retrieving (returning pups to the nest). **B.** Probabilities of behavioural expression in naïve (N) males and in fathers with five (5D) or 27 days of experience (27D). Aggression declined immediately from N to 5D fathers. Although fathers occasionally still ignored pups, the probability of retrieval was very high. Letters indicate statistical differences between the groups of males.

Figure 2. Aromatase expression in the paternal brain. Sagittal view (**A**) and cross sections

(**B**) of the brain regions analysed here (Bregma coordinates: 0.98 mm; 0.14 mm; and -1.34

mm). Positions and extensions of nuclei/areas are illustrative only and do not indicate

anatomical relationships. The connectivity denoted in (A) shown by the arrows are exemplars

for main information pathways (7,85). Representative images from regions highlighted in (B),

showing aromatase immunostaining (arrows) in naïve and experienced fathers (27D) in the

left (L) and right (R) hemisphere. Scale bars in all images indicate 100 µm.

Figure 3. Aromatase expression differentially depends on paternal experience and brain hemisphere. **A.** Standardised cell counts (light grey circles) and posterior means (± 95% confidence interval, white circles) for each investigated nucleus/area separately for the left and right sides of the brain and levels of pup-experience (N, 5D, 27D). **B.** Hemisphere contrast (left – right, with 95% confidence intervals) of the standardised counts of aromatase positive cells for each brain nucleus/area and experimental group. Contrasts with credible intervals that exclude zero are highlighted in purple ($p<0.05$, see text for description). We found evidence for lateralisation of aromatase cell counts both independent of (BNST, MPOA, and most likely NAC) and dependent on experience (most LS and anterior PIR (Bregma 0.98)). **C.** Experience contrast (5D–N; 27D–N, 27D–5D, with 95% confidence intervals) for the standardised counts of aromatase positive cells, separately for each hemisphere and brain nucleus/area. We found aromatase positive cells in posterior PIR (Bregma -1.34), CAM, MAM, and VMH increased with experience level. Expression of aromatase in the LS and anterior PIR increased only in the right hemisphere. No changes were observed in the NAC and BNST. Contrasts with credible intervals that exclude zero are highlighted in purple ($p<0.05$, see text for description). Sample sizes described in text.

Figure 4. Residual correlations among brain nuclei/areas viewed as a network. Each nucleus/area at each Bregma coordinate (see Table 1) is represented by a node, with edges representing significant partial correlations (solid for $p< 0.05$, dashed for $p< 0.10$).

Figure 5. Effects of paternal experience on aromatase expression in areas of the limbic system. Summary of the changes in aromatase expression in 5D (**A**) and 27D (**B**) males in the left and right hemispheres of the brain nuclei/areas analysed in the current study. **A.** Pup experience in first-litter fathers (5D group) increased the numbers of aromatase positive cells (indicated by light purple) relative to naïve males in the amygdala (CAM and MAM), hypothalamus (VMH), and in posterior piriform cortex (PIR, half shading indicates these partial changes). **B.** Second-litter fathers (27D group) had further increased aromatase expression in the same areas as 5D males (indicated as darker purple), i.e., the amygdala (CAM and MAM), hypothalamus (VMH), and posterior piriform cortex (PIR) in both sides of the brain. In the right hemisphere, the number of aromatase positive cells increased (indicated by blue) in the anterior PIR, lateral septum (LS) relative to naïve and 5D males, and potentially in the MPOA relative to 5D males only.

References

1. Bales KL, Saltzman W. Fathering in rodents: Neurobiological substrates and consequences for offspring. Hormones and Behavior. 2016 Jan;77:249–59.

2. Brunton PJ, Russell JA. The expectant brain: adapting for motherhood. Nat Rev Neurosci. 2008 Jan;9(1):11–25.

3. Dulac C, O'Connell LA, Wu Z. Neural control of maternal and paternal behaviors. Science. 2014 Aug 15;345(6198):765–70.

4. Guoynes C, Marler C. Paternal Behavior from a Neuroendocrine Perspective. In: Oxford Research Encyclopedia of Neuroscience [Internet]. 2020 [cited 2023 May 4]. Available from: https://oxfordre.com/neuroscience/display/10.1093/acrefore/9780190264086.001.0001/acrefore-9780190264086-e-9

5. Horrell ND, Acosta MC, Saltzman W. Plasticity of the paternal brain: Effects of fatherhood on neural structure and function. Dev Psychobiol. 2021 Jan 22;dev.22097.

6. Lonstein JS, De Vries GJ. Sex differences in the parental behavior of rodents. Neurosci Biobehav Rev. 2000 Aug;24(6):669–86.

7. Numan M. The Parental Brain: Mechanisms, Development, and Evolution. Oxford University Press; 2020.

8. Numan M, Insel TR. The neurobiology of parental behavior. New York: Springer; 2003. 418 p. (Hormones, brain, and behavior).

9. Rilling JK, Mascaro JS. The neurobiology of fatherhood. Curr Opin Psychol. 2017 Jun;15:26–32.

10. Woodroffe R, Vincent A. Mother's little helpers: Patterns of male care in mammals. Trends in Ecology & Evolution. 1994 Aug 1;9(8):294–7.

11. Fahrbach SE, Pfaff DW. Hormonal and Neural Mechanisms Underlying Maternal Behavior in the Rat. In: Pfaff DW, editor. The Physiological Mechanisms of Motivation [Internet]. New York, NY: Springer; 1982 [cited 2023 May 4]. p. 253–85. Available from: https://doi.org/10.1007/978-1-4612-5692-2_9

12. Rosenblatt JS, Siegel HI. Factors Governing the Onset and Maintenance of Maternal Behavior among Nonprimate Mammals. In: Gubernick DJ, Klopfer PH, editors. Parental Care in Mammals [Internet]. Boston, MA: Springer US; 1981 [cited 2023 May 4]. p. 13–76. Available from: https://doi.org/10.1007/978-1-4613-3150-6_2

13. Akther S, Huang Z, Liang M, Zhong J, Fakhrul AAKM, Yuhi T, et al. Paternal Retrieval Behavior Regulated by Brain Estrogen Synthetase (Aromatase) in Mouse Sires that Engage in Communicative Interactions with Pairmates. Front Neurosci [Internet]. 2015 Dec 15 [cited 2021 May 24];9. Available from: http://journal.frontiersin.org/Article/10.3389/fnins.2015.00450/abstract

14. Ehret G, Jürgens A, Koch M. Oestrogen receptor occurrence in the male mouse brain: modulation by paternal experience. Neuroreport. 1993 Sep 10;4(11):1247–50.

552 15. Ehret G, Koch M, Haack B, Markl H. Sex and parental experience determine the onset of
553   an instinctive behavior in mice. Naturwissenschaften. 1987 Jan;74(1):47–47.

554 16. Ehret G, Buckenmaier J. Estrogen-receptor occurrence in the female mouse brain: Effects
555   of maternal experience, ovariectomy, estrogen and anosmia. Journal of Physiology-Paris.
556   1994 Jan 1;88(5):315–29.

557 17. Ehret G, Koch M. Ultrasound-induced Parental Behaviour in House Mice is Controlled by
558   Female Sex Hormones and Parental Experience. Ethology. 1989;80(1–4):81–93.

559 18. Elwood RW. Inhibition of infanticide and onset of paternal care in male mice (Mus
560   musculus). Journal of Comparative Psychology. 1985;99:457–67.

561 19. Koch M, Ehret G. Estradiol and parental experience, but not prolactin are necessary for
562   ultrasound recognition and pup-retrieving in the mouse. Physiol Behav. 1989
563   Apr;45(4):771–6.

564 20. Stolzenberg DS, Rissman EF. Oestrogen-independent, experience-induced maternal
565   behaviour in female mice. J Neuroendocrinol. 2011 Apr;23(4):345–54.

566 21. Tachikawa KS, Yoshihara Y, Kuroda KO. Behavioral Transition from Attack to Parenting
567   in Male Mice: A Crucial Role of the Vomeronasal System. Journal of Neuroscience. 2013
568   Mar 20;33(12):5120–6.

569 22. Koch M, Ehret G. Parental behavior in the mouse: effects of lesions in the
570   entorhinal/piriform cortex. Behavioural Brain Research. 1991 Jan;42(1):99–105.

571 23. Numan M. Hypothalamic neural circuits regulating maternal responsiveness toward
572   infants. Behav Cogn Neurosci Rev. 2006 Dec;5(4):163–90.

573 24. Bridges RS. Neuroendocrine regulation of maternal behavior. Front Neuroendocrinol.
574   2015 Jan;36:178–96.

575 25. Ehret G, Koch M, Jirikowski GF. Hormone, ZNS und Fortpflanzungsverhalten.
576   Allergologie. 1991;14:383–94.

577 26. Kohl J, Dulac C. Neural control of parental behaviors. Curr Opin Neurobiol. 2018
578   Apr;49:116–22.

579 27. Poindron P, Lévy F, Krehbiel D. Genital, olfactory, and endocrine interactions in the
580   development of maternal behaviour in the parturient ewe. Psychoneuroendocrinology.
581   1988;13(1–2):99–125.

582 28. Romero-Morales L, Martínez-Torres M, Cárdenas M, Álvarez C, Carmona A, Cedillo B,
583   et al. An increase in estradiol facilitates the onset of paternal behavior in the dwarf
584   hamster (Phodopus campbelli). Hormones and Behavior. 2018 Mar;99:35–40.

585 29. Rosenblatt JS, Wagner CK, Morrell JI. Hormonal priming and triggering of maternal
586   behavior in the rat with special reference to the relations between estrogen receptor
587   binding and ER mRNA in specific brain regions. Psychoneuroendocrinology. 1994;19(5–
588   7):543–52.

30. Rosenblatt JS, Ceus K. Estrogen Implants in the Medial Preoptic Area Stimulate Maternal
Behavior in Male Rats. Hormones and Behavior. 1998 Feb 1;33(1):23–30.

31. Siegel HI, Rosenblatt JS. Estrogen-induced maternal behavior in hysterectomized-
ovariectomized virgin rats. Physiology & Behavior. 1975 Apr 1;14(4):465–71.

32. Koch M. Effects of treatment with estradiol and parental experience on the number and
distribution of estrogen-binding neurons in the ovariectomized mouse brain.
Neuroendocrinology. 1990 May;51(5):505–14.

33. Li Y, Lian Z, Wang B, Tai F, Wu R, Hao P, et al. Natural variation in paternal behavior is
associated with central estrogen receptor alpha and oxytocin levels. J Comp Physiol A
Neuroethol Sens Neural Behav Physiol. 2015 Mar;201(3):285–93.

34. Koch M, Ehret G. Immunocytochemical localization and quantitation of estrogen-binding
cells in the male and female (virgin, pregnant, lactating) mouse brain. Brain Res. 1989
Jun 5;489(1):101–12.

35. Nilsson ME, Vandenput L, Tivesten Å, Norlén AK, Lagerquist MK, Windahl SH, et al.
Measurement of a Comprehensive Sex Steroid Profile in Rodent Serum by High-Sensitive
Gas Chromatography-Tandem Mass Spectrometry. Endocrinology. 2015 Jul
1;156(7):2492–502.

36. Cornil CA, Seredynski AL, de Bournonville C, Dickens MJ, Charlier TD, Ball GF, et al.
Rapid control of reproductive behaviour by locally synthesised oestrogens: focus on
aromatase. J Neuroendocrinol. 2013 Nov;25(11):1070–8.

37. Cornil CA, Ball GF, Balthazart J. Functional significance of the rapid regulation of brain
estrogens: Where do the estrogens come from? Brain Res. 2006 Dec 18;1126(1):2–26.

38. Garcia-Segura LM. Aromatase in the Brain: Not Just for Reproduction Anymore. Journal
of Neuroendocrinology. 2008;20(6):705–12.

39. Trainor BC, Bird IM, Alday NA, Schlinger BA, Marler CA. Variation in Aromatase
Activity in the Medial Preoptic Area and Plasma Progesterone Is Associated with the
Onset of Paternal Behavior. Neuroendocrinology. 2003;78(1):36–44.

40. Trainor BC, Marler CA. Testosterone promotes paternal behaviour in a monogamous
mammal via conversion to oestrogen. Proc R Soc Lond B. 2002 Apr 22;269(1493):823–9.

41. Foidart A, Harada N, Balthazart J. Aromatase-immunoreactive cells are present in mouse
brain areas that are known to express high levels of aromatase activity. Cell Tissue Res.
1995 Jun;280(3):561–74.

42. Schleicher G, Stumpf WE, Morin JK, Drews U. Sites of aromatization of [3H]testosterone
in forebrain of male, female and androgen receptor-deficient Tfm mice: an
autoradiographic study. Brain Res. 1986 Nov 12;397(2):290–6.

43. Stanić D, Dubois S, Chua HK, Tonge B, Rinehart N, Horne MK, et al. Characterization of
aromatase expression in the adult male and female mouse brain. I. Coexistence with
oestrogen receptors α and β, and androgen receptors. PLoS One. 2014;9(3):e90451.

44. Charitidi K, Canlon B. Estrogen receptors in the central auditory system of male and
female mice. Neuroscience. 2010 Feb 3;165(3):923–33.

45. Paxinos G, Franklin K. The Mouse Brain in Stereotaxic Coordinates. 2nd ed. San Diego:
Academic Press; 2001.

46. R Core Team. R: A language and environment for statistical computing. R Foundation for
Statistical Computing, Vienna, Austria. URL https://www.R-project.org/. 2022;

47. van Buuren S, Groothuis-Oudshoorn K. mice: Multivariate Imputation by Chained
Equations in R. Journal of Statistical Software. 2011 Dec 12;45:1–67.

48. Bürkner PC. brms: An R Package for Bayesian Multilevel Models Using Stan. Journal of
Statistical Software. 2017 Aug 29;80:1–28.

49. Vehtari A, Gelman A, Simpson D, Carpenter B, Bürkner PC. Rank-Normalization,
Folding, and Localization: An Improved R̂ for Assessing Convergence of MCMC (with
Discussion). Bayesian Analysis. 2021 Jun;16(2):667–718.

50. Lenth R. _emmeans: Estimated Marginal Means, aka Least-Squares Means_. R package
version 1.8.0, <https://CRAN.R-project.org/package=emmeans>. 2022;

51. Makowski D, Ben-Shachar MS, Lüdecke D. bayestestR: Describing Effects and their
Uncertainty, Existence and Significance within the Bayesian Framework. Journal of Open
Source Software. 2019 Aug 13;4(40):1541.

52. Wickham H. ggplot2: Elegant Graphics for Data Analysis. New York: Springer-Verlag;
2016.

53. de la Fuente A, Bing N, Hoeschele I, Mendes P. Discovery of meaningful associations in
genomic data using partial correlation coefficients. Bioinformatics. 2004 Dec
12;20(18):3565–74.

54. Shipley B. Cause and Correlation in Biology: A User's Guide to Path Analysis, Structural
Equations, and Causal Inference in R. Cambridge, UK: Cambridge University Press;
2000. 317 p.

55. Kalisch M, Mächler M, Colombo D, Maathuis MH, Bühlmann P. Causal Inference Using
Graphical Models with the R Package pcalg. Journal of Statistical Software. 2012 May
17;47:1–26.

56. Chen PB, Hu RK, Wu YE, Pan L, Huang S, Micevych PE, et al. Sexually Dimorphic
Control of Parenting Behavior by the Medial Amygdala. Cell. 2019 Feb 21;176(5):1206-
1221.e18.

57. Dwyer J, Kelly DA, Bergan J. Brain-Wide Synaptic Inputs to Aromatase-Expressing
Neurons in the Medial Amygdala Suggest Complex Circuitry for Modulating Social
Behavior. eNeuro [Internet]. 2022 Mar 1 [cited 2023 May 4];9(2). Available from:
https://www.eneuro.org/content/9/2/ENEURO.0329-21.2021

58. Hashikawa K, Hashikawa Y, Falkner A, Lin D. The neural circuits of mating and fighting
in male mice. Curr Opin Neurobiol. 2016 Jun;38:27–37.

59. Lee H, Kim DW, Remedios R, Anthony TE, Chang A, Madisen L, et al. Scalable Control
of Mounting and Attack by ESR1+ Neurons in the Ventromedial Hypothalamus. Nature.
2014 May 29;509(7502):627–32.

60. Sheehan T, Paul M, Amaral E, Numan MJ, Numan M. Evidence that the medial amygdala
projects to the anterior/ventromedial hypothalamic nuclei to inhibit maternal behavior in
rats. Neuroscience. 2001;106(2):341–56.

61. Unger EK, Burke KJ, Yang CF, Bender KJ, Fuller PM, Shah NM. Medial Amygdalar
Aromatase Neurons Regulate Aggression in Both Sexes. Cell Reports. 2015 Feb
3;10(4):453–62.

62. Yang T, Yang CF, Chizari MD, Maheswaranathan N, Burke KJ, Borius M, et al. Social
Control of Hypothalamus-Mediated Male Aggression. Neuron. 2017 Aug 16;95(4):955-
970.e4.

63. Trainor BC, Finy MS, Nelson RJ. Paternal aggression in a biparental mouse: parallels
with maternal aggression. Horm Behav. 2008 Jan;53(1):200–7.

64. Getz LL, Carter CS, Gavish L. The mating system of the prairie vole, Microtus
ochrogaster: Field and laboratory evidence for pair-bonding. Behav Ecol Sociobiol. 1981
Jun 1;8(3):189–94.

65. Devidze N, Mong JA, Jasnow AM, Kow LM, Pfaff DW. Sex and estrogenic effects on
coexpression of mRNAs in single ventromedial hypothalamic neurons. Proceedings of the
National Academy of Sciences. 2005 Oct 4;102(40):14446–51.

66. Malikov V, Madeira MD. Regulation of ERα protein expression by 17β-estradiol in
cultured neurons of hypothalamic ventromedial nucleus. Neurochem Res. 2013
Jan;38(1):82–9.

67. Honda SI, Wakatsuki T, Harada N. Behavioral Analysis of Genetically Modified Mice
Indicates Essential Roles of Neurosteroidal Estrogen. Front Endocrinol (Lausanne). 2011
Sep 26;2:40.

68. Besnard A, Leroy F. Top-down regulation of motivated behaviors via lateral septum sub-
circuits. Mol Psychiatry. 2022 Aug;27(8):3119–28.

69. Menon R, Süß T, Oliveira VE de M, Neumann ID, Bludau A. Neurobiology of the lateral
septum: regulation of social behavior. Trends in Neurosciences. 2022 Jan 1;45(1):27–40.

70. Rizzi-Wise CA, Wang DV. Putting Together Pieces of the Lateral Septum: Multifaceted
Functions and Its Neural Pathways. eNeuro. 2021 Dec 2;8(6):ENEURO.0315-21.2021.

71. Fleischer S, Slotnick BM. Disruption of maternal behavior in rats with lesions of the
septal area. Physiology & Behavior. 1978 Aug 1;21(2):189–200.

72. Slotnick BM, Nigrosh BJ. Maternal behavior of mice with cingulate cortical, amygdala,
or septal lesions. Journal of Comparative and Physiological Psychology. 1975;88(1):118.

73. Risold PY, Swanson LW. Connections of the rat lateral septal complex. Brain Res Brain
Res Rev. 1997 Sep 19;24(2–3):115–95.

74. Shinohara Y, Hirase H, Watanabe M, Itakura M, Takahashi M, Shigemoto R. Left-right
asymmetry of the hippocampal synapses with differential subunit allocation of glutamate
receptors. Proceedings of the National Academy of Sciences. 2008 Dec 9;105(49):19498–
503.

75. Xiao L, Jordan CL. Sex Differences, Laterality, and Hormonal Regulation of Androgen
Receptor Immunoreactivity in Rat Hippocampus. Hormones and Behavior. 2002
Nov;42(3):327–36.

76. Shipton OA, El-Gaby M, Apergis-Schoute J, Deisseroth K, Bannerman DM, Paulsen O,
et al. Left–right dissociation of hippocampal memory processes in mice. Proc Natl Acad
Sci USA. 2014 Oct 21;111(42):15238–43.

77. Shinohara Y, Hosoya A, Yamasaki N, Ahmed H, Hattori S, Eguchi M, et al. Right-
hemispheric dominance of spatial memory in split-brain mice. Hippocampus.
2012;22(2):117–21.

78. Ehret G. Hemisphere Dominance of Brain Function--Which Functions Are Lateralized
and Why? In: 23 problems in systems neuroscience. New York, NY, US: Oxford
University Press; 2006. p. 44–61. (Computational neuroscience).

79. Palomero-Gallagher N, Amunts K. A short review on emotion processing: a lateralized
network of neuronal networks. Brain Struct Funct. 2022;227(2):673–84.

80. Wu Z, Autry AE, Bergan JF, Watabe-Uchida M, Dulac CG. Galanin neurons in the
medial preoptic area govern parental behaviour. Nature. 2014 May;509(7500):325–30.

81. O'Connell LA, Hofmann HA. Evolution of a Vertebrate Social Decision-Making
Network. Science. 2012 Jun;336(6085):1154–7.

82. Ehret G. Left hemisphere advantage in the mouse brain for recognizing ultrasonic
communication calls. Nature. 1987 Jan;325(6101):249–51.

83. Geissler DB, Schmidt HS, Ehret G. Knowledge About Sounds—Context-Specific
Meaning Differently Activates Cortical Hemispheres, Auditory Cortical Fields, and
Layers in House Mice. Frontiers in Neuroscience [Internet]. 2016 [cited 2023 May 4];10.
Available from: https://www.frontiersin.org/articles/10.3389/fnins.2016.00098

84. Marlin BJ, Mitre M, D'amour JA, Chao MV, Froemke RC. Oxytocin enables maternal
behaviour by balancing cortical inhibition. Nature. 2015 Apr 23;520(7548):499–504.

85. Kohl J, Autry AE, Dulac C. The neurobiology of parenting: A neural circuit perspective.
Bioessays. 2017 Jan;39(1):1–11.