Impact of High Fat Diet and Sex in a Rabbit Model of Carpal Tunnel Syndrome

Tomoyuki Kuroiwa[1], Hayman Lui[1], Koichi Nakagawa[1], Naoya Iida[1], Claire Desrochers[1], Rou Wan[1], Elameen Adam[1], Dirk Larson[2], Peter Amadio[1], Anne Gingery[1,3]

1. Department of Orthopedic Surgery, Mayo Clinic, Rochester, Minnesota, United States of America.
2. Department of Quantitative Health Sciences, Mayo Clinic, Rochester, Minnesota, United States of America.
3. Department of Biochemistry and Molecular Biology, Mayo Clinic, Rochester, Minnesota, United States of America

Corresponding authors

Peter C Amadio, MD

Department of Orthopedic Surgery, Mayo Clinic

200 First St. SW, Rochester, Minnesota, United States of America

Tel.: +1-507-201-2220

E-mail: pamadio@mayo.edu

Anne Gingery, PhD

Department of Orthopedic Surgery, Mayo Clinic

200 First St. SW, Rochester, Minnesota, United States of America

Tel.: +1-507-293-2105

E-mail: gingery.anne@mayo.edu

Abstract

Carpal tunnel syndrome (CTS) is a common musculoskeletal disorder, characterized by fibrosis of the subsynovial connective tissue (SSCT) mediated by transforming growth factor beta (TGF-β). Risk factors for CTS include metabolic dysfunction and age. Additionally, the incidence of CTS is higher in women. In this study we hypothesized that a high-fat diet (HFD), a common driver of metabolic dysfunction, would promote SSCT fibrosis found in CTS and that this response would be sex dependent. To test this, we examined the effects of HFD and sex on SSCT fibrosis using our established rabbit model of CTS. Forty-eight (24 male, 24 female) adult rabbits were divided into four groups including HFD or standard diet with and without CTS induction. SSCT was collected for histological and gene expression analysis. HFD promoted SSCT thickening and upregulated profibrotic genes, including TGF-β. Fibrotic genes were differentially expressed in males and females. Interestingly while the prevalence of CTS is greater in women than in men, the converse is observed in the presence of metabolic dysfunction. This work recapitulates this clinical observation and begins to elucidate the sex-based differences found in SSCT fibrosis. This knowledge should drive further research and may lead to metabolic and sex specific therapeutic strategies for the treatment of patients with CTS.

Key words

Carpal tunnel syndrome, metabolic dysfunction, metabolic syndrome, sex differences, high-fat diet, rabbit model, fibrosis, subsynovial connective tissue, transforming growth factor beta

Introduction

Carpal tunnel syndrome (CTS), is the most common compression neuropathy with a prevalence of nearly 5% in adults [1]. Despite this prevalence, in most cases the underlying cause of CTS is unknown. Surgery is the only effective curative treatment [2]. A better understanding of the etiology of CTS is important, as this may lead to new nonsurgical treatments, targeting the underlying mechanism of disease.

In addition to deformation of the median nerve, the main pathological feature of CTS is fibrosis of the subsynovial connective tissue (SSCT), which is driven by transforming growth factor beta (TGF-β)-mediated signaling [3]. A significant risk factor for CTS is metabolic syndrome (MetS) [4], which has been proposed to induce systemic inflammation that promotes fibrosis in various tissues and organs [5]. Multiple studies have also reported that both MetS and high fat diet (HFD) promote TGF-β-mediated fibrosis [6-13], as is found in CTS. Therefore, it is reasonable to explore the role of metabolic dysfunction in CTS.

Another consideration when evaluating CTS etiology is the sex difference in the prevalence of CTS, which occurs twice as frequently in women as in men [14]. Interestingly, men with metabolic dysfunction are more likely to have CTS than are women with metabolic dysfunction [15,16]. Based on this information, we hypothesized that there is a sex difference in the relationship between CTS and metabolic dysfunction. We also hypothesized that HFD [17-19] drives metabolic dysfunction and induces SSCT fibrosis in the carpal tunnel in a sex dependent manner. To test this hypothesis, we conducted a comparative study to evaluate SSCT fibrosis in mature male and female rabbits who were fed a HFD or normal diet, with or without CTS induction [20-22]

Material and Methods

Animals

This study was approved by Mayo Clinic Institutional Animal Care and Use Committee (IACUC A5660), and all procedures were performed in accordance with the National Institutes of Health Guide for the Care and Use of Laboratory Animals. 48 retired breeder New Zealand White rabbits (24 females, 24 males) were used to represent middle-aged animals, as CTS is an age-related disease with its peak incidence in humans aged 45-55. For this reason, we used mature rabbits with an average age of 2.6 (0.9 - 4.3) years.

Rabbits were randomly divided into four groups of 12 (6 females and 6 males in each group). The groups were fed a standard diet (STD) or a high-fat diet (HFD), and received a CTS induction dextrose injections (CTS) or saline injections (Sham) as previously described [23] Groups were designated as follows: STD+Sham, STD+CTS, HFD+Sham, and HFD+CTS (Figure 1). Rabbits in the STD groups were acclimated for a week on STD and then received the CTS or Sham injections as described below. STD rabbits continued the STD for an additional 6 weeks prior to sacrifice. HFD group rabbits were acclimated for a week and fed HFD for 12 weeks prior to injections to develop metabolic dysfunction, based on previous studies of HFD in rabbits [24-26]. The HFD rabbits then received CTS or Sham injections and continued the HFD for an additional 6 weeks prior to sacrifice. (Figure 1). In all groups, immediately after sacrifice, one forepaw was harvested for histology and the other for SSCT gene expression analysis. Metabolic dysfunction was assessed by measuring, hyperglycemia, hypercholesterolemia, and hypertriglyceridemia dysfunction as described below [27]. One STD-fed female and one HFD-fed

female rabbit died during the study. The former died from causes unrelated to the study design, whereas the latter died from liver failure, associated with the HFD.

Diet

The diet for the STD group was a standard rabbit diet (Laboratory Rabbit Diet HF 5326, CA). This diet included 12% water, 14.5% protein, 23% fiber, and 3.8% fat. The HFD (Teklad Custom Diet TD.200785, ENVIGO, IN) was the STD supplemented with 0.5% cholesterol and 4% peanut oil, as previously reported [20,21,25,26]. Water and food were not restricted during this study.

CTS induction

Using our established rabbit model of CTS, a small incision was made with direct vision while under anesthesia. For CTS and sham, the carpal tunnels of both forepaws were injected with either 0.1 ml of 10% dextrose solution (CTS) or 0.1 ml of 0.9% saline (Irrigation, Baxter, IL) (Sham), using a 30-gauge needle. As per the protocol two injections were given 1 week apart. Each injection was made inside the synovium around the middle digit flexor digitorum superficialis tendon, as described previously [22,28].

Metabolic Dysfunction Assessment

Blood samples were taken at sacrifice. Rabbits were fasted overnight and blood was taken from the marginal ear vein. Blood samples were centrifuged at 3,000 rpm for 20 minutes at 4 °C, and the supernatant was collected as serum. Samples were sent to IDEXX BioAnalytics (Columbia, MO) for analysis of total cholesterol, triglyceride, glucose, and insulin levels in the serum (IDEXX BioAnalytics Case # 82497-2021).

Histology

The tissue was harvested so that an axial section at the wrist level was visible after bone removal. The SSCT was fixed in a 4% paraformaldehyde solution for approximately one week, embedded in paraffin, and sectioned into 4-μm-thick slices. Sections were stained with hematoxylin-eosin and observed under a light microscope (All-in-One Fluorescence Microscope BZ-X800, Keyence, IL, RRID:SCR_019119).

Image analysis

Tissue images were evaluated using ImageJ software (ImageJ, National Institutes of Health; http://imagej.nih.gov/ij/, RRID: SCR_003070) by two blinded observers. The distance between the flexor digitorum superficialis (FDS) and flexor digitorum profundus (FDP) was measured and determined to be the SSCT thickness, as described in previous studies [22,29] (Figure 2). The average values of the thickness measured by two observers were used for analysis.

Quantitative real-time polymerase chain reaction (PCR)

After sacrifice, total RNA was extracted from the SSCT of one forepaw using Trizol reagent (Invitrogen Life Technologies, NY). The cDNA was obtained from equal amounts of total RNA by reverse transcription using an iScriptTM cDNA Synthesis Kit (Bio-Rad, CA). Gene expression of 96 fibrosis-associated genes were evaluated using a PCR array kit (RT² Profiler™ PCR Array Rabbit Fibrosis PANZ-120Z, Qiagen) by quantitative real-time PCR using a Thermal Cycler (Bio-Rad Laboratories, CA, RRID:SCR_019152). The average values of the results in triplicate were used for analysis. Eighty-four fibrosis-related genes were evaluated. Data was normalized to the geo-mean of housekeeping genes Acta2, Actb, Gapdh, Ldha, and LOC100346936.

Statistics

Sample size and power

Based on previous studies using this model, a sample size of 6 rabbits per group is required to achieve 80% power to detect a difference in means of 0.15 mm between groups in the FDS-FDP distance, assuming an overall within group standard deviation of 0.08 mm[22]. Additionally, in order to perform separate analyses within each sex, each group was fully powered with 6 males and 6 females, for a total of 12 rabbits per group (48 rabbits total).

Analysis

Separate analyses were performed for males and females. Within each sex subset, the effects of CTS induction (no vs yes) and diet (HFD vs STD) on SSCT thickness and gene expression level was evaluated using two-way analysis of variance (ANOVA). For blood test results, mean values were calculated for each diet group and sex and compared using two-way ANOVA. P-values less than 0.05 were considered statistically significant.

Results

Metabolic dysfunction

Blood analyses were completed at sacrifice. The results showed no significant interactions between diet and sex (Figure 2). HFD female and male rabbits had significantly higher cholesterol, triglycerides, and insulin levels and significant lower glucose levels as compared to STD rabbits (Figure 2).

SSCT histology

SSCT thickness was measured as a marker of fibrosis (Figure 3). A representative image of hematoxylin-eosin SSCT measurement is shown in Figure 3a. The overall average SSCT

thickness was 0.16 mm, 0.31 mm, 0.33 mm, and 0.33 mm in the STD+Sham, the STD+CTS, the HFD+Sham, and the HFD+CTS groups, respectively. The analysis showed that there was a significant interaction between diet and CTS induction and that both main effects of induction and diet were significant in affecting SSCT thickness (Figure 3b). In male rabbits, both diet and CTS induction resulted in significantly thicker SSCT (Figure 3c). In contrast, CTS induction alone in female rabbits was found to be the main driver of SSCT fibrosis. (Figure 3d).

SSCT Gene expression

Scatter plots of gene expression changes due to the effects of CTS induction or diet are shown in Figure 4a-d. Many fibrosis-promoting genes, including Itga1, Itgb1, Tgfbr1, Ilk, Stat6, and Vegfa were upregulated in the combination of CTS induction and HFD. Significant main effects of induction and diet were found in 45 and 17 of the 84 fibrotic genes assessed, respectively. In particular, many fibrosis-promoting genes, including Il1a, Itgav, Tgfb3, Tgfbr1, Timp1, and Smad7 were upregulated by CTS induction alone. Furthermore, it was observed that HFD fibrosis-promoting genes such as Ctgf and Cxcr4 were upregulated, and genes such as Dcn and Ifng were down-regulated in rabbits treated with HFD. (Figure 4e).

In gender subanalysis, it was observed that the combination of male sex and a HFD resulted in 8 significantly upregulated genes, including Cul1, Cxcr4, Eng, Fli1, Prkcd, Stat6, Tgfbr1, and Timp4. In contrast, female interaction between sex and diet demonstrated only one upregulated gene (Il1b). No down-regulated genes were observed in HFD in both sexes. In males, significant main effects of CTS induction and diet were found in 28 and 3 significantly regulated genes, respectively. In females, significant main effects of CTS induction and diet were found in 4 and

13 genes, respectively. Of the 46 genes that demonstrated significant main effects, only two genes had the same main effect in both males and females (Itgav and Tgfb3)(Figure 5).

Discussion

In this study both CTS induction and HFD significantly increased SSCT thickness. All rabbits exhibited induced hypercholesterolemia with HFD. Both CTS induction and HFD upregulated many fibrosis-inducing genes such as Tgfbr1, the receptor protein of the TGF-β family, as well as fibrosis-associated integrin-related genes (Itga1, Itgb1, Ilk) [3,30-32], Stat 6 [33], and Vegfa[34,35].

In this study, we used a HFD rabbit model as previously described [21,24-26]. Similar to Morelli et. Al [21]., the rabbits used in this study developed hypercholesterolemia and hypertriglyceridemia. However, unlike previous reports, we did not find a significant increase in blood glucose levels. This may have been due to age differences as the present study used middle-aged rabbits compared to young rabbits in previous studies. As CTS and metabolic dysfunction is more prevalent in older patients, we selected older animals to more accurately represent the clinical condition studied. These results are consistent with trends observed in other HFD-fed animal studies using both young and aged animals who have previously observed n significant elevation in blood glucose levels in young animals only [36,37].

It is well established that metabolic dysfunction promotes fibrosis in many tissues/organs [5,38]. Clinically, metabolic dysfunction has been reported as one of the risk factors for CTS[4,39-41], and the results reported here provide support for this risk factor driving SSCT fibrosis. Interestingly, subanalysis of gender showed that the HFD induced SSCT thickening in male rabbits only. Previous studies have reported that mature male rats and mice were also more vulnerable to the effects of the HFD than females [42,43]. In addition clinical studies have reported that metabolic

dysfunction is a risk factor for CTS [4,39-41], and that the incidence is higher in men[15,16]. This is consistent with our finding that both gender and HFD significantly contributes to inducing SSCT thickening and fibrosis.

Ctgf is a known mediator of TGF-β induced fibrosis [44]. In this study, significantly increased expression of Ctgf was observed in the HFD rabbits. This same trend has been observed in previous HFD animal studies[45-47] and in analysis of SSCT tissue of patients with CTS[3,48]. Moreover, Tgfbr1 and integrin-related genes were observed to be significantly upregulated in both CTS induction and HFD fed animals. Tgfbr1 encodes a receptor that binds to TGF-β protein, promotes TGF-β signaling, and has also been found to be upregulated with metabolic syndrome [11,49,50]. A subset of the integrin family has been reported to play an important role in the activation of latent TGFB1, thereby promoting the fibrosis process in models of various organ fibrosis [31,32,51,52]. The significant upregulation of these genes is consistent with the results of the SSCT thickness assessment.

In contrast, the expression of anti-fibrotic genes such asIfng[53,54] and Dcn[55-57] were noted to be significantly downregulated in HFD rabbits. Interferon-γ, encoded by Ifng, is an immune-regulatory cytokine that also serves as a strong anti-fibrotic agent[54]. IFN-γ also inhibits fibroblast proliferation and collagen deposition and downregulates Tgfb expression[53]. The combination of both upregulation of fibrotic genes and downregulation of anti-fibrotic genes in HFD rabbits suggests that metabolic dysfunction induces gene expression changes that may drive SSCT fibrosis.

Gene expression levels also differed between males and females. In male rabbits, CTS induction significantly upregulated Cav1 and Smad7. Cav1 is overexpressed in adipose tissue of obese

patients and is positively correlated with body fat and body mass index[58]. Cav1 has also been reported to suppress fibrosis in various organs via suppression of TGF-β/Smad signaling [59,60]. Smad7, a Smad inhibitory protein, is a key regulator that binds to regulatory SMADs and acts in to inhibit TGF-β signaling. Smad7 overexpression has been shown to antagonize TGF-β-mediated fibrosis and inflammation [61,62]. Therefore, the observed elevation in Cav1 and Smad7 expression in male rabbits may be a compensatory response to ongoing fibrosis in the SSCT.

Furthermore, Dcn, another negative regulator of TGF-β signaling, was significantly down regulated in HFD males only. Decorin, a small leucine-rich proteoglycan encoded by Dcn, regulates collagen fibrillogenesis during development. It exerts a protective effect against fibrosis in various tissues by directly blocking the bioactivity of Tgfb1[55-57]. The loss of this important TGF-β regulator in conditions of metabolic dysfunction may contribute to increased SSCT fibrosis in males.

Another gene that was regulated by diet in males was plasminogen activator encoded by Plat. Plat is a component of the plasminogen system that is important for proteolytic remodeling of extracellular matrix. Plasminogen activators can act to attenuate the action of plasminogen activator inhibitors (PAI-1) that can drive fibrosis and extracellular matrix deposition [63]. Expression of Plat could be a compensatory mechanism to attenuate fibrosis. Plat also has been shown to limit activation of Tgfb1[64,65].

In contrast, HFD significantly upregulated Mmp2 and downregulated Smad ubiquitin regulatory factor 2 (Smurf2) in females only.. MMP2 has been shown to have anti-fibrotic effects[66,67] and Smurf2 negatively regulates TGF-β signaling by ubiquitinating TGF-β receptors (TβR) and Smad proteins. Together, these significant disparities in gene expression may provide a

mechanistic explanation for the disparate response to CTS induction and diet in males and females.

There are some limitations of this study that are important to note. In this study, female rabbits were housed 2 to a cage, whereas male rabbits were housed 1 to a cage, as required by our IACUC. This difference in housing may have caused differences in exercise and other behaviors. HFD rabbits were maintained on HFD for 12 weeks, in contrast the CTS or sham induced STD rabbits which were treated after the standard 1 week of acclimation for all animals. As noted, all rabbits were housed in a standard manner, except for the special diet fed to the HFD group. We believe that the effect of the difference in length of housing in our facility is likely to be negligible, as the duration of housing in our facility for the treatment phase was identical in all four groups. Our future work includes assessment of sex differences in the gene and protein regulation of Cav1, Dcn, Mmp2, Plat, Smad7, Smurf2 in human SSCT biopsies and future mechanistic studies.

In summary, the effect of HFD and its interaction with sex on SSCT fibrosis in a CTS animal model was studied. HFD resulted in increased SSCT thickness compared to STD rabbits and upregulation of fibrosis related genes (Tgfbr1, Itga1, Itgb1, Ilk, and Ctgf) which are key activators of TGF-β signaling. This is consistent with previous analysis of the SSCT in CTS patients. In addition, the combination of HFD and CTS induction in males resulted in significantly greater SSCT fibrosis when compared to females. This is supportive of previous reports where men with metabolic dysfunction are more likely to develop CTS [15,16]For the first time, this study has shown a potential mechanism of the differential sex-based development of CTS in patients with metabolic dysfunction.

Acknowledgments

This study was supported by Mayo Clinic and NIH-NIAMS Grant R01AR076347.

Author Contribution

T. Kuroiwa, A. Gingery, and P. Amadio formulated the concept and designed the research; T. Kuroiwa, A. Gingery, H. Lui, K. Nakagawa, R. Wan, and E. Adam performed the experiments; T. Kuroiwa, N, Iida, and C. Desrochers acquired the data; T. Kuroiwa, D. Larson, PC. Amadio and A, Gingery analyzed and interpreted the data. All authors were involved in drafting and revising the manuscript.

Conflicts of Interest:

The authors declare no conflicts of interest.

References

1.   Atroshi I, Gummesson C, Johnsson R, Ornstein E, Ranstam J, Rosen I. Prevalence of carpal tunnel syndrome in a general population. *Jama.* 1999;282(2):153-158.
2.   Gerritsen AA, de Krom MC, Struijs MA, Scholten RJ, de Vet HC, Bouter LM. Conservative treatment options for carpal tunnel syndrome: a systematic review of randomised controlled trials. *Journal of neurology.* 2002;249(3):272-280.
3.   Gingery A, Yang TH, Passe SM, An KN, Zhao C, Amadio PC. TGF-β signaling regulates fibrotic expression and activity in carpal tunnel syndrome. *J Orthop Res.* 2014;32(11):1444-1450.
4.   Balci K, Utku U. Carpal tunnel syndrome and metabolic syndrome. *Acta neurologica Scandinavica.* 2007;116(2):113-117.
5.   Catrysse L, van Loo G. Inflammation and the Metabolic Syndrome: The Tissue-Specific Functions of NF-κB. *Trends Cell Biol.* 2017;27(6):417-429.
6.   Aihara K, Ikeda Y, Yagi S, Akaike M, Matsumoto T. Transforming Growth Factor-β1 as a Common Target Molecule for Development of Cardiovascular Diseases, Renal Insufficiency and Metabolic Syndrome. *Cardiol Res Pract.* 2010;2011:175381.
7.   Carmiel-Haggai M, Cederbaum AI, Nieto N. A high-fat diet leads to the progression of non-alcoholic fatty liver disease in obese rats. *Faseb j.* 2005;19(1):136-138.
8.   Ding S, Yu L, An B, Zhang G, Yu P, Wang Z. Combination effects of airborne particulate matter exposure and high-fat diet on hepatic fibrosis through regulating the ROS-endoplasmic reticulum stress-TGFβ/SMADs axis in mice. *Chemosphere.* 2018;199:538-545.
9.   Hong SH, Kang M, Lee KS, Yu K. High fat diet-induced TGF-β/Gbb signaling provokes insulin resistance through the tribbles expression. *Sci Rep.* 2016;6:30265.
10.   Koncarevic A, Kajimura S, Cornwall-Brady M, et al. A novel therapeutic approach to treating obesity through modulation of TGFβ signaling. *Endocrinology.* 2012;153(7):3133-3146.
11.   Lin E, Kuo PH, Liu YL, Yang AC, Tsai SJ. Transforming growth factor-β signaling pathway-associated genes SMAD2 and TGFBR2 are implicated in metabolic syndrome in a Taiwanese population. *Sci Rep.* 2017;7(1):13589.
12.   Park YH, Oh EY, Han H, et al. Insulin resistance mediates high-fat diet-induced pulmonary fibrosis and airway hyperresponsiveness through the TGF-β1 pathway. *Experimental & molecular medicine.* 2019;51(5):1-12.
13.   Singh AK, Kari JA. Metabolic syndrome and chronic kidney disease. *Current opinion in nephrology and hypertension.* 2013;22(2):198-203.
14.   Mondelli M, Giannini F, Giacchi M. Carpal tunnel syndrome incidence in a general population. *Neurology.* 2002;58(2):289-294.
15.   Nathan PA, Keniston RC, Meadows KD, Lockwood RS. The relationship between body mass index and the diagnosis of carpal tunnel syndrome. *Muscle Nerve.* 1994;17(12):1491-1493.
16.   Nathan PA, Keniston RC, Myers LD, Meadows KD. Obesity as a risk factor for slowing of sensory conduction of the median nerve in industry. A cross-sectional and longitudinal study involving 429 workers. *Journal of occupational medicine : official publication of the Industrial Medical Association.* 1992;34(4):379-383.
17.   Riccardi G, Giacco R, Rivellese AA. Dietary fat, insulin sensitivity and the metabolic syndrome. *Clinical nutrition (Edinburgh, Scotland).* 2004;23(4):447-456.
18.   Roopchand DE, Carmody RN, Kuhn P, et al. Dietary Polyphenols Promote Growth of the Gut Bacterium Akkermansia muciniphila and Attenuate High-Fat Diet-Induced Metabolic Syndrome. *Diabetes.* 2015;64(8):2847-2858.

19. Cani PD, Amar J, Iglesias MA, et al. Metabolic endotoxemia initiates obesity and insulin resistance. *Diabetes.* 2007;56(7):1761-1772.

20. Lozano WM, Arias-Mutis OJ, Calvo CJ, Chorro FJ, Zarzoso M. Diet-Induced Rabbit Models for the Study of Metabolic Syndrome. *Animals : an open access journal from MDPI.* 2019;9(7).

21. Morelli A, Comeglio P, Filippi S, et al. Mechanism of action of phosphodiesterase type 5 inhibition in metabolic syndrome-associated prostate alterations: an experimental study in the rabbit. *Prostate.* 2013;73(4):428-441.

22. Yoshii Y, Zhao C, Schmelzer JD, Low PA, An KN, Amadio PC. Effects of hypertonic dextrose injections in the rabbit carpal tunnel. *J Orthop Res.* 2011;29(7):1022-1027.

23. Yoshii Y, Zhao C, Schmelzer JD, Low PA, An KN, Amadio PC. The effects of hypertonic dextrose injection on connective tissue and nerve conduction through the rabbit carpal tunnel. *Archives of physical medicine and rehabilitation.* 2009;90(2):333-339.

24. Marchiani S, Vignozzi L, Filippi S, et al. Metabolic syndrome-associated sperm alterations in an experimental rabbit model: relation with metabolic profile, testis and epididymis gene expression and effect of tamoxifen treatment. *Molecular and cellular endocrinology.* 2015;401:12-24.

25. Filippi S, Vignozzi L, Morelli A, et al. Testosterone partially ameliorates metabolic profile and erectile responsiveness to PDE5 inhibitors in an animal model of male metabolic syndrome. *J Sex Med.* 2009;6(12):3274-3288.

26. Maneschi E, Morelli A, Filippi S, et al. Testosterone treatment improves metabolic syndrome-induced adipose tissue derangements. *The Journal of endocrinology.* 2012;215(3):347-362.

27. Alberti KG, Eckel RH, Grundy SM, et al. Harmonizing the metabolic syndrome: a joint interim statement of the International Diabetes Federation Task Force on Epidemiology and Prevention; National Heart, Lung, and Blood Institute; American Heart Association; World Heart Federation; International Atherosclerosis Society; and International Association for the Study of Obesity. *Circulation.* 2009;120(16):1640-1645.

28. Oh S, Ettema AM, Zhao C, et al. Dextrose-induced subsynovial connective tissue fibrosis in the rabbit carpal tunnel: A potential model to study carpal tunnel syndrome? *Hand (New York, NY.* 2008;3(1):34-40.

29. Yoshii Y, Zhao C, Schmelzer JD, Low PA, An KN, Amadio PC. Effects of multiple injections of hypertonic dextrose in the rabbit carpal tunnel: a potential model of carpal tunnel syndrome development. *Hand (New York, NY.* 2014;9(1):52-57.

30. Henderson NC, Sheppard D. Integrin-mediated regulation of TGFβ in fibrosis. *Biochim Biophys Acta.* 2013;1832(7):891-896.

31. Munger JS, Huang X, Kawakatsu H, et al. The integrin alpha v beta 6 binds and activates latent TGF beta 1: a mechanism for regulating pulmonary inflammation and fibrosis. *Cell.* 1999;96(3):319-328.

32. Wang B, Dolinski BM, Kikuchi N, et al. Role of alphavbeta6 integrin in acute biliary fibrosis. *Hepatology.* 2007;46(5):1404-1412.

33. Li J, Yang Y, Li Q, et al. STAT6 contributes to renal fibrosis by modulating PPARα-mediated tubular fatty acid oxidation. *Cell Death Dis.* 2022;13(1):66.

34. Yamanaka Y, Gingery A, Oki G, Yang TH, Zhao C, Amadio PC. Blocking fibrotic signaling in fibroblasts from patients with carpal tunnel syndrome. *Journal of cellular physiology.* 2018;233(3):2067-2074.

35. Zhang J, Chu M. Differential roles of VEGF: Relevance to tissue fibrosis. *Journal of cellular biochemistry.* 2019;120(7):10945-10951.

36. de Castro UG, dos Santos RA, Silva ME, de Lima WG, Campagnole-Santos MJ, Alzamora AC. Age-dependent effect of high-fructose and high-fat diets on lipid metabolism and lipid accumulation in liver and kidney of rats. *Lipids Health Dis.* 2013;12:136.

37. Nishikawa S, Yasoshima A, Doi K, Nakayama H, Uetsuka K. Involvement of sex, strain and age factors in high fat diet-induced obesity in C57BL/6J and BALB/cA mice. *Experimental animals.* 2007;56(4):263-272.

38. Duan Y, Zeng L, Zheng C, et al. Inflammatory Links Between High Fat Diets and Diseases. *Front Immunol.* 2018;9:2649.

39. Becker J, Nora DB, Gomes I, et al. An evaluation of gender, obesity, age and diabetes mellitus as risk factors for carpal tunnel syndrome. *Clin Neurophysiol.* 2002;113(9):1429-1434.

40. Hendriks SH, van Dijk PR, Groenier KH, Houpt P, Bilo HJ, Kleefstra N. Type 2 diabetes seems not to be a risk factor for the carpal tunnel syndrome: a case control study. *BMC musculoskeletal disorders.* 2014;15:346.

41. Onder B, Yalçın E, Selçuk B, Kurtaran A, Akyüz M. Carpal tunnel syndrome and metabolic syndrome co-occurrence. *Rheumatol Int.* 2013;33(3):583-586.

42. Chukijrungroat N, Khamphaya T, Weerachayaphorn J, Songserm T, Saengsirisuwan V. Hepatic FGF21 mediates sex differences in high-fat high-fructose diet-induced fatty liver. *Am J Physiol Endocrinol Metab.* 2017;313(2):E203-e212.

43. Hwang LL, Wang CH, Li TL, et al. Sex differences in high-fat diet-induced obesity, metabolic alterations and learning, and synaptic plasticity deficits in mice. *Obesity (Silver Spring).* 2010;18(3):463-469.

44. Lipson KE, Wong C, Teng Y, Spong S. CTGF is a central mediator of tissue remodeling and fibrosis and its inhibition can reverse the process of fibrosis. *Fibrogenesis & tissue repair.* 2012;5(Suppl 1):S24.

45. Logvinov SV, Naryzhnaya NV, Kurbatov BK, et al. High carbohydrate high fat diet causes arterial hypertension and histological changes in the aortic wall in aged rats: The involvement of connective tissue growth factors and fibronectin. *Exp Gerontol.* 2021;154:111543.

46. Tan JT, McLennan SV, Williams PF, et al. Connective tissue growth factor/CCN-2 is upregulated in epididymal and subcutaneous fat depots in a dietary-induced obesity model. *Am J Physiol Endocrinol Metab.* 2013;304(12):E1291-1302.

47. Yoshino J, Patterson BW, Klein S. Adipose Tissue CTGF Expression is Associated with Adiposity and Insulin Resistance in Humans. *Obesity (Silver Spring).* 2019;27(6):957-962.

48. Chikenji T, Gingery A, Zhao C, et al. Transforming growth factor-β (TGF-β) expression is increased in the subsynovial connective tissues of patients with idiopathic carpal tunnel syndrome. *J Orthop Res.* 2014;32(1):116-122.

49. Li J, Yue S, Fang J, et al. MicroRNA-10a/b inhibit TGF-β/Smad-induced renal fibrosis by targeting TGF-β receptor 1 in diabetic kidney disease. *Molecular therapy Nucleic acids.* 2022;28:488-499.

50. Suo XG, Wang F, Xu CH, et al. Targeted inhibition of TGF-β type I receptor by AZ12601011 protects against kidney fibrosis. *Eur J Pharmacol.* 2022;929:175116.

51. Hahm K, Lukashev ME, Luo Y, et al. Alphav beta6 integrin regulates renal fibrosis and inflammation in Alport mouse. *The American journal of pathology.* 2007;170(1):110-125.

52. Liu S, Kapoor M, Denton CP, Abraham DJ, Leask A. Loss of beta1 integrin in mouse fibroblasts results in resistance to skin scleroderma in a mouse model. *Arthritis and rheumatism.* 2009;60(9):2817-2821.

53. Emmez H, Kardes O, Dogulu F, Kurt G, Memis L, Baykaner MK. Role of antifibrotic cytokine interferon-gamma in the prevention of postlaminectomy peridural fibrosis in rats. *Neurosurgery.* 2008;62(6):1351-1357; discussion 1357-1358.

54. Hasaneen N, Vu T, Fusiak T, Foda H, Condos R, Smaldone G. Anti-fibrotic role of inhaled interferon-γ in idiopathic pulmonary fibrosis. *European Respiratory Journal.* 2015;46(suppl 59):PA2034.

55. Baghy K, Dezso K, Laszlo V, et al. Ablation of the decorin gene enhances experimental hepatic fibrosis and impairs hepatic healing in mice. *Lab Invest.* 2011;91(3):439-451.

56.   Ferdous Z, Wei VM, Iozzo R, Höök M, Grande-Allen KJ. Decorin-transforming growth factor-interaction regulates matrix organization and mechanical characteristics of three-dimensional collagen matrices. *The Journal of biological chemistry.* 2007;282(49):35887-35898.

57.   Yan W, Wang P, Zhao CX, Tang J, Xiao X, Wang DW. Decorin gene delivery inhibits cardiac fibrosis in spontaneously hypertensive rats by modulation of transforming growth factor-beta/Smad and p38 mitogen-activated protein kinase signaling pathways. *Human gene therapy.* 2009;20(10):1190-1200.

58.   Catalán V, Gómez-Ambrosi J, Rodríguez A, et al. Expression of caveolin-1 in human adipose tissue is upregulated in obesity and obesity-associated type 2 diabetes mellitus and related to inflammation. *Clinical endocrinology.* 2008;68(2):213-219.

59.   Shihata WA, Putra MRA, Chin-Dusting JPF. Is There a Potential Therapeutic Role for Caveolin-1 in Fibrosis? *Frontiers in pharmacology.* 2017;8:567.

60.   Zhang GY, He B, Liao T, et al. Caveolin 1 inhibits transforming growth factor-β1 activity via inhibition of Smad signaling by hypertrophic scar derived fibroblasts in vitro. *Journal of dermatological science.* 2011;62(2):128-131.

61.   Nakao A, Afrakhte M, Moren A, et al. Identification of Smad7, a TGFbeta-inducible antagonist of TGF-beta signalling. *Nature.* 1997;389(6651):631-635.

62.   Yan X, Liu Z, Chen Y. Regulation of TGF-beta signaling by Smad7. *Acta biochimica et biophysica Sinica.* 2009;41(4):263-272.

63.   Rahman FA, Krause MP. PAI-1, the Plasminogen System, and Skeletal Muscle. *Int J Mol Sci.* 2020;21(19).

64.   Lemaire R, Burwell T, Sun H, et al. Resolution of Skin Fibrosis by Neutralization of the Antifibrinolytic Function of Plasminogen Activator Inhibitor 1. *Arthritis Rheumatol.* 2016;68(2):473-483.

65.   Kanno Y. The Role of Fibrinolytic Regulators in Vascular Dysfunction of Systemic Sclerosis. *Int J Mol Sci.* 2019;20(3).

66.   Onozuka I, Kakinuma S, Kamiya A, et al. Cholestatic liver fibrosis and toxin-induced fibrosis are exacerbated in matrix metalloproteinase-2 deficient mice. *Biochemical and biophysical research communications.* 2011;406(1):134-140.

67.   Radbill BD, Gupta R, Ramirez MC, et al. Loss of matrix metalloproteinase-2 amplifies murine toxin-induced liver fibrosis by upregulating collagen I expression. *Digestive diseases and sciences.* 2011;56(2):406-416.

Figure legends

Figure 1. Flow diagram of the study. Rabbits were divided into four groups - high fat diet (HFD) or standard diet (STD) and CTS induction (CTS) or Sham. Rabbits were sacrificed 6 weeks after the first injection. Blood tests (BT) were performed at the time of sacrifice.

Figure 2. Sex and diet treatment differences in cholesterol, triglycerides, glucose, and insulin levels of rabbits at sacrifice were measured. No significant interactions were found between high fat diet (HFD) or standard diet (STD) and sex. Diet significantly regulated cholesterol, triglycerides, glucose, and insulin levels. Significant sex differences were found in insulin levels. Upper whisker, maximum value except outlier; box top, 75th percentile; central line, median; box bottom, 25th percentile; lower whisker, minimum value except outlier; small circle, outlier. P values for the main or interaction comparisons (diet and sex) are shown. NS indicates not significant.

Figure 3. SSCT Thickness. (a) Representative histological cross-sectional images of the carpal tunnel inlet (H&E staining) in STD+Sham and HFD+Sham groups. Subsynovial connective tissues (SSCT) thickness between flexor digitorum superficialis (FDS2) and flexor digitorum profundus (FDP). (b) Average SSCT thickness in CTS and Sham induced group treated with high fat diet (HFD) or standard diet (STD). The interaction between diet and CTS induction was significant. (c) Assessment by sex showed that in male rabbits there were significant interaction between diet and CTS induction. (d). Female rabbits CTS induction was the primary driver of SSCT thickness. Means ± standard errors are depicted. P values for the main or interaction comparisons (CTS induction and diet) are shown. *, and ** indicate significant differences between the respective groups with p value < 0.05, and < 0.01, respectively. NS indicates not significant.

Figure 4. SSCT gene expression of CTS induction and diet. Scatter plots (a-d) Yellow dots indicate genes that showed a significant 1.5-fold or greater change relative to STD+Sham, HFD+Sham, or STD+CTS group. Blue dots indicate genes that showed a 0.5-fold or less change. $P < 0.05$ was considered a significant difference. (e) Fold gene expression as compared to sham controls for select genes. Expression values were normalized to STD+Sham. Interaction indicates the significance of the interaction between CTS induction and diet. Means ± standard errors are depicted. P values for the main or interaction comparisons (CTS induction and diet) are shown. *, and ** indicate significant differences between the respective groups.

Figure 5. SSCT gene expression of CTS induction and diet for females and males Fold changes normalized to male STD+Sham or female STD+Sham groups. Induction and diet indicate the significance of the main effects of the CTS induction and HFD, respectively. Means ± standard errors are depicted. P values for the main or interaction comparisons (CTS induction and diet) are shown. *, and ** indicate significant differences between the respective groups.

Fig 1

Fig 2

Fig 3

a

b

SSCT Thickness

Interaction p < 0.05
Diet p < 0.01
Induction p < 0.05

c

Male

Interaction p < 0.05
Diet p < 0.001
Induction NS

d

Female

Interaction NS
Diet NS
Induction p < 0.01

Fig 4

Fig 4e

Fig 5