A binding site for phosphoinositide modulation of voltage-gated sodium channels described by multiscale simulations

Yiechang Lin*#, Elaine Tao*#, James Champion, Ben Corry#

Research School of Biology, Australian National University, Canberra.

These authors contributed equally.

#*yiechang.lin@anu.edu.au, elaine.tao@anu.edu.au, ben.corry@anu.edu.au*

Voltage gated sodium channels (Na_v) are membrane proteins which open to facilitate the inward flux of sodium ions into excitable cells. In response to stimuli, Na_v channels undergo a transition from the resting, closed state to an open state which allows ion influx, before rapidly inactivating. Dysregulation of this functional cycle due to mutations leads to diseases including epilepsy, pain conditions and cardiac disorders, making Na_v channels a significant pharmacological target. Phosphoinositides are important lipid cofactors for ion channel function. The phosphoinositide PI(4,5)P_2 decreases Na_v1.4 activity by increasing the difficulty of channel opening, accelerating fast activation and slowing recovery from fast inactivation. Using multiscale molecular dynamics simulations, we show that PI(4,5)P_2 binds stably to inactivated Na_v at a conserved site within the DIV S4-S5 linker, which couples the voltage sensing domain (VSD) to the pore. As the Na_v C-terminal domain is proposed to also bind here during recovery from inactivation, we hypothesise that PI(4,5)P_2 prolongs inactivation by competing to bind to this site. In atomistic simulations, PI(4,5)P_2 reduces the mobility of both the DIV S4-S5 linker and the DIII-IV linker, responsible for fast inactivation, slowing the conformational changes required for the channel to recover to the resting state. We further show that in a resting state Na_v model, phosphoinositides bind to VSD gating charges, which may anchor them and impede VSD activation. Our results provide a mechanism by which phosphoinositides alter the voltage dependence of activation and the rate of recovery from inactivation, an important step for the development of novel therapies to treat Na_v-related diseases.

Introduction

Voltage-gated sodium (Na_v) channels are critical to the regulation of brain activity, cardiac rhythm, and muscle contraction. Expressed in the membranes of excitable cells, Na_v channels respond to membrane depolarization to open a pore that facilitates the selective flow of sodium current into the cell, initiating the action potential. In mammals, the Na_v channel family consists of nine subtypes (Na_v 1.1-1.9), distributed throughout the central and peripheral nervous system, as well as in cardiac and skeletal muscle (Yu and Catterall, 2003). The Na_v1.4 subtype is predominantly expressed in skeletal myofibers, where it initiates muscle contraction. Genetic mutations in this subtype are associated with various motor dysfunctions, such as both hyperkalaemic and hypokalaemic periodic paralyses (Jurkat-Rott et al., 2000; Mantegazza et al., 2021; Venance et al., 2006). Na_v1.7 is found in peripheral sensory neurons and is responsible for nociception. Several pain disorders, such as inherited erythromelalgia and small fibre neuropathy arise from gain-of-function Na_v1.7 mutations (Dib-Hajj et al., 2013). Both Na_v subtypes have been investigated as promising pharmacological targets for the treatment of myopathy and pain conditions (Dib-Hajj et al., 2013; Ghovanloo et al., 2021).

Structurally, Na$_v$ channels consist of four homologous domains (DI-DIV) arranged in a domain-
swapped configuration (Fig. 1A-B). Each domain comprises of six transmembrane helices (S1-
S6). The central pore domain is formed by S5, S6 helices and selectivity filter (SF), while the four
peripheral voltage-sensing domains (VSDs) are formed by S1-S4 (de Lera Ruiz and Kraus, 2015;
Pan et al., 2018). The pore domain also features lateral fenestrations that provide a pathway for
the access of small molecules to the pore via the membrane (Gamal El-Din et al., 2018) and have
been shown in computational studies to be accessible to lipid tails (Boiteux et al., 2014; Martin
and Corry, 2014; Raju et al., 2013; Tao and Corry, 2022). Additionally, the C-terminal domain
(CTD) extends from the DIV S6 helix into the cytoplasm, where it is thought to associate with the
DIII-IV and DIV S4-S5 linkers in the resting state (Clairfeuille et al., 2019).

Na$_v$ channels adopt distinct functional states during the cycle of membrane depolarization and
repolarization (Fig. 1C). In the resting state, the pore is closed and VSD S4 helices are in the
down, deactivated state. Membrane depolarization triggers the asynchronous transition of VSD I-
III S4 helices to the up, activated conformation, causing Na$_v$ channels to open (Capes et al., 2013;
Cha et al., 1999; Goldschen-Ohm et al., 2013). During prolonged depolarization, VSD-IV moves
upwards, causing the channels to adopt a fast inactivated state in which a three-residue (IFM)
hydrophobic motif located on the intracellular linker between DIII and DIV (DIII-IV linker)
allosterically closes the pore (Yan et al., 2017). The CTD is proposed to bind at the S4-S5 linker
of VSD-IV and sequester the DIII-IV linker during the resting state. During fast inactivation, the
CTD dissociates from VSD-IV and releases the DIII-IV linker to allow IFM binding (Clairfeuille *et*
al., 2019). Upon repolarization, the VSDs deactivate, the IFM motif disassociates, the CTD
rebinds to VSD-IV and the pore returns to the resting state.

Phosphoinositides (PIPs) are important cellular signaling molecules found on the cytoplasmic
leaflet of the mammalian cell membrane. They can exist in seven forms, with phosphorylation
possible at one (PIP1), two (PIP2) or all three (PIP3) positions on the inositol ring, at the 3', 4'
and/or 5' carbons. PIPs, particularly PI(4,5)P$_2$, featuring phosphates at the 4' and 5' carbon
positions (Fig. 1D), are known to bind and modulate the activity of numerous ion channels families
(Hille et al., 2015). These include voltage-gated ion channels, some of which have been resolved
with PI(4,5)P$_2$ bound (Gao et al., 2021; Sun and MacKinnon, 2020). PIP is known to interact with
the VSDs of different potassium channels, to stabilize the positive gating charges and support the
voltage-sensing mechanism (Schmidt et al., 2006). By also binding at the VSD-pore interface in
channels such as K$_v$7.1, PIP is proposed to facilitate coupling of VSD movement to pore opening
(Kasimova et al., 2015; Ma et al., 2022; Sun and MacKinnon, 2020). PI(4,5)P$_2$ also forms specific
interactions with VSD-II of Cav2.2 in the down-state, making channel activation more difficult (Gao
et al., 2021). Although PI(4,5)P$_2$ is known to bind to numerous voltage gated ion channels, its
effects on channel gating and function are complex and yet to be fully elucidated.

Recent experiments show that Na$_v$1.4 channel kinetics and voltage dependence are modulated
by PI(4,5)P$_2$ (Gada et al., 2023). PI(4,5)P$_2$ inhibits Na$_v$1.4 by causing a depolarizing shift in voltage
dependence of activation, accelerating transition to the inactivated state and slowing recovery
from inactivation, resulting in reduced peak current and suppression of late current (summarized
in Fig. 1C). While this is likely to occur via a direct interaction with Na$_v$1.4, the structural basis of
PI(4,5)P$_2$ modulation remains to be understood.

Here, we used a combination of coarse-grained and atomistic MD simulations to identify a putative
PIP binding site to inactivated $Na_v1.4$ in VSD-IV and the DIII-IV linker. We analyze the atomistic
level interactions between the positively charged residues at this site with $PI(4,5)P_2$ and $PI(4)P$,
comparing this with structurally resolved PIP binding sites in the related ion channels $Ca_v2.2$ and
$K_v7.1$. Consistent with the sequence conservation at the identified site, we find that PIPs also bind
to Nav1.7 in coarse-grained simulations, with notable differences dependent on VSD
conformation states. This work provides insight into how PIPs can negatively regulate Na_v
channels, a first step for the potential development of PIP-analogue sodium channel inhibitors.

Fig. 1 Structure of Na_v and modulation by phosphoinositides (A) Na_v channel topology featuring transmembrane
helices (S1-S6), the selectivity filter (SF) and the DIII-IV linker (containing the IFM motif) located between DIII and DIV.
(B) $Na_v1.4$ structure (6agf) showing the four domain-swapped voltage sensor domains (VSDs I-IV), pore and DIII-IV
linker on the intracellular side. **(C)** Summary of $PI(4,5)P_2$ effects on transitions between Na_v channel functional states
(Gada *et al.*, 2023) **(D)** Structure of the $PI(4,5)P_2$ headgroup with the 4'- and 5'- phosphates indicated

Results

Fig. 2 Lipid fingerprint and binding of all PIP types to Na$_v$1.4 (A**) Na$_v$1.4 embedded in a 360 Å x 360 Å model mammalian membrane containing 63 lipid species. **(B) Lipid depletion enrichment index of lipids around Na$_v$1.4 grouped into 12 headgroup types **(C)** Na$_v$1.4, shown from the intracellular (left) and membrane (right) sides, colored by PIP occupancy (darker purple = greater PIP occupancy) **(D)** Distribution of PIP binding occupancies (left) and occupancy of lipid species at 4 residues with the highest PIP occupancy.

To investigate how diverse lipid species interact with Na$_v$1.4, we carried out coarse-grained simulations of Na$_v$1.4 (PDB ID: 6agf, inactivated state) embedded in a complex mammalian membrane for 16 µs in triplicate (Fig. 2A). Glycosphingolipid (GM), phosphoinositide (PIP) and diacylglycerol (DAG) were highly enriched around Nav1.4 (Fig. 2B). Additionally, we observed modest enrichment of lysophosphatidylcholine (LPC), phosphatidylinositol (PI), phosphatidylserine (PS) and phosphatidylethanolamine (PE) and slight depletion of ceramide (CER), sphingomyelin (SM), phosphatidylcholine (PC) and cholesterol (CL). To investigate specific interactions, we generated z-density maps and calculated the per-residue occupancy of the 12 different lipid types (Fig. 2C, Fig. S1-2). Binding residues of interest were identified by constructing occupancy distributions by residue for each lipid type and identifying outlying values with high occupancies (Fig. S3). DG lipids form significant interactions within the lateral fenestrations of Na$_v$1.4 (Fig. S2). LPC and PI also frequently interact with different VSD residues (Fig. S1).

Given our interest in understanding the modulation of Na$_v$1.4 by PIPs, we focus on their interactions for the remainder of this manuscript. Despite the very low concentration of PIPs (0.5% each of PIP1, PIP2 and PIP3) in the mammalian membrane, they are highly enriched around Na$_v$1.4, particularly near the DIII-IV linker and the VSD-IV (Fig. 2C). Contact analysis revealed a putative PIP binding site involving the K1330 residue of the DIII-IV linker and residues R1463, K1466 and R1469 in the DIV S4-S5 linker, which connects the pore and VSD-IV (Fig. 2D). Across all three replicates, only residues within this site were occupied by PIPs for more than 80% (Quartile 3 + 1.5 x Interquartile range) of simulation time on average (Fig. 2D). We note that these

126 residues have higher occupancies for PIP compared to other lipids, including other negatively
127 charged phospholipids (PA, PS and PI) (Fig. 2D).

Fig. 3 Binding of different PIP species in enriched PIP simulations (A) Enriched PIP simulation system, with Na$_v$1.4
embedded in a POPC membrane (transparent gray) and 5% each of PIP1 (blue), PIP2 (purple) and PIP3 (pink) added
to the lower leaflet. **(B)** Representative snapshots showing PIP binding to VSD-IV and the DIII-IV linker. Na$_v$1.4 is shown
in white and PIPs are colored according to residue name (PIP1 in blue, PIP2 in purple and PIP3 in pink) **(C)** A frequency
distribution showing interaction times for each PIP species, defined as the length of a continuous period in which a PIP
was within 0.7 nm of two VSD IV binding site residues. **(D)** Frequency plots showing number of positive residues
interacting with bound PIP in the DIII-IV linker (vertical) and VSD-IV (horizontal). **(E)** Minimum distance between binding
residues on Na$_v$1.4 and bound PIPs lipid across simulation time for the 5 longest binding events, colored by distance
and the type of PIP bound.

Simulations in a complex mammalian membrane showed that PIP species bind specifically and
selectively to Na$_v$1.4 in the presence of other negatively charged lipids and at low, physiological
concentrations. However, the large membrane required, and long PIP binding durations
prevented sampling of large numbers of binding and unbinding events. To address this, we carried
out additional simulations where Na$_v$1.4 was placed in a smaller POPC membrane with a 5%
concentration of each PIP species in the lower leaflet (Fig. 3A).

Across ten replicates of these enriched PIP simulations, each carried out for 80 μs, we observed
all three PIP species binding at the identified site on VSD-IV and DIII-IV linker (Fig. 3B). PIPs can
approach and bind to this site from either side of the VSD, however, PIP1 only forms stable
interactions when it approaches and binds from the VSD-I side while PIP2 and PIP3 usually bind

from the VSD-III side (Fig. 3B). These enriched PIP simulations also revealed additional positively
charged residues in the DIII-IV linker (K1329, K1333 and K1352) and DIV S4 (R1460) which
support binding.

There were 156 PIP binding/unbinding events with duration greater than 2 μs occurring the
identified site (Fig. 3C). Of these, 43 were with PIP1, 44 with PIP2 and 33 with PIP3. The number
of short-term (2-10 μs) interactions decreased with headgroup charge. That is, PIP1 formed the
greatest number of short-term interactions while PIP3 had the fewest. Of the 31 binding events
with duration greater than 10 μs, 7 were with PIP1, 15 with PIP2 and 9 with PIP3 (Fig. S3-4).

When we analyzed interactions occurring during 2-80+ μs binding events, we found that the
number of interacting basic residues changes depending on the PIP headgroup charge (Fig. 3D).
PIP1 (headgroup charge: -3e) binding is most frequently coordinated by two positive charges in
VSD-IV and zero or one residue in the DIII-IV linker. PIP2 (headgroup charge: -5e) binding most
frequently involves one or two interactions from the DIII-IV linker and three from VSD-IV for a total
of five interactions. Interestingly, despite its greater negative charge, PIP3 (headgroup charge: -
7e) has interacts similarly to PIP1, and has fewer interactions than PIP2.

The minimum distance between the interacting PIP headgroup and each binding residue across
simulation time is shown for the five 40+ μs PIP binding events (Fig. 3E). Of these, three are PIP2
binding events which almost span the entire 80 μs of simulation. One 40+ μs binding event each
for PIP1 and PIP3 were also observed. The stable PIP1 binding event observed involved
interactions with R1460 and R1469 in VSD-IV, as well as fluctuating interactions with the four
residues of the DIII-IV linker (K1329, K1330, K1333, K1352). The three long term PIP2 binding
events observed were similar to each other, mainly stabilized by R1460, R1463 and R1469. As
with PIP1, the number and identity of interacting DIII-IV linker residues varied across the span of
each simulation and between replicates, owing to the flexibility of the linker. Like PIP2, PIP3
binding was characterized by stable interactions with R1460, R1463 and R1469 (VSD-IV) as well
as K1329 and K1330 (DIII-IV linker).

Fig. 4 PI(4,5)P₂ and PI(4)P binding to Na$_v$1.4 in atomistic simulations (A) Representative snapshots of PI(4,5)P₂ bound from the VSD-I side (purple stick) and PI(4)P bound from the VSD-III side (cyan stick), with six basic residues forming the binding site located on the DIII-IV linker (orange VDW representation) and VSD-IV S4-S5 linker (shown in green VDW representation) visualized from the intracellular face of the protein **(B)** Proportion of frames where each of the binding site residues were identified to be within 4.5 Å with the different headgroup regions, P4, P5 and PO4, for PI(4,5)P₂ (left) and PI(4)P (right) **(C)** Comparison of RMSF per carbon-alpha for simulations with and without bound PI(4,5)P₂, showing residues on the S4-S5 linker and DIV linker with significant differences in mobility (p-value < 0.05) **(D)** Interaction network plots between the PIP headgroup and basic binding residues on DIII-IV linker (orange) and DIV S4-S5 linker (green), generated by ProLIF – showing the dominant interactions across simulations of PI(4,5)P₂ and PI(4)P.

Coarse-grained simulations enhance sampling by reducing system size, enabling larger time steps and construction of a smoother energy surface. However, this loss of resolution prevents distinction between phosphate positions on the inositol group and does not permit analysis of protein conformational changes induced by PIP binding. To address this, we backmapped representative snapshots from our PIP enriched simulations, where we observed stable, long-term binding events between Na$_v$1.4 and PIP1/PIP2. The coarse-grained PIP1 and PIP2 molecules were converted to the atomistic PI(4)P and PI(4,5)P₂ respectively. For each system, 7.5 μs of atomistic simulations were performed (5 replicates, 1.5 μs each).

In these simulations, the PI(4,5)P₂ headgroup was stable at the binding site identified in coarse-grain simulations (RMSD <2.2 Å) (Fig. 4A, Fig. S6). PI(4,5)P₂ binding is predominantly coordinated by R1469 on the DIV S4-S5 linker, as well as R1466 and R1463. In one replicate, PI(4,5)P₂ associates with R1460 via the phosphate group (PO4) that connects the headgroup to the PIP tails. The PI(4,5)P₂ headgroup also forms electrostatic interactions with K1329 and K1330 in the DIII-IV linker, but forms few contacts with K1333 (Fig. 4B, Fig. S7). This loop portion of the

DIII-IV linker is highly flexible (Fig. S6), thus the lysine sidechains are able to flip between binding
the PI(4,5)P$_2$ headgroup and facing the intracellular space.

The 4'-phosphate formed interactions with residues belonging to the DIII-IV linker (K1329 and
K1333) and DIV S4-S5 linker (K1666 and R1469) with similar frequency, in 55-60% of simulation
frames (Fig. 4B). In contrast, the 5'-phosphate formed contacts with three VSD-IV S4-S5 residues
(R1463, K1466 and R1469) in 70-98% of simulation frames and DIII-IV linker residues in 37-48%
of frames. Taken together, this data suggests that although the headgroup is flexible when bound,
the 5'-phosphate is more important for coordinating VSD-IV S4-S5 residues while the 4'-
phosphate associates with both regions of the binding site.

Atomistic simulations of PI(4)P bound from the VSD-III side show that this is also a stable pose
(RMSD <2 Å), where the headgroup interacts with the same positively charged residues as seen
for PI(4,5)P$_2$ (Fig. 4A, Fig. S6). The residues on the S4-S5 linker, R1463, K1466 and R1469,
predominantly bind to the 4'-phosphate (Fig. 4B). Due to the more buried location of PI(4)P
binding, the PO4 phosphate can associate more with DIII-IV linker lysines (Fig. 4B), however, the
absence of the 5'-phosphate leads to a reduced number of total electrostatic interactions (Fig.
S8).

To investigate structural changes that might occur in the presence of PI(4)P/PI(4,5)P$_2$, we also
carried out simulations of the inactivated Na$_v$1.4 structure without any PIPs bound for comparison.
Although the DIII-IV linker remains bound throughout simulations both with and without PIPs,
residues P1336 and R1337 in the DIII-IV linker downstream of the IFM motif are significantly less
mobile with PI(4,5)P$_2$ present (Fig. 4C). Additionally, seven residues belonging to the DIV S4-S5
linker, including binding residues K1466 and R1469, also have significantly lower mobility when
PI(4,5)P$_2$ is bound (Fig. 4C). In the PI(4)P bound simulations, there were no significant differences
in DIII-IV linker or S4-S5 linker mobility.

Fig. 5 Comparison of the identified phosphoinositide binding site to Na$_v$ subtypes and other ion channels (A)
Binding poses of PI(4,5)P$_2$ (in purple) and PI(4)P (in cyan) aligned with two other tetrameric channels structures K$_v$7.1
(6v01, in yellow) and Ca$_v$2.2 (7mix, in red) that were resolved with PI(4,5)P$_2$ at their respective VSDs. **(B)** Sequence
alignment of the S4 helix and S4-S5 linker of the four domains of Na$_v$1.4, compared to VSD-II of Ca$_v$2.2 and one of the
four identical VSDs of K$_v$7.1; residues colored by amino acid class; purple boxes indicate PI(4,5)P$_2$ binding residues
(identified with 5 Å of the headgroup). **(C)** Sequence alignment of the nine human Na$_v$ channel subtypes shows high
sequence similarity in the S4 helix, S4-S5 linker and DIII-IV linker regions.

The PIP binding poses seen in our simulations is similar to resolved binding poses for PI(4,5)P$_2$
in cryo-EM structures of Ca$_v$2.2 and K$_v$7.1 (Fig. 5A). A sequence alignment of Na$_v$1.4 VSDs show
that there are more positively charged residues present in the S4/S4-S5 linker regions of DIV
compared to the other domains (Fig. 5B). PIP is bound at similar VSD residues on both these ion
channels, with PIP forming interactions with a gating charge further up the S4 helix in Ca$_v$2.2 due
to the VSD being in a different state. Additionally, the high sequence similarity in the S4/S5-S5
linker and DIII-IV linker regions between the nine human Na$_v$ channel subtypes suggests
conservation of the PIP binding site (Fig. 5C).

Fig. 6 PIP binding to Nav1.7 with different VSD states in coarse-grained simulations (A) Atomistic representation
of the three different Nav1.7 structures simulated: (1) the inactivated state (blue, PDB ID: 6j8g) with the VSDs all in the
activated, up state (2) the Nav1.7-NavPas chimera (pink, PDB ID: 6nt4) with the Nav1.7 VSD-IV in the deactivated,
down state and a bound NavPas CTD (3) a Nav1.7 resting state model (orange, model generation detailed in Table.
S2) with all four VSDs in the deactivated, down state. Panel insets show the different conformations of VSD-IV (left)
and VSD-II (right) across different structures. The inactivation switch formed by the CTD and VSD-IV S4-S5 linker
proposed by Clairfeuille et al. is shown (middle). **(B)** Occupancy of all PIPs (PIP1, PIP2, PIP3) at binding site residues
in the three systems **(C)** Distribution of PIP binding durations at the identified site **(D)** Representative simulation
snapshot of PIP binding at DIV S4-S5 linker in the Nav1.7-NavPas system. The CTD (dark pink) prevents PIP access
to DIII-IV linker lysines **(E)** PIP occupancy at the bottom three gating charges on VSD I-III in the inactivated (blue) and
resting state model (orange) simulations. For VSD-IV, PIP occupancy in the Nav1.7-NavPas system (pink) is also
shown. (F) Representative simulation snapshot showing PIP (purple) binding at the gating charges (orange) in the
resting state model simulations.

To assess possible state- and subtype-dependent differences in PIP binding, we performed
embedded three structures of Nav1.7 with different VSD conformations in a PIP enriched
membrane (Fig. 6A). The inactivated Nav1.7 structure (blue, PDB ID: 6j8g) contains all VSDs in
the activated/up state and a bound DIII-IV linker. We also simulated a NavPas (American
cockroach) chimera structure with a human Nav1.7 VSD-IV in the deactivated/down state (VSDs
I-III are activated, and from NavPas) (pink, PDB ID: 6nt4). This structure features a dissociated
DIII-IV linker and a resolved C-terminal domain bound to the DIV S4-S5 linker (forming the

'inactivation switch') at residues identified to form part of our PIP binding site. To assess PIP
binding at VSD I-III in the deactivated/down state, we modelled the $Na_v1.7$ resting state based on
templates structures in which different VSDs have been captured in deactivated states with the
aid of toxins (detailed in Table S2). These structures feature three or more of the gating charge
residues below the hydrophobic constriction site (HCS) and displacement of the S4-S5 linker (Fig.
S9). Given that no resting state mammalian Na_v channel structure has been resolved, it is possible
that the modelled VSDs may not reflect the fully deactivated state.

In triplicate 50 us coarse-grained simulations, PIPs bind to the analogous site to that seen in
$Na_v1.4$ in the inactivated $Na_v1.7$ structure, interacting with residues belonging to both the DIII-IV
linker and VSD-IV for durations comparable to Nav1.4, (Fig. 6B-C, Fig. S10). Binding of PIP to
the DIV S4-S5 linker to the deactivated VSD-IV in the Nav1.7-NavPas chimera and resting state
model was also observed (Fig. 6B-C). However, in the Nav1.7-NavPas chimera, the PIP bound
at the S4-S5 linker cannot simultaneously associate with the DIII-IV linker (Fig. S10), due to its
sequestration by the C-terminal domain, which moves the lysines away from the binding VSD-IV
residues (Fig. 6D). Instead, K1491, K1492 and K1495 (on the DIII-IV linker) are occupied by
different PIPs on the other side of the VSD. In the resting state model, which features an unbound
DIII-IV linker PIPs binding at the DIV S4-S5 residues also do not associate with the DIII-IV linker
(Fig S10). Comparison of binding durations at this site across the three systems reveals a greater
number of long PIP interactions ($> 20 \mu s$) with inactivated Nav1.7 (5) compared to Nav1.7-NavPas
(3) or the resting state model (2) where VSD-IV is down, and the DIII-IV linker is dissociated.

There are state-dependent differences in PIP occupancy at the gating charges in each VSD (Fig.
6E-F, Fig. S11). PIP can associate with the lowest three gating charges VSD-I of the resting state
model, but not in the inactivated state (Fig. 6E). This is likely due to the large displacement in the
DI-S4 helix, which moves down three helical turns in the resting state model so that all gating
charges are below the HCS (Fig. S9). In VSD II and III, the PIP interaction differences between
inactivated and resting state are present but less pronounced, owing to a smaller difference in the
relative displacement between the gating charges between states (Fig. S9). PIP occupancy is
also higher at VSD-IV when it is in the deactivated conformation, however, the presence of a
bound CTD in the Nav1.7-NavPas chimera reduces this.

Discussion

Recently, $PI(4,5)P_2$ was shown to be a negative regulator of $Na_v1.4$, modulating channel kinetics
and voltage dependence. Presence of $PI(4,5)P_2$ causes a depolarizing shift in the voltage
dependence of activation, making $Na_v1.4$ opening more difficult in a physiological context.
Additionally, it stabilizes the inactivated state of $Na_v1.4$, marked by both shortened inactivation
times and slowed recovery from the inactivation.

Using a multiscale simulation approach, we identified a putative PIP binding site comprised of
positively charged residues belonging to the S4 helix/S4-S5 linker of VSD-IV (R1460, R1463,
K1466, R1469) and DIII-IV linker (K1329, K1330, K1333, K1352). Coarse grained simulations of
$Na_v1.4$ embedded in a complex membrane showed that PIP interacts with residues belonging to
VSD-IV and the DIII-IV linker. In coarse grained enriched PIP simulations, PIP2 formed longer

duration interactions with Na$_v$1.4 than PIP1 and PIP3, supported by a greater number of charged
interactions. Atomistic simulations verified the stability of PI(4,5)P$_2$ (the most common PIP2
species in the plasma membrane) at this site and showed that the binding of PI(4,5)P$_2$ reduces
the mobility of some DIV S4-S5 and DIII-IV linker residues. Simulations of Na$_v$1.7 with VSDs in
different conformational states showed that the PIP binding site is conserved in Na$_v$1.7 and that
PIP interactions at VSD gating charges are functional state dependent, with more interactions
being formed when the VSDs are deactivated.

The DIII-IV linker, CTD and S4-S5 linkers all play key roles throughout the Na$_v$ conformational
cycle. Mutation of the IFM inactivation motif as well as other residues in the DIII-IV linker alter fast
inactivation and recovery from fast inactivation (McPhee et al., 1998; West et al., 1992). While the
precise role of the Na$_v$ C-terminal domain and its conformation during the Na$_v$ activation cycle
remain elusive, it is likely to be important for coordinating fast inactivation (Clairfeuille *et al.*, 2019;
Mantegazza et al., 2001). CTD binding to DIV S4-S5 and sequestration of the DIII-IV linker is
proposed to occur in the resting state. After the pore opens, activation of VSD-IV is thought to
cause CTD dissociation, releasing the DIII-IV linker to allow fast inactivation. Residues in 'switch
1' of the CTD-binding site on the DIV S4-S5 stably bind PIP in our simulations of inactivated
Nav1.4 and Nav1.7. We hypothesize that PIP binding at this location makes it more difficult for
the CTD to reassociate with VSD-IV, a conformational change which is required during recovery
from inactivation.

More generally, the S4-S5 linkers in all four domains couple the VSD to the pore helices and
adopt different orientations depending on VSD activation states. When the VSD is activated (in
the open and inactivated states), the S4-S5 linkers lie parallel to the membrane. In the resting
state, when the voltage sensor is deactivated, the S4-S5 linkers move downward below the plane
of the membrane. We propose that PIP binding at the identified site could additionally stabilize
both the DIV S4-S5 linker and DIII-IV linker to favor the inactivated state. Although the recovery
from fast inactivation occurs on the order of several milliseconds (Bezanilla and Armstrong, 1977),
and is beyond atomistic simulation timescales, we observed statistically significant reductions in
the RMSF of several DIII-IV linker and DIV S4-S5 linker residues when PI(4,5)P$_2$ was bound in
1.5 μs atomistic simulations. Reduction in the mobility of the DIII-IV linker may slow the
dissociation of the upstream IFM motif and stabilization of the S4-S5 linker prevents the downward
movement required for the channel to transition back to the resting state.

The PIP binding residues identified here are conserved in Na$_v$1.1-1.9 (Fig. 5D), suggestive of a
shared binding site and mechanism for PIP-mediated modulation across subtypes. Mutations at
these conserved residues in other subtypes lead to various gain-of-function diseases (Table S1).
For example, analogous to the Na$_v$1.4 R1469 residue, the R1642C mutation (in Na$_v$1.3) leads to
developmental epileptic encepahlopathy (Zaman et al., 2018), and R1644C/H mutations (in
Na$_v$1.5) cause cardiac arrythmias (Frustaci et al., 2005). These diseases are associated with
accelerated rates of channel recovery from inactivation, consistent with our observations that an
interaction between PI(4,5)P$_2$ and the residue corresponding to R1469 in other Na$_v$ subtypes
could be important for prolonging the fast-inactivated state.

The PIP binding site identified here harbors sequence and structural similarity to PI(4,5)P$_2$ binding
sites found in other cation channels (Fig. 5B). For example, PI(4,5)P$_2$ is resolved at a similar site
near the VSD and S4-S5 linker in a recent cryo-EM structure of K$_v$7.1, where the phosphate

headgroup forms analogous contacts to R249 and R243 (PDB ID: 6v01) (Sun and MacKinnon,
2020). Despite differences in the role of PI(4,5)P$_2$, which negatively regulates Na$_v$1.4 but is
required for K$_v$7.1 pore opening, the binding site appears to be conserved. Based on the PI(4,5)P$_2$
binding site, a structurally similar compound was developed as an activator of K$_v$7 channels and
proposed to be a future antiarrhythmic therapy (Liu et al., 2020).

PI(4,5)P$_2$ also binds to the down, deactivated state of VSD-II in Ca$_v$2.2 (PDB ID: 7mix) (Gao *et*
al., 2021). In this structure, the PI(4,5)P$_2$ headgroup interacts with two VSD-II gating charges,
R584 and K587. Compared to the positioning of PI(4,5)P$_2$ in our simulations of Na$_v$1.4 with an
activated VSD-IV, the headgroup associates further up the S4 helix in Ca$_v$2.2 due to the VSD
being in a deactivated state. This is also seen in K$_v$7.1 which contains an extended GGT loop in
the S4-S5 linker which prevents PI(4,5)P$_2$ binding in the VSD-down state (Mandala and
MacKinnon, 2023). In our coarse-grained simulations of the resting Na$_v$1.7 model, we observe a
similar state-dependent difference in PI(4,5)P$_2$ interactions with the deactivated states of each
VSD. Since activation of VSDs I-III are known to be coupled to channel opening [], we propose
that PIP binding at these VSDs impedes their ability to activate and thus increases the voltage
threshold required of opening.

The leftward shift in voltage dependence of inactivation is less pronounced when PI(4,5)P$_2$ is
converted to PI(4)P rather than completely dephosphorylated to PI (Gada *et al.*, 2023). This
suggests that PI(4)P may play a compensatory role when PI(4,5)P$_2$ is not present. This is
supported by our simulations which show that PI(4)P can also stably occupy the identified binding
site, albeit with shorter duration and form less electrostatic interactions compared to PI(4,5)P$_2$.
Our atomistic simulations also showed that the 5'-phosphate is more important than the 4'-
phosphate for forming interactions with the DIV S4-S5 linker residues of the inactivation switch.
These factors might suggest that PI(4,5)P$_2$ binding is preferred over PI(4)P at this site and can
better compete to bind over the CTD, suggesting that PI(4,5)P$_2$ is more effective at stabilizing the
inactivated state and preventing recovery to the resting state.

Simulations using the Martini2.2 forcefield have previously been used to investigate lipid-protein
interactions and successfully predict specific lipid binding sites, including the PIP binding site on
Kir channels (Stansfeld et al., 2009). While Martini2.2 simulations explicitly represent all three PIP
species (PIP1, PIP2 and PIP3) present in the mammalian membrane, it does not distinguish
between phosphate positions within each charge state (e.g. PI(3,4)P$_2$ vs PI(4,5)P$_2$). In this work,
we employed atomistic interactions to complement and strengthen findings from coarse grain
simulations, allowing us to identify the specific contribution of each PI(4,5)P$_2$ phosphate group to
binding as well as investigate the conformational changes associated with PI(4,5)P$_2$ binding.

Fig. 7 Proposed mechanism of PIP effects on the sodium channel functional cycle

Conclusion

Using multiscale simulations, we show that PI(4,5)P$_2$ binds stably to inactivated structures of Na$_v$1.4 and Na$_v$1.7 at a conserved site within the DIV S4-S5 linker. As the CTD is proposed to also bind here during recovery from inactivation, we hypothesize that PI(4,5)P$_2$ competes with the CTD to bind to this site, prolonging inactivation. At this site, PI(4,5)P$_2$ simultaneously binds to the DIII-IV linker which is responsible for allosterically blocking the pore during fast inactivation (Fig. 7). Its binding reduces the mobility of both the DIV S4-S5 and DIII-IV linkers, potentially slowing the conformational changes required for the channel to recover to the resting state. We also propose that in the resting state, PIPs form additional interactions with S4 gating charges, particularly in VSD-1, anchoring them to the membrane in a way which may make the upwards movement required for their activation more difficult. Our results provide insight into how sodium channels are modulated by phosphoinositides, an important step for the development of novel therapies to treat Nav-related diseases.

Methods

Coarse grained simulations

Coarse-grained simulations of Na$_v$1.4 embedded in a complex mammalian membrane were carried out to investigate lipid-protein interactions. The inactivated Na$_v$1.4 alpha subunit (PDB ID: 6agf) (Pan *et al.*, 2018) was coarse-grained using the CHARMM-GUI Martini Maker (Jo et al., 2008; Qi et al., 2015) and embedded in a 360 Å x 360 Å complex membrane using insane.py (Wassenaar et al., 2015). The composition of the complex mammalian membrane is as reported in Ingólfsson et al. (2017). Three replicate simulations, each with different starting coordinates, were carried out for 16 μs each.

To better sample binding events, we also carried out PIP-enriched simulations in which $Na_v1.4$
was embedded in a 160 Å x 160 Å POPC membrane with 5% of each PIP species (PIP1, PIP2,
PIP3) added to the cytoplasmic leaflet using insane.py. Ten replicate simulations, each with
different starting coordinates, were carried out for 80 μs each.

To explore possible state and subtype dependent differences in PIP binding, the inactivated
$Na_v1.7$ structure (PDB ID: 6j8g) (Shen et al., 2019) and the $Na_v1.7$-NavPas chimera with CTD
bound and VSD-IV in the deactivated state (PDB ID: 6nt4) (Clairfeuille *et al.*, 2019) were also
coarse-grained and simulated in PIP-enriched membranes (same protocol as above) for three
replicates of 50 μs each. Additionally, a model $Na_v1.7$ with all four VSDs in the deactivated state
using Modeller (Šali and Blundell, 1993). The template and structural information for this model
are detailed in Table S2. In brief, VSD-I down was modelled from 7xve (Huang et al., 2022), VSD-
II and VSD-III were both modelled from the deactivated VSD-II from 7k48 (Wisedchaisri et al.,
2021), and VSD-IV and the unbound DIII-IV linker were modelled from the corresponding regions
in 6nt4 (Clairfeuille *et al.*, 2019). Adjacent S5/S6 regions to each VSD were also modelled from
each specified template to ensure proper contacts between the pore domain and VSDs. The CTD
was not included in the model.

All systems were solvated and ionized with 150 mM NaCl. All coarse-grained simulations were
carried out with GROMACS 2022 (Bauer et al., 2023) using the Martini2.2 forcefield (de Jong et
al., 2013). Energy minimization was carried out on each system using the steepest descent
method for 1,000 steps. Following this, equilibration in the constant pressure, constant volume
(NVT) ensemble at 1 atm for 10 ps was carried out with backbone position restraints (1,000 kJ
$mol^{-1}nm^{-2}$) using a 2-fs timestep. Following this, constant pressure and temperature (NPT)
equilibration simulations were carried out, using 5-, 10-, and 20-fs timesteps in sequence, with
each running for 5,000 steps. 1 atm pressure was maintained using a Berendsen barostat with
semi-isotropic conditions. Production simulations were carried out in the NPT ensemble, kept at
a temperature of 310 K using the Nose-Hoover thermostat (Evans and Holian, 1985) and a
pressure of 1 bar using the Parrinello-Rahman barostat (Parrinello and Rahman, 1981). A time
step of 20 fs was used. During production simulations, the backbone beads were weakly
restrained to their starting coordinates using a force constant of 10 kJ $mol^{-1}nm^{-2}$.

Atomistic simulations

Atomistic simulations were performed to characterize atomic interactions between $Na_v1.4$
residues and the bound $PI(4,5)P_2$ headgroup. Frames from a stable PIP2 binding event (from
replicate 1 of enriched PIP simulations) were clustered using a selection of the bound PIP2
headgroup beads (C1 C2 C3 PO4 P1 P2) and binding residues K1329, K1330, K1333, K1463,
K1466 and R1469 with an RMSD cutoff of 2.5 Å. The protein and bound PIP2 were extracted
from the representative frame of the cluster and backbone beads of the coarse-grained VSD-IV
were aligned to the corresponding carbon-alpha atoms in the original cryo-EM structure of $Na_v1.4$
(Pan *et al.*, 2018). PIP2 was backmapped to atomistic coordinates of SAPI24 (the CHARMM lipid
for $PI(4,5)P_2$ with protonation at the 4-position phosphate) and the protein was replaced with the
6agf structure. The system was embedded in a 140 Å x 140 Å POPC membrane, solvated and
0.15 M NaCl added using the CHARMM-GUI Membrane Builder (Jo *et al.*, 2008; Lee et al., 2016;
Wu et al., 2014). An identical system was set up with $Na_v1.4$ in a POPC membrane without PIP.

Atomistic simulations were performed with Amber20 (Case et al., 2020), using the CHARMM36m
(Huang et al., 2017) and TIP3P water (Jorgensen et al., 1983) forcefields. Equilibration steps
were performed (minimisation, heating, pressurising), with 5 kJ/mol restraints on the protein
backbone, followed by 24 ns of gradually reducing restraints. Five replicates of unrestrained
production equilibrium simulations were performed, run for 1.5 μs each. The temperature was
set at 310 K using the Langevin thermostat (Loncharich et al., 1992) and at a collision frequency
of 5 ps^{-1}. Pressure was set at 1 bar using the Monte Carlo barostat (Åqvist et al., 2004) with
anisotropic scaling and relaxation time of 1 ps. 12 Å van der Waals cut-off and hydrogen bond
SHAKE constraints were used. Hydrogen mass repartitioning was used to enable a 4 fs timestep
(Hopkins et al., 2015).

Analysis

Simulations were visualised and protein image figures produced using Visual Molecular Dynamics
(VMD) (Humphrey et al., 1996).

Coarse grain lipid-protein interactions were characterized using in-house python scripts which
used the numpy, MDAnalysis (Michaud-Agrawal et al., 2011) and pandas libraries (as done
previously in Lin et al. (2022)). A cut-off of 0.7 nm was used to define interactions between lipids
and protein residues. Distance heatmaps were generated based on the minimum distance
between a PIP bead and the side chain (SC2) bead of the arginine or lysine residue interacting.
Binding durations were calculated by first counting the number of interactions between each PIP
and VSD-IV binding residues (R1460, R1463, K1466 and R1469). A binding event is defined as
the time between the first and last time that PIP interacts with two or more binding residues on
VSD-IV, provided that it interacts with a minimum of one binding residue between this period.

For atomistic simulations, MDAnalysis (Michaud-Agrawal *et al.,* 2011) was used to calculate
RMSD of various parts of the protein and PIP headgroup, and RMSF of the carbon-alphas.
Statistical significance in RMSF was assessed using student's t-test. ProLIF (Bouysset and
Fiorucci, 2021) was used to compute electrostatic interactions between the protein and binding
residues. Representative snapshot of PIP2 binding was generated using the WMC Clustering
Tool in VMD to identify the top cluster of the PIP2 headgroup (RMSD cutoff of 3 Å), then
subsequently cluster the six binding residues to identify the most representative binding
configuration. Trajectories were strided every 1 ns and the first 250 ns of simulations was
discarded as equilibration time for analyses of RMSF, ProLIF interactions and clustering.

ClustalOmega (Goujon et al., 2010; Sievers et al., 2011) and JalView (Waterhouse et al., 2009)
were used to generate and visualize sequence alignments. Structural representations of $K_v7.1$,
$Ca_v2.2$ and $Na_v1.4$ structures were created in VMD by aligning each of the S4 helices on the VSD
where the PIP was bound.

Acknowledgements

482 This research was undertaken with the assistance of resources and services from the National
483 Computational Infrastructure (NCI), which is supported by the Australian Government. This
484 research is also supported by Australian Government Research Training Program (RTP)
485 Scholarships. Y.L. is supported by a 2023 NCI HPC-AI Talent Program Scholarship.

References

Åqvist, J., Wennerström, P., Nervall, M., Bjelic, S., and Brandsdal, B.O. (2004). Molecular dynamics simulations of water and biomolecules with a Monte Carlo constant pressure algorithm. Chemical physics letters *384*, 288-294.

Bauer, P., Hess, B., and Lindahl, E. (2023). GROMACS 2022.5 Manual (2022.5).

Bezanilla, F., and Armstrong, C.M. (1977). Inactivation of the sodium channel. I. Sodium current experiments. The Journal of general physiology *70*, 549.

Boiteux, C., Vorobyov, I., and Allen, T.W. (2014). Ion conduction and conformational flexibility of a bacterial voltage-gated sodium channel. Proceedings of the National Academy of Sciences *111*, 3454-3459.

Bouysset, C., and Fiorucci, S. (2021). ProLIF: a library to encode molecular interactions as fingerprints. Journal of Cheminformatics *13*, 1-9.

Capes, D.L., Goldschen-Ohm, M.P., Arcisio-Miranda, M., Bezanilla, F., and Chanda, B. (2013). Domain IV voltage-sensor movement is both sufficient and rate limiting for fast inactivation in sodium channels. Journal of General Physiology *142*, 101-112.

Case, D.A., Belfon, K., Ben-Shalom, I.Y., Brozell, S.R., Cerutti, D.S., T.E. Cheatham, I., Cruzeiro, V.W.D., Darden, T.A., Duke, R.E., Giambasu, G., et al. (2020). AMBER 2020.

Cha, A., Ruben, P.C., George, A.L., Fujimoto, E., and Bezanilla, F. (1999). Voltage sensors in domains III and IV, but not I and II, are immobilized by Na+ channel fast inactivation. Neuron *22*, 73-87.

Clairfeuille, T., Cloake, A., Infield, D.T., Llongueras, J.P., Arthur, C.P., Li, Z.R., Jian, Y., Martin-Eauclaire, M.-F., Bougis, P.E., and Ciferri, C. (2019). Structural basis of α-scorpion toxin action on Nav channels. Science *363*, eaav8573.

de Jong, D.H., Singh, G., Bennett, W.D., Arnarez, C., Wassenaar, T.A., Schafer, L.V., Periole, X., Tieleman, D.P., and Marrink, S.J. (2013). Improved parameters for the martini coarse-grained protein force field. Journal of chemical theory and computation *9*, 687-697.

de Lera Ruiz, M., and Kraus, R.L. (2015). Voltage-Gated Sodium Channels: Structure, Function, Pharmacology, and Clinical Indications. Journal of Medicinal Chemistry *58*, 7093-7118. 10.1021/jm501981g.

Dib-Hajj, S.D., Yang, Y., Black, J.A., and Waxman, S.G. (2013). The NaV1. 7 sodium channel: from molecule to man. Nature Reviews Neuroscience *14*, 49-62.

Evans, D.J., and Holian, B.L. (1985). The nose–hoover thermostat. The Journal of chemical physics *83*, 4069-4074.

Frustaci, A., Priori, S.G., Pieroni, M., Chimenti, C., Napolitano, C., Rivolta, I., Sanna, T., Bellocci, F., and Russo, M.A. (2005). Cardiac histological substrate in patients with clinical phenotype of Brugada syndrome. Circulation *112*, 3680-3687.

Gada, K.D., Kamuene, J.M., Chandrashekar, A., Kissell, R.C., Yauch, A.K., and Plant, L.D. (2023). PI (4, 5) P2 regulates the gating of NaV1. 4 channels. Journal of General Physiology *155*, e202213255.

Gamal El-Din, T.M., Lenaeus, M.J., Zheng, N., and Catterall, W.A. (2018). Fenestrations control resting-state block of a voltage-gated sodium channel. Proceedings of the National Academy of Sciences *115*, 13111-13116.

Gao, S., Yao, X., and Yan, N. (2021). Structure of human Cav2. 2 channel blocked by the painkiller ziconotide. Nature *596*, 143-147.

Ghovanloo, M.-R., Choudhury, K., Bandaru, T.S., Fouda, M.A., Rayani, K., Rusinova, R., Phaterpekar, T., Nelkenbrecher, K., Watkins, A.R., and Poburko, D. (2021). Cannabidiol inhibits the skeletal muscle Nav1. 4 by blocking its pore and by altering membrane elasticity. Journal of General Physiology *153*, e202012701.

Goldschen-Ohm, M.P., Capes, D.L., Oelstrom, K.M., and Chanda, B. (2013). Multiple pore conformations
driven by asynchronous movements of voltage sensors in a eukaryotic sodium channel. Nature
communications *4*, 1350.
Goujon, M., McWilliam, H., Li, W., Valentin, F., Squizzato, S., Paern, J., and Lopez, R. (2010). A new
bioinformatics analysis tools framework at EMBL–EBI. Nucleic acids research *38*, W695-W699.
Hille, B., Dickson, E.J., Kruse, M., Vivas, O., and Suh, B.-C. (2015). Phosphoinositides regulate ion
channels. Biochimica Et Biophysica Acta (BBA)-Molecular and Cell Biology of Lipids *1851*, 844-856.
Hopkins, C.W., Le Grand, S., Walker, R.C., and Roitberg, A.E. (2015). Long-time-step molecular dynamics
through hydrogen mass repartitioning. Journal of chemical theory and computation *11*, 1864-1874.
Huang, G., Wu, Q., Li, Z., Jin, X., Huang, X., Wu, T., Pan, X., and Yan, N. (2022). Unwinding and spiral
sliding of S4 and domain rotation of VSD during the electromechanical coupling in Nav1. 7. Proceedings
of the National Academy of Sciences *119*, e2209164119.
Huang, J., Rauscher, S., Nawrocki, G., Ran, T., Feig, M., De Groot, B.L., Grubmüller, H., and MacKerell Jr,
A.D. (2017). CHARMM36m: an improved force field for folded and intrinsically disordered proteins.
Nature methods *14*, 71-73.
Humphrey, W., Dalke, A., and Schulten, K. (1996). VMD: visual molecular dynamics. Journal of molecular
graphics *14*, 33-38.
Ingólfsson, H.I., Carpenter, T.S., Bhatia, H., Bremer, P.-T., Marrink, S.J., and Lightstone, F.C. (2017).
Computational lipidomics of the neuronal plasma membrane. Biophysical journal *113*, 2271-2280.
Jo, S., Kim, T., Iyer, V.G., and Im, W. (2008). CHARMM-GUI: a web-based graphical user interface for
CHARMM. Journal of computational chemistry *29*, 1859-1865.
Jorgensen, W.L., Chandrasekhar, J., Madura, J.D., Impey, R.W., and Klein, M.L. (1983). Comparison of
simple potential functions for simulating liquid water. The Journal of chemical physics *79*, 926-935.
Jurkat-Rott, K., Mitrovic, N., Hang, C., Kouzmenkine, A., Iaizzo, P., Herzog, J., Lerche, H., Nicole, S., Vale-
Santos, J., and Chauveau, D. (2000). Voltage-sensor sodium channel mutations cause hypokalemic
periodic paralysis type 2 by enhanced inactivation and reduced current. Proceedings of the National
Academy of Sciences *97*, 9549-9554.
Kasimova, M.A., Zaydman, M.A., Cui, J., and Tarek, M. (2015). PIP2-dependent coupling is prominent in
Kv7. 1 due to weakened interactions between S4-S5 and S6. Scientific reports *5*, 1-9.
Lee, J., Cheng, X., Swails, J.M., Yeom, M.S., Eastman, P.K., Lemkul, J.A., Wei, S., Buckner, J., Jeong, J.C.,
and Qi, Y. (2016). CHARMM-GUI input generator for NAMD, GROMACS, AMBER, OpenMM, and
CHARMM/OpenMM simulations using the CHARMM36 additive force field. Journal of chemical theory
and computation *12*, 405-413.
Lin, Y.C., Buyan, A., and Corry, B. (2022). Characterizing the lipid fingerprint of the mechanosensitive
channel Piezo2. Biophysical Journal *121*, 369a.
Liu, Y., Xu, X., Gao, J., Naffaa, M.M., Liang, H., Shi, J., Wang, H.Z., Yang, N.-D., Hou, P., and Zhao, W.
(2020). A PIP2 substitute mediates voltage sensor-pore coupling in KCNQ activation. Communications
biology *3*, 385.
Loncharich, R.J., Brooks, B.R., and Pastor, R.W. (1992). Langevin dynamics of peptides: The frictional
dependence of isomerization rates of N-acetylalanyl-N'-methylamide. Biopolymers: Original Research on
Biomolecules *32*, 523-535.
Ma, D., Zhong, L., Yan, Z., Yao, J., Zhang, Y., Ye, F., Huang, Y., Lai, D., Yang, W., and Hou, P. (2022).
Structural mechanisms for the activation of human cardiac KCNQ1 channel by electro-mechanical
coupling enhancers. Proceedings of the National Academy of Sciences *119*, e2207067119.
Mandala, V.S., and MacKinnon, R. (2023). The membrane electric field regulates the PIP2-binding site to
gate the KCNQ1 channel. Proceedings of the National Academy of Sciences *120*, e2301985120.
Mantegazza, M., Cestèle, S., and Catterall, W.A. (2021). Sodium channelopathies of skeletal muscle and
brain. Physiological Reviews *101*, 1633-1689.

Mantegazza, M., Yu, F.H., Catterall, W.A., and Scheuer, T. (2001). Role of the C-terminal domain in inactivation of brain and cardiac sodium channels. Proceedings of the National Academy of Sciences *98*, 15348-15353.

Martin, L.J., and Corry, B. (2014). Locating the route of entry and binding sites of benzocaine and phenytoin in a bacterial voltage gated sodium channel. PLoS computational biology *10*, e1003688.

McPhee, J.C., Ragsdale, D.S., Scheuer, T., and Catterall, W.A. (1998). A critical role for the S4-S5 intracellular loop in domain IV of the sodium channel α-subunit in fast inactivation. Journal of Biological Chemistry *273*, 1121-1129.

Michaud-Agrawal, N., Denning, E.J., Woolf, T.B., and Beckstein, O. (2011). MDAnalysis: a toolkit for the analysis of molecular dynamics simulations. Journal of computational chemistry *32*, 2319-2327.

Pan, X., Li, Z., Zhou, Q., Shen, H., Wu, K., Huang, X., Chen, J., Zhang, J., Zhu, X., and Lei, J. (2018). Structure of the human voltage-gated sodium channel Nav1. 4 in complex with β1. Science *362*, eaau2486.

Parrinello, M., and Rahman, A. (1981). Polymorphic transitions in single crystals: A new molecular dynamics method. Journal of Applied physics *52*, 7182-7190.

Qi, Y., Ingólfsson, H.I., Cheng, X., Lee, J., Marrink, S.J., and Im, W. (2015). CHARMM-GUI martini maker for coarse-grained simulations with the martini force field. Journal of chemical theory and computation *11*, 4486-4494.

Raju, S., Barber, A.F., LeBard, D.N., Klein, M.L., and Carnevale, V. (2013). Exploring volatile general anesthetic binding to a closed membrane-bound bacterial voltage-gated sodium channel via computation. PLoS computational biology *9*, e1003090.

Šali, A., and Blundell, T.L. (1993). Comparative protein modelling by satisfaction of spatial restraints. Journal of molecular biology *234*, 779-815.

Schmidt, D., Jiang, Q.-X., and MacKinnon, R. (2006). Phospholipids and the origin of cationic gating charges in voltage sensors. Nature *444*, 775-779.

Shen, H., Liu, D., Wu, K., Lei, J., and Yan, N. (2019). Structures of human Nav1. 7 channel in complex with auxiliary subunits and animal toxins. Science *363*, 1303-1308.

Sievers, F., Wilm, A., Dineen, D., Gibson, T.J., Karplus, K., Li, W., Lopez, R., McWilliam, H., Remmert, M., and Söding, J. (2011). Fast, scalable generation of high-quality protein multiple sequence alignments using Clustal Omega. Molecular systems biology *7*, 539.

Stansfeld, P.J., Hopkinson, R., Ashcroft, F.M., and Sansom, M.S. (2009). PIP2-binding site in Kir channels: definition by multiscale biomolecular simulations. Biochemistry *48*, 10926-10933.

Sun, J., and MacKinnon, R. (2020). Structural basis of human KCNQ1 modulation and gating. Cell *180*, 340-347. e349.

Tao, E., and Corry, B. (2022). Characterizing fenestration size in sodium channel subtypes and their accessibility to inhibitors. Biophysical Journal *121*, 193-206.

Venance, S.L., Cannon, S., Fialho, D., Fontaine, B., Hanna, M., Ptacek, L., Tristani-Firouzi, M., Tawil, R., and Griggs, R. (2006). The primary periodic paralyses: diagnosis, pathogenesis and treatment. Brain *129*, 8-17.

Wassenaar, T.A., Ingólfsson, H.I., Bockmann, R.A., Tieleman, D.P., and Marrink, S.J. (2015). Computational lipidomics with insane: a versatile tool for generating custom membranes for molecular simulations. Journal of chemical theory and computation *11*, 2144-2155.

Waterhouse, A.M., Procter, J.B., Martin, D.M., Clamp, M., and Barton, G.J. (2009). Jalview Version 2—a multiple sequence alignment editor and analysis workbench. Bioinformatics *25*, 1189-1191.

West, J.W., Patton, D.E., Scheuer, T., Wang, Y., Goldin, A.L., and Catterall, W.A. (1992). A cluster of hydrophobic amino acid residues required for fast Na (+)-channel inactivation. Proceedings of the National Academy of Sciences *89*, 10910-10914.

Wisedchaisri, G., Tonggu, L., El-Din, T.M.G., McCord, E., Zheng, N., and Catterall, W.A. (2021). Structural
basis for high-affinity trapping of the NaV1. 7 channel in its resting state by tarantula toxin. Molecular
cell *81*, 38-48. e34.
Wu, E.L., Cheng, X., Jo, S., Rui, H., Song, K.C., Dávila-Contreras, E.M., Qi, Y., Lee, J., Monje-Galvan, V., and
Venable, R.M. (2014). CHARMM-GUI membrane builder toward realistic biological membrane
simulations. Wiley Online Library.
Yan, Z., Zhou, Q., Wang, L., Wu, J., Zhao, Y., Huang, G., Peng, W., Shen, H., Lei, J., and Yan, N. (2017).
Structure of the Nav1. 4-β1 complex from electric eel. Cell *170*, 470-482. e411.
Yu, F.H., and Catterall, W.A. (2003). Overview of the voltage-gated sodium channel family. Genome
biology *4*, 1-7.
Zaman, T., Helbig, I., Božović, I.B., DeBrosse, S.D., Bergqvist, A.C., Wallis, K., Medne, L., Maver, A.,
Peterlin, B., and Helbig, K.L. (2018). Mutations in SCN3A cause early infantile epileptic encephalopathy.
Annals of neurology *83*, 703-717.

640