Short title: Probing leaf plastid volumes in situ

Correspondence:

Hans-Henning Kunz

Department for Plant Biochemistry LMU Munich,

Großhadernerstr. 2-4,

82152 Planegg-Martinsried, Germany

Email: kunz@lmu.de

Probing the *in-situ* volumes of Arabidopsis leaf plastids using 3D confocal and scanning electron microscopy.

Jan Knoblauch[a], Rainer Waadt[c], Asaph B. Cousins[a], Hans-Henning Kunz[a,b]*

[a] School of Biological Sciences, Washington State University, PO Box 644236, Pullman, WA 99164-4236, USA

[b] LMU Munich, Plant Biochemistry, Großhadernerstr. 2-4, 82152 Planegg-Martinsried, Germany

[c] Institute of Plant Biology and Biotechnology, University of Münster, Schlossplatz 7, 48149 Münster, Germany

*Corresponding author: Hans-Henning Kunz, kunz@lmu.de

The author responsible for distribution of materials integral to the findings presented in this article in accordance with the policy described in the Instructions for Authors (https://academic.oup.com/plphys/pages/General-Instructions) is Hans-Henning Kunz.

Keywords:

One-sentence summary

This work describes and compares three different strategies to obtain accurate volumes of leaf plastids from Arabidopsis, the most widely used model plant.

Footnotes:

Author Contributions

- H.-H.K. designed research. J.K. performed most experiments and analyzed data. R.W. designed and cloned the envelope fluorescence marker construct. A.B.C. analyzed and discussed data. All authors wrote and assisted in editing the manuscript.

Funding

- J.K.'s work was funded by an NSF Career Award (IOS-1553506) and a third call ERA-CAPS grant from NSF (IOS-1847382) to H.-H.K. R.W. received funding from the Deutsche

Forschungsgemeinschaft (DFG; WA 3768/4-1). H.-H.K. is funded by the Deutsche Forschungsgemeinschaft (DFG) (SFB-TR 175, project B09).

Abstract

Leaf plastids harbor a plethora of biochemical reactions including photosynthesis, the most important pathway on earth. Scientists are eager to unveil the physiological processes within the organelle but also their interconnection with the rest of the plant cell. An increasingly important feature of this venture is to use experimental data in the design of metabolic models. A remaining obstacle has been the limited in situ volume information of plastids and other cell organelles. To fill this gap for chloroplasts, we established three microscopy protocols delivering *in situ* volumes based on: 1) chlorophyll florescence emerging from the thylakoid membrane, 2) a CFP marker embedded in the envelope, and 3) calculations from serial block-face scanning electron microscopy (SBFSEM). The obtained data were corroborated by comparing wild-type data with two mutant lines affected in the plastid division machinery known to produce small and large mesophyll chloroplasts, respectively. Furthermore, we also determined the volume of the much smaller guard cell plastids. Interestingly, their volume is not governed by the same components of the division machinery which defines mesophyll plastid size. Based on our three approaches the average volume of a mature Col-0 wild-type mesophyll chloroplasts is 93 μm^3. Wild-type guard cell plastids are approximately 18 μm^3. Lastly, our comparative analysis shows that the chlorophyll florescence analysis can accurately determine chloroplast volumes, providing an important tool to research groups without access to transgenic marker lines expressing genetically encoded fluorescence proteins or high-end costly SBFSEM equipment.

Introduction

Photosynthesis, the light-driven CO_2 fixating pathway, is the foundation of life and global food production. In land plants, this pathway is housed in the chloroplast, a specialized plastid-type of endosymbiotic origin located in mesophyll or bundle sheath leaf cells. Chloroplasts are only one of several plastid-types found in the various diverse plant tissues (Choi et al., 2021), with the proplastid representing the most basic undifferentiated precursor organelle (Jarvis and López-Juez, 2013). Through highly coordinated gene expression involving the nuclear and the organellar genome, proplastids develop into chloroplasts, amyloplasts, chromoplasts etc., all with distinct morphologies and varying sizes (Liebers et al., 2017; Sun et al., 2017). Depending on plant age some plastid types can interconvert (Jarvis and López-Juez, 2013). The number of plastids can surpass 100 per cell in *Arabidopsis thaliana*, contributing to about a quarter of the total cell volume (Crumpton-Taylor et al., 2012; Unal et al., 2020). The abundance of genetically identical plastids occurs through binary fission facilitated by a complex contractile FtsZ ring inside the organelle and additional plastid-dividing (PD) rings that contain proteins anchored or associated with the inner and outer envelope membrane (Yoshida, 2018; Osteryoung and Pyke, 2014; Chen et al., 2018). The discovery of this intricate machinery was investigated through several loss and gain-of-function mutants of FtsZ and PD ring components. These mutants represent invaluable research tools to understanding both organelle fission and the significance of plastid abundance and size on basic physiological responses such as light stress avoidance (Dutta et al., 2017).

Plastids carry out general and highly specialized biochemical reactions, many yielding phytohormones or their respective precursors, which are critical for plant development and stress response (Bittner et al., 2022). Unsurprisingly, understanding plastid and chloroplast function has been the focus of many scientists interested in a wide range of topics, from photosynthesis to the

importance of plastids for plant environmental interactions (Kleine et al., 2021). In recent years,

computational modeling of energy/metabolic flux has given new insights into the complex inner

workings of the organelle (Krantz et al., 2021; Fürtauer et al., 2018). This modeling has been added

by non-aqueous fractionation to determine how these organelles interact (Klie et al., 2011;

Fürtauer et al., 2016; Höhner et al., 2021). Nevertheless, the efficacy of these models would be

further improved with precise determination of organellar volumes. This is especially important

for *A. thaliana*, arguably the most studied plant worldwide and the primary model plant system for

elucidating the molecular, structural, and biochemical control of energy/metabolic fluxes

(Woodward and Bartel, 2018).

Chloroplast dimensions and volumes are most often inferred using two-dimensional (2D)

imaging techniques (Kunz et al., 2014; Aranda-Sicilia et al., 2016; Unal et al., 2020). Transmission

electron microscopy (TEM) has been the primary method for obtaining these 2D images. However,

TEM requires fixation which can result in tissue, cellular, and organellar shrinkage. For instance,

spinach chloroplasts loose about 30% of their volume during the fixation procedure (Winter et al.,

1994). Also, TEM imaging is error-prone since optimal imaging quality requires 60-80 nm thick

sections and it is impossible to know what plane of the chloroplast is visible or the angle of the

section. This means it is unclear if a given chloroplast image is a glancing section or cuts through

the center, making accurate volume calculations challenging with a bias towards underestimating

volumes. This uncertainty leads to considerable variation in estimated chloroplast volumes and

requires large time-consuming datasets to approximate accurate chloroplast volumes even within

a single cell. A recent study using wheat and chickpea demonstrated that the 2D approach of

estimating chloroplast volumes is inaccurate and prone to volume underestimations (Harwood et

al., 2019).

104 The recent application of technologies to create three-dimensional (3D) representations of

105 leaf anatomy, including the serial block-face scanning electron microscopy (SBFSEM), has

106 introduced alternative ways to address ongoing uncertainty in chloroplast volumes (Denk and

107 Horstmann, 2004). In short, SBFSEM employs automated collection of serial surface images from

108 a resin-embedded sample block. This occurs via an internal ultramicrotome that cuts a 40-80 nm

109 thin section. The newly exposed surface of the truncated sample block is scanned to generate the

110 next SEM image. However, the SBFSEM technology requires significant specialized

111 instrumentation, and similar with TEM has potential difficulties with sample fixation and

112 preparation that may result in inferior images and data misinterpretation.

113 Live imaging of leaf tissue using confocal microscopy can avoid the errors associated with

114 fixation, such as chloroplast shrinkage. Confocal microscopy does, however, present its own

115 challenges and limitations. While there is no risk of chloroplast deformation due to fixation, the

116 relatively long wavelength of light drastically lowers the achievable imaging resolution compared

117 to electron microscopy. Additionally, since most confocal microscopes utilize a pinhole to image

118 optical sections inside the sample tissue, the fluorescence is scattered by the tissue it passes through

119 before reaching the objective, lowering resolution and limiting the depth accurate imaging can be

120 done to. As long as this is taken into consideration, however, confocal can be powerful tool as it

121 also allows for colocalizing several fluorophores within the sample, allowing for easy visualization

122 of one or more structures of interest.

123 In this study, we use three different 3D imaging techniques to measure leaf chloroplast

124 volumes in *A. thaliana* (Figure 1A). Two protocols employ confocal microscopy z-stacks using

125 chlorophyll fluorescence or a chloroplast envelope marker, respectively, as easy to replicate, more

126 accessible methods. The third approach is based on SBFSEM and 3D reconstruction. To validate

127   our assays, we used three different *A. thaliana* genotypes: 1) Col-0 as a wild-type control, 2) *35s-*

128   *PDV1 35s-PDV2*, which has more, smaller chloroplasts, and 3) *arc5-2*, having fewer but gigantic

129   chloroplasts per cell (Osteryoung, 2017). The *35s-PDV1 35s-PDV2* and *arc5-2* are chloroplast

130   division mutants and were chosen to quantify the accuracy of each volume determination

131   approach.

Results & Discussion

Mesophyll chloroplast volume measurements using chlorophyll fluorescence

For plastid volume calculations three different, wild type and previously described genotypes were

selected. *35s-PDV1 35s-PDV2* has smaller chloroplasts caused by the overexpression of the outer

envelope PLASTID DIVISION1 (*PDV1*) and PLASTID DIVISION2 (*PDV2*) proteins, which

recruit the ARC5/DRB5P ring during chloroplast division (Okazaki et al., 2009; Dutta et al., 2017).

Conversely, *arc5-2* is a T-DNA insertion loss of function mutant that exhibits a low number of

gigantic chloroplasts per cell. The *ARC5* locus encodes one of the outer envelope membrane

proteins responsible for assembling the most outer PD ring (Robertson et al., 1996; Miyagishima

et al., 2006). Col-0 was used as a wildtype control. All genotypes exhibited a similar green leaf

color and were indistinguishable from controls with regards to their growth rate and appearance

(Figure 1B). The reported chloroplast phenotypes became visible in the micrographs (Figure 1C-

D). For all experiments, leaf discs were collected from the first three mature true leaves. Three

separate grow-outs per genotype were utilized to test the consistency of our results. Initially,

chloroplast volumes were calculated based on confocal microscopy z-stacks of chlorophyll

fluorescence (supplemental movies 1-3), which emerges from the thylakoid membrane (Figure

1E). Across all three grow-outs, Col-0 mesophyll chloroplasts had an average volume of 88.24 ±

1.58 μm^3. *35s-PDV1 35s-PDV2* showed a clear trend towards slightly smaller mesophyll

chloroplast volumes with an average volume of 60.13 ± 1.05 μm³. Lastly, *arc5-2* chloroplast

volumes were significantly greater (one-way ANOVA, *p*<0.05) as in wild-type and *35s-PDV1 35s-*

PDV2 plants, averaging at 1538 ± 145 μm³. Frequency distribution plots of the combined volume

data collected on all three genotypes are shown in Supplemental Figure 1A-B. Comparing different

grow-outs, minor differences in chloroplast volumes can be observed within each genotype. For

all three genotypes, the second grow-out season gave rise to slightly lower average chloroplast

volumes than the first and third season indicating minor seasonal effects. However, statistical

analysis within each genotype did not indicate significant differences between grow-outs (one-

way ANOVA, *p*<0.05).

On average, Col-0 contained 74 ± 3 chloroplasts per mesophyll cell (Figure 1F). This is in

the middle range compared to reports by others suggesting averages of 60, 76 (± 5) or between 80

and 120 chloroplasts per cell (Kinsman and Pyke, 1998; Okazaki et al., 2009; Crumpton-Taylor et

al., 2012). In line with the literature, *35s-PDV1 35s-PDV2* and *arc5-2* mesophyll cells contained

on average 92 ± 5 and 5 ± 0.5 chloroplasts, respectively (Miyagishima et al., 2006; Okazaki et al.,

2009). The differences between studies can be related to different mesophyll cell types,

developmental states, or to local growth conditions which were all reported to affect plastid

numbers (Antal et al., 2013).

Guard cell plastid volume measurements using chlorophyll fluorescence

Guard cell chloroplasts are known to be smaller than mesophyll chloroplasts (Pyke and Leech,

1994). Thus, we also assayed guard cell plastid volumes to test the feasibility of our protocol across

different cell types. When comparing guard cell chloroplast volumes, no differences between

genotypes were observed. For Col-0 chloroplast volumes averaged at 17.69 ± 0.21 μm³, while

volumes for *35s-PDV1 35s-PDV2*, and *arc5-2* were 18.04 ± 0.26 μm³, and 17.26 ± 0.28 μm³

respectively (Figure 1G). Comparing the genotypes by one-way ANOVA showed no statistically

significant difference ($p<0.05$). Supplemental figure 2 shows an overlapping frequency

distribution for the three genotypes' guard cell plastid volumes. When comparing the three grow-

outs individually the same pattern can be seen. In summary, our data are in line with previous

reports on *arc5* mutants showing that loss of *ARC5* affects mesophyll but not guard plastid sizes

(Pyke and Leech, 1994).

As shown above, guard cell chloroplast volumes are similar between Col-0 and the two

mutants affected in mesophyll chloroplast division. Similarly, the number of plastids per guard

cell was not significantly different (Figure 1H): Col-0 guard cells contained 4.5 ± 1.0 chloroplasts

on average, which aligns with previous research done on this (Fujiwara et al., 2019), while both

35s-PDV1 35s-PDV2 and *arc5-2* contained only 3.6 ± 0.83 chloroplasts on average. Some

previous studies reported immature, non-fluorescing plastids in *arc5-2* guard cells (Fujiwara et al.,

2018). Since no brightfield images were acquired in this study, we cannot comment on this

observation as we only visualized fluorescing plastids. Also, giant chloroplasts were previously

found in rare instances in *arc5*-2 guard cells (Fujiwara et al., 2018)*, which we did not encounter.

Mesophyll chloroplast volume measurements using a plastid outer envelope fluorescence

marker protein

Chlorophyll fluorescence emerges from the thylakoid membrane pigments and thus may not

effectively represent the entire volume of the chloroplast. Therefore, a cyan-fluorescence protein

(CFP) envelope marker localized to the outer chloroplast envelope was employed. Initially, a Col-

0 wild-type line with robust CFP signal was isolated using confocal microscopy (Figure 2A-C).

Since strong overexpression of envelope proteins can result in membrane alterations, we ensured

that this did not occur in the lines selected for this study (Breuers et al., 2012). Subsequently, the

marker gene was introgressed into the *35s-PDV1 35s-PDV2* and *arc5-2* mutants with known

plastid volume differences. Once homozygous F3 plants were available, we acquired z-stacks of

both, the envelope CFP marker and chlorophyll fluorescence signals in parallel from all three

genotypes. As expected, average chloroplast volumes calculated using the envelope marker was

slightly higher than that calculated using chlorophyll fluorescence (Figure 2D). For Col-0 the

average volume for chlorophyll fluorescence and the envelope marker were 78.9 $\mu m^3 \pm 2.6$ and

94.7 ± 2.9 μm^3 respectively, for *35s-PDV1 35s-PDV2* they were 55.6 $\mu m^3 \pm 2.4$ and 63.1 ± 2.0

μm^3 respectively, and for *arc5-2* the values were 1714.9 $\mu m^3 \pm 222.7$ and 2232.0 ± 330.0 μm^3,

respectively. Statistical analyses comparing chlorophyll fluorescence and envelope marker

volumes within each genotype, did not show a significant difference in values ($p<0.05$), albeit a

trend was clear. When comparing different genotypes, the same observation emerged i.e.,

statistically significant higher volumes exist only between *arc5-2* and the two other lines. Col-0

volumes based on chlorophyll fluorescence reflect to the lower end found in the three, independent

grow-out experiment previously (Figure 1G). Seasonal effects are the most likely explanation.

Chloroplast volume determination using serial block-face scanning electron microscopy

When it comes to the application in plant tissues, serial block-face scanning electron microscopy

(SBFSEM) is still a relatively new imaging technique. In fact, for *A. thaliana* mesophyll cell

organelles there are no published volume data yet. Thus, we performed SBFSEM as an orthogonal,

more high-resolution method for chloroplast analysis. Imaging was done for the same three

genotypes as described above. Since the tissue has to be fixed, dehydrated and embedded, osmotic

alterations may have an influence on organelle volume. The protocol used here and described in

the materials and methods section was developed as part of a study to improve SBFSEM protocols

and analysis in plant tissues (Mullendore et al. in preparation). This includes a machine learning

algorithm "ANATOMICS MLT" which was used for this study to auto-label large image data sets.

Figure 3A shows chloroplasts labeled by ANATOMICS MLT. The average chloroplast volume

determined for Col-0 was 92.9 ± 1.3 μm^3. For *35s-PDV1 35s-PDV2*, and *arc5-2* the volumes were

69.4 ± 0.8 μm^3, and 1970 ± 759 μm^3, respectively (Figure 3B-C). As for all previous imaging

techniques used, only trends in chloroplast volumes difference could be identified between *35s-*

PDV1 35s-PDV2 and Col-0, while both genotypes were significantly different from *arc5-2*. A full

image stack reconstruction for each genotype can be found as supplemental movies 4-6.

Protocol comparisons and recommendations

In this study, we set out to gather accurate volume information from mesophyll and guard cell

plastids of the model plant *A. thaliana*. In addition, we wanted to test the feasibility of employing

standard confocal microscopy, available in most biology departments, to determine organellar

volumes. When comparing between imaging approaches, chloroplast volumes were relatively

consistent (about 93 μm^3 in wild-type plants), with volumes derived by an outer envelope marker

protein showing a slightly higher average between 10-20% when compared to both chlorophyll

fluorescence and SBFSEM measurements (Figure 4A-C, Supplemental Table 1). However, when

running a one-way ANOVA comparing chloroplast volumes measured by each of the three

imaging techniques, no statistically significant difference ($p<0.05$) was found within either of the

genotypes, regardless of the imaging approach applied. Slightly higher average volumes using data

from the envelope marker are to be expected, as it defines the outer boundary of a chloroplast and

thus encompasses envelope membranes, inter membrane space, and the stroma. In contrast,

chlorophyll fluorescence primarily represents the volume determined by thylakoid membranes.

One limitation we observed when subtracting the chlorophyll fluorescence from the outer

envelope-based volume is that the obtained values are too low (≈ 16 μm^3) to accurately reflect the

stromal volume. According to the literature the stroma is expected to occupy about 50% of a plastid

(Antal et al., 2013). This shortcoming can be explained by the insufficient resolution of confocal

micrographs which does not allow to accurately resolve the stroma between thylakoid membranes.

The critical advantage of SBFSEM is the much higher resolution compared to confocal or

super-resolution microscopy, but much faster acquisition time than focused ion beam

methodologies. Therefore, it allows to image large tissue areas with hundreds or thousands of cells

within reasonable time (days) while allowing to acquire surface areas and volumes of organelles

that are too small to image by light microscopy-based methods (e.g., confocal) such as the ER or

Golgi apparatus. It is however critical that appropriate protocols are used to maintain the volume

of organelles. Chloroplasts are large enough to serve as a tool to compare volumes taken by

confocal microscopy and SBFSEM. It is reasonable to assume that if chloroplast volumes between

in situ and embedded tissue are similar, other organelle volumes in the embedded tissue will be

accurate as well.

The difference in chloroplast volumes between chlorophyll fluorescence and SBFSEM

images is surprisingly small and not statistically significant, showing that our protocol provides

excellent maintenance of the tissue. All chloroplasts in our SBFSEM micrographs appear intact

(no sharp edges which are usually an indication of shrinkage) and undamaged. Nevertheless,

fixative and buffers concentrations as well as the dehydration procedure may have to be tailored

to other specimen types such as stems and roots or other plant species. Stains can also be adjusted

to fit organelles or structures of interest. Usually this requires a fair amount of strategic trial and

error. However, SBFSEM currently allows standard resolutions of down the 10 nm at comparably

high acquisition speeds which is a massive improvement over other available technology. As we

show in this study, accurate quantitative anatomical data comparable to in situ studies can be

achieved.

The chloroplast division mutants used as a proof of concept in this study demonstrates the

ability and limitations to quantify volume differences. While each method confirmed the drastic

mesophyll plastid size variation between Col-0 and *arc5-2* plants, the expectedly smaller

differences between Col-0 and *35s-PDV1 35s-PDV2* were only revealed by trend but not with

sufficient statical power to allow strong conclusions regardless of the imaging applied. Volume

differences between mesophyll and guard cell chloroplasts were also successfully resolved.

Interestingly, while mesophyll chloroplast volumes varied among genotypes, guard cell plastids

had consistent volumes and numbers throughout. This provides more evidence that guard cell and

mesophyll plastid division are governed by distinct genetic programs.

When comparing our results with the literature we find quite stark differences. While we

determined the Col-0 wild-type mesophyll chloroplast using orthogonal approaches at a size of $\approx$

93 μm^3 most publications refer to old volume studies on other species and an average value of $\approx$

31 μm^3 (Antal et al., 2013; Nobel, 2020). Although mesophyll chloroplast sizes seem highly

species- and to a minor degree daytime-dependent the values from spinach, pea, tobacco, wheat,

poplar are in the range from 15 to 35 μm^3 and appear to be rather underestimations (Antal et al.,

2013; Nobel, 2020). At least side by side in a light microscope *A. thaliana* and pea chloroplast do

not seem to drastically differ in size (Schulz et al., 2004). Recent 3D assays describe wheat

chloroplasts at 114.6 $\pm$ 21.5, chickpea at 22.4 $\pm$ 10.2 μm^3, and rice at 47 μm^3 (Oi et al., 2017;

Harwood et al., 2020). For *A. thaliana* only one study on cotyledon development was found to

report 3D derived plastid volume data (Pipitone et al., 2021). Albeit only three plastids were

assayed by SBF-SEM the obtained value of 112.14 μm^3 ($\pm$4.3) is roughly in the same range as the

93 μm^3 we recorded from mesophyll chloroplasts in mature true leaves. In conjunction, both

studies emphasize that *A. thaliana* chloroplasts are far bigger than the often-suggested average

plastid volume of 31 μm^3. This needs to be taken into account when organelle volumes are

employed in flux models. Additionally, because of the wide range of plastids sizes reported from

different species generalizing metabolic flux assumptions can be problematic. Fortunately, as we

show through our study rapid z-stack recordings based on chlorophyll fluorescence using a

standard confocal microscope gives sufficiently accurate volume information to survey this data

point for any given plant species of interest. Since we obtained very similar values regardless of

the approach employed, we can state with high confidence that a volume of $\approx$ 93 μm^3 reflects the

natural in situ situation in the model plant Arabidopsis. Moreover, the fixation-staining protocol

we utilized for SBFSEM is well-suited for plastids. The low deviation from values obtained via

fixation-staining-free confocal microscopy confirms that no shrinkage occurred in our specimens.

Conclusions

Overall, comparing chloroplast volumes from plastid division mutants acquired through different

imaging methods shows relatively close overlap of volumes. This suggests that even though z-

stack confocal micrographs have much lower resolution than SBFSEM, they still give fairly

accurate volume data. Between the two confocal microscopy methods, using an envelope marker

will yield more accurate total chloroplast volumes. Nevertheless, for a much easier and faster

estimation, chlorophyll fluorescence delivers high replicate numbers and is thus quite accurate.

Moreover, it does not require cloning, transformation or introgression of a fluorescence protein

encoding transgene. SBFSEM yields the most accurate volume data due to its much higher

resolution. However, lengthy sample preparation procedure optimization might be necessary to

prevent artifacts resulting in skewed volumes. Researchers are advised to consider our conclusions

for balanced and informed decision-making which answer the question at hand with the best

equipment available to them.

Materials and Methods

Plant Material

A. thaliana plants used: Columbia-0 as a control, *arc5-2* (SAIL_71_D11), a giant chloroplast

mutant due to a knockout of the *ARC5* chloroplast division gene, and 35*S-PDV1* 35*S-PDV2* with

smaller chloroplasts caused by overexpression of *PDV1* and *PDV2* (Miyagishima et al., 2006;

Okazaki et al., 2009; Dutta et al., 2017). These plants were grown in a Conviron growth chamber

(Winnipeg, Man, Canada) using a 16/8-hour light/dark cycle at 150 µmol photons $m^{-2}s^{-1}$,

21°C/19°C day/night cycle and 60-80% humidity on soil. Approximately, 100 seeds per genotype

were sown into one 1.15-quart pot with soil (Sungro Professional Growing Mix #1, Sun Gro

Horticulture, Agawam, MA, USA), and after a week, the ten largest seedlings were transferred

into individual 0.59-pint pots and grown for 4 additional weeks before imaging.

To target a fluorescent marker to the outer chloroplast envelope surface, the coding sequence (192

bp) of the outer envelope protein 7.1 (OEP7; At3g52420) without its Stop codon and with a Gly-

Gly-Ser-Gly-linker at the 3'-end was first amplified using Q5 High-Fidelity DNA Polymerase

(New England Biolabs) and the primers GGB_OEP7_F –

AACAGGTCTCAAACAATGGGAAAAACTTCGGGAGC and GGC_OEP7_GGSG_R –

AACAGGTCTCTAGCCTCCAGATCCTCCCAAACCCTCTTTGGATGTGG, followed by

FastDigest Eco31I (Thermo Fisher Scientific) digestion, and ligation into the GreenGate module

backbone pGGB (Lamproupolos et al., 2013). Together with previously described GreenGate

modules (Lamproupolos et al., 2013; Waadt et al., 2017), OEP7, the orange fluorescing

mNectarine (Johnson et al., 2009) without Start codon, and the cyan fluorescing mTurquoise

(Goedhart et al., 2010) with a 5'-end Gly-Ser-linker were ligated into the plant expression vector

pGGZ003 yielding the plasmid pGGZ-RW105 (pGGZ003-pUBQ10-OEP7-GGSG-

mNectarine_ATG-GSL-mTurquoise-tHSP18.2M-hygR). This construct was then transformed

into Col-0 wild-type by floral dip, and positive transformants were selected through germination

on hygromycin (15 μg ml^{-1}, Chem-Impex, Wood Dale, IL, USA). Subsequently, the envelope

marker was introgressed into *arc5-2* and *35s-PDV1 35s-PDV2* mutant lines.

Confocal Laser Scanning Microscopy

Mesophyll Chloroplast Fluorescence Volume Measurements

For mesophyll imaging, fresh leaf disks of 1 cm diameter were taken from the center of a mature

leaf and imaged with a HC PL APO CS2 63x/1.20 NA water-immersion objective on a Leica SP8

confocal laser-scanning microscope (Leica Biosystems, Deer Park, IL, USA). Chlorophyll was

excited using a 405 nm pulsed laser and the fluorescence was collected using a HyD detector set

to 650-720 nm. The gain was optimized for each stack, using the brightness indicator. Additional

imaging parameters were: scanning speed 100 Hz, zoom 2, line average 2, pinhole 1, and z-stacks

system optimized with 0.305-μm-thick optical sections for volume analysis. Number of z-sections

varied, and stacks were later cropped to exclude images where the fluorescence border could not

be accurately identified.

Guard Cell Chloroplast Volume Measurements

For imaging of guard cell chloroplasts, the abaxial side of the leaf was used as there are more

stomata present. The abaxial side of a leaf was glued to a glass slide using medical adhesive

(Hollister, Libertyville, IL, USA). After allowing the glue to dry, leaf tissue was removed by

scraping with a razor blade until the lower epidermis became exposed (Azoulay-Shemer et al.,
2016). Guard cells turgescence was determined visually and used to checked for intactness, and
guard cell chloroplasts were then imaged using a HC PL APO CS2 63x/1.40 oil-immersion
objective. Z-stacks were taken using 0.255-µm-thick optical sections. Otherwise, images were
taken as described above.

mTurquoise Imaging

Plants with chloroplast envelopes tagged with mTurquoise were also imaged using a HC PL APO
CS2 63x/1.20 NA water-immersion objective on a Leica SP8 confocal laser-scanning microscope
(Leica Biosystems, Deer Park, IL, US). Dual channels were used to image mTurquoise
simultaneously with chlorophyll fluorescence. A HyD detector set to 460-520 was used to collect
mTurquoise fluorescence, while a HyD detector set to 650-720 nm was used for chlorophyll
fluorescence. All other imaging parameters were as described above.

Chloroplast Volume Analysis

Chloroplast reconstruction and analysis was done using ImageJ software, utilizing the Bio-Formats
plugin to import .lif files (https://github.com/ome/bioformats). After importing a z-stack, and
manually thresholding to optimize chloroplast outlines, a Median Filter (Radius: 2.0) was applied
for smoothing and subtract (~10), a global brightness reduction tool, was used to remove
background noise. Chloroplast volumes were extracted with the 3D Objects Counter function in
ImageJ, and the resulting labels were checked to ensure only volumes of complete chloroplasts
were counted.

Serial Block Face Scanning Electron Microscopy (SBFSEM)

Fixation Protocol

Arabidopsis leaves were removed from the plant, cut into 2 mm x 2 mm squares, and put into a fixative solution containing 4% glutaraldehyde, 2 mM $CaCl_2$ in 0.1 M cacodylate buffer (pH 6.8) for 6 hours at room temperature. Samples were then microwaved at 300W at a 35°C maximum temperature limit for 2 min and then washed 3x for 10 minutes in 0.1 M cacodylate buffer followed by a post-fixation with 1.5% KFeCN, 2% OsO_4, 2 mM $CaCl_2$, in 0.15 M cacodylate buffer overnight at 4°C. After washing 3x for 10 min in double distilled water (ddH_2O) at room temperature, samples were incubated in 0.2% gallic acid for 1 hour at room temperature, and then washed 3x for 10 minutes in ddH_2O. A secondary post-fixation was done in 2% OsO_4 for 3 hours at room temperature, and after washing 3x for 10 min in ddH_2O, a 2% uranyl acetate incubation was applied overnight at 4°C. Samples were washed 3x for 10 minutes in ddH_2O at room temperature, then stained with Walton's lead aspartate (Walton, 1979) at 60°C for one hour and finally washed 3 times for 10 min in ddH_2O at room temperature.

Samples were dehydrated with an acetone series using freshly mixed acetone solutions. 10% steps were done between 10%-50% (v/v) acetone for 10 min at room temperature. A second exchange of 50% acetone was incubated at -20°C for 1 hour. For the remainder of the dehydration series, exchanges were performed at -20°C overnight in 60%, 70% ,80% ,90% ,100% ,100%, 100% (v/v) acetone. After the third 100% acetone treatment, samples were incubated overnight and then moved to room temperature to acclimate for ~30 min before the final dehydration in 100% acetone two times for 10 minutes at room temperature.

Samples were infiltrated in Spurr's resin (Sigma Aldrich, St. Louis, MO, USA) in 3:1, 2:1, 1:1, 1:2, 1:3, 100% acetone:hard Spurr resin without the hardener of the resin (DMAE) to prevent

premature polymerization, each overnight at room temperature on a rotator. This was followed by

two exchanges of 100% hard Spurr's resin with DMAE, overnight at room temperature on a

rotator, with the lids removed. Finally, samples were microwaved at 100W and 40°C for 1 hour

and subsequently put into a 70°C oven to polymerize.

Imaging

To prepare samples for SBFSEM, the embedded tissue was trimmed and sectioned using a Leica

EM UC7 ultramicrotome. Ultrathin sections (~70 nm) were taken and checked for quality using a

FEI Technai G2 20 Twin (Thermo Fisher, Waltham, MA, USA). Samples were then transferred to

a SBFSEM stub, which is the sample holder, trimmed on the ultramicrotome using a glass knife,

after which a Technics Hummer V Sputter Coater was used to apply 10 nm of gold coating. For

imaging, an Apreo VolumeScope SEM (Thermo Fisher, Waltham, MA,USA) with the VS-DBS:

LoVac lens-mounted BSED detector was used. Imaging conditions were set to 2 kV accelerating

voltage, 50 Pa chamber pressure, a beam current of 0.10 nA, a pixel size of 20 x 20 nm, and a

dwell time of 3 μs.

Image Processing

The SBFSEM image stacks were processed for volume analysis using software specifically

designed to work with SBFSEM stacks, Amira (Thermo Fisher, Waltham, MA, USA). Partial

stacks (~500 images) were imported into Amira, and a gaussian filter (XY, standard deviation 1,

1, kernel size factor 2) was used to remove noise. Images are then inverted, resampled (Lanczos

Filter, Voxel Size: x= 40, y= 40, z= 40), and then exported as a sequence of 2D tiff files. This

process was repeated until the entire SBFSEM stack had been processed and exported as described

above. The complete stack of processed images was then imported into Amira, cropped around the

area of interest and contrast matched (XY planes, mean & variance) using the best contrasted

image within the stack as a reference. This made the contrast consistent for all images throughout

the SBFSEM stack. Auto align slices (Rigid, Align and Resample) was run to automatically align

all SBFSEM slices. Images were cropped again to remove overlapping edges created by the

alignment, and a median filter (XY planes, 3 iterations, Iterative) was used to remove remaining

noise. Finally, the stack was exported as a set of 2D tiff files. The fully processed stack was then

imported into ImageJ as an image sequence, manually contrasted, and saved as a tiff, creating a

3D tiff stack.

Volume Analysis

Volume analysis of SBFSEM stacks was done using Anatomics MLT

(https://github.com/ajbrookhouse/WSU_PlantBio_ML), a machine learning algorithm designed to

analyze volumes, object counts, and surface areas of features within SBFSEM stacks. Amira was

used to label all chloroplasts within a sequence of 10 processed images with the Segmentation tool

by using the brush, then images and labels were exported as separate 3D tiff files. These stacks

were then used in the train function of Anatomics MLT (Training Config: Instance.yaml, 100,000

iterations) to train the program to label chloroplasts. The model was then used in the Auto-Label

section to label full image stacks. Finally, the Output Tools tab was designed to analyze data from

the created labels and can be used to create mesh or point clouds to visually display the labels.

This was used to calculate chloroplast volumes and create Point Clouds of the labels which were

viewed and overlayed with original images to check for labeling accuracy using the Visualize tab.

Statistical Analysis

Graphing and statistical analysis was done using GraphPad software (Dotmatics, Boston, MA,

USA). Means as well as standard deviation and error were calculated. One-way ANOVAs were

used to determine statistical significance between biological replicates, genotypes, and imaging

techniques. A 2-way ANOVA was run to analyze volume data within the chloroplast fluorescence

dataset, comparing each growth cycle within a genotype to the volumes of chloroplasts between

genotypes. A 3-way ANOVA was also used to compare chloroplast volumes of genotypes within

each imaging method and to compare volumes of chloroplasts of each genotype between imaging

techniques.

Accession numbers

arc5-2 (SAIL_71_D11, S870785), *35s-PDV1 35s-PDV2, ARC5* (AT3G19720), *PDV1*

(AT5G53280), *PDV2* (AT2G16070), *OEP7.1* (At3g52420)

Supplemental Data

Supplemental Figure S1. A, Frequency distribution of all mesophyll chloroplast volumes between

35s-PDV1 35s-PDV2 and Col-0 chloroplast within mesophyll cells show a distribution of *35s-*

PDV1 35s-PDV2 volumes slightly lower than Col-0, not at a statistically significant level. B, a

frequency distribution for *arc5-2* shows that volumes below 1000 μm^3 are the most common,

however values reach as high as 7000 μm^3.

Supplemental Figure S2. Frequency distribution of guard cell chloroplast volumes shows very

similar distribution and overall volumes between all three genotypes.

Supplemental Table 1: Summary of chloroplast volumes in µm3 calculated using image stacks

from each method employed. For each genotype the average and standard error are presented.

Supplemental Movie 1-3: Video reconstructions showing confocal Z-stacks of each genotype, *35s-*

PDV1 35s-PDV2, Col-0 and *arc5-2*, respectively.

Supplemental Movie 4: Video reconstruction of scanning through a Col-0 SBFSEM image stack.

Chloroplasts are labeled using the machine learning algorithm and displayed in cyan.

Supplemental Movie 5: Video reconstruction of scanning through a *35s-PDV1 35s-PDV2*

SBFSEM image stack.

Supplemental Movie 6: Video reconstruction of scanning through an *arc5-2* SBFSEM image stack.

Acknowledgements

We thank Dr. Katherine W Osteryoung from Michigan State University for providing the original

mutant germplasms used in this project. J.K. received the WSU Franceschi training grant to

support the microscopy work. J.K. thanks his graduate committee members Drs. Andrew

McCubbin and Helmut Kirchhoff (both at WSU) for their support and insights.

Figure Legends

Figure 1: Confocal microscopy reveals changes in gain and loss of function mutants in

components of the plastid division machinery that are restricted to mesophyll cells. A, the

diagram of the Arabidopsis plant shows the size of the plants when imaging and sample preparation

was done. The * indicates the leaves used for imaging, and circles indicate the location on each

leaf. From here, these samples were either used immediately for imaging using confocal

microscopy or fixed and prepared for SBFSEM. **B**, No apparent changes in plant growth or

appearance were observed in long-day cultivated genotypes used in this study, *35s-PDV1 35s-*

PDV2, Col-0, *arc5-2* (scale bars = 2 cm). **C**, Confocal images of chlorophyll fluorescence for each

genotype show clear size differences in mesophyll chloroplasts (scale bars = 25 µm). **D**, Confocal

images of guard cell chloroplasts taken using chlorophyll fluorescence shows similar chloroplast

volumes within guard cells of the three genotypes investigated (scale bars = 25 μm). **E**,

Comparison of mesophyll chloroplast volumes deduced from z-stack of chlorophyll fluorescence

recordings. Individual grow-outs are shown for each genotype. Statistical analysis done by one-

way ANOVA with Tukey's multiple comparison test ($p<0.05$) showed no significant difference

between Col-0 and *35s-PDV1 35s-PDV2* chloroplast volumes, neither in the full dataset, nor

between any individual repetitions. Significant differences were observed between *arc5-2* and *35s-*

PDV1 35s-PDV2 but also between *arc5-2* and Col-0 mesophyll chloroplast volumes. Letters above

each sample in the graph indicate groups of significance (±SE). **F**, Chloroplast numbers per

mesophyll cell were counted and show an inverse relationship to that of chloroplast volumes

observed between the genotypes. *35s-PDV1 35s-PDV2* has the highest average number of

chloroplasts per cell with around 90 chloroplasts, followed by Col-0 averaging slightly fewer, at

around 70 chloroplasts per cell. Finally, *arc5-2* mutants showed by far the lowest chloroplast count

per cell with an average of 5 chloroplasts (±SE). **G**, Guard cell chloroplast volumes calculated

based on z-stacks of chlorophyll fluorescence recordings, in three separate grow outs, statistical

analysis shows no significant difference ($p<0.05$) between any of the genotypes, neither within the

whole dataset nor within individual growing cycles. Letters above each data set again show groups

of significance (±SE). **H**, Guard cell chloroplast counts show that, unlike the similarities in

volumes, a slight difference in average chloroplast number per guard cell can be observed. Col-0

guard cells contain on average 4.5 chloroplasts, while both the *35s-PDV1 35s-PDV2* and *arc5-2*

genotypes contain only around 3.5 chloroplasts per guard cell (±SE).

Figure 2: Comparing mesophyll chloroplast volumes deduced from chlorophyll fluorescence and an envelope membrane located fluorescent marker. Comparison of **A**, chlorophyll fluorescence, **B**, the outer chloroplast envelope labeled with CFP visualized in **C**, as an overlay in Col-0 (scale bar = 10 µm). **D**, comparison between chloroplast volumes extracted by chlorophyll fluorescence and mTurquoise envelope marker shows slightly higher volumes in each genotype when calculated using the mTurquoise marker. A statistically significant difference ($p<0.05$) in chloroplast volumes was only found between Col-0 and *35s-PDV1 35s-PDV2* when compared to *arc5-2* mutants. No significant difference is observed between imaging methods when looking at individual genotypes (±SE).

Figure 3: Chloroplast volume analysis using SBFSEM. A, Reconstructions of Col-0 chloroplasts by using SBFSEM. Green objects represent chloroplasts labeled using the machine learning algorithm, while in grey a surface was created by thresholding, representing mostly the cell wall and some other high contrast objects within the cells. **B**, deduced chloroplast volumes measured by SBFSEM imaging. Statistical significance ($p<0.05$) in chloroplast volumes is shown by numbers above each dataset in the graph. A significant difference in chloroplast volumes can only be seen between Col-0 and *arc5-2*, as well as between *35s-PDV1 35s-PDV2*, and *arc5-2*. **C**, A frequency distribution of Col-0 and *35s-PDV1 35s-PDV2* shows a shift in volume distribution between the two genotypes (±SE).

Figure 4: Comparison of chloroplast volumes measured by imaging type. A-C, comparison of mesophyll chloroplast volumes extracted using confocal imaging of chlorophyll fluorescence (Chl

a), CFP envelope marker, and SBFSEM. The means of the data collected using each imaging

method was taken for all genotypes. Each pair of imaging methods is compared by plotting the

means (±SE). An identity line was added to highlight differences in means between imaging types

in each graph. This shows overall similar volumes and distributions between the methods, with

data extracted using the CFP envelope marker showing the largest volumes and chlorophyll

fluorescence having the lowest. A one-way ANOVA done for statistical analysis only showing a

significant difference ($p<0.05$) when comparing *35s-PDV1 35s-PDV2* and Col-0 volumes acquired

by either method to *arc5-2*. No significant difference was found between imaging methods for any

genotype.

Conflict of interest statement. None declared.

References

Antal, T. K., Kovalenko, I. B., Rubin, A. B., & Tyystjärvi, E. (2013). Photosynthesis-related

quantities for education and modeling. *Photosynthesis Research*, *117*(1–3), 1–30.

https://doi.org/10.1007/s11120-013-9945-8

Aranda-Sicilia, M. N., Aboukila, A., Armbruster, U., Cagnac, O., Schumann, T., Kunz, H.-H.,

Jahns, P., Rodríguez-Rosales, M. P., Sze, H., & Venema, K. (2016). Envelope K$^+$/H$^+$

Antiporters AtKEA1 and AtKEA2 Function in Plastid Development. *Plant Physiology*, *172*(1),

441–449. https://doi.org/10.1104/pp.16.00995

Bittner, A., Cieśla, A., Gruden, K., Lukan, T., Mahmud, S., Teige, M., Vothknecht, U. C., &

Wurzinger, B. (2022). Organelles and phytohormones: A network of interactions in plant stress

responses. *Journal of Experimental Botany*, *73*(21), 7165–7181.

https://doi.org/10.1093/jxb/erac384

Breuers, F. K. H., Bräutigam, A., Geimer, S., Welzel, U. Y., Stefano, G., Renna, L., Brandizzi,

F., & Weber, A. P. M. (2012). Dynamic Remodeling of the Plastid Envelope Membranes – A

Tool for Chloroplast Envelope in vivo Localizations. *Frontiers in Plant Science*, *3*.

https://doi.org/10.3389/fpls.2012.00007

Chen, C., MacCready, J. S., Ducat, D. C., & Osteryoung, K. W. (2018). The Molecular

Machinery of Chloroplast Division. *Plant Physiology*, *176*(1), 138–151.

https://doi.org/10.1104/pp.17.01272

Choi, H., Yi, T., & Ha, S.-H. (2021). Diversity of Plastid Types and Their Interconversions.

Frontiers in Plant Science, *12*, 692024. https://doi.org/10.3389/fpls.2021.692024

Crumpton-Taylor, M., Grandison, S., Png, K. M. Y., Bushby, A. J., & Smith, A. M. (2012).

Control of Starch Granule Numbers in Arabidopsis Chloroplasts. *Plant Physiology*, *158*(2),

905–916. https://doi.org/10.1104/pp.111.186957

Denk, W., & Horstmann, H. (2004). Serial Block-Face Scanning Electron Microscopy to

Reconstruct Three-Dimensional Tissue Nanostructure. *PLoS Biology*, *2*(11), e329.

https://doi.org/10.1371/journal.pbio.0020329

Dutta, S., Cruz, J. A., Imran, S. M., Chen, J., Kramer, D. M., & Osteryoung, K. W. (2017).

Variations in chloroplast movement and chlorophyll fluorescence among chloroplast division

mutants under light stress. *Journal of Experimental Botany*, *68*(13), 3541–3555.

https://doi.org/10.1093/jxb/erx203

Fujiwara, M. T., Yasuzawa, M., Kojo, K. H., Niwa, Y., Abe, T., Yoshida, S., Nakano, T., & Itoh,

R. D. (2018). The Arabidopsis arc5 and arc6 mutations differentially affect plastid morphology

in pavement and guard cells in the leaf epidermis. *PLOS ONE*, *13*(2), e0192380.

https://doi.org/10.1371/journal.pone.0192380

Fürtauer, L., Weckwerth, W., & Nägele, T. (2016). A Benchtop Fractionation Procedure for

Subcellular Analysis of the Plant Metabolome. *Frontiers in Plant Science*, *7*.

https://doi.org/10.3389/fpls.2016.01912

Fürtauer, L., Weiszmann, J., Weckwerth, W., & Nägele, T. (2018). Mathematical Modeling

Approaches in Plant Metabolomics. In C. António (Ed.), *Plant Metabolomics* (Vol. 1778, pp.

329–347). Springer New York. https://doi.org/10.1007/978-1-4939-7819-9_24

Goedhart, J., van Weeren, L., Hink, M. A., Vischer, N. O. E., Jalink, K., & Gadella, T. W. J.

(2010). Bright cyan fluorescent protein variants identified by fluorescence lifetime screening.

Nature Methods, *7*(2), 137–139. https://doi.org/10.1038/nmeth.1415

Harwood, R., Goodman, E., Gudmundsdottir, M., Huynh, M., Musulin, Q., Song, M., &

Barbour, M. M. (2020). Cell and chloroplast anatomical features are poorly estimated from 2D

cross-sections. *New Phytologist*, *225*(6), 2567–2578. https://doi.org/10.1111/nph.16219

Jarvis, P., & López-Juez, E. (2013). Biogenesis and homeostasis of chloroplasts and other

plastids. *Nature Reviews Molecular Cell Biology*, *14*(12), 787–802.

https://doi.org/10.1038/nrm3702

Johnson, D. E., Ai, H., Wong, P., Young, J. D., Campbell, R. E., & Casey, J. R. (2009). Red

Fluorescent Protein pH Biosensor to Detect Concentrative Nucleoside Transport. *Journal of*

Biological Chemistry, *284*(31), 20499–20511. https://doi.org/10.1074/jbc.M109.019042

Kinsman, E. A., & Pyke, K. A. (1998). Bundle sheath cells and cell-specific plastid development

in *Arabidopsis* leaves. *Development*, *125*(10), 1815–1822.

https://doi.org/10.1242/dev.125.10.1815

Kleine, T., Nägele, T., Neuhaus, H. E., Schmitz-Linneweber, C., Fernie, A. R., Geigenberger, P.,

Grimm, B., Kaufmann, K., Klipp, E., Meurer, J., Möhlmann, T., Mühlhaus, T., Naranjo, B.,

Nickelsen, J., Richter, A., Ruwe, H., Schroda, M., Schwenkert, S., Trentmann, O., … Leister,

D. (2021). Acclimation in plants – the Green Hub consortium. *The Plant Journal*, *106*(1), 23–

40. https://doi.org/10.1111/tpj.15144

Klie, S. (2011). Analysis of the compartmentalized metabolome – a validation of the non-

aqueous fractionation technique. *Frontiers in Plant Science*, *2*.

https://doi.org/10.3389/fpls.2011.00055

Koniger, M., Delamaide, J. A., Marlow, E. D., & Harris, G. C. (2008). Arabidopsis thaliana

leaves with altered chloroplast numbers and chloroplast movement exhibit impaired

adjustments to both low and high light. *Journal of Experimental Botany*, *59*(9), 2285–2297.

https://doi.org/10.1093/jxb/ern099

Krantz, M., Zimmer, D., Adler, S. O., Kitashova, A., Klipp, E., Mühlhaus, T., & Nägele, T.

(2021). Data Management and Modeling in Plant Biology. *Frontiers in Plant Science*, *12*,

717958. https://doi.org/10.3389/fpls.2021.717958

Kunz, H.-H., Gierth, M., Herdean, A., Satoh-Cruz, M., Kramer, D. M., Spetea, C., & Schroeder,

J. I. (2014). Plastidial transporters KEA1, -2, and -3 are essential for chloroplast

osmoregulation, integrity, and pH regulation in *Arabidopsis*. *Proceedings of the National*

Academy of Sciences, *111*(20), 7480–7485. https://doi.org/10.1073/pnas.1323899111

Lampropoulos, A., Sutikovic, Z., Wenzl, C., Maegele, I., Lohmann, J. U., & Forner, J. (2013).

GreenGate—A Novel, Versatile, and Efficient Cloning System for Plant Transgenesis. *PLoS*

ONE, *8*(12), e83043. https://doi.org/10.1371/journal.pone.0083043

Liebers, M., Grübler, B., Chevalier, F., Lerbs-Mache, S., Merendino, L., Blanvillain, R., &

Pfannschmidt, T. (2017). Regulatory Shifts in Plastid Transcription Play a Key Role in

Morphological Conversions of Plastids during Plant Development. *Frontiers in Plant Science*,

8. https://doi.org/10.3389/fpls.2017.00023

Miyagishima, S., Froehlich, J. E., & Osteryoung, K. W. (2006). PDV1 and PDV2 Mediate

Recruitment of the Dynamin-Related Protein ARC5 to the Plastid Division Site. *The Plant Cell*,

18(10), 2517–2530. https://doi.org/10.1105/tpc.106.045484

Nobel, P. S. (2020). Water. In *Physicochemical and Environmental Plant Physiology* (pp. 53–

115). Elsevier. https://doi.org/10.1016/B978-0-12-819146-0.00002-X

Oi, T., Enomoto, S., Nakao, T., Arai, S., Yamane, K., & Taniguchi, M. (2017). Three-

dimensional intracellular structure of a whole rice mesophyll cell observed with FIB-SEM.

Annals of Botany, *120*(1), 21–28. https://doi.org/10.1093/aob/mcx036

Okazaki, K., Kabeya, Y., Suzuki, K., Mori, T., Ichikawa, T., Matsui, M., Nakanishi, H., &

Miyagishima, S. (2009). The PLASTID DIVISION1 and 2 Components of the Chloroplast

Division Machinery Determine the Rate of Chloroplast Division in Land Plant Cell

Differentiation. *The Plant Cell*, *21*(6), 1769–1780. https://doi.org/10.1105/tpc.109.067785

Osteryoung, K. W., & Pyke, K. A. (2014). Division and Dynamic Morphology of Plastids.

Annual Review of Plant Biology, *65*(1), 443–472. https://doi.org/10.1146/annurev-arplant-

050213-035748

Pipitone, R., Eicke, S., Pfister, B., Glauser, G., Falconet, D., Uwizeye, C., Pralon, T., Zeeman, S.

C., Kessler, F., & Demarsy, E. (2021). A multifaceted analysis reveals two distinct phases of

chloroplast biogenesis during de-etiolation in Arabidopsis. *ELife*, *10*, e62709.

https://doi.org/10.7554/eLife.62709

Pyke, K. A., & Leech, R. M. (1994). A Genetic Analysis of Chloroplast Division and Expansion

in Arabidopsis thaliana. *Plant Physiology*, *104*(1), 201–207.

https://doi.org/10.1104/pp.104.1.201

Robertson, E. J., Rutherford, S. M., & Leech, R. M. (1996). Characterization of Chloroplast

Division Using the Arabidopsis Mutant arc5. *Plant Physiology*, *112*(1), 149–159.

https://doi.org/10.1104/pp.112.1.149

Schulz, A., Knoetzel, J., Scheller, H. V., & Mant, A. (2004). Uptake of a Fluorescent Dye as a

Swift and Simple Indicator of Organelle Intactness: Import-competent Chloroplasts from Soil-

grown *Arabidopsis*. *Journal of Histochemistry & Cytochemistry*, *52*(5), 701–704.

https://doi.org/10.1177/002215540405200514

Sun, T., Yuan, H., Cao, H., Yazdani, M., Tadmor, Y., & Li, L. (2018). Carotenoid Metabolism in

Plants: The Role of Plastids. *Molecular Plant*, *11*(1), 58–74. https://doi.org/DOI:

10.1016/j.molp.2017.09.010

Unal, D., García-Caparrós, P., Kumar, V., & Dietz, K.-J. (2020). Chloroplast-associated

molecular patterns as concept for fine-tuned operational retrograde signalling. *Philosophical*

Transactions of the Royal Society B: Biological Sciences, *375*(1801), 20190443.

https://doi.org/10.1098/rstb.2019.0443

Waadt, R., Krebs, M., Kudla, J., & Schumacher, K. (2017). Multiparameter imaging of calcium

and abscisic acid and high-resolution quantitative calcium measurements using R-GECO1-

mTurquoise in Arabidopsis. *New Phytologist*, *216*(1), 303–320.

https://doi.org/10.1111/nph.14706

Winter, H., Robinson, D. G., & Heldt, H. W. (1994). Subcellular volumes and metabolite

concentrations in spinach leaves. *Planta*, *193*(4), 530–535.

https://doi.org/10.1007/BF02411558

Woodward, A. W., & Bartel, B. (2018). Biology in Bloom: A Primer on the *Arabidopsis thaliana*

Model System. *Genetics*, *208*(4), 1337–1349. https://doi.org/10.1534/genetics.118.300755

Yoshida, Y. (2018). Insights into the Mechanisms of Chloroplast Division. *International Journal*

of Molecular Sciences, *19*(3), 733. https://doi.org/10.3390/ijms19030733

Figure 1: Confocal microscopy reveals changes in gain and loss of function mutants in components of the plastid division machinery that are restricted to mesophyll cells. A, the diagram of the Arabidopsis plant shows the size of the plants when imaging and sample preparation was done. The * indicates the leaves used for imaging, and circles indicate the location on each leaf. From here, these samples were either used immediately for imaging using confocal microscopy or fixed and prepared for SBFSEM. **B**, No apparent changes in plant growth or appearance were observed in long-day cultivated genotypes used in this study, *35s-PDV1 35s-PDV2*, Col-0, *arc5-2* (scale bars = 2 cm). **C**, Confocal images of chlorophyll fluorescence for each genotype show clear size differences in mesophyll chloroplasts (scale bars = 25 μm). **D**, Confocal images of guard cell chloroplasts taken using chlorophyll fluorescence shows similar chloroplast volumes within guard cells of the three genotypes investigated (scale bars = 25 μm). **E**, Comparison of mesophyll chloroplast volumes deduced from z-stack of chlorophyll fluorescence recordings. Individual grow-outs are shown for each genotype. Statistical analysis done by one-way ANOVA with Tukey's multiple comparison test ($p<0.05$) showed no significant difference between Col-0 and *35s-PDV1 35s-PDV2* chloroplast volumes, neither in the full dataset, nor between any individual repetitions. Significant differences were observed between *arc5-2* and *35s-PDV1 35s-PDV2* but also between *arc5-2* and Col-0 mesophyll chloroplast volumes. Letters above each sample in the graph indicate groups of significance (±SE). **F**, Chloroplast numbers per mesophyll cell were counted and show an inverse relationship to that of chloroplast volumes observed between the genotypes. *35s-PDV1 35s-PDV2* has the highest average number of chloroplasts per cell with around 90 chloroplasts, followed by Col-0 averaging slightly fewer, at around 70 chloroplasts per cell. Finally, *arc5-2* mutants showed by far the lowest chloroplast count per cell with an average of 5 chloroplasts

(±SE). **G**, Guard cell chloroplast volumes calculated based on z-stacks of chlorophyll fluorescence recordings, in three separate grow outs, statistical analysis shows no significant difference ($p<0.05$) between any of the genotypes, neither within the whole dataset nor within individual growing cycles. Letters above each data set again show groups of significance (±SE). **H**, Guard cell chloroplast counts show that, unlike the similarities in volumes, a slight difference in average chloroplast number per guard cell can be observed. Col-0 guard cells contain on average 4.5 chloroplasts, while both the *35s-PDV1 35s-PDV2* and *arc5-2* genotypes contain only around 3.5 chloroplasts per guard cell (±SE).

Figure 2: Comparing mesophyll chloroplast volumes deduced from chlorophyll fluorescence and an envelope membrane located fluorescent marker. Comparison of **A**, chlorophyll fluorescence, **B**, the outer chloroplast envelope labeled with CFP visualized in **C**, as an overlay in Col-0 (scale bar = 10 μm). **D**, comparison between chloroplast volumes extracted by chlorophyll fluorescence and mTurquoise envelope marker shows slightly higher volumes in each genotype when calculated using the mTurquoise marker. A statistically significant difference ($p<0.05$) in chloroplast volumes was only found between Col-0 and *35s-PDV1 35s-PDV2* when compared to *arc5-2* mutants. No significant difference is observed between imaging methods when looking at individual genotypes (±SE).

Figure 3: Chloroplast volume analysis using SBFSEM. A, Reconstructions of Col-0 chloroplasts by using SBFSEM. Green objects represent chloroplasts labeled using the machine learning algorithm, while in grey a surface was created by thresholding, representing mostly the cell wall and some other high contrast objects within the cells. **B**, deduced chloroplast volumes measured by SBFSEM imaging. Statistical significance ($p<0.05$) in chloroplast volumes is shown by numbers above each dataset in the graph. A significant difference in chloroplast volumes can only be seen between Col-0 and *arc5-2*, as well as between *35s-PDV1 35s-PDV2*, and *arc5-2*. **C**, A frequency distribution of Col-0 and *35s-PDV1 35s-PDV2* shows a shift in volume distribution between the two genotypes (±SE).

Figure 4: Comparison of chloroplast volumes measured by imaging type. A-C, comparison of mesophyll chloroplast volumes extracted using confocal imaging of chlorophyll fluorescence (Chl *a*), CFP envelope marker, and SBFSEM. The means of the data collected using each imaging method was taken for all genotypes. Each pair of imaging methods is compared by plotting the means (±SE). An identity line was added to highlight differences in means between imaging types in each graph. This shows overall similar volumes and distributions between the methods, with data extracted using the CFP envelope marker showing the largest volumes and chlorophyll fluorescence having the lowest. A one-way ANOVA done for statistical analysis only showing a significant difference ($p<0.05$) when comparing *35s-PDV1 35s-PDV2* and Col-0 volumes acquired by either method to *arc5-2*. No significant difference was found between imaging methods for any genotype.