Intestinal LKB1 loss drives a pre-malignant program along the serrated cancer

pathway

Short title: LKB1 loss drives serrated cancer predisposition

Plugge S.F.[#,1], Ma H.[#,1,2], van der Vaart J.Y. [#,1], Sprangers J.[1], Morsink F.H.M.[2], Xanthakis D.[1], Jamieson C.[1],

Stroot, K.B.W.[1], Keijzer A.R.[1], Margaritis T[3]., Candelli T.[3], Straver R.[1], de Ridder J.[1], Holstege F.C.P.[3, 4], de

Leng W.W.J.[2], Offerhaus G.J.A.[2], Merenda A.[*,1], Maurice M.M.[*,1]

1. Center for Molecular Medicine and Oncode Institute, University Medical Center Utrecht, Utrecht, The

Netherlands

2. Department of Pathology, University Medical Center Utrecht, Utrecht, The Netherlands

3. Princess Maxima Center for Pediatric Oncology, Utrecht, Utrecht, The Netherlands

4. Center for Molecular Medicine, University Medical Center Utrecht, Utrecht, The Netherlands

[#]These authors contributed equally

*Corresponding authors (a.merenda@huborganoids.nl & m.m.maurice@umcutrecht.nl)

Grant support

This work is part of the Oncode Institute, which is partly financed by the Dutch Cancer Society. This work

was supported by the China Scholarship Council (to H.M.), the Dutch Cancer Society KWF Grant 13112 (to

M.M.M.), Zon-MW TOP grant 91218050 (to M.M.M.) and the Netherlands Organization for Scientific

Research (NWO) Gravitation programme IMAGINE! (project number 24.005.009) (to M.M.M.).

Conflict of interest

M.M.M. is co-founder and shareholder of Laigo Bio and inventor on patents related to membrane protein

degradation; J.R. is co-founder of Cyclomics BV., a genomics company. All other authors declare to have no

conflict of interest.

Data availability

The scRNA-seq and bulk RNA-seq data generated in this study will be uploaded to Gene Expression

Omnibus (GEO) upon publication. If needed prior publication, data can be requested via the corresponding

author.

Author contributions

S.F.P, H.M., J.Y.V., A.M. and M.M.M., conceived and designed the experiments. S.F.P, H.M., J.Y.V., A.M.,

D.X., F.H.M.M., C.J., K.B.W.S and A.R.K performed the experiments. S.F.P, H.M., A.M., J.Y.V., J.S., T.M.,

T.C., R.S., J.R., F.C.P.H., G.J.A.O. and M.M.M. analyzed the data. W.W.J.L. and G.J.A.O. provided patient

material. S.F.P, H.M., A.M., J.Y.V. and M.M.M. wrote the manuscript, which was reviewed by all authors.

Abstract

<u>Background & Aims</u>

Heterozygous inactivating mutations of Serine Threonine Kinase 11 (STK11)/Liver Kinase B1 (LKB1) are

causative to the Peutz-Jeghers syndrome (PJS), a hereditary disease characterized by gastrointestinal

hamartomatous polyposis and increased cancer susceptibility. While LKB1 loss-induced polyp formation has

been ascribed to non-epithelial tissues, how LKB1 deficiency increases cancer risk of patients by altering the

phenotypical landscape and hierarchical organization of epithelial tissues remains poorly understood.

<u>Methods</u>

Using CRISPR/Cas9, we generated heterozygous and homozygous Lkb1-deficient mouse small intestinal

and human colon organoids. These organoids were characterized by an integrated approach that combines

imaging, bulk and single-cell RNA sequencing and growth factor dependency assays. Our findings were

validated in human PJS-derived tissues using immunohistochemistry and linked to colorectal cancer profiles

using the TCGA cancer database.

<u>Results</u>

Our results reveal that heterozygous Lkb1 loss is sufficient to push intestinal cells into a premalignant

transcriptional program associated with serrated colorectal cancer, which is further amplified by loss-of-

heterozygosity. This altered epithelial growth state associates with persistent features of regeneration and

enhanced EGFR ligand and receptor expression, conferring niche-independent growth properties to Lkb1-

deficient organoids. Moreover, our newly generated LKB1-mutant signature is enriched in sporadic serrated

colorectal cancer, and synergistic cooperation of Lkb1-deficiency with mutant Kras was experimentally

confirmed by assessing organoid growth properties and transcriptomes.

<u>Conclusions</u>

Heterozygous loss of LKB1 pushes intestinal cells into a chronic regenerative state which is amplified upon

loss-of-heterozygosity. Lkb1-deficiency thereby generates fertile ground for serrated colorectal cancer

formation in the intestine, potentially explaining the increased cancer risk observed in PJS.

Keywords: LKB1; Peutz-Jeghers Syndrome; Serrated tumors; Colorectal cancer; Organoids; Regeneration

Introduction

Peutz-Jeghers syndrome (PJS) is a hereditary genetic disorder affecting approximately 1:50.000-1:200.000 individuals worldwide[1]. PJS is characterized by the formation of benign hamartomatous polyps throughout the gastrointestinal tract, which necessitates periodical surgical resection. Additionally, patients carry an increased risk of developing early-onset epithelial tumors, particularly within the gastrointestinal tract (up to 66%), breast (up to 54%), pancreas (up to 36%) and gonads[1–3], resulting in a cumulative lifetime risk of 93%[3]. Moreover, PJS patients are predisposed to small intestinal cancers that rarely develop in the general population (0.03% compared to 13% in PJS)[2,4].

PJS is caused by heterozygous inactivating mutations in the tumor suppressor gene Liver Kinase B1 (*LKB1/STK11*)[5]. Of note, inactivation of *LKB1* is also linked to the formation of sporadic cancers such as lung carcinoma, melanoma, pancreatic and cervical cancers[6,7]. The *LKB1* gene encodes for a serine/threonine kinase involved in diverse cellular processes, including energy metabolism, cell adhesion, DNA methylation and apicobasal polarity[8–11]. Loss of heterozygosity (LOH) of *LKB1* is a driving force of cancer development, since LOH occurs in over 64% of PJS-derived carcinomas[12,13].

Recent studies using murine models revealed that *Lkb1* loss in non-epithelial lineages suffices to induce the development of GI polyps resembling those observed in PJS patients[14–17]. This led to a reevaluation of *Lkb1*'s role in non-epithelial tissues, with an emphasis on polyposis formation. The relationship between polyp formation and cancer predisposition in PJS patients however remains highly debated, as LOH and dysplastic transformation of PJS polyps appear to be very rare compared to other polyposis syndromes like familial adenomatous polyposis (FAP)[12,13,18,19].

Since *Lkb1* loss mediates predisposition to carcinoma development in multiple epithelial tissues irrespective of polyp formation, we hypothesize that epithelial-intrinsic alterations may sensitize these tissues to malignant transformation. Notably, LKB1 deficiency in the intestinal epithelium itself does not mediate polyp formation but was shown to alter secretory cell fate specification[14,20,21]. Despite these advancements, how alterations in intestinal epithelial organization due to *LKB1* predispose for cancer development in PJS patients remains poorly understood.

Here, we model *mono*- and *bi*-allelic inactivation of *Lkb1* in mouse and human intestinal organoids to investigate how *Lkb1* loss affects the intestinal epithelial cellular and transcriptional landscape. We uncover that *Lkb1* loss promotes features of epithelial transformation linked to the serrated pathway for intestinal carcinogenesis. These features include loss of classical adult stem cell populations, activation of regeneration-related transcriptional programs, intestinal-to-gastric cellular transdifferentiation, and the acquirement of niche-independent growth properties. Our findings argue that *Lkb1* loss mediates epithelium-wide activation of a pre-cancer program, providing an explanation for the increased cancer risk observed in PJS patients. By linking *Lkb1* loss to a major pathway for colorectal cancer (CRC) development, these findings pave a path for the development of new strategies for prevention, surveillance and precision treatment of PJS tumors and other malignancies harboring *LKB1*-inactivating mutations.

Materials and Methods

Patient material

Colorectal tissue collection for the generation of healthy colon organoids was performed according to the

guidelines of the European Network of Research Ethics Committees (EUREC). Formalin-Fixed Paraffin-

Embedded (FFPE) intestinal tissue samples were obtained from pathology archives. For details see

supplementary methods.

Organoid culture

Both mouse small intestinal organoids and human colon organoids were established and cultured as

previously described[22,23]. Organoids were grown in Matrigel (Corning) or BME (Cultrex) and passaged every

6-8 days; their medium was refreshed every 2-3 days. For medium composition see supplementary

materials.

Gene editing of organoids

For generation of mutant and overexpression organoids, electroporation of organoids was performed as

previously described[24]. For further details, plasmids and organoid selection see supplementary materials.

Imaging of organoids

Organoids were fixed and stained 3-4 days after passaging. For detailed protocols on immunofluorescence,

immunohistochemistry and electron microscopy see supplementary materials.

Bulk RNA sequencing

Organoids were lysed in RLT lysis buffer (Qiagen) 3-5 days after passaging. Total RNA was processed for

sequencing using Illumina Truseq Stranded mRNA kit and sequenced. For details see supplementary

materials.

Single cell RNA sequencing

Single cells from organoid were harvested using TrypLE, four days after passaging. Cells were filtered and

scRNA-seq was performed according to the Sort-seq protocol[25]. For details see supplementary materials.

Results

Lkb1 bi-allelic loss results in phenotypic heterogeneity and increased self-renewal potential of small intestinal organoids.

To study the role of _Lkb1_ loss in the intestinal epithelium, we employed CRISPR/Cas9 technology to generate _Lkb1_-mutant mouse small intestinal organoid (mSIO) lines. Targeted organoids were cultured transiently in WENR (see supplementary materials) to enhance outgrowth efficiency[26]. Under these conditions, organoids acquire a cystic morphology, due to the expansion of stem cells[22,27]. To obtain genotypically clonal _Lkb1_-mutant cultures, we grew organoids from single cells. Sequencing analysis confirmed the generation of homozygous mutant clones carrying bi-allelic truncating indels (Figure 1A, _Lkb1$^{-/-}$_). In addition, we obtained a heterozygous clone that carried an out-of-frame insertion in one allele and an in-frame deletion (p.P38_R39del) in the other allele. The two deleted amino acids are located outside essential protein regions (Figure 1A, _Lkb1$^{+/-}$_) and allowed for protein expression (Figure 1B), predicting retained protein function. Subsequently, organoids were transferred to ENR medium, allowing for formation of typical budding structures resembling the _in vivo_ crypt-villus axis[22]. Clones that transitioned into budding organoids over four passages comprised all genotypes, including wild-type (WT), _Lkb1$^{+/-}$_ or _Lkb1$^{-/-}$ (Lkb1$^{bud-/-}$)_. Strikingly, a subset of clones that retained cystic morphology were exclusively genotyped as _Lkb1$^{-/-}$_ (_Lkb1$^{cys-/-}$_) (Figure S1A). In line with Lkb1's role in AMPK-mTOR signaling, we observed reduced phosphorylated Ampk (pAmpk), a direct substrate of Lkb1 kinase activity[10], along with upregulated mTor signaling, indicated by increased phosphorylated 4E-BP1 in _Lkb1$^{-/-}$_ organoids (Figure 1C-D). As expected, _Lkb1$^{+/-}$_ organoids displayed an intermediate phenotype, consistent with mono-allelic expression (Figure 1C-D). Furthermore, _Lkb1$^{bud-/-}$_ organoid cultures displayed a notable accumulation of dead cells at their basal side (Figure 1E), validated by cleaved caspase-3 staining (Figure 1F). This observation differed from WT and _Lkb1$^{+/-}$_ organoids for which dying cells were apically extruded and shed into the lumen, as shown previously (Figure 1F)[22]. Mislocalization of dead cells in _Lkb1$^{bud-/-}$_ organoids was corrected by wild-type LKB1 overexpression, indicating that this is a consequence of _Lkb1_ loss (Figure S1B-C).

To assess the stability of budding and cystic _Lkb1$^{-/-}$_ organoid phenotypes, we generated a _Lkb1$^{-/-}$_ bulk population (100% knockout score). One week after transfer to ENR medium, the cystic-to-budding ratio was 1:1 (Figure S1D). We next expanded nine budding and nine cystic clonal organoids lines and tracked their morphology. After two passages, ~70% of _Lkb1$^{bud-/-}$_ clones retained budding morphology, while all _Lkb1$^{cys-/-}$_ clones remained cystic (Figure S1E). Thus, homozygous _Lkb1_ loss leads to two distinct morphologies, with some budding-to-cystic conversion. Of note, reversal of cystic morphology by reintroduction of LKB1 expression was not possible, as LKB1$^+$/mCherry$^+$ cells were rapidly lost over time (Figure S1F-G), potentially due to competition by _Lkb1$^{cys-/-}$_ cells[28]. Next, we examined whether _Lkb1_ loss alters the ability to reconstitute organoids from single cells, as a proxy for their self-renewal capacity. While WT, _Lkb1$^{+/-}$_ and _Lkb1$^{bud-/-}$_ organoids displayed similar outgrowth rates, _Lkb1$^{cys-/-}$_ organoids carried a significantly improved capacity to form organoids (Figure 1G).

To rule out that the observed phenotypes are caused by unintended gRNA-induced mutations rather than _Lkb1_ loss, we employed an alternative, non-overlapping _Lkb1_-targeting gRNA (gRNA2). Targeting _Lkb1_ with gRNA2 led to similar morphologic changes, including shedding of dead cells at the basal side and a divergent budding and cystic morphology for _Lkb1$^{-/-}$_ (Figure S2). Moreover, newly generated _Lkb1$^{cys-/-}$_ clones

also displayed superior reconstitution potential compared to their *Lkb1^bud-/-^* counterparts (Figure S2C). We
conclude that *Lkb1*-deficiency results in specific organoid phenotypes related to loss of one or both alleles of
Lkb1.

***Lkb1* loss alters the number, morphology and positioning of secretory cells.**

To investigate whether our *Lkb1*-mutant organoid lines serve as representative models for intestinal
epithelial *Lkb1* loss *in vivo*, we examined alterations in localization and morphology of Paneth (Lyz1^+^) and
goblet (Muc2^+^) cells previously reported for the *Lkb1*-mutant mouse intestine[20,21]. Compared to WT
organoids, Lyz1^+^ cells in *Lkb1^+/-^* and *Lkb1^bud-/-^* organoids were scattered beyond the crypt region into the
villus domain (Figure 2A, S3A). Additionally, in both WT and *Lkb1^+/-^* organoids, Muc2^+^ cells were sparsely
located to the villus region, while *Lkb1^bud-/-^* organoids displayed an increase in Muc2^+^ cells that were found
scattered throughout the crypt-villus axis (Figure 2B). These observations resemble reported aberrancies in
Paneth and goblet cell numbers and localization in *Lkb1*-deficient mouse intestine *in vivo*[20]. Conversely,
*Lkb1^cys-/-^* organoids lacked Lyz1^+^ cells but did form Muc2^+^ cells that are usually lacking in cystic organoids[27],
indicating that secretory lineage specification is also altered within these organoids (Figure 2A-B).

We next compared these findings to the intestinal tissue of human PJS patients. The number and
localization of MUC2^+^ cells per intestinal crypt were similar in healthy and PJS patient-derived tissues (Figure
2C-D, S3B). Compared to healthy donors, however, LYZ^+^ cells were mislocalized within the PJS mucosa
(Figure 2E-F, S3C). Thus, heterozygous loss of *LKB1* in PJS intestines leads to an altered LYZ^+^ cell
distribution, similar to our observations in *Lkb1^+/-^* organoids.

To examine *Lkb1* loss-induced alterations in secretory cell morphology at the ultrastructural level, we
employed electron microscopy. In WT intestinal organoids, Paneth cells displayed a typical apical
accumulation of spherically-shaped secretory granules filled with electron-dense content (Figure 2G)[22], while
*Lkb1^+/-^* Paneth cells carried granules with an abnormal 'sausage-shaped' electron-dense core (Figure 2G). In
*Lkb1^bud-/-^* Paneth cell granules, the electron-dense core was surrounded by a peripheral electron-lucent halo
(Figure 2G). WT goblet cells were columnar-shaped and contained typical large translucent mucus-filled
granules (Figure 2H)[22,29]. *Lkb1^+/-^* mucus-secreting cells however were rounded and flattened while *Lkb1^bud-/-^*
goblet-like cells displayed granules with an unusual electron-dense core (Figure 2H). In addition, in *Lkb1^cys-/-^*
organoids the mucus-like granule content was frequently lost during EM preparation, suggesting an altered
mucin composition and explaining the limited number of observed Muc2^+^ cells (Figure 2H). Morphologically,
Paneth- and goblet-like cells in all *Lkb1*-mutant organoids looked like 'intermediate' cells that were proposed
previously to represent transition states between undifferentiated and mature cells[29], suggesting that *Lkb1*
loss may interfere with terminal differentiation of secretory cells. Furthermore, although previous studies[11]
identified roles of *LKB1* in apical brush border formation using 2D human cell lines, our EM images showed
no obvious brush border defects in any of the *Lkb1*-mutant organoids (Figure S3D).

Our findings thus reveal that *Lkb1* loss in budding organoids induces alterations in secretory cell
number, localization and morphology similar to *in vivo* mouse models and PJS patient intestines, validating
these organoids as tools for modelling epithelial alterations in PJS. While *Lkb1^cys-/-^* organoids are more
difficult to directly link to PJS epithelial organization *in vivo*, they may represent an intermediate state in
cancer progression. We therefore included all clonal organoid variants for further characterization.

215 *Lkb1* **mono- and bi-allelic loss induces incremental expression of a regenerative gene signature.**

216 To examine how *Lkb1* loss affects the epithelial landscape, we determined transcriptional changes using

217 bulk mRNA sequencing. Compared to WT organoids, we observed differential expression of 1298, 2312 and

218 7816 genes in *Lkb1*[+/-], *Lkb1*[bud-/-] and *Lkb1*[cys-/-] organoids, respectively (Table S1).

219 Using gene set enrichment analysis (GSEA) we examined transcriptional gene profiles of individual

220 intestinal cell types in *Lkb1*-mutant organoids[30]. For clarity's sake, in the following sections we will use '*Lkb1*-

221 mutant' to refer to all genotypes (both mono-allelic and bi-allelic deficient). All *Lkb1*-mutant clones showed

222 significant downregulation of gene sets linked to mature Paneth cells, whereas goblet cell gene sets were

223 enriched within *Lkb1*[bud-/-] organoids in comparison to WT organoids, again suggesting alterations in secretory

224 cell lineage specification (Figure 3A). Noticeably, the *Lgr5*[+] adult stem cell signature[30] was downregulated in

225 all *Lkb1*-mutant clones (Figure 3A, S4A), reminiscent of fetal intestinal spheroids that rely on *Lgr5*-negative

226 progenitors[31]. Indeed, we uncovered a gradual enrichment of an embryonic spheroid gene signature[31] from

227 WT to *Lkb1*[+/-] to *Lkb1*[bud-/-] organoids, correlating with the number of lost *Lkb1* alleles (Figure 3A-B). This

228 transition was further enhanced in *Lkb1*[cys-/-] versus *Lkb1*[bud-/-] organoids (Figure S4A-B). Previous studies

229 linked transient upregulation of a fetal-like intestinal gene signature to epithelial remodeling and collagen

230 deposition during injury repair[32–36]. In line, all *Lkb1*-mutant organoid clones exhibited enrichment of gene

231 signatures associated with intestinal regeneration following damage or infection (Figure 3A, S4A).

232 To confirm that *Lkb1* deficiency rather than organoid morphology mediates activation of a

233 regenerative program in Lkb1[cys-/-] organoids, we analyzed transcriptomes of *Apc*[-/-] and WT Wnt-treated cystic

234 organoids[27]. *Lkb1*[cys-/-] organoids were significantly enriched for expression of regenerative gene sets

235 compared to *Apc*[-/-] and Wnt-treated organoids (Figure S4C). These findings confirm that *Lkb1* loss itself

236 promotes a regenerative state, although morphology may contribute to the enhancement of this phenotype.

237 Furthermore, all *Lkb1*-mutant clones, and control fetal intestinal organoids (E16.5), displayed

238 increased expression of the fetal marker Anxa1[32], which was absent in WT adult intestinal organoids (Figure

239 3C, S4D). These findings indicate that *Lkb1* deficiency converts the epithelium into a primed regenerative

240 state that is already apparent upon monoallelic *Lkb1* loss, which mirrors the PJS intestinal epithelium.

241

242 *Lkb1*-**mutant organoids show a Yap-induced regenerative state and display intra-epithelial self-**

243 **sufficiency for Egf.**

244 Over recent years, Yap signaling was placed central to intestinal regeneration[37]. Additionally, *LKB1* was

245 identified as a negatively regulator of the Hippo-Yap pathway[38]. We therefore wondered if *Lkb1* loss may

246 activate Yap signaling in intestinal epithelial organoids. Indeed, we identified enhanced expression of

247 multiple Yap-regulated gene signatures[37,39] in *Lkb1*-mutant organoids (Figure 4A, S5A). Notably, Yap target

248 gene expression increased with the number of lost *Lkb1* alleles (normalized enrichment scores of 2.01 and

249 2.68, respectively). Yap target gene expression was further enhanced in *Lkb1*[cys-/-] organoids compared

250 *Lkb1*[bud-/-] organoids (Figure S5A), which is consistent with their enhanced regenerative transcriptional

251 program. Sustained YAP signaling may explain their related cystic morphology (Figure 1E) and Paneth cell

252 deficiency (Figure 2A), as YAP signaling needs to be suppressed to induce symmetry breaking and Paneth

253 cell formation in mSIOs (Figure S5A)[40]. Immunofluorescence staining confirmed increased Yap protein

254 expression and nuclear localization in both *Lkb1*[bud-/-] and *Lkb1*[cys-/-] organoids (Figure 4B-C, S5B-D).

YAP pathway activation is well-known to enhance EGF family ligand expression during
regeneration[37,41]. In line, *Lkb1*-mutant organoids showed upregulated expression of Egf ligands and receptor
family members, correlating with *Lkb1* allelic loss and alterations in organoid morphology (Figure 4D-E, S5E-
F). Given the dependency of healthy intestinal organoids on Egf, we examined if *Lkb1*-mutant organoids may
grow without Egf. Unlike WT organoids, all *Lkb1*-mutant organoids grew out without Egf supplementation
(Figure 4F-G, S5G-H). To ensure this was due to endogenous ligand secretion and not downstream pathway
activation, we treated the organoids with Gefitinib, an EGFR inhibitor. All organoid lines remained sensitive
to EGFR inhibition (Figure 4H, S5I), confirming endogenous EGF production as a driver of growth.
Interestingly, increased Egf ligand expression was linked previously to enhanced secretory cell
differentiation[42,43]. Indeed, increased secretory cell differentiation in *Lkb1*-mutant organoids (Figure 2)
correlates with increased Egf ligand expression (Figure 4D).
These results demonstrate that *Lkb1* loss induces a Yap-induced regenerative state leading to EGF
ligand self-sufficiency. As niche-independent growth is a hallmark of cancer progression[44], these results
suggest that *Lkb1*-mutant epithelial cells carry a growth advantage which may promote tumorigenesis
initiation.
270
Lkb1-deficiency mediates reprogramming of intestinal stem cells into a regenerative state.
To analyze how *Lkb1* loss alters intestinal lineage specification and cellular hierarchies, we subjected WT
and *Lkb1*-mutant organoids to single cell RNA sequencing. We included mouse fetal intestinal organoids, to
explore a potential overlap with fetal gene expression. Unsupervised clustering identified 19 distinct clusters
(Figure S6A). *Lkb1*$^{cys-/-}$ organoids and fetal cells clearly separated, while *Lkb1*$^{bud-/-}$ and *Lkb1*$^{+/-}$ organoids
more closely resembled WT cells (Figure 5A). These findings indicate that although *Lkb1*$^{cys-/-}$ organoids
express a fetal-like gene program they do not fully convert into a fetal state (Figure 5A). Furthermore,
Lkb1$^{cys-/-}$ organoids harbor a homogeneous cell population distinct from budding organoid clusters, lacking a
known intestinal cellular identity. We therefore annotated cell types only in budding organoid lines, using
intestinal cell type gene signatures (Figure 5B, S6B-I)[30]. As the gene signatures for Paneth and goblet cells
mapped to the same cluster (Figure S6E, S6H), we marked this cluster as secretory progenitors, similar to
previous reports[45]. We reclustered all cells from budding organoids (WT, *Lkb1*$^{+/-}$ and *Lkb1*$^{bud-/-}$) for further
analyses (Figure 5C-D). Cell type fraction analysis revealed an increased fraction of secretory progenitor
cells in *Lkb1*$^{bud-/-}$ organoids (Figure 5E), confirming alterations in secretory cell differentiation. This increase
corresponded with a decrease in enterocyte numbers, suggesting a shift in differentiation from absorptive
towards secretory lineages (Figure 5E). Furthermore, we identified 'intermediate' cells in *Lkb1*$^{+/-}$ and *Lkb1*$^{bud-}$
$^{/-}$ organoids that co-express features of Paneth and goblet cells (Figure 5F), in line with our ultrastructural
morphological analysis.
Notably, *Anxa1* expression in *Lkb1*$^{bud-/-}$ organoids was mainly enhanced within stem and transit
amplifying (stem/TA) cell clusters, indicating that these progenitor populations drive the epithelium's
regenerative state (Figure 5G). Reclustering of stem/TA cell populations showed distinct clusters for WT,
Lkb1$^{+/-}$ and *Lkb1*$^{bud-/-}$ stem/TA cells (Figure S6J-K). GSEA on genes differentially expressed between WT
stem cells and *Lkb1*$^{+/-}$ *and Lkb1*$^{bud-/-}$ stem cells (Table S2) revealed enrichment of goblet cell gene expression
within *Lkb1*$^{+/-}$ *and Lkb1*$^{bud-/-}$ stem cell populations (Figure 5H-I). Furthermore, *Lkb1*$^{+/-}$ *and Lkb1*$^{bud-/-}$ stem cells
displayed enrichment of regenerative and Yap gene signatures, depletion of the *Lgr5*$^+$ stem cell signature,

and increased expression of Anxa1 (Figure 5H, J). These findings support a model in which *Lkb1* loss
induces a regenerative response originating from the stem cell compartment.
In conclusion, our results argue that *Lkb1*-mutant stem/TA cells reside in a regenerative state and
preferentially differentiate towards secretory cell lineages.
300

Lkb1 loss drives a pre-malignant program along the serrated pathway which is amplified by
additional Kras mutations.
Next, we aimed to obtain insight in how transcriptional reprogramming induced by *Lkb1* loss may promote
cancer predisposition. CRC mainly develops via two pathways: the classical adenoma-carcinoma sequence,
initiated by bi-allelic *APC* mutations, and the serrated cancer pathway, initiated by *BRAF* or *KRAS*
mutations[46,47]. We examined expression of gene sets linked to these CRC routes to assess if *Lkb1* loss
drives entry into one of these pathways. *Lkb1*-mutant organoids displayed significant enrichment of genes
that mark neoplastic populations of serrated-specific cells (SSC[46], iCMS3[48]), which increased with the
number of lost *Lkb1* alleles (normalized enrichment scores of 1.70 and 2.16, respectively) (Figure 6A, S7A).
Single cell analysis confirmed this trend, attributing the increase of the serrated gene program to *Lkb1*$^{+/-}$ *and*
Lkb1$^{bud-/-}$ stem cell populations (Figure S8). This increase was accompanied by the loss of adenoma-specific
genes (ASC[46], iCMS2[48]) (Figure 6A, S7A) and an upregulation of expression of gastric markers like *Aqp5*,
Tacstd2 [*Trop2*], *Tff2*, and *Msln* (Figure 6B, S7B, Table S1), indicating the occurrence of intestinal-to-gastric
cellular transdifferentiation[46]. Notably, *LKB1* expression levels were significantly decreased within sessile
serrated lesions (SSLs)[46] compared to pre-cancerous conventional adenomas (AD)[46] (Figure 6C), supporting
the observation that a decrease in LKB1 function accommodates SSL formation.
To investigate the susceptibility of an *Lkb1*-mutant epithelial state to early driver mutations found in
different CRC subtypes, we introduced *Apc* (Figure S7C) or *Kras* mutations (Figure S7D) in *Lkb1*-mutant
organoids. *Kras* mutations synergistically enhanced *Lkb1*$^{bud-/-}$ organoid formation from single cells, while *Apc*
mutations did not (Figure 6D-E, S7E-F). Bulk mRNA sequencing of double-mutant organoids showed that
Lkb1 loss acts synergistically with Kras^{G12D} mutations to drive a stepwise SSC transcriptomic shift that
correlated with allelic loss (Table S3, Figure 6F, S7G). By contrast, no synergism of *Lkb1* loss with ASC
signature expression or with *Apc*$^{-/-}$ mutations was observed (Figure 6G, S7H).
These results indicate that *Lkb1* loss drives a state of cancer predisposition along the serrated CRC
pathway, which is amplified by *Kras* mutations in *Lkb1*$^{+/-}$ and *Lkb1*$^{bud-/-}$ organoids. Furthermore, *Lkb1*$^{cys-/-}$
organoids achieve an advanced serrated state without additional oncogenic mutations. Thus, loss of *LKB1*
primes the intestinal epithelium for progression along the serrated pathway.
328

LKB1-mutant phenotypes are conserved in human colon organoids and correlate with serrated
features in sporadic CRC.
To examine if the Lkb1-mutant phenotypes observed in mSIOs can be translated to the human colon
epithelium, we obtained one *LKB1*$^{+/-}$ and two *LKB1*$^{-/-}$ clonal human colon organoid (HCO) lines for which we
confirmed a decrease in LKB1 protein expression. In line, pAMPK levels were decreased in *LKB1*$^{-/-}$, but
higher in *LKB1*$^{+/-}$ HCOs, indicating retained function of the in-frame edited *LKB1* allele (Figure 7A-D).
Furthermore, progressive loss of *LKB1* induced enrichment of secretory gene signatures[49] and multiple
regenerative gene sets[31–34,37,39], similar to our observations in mSIOs (Figure 7E, Table S4). *LKB1* loss also

induced EGF self-sufficiency in HCOs, indicated by an increased expression of EGF receptor family
members (in both *LKB1*[+/-] and *LKB1*[-/-]; Figure 7F) and EGF ligands (in *LKB1*[-/-]; Figure 7G) as well as an
increased tolerance for EGF withdrawal (Figure 7H-I). Notably, all *LKB1*-mutant HCOs retained sensitivity to
Gefitinib treatment (Figure 7I). Finally, *LKB1*-mutant HCOs also displayed enrichment of gene signatures
linked to the serrated pathway (Figure 7J).

To assess how these findings correlate with human CRC development, we derived an *LKB1*-mutant
gene signature from our HCO models using a directional strategy, taking gene dosage as a covariate, and
assessed its correlation with human CRC datasets. Expression of the *LKB1*-mutant gene signature was
found enriched within human SSLs compared to tubular adenomas[50] (Figure 7K), within mucinous CRC
(Figure 7L), a subtype linked to SSLs[51], and in serrated consensus molecular subtypes CMS1 and CMS4
(Figure 7M)[47].

Together, these data underscore the relevance of our organoid-derived models. Our findings show
that *LKB1* loss-induced phenotypes are conserved among species in the gastrointestinal tract and display
shared identity with SSLs in human patients.

Discussion

Over recent years, *Lkb1*-deficient mouse models have faithfully mimicked intestinal hamartomatous
polyposis, a hallmark of PJS disease, suggesting that non-epithelial loss of *LKB1* function drives PJS
polyposis[14–17]. The relationship between polyposis and epithelial cancer predisposition in PJS however
remains highly debated and does not explain extra-intestinal cancer development[2,12,13,18]. Furthermore, the
impact of heterozygous and homozygous *Lkb1* loss on epithelial tissue organization at the single cell level
has not been addressed.

We employed mSIOs and HCOs to understand how *Lkb1* loss affects epithelial cell identities and
drives a pre-malignant state for CRC. By generating heterozygous and homozygous *Lkb1* knockouts, we
aimed to model cancer predisposition and development in PJS patients, as over 64% of PJS-derived
carcinomas present inactivation of the second gene copy[12,13]. Our findings argue that heterozygous *Lkb1*
loss is sufficient to push cells into a premalignant program along the serrated CRC pathway, independent of
the micro-environment. This premalignant program is defined by a sustained regenerative response, niche-
independent growth, and aberrant secretory cell differentiation, which is further amplified upon loss of the
second allele. If this epithelial-intrinsic phenotypic shift can be fully attributed to the known functions of
LKB1[10,38], or whether it is further amplified by LKB1[+/-] mesenchyme-induced polyposis in PJS patients
requires further investigation.

Our organoid-based model validates the link between *Lkb1* loss and aberrant secretory cell
specification *in vivo*. We observed that *Lkb1*-deficiency mediates an increase in goblet cells and mucus
formation and the generation of secretory cells with 'intermediate' features, both morphologically and
transcriptionally. Supporting this, our newly generated *LKB1*-mutant gene signature correlated with
transcriptional profiles of mucinous subtypes of sporadic CRC. Similar secretory cell alterations induced by
Lkb1 loss may occur in other epithelial tissues, as mucinous carcinomas are common in PJS patients[52,53].
Although the underlying mechanism requires further investigation, these findings indicate a link between
secretory cell alterations and cancer subtype progression.

Transcriptomic analysis revealed that *Lkb1* loss induces a dosage-dependent expression of a
regenerative epithelial program. This tissue repair program was activated in standard culture conditions,
without external damage, suggesting an inherent cellular response upon *Lkb1* loss. We anticipate that, in
PJS patients, external injury events that inflict on the *LKB1*-mutant intestine may enhance this response and
lock the epithelium into a chronic regenerative state, which may contribute to increased tumor development.
The regenerative state also showed increased *Egf* receptor family and ligand expression, endowing niche-
independent properties, which is regarded as a hallmark of cancer[44]. The activation of regenerative
programs has been extensively linked to an increased risk of tumorigenesis in many epithelial organs[54–56],
including pancreas and breast. Thus, induction of a chronic regenerative state by *Lkb1* loss may potentially
explain the increased risk of cancer development in other organs in PJS patients as well.

Importantly, we uncovered that *Lkb1* loss predisposes for the serrated CRC pathway, accompanied
by common serrated features like gastric metaplasia, a regenerative gene program and classical stem cell
depletion[46,47]. Furthermore, elevated expression of our newly generated *LKB1*-mutant gene signature was
significantly enriched in SSLs compared to classical adenomas[50]. These findings suggest that PJS cancer
pathways differ from the conventional APC-mutant adenocarcinomas in FAP[57].

A recently presented model proposed that different environmental triggers like microbial dysbiosis
and subsequent MAPK pathway activation drive SSL formation[58]. While we currently do not know how
external stressors (such as microbial dysbiosis) may affect tumor development in PJS patients, we observed
that introduction of oncogenic *Kras* in *Lkb1*-mutant organoids amplifies serrated features and enhances
organoid-forming capacity. Furthermore, our results reveal that homozygous *Lkb1* loss induces two organoid
phenotypes with either a budding or a cystic morphology accompanied by strongly induced serrated
features. Together, our findings suggest that the PJS epithelium is sensitized to events that synergistically
drive cells into a premalignant serrated state, such as intestinal damage or activating mutations in MAPK
pathway components. Furthermore, as clinical management of PJS patients involves life-long monitoring for
polyp resection and (pre)cancerous lesion screening, our results support a rationale for investigating SSL
prevalence in PJS cohorts. The malignancy of such lesions may be characterized by markers identified in
this study, such as ANXA1.

Together, our study reveals that intestinal organoids are valuable models for *LKB1* mutation effects
in PJS patients. Our findings establish a new function of *LKB1*, relating its loss to the induction of a
premalignant serrated program which provides a potential explanation for increased cancer incidence in PJS
patients. The presented models may provide a starting point to uncover how additional mutations and
environmental stressors interact with *LKB1* mutations and how stromal and immune components contribute
to malignant growth of the PJS epithelium. Ultimately, in-depth knowledge on the pathways and processes
involved in driving PJS epithelia into a pre-malignant state may help to design strategies that re-establish
epithelial homeostasis and reduce the incidence of cancer development in PJS patients.

References

1. Beggs AD, Latchford AR, Vasen HFA, et al. Peutz-Jeghers syndrome: a systematic review and recommendations for management. Gut 2010;59:975–986.

2. Lier MGF Van, Wagner A, Mathus-Vliegen EMH, et al. High cancer risk in peutz-jeghers syndrome: A systematic review and surveillance recommendations. Am J Gastroenterol 2010;105:1258–1264.

3. Giardiello FM, Brensinger JD, Tersmette AC, et al. Very high risk of cancer in familial Peutz-Jeghers syndrome. Gastroenterology 2000;119:1447–1453.

4. Tomasetti C, Vogelstein B. Variation in cancer risk among tissues can be explained by the number of stem cell divisions. Science (80-) 2015;347:78–81.

5. Hemminki A, Markie D, Tomlinson I, et al. A serine/threonine kinase gene defective in Peutz–Jeghers syndrome. Nat 1998 3916663 1998;391:184–187.

6. Sanchez-Cespedes M. A role for LKB1 gene in human cancer beyond the Peutz-Jeghers syndrome. Oncogene 2007;26:7825–7832.

7. Giardiello FM, Trimbath JD. Peutz-Jeghers syndrome and management recommendations. Clin Gastroenterol Hepatol 2006;4:408–415.

8. **Pierce SE, Granja JM**, Corces MR, et al. LKB1 inactivation modulates chromatin accessibility to drive metastatic progression. Nat Cell Biol 2021;23:915–924.

9. Kline ER, Shupe J, Gilbert-Ross M, et al. LKB1 Represses Focal Adhesion Kinase (FAK) Signaling via a FAK-LKB1 Complex to Regulate FAK Site Maturation and Directional Persistence. J Biol Chem 2013;288:17663.

10. Shackelford DB, Shaw RJ. The LKB1-AMPK pathway: Metabolism and growth control in tumour suppression. Nat Rev Cancer 2009;9:563–575.

11. Baas AF, Kuipers J, Wel NN Van Der, et al. Complete Polarization of Single Intestinal Epithelial Cells upon Activation of LKB1 by STRAD. Cell 2004;116:457–466.

12. **Entius MM, Keller JJ**, Westerman AM, et al. Molecular genetic alterations in hamartomatous polyps and carcinomas of patients with Peutz-Jeghers syndrome. J Clin Pathol 2001;54:126–131.

13. Leng WWJ De, Westerman AM, Weterman MAJ, et al. Cyclooxygenase 2 expression and molecular alterations in Peutz-Jeghers hamartomas and carcinomas. Clin Cancer Res 2003;9:3065–72.

14. Cotton JL, Dang K, Hu L, et al. PTEN and LKB1 are differentially required in Gli1-expressing mesenchymal cells to suppress gastrointestinal polyposis. Cell Rep 2022;40:111125.

15. Poffenberger MC, Metcalfe-Roach A, Aguilar E, et al. LKB1 deficiency in T cells promotes the development of gastrointestinal polyposis. Science (80-) 2018;361:406–411.

16. Ollila S, Domènech-Moreno E, Laajanen K, et al. Stromal Lkb1 deficiency leads to gastrointestinal tumorigenesis involving the IL-11-JAK/STAT3 pathway. J Clin Invest 2018;128:402–414.

17. Katajisto P, Vaahtomeri K, Ekman N, et al. LKB1 signaling in mesenchymal cells required for
suppression of gastrointestinal polyposis. Nat Genet 2008;40:455–459.

18. Korsse SE, Biermann K, Offerhaus GJA, et al. Identification of molecular alterations in gastrointestinal
carcinomas and dysplastic hamartomas in Peutz-Jeghers syndrome. Carcinogenesis 2013;34:1611–
1619.

19. Fodde R. The APC gene in colorectal cancer. Eur J Cancer 2002;38:867–871.

20. **Shorning BY, Zabkiewicz J**, McCarthy A, et al. Lkb1 deficiency alters goblet and paneth cell
differentiation in the small intestine. Morty RE, ed. PLoS One 2009;4:e4264.

21. **Gao Y, Yan Y**, Tripathi S, et al. LKB1 Represses ATOH1 via PDK4 and Energy Metabolism and
Regulates Intestinal Stem Cell Fate. Gastroenterology 2020;158:1401.

22. Sato T, Vries RG, Snippert HJ, et al. Single Lgr5 stem cells build crypt-villus structures in vitro without
a mesenchymal niche. Nature 2009;459:262–5.

23. **Wetering M van de, Francies HE, Francis JM**, et al. Prospective Derivation of a Living Organoid
Biobank of Colorectal Cancer Patients. Cell 2015;161:933–945.

24. **Fujii M, Matano M**, Nanki K, et al. Efficient genetic engineering of human intestinal organoids using
electroporation. Nat Protoc 2015 1010 2015;10:1474–1485.

25. **Muraro MJ, Dharmadhikari G**, Grün D, et al. A Single-Cell Transcriptome Atlas of the Human
Pancreas. Cell Syst 2016;3:385.

26. **Pleguezuelos-Manzano C, Puschhof J**, Brink S van den, et al. Establishment and Culture of Human
Intestinal Organoids Derived from Adult Stem Cells. Curr Protoc Immunol 2020;130.

27. **Merenda A, Fenderico N**, Maurice MM. Wnt Signaling in 3D: Recent Advances in the Applications of
Intestinal Organoids Trends in Cell Biology. Trends Cell Biol 2020;30.

28. Krotenberg Garcia A, Fumagalli A, Le HQ, et al. Active elimination of intestinal cells drives oncogenic
growth in organoids. Cell Rep 2021;36:109307.

29. Troughton WD, Trier JS. Paneth and goblet cell renewal in mouse duodenal crypts. J Cell Biol
1969;41:251–268.

30. **Haber AL, Biton M, Rogel N**, et al. A single-cell survey of the small intestinal epithelium. Nature
2017;551:333–339.

31. Mustata RC, Vasile G, Fernandez-Vallone V, et al. Identification of Lgr5-Independent Spheroid-
Generating Progenitors of the Mouse Fetal Intestinal Epithelium. Cell Rep 2013;5:421–432.

32. **Yui S, Azzolin L**, Maimets M, et al. YAP/TAZ-Dependent Reprogramming of Colonic Epithelium
Links ECM Remodeling to Tissue Regeneration. Cell Stem Cell 2018;22:35-49.e7.

33. **Nusse YM, Savage AK**, Marangoni P, et al. Parasitic helminths induce fetal-like reversion in the
intestinal stem cell niche. Nature 2018;559:109–113.

34. Ayyaz A, Kumar S, Sangiorgi B, et al. Single-cell transcriptomes of the regenerating intestine reveal a
revival stem cell. Nature 2019;569:121–125.

35. **Kobayashi S, Ogasawara N**, Watanabe S, et al. Collagen type I-mediated mechanotransduction
controls epithelial cell fate conversion during intestinal inflammation. Inflamm Regen 2022;42:49.

36. **Qu M, Xiong L, Lyu Y**, et al. Establishment of intestinal organoid cultures modeling injury-associated
epithelial regeneration. Cell Res 2021;31:259–271.

37. Gregorieff A, Liu Y, Inanlou MR, et al. Yap-dependent reprogramming of Lgr5+ stem cells drives
intestinal regeneration and cancer. Nature 2015;526:715–718.

38. Mohseni M, Sun J, Lau A, et al. A genetic screen identifies an LKB1-MARK signalling axis controlling
the Hippo-YAP pathway. Nat Cell Biol 2014;16:108–117.

39. **Pikkupeura LM, Bressan RB, Guiu J**, et al. Transcriptional and epigenomic profiling identifies YAP
signaling as a key regulator of intestinal epithelium maturation. 2023.

40. **Serra D, Mayr U, Boni A**, et al. Self-organization and symmetry breaking in intestinal organoid
development. Nature 2019;569:66–72.

41. Zhang J, Ji J-Y, Yu M, et al. YAP-dependent induction of amphiregulin identifies a non-cell-
autonomous component of the Hippo pathway. Nat Cell Biol 2009.

42. Heuberger J, Kosel F, Qi J, et al. Shp2/MAPK signaling controls goblet/paneth cell fate decisions in
the intestine. Proc Natl Acad Sci 2014;111:3472–3477.

43. Wang LX, Zhu F, Li JZ, et al. Epidermal growth factor promotes intestinal secretory cell differentiation
in weaning piglets via Wnt/β-catenin signalling. Animal 2020;14:790–798.

44. Hanahan D, Weinberg RA. Hallmarks of Cancer: The Next Generation. Cell 2011;144:646–674.

45. **Malagola E, Vasciaveo A**, Ochiai Y, et al. Isthmus progenitor cells contribute to homeostatic cellular
turnover and support regeneration following intestinal injury. Cell 2024;187:3056-3071.e17.

46. **Chen B, Scurrah CR**, McKinley ET, et al. Differential pre-malignant programs and microenvironment
chart distinct paths to malignancy in human colorectal polyps. Cell 2021;184:6262-6280.e26.

47. Aiderus A, Barker N, Tergaonkar V. Serrated colorectal cancer: preclinical models and molecular
pathways. Trends in Cancer 2024;10:76–91.

48. Joanito I, Wirapati P, Zhao N, et al. Single-cell and bulk transcriptome sequencing identifies two
epithelial tumor cell states and refines the consensus molecular classification of colorectal cancer.
Nat Genet 2022;54:963–975.

49. **Wang Y, Song W**, Wang J, et al. Single-cell transcriptome analysis reveals differential nutrient
absorption functions in human intestine. J Exp Med 2020;217.

50. **Sousa E Melo F De, Wang X**, Jansen M, et al. Poor-prognosis colon cancer is defined by a
molecularly distinct subtype and develops from serrated precursor lesions. Nat Med 2013;19:614–

618.

51.   Tuppurainen K, Mäkinen JM, Junttila O, et al. Morphology and microsatellite instability in sporadic
serrated and non-serrated colorectal cancer. J Pathol J Pathol 2005;207:285–294.

52.   **Collet L, Ghurburrun E**, Meyers N, et al. Kras and Lkb1 mutations synergistically induce intraductal
papillary mucinous neoplasm derived from pancreatic duct cells. Gut 2020;69:704–714.

53.   Osoegawa A, Kometani T, Nosaki K, et al. LKB1 Mutations Frequently Detected in Mucinous
Bronchioloalveolar Carcinoma. Jpn J Clin Oncol 2011;41:1132–1137.

54.   Jiang Z, White RA, Wang TC. Adult Pancreatic Acinar Progenitor-like Populations in Regeneration
and Cancer. Trends Mol Med 2020;26:758–767.

55.   Zou G, Park J-I. Wnt signaling in liver regeneration, disease, and cancer. Clin Mol Hepatol
2023;29:33–50.

56.   Greten FR, Grivennikov SI. Inflammation and Cancer: Triggers, Mechanisms, and Consequences.
Immunity 2019;51:27–41.

57.   **Li J, Wang R, Zhou X**, et al. Genomic and transcriptomic profiling of carcinogenesis in patients with
familial adenomatous polyposis. Gut 2020;69:1283–1293.

58.   IJspeert JEG, Dekker E. Gastric metaplasia could initiate the serrated neoplasia pathway in CRC. Nat
Rev Gastroenterol Hepatol 2022;19:217–218.

Figure legends

Figure 1. *Lkb1*-mutant organoids display alterations in phenotype and reconstitution capacity.
A) Sequences of *Lkb1*-mutant organoids obtained with gRNA1 showing indels/deletions detected in exon 1 of *Lkb1*. *Lkb1*[+/-] shows two amino acids deletion outside essential Lkb1 domains. PAM sequences are underlined. **B-D)** Western blot of Lkb1 **(B)**, (p)AMPK **(C)**, and 4E-BP1 **(D)** in WT and *Lkb1*-mutant organoids. **E)** Brightfield images of *Lkb1*-mutant organoid clones. Scale bar = 100 µM. **F)** Immunofluorescent images of cleaved caspase-3 (CC3) and F-actin in *Lkb1*-mutant and WT organoids. Intraluminal (red arrowheads) and extraluminal CC3[+] cells (orange arrowheads) are indicated. Scale bar = 50 µm. **G)** Organoid reconstitution assay from single cells. One-way ANOVA (Dunnett) was applied. ** = p < 0.05. N* = 16.

Figure 2. *Lkb1* loss mediates alterations in the number, location and ultrastructural morphology of secretory cells.
A) Immunofluorescence staining of Lysozyme1 (Lyz1) and F-actin in WT and *Lkb1*-mutant small intestinal organoids. Scale bar = 50 µm. White arrows indicate mislocalized Lyz1[+] cells. **B)** Immunohistochemistry staining of Mucin 2 (Muc2) in WT and *Lkb1*-mutant organoids. Scale bar = 100 µm. **C&E)** Immunohistochemistry staining of MUC2 **(C)** and LYZ **(E)** in healthy and non-transformed PJS epithelium. Orange arrowheads indicate mislocalized LYZ[+] cells. Scale bar = 100 µm. Shown images are zoom-in of Figure S3B-C. **D&F)** The average position and number of MUC2+ cells **(D)** and LYZ+ cells **(F)** in healthy and PJS epithelia. Each dot represents a single crypt. Unpaired t-test was applied. *ns = not significant, **** = p < 0.0001*. **G-H)** Electron microscopy images Paneth **(G)** and goblet **(H)** cells in WT and *Lkb1*-mutant organoids. Scale bar upper panel = 5 µm, lower panel = 1 µm.

Figure 3. *Lkb1* loss mediates increased expression of a regenerative gene signature in the intestinal epithelium.
A) GSEA of *Lkb1*-mutant versus WT organoids for cell types[30], colitis[32], injury[36], parasite[33], fetal spheroid[31], collagen[35] and revival stem cell[34] gene sets. Heatmap displays normalized enrichment scores (NES). ns = not significant, * = FDR < 0.05, ** = FDR ≤ 0.01, *** = FDR ≤ 0.001, **** = FDR ≤ 0.0001. **B)** Heatmap of z-score transformed expression values of *Lgr5*[+] stem cell genes (Stem)[30] enriched in WT organoids and fetal genes (Fetal)[31] enriched in *Lkb1*-mutant organoids. Marker genes are highlighted. N = 3. **C)** Immunofluorescence staining of Anxa1, DAPI and F-actin in WT and *Lkb1*-mutant organoids. Fetal organoids from E16.5 mice are shown as a positive control. Scale bar = 50 µm**.**

Figure 4. *Lkb1* loss accommodates Yap pathway activation and mediates upregulation of EGF family receptors and their ligands to drive niche independency.
A) GSEA of *Lkb1*-mutant versus WT organoids for Yap gene sets[37,39]. Heatmap displays normalized enrichment scores (NES). ns = not significant, * = FDR < 0.05, *** = FDR ≤ 0.001, **** = FDR ≤ 0.0001. **B)** Immunofluorescence staining for Yap, DAPI and F-actin in WT and *Lkb1*-mutant organoids. Scale bar = 172 µm. **C)** Quantification of the nuclear fraction of Yap. *N* = 13-19. One-way ANOVA was applied. **D-E)** Cumulative normalized counts of *Egfr* ligands **(D)** and *Egf* family receptors **(E)** in WT and *Lkb1*-mutant organoids. One-way ANOVA was applied. *N* = 3. **F)** Brightfield images of WT and *Lkb1*-mutant organoids

grown in the presence (ENR) and absence (NR) of Egf. Scale bar = 500 µm. **G)** Surface area covered by WT

and *Lkb1*-mutant organoids grown in ENR or NR. *N* = 2-3. Two-way ANOVA was applied. **H)** Brightfield

images of WT and *Lkb1*-mutant organoids cultured in NR and treated with DMSO or 100 nM Gefitinib. Scale

bar = 500 µm. * = *p < 0.05,* and **** = *p < 0.0001.*

Figure 5. *Lkb1* loss modifies intestinal stem cell identity.

A-B) UMAP of scRNA-seq data from WT Adult, WT fetal and *Lkb1*-mutant organoids. Colors indicate the

different organoids lines **(A)** and the different cell types **(B)**. **C-D)** UMAP of scRNA-seq data from reclustered

WT Adult, *Lkb1*[+/-] and *Lkb1*[bud-/-] organoids. Colors indicate the different organoids lines **(C)** and the different

cell types **(D)**. **E)** Percentage of cell types for each organoid line. **F)** Scatter plot showing module scores for

Paneth[30] and goblet[30] cell signatures across WT Adult, *Lkb1*[+/-] and *Lkb1*[bud-/-] cells. Color indicates the origin

of the cell. **G)** Expression of *Anxa1* onto the UMAP plot. **H)** GSEA on genes differentially expressed between

WT and *Lkb1*-mutant stem cells for cell types[30], colitis[32], embryonic spheroid[31], Yap regeneration[37], Yap

overexpression[39] and revival stem cell[34] gene sets. Heatmap displays normalized enrichment scores (NES).

ns = not significant, * = FDR < 0.05, *** = FDR ≤ 0.001, **** = FDR ≤ 0.0001. **I-J)** Violin plot showing module

scores for goblet cell gene signature[30] **(I)** or *Anxa1* **(J)** across WT, *Lkb1*[+/-] and *Lkb1*[bud-/-] stem cells. ns = not

significant, **** = p ≤ 0.0001 using one-way ANOVA with Wilcoxon test.

Figure 6. *Lkb1* loss drives a pre-malignant program along the serrated pathway that is further

amplified by *Kras* mutations.

A) GSEA of *Lkb1*-mutant versus WT organoids for serrated-specific (SSC)[46], adenoma-specific (ASC)[46],

iCMS2[48] and iCMS3[48] genesets. Heatmap displays normalized enrichment scores (NES). ns = not

significant, ** = FDR < 0.01, *** = FDR ≤ 0.001, **** = FDR ≤ 0.0001. **B)** Heatmap of z-score transformed

expression values of ASC[46] genes enriched in WT organoids and SSC[46] genes enriched in *Lkb1*-mutant

organoids. N = 3. **C)** Boxplot showing log2 transformed *LKB1* expression in pseudobulk samples of pre-

malignant adenoma (AD) and sessile serrated lesions (SSL)[46]. **D-E)** Organoid reconstitution assay from

single cells for *Kras*-mutant and *Kras*-mutant/*Lkb1*-mutant organoids **(D)** or *Apc*-deficient and *Apc/Lkb1*-

mutant organoids **(E)**. *N* = 8 **F-G)** Gene Set Variation Analysis (GSVA) analysis of SSC[46] genes **(F)** and

ASC[46] genes **(G)** for WT, *Apc* mutant, *Kras* mutant, *Lkb1*-mutant, *Apc/Lkb1* double mutant and *Kras/Lkb1*

double mutant organoids. *N* = 3. One-way ANOVA with Wilcoxon test was applied. *ns = not significant, * = p*

< 0.05, ** = *p < 0.01* and **** = *p < 0.0001.*

Figure 7. *LKB1* loss phenotypes are conserved in human colon epithelium and sporadic CRC.

A) CRISPR/Cas9-induced sequence alterations in exon 1 of the *LKB1* gene in human colon organoids

(HCOs). *LKB1*[+/-] HCOs acquired a two amino acid deletion in one allele, located outside known functional

LKB1 domains. PAM sequences are underlined. **B)** Brightfield images of WT and LKB1-mutant HCOs. Scale

bar = 500 µm. **C-D)** Western blot of LKB1 **(C)** and (p)AMPK **(D)** in WT and *LKB1*-mutant HCOs. **E)** Gene Set

Variation Analysis (GSVA) analysis of signatures[31–34,37,39,49] enriched in *LKB1*-mutant HCOs. *N* = 3-6. One-

way ANOVA with Wilcoxon test was applied. *ns = not significant, ** = p < 0.05, ** = p < 0.01, *** = p < 0.001*

and **** = *p < 0.0001.* **F-G)** Cumulative normalized counts of EGF receptors **(F)** and EGF ligands **(G)** in WT

and *LKB1*-mutant HCOs. One-way ANOVA was applied. *N* = 3-6. **H)** Brightfield images of WT and *LKB1*-

mutant HCOs grown in the absence of EGF. Scale bar = 500 μm. **I)** Single cell outgrowth assay with and

without EGF. One-way ANOVA (Dunnett) was applied. N = 3. **J)** Heatmap of z-score transformed expression

values of ASC[46] genes enriched in WT organoids and SSC[46] genes enriched in *LKB1*[-/-] organoids. N = 3-6.

K-M) Box plots showing log2 Z-scores for the *LKB1*-mutant signature over the bulk RNA sequencing

datasets from Sousa E Melo *et al*[50]. **(K)** or TCGA **(L-M)** grouped as tubular (N = 7) or serrated (N = 6) **(K)**,

mucinous (N = 68) or non-mucinous (N = 437) **(L)** or consensus molecular subtype (CMS1; N = 63, CMS2; N

= 125, CMS3; N = 45, CMS 4; N = 73) **(M)** Welch's unequal variances t-test or one-way ANOVA with

Wilcoxon test was applied. *ns = not significant, * = p < 0.05, ** = p < 0.01 and **** = p < 0.0001*.

A

Lkb1 WT
CCAGCCGCGC---CGCAAACGCGCCAAGCTCAT
GGTCGGCGCG--GCGTTTGCGCGGTTCGAGTA

Lkb1+/-
(p.R40PfsX119)   +1
CCAGCCGCGCC--CGCAAACGCGCCAAGCTCAT

Lkb1+/-
(p.P38_R39del)   -6
CCAG------CGCAAACGCGCCAAGCTCAT

Lkb1+/-

Lkb1 WT
CCAGCCGCGC---CGCAAACGCGCCAAGCTCAT
GGTCGGCGCG--GCGTTTGCGCGGTTCGAGTA

Lkb1bud-/-
(p.R40TfsX49)   +1
CCAGCCGCGC---CACAAACGCGCCAAGCTCAC

Lkb1bud-/-
(p.R40PfsX119)   +1
CCAGCCGCGCC--CGCAAACGCGCCAAGCTCAT

Lkb1bud-/-

Lkb1 WT
CCAGCCGCGC---CGCAAACGCGCCAAGCTCAT
GGTCGGCGCG--GCGTTTGCGCGGTTCGAGTA

Lkb1cys-/-
(p.P40PfsX119)   +1
CCAGCCGCGCC--CGCAAACGCGCCAAGCTCAT

Lkb1cys-/-
(p.K41AfsX50)   +2
CCAGCCGCGCG--CGCAAACGCGCCAAGCTCAT

Lkb1cys-/-

B

Lkb1
Gapdh

WT, Lkb1+/-, Lkb1bud-/-, Lkb1cys-/-

C

pAmpk
Ampk
Actin

WT, Lkb1+/-, Lkb1bud-/-, Lkb1cys-/-

D

4E-BP1
Actin

WT, Lkb1+/-, Lkb1bud-/-, Lkb1cys-/-

E

WT | Lkb1+/- | Lkb1bud-/- | Lkb1cys-/-

F

Cleaved caspase 3
F-actin

Max. projection | Single plane

WT | Lkb1+/- | Lkb1bud-/-

G

Luminescence (RLU)

WT | Lkb1+/- | Lkb1bud-/- | Lkb1cys-/-

A

WT | Lkb1^+/- | Lkb1^bud-/- | Lkb1^cys-/-

Lyz1
F-actin

B

WT | Lkb1^+/- | Lkb1^bud-/- | Lkb1^cys-/-

Muc2

C

Healthy donor | PJS non-transformed

MUC2

D

MUC2^+cells/total cells

ns

Healthy PJS

MUC2^+ position/total MUC2^+ cells

ns

Healthy PJS

E

Healthy donor | PJS non-transformed

LYZ

F

LYZ^+ cells/total cells

ns

Healthy PJS

Average LYZ+ position

****

Healthy PJS

G

Paneth cells

WT

Lkb1^+/-

Lkb1^bud-/-

H

Goblet cells

WT

Lkb1^+/-

Lkb1^bud-/-

Lkb1^cys-/-

A

	Lkb1$^{+/-}$ vs. WT	Lkb1$^{bud\ -/-}$ vs. WT
Yap regeneration[37]	****	****
Yap overexpression[39]	****	****

NES

WT ◄ -2 — 0 — 2 ► *Lkb1*-deficient

B

	WT	Lkb1$^{+/-}$	Lkb1$^{bud\ -/-}$
Yap – zoom			
Yap DAPI F-actin			

C Yap quantification

Fold increase (nuclear fraction Yap)

ns, ****

WT, Lkb1$^{+/-}$, Lkb1$^{bud\ -/-}$

D Egf family ligands

Cumulative normalized counts

****, *

WT, Lkb1$^{+/-}$, Lkb1$^{bud\ -/-}$

E Egf family receptors

Cumulative normalized counts

****, ****

WT, Lkb1$^{+/-}$, Lkb1$^{bud\ -/-}$

F

	+ Egf	– Egf
WT		
Lkb1$^{+/-}$		
Lkb1$^{bud\ -/-}$		

G Quantification Egf depletion

Surface area organoids

*, ns, ns

WT ENR, WT NR, Lkb1$^{+/-}$ ENR, Lkb1$^{+/-}$ NR, Lkb1$^{bud\ -/-}$ ENR, Lkb1$^{bud\ -/-}$ NR

H

	DMSO	Gefitinib
WT		
Lkb1$^{+/-}$		
Lkb1$^{bud\ -/-}$		

A

Lkb1 WT
TTATGCCGCAGGGCGGTCAAGATCCTCAAGAAG
AATACGGCGTCCCGCCAGTTCTAGGAGTTCTTC

Lkb1cys -/- (Cl.5)
(p.A76RfsX110)
TTATGCCGCA-GGCGGTCAAGATCCTCAAGAAG

Lkb1bud -/- (Cl.10)
(p.A76RfsX110)
TTATGCCGCA-GGCGGTCAAGATCCTCAAGAAG

Lkb1bud -/- (Cl.11)
(p.A76RfsX110)
TTATGCCGCA-GGCGGTCAAGATCCTCAAGAAG

Lkb1cys -/- (Cl.13)
(p.A76RfsX110)
TTATGCCGCA-GGCGGTCAAGATCCTCAAGAAG

B

	WT	Lkb1cys -/- (Cl.5)	Lkb1bud -/- (Cl.10)	Lkb1bud -/- (Cl.11)	Lkb1cys -/- (Cl.13)
Lkb1					
Gapdh					

C

Luminescence (RLU)

ns
ns
*
**

WT, Lkb1bud -/- (Cl.10), Lkb1bud -/- (Cl.11), Lkb1cys -/- (Cl.5), Lkb1cys -/- (Cl.13)

D

WT	Lkb1cys -/- (Cl.5)	Lkb1bud -/- (Cl.10)	Lkb1bud -/- (Cl.11)	Lkb1cys -/- (Cl.13)

A

	WT	Lkb1+/-	Lkb1bud-/-
Max. projection			
First plane			
Middle planes			
Last plane			

Lyz1
F-actin

B MUC2

Healthy donor

PJS non-transformed

C LYZ

Healthy donor

PJS non-transformed

D

WT	Lkb1+/-	Lkb1bud-/-	Lkb1cys-/-

Figure S5

A

Lkb1$^{cys-/-}$ vs. Lkb1$^{bud-/-}$

Yap regneration[30] ****
Yap overexpression[38] ****

NES

-4 -2 0 2 4

Lkb1$^{bud-/-}$ *Lkb1$^{cys-/-}$*

B

Yap DAPI F-actin | Yap - zoom | Yap DAPI F-actin

Lkb1$^{cys-/-}$

C

WT | Lkb1$^{+/-}$ | Lkb1$^{bud-/-}$

Yap DAPI F-actin

D

Yap quantification

****

Fold increase (nuclear fraction YAP)

Lkb1$^{bud-/-}$ Lkb1$^{cys-/-}$

E

Egf family ligands

***

Cumulative normalized counts

Lkb1$^{bud-/-}$ Lkb1$^{cys-/-}$

F

Egf family receptors

****

Cumulative normalized counts

Lkb1$^{bud-/-}$ Lkb1$^{cys-/-}$

G

Lkb1$^{cys-/-}$

- Egf

+ Egf

H

Quantification Egf depletion

ns

Surface area organoids

Lkb1$^{cys-/-}$ ENR   Lkb1$^{cys-/-}$ NR

I

Lkb1$^{cys-/-}$

Gefinitib

DMSO