Intestinal LKB1 loss drives a pre-malignant program along the serrated cancer

pathway

Plugge S.F.[#,1], Ma H.[#,1,2], van der Vaart J.Y.[1], Sprangers J.[1], Morsink F.H.M.[2], Xanthakis D.[1], Jamieson C.[1],

Keijzer A.R.[1], Margaritis T.[3]., Candelli T.[3], Straver R.[1], de Ridder J.[1], Holstege F.C.P.[3, 4], de Leng W.W.J.[2],

Offerhaus G.J.A[2], Merenda A.*[,1], Maurice M.M.*,[1]

1. Center for Molecular Medicine and Oncode Institute, University Medical Center Utrecht, Utrecht, The

Netherlands

2. Department of Pathology, University Medical Center Utrecht, Utrecht, The Netherlands

3. Princess Maxima Center for Pediatric Oncology, Utrecht, Utrecht, The Netherlands

4. Center for Molecular Medicine, University Medical Center Utrecht, Utrecht, The Netherlands

[#]These authors contributed equally

*Corresponding authors

Abstract

Peutz-Jeghers syndrome (PJS) is a familial disorder caused by heterozygous inactivating Liver Kinase B1

(*LKB1)* mutations that promote gastrointestinal polyposis and enhance cancer susceptibility. How *LKB1*-

deficiency alters the phenotypical landscape and hierarchical organization of epithelial tissues to mediate

increased cancer risk remains poorly understood. Here, we employ small intestinal organoids to investigate

these issues for heterozygous and homozygous *Lkb1* epithelial loss. We show that *Lkb1* loss causes an

allele dosage-dependent activation of a transcriptional program for tissue repair that is already induced in

stem cell populations and mediates alterations in secretory cell type morphology and positioning.

Furthermore, this shift towards a regenerative state omits the need for EGF supplementation, thus inducing

niche-independent properties. Strikingly, we uncover that heterozygous loss of *Lkb1* is sufficient to push the

epithelium into a premalignant program for colorectal carcinogenesis along the serrated pathway, which is

further amplified by loss-of-heterozygosity (LOH) or Kras mutations. We conclude that persistent

upregulation of a regenerative program due to *LKB1* loss alters cellular hierarchy and induces niche

independency, which predisposes PJS epithelium to uncontrolled growth along the serrated pathway.

Introduction

The Peutz-Jeghers syndrome (PJS) is a hereditary genetic disorder that occurs in approximately 1:50.000 -

1:200.000 individuals worldwide (Beggs *et al.*, 2010). PJS patients suffer from a life-long development of

benign hamartomatous polyps throughout their gastrointestinal tract for which they need periodical surgical

resection. Moreover, patients carry an increased risk of developing early-onset epithelial tumors, particularly

within the gastrointestinal (GI) tract (up to 66%), breast (up to 54%), pancreas (up to 36%) and gonads

(Giardiello *et al.*, 2000; Hearle *et al.*, 2006; Beggs *et al.*, 2010; Van Lier *et al.*, 2010a), resulting in a

cumulative lifetime risk of 93% (Giardiello *et al.*, 2000). Of note, PJS patients are predisposed to

malignancies of the small intestine (frequency 13%), a condition that is otherwise very rare (0.03%) (Van Lier

et al., 2010a; Tomasetti and Vogelstein, 2015).

Heterozygous inactivating mutations in the tumor suppressor gene Liver Kinase B1 (*LKB1*) were
identified as the main cause of PJS (Hemminki *et al.*, 1998; Jenne *et al.*, 1998). Furthermore, *LKB1*
mutations are causally linked to a broad range of sporadic cancer cases, including lung carcinoma,
melanoma, pancreatic and cervical cancers (Giardiello and Trimbath, 2006; Sanchez-Cespedes, 2007).
LKB1 encodes a serine/threonine kinase implicated in the regulation of a variety of critical cellular processes,
including energy metabolism, cell adhesion, DNA methylation and apicobasal polarity (Shackelford and
Shaw, 2009; Kline *et al.*, 2013; Pierce *et al.*, 2021). Loss of heterozygosity (LOH) in this gene is regarded as
a driving force of cancer development, as over 64% of PJS-derived carcinomas present with inactivation of
the second copy of the gene (Entius *et al.*, 2001; De Leng *et al.*, 2003).

Over recent years, modelling of PJS in murine models revealed that *Lkb1* loss in either
mesenchymal or T cell lineages induces development of gastro-intestinal polyps with strong resemblance to
those observed in PJS patients. (Katajisto *et al.*, 2008; Ollila *et al.*, 2018; Poffenberger *et al.*, 2018; Cotton *et
al.*, 2022). These results have shifted focus to non-epithelial roles of *Lkb1* in driving PJS disease
development, with an emphasis on processes underlying polyposis. If and how polyp formation in PJS
patients links to cancer predisposition however remains a heavily debated issue, as LOH and dysplastic
transformation of PJS polyps were reported to be very rare (Shepherd, Bussey and Jass, 1987; Entius *et al.*,
2001; De Leng *et al.*, 2003; Korsse *et al.*, 2013). These observations differ from other polyposis syndromes,
such as familial adenomatous polyposis (FAP), where epithelial loss of the underlying germline mutation in
the adenomatous polyposis coli gene APC and its role in cancer development has been well described
(Fodde, 2002).

Since *LKB1* loss mediates predisposition to carcinoma development in multiple epithelial tissues
irrespective of polyp formation, we hypothesized that epithelial-intrinsic alterations may sensitize these
tissues to cancer formation. Previous studies indicate that intestinal epithelial loss of *LKB1* does not lead to
polyp formation while alterations in secretory cell fate specification were identified (B Y Shorning *et al.*, 2009;
Gao *et al.*, 2020; Cotton *et al.*, 2022; Kim *et al.*, 2022). Despite these advancements, how epithelial *LKB1*
loss changes the phenotypic landscape of the intestinal epithelium and how these alterations may link to
cancer predisposition in PJS patients remains poorly understood.

Here, we model *mono-* and *bi-*allelic inactivation of *Lkb1* in small intestinal organoids to investigate
how *Lkb1* loss alters the dynamics of cellular identities and lineage relationships of the intestinal epithelium.
We uncover that *Lkb1* loss promotes features of epithelial transformation linked to the serrated pathway for
intestinal carcinogenesis. These features include loss of classical adult stem cell populations, activation of
transcriptional programs related to regeneration and intestinal-to-gastric cellular transdifferentation, as well
as the acquirement of niche-independent growth properties. Our findings argue that *Lkb1* loss mediates
epithelium-wide activation of a pre-cancer program akin to sporadic serrated precursor lesions, providing an
explanation for the increased cancer risk observed in PJS patients. By linking *Lkb1* loss to a major pathway
for colorectal cancer development, these findings pave a path for the development of new strategies for
prevention, surveillance and precision treatment of PJS tumors as well as other malignancies harboring
LKB1 inactivating mutations.

Results

***Lkb1* bi-allelic loss results in phenotypic heterogeneity and increased self-renewal potential of small**

intestinal organoids.

To study the role of *Lkb1* loss in the intestinal epithelium, we employed CRISPR/Cas9 technology to

generate *Lkb1*-deficient mouse small intestinal organoid lines. To obtain homogenous organoid cultures that

are heterozygous or homozygous for *Lkb1* loss, we isolated single organoids from bulk cultures and

expanded these as individual clones. After genotyping, we selected one *Lkb1$^{+/-}$* and two *Lkb1$^{-/-}$* clonal lines

for further characterization (Figure S1A, Figure 1A-B). *Lkb1$^{+/-}$* organoids were morphologically similar to wild

type (WT) organoids, with typical budding structures that reflect a crypt-villus organization (Sato *et al.*, 2009)

(Figure 1A). Remarkably, the introduction of homozygous *Lkb1* loss resulted in phenotypic bifurcation, with

clonal lines displaying either a budding or a cystic growth morphology (Figure 1A). Notably, from day 3 after

passaging, budding *Lkb1$^{-/-}$* (*Lkb1$^{bud\ -/-}$*) organoids accumulated clusters of dead cells in their periphery (Figure

1A), as indicated by their positive staining for cleaved caspase 3 (Figure S1B). These observations are

different from *Lkb1$^{+/-}$* and WT organoids that mainly accumulate dead cells in their lumen (Figure 1SB) (Sato

et al., 2009).

Next, we tested whether *Lkb1* loss alters the ability of intestinal organoids to reconstitute organoids

from single cells, as a proxy for their self-renewal capacity. While budding WT and *Lkb1*-mutant organoids

lines displayed similar outgrowth rates, cystic *Lkb1$^{-/-}$* (*Lkb1$^{cys\ -/-}$*) organoids carried a significantly improved

capacity to form organoids upon seeding of single cells (Figure 1C). Regardless of these observations, both

homozygous mutant morphologies displayed a similar increase in the number of proliferative cells in

comparison to heterozygous and wild-type organoids, as shown by EdU incorporation (Figure 1D, E). To rule

out possible CRISPR off-target effects, we used an alternative gRNA to introduce *Lkb1* mutations, which

resulted in an identical phenotypic heterogeneity of *Lkb1$^{-/-}$* organoid lines (Figure S1C-E). Moreover, these

newly generated *Lkb1$^{cys\ -/-}$* clones also displayed superior reconstitution potential compared to their

homozygous budding counterparts (Figure S1F). These findings reveal that biallelic *Lkb1* loss results in

phenotypic heterogeneity within the intestinal epithelium, and that homozygous *Lkb1*-mutant organoids with

cystic morphology display an increased recovery potential.

***Lkb1* loss alters the number, morphology and positioning of secretory cells.**

To investigate whether our *Lkb1*-mutant organoid lines serve as accurate models for intestinal epithelial loss

of *Lkb1 in vivo*, we examined alterations in localization and morphology of Paneth (Lyz$^+$) and goblet (Muc2$^+$)

cells that were previously reported for the *Lkb1*-mutant mouse intestine (Boris Y. Shorning *et al.*, 2009; Gao

et al., 2020). Compared to WT organoids, the localization of Lyz1$^+$ Paneth cells in *Lkb1$^{+/-}$* and *Lkb1$^{bud\ -/-}$*

organoids was not restricted to the bottom of the crypt but displayed a more scattered distribution that

reached into the villus domain (Figure 2A). In addition, while Muc2$^+$ cells in WT and *Lkb1$^{+/-}$* organoids were

present in modest numbers and always located to the villus region, Muc2$^+$ cell numbers were increased in

Lkb1$^{bud\ -/-}$ organoids and found scattered throughout the crypt-villus axis (Figure 2B). These observations are

highly reminiscent of the reported aberrancies in Paneth and Goblet cell numbers and localization in the

Lkb1-deficient mouse intestine *in vivo* (B Y Shorning *et al.*, 2009). By contrast, *Lkb1$^{cys\ -/-}$* organoids were

devoid of Paneth cells but did allow for the formation of Muc2$^+$ cells that are usually lacking in cystic

organoids (Sato, Es, *et al.*, 2011), indicating that secretory lineage specification is also altered within these

organoids (Figure 2A-B).

We next investigated how these findings relate to intestinal tissue organization of human PJS

patients. While the average number and location of MUC2$^+$ cells per intestinal crypt was similar for healthy

donors and PJS patients (Figure 2C, Figure S2A-B), LYZ1$^+$ Paneth cells were clearly mislocalized within the

PJS mucosa (Figure 2D). In the healthy intestine, Paneth cells localized strictly to the crypt base in the

healthy intestine, as expected (Barker *et al.*, 2007). In PJS epithelia, however, Paneth cells frequently

scattered towards higher regions of the crypt (Figure 2D). Thus, heterozygous loss of *LKB1* in PJS patient-

derived intestines leads to mislocalisation of Paneth cells but does not alter goblet cell distribution, similar to

our observations in *Lkb1$^{+/-}$* organoids.

To examine *Lkb1* loss-induced alterations in secretory cell morphology at the ultrastructural level, we

employed electron microscopy. In WT intestinal organoids, Paneth cells displayed a typical morphology with

prominent apical accumulation of spherically-shaped secretory granules that are uniformly filled with

electron-dense content (Figure 2E) (Satoh *et al.*, 1990; Sato *et al.*, 2009). By contrast, secretory granules of

Lkb1$^{+/-}$ Paneth cells commonly contained an abnormally 'sausage-shaped' electron-dense core (Figure 2E).

In *Lkb1$^{bud-/-}$* organoids, the content of Paneth cell-secretory granules was even more aberrant, displaying a

small electron-dense core surrounded by a peripheral electron-lucent halo (Figure 2E). In WT intestinal

organoids, columnar-shaped goblet cells were identified with large translucent granules filled with mucus

(Figure 2F) (Troughton and Trier, 1969; Bansil and Turner, 2018). In *Lkb1*-deficient clones (*Lkb1$^{+/-}$* and

Lkb1$^{bud-/-}$), mucus-secreting cells displayed a more round and flattened morphology compared to WT goblet

cells (Figure 2F). In addition, in *Lkb1$^{cys-/-}$* organoids the content of these mucus-like granules was frequently

lost during EM preparation, suggesting an altered mucin composition, which might also explain the limited

number of identified Muc2$^+$ cells (Figure 2F). Furthermore, previous studies identified roles of *LKB1* in apical

brush border formation, using 2D human (cancer) cell lines (Baas *et al.*, 2004). Inspection of our EM images

however did not reveal any evidence for loss of these subcellular structures in *Lkb1*-deficient organoids

(Figure S2C).

Taken together, our findings reveal that engineering *Lkb1* loss in organoids induces highly similar

alterations in secretory cell number, localization and morphology as those observed in *in vivo* mouse models

and the intestine of PJS patients. These results thus confirm the suitability and relevance of organoids for

modelling epithelial alterations in PJS. Furthermore, our results indicate that *Lkb1* loss-induced alterations in

cell lineage specification and positioning are intrinsic to the epithelium and do not rely on interactions with

the mesenchyme.

***Lkb1* mono- and bi-allelic loss induces incremental expression of a regenerative gene signature**

To examine how *Lkb1* loss affects epithelial organization, we determined transcriptional changes

induced by *Lkb1* loss in organoids using bulk mRNA sequencing. Compared to WT organoids, we observed

differential expression of 1298, 2312 and 7816 genes in *Lkb1$^{+/-}$*, *Lkb1$^{bud-/-}$* and *Lkb1$^{cys-/-}$* organoids,

respectively (Supplementary Table 1). Using gene set enrichment analysis (GSEA) we first examined the

consequences of *Lkb1* loss on well-defined gene signatures derived from individual intestinal cell types

(Haber *et al.*, 2017). In agreement with our earlier results, all *Lkb1*-deficient clones showed significant

downregulation of gene sets linked to mature Paneth cells, whereas goblet cell gene sets were enriched

within both budding and cystic *Lkb1*[-/-] organoids in comparison to WT organoids, again suggesting

alterations in secretory cell lineage specification (Figure 3A).

Noticeably, the *Lgr5*[+] adult stem cell signature (Haber *et al.*, 2017) was downregulated in all *Lkb1*-

deficient clones (Figure 3A). These data are reminiscent of the previously reported growth state of fetal

intestinal spheroids that relies on proliferation of *Lgr5*-negative progenitors (Mustata *et al.*, 2013). Indeed, we

uncovered a gradual enrichment of an embryonic spheroid gene signature in *Lkb1*-deficient organoids

(Mustata *et al.*, 2013), that correlated with the number of lost *Lkb1* alleles (Figure 3A-B). The transition from

adult stem cell to a fetal spheroid signature was most striking for *Lkb1*[cys -/-] organoids compared to *Lkb1*[bud -/-]

organoids (Figure 3B). Previous studies indicated that a transient upregulation of a fetal-like intestinal gene

signature associates with epithelial remodeling during injury repair and wound healing (Fordham *et al.*, 2013;

Nusse *et al.*, 2018; Yui, Azzolin, Maimets, Pedersen, Robert P Fordham, *et al.*, 2018; Wang *et al.*, 2019). In

line with these observations, *Lkb1*-deficient organoid clones also exhibited enrichment of gene signatures

associated with intestinal regeneration following damage or infection response (Figure 3A). Furthermore,

Lkb1-deficient clones, as well as control fetal intestinal organoids (E16.5), showed an increased expression

of the fetal marker Anxa1, while this marker was not detected in WT adult intestinal organoids (Figure 3C)

(Yui, Azzolin, Maimets, Pedersen, Robert P. Fordham, *et al.*, 2018). In addition, a dysplastic area of human

PJS patient-derived duodenal epithelial tissue showed patches of cells that stain positive for ANXA1 protein

expression, while non-polyp and healthy duodenum epithelial cells remained negative (Figure 3D). Together,

these findings indicate that *Lkb1* loss converts the epithelium into a permanent regenerative state. Strikingly,

this conversion towards a tissue repair growth state is already apparent upon monoallelic *Lkb1* loss, a

representative condition for the PJS intestinal epithelium.

***Lkb1*-deficient organoids show a Yap-induced regenerative state that promotes intra-epithelial self-**

sufficiency for Egf.

Over recent years, Yap signaling was placed central to intestinal regeneration (Gregorieff *et al.*, 2015).

Additionally, *LKB1* was uncovered previously as a negatively regulator of the Hippo-Yap pathway in an

siRNA kinome screen (Mohseni *et al.*, 2014). We therefore wondered if *Lkb1* loss may activate Yap signaling

as part of the induced regenerative program in intestinal organoids. Indeed, we identified enhanced

expression of a Yap-regulated gene signature linked to intestinal regeneration in *Lkb1*-mutant organoids

(Figure 3A) (Gregorieff *et al.*, 2015). Notably, increased Yap target gene expression was already identified in

heterozygous *Lkb1*-mutant conditions and progressively increased with loss of both *Lkb1* alleles (Figure S3).

Again, *Lkb1*[cys -/-] organoids showed the strongest induction of Yap target genes, in line with their robust

expression of a regenerative transcriptional program (Figure S3). Immunofluorescence staining further

confirmed an increase in Yap protein expression in both budding and cystic *Lkb1*[-/-] organoids, and a

prominent nuclear localization of Yap, particularly within *Lkb1*[cys -/-] organoids (Figure 4A-B).

YAP pathway activation is well-known to induce enhanced expression of EGF family ligands,

particularly in the context of regeneration (Zhang *et al.*, 2009; Ren *et al.*, 2010; Yang *et al.*, 2012; Gregorieff

et al., 2015). We therefore wondered whether *Lkb1*-deficient organoids display altered state of EGF pathway

dependency. Noticeably, we found that a combined set of the *Egf* ligand family were upregulated in *Lkb1*-

deficient organoids, in a gradual manner along with *Lkb1* allelic loss and alterations in organoid morphology

(Figure 5A). Furthermore, these changes were accompanied with an elevated expression of *Egf* receptor

family members (Figure 5B). As survival and growth of healthy homeostatic adult intestinal organoids
depends on supplementation with Egf, we investigated whether the enhanced expression of *Egf* family
ligands and its receptors in *Lkb1*-deficient organoids mediates niche factor self-sufficiency, by omitting Egf
from the culture medium. In contrast to WT organoids, all *Lkb1*-deficient organoids were capable of growing
without Egf supplementation (Figure 5C). In support of these findings, we confirmed that organoid survival
still depended on EGFR signaling, since *Lkb1*-deficient organoids were sensitive to Gefitinib, a potent EGFR
inhibitor (Figure 5D).

Together, these results demonstrate that *Lkb1* loss induces a constitutive Yap-induced regenerative
state which drives epithelial self-sufficiency for EGF ligand production. As niche-independent growth
properties are considered a hallmark of cancer progression (Fujii *et al.*, 2016), these results provide a
potential explanation for a growth advantage of *Lkb1*-deficient epithelial cells and the initiation of
tumorigenesis.

Lkb1-deficiency mediates reprogramming of intestinal stem cells into a regenerative state.
To perform an in-depth analysis of how *Lkb1* loss alters intestinal lineage specification and cellular
hierarchies, we subjected WT and *Lkb1*-deficient organoids to single cell RNA sequencing. We included an
analysis of mouse fetal intestinal organoids, to explore a potential overlap with fetal gene expression.
Unsupervised clustering of single cells identified 19 distinct clusters (Figure S4A). A clear separation of
cellular profiles derived from *Lkb1*$^{cys\ -/-}$ organoids, fetal and WT cells was observed, while *Lkb1*$^{bud\ -/-}$ and
Lkb1$^{+/-}$ organoids displayed closer resemblance to WT cells (Figure 6A). The separation of *Lkb1*$^{cys\ -/-}$ and fetal
cell clusters in UMAP indicates that *Lkb1*$^{cys\ -/-}$ organoids do not fully convert to a fetal state but only show
partial expression of a fetal-like gene program (Figure 6A). Furthermore, *Lkb1*$^{cys\ -/-}$ organoids are composed
of a rather homogenous population of cells that are different from all budding organoid clusters. We therefore
restricted annotation of intestinal cell types to budding organoid lines, using publicly available intestinal cell
type gene signatures (Figure 6C, Figure S4B-I) (Haber *et al.*, 2017). Cell type fraction analysis revealed that
Lkb1$^{bud\ -/-}$ organoids comprise an increased fraction of goblet cells compared to WT organoids, confirming
alterations in secretory cell differentiation (Figure 6B). Interestingly, the increase in goblet cells was
accompanied with a decrease in enterocytes, suggesting that the preferred differentiation trajectory in *Lkb1*-
deficient budding organoids is shifted from absorptive towards secretory lineages (Figure 6C). The
previously observed morphological aberrancies in secretory lineage specification were confirmed at the
transcriptional level, as *Lkb1*-deficient organoids displayed the occurrence of 'hybrid' cells that co-express
features of both Paneth and goblet cells (Figure 6D). Moreover, expression of Egf ligands was specifically
increased in *Lkb1*-deficient enteroendocrine and goblet secretory cells (Figure 6E) suggesting that these
cells might be responsible for the observed niche-independency of *Lkb1*-mutant organoids.

As expected, expression of the fetal and regenerative marker *Anxa1* is highly abundant in both fetal
and *Lkb1*$^{cys\ -/-}$ organoids (Figure 6F). Notably, we found that *Anxa1* expression in *Lkb1*$^{bud\ -/-}$ organoids is
induced mainly within stem and transit amplifying (stem/TA) cell clusters, indicating that this population is the
main contributor to the observed epithelial conversion into a regenerative state (Figure 6A, F). To further test
this assumption, we performed reclustering of stem/TA cell populations including all *Lkb1*$^{cys\ -/-}$ cells.
Unsupervised clustering indicates distinct clusters for WT, *Lkb1*$^{+/-}$ and *Lkb1*$^{bud\ -/-}$ stem/TA cells (Figure S4J-
K). To examine the effect of *Lkb1* loss on intestinal stem cells, we performed GSEA on genes differentially

expressed between WT stem cells and *Lkb1*-deficient stem cells (Supplementary Table 2). We observed

enrichment of goblet cell gene expression within *Lkb1*-deficient stem cell populations compared to WT

organoids (Figure 6G-H). This suggests that the capacity for preferential differentiation towards secretory

cells is already present in *Lkb1*-deficient stem cells. Furthermore, we identified an enrichment of four out of

six previously reported fetal and regenerative gene signatures, a depletion of the *Lgr5*[+] stem cell signature

and an increased expression of Anxa1 in *Lkb1*-deficient stem cells (Figure 6G, I). These findings strengthen

a model in which induction of a regenerative response by *Lkb1* loss originates from the stem cell

compartment of the intestinal epithelium.

In conclusion, our results argue that *Lkb1*-deficient stem/TA cells reside in a regenerative state and

preferentially differentiate towards secretory rather than absorptive cell lineages. As a result, secretory cells

start secreting EGF ligands to drive niche-independent growth of *Lkb1*-deficient epithelium.

Lkb1 loss drives a pre-malignant program along the serrated pathway which is amplified by

additional _Kras_ mutations.

Next, we aimed to obtain insight in how transcriptional reprogramming induced by *Lkb1* loss may promote

cancer predisposition. Colorectal cancer (CRC) develops mainly via two pathways that involve distinct

originating lesions and mutational events (Chen *et al.*, 2021). The classical adenoma-carcinoma sequence is

initiated by bi-allelic mutations in the tumor suppressor APC which drives formation of stem cell-enriched

adenomatous polyps that collect additional mutations (e.g. in KRAS and p53) to develop into microsatellite-

stable CRC characterized by chromosomal instability (CIN) (Fearon and Vogelstein, 1990). Recently, an

alternative pathway emerged, in which colorectal cancer arises from serrated precursor polyps that are

initiated by BRAF or KRAS mutations and display combined features of fetal gene expression, differentiated

cells and gastric metaplasia (Patai *et al.*, 2013; Chen *et al.*, 2021). Whether and how *Lkb1* loss drives entry

into one of these pathways for CRC malignant progression is not known.

To investigate this issue, we examined expression of gene sets linked to the conventional and

serrated pathways of CRC development in *Lkb1*-deficient organoids (Chen *et al.*, 2021). Strikingly, all *Lkb1*-

deficient organoids display significant enrichment of genes that were previously found to mark neoplastic

populations of serrated-specific cells (SSC) (Figure 7A). These effects gradually increased with the number

of lost *Lkb1* alleles, along with a simultaneous decrease in the expression of adenoma-specific genes (ASC)

(Figure 7B). Single cell analysis further revealed that the serrated gene program was selectively increased in

Lkb1-deficient stem cell populations (Figure S5). Furthermore, *Lkb1*[cys -/-] organoids lose expression of the

intestinal transcription factor *Cdx2* and upregulate expression of common gastric markers like *Aqp5*, *Tacstd2*

[*Trop2*], *Tff2*, and *Msln* (Figure 7B, Supplementary Table 1), indicating that *Lkb1* loss promotes intestinal-to-

gastric cellular transdifferentiation, often referred to as gastric metaplasia (Balbinot *et al.*, 2018; Chen *et al.*,

2021).

To investigate how an *Lkb1*-deficient epithelial state cooperates with common cancer mutations in

CRC, we introduced *Apc* or *Kras* mutations in *Lkb1*-mutant organoids. Truncating *Apc* mutations are linked

to the classical adenoma sequence, while *Kras* mutations are observed in both pathways (Fearon and

Vogelstein, 1990; Patai *et al.*, 2013). As expected, based on earlier reports, introduction of a truncating *Apc*

mutation induced cystic morphologies of all organoids (Figure S6A-B) (Sato, Stange, *et al.*, 2011; Dow *et al.*,

2015). Surprisingly, however, introduction of a G12D *Kras* mutation in *Lkb1*-deficient organoids also

consistently mediated transition to a cystic morphology, while single *Kras*-mutant organoids maintained their

crypt-villus organization (Figure S6C-D). Functionally, *Kras* mutations conveyed synergistic effects on the

ability of single *Lkb1*-deficient cells to reconstitute organoid growth (Figure 7C). By contrast, *Apc* mutations

did not add any significant increase to the reconstitution capacity of *Lkb1*$^{-/-}$ organoids (Figure 7D).

To gain insight in how *Apc* or *Kras* mutations altered the phenotypic landscape of *Lkb1*-deficient

organoids, we examined transcriptional expression profiles of all organoid lines using bulk mRNA

sequencing (Supplementary Table 3). With the introduction of mutant *Apc,* a slight increase was observed in

fetal and SSC gene programs in both *Lkb1*$^{+/-}$ and *Lkb1*$^{-/-}$ organoids, but the expression of stem cell and ASC

genes remained high for this mutational combination (Figure 7E, Figure S6E). By contrast, introduction of

mutant *Kras* in *Lkb1*$^{+/-}$ or *Lkb1*$^{bud\,-/-}$ backgrounds induced a strong shift towards expression of fetal and SSC

genes, while stem and ASC genes were downregulated, to a similar extent as observed for *Lkb1*$^{cys\,-/-}$

organoids (Figure 7E, Figure S6E). In *Lkb1*$^{cys\,-/-}$ organoids, introduction of mutant *Kras* did not further

enhance the already strong serrated features, while introduction of truncated *Apc* in these organoids clearly

suppressed the serrated state. Quantification of these findings by GSEA confirmed a significant amplification

of serrated signature by the introduction of *Kras* mutations in both *Lkb1*$^{+/-}$ and *Lkb1*$^{bud\,-/-}$ organoids, and

downregulation of the serrated pathway by introduction of mutant Apc (Figure 7F). We conclude that *Lkb1*

mutations synergize with mutant *Kras,* but not *Apc,* to drive expression of the pre-malignant serrated gene

program.

To investigate the link between *Lkb1* loss and advanced serrated colorectal tumors, we examined

the TCGA colorectal cancer database. First, we observed that *LKB1*-mutant tumors are significantly enriched

for mutations associated with serrated cancers, including *BRAF*, *RNF43* and *PIK3CA* mutations, while *APC*

mutations were significantly enriched in *LKB1* WT tumors (Figure 7G) (Day *et al.*, 2013; Robertson *et al.*,

2019). Mutations found in both classical and serrated adenocarcinomas, such as mutant *KRAS* and *TP53*,

were not significantly altered between *LKB1* WT and mutant tumors (Figure 7G). We next scored TCGA

colorectal samples for expression of a gene set that was upregulated in all *Lkb1*-deficient organoids.

Strikingly, expression of this *Lkb1*-deficient gene signature correlated with a mucinous histological CRC

subtype that is often found in serrated adenocarcinomas (Figure 7H) (Tuppurainen *et al.*, 2005). In addition,

among the consensus molecular subtypes (CMS), the serrated subtypes CMS1 and CMS4 carry significantly

higher expression of the *Lkb1*-deficient gene signature compared to the classical subtype CMS2 (Figure 7I)

(Nguyen, Goel and Chung, 2020). A recent study by Joanito *et al.* redefined the CMS classification and

found two epithelial intrinsic cell states in CRC samples (Joanito *et al.*, 2022). The iCMS2 state shows more

characteristics of classical CRCs, while the iCMS3 state correlates with features of serrated CRCs. Using

GSEA, we also found enrichment of iCMS3 transcriptional features in our *Lkb1*-deficient organoids, while the

iCMS2 state was downregulated (Figure S6F).

These results indicate that *Lkb1* loss drives cancer predisposition along the serrated pathway of

CRC development. *Kras* activating mutations cooperate with *Lkb1* loss by amplifying the serrated state.

Furthermore, upon homozygous *Lkb1* loss, a fraction of mutant cells grows out into *Lkb1*$^{cys\,-/-}$ organoids that

achieve an advanced serrated state without additional oncogenic mutations. Together, these findings

suggest that loss of *LKB1* converts intestinal epithelial cells in a pre-malignant state that primes for

progression along the serrated pathway.

Discussion

Over recent years, a variety of *Lkb1*-deficient mouse models have faithfully mimicked intestinal hamartomatous polyp formation as a hallmark for PJS disease. The combined findings have promoted a view in which non-epithelial functions of *LKB1* contribute to PJS disease development (Katajisto *et al.*, 2008; Ollila *et al.*, 2018; Poffenberger *et al.*, 2018; Cotton *et al.*, 2022). Despite this progress, the relationship between *LKB1* loss-induced hamartomatous polyposis and epithelial cancer predisposition has remained highly debated and, moreover, does not explain cancer development in extra-intestinal epithelia (Shepherd, Bussey and Jass, 1987; Entius *et al.*, 2001; De Leng *et al.*, 2003; Van Lier *et al.*, 2010b; Korsse *et al.*, 2013). Furthermore, how heterozygous and homozygous loss of *LKB1* mediates dose-dependent alterations in PJS epithelial tissue organization at the single cell level has not been addressed, leaving the underlying cancer driver mechanisms and pathways obscure.

Here, we employed mouse small intestinal organoids, an epithelium-only culture system, to obtain an in-depth understanding of how *Lkb1* loss affects intestinal epithelial homeostatic cell identities and examine how these alterations drive the epithelium into a pre-malignant state for colorectal cancer formation. By generating both heterozygous and homozygous knockouts of *Lkb1*, we aimed to model aspects of cancer predisposition and development in PJS patients, as over 64% of PJS-derived carcinomas present inactivation of the second copy of the gene (Entius *et al.*, 2001; De Leng *et al.*, 2003). Our findings argue that heterozygous *Lkb1* loss in the intestinal epithelium is sufficient to push cells into a premalignant programme for tumour development along the serrated colorectal cancer pathway, by a mechanism independent of the micro-environment. Furthermore, expression of the serrated premalignant program is amplified upon loss of the second allele. These results thus reveal that heterozygous *Lkb1*-mutant PJS epithelia already have undergone alterations in phenotypic cellular identities in comparison to healthy individuals, and supports the view that loss of the second allele is a driver of cancer formation.

Our organoid-based model further confirms the previously described link between *Lkb1* loss and aberrant secretory cell specification *in vivo* in mice, indicating that our model faithfully reproduces the *in vivo* situation. We observed an increase in goblet cell numbers and mucus formation and identified secretory cells that display 'hybrid' features, based both on morphology and transcriptome analysis. In line with these findings, we show that expression of our newly identified *Lkb1*-deficient gene signature correlates with a mucinous histological subtype of sporadic colorectal cancer cases. Together, these findings indicate a common link between early premalignant alterations in secretory cell type specification and colorectal cancer subtype progression. Similar *Lkb1* loss-induced alterations in secretory cell identities may play a role in other epithelial tissues, as mucinous carcinomas in different organs were frequently reported in PJS patients (Sato *et al.*, 2001; Osoegawa *et al.*, 2011; Collet *et al.*, 2020). If and how aberrant secretory cell specification may facilitate cancer development however remains an issue for further investigation.

Besides the observed alteration in secretory cell identities, transcriptomic analysis revealed that *Lkb1* loss induces an allele dosage-dependent expression of a regenerative epithelial program. Notably, activation of this tissue repair program occurred in regular culture conditions where no intestinal epithelial damage was induced. We conclude that the *Lkb1*-deficient epithelium resides in a persistent regenerative state, a condition that is worsened by loss-of-heterozygosity. A chronic state of regeneration may contribute to an increase in tumour development as exemplified by the increased incidence of colorectal cancer formation in patients suffering from inflammatory bowel disease (Lutgens *et al.*, 2013). As part of the induced

regenerative/YAP signaling-dependent program, the *Lkb1*-deficient epithelium showed an increase in

transcriptional expression of the *Egf* receptor family and its ligands, which allowed them to grow in the

absence of Egf supplementation, a condition that is not tolerated by healthy small intestinal organoids. These

findings thus reveal that the regenerative growth state induced by *Lkb1* loss endows the intestinal epithelium

with niche-independent properties, which is regarded as a hallmark of cancer (Hanahan and Weinberg,

2011). Furthermore, these results provide an explanation of the previously reported expanded proliferative

zone observed in the intestines of PJS patients (De Leng *et al.*, 2007)

Importantly, we uncovered that *Lkb1* loss predisposes the intestinal epithelium to express a gene

program related to the serrated pathway for colorectal cancer formation. In support of these observations,

Lkb1 loss induced common features of early serrated lesions such as gastric metaplasia, induction of a

regenerative gene program and depletion of classical adult stem cells (Chen *et al.*, 2021). Furthermore,

inspection of a large set of sporadic colorectal cancer cases revealed that *LKB1* mutations frequently co-

occur with driver mutations linked to the serrated colorectal cancer pathway. Additionally, we observed

elevated expression of our *Lkb1*-deficient gene signature in the serrated colorectal cancer subtypes CMS1

and CMS4. Together, these findings thus suggest that the pathways employed for colorectal cancer

development in PJS patients are strikingly different from the well-studied cancer driver events in familial

adenomatous polyposis (FAP) patients that develop conventional *APC*-mutant adenocarcinomas (Li *et al.*,

2020).

Based on the work of Chen and colleagues (Chen *et al.*, 2021) a model was proposed in which

different environmental triggers, such as microbial dysbiosis and a subsequent MAPK pathway activation

may drive the formation of sessile serrated lesions (IJspeert and Dekker, 2022; Shields *et al.*, 2022). Our

findings indicate that different types of insults may induce a shift of the *Lkb1*-mutant epithelium towards full-

blown expression of the premalignant serrated program. This is exemplified by introduction of oncogenic

mutant *Kras* in *Lkb1*-deficient organoids, which amplified serrated features and synergistically cooperated

with *Lkb1* loss for organoid-forming capacity. Alternatively, we found that *Lkb1* loss-of-heterozygosity

induced two distinct organoid phenotypes that displayed either a crypt-villus organization or a cystic

phenotype with strongly induced serrated features. Since both $Lkb1^{-/-}$ organoid types are genomically

identical, we hypothesize that an unknown external stressor may induce this switch to a highly serrated

cystic phenotype. Although the responsible molecular trigger remains to be identified, our findings indicate

that $Lkb1^{-/-}$ cells are sensitized to undergo such a phenotypic switch and adopt a pre-malignant growth state.

Based on our findings, we conclude that the *Lkb1*-mutant epithelium is primed for the sessile serrated cancer

pathway and not the conventional adenoma-to-carcinoma sequence. In line with this reasoning, the

introduction of *Apc* mutations in *Lkb1*-deficient organoids did not display synergistic effects on pre-malignant

transcriptional features or organoid reconstitution capacity. Together, our findings suggest that the PJS

epithelium is sensitized to events that synergistically drive cells into a premalignant serrated state, such as

intestinal damage or activating mutations in MAPK pathway components.

PJS patients undergo extensive and life-long monitoring in the clinic, to resect obstructive polyps and

screen for (pre)cancerous lesions. Our results argue that monitoring of PJS patients will require specific

screening for the occurrence of serrated lesions that potentially may occur in non-polyp regions as well.

Serrated lesions are known to be frequently overlooked as their diagnosis relies on the existence of an

architecturally distorted serrated crypt that are difficult to identify with regular screening methods (Rosty *et*

al., 2012). Diagnosed serrated lesions may be confirmed by various biomarkers including an increased
MUC5AC expression linked to the absence of CDX2 (Chen *et al.*, 2021), or the usage of additional markers
identified in this study such as ANXA1.
Taken together, our study reveals that intestinal organoids comprise a valuable model for *LKB1*
mutation-induced alterations in epithelial organization and growth in PJS patients. Our findings further
establish that induction of a premalignant sessile serrated program is a hallmark of PJS disease and
provides an explanation for increased cancer incidence in PJS patients. The presented models may provide
a starting point to uncover which and how additional mutations and environmental stressors synergistically
cooperate with *LKB1* mutations, and how interactions with stromal and immune components contribute to the
initiation of PJS malignant growth of the PJS epithelium. Ultimately, in-depth knowledge on the pathways
and processes involved in driving PJS epithelia into a pre-malignant state may help to design strategies that
re-establish epithelial homeostasis and reduce the incidence of cancer development in PJS patients.

References

Ayyaz, A. *et al.* (2019) 'Single-cell transcriptomes of the regenerating intestine reveal a revival stem cell', *Nature*. Springer US, 569(7754), pp. 121–125. doi: 10.1038/s41586-019-1154-y.

Baas, A. F. *et al.* (2004) 'Complete Polarization of Single Intestinal Epithelial Cells upon Activation of LKB1 by STRAD', *Cell*. Cell Press, 116(3), pp. 457–466. doi: 10.1016/S0092-8674(04)00114-X.

Balbinot, C. *et al.* (2018) 'The Cdx2 homeobox gene suppresses intestinal tumorigenesis through non-cell-autonomous mechanisms', *Journal of Experimental Medicine*, 215(3), pp. 911–926. doi: 10.1084/jem.20170934.

Bansil, R. and Turner, B. S. (2018) 'The biology of mucus: Composition, synthesis and organization', *Advanced Drug Delivery Reviews*. Elsevier, 124, pp. 3–15. doi: 10.1016/J.ADDR.2017.09.023.

Barker, N. *et al.* (2007) 'Identification of stem cells in small intestine and colon by marker gene Lgr5', *Nature 2007 449:7165*. Nature Publishing Group, 449(7165), pp. 1003–1007. doi: 10.1038/nature06196.

Beggs, A. D. *et al.* (2010) 'Peutz-Jeghers syndrome: a systematic review and recommendations for management.', *Gut*. England, 59(7), pp. 975–986. doi: 10.1136/gut.2009.198499.

Butler, A. *et al.* (2018) 'Integrating single-cell transcriptomic data across different conditions, technologies, and species', *Nature Biotechnology*, 36(5), pp. 411–420. doi: 10.1038/nbt.4096.

Candelli, T. *et al.* (2018) 'Sharq, A versatile preprocessing and QC pipeline for Single Cell RNA-seq', *bioRxiv*, p. 250811. doi: 10.1101/250811.

Chen, B. *et al.* (2021) 'Differential pre-malignant programs and microenvironment chart distinct paths to malignancy in human colorectal polyps', *Cell*. The Authors, 184(26), pp. 6262-6280.e26. doi: 10.1016/j.cell.2021.11.031.

Collet, L. *et al.* (2020) 'Kras and Lkb1 mutations synergistically induce intraductal papillary mucinous neoplasm derived from pancreatic duct cells', *Gut*. BMJ Publishing Group, 69(4), pp. 704–714. doi: 10.1136/GUTJNL-2018-318059.

Cotton, J. L. *et al.* (2022) 'PTEN and LKB1 are differentially required in Gli1-expressing mesenchymal cells to suppress gastrointestinal polyposis', *Cell Reports*, 40(3), p. 111125. doi: 10.1016/J.CELREP.2022.111125.

Day, F. L. *et al.* (2013) 'PIK3CA and PTEN gene and exon mutation-specific clinicopathologic and molecular associations in Colorectal cancer', *Clinical Cancer Research*. American Association for Cancer Research, 19(12), pp. 3285–3296. doi: 10.1158/1078-0432.CCR-12-3614/85568/AM/PIK3CA-AND-PTEN-GENE-AND-EXON-MUTATION-SPECIFIC.

Dow, L. E. *et al.* (2015) 'Apc Restoration Promotes Cellular Differentiation and Reestablishes Crypt Homeostasis in Colorectal Cancer', *Cell*. Cell Press, 161(7), pp. 1539–1552. doi: 10.1016/j.cell.2015.05.033.

Entius, M. M. *et al.* (2001) 'Molecular genetic alterations in hamartomatous polyps and carcinomas of patients with Peutz-Jeghers syndrome', *Journal of clinical pathology*, 54(2), pp. 126–131. doi:

10.1136/jcp.54.2.126.

Fearon, E. R. and Vogelstein, B. (1990) 'A Genetic Model for Colorectal Tumorigenesis', *Cell*, 61, pp. 759–
767.

Fodde, R. (2002) 'The APC gene in colorectal cancer', *European Journal of Cancer*. Pergamon, 38(7), pp.
867–871. doi: 10.1016/S0959-8049(02)00040-0.

Fordham, R. P. *et al.* (2013) 'Transplantation of expanded fetal intestinal progenitors contributes to colon
regeneration after injury', *Cell Stem Cell*. The Authors, 13(6), pp. 734–744. doi: 10.1016/j.stem.2013.09.015.

Fujii, M. *et al.* (2015) 'Efficient genetic engineering of human intestinal organoids using electroporation',
Nature Protocols 2015 10:10. Nature Publishing Group, 10(10), pp. 1474–1485. doi:
10.1038/nprot.2015.088.

Fujii, M. *et al.* (2016) 'A Colorectal Tumor Organoid Library Demonstrates Progressive Loss of Niche Factor
Requirements during Tumorigenesis', *Cell Stem Cell*. Cell Press, 18(6), pp. 827–838. doi:
10.1016/J.STEM.2016.04.003.

Gao, Y. *et al.* (2020) 'LKB1 Represses ATOH1 via PDK4 and Energy Metabolism and Regulates Intestinal
Stem Cell Fate', *Gastroenterology*. W.B. Saunders, 158(5), p. 1401. doi: 10.1053/j.gastro.2019.12.033.

Giardiello, F. M. *et al.* (2000) 'Very high risk of cancer in familial Peutz-Jeghers syndrome.',
Gastroenterology. United States, 119(6), pp. 1447–1453. doi: 10.1053/gast.2000.20228.

Giardiello, F. M. and Trimbath, J. D. (2006) 'Peutz-Jeghers syndrome and management recommendations.',
Clinical gastroenterology and hepatology : the official clinical practice journal of the American
Gastroenterological Association. United States, 4(4), pp. 408–415. doi: 10.1016/j.cgh.2005.11.005.

Gregorieff, A. *et al.* (2015) 'Yap-dependent reprogramming of Lgr5+ stem cells drives intestinal regeneration
and cancer', *Nature*, 526(7575), pp. 715–718. doi: 10.1038/nature15382.

Haber, A. L. *et al.* (2017) 'A single-cell survey of the small intestinal epithelium', *Nature*. Nature Publishing
Group, 551(7680), pp. 333–339. doi: 10.1038/nature24489.

Hafemeister, C. and Satija, R. (2019) 'Normalization and variance stabilization of single-cell RNA-seq data
using regularized negative binomial regression', *Genome Biology*. Genome Biology, 20(1), pp. 1–15. doi:
10.1186/s13059-019-1874-1.

Hanahan, D. and Weinberg, R. A. (2011) 'Hallmarks of Cancer: The Next Generation', *Cell*. Cell Press,
144(5), pp. 646–674. doi: 10.1016/J.CELL.2011.02.013.

Hanemaaijer, E. S. *et al.* (2021) 'Single-cell atlas of developing murine adrenal gland reveals relation of
Schwann cell precursor signature to neuroblastoma phenotype', *Proceedings of the National Academy of*
Sciences of the United States of America, 118(5). doi: 10.1073/pnas.2022350118.

Hearle, N. *et al.* (2006) 'Frequency and spectrum of cancers in the Peutz-Jeghers syndrome.', *Clinical*
cancer research : an official journal of the American Association for Cancer Research. United States,
12(10), pp. 3209–3215. doi: 10.1158/1078-0432.CCR-06-0083.

Hemminki, A. *et al.* (1998) 'A serine/threonine kinase gene defective in Peutz–Jeghers syndrome', *Nature*
1998 391:6663. Nature Publishing Group, 391(6663), pp. 184–187. doi: 10.1038/34432.

IJspeert, J. E. G. and Dekker, E. (2022) 'Gastric metaplasia could initiate the serrated neoplasia pathway in
CRC', *Nature Reviews Gastroenterology and Hepatology*, 19(4), pp. 217–218. doi: 10.1038/s41575-022-
00592-z.

Jenne, D. E. *et al.* (1998) 'Peutz-Jeghers syndrome is caused by mutations in a novel serine
threoninekinase', *Nature Genetics 1998 18:1*. Nature Publishing Group, 18(1), pp. 38–43. doi:
10.1038/ng0198-38.

Joanito, I. *et al.* (2022) 'Single-cell and bulk transcriptome sequencing identifies two epithelial tumor cell
states and refines the consensus molecular classification of colorectal cancer', *Nature Genetics*, 54(7), pp.
963–975. doi: 10.1038/s41588-022-01100-4.

Katajisto, P. *et al.* (2008) 'LKB1 signaling in mesenchymal cells required for suppression of gastrointestinal
polyposis', *Nature genetics*. Nat Genet, 40(4), pp. 455–459. doi: 10.1038/NG.98.

Kim, Y. *et al.* (2022) 'LKB1 in Intestinal Epithelial Cells Regulates Bile Acid Metabolism by Modulating
FGF15/19 Production', *Cmgh*, 13(4), pp. 1121–1139. doi: 10.1016/j.jcmgh.2021.12.017.

Kline, E. R. *et al.* (2013) 'LKB1 Represses Focal Adhesion Kinase (FAK) Signaling via a FAK-LKB1 Complex
to Regulate FAK Site Maturation and Directional Persistence', *The Journal of Biological Chemistry*. American
Society for Biochemistry and Molecular Biology, 288(24), p. 17663. doi: 10.1074/JBC.M112.444620.

Korsse, S. E. *et al.* (2013) 'Identification of molecular alterations in gastrointestinal carcinomas and
dysplastic hamartomas in Peutz-Jeghers syndrome', *Carcinogenesis*. Oxford Academic, 34(7), pp. 1611–
1619. doi: 10.1093/CARCIN/BGT068.

De Leng, W. W. J. *et al.* (2003) 'Cyclooxygenase 2 expression and molecular alterations in Peutz-Jeghers
hamartomas and carcinomas.', *Clinical cancer research : an official journal of the American Association for*
Cancer Research. United States, 9(8), pp. 3065–3072.

De Leng, W. W. J. *et al.* (2007) 'Peutz–Jeghers syndrome polyps are polyclonal with expanded progenitor
cell compartment', *Gut*. BMJ Publishing Group, 56(10), pp. 1475–1476. doi: 10.1136/GUT.2007.128132.

Li, J. *et al.* (2020) 'Genomic and transcriptomic profiling of carcinogenesis in patients with familial
adenomatous polyposis', *Gut*, 69, pp. 1283–1293. doi: 10.1136/gutjnl-2019-319438.

Van Lier, M. G. F. *et al.* (2010a) 'High cancer risk in peutz-jeghers syndrome: A systematic review and
surveillance recommendations', *American Journal of Gastroenterology*. Nature Publishing Group, 105(6), pp.
1258–1264. doi: 10.1038/ajg.2009.725.

Van Lier, M. G. F. *et al.* (2010b) 'High cancer risk in peutz-jeghers syndrome: A systematic review and
surveillance recommendations', *American Journal of Gastroenterology*. Nature Publishing Group, 105(6), pp.
1258–1264. doi: 10.1038/ajg.2009.725.

Lutgens, M. W. M. D. *et al.* (2013) 'Declining Risk of Colorectal Cancer in Inflammatory Bowel DiseaseAn

Updated Meta-analysis of Population-based Cohort Studies', *Inflammatory Bowel Diseases*. Oxford
Academic, 19(4), pp. 789–799. doi: 10.1097/MIB.0B013E31828029C0.

Mohseni, M. *et al.* (2014) 'A genetic screen identifies an LKB1-MARK signalling axis controlling the Hippo-
YAP pathway', *Nature Cell Biology*. Nature Publishing Group, 16(1), pp. 108–117. doi: 10.1038/ncb2884.

Muraro, M. J. *et al.* (2016) 'A Single-Cell Transcriptome Atlas of the Human Pancreas', *Cell Systems*.
Elsevier, 3(4), p. 385. doi: 10.1016/J.CELS.2016.09.002.

Mustata, R. C. *et al.* (2013) 'Identification of Lgr5-Independent Spheroid-Generating Progenitors of the
Mouse Fetal Intestinal Epithelium', *Cell Reports*, 5(2), pp. 421–432. doi: 10.1016/j.celrep.2013.09.005.

Nguyen, L. H., Goel, A. and Chung, D. C. (2020) 'Pathways of Colorectal Carcinogenesis', *Gastroenterology*.
Elsevier, Inc, 158(2), pp. 291–302. doi: 10.1053/j.gastro.2019.08.059.

Nusse, Y. M. *et al.* (2018) 'Parasitic helminths induce fetal-like reversion in the intestinal stem cell niche',
Nature. Springer US, 559(7712), pp. 109–113. doi: 10.1038/s41586-018-0257-1.

Ollila, S. *et al.* (2018) 'Stromal Lkb1 deficiency leads to gastrointestinal tumorigenesis involving the IL-11-
JAK/STAT3 pathway', *The Journal of clinical investigation*. J Clin Invest, 128(1), pp. 402–414. doi:
10.1172/JCI93597.

Osoegawa, A. *et al.* (2011) 'LKB1 Mutations Frequently Detected in Mucinous Bronchioloalveolar
Carcinoma', *Japanese Journal of Clinical Oncology*. Oxford Academic, 41(9), pp. 1132–1137. doi:
10.1093/JJCO/HYR102.

Patai, Á. V. *et al.* (2013) 'Serrated pathway: Alternative route to colorectal cancer', *World Journal of*
Gastroenterology, 19(5), pp. 607–615. doi: 10.3748/wjg.v19.i5.607.

Pierce, S. E. *et al.* (2021) 'LKB1 inactivation modulates chromatin accessibility to drive metastatic
progression', *Nature Cell Biology*, 23(8), pp. 915–924. doi: 10.1038/s41556-021-00728-4.

Poffenberger, M. C. *et al.* (2018) 'LKB1 deficiency in T cells promotes the development of gastrointestinal
polyposis', *Science*. American Association for the Advancement of Science, 361(6400), pp. 406–411. doi:
10.1126/SCIENCE.AAN3975.

Qu, M. *et al.* (2021a) 'Establishment of intestinal organoid cultures modeling injury-associated epithelial
regeneration', *Cell Research*. Springer US, 31(3), pp. 259–271. doi: 10.1038/s41422-020-00453-x.

Qu, M. *et al.* (2021b) 'Establishment of intestinal organoid cultures modeling injury-associated epithelial
regeneration', *Cell Research*. Springer US, (October 2020). doi: 10.1038/s41422-020-00453-x.

Ran, F. A. *et al.* (2013) 'Genome engineering using the CRISPR-Cas9 system', *Nature Protocols*, 8(11), pp.
2281–2308. doi: 10.1038/nprot.2013.143.

Ren, F. *et al.* (2010) 'Hippo signaling regulates Drosophila intestine stem cell proliferation through multiple
pathways', *Proceedings of the National Academy of Sciences of the United States of America*. National
Academy of Sciences, 107(49), pp. 21064–21069. doi:
10.1073/PNAS.1012759107/SUPPL_FILE/PNAS.201012759SI.PDF.

Robertson, D. J. *et al.* (2019) 'Terminology, Molecular Features, Epidemiology, and Management of Serrated
Colorectal Neoplasia', *Gastroenterology*. Elsevier, 157(4), pp. 949-966.e4. doi:
10.1053/J.GASTRO.2019.06.041.

Rosty, C. *et al.* (2012) 'Serrated polyps of the large intestine: current understanding of diagnosis,
pathogenesis, and clinical management', *Journal of Gastroenterology 2012 48:3*. Springer, 48(3), pp. 287–
302. doi: 10.1007/S00535-012-0720-Y.

Sanchez-Cespedes, M. (2007) 'A role for LKB1 gene in human cancer beyond the Peutz-Jeghers
syndrome.', *Oncogene*. England, 26(57), pp. 7825–7832. doi: 10.1038/sj.onc.1210594.

Sato, N. *et al.* (2001) 'STK11/LKB1 Peutz-Jeghers Gene Inactivation in Intraductal Papillary-Mucinous
Neoplasms of the Pancreas', *The American Journal of Pathology*. Elsevier, 159(6), pp. 2017–2022. doi:
10.1016/S0002-9440(10)63053-2.

Sato, T. *et al.* (2009) 'Single Lgr5 stem cells build crypt–villus structures in vitro without a mesenchymal
niche', *Nature*. Macmillan Publishers Limited. All rights reserved, 459(7244), pp. 262–265.

Sato, T., Stange, D. E., *et al.* (2011) 'Long-term Expansion of Epithelial Organoids From Human Colon,
Adenoma, Adenocarcinoma, and Barrett's Epithelium', *Gastroenterology*. W.B. Saunders, 141(5), pp. 1762–
1772. doi: 10.1053/J.GASTRO.2011.07.050.

Sato, T., Es, J. H. Van, *et al.* (2011) 'Paneth cells constitute the niche for Lgr5 stem cells in intestinal crypts',
Nature, 469(7330), pp. 415–418. doi: 10.1038/nature09637.Paneth.

Satoh, Y. *et al.* (1990) 'Ultrastructure of Paneth cells in the intestine of various mammals', *Journal of Electron
Microscopy Technique*. John Wiley & Sons, Ltd, 16(1), pp. 69–80. doi: 10.1002/JEMT.1060160109.

Shackelford, D. B. and Shaw, R. J. (2009) 'The LKB1-AMPK pathway: Metabolism and growth control in
tumour suppression', *Nature Reviews Cancer*. Nature Publishing Group, 9(8), pp. 563–575. doi:
10.1038/nrc2676.

Shepherd, N. A., Bussey, H. J. R. and Jass, J. R. (1987) 'Epithelial misplacement in Peutz-Jeghers Polyps A
Diagnostic Pitfall', *The American Journal of Surgical Pathology*, 11(10), pp. 743–749.

Shields, C. E. D. *et al.* (2022) 'Sensitive To Immune Checkpoint Therapy', 11(7), pp. 1792–1807. doi:
10.1158/2159-8290.CD-20-0770.Bacterial-driven.

Shorning, Boris Y. *et al.* (2009) 'Lkb1 deficiency alters goblet and paneth cell differentiation in the small
intestine', *PLoS ONE*, 4(1), pp. 1–8. doi: 10.1371/journal.pone.0004264.

Shorning, B Y *et al.* (2009) 'Lkb1 Deficiency Alters Goblet and Paneth Cell Differentiation in the Small
Intestine', *PLoS ONE*, 4(1), p. 4271. doi: 10.1371/journal.pone.0004264.

Tomasetti, C. and Vogelstein, B. (2015) 'Variation in cancer risk among tissues can be explained by the
number of stem cell divisions', *Science*. American Association for the Advancement of Science, 347(6217),
pp. 78–81. doi: 10.1126/SCIENCE.1260825.

Troughton, W. D. and Trier, J. S. (1969) 'Paneth and goblet cell renewal in mouse duodenal crypts.', *The

Journal of cell biology, 41(1), pp. 251–268. doi: 10.1083/jcb.41.1.251.

Tuppurainen, K. *et al.* (2005) 'Morphology and microsatellite instability in sporadic serrated and non-serrated
colorectal cancer', *Journal of Pathology J Pathol*, 207, pp. 285–294. doi: 10.1002/path.1850.

Wang, Y. *et al.* (2019) 'Long-Term Culture Captures Injury-Repair Cycles of Colonic Stem Cells', *Cell*.
Elsevier Inc., 179(5), pp. 1144-1159.e15. doi: 10.1016/j.cell.2019.10.015.

Yang, N. *et al.* (2012) 'TAZ induces growth factor-independent proliferation through activation of EGFR
ligand amphiregulin', *Cell Cycle*. Taylor and Francis Inc., 11(15), pp. 2922–2930. doi: 10.4161/CC.21386.

Yui, S., Azzolin, L., Maimets, M., Pedersen, M. T., Fordham, Robert P, *et al.* (2018) 'YAP/TAZ-Dependent
Reprogramming of Colonic Epithelium Links ECM Remodeling to Tissue Regeneration', *Cell stem cell*.
2017/12/14. Cell Press, 22(1), pp. 35-49.e7. doi: 10.1016/j.stem.2017.11.001.

Yui, S., Azzolin, L., Maimets, M., Pedersen, M. T., Fordham, Robert P., *et al.* (2018) 'YAP/TAZ-Dependent
Reprogramming of Colonic Epithelium Links ECM Remodeling to Tissue Regeneration', *Cell Stem Cell*.
Elsevier Inc., 22(1), pp. 35-49.e7. doi: 10.1016/j.stem.2017.11.001.

Zhang, J. *et al.* (2009) 'YAP-dependent induction of amphiregulin identifies a non-cell-autonomous
component of the Hippo pathway', *Nature Cell Biology*. doi: 10.1038/ncb1993.

    Figure legends

    Figure 1. *Lkb1*-deficient organoids show heterogeneous morphology and viability.

    A) Representative brightfield images of mouse small intestine *Lkb1*-deficient organoid clones. Lower panel

    images show zoom-in images of boxed area in upper panel images. Scale = 100 µM **B)** Western blot

    analysis shows loss of Lkb1 expression in heterozygous CRISPR targeted clones and complete loss of Lkb1

    protein levels in homozygously targeted clones. **C)** Cell reconstitution assay from single cells performed at

    day 7 identifies increased organoid forming capacities in *Lkb1*$^{cys-/-}$. One-way ANOVA (Dunnett) was applied

    for analysis. * = *p < 0.05.* **D)** EdU incorporation experiment marking the proliferative compartments in wild-

    type (WT) and *Lkb1*-deficient organoids. **E)** Quantification of EdU$^+$ cells over the total number of cells (DAPI$^+$

    cells) of *Lkb1*-deficient organoids, *n*=1000.

    Figure 2. *Lkb1* loss alters the number, location and granule composition of secretory cells. A)

    Representative images of immunostaining for Lysozyme1 (Lyz1) to detect Paneth cells in *Lkb1*-deficient

    clones. Images are max projections of z-stack. Scale bar = 25 µm. **B)** Immunostaining of Mucin 2 (Muc2) to

    visualize goblet cells in *Lkb1*-deficient clones. Scale bar =100 µm. **C-D)** Representative images of Mucin 2

    (C) and Lysozyme 1 **(D)** staining to visualize the localization of goblet and Paneth cells respectively in

    healthy intestinal epithelium and PJS epithelium. Arrows point towards mislocalized Paneth cells. Scalebar =

    100 µm. **E)** Electron microscopy images comparing the ultrastructure of Paneth cells and their secretory

    granules in wild-type organoids and *Lkb1*-deficient organoids. Scale bar upper panels = 5 µm, scale bar

    lower panels = 1 µm. **F)** Electron microscopy images comparing the ultrastructure of goblet cells and their

    secretory granules in wild-type organoids and *Lkb1*-deficient organoids. Scale bar upper panels = 5 µm,

    scale bar lower panels = 1 µm.

    Figure 3. *Lkb1* loss upregulates a regenerative gene signature in the intestinal epithelium. A) *Lkb1*-

    deficient versus wild-type (WT) organoid derived bulk mRNA-seq sequencing data was analysed by gene set

    enrichment analysis (GSEA) for cell types (Haber *et al.*, 2017), colitis (Yui, Azzolin, Maimets, Pedersen,

    Robert P. Fordham, *et al.*, 2018), injury (Qu *et al.*, 2021a), parasite (Nusse *et al.*, 2018), fetal spheroid

    (Mustata *et al.*, 2013), Yap regeneration (Gregorieff *et al.*, 2015) and revival stem cell (Ayyaz *et al.*, 2019)

    gene sets. The normalized enrichment score (NES) is depicted on the X-axis and the false discovery rates

    (FDRs) is indicated by the size of the dots. The color of the lines reflects whether a signature is enriched in

    WT (red) or *Lkb1*-deficient (blue) organoids. The shape of the dot (open/closed) indicates whether a

    signature is significantly enriched (FDR < 0.05). **B)** Heatmap of z-score transformed expression values of

    stem cell genes enriched in WT organoids (top cluster) and fetal genes enriched in *Lkb1*-deficient organoids

    (bottom cluster). Key marker genes are highlighted. **C)** Immunofluorescent staining to detect expression of

    the fetal marker Anxa1 in *Lkb1*-deficient clones. Embryonic spheroids from E16.5 mice were used as positive

    control. **D)** Representative images of immunohistochemistry staining of the fetal marker ANXA1 in healthy

    intestinal epithelium, PJS normal epithelium and PJS dysplastic tissue. High magnification of the area

    indicated by black arrows is shown in the upper right corner. Scale bar = 100 µm

Figure 4. *Lkb1*-deficient organoids show a Yap-induced regenerative state.

A) Immunostaining of Yap in wild-type (WT) and *Lkb1*-deficient organoids. Nuclei are visualized by DAPI staining. High magnification of the area indicated by black arrows is shown in the upper right corner. Scale bar = 20 µm. **B)** Quantification of the nuclear fraction of Yap. Dots represent individual organoids. N=5 for WT organoids, N=7 for *Lkb1*$^{+/-}$ organoids, N=7 for *Lkb1*$^{bud\,-/-}$ organoids and N=7 for *Lkb1*$^{cys\,-/-}$. One-way ANOVA was applied for analysis and statistical significance was defined as *= p < 0.05, ** = p < 0.01 and **** = p < 0.0001.*

Figure 5. *Lkb1*-deficient organoids upregulate EGF family receptors and its ligands for niche independency.

A-B) Average expression of *Egfr* binding ligands **(A)** and *Egf* family receptors **(B)** in wild-type (WT) and *Lkb1*-deficient organoids. One-way ANOVA was applied for analysis and statistical significance was defined as *= p < 0.05, ** = p < 0.01 and **** = p < 0.0001.* **C)** Representative bright field images of WT and *Lkb1*-deficient organoids grown in the presence (ENR) and absence (NR) of Egf. Organoids were incubated for 7 days in indicated medium and split on day 4. Scale bar = 500 µm. **D)** Representative bright field images of WT and *Lkb1*-deficient organoids cultured in NR and treated with DMSO or 100 nM Gefitinib. Organoids were incubated for 9 days in indicated medium and split on day 6. Scale bar = 500 µm.

Figure 6. *Lkb1*-deficiency redefines intestinal stem cell identity.

A-B) UMAP visualization of scRNA-seq data from wild-type, fetal and *Lkb1*-deficient organoids. Colors indicate the different organoids lines **(A)** and the different cell types **(B)**. **C)** Fraction of cell types for each organoid line. Each color indicates a different cell type. **D)** Scatter plot showing module scores for a Paneth gene signature (Haber *et al.*, 2017) on the y-axis and for a goblet cell signature (Haber *et al.*, 2017) on the x-axis across all cells. Color/shape indicates the origin of the single cell. **E)** Violin plot showing a module score for Egf family ligands within the goblet and enteroendocrine (EEC) cluster. Colors reflect different organoid lines. **F)** Expression of the fetal marker *Anxa1* is overlaid onto the UMAP plot for the different organoid lines. **G)** GSEA was performed for cell types (Haber *et al.*, 2017), colitis (Yui, Azzolin, Maimets, Pedersen, Robert P. Fordham, *et al.*, 2018), embryonic spheroid (Mustata *et al.*, 2013), Yap regeneration (Gregorieff *et al.*, 2015) and revival stem cell (Ayyaz *et al.*, 2019) gene sets. The normalized enrichment score (NES) is depicted on the X-axis and the false discovery rates (FDRs) are indicated by the size of the dots. The color of the lines reflects whether a signature is enriched in WT (red) or *Lkb1*-deficient (blue) organoids. The shape of the dot (open/closed) indicates whether a signature is significantly enriched (FDR < 0.05). **H)** Violin plot showing module scores for a goblet cell gene signature across WT stem cells, *Lkb1*$^{+/-}$ stem cells, *Lkb1*$^{bud\,-/-}$ stem cells and all *Lkb1*$^{cys\,-/-}$ cells. **I)** Violin plot showing expression of the regenerative marker *Anxa1* across WT stem cells, *Lkb1*$^{+/-}$ stem cells, *Lkb1*$^{bud\,-/-}$ stem cells and all *Lkb1*$^{cys\,-/-}$ cells.

Figure 7. *Lkb1* loss drives a pre-malignant program along the serrated pathway which can be amplified by additional *Kras* mutations. A) GSEA was performed for a serrated-specific gene set from (Chen *et al.*, 2021). The normalized enrichment score (NES) is depicted on the X-axis and the false discovery rate (FDR) is indicated by the size of the dots. The shape of the dot (open/closed) indicates whether a signature is significantly enriched (FDR < 0.05). **B)** Heatmap of z-score transformed expression

values of adenoma-specific genes (ASC) enriched in WT organoids (top cluster) and serrated-specific genes

(SSC) enriched in *Lkb1*-deficient organoids (bottom cluster). **C)** Cell reconstitution assay from single cells

performed at day 7 from trypsinization is shown for WT, *Lkb1*-deficient, *Apc*-deficient and *Apc/Lkb1*-deficient

organoids. **D)** Cell reconstitution assay from single cells performed at day 7 from trypsinization is shown for

WT, *Lkb1*-deficient, *Kras*-mutant and *Kras*-mutant/*Lkb1*-deficient organoids. **E)** Heatmap of z-score

transformed expression values of stem cell genes enriched in control organoids (top cluster) and fetal genes

enriched in *Lkb1*-deficient organoids (bottom cluster). **F)** GSEA of a serrated-specific gene set (Chen *et al.*,

2021) was performed on *Lkb1*-deficient and *Apc/Lkb1*-deficient or *Kras*-mutant/*Lkb1*-deficient organoids. The

normalized enrichment score (NES) is depicted on the X-axis and the FDR is indicated by the size of the

dots. The color of the lines reflect whether a signature is enriched in *Lkb1*-deficient (red) or *Lkb1*-

deficient/*Kras*-mutant (blue) organoids. **G)** Percentage of samples mutated for *BRAF, RNF43, PIK3CA, APC,*

KRAS or *TP53* in *LKB1* WT and *LKB1*-mutant colorectal tumors according to cBioPortal. **H, I)** Box plots

showing log2 Z-scores for the *Lkb1*-deficient signature (genes significantly upregulated in all deficient clones

compared to control organoids) over the colon adenocarcinoma TCGA bulk RNA sequencing dataset

grouped by pathology classification as mucinous or non-mucinous **(H)** or consensus molecular subtype **(I)**.

Figure S1. *Lkb1* loss induced by different gRNAs results in similar heterogeneous phenotypes.

A) Sequences of *Lkb1*-deficient organoids obtained with gRNA1 showing indels/deletions detected in Exon 1

of the *Lkb1* gene. PAM sequences are underlined. **B)** Immunofluorescent images showing cleaved caspase

3 staining marking apoptotic cells in *Lkb1*-deficient and control organoids. Scale bar = 33 μm. **C)** Sequences

of *Lkb1*-deficient organoids obtained with gRNA1 showing indels/deletions detected in Exon 1 of the *Lkb1*

gene. PAM sequences are underlined. **D)** Representative images of *Lkb1*-deficient clones showing

heterogeneous morphologies. **E)** Western blot analysis of Lkb1 expression in each clone. **F)** Cell

reconstitution assay from single cells performed at day 7 from trypsinization.

Figure S2. Goblet cell number and positioning are unaltered in PJS epithelia and *Lkb1*-deficient

organoids display proper brush border formation. A-B) The average number **(A)** and position **(B)** of

goblet cells healthy and PJS epithelia. Unpaired t-test was applied for analysis. **C)** Electron microscopy

images of WT and *Lkb1*-deficient organoids. Arrows indicate brush border formation in indicated organoid

lines. A zoom-in of the area indicated by the red arrows is shown in the bottom panel. Scale bar = 2 μm.

Figure S3. Yap target gene expression progressively increases with loss of both *Lkb1* alleles.

Heatmap of z-score transformed expression values of Yap regulated regenerative genes. Key marker genes

are highlighted.

Figure S4. Identification of intestinal cell types of WT and *Lkb1*-deficient organoids.

A) UMAP visualization of scRNA-seq data from wild-type, fetal and *Lkb1*-deficient organoids. Colors indicate

different clusters as defined by Seurat analysis **B)** Expression of known markers for different cell populations

in WT and *Lkb1*-deficient organoids. **C-I)** Module scores for enterocyte **(C)**, enteroendocrine **(D)**, goblet **(E)**,

stem **(F)**, transit amplifying (G1) **(G)**, transit amplifying (G2) **(H)** and tuft **(I)** cell gene signatures overlaid on

the UMAP plot of *Lkb1*-deficient and control organoids. **J-K)** Reclustering of stem cell populations from WT,

Lkb1$^{+/-}$, *Lkb1$^{-/-\ bud}$* and *Lkb1$^{-/-\ cys}$* organoids.

Figure S5. Induction of serrated genes in *Lkb1*-deficient organoids is reflected at the single cell level.

A) Module score for a serrated-specific (SSC) gene set on the UMAP plot of *Lkb1*-deficient and WT

organoids. **B)** Violin plot showing a module score for a serrated-specific gene set across WT adult, WT fetal,

Lkb1$^{+/-}$, *Lkb1$^{bud\ -/-}$* and *Lkb1$^{cys\ -/-}$* organoids. **C)** GSEA was performed for a serrated-specific gene set from

(Chen *et al.*, 2021). The normalized enrichment score (NES) is depicted on the Y-axis and the false

discovery rate (FDR) is indicated by the size of the dots. Line colors reflects enrichment of the signature in

WT or *Lkb1*-deficient stem cells. Dot shape (open/closed) indicates significant enrichment (FDR < 0.05). **D)**

Module score for a serrated-specific gene set on the UMAP plot of WT stem cells, *Lkb1$^{+/-}$* stem cells, *Lkb1^{bud}*

$^{-/-}$ stem cells and all *Lkb1$^{cys\ -/-}$* cells. **E)** Violin plot showing module scores of a serrated-specific gene set for

WT stem cells, *Lkb1$^{+/-}$* stem cells, *Lkb1$^{bud\ -/-}$* stem cells and all *Lkb1$^{cys\ -/-}$* cells.

Figure S6. *Lkb1* loss drives expression of a pre-malignant program along the serrated pathway which

can be amplified by additional *Kras* mutations. A) Representative images of *Apc* and *Apc/Lkb1*-deficient

organoids. **B)** Representative images of *Kras*-mutant and *Kras*-mutant/*Lkb1*-deficient organoids. **C)** Heatmap

of z-score transformed expression values of stem cell genes enriched in control organoids (top cluster) and

fetal genes enriched in *Lkb1*-deficient organoids (bottom cluster). **D)** GSEA of iCMS2 and iCMS3 gene

signatures (Joanito *et al.*, 2022) was performed on *Lkb1*-deficient organoids compared to control organoids.

The normalized enrichment score (NES) is depicted on the X-axis and the FDR is indicated by the size of the

dots. Line color reflects signature enrichment in *Lkb1*-deficient (red) or *Lkb1*-deficient/*Kras*-mutant (blue)

organoids. Dot shape (open/closed) indicates significant enrichment (FDR < 0.05).

Materials and Methods
755
Organoid culture
Small intestinal mouse organoids were established from crypts isolated from the small intestine of wild-type
mice, as previously described (Sato *et al.*, 2009). Basal culture medium for mouse organoids contained
advanced DMEM/F12 (Gibco) supplemented with 1% Penicillin-Streptomycin (Sigma-Aldrich), 10 mM
HEPES (Gibco), 1% Ultraglutamine 1 (Lonza), 1% B27 (Sigma-Aldrich), 0.5% N2 (Sigma-Aldrich) and 1 mM
N-acetylcysteine (Sigma- Aldrich). Basal culture medium was further supplemented with murine recombinant
EGF (Peprotech), 2.5% R-spondin1 conditioned medium (homemade) and 10% Noggin conditioned medium
(homemade) to obtain the complete ENR (EGF-Noggin-Rspondin) medium. Organoids were grown in
Matrigel (Corning) or BME (Cultrex) and passaged every 6-8 days as previously described (Sato *et al.*,
2009); their medium was refreshed every 2-3 days. For growth factor withdrawal experiments, EGF was
removed from organoid culture medium and organoids were monitored for a total of 16 days. Bright field
images were acquired with the Invitrogen™ EVOS™ FL Color Imaging System (Thermo Fisher Scientific).
768
769 **Vectors and sgRNAs**
770 The pSpCas9(BB)-2A-Puro plasmid was obtained from Addgene (48139). Single guide RNAs (sgRNAs) and
771 are listed in **Table S4**. Cloning of sgRNAs into the The pSpCas9(BB)-2A-Puro plasmid was performed
772 according to previously published protocols (Ran *et al.*, 2013) and vector sequences conformed by Sanger
773 sequencing.
774
Organoid electroporation
For electroporation, mouse organoids were expanded in ENR medium without antibiotics and supplemented
with 50% Wnt3a-conditioned medium (homemade) and 1mM Nicotinamide (Sigma-Aldrich). Electroporation
of organoids was performed as previously described using a NEPA21 (NEPAGENE) electroporator (Fujii *et*
al., 2015). To obtain *Lkb1*-deficient organoids, a total of 20 μg of DNA (10 μg gRNA, 7.2 μg of pPB-CAG-
rtTA_Hygromycin and 2.8 μg PiggyBac transposase) was used. To obtain *Apc* and *Kras* mutant organoids a
total of 10 μg DNA (gRNA and gRNA with oligonucleotide respectively) was used. 100 μg/mL Hygromycin B
was used to select *Lkb1*-deficient organoids and Noggin and Egf were omitted from the culture medium to
select for *Apc* and *Kras* mutant organoids respectively. After selection, single organoids were manually
picked for clonal expansion and then genotyped. Genomic DNA was isolated using the DNA micro kit
(Qiagen) according to the manufacturer's instructions and PCR amplification was performed using the
GoTaq Flexi DNA polymerase kit (Promega). Organoid genotype was confirmed by Sanger sequencing.
Primers are listed in **Table S5.**
788
Protein extraction and Western blotting
Organoids were washed in cold PBS and collected in cell recovery solution (Corning) according to the
manufacturer's instructions. Whole cell lysates were obtained by direct lysis in RIPA buffer (50 mM Tris- HCl
pH 8.0, 150 mM NaCl, 0.1% SDS, 0.5% Na-Deoxycholate, 1% NP-40) containing complete protease
inhibitors (Sigma-Aldrich) for 30 minutes on ice. Proteins were run on SDS–PAGE gels and transferred to

PVDF membranes, followed by blocking with Odyssey blocking buffer (LI-COR). The antibodies used are
described in **Table S6**. Blots were imaged with the Amersham Typhoon (GE Healthcare Life Sciences).
796

Immunofluorescence

Organoids were seeded in 15-well Ibidi chambers after passaging and fixed 3-4 days later in 4%
Paraformaldehyde (Sigma-Aldrich) for 60 minutes at room temperature. After fixation organoids
were washed with cold PBS and permeabilized with PBD0.2T (48,5 mL PBS, 1 mL 10%Triton, 0,5mL DMSO
and 0,5 g BSA) for 20 minutes at room temperature, after which they were incubated with primary antibodies
diluted in PBD0.2T buffer at 4°C overnight. Washes were performed PBD0.2T buffer and secondary
antibodies incubated 2-3 hours at room temperature. All antibodies are described in **Table S7**. Quantification
of the nuclear fraction of Yap was analyzed using the JACoP Fiji plugin. EdU incorporation assays were
performed by culturing organoids in ENR medium containing 10 µM Edu for 6 hours to cells, after which
single organoids were harvested, fixed in 4% PFA and EdU-labeling was performed according to
manufacturer instructions (Click-iT EdU Cell Proliferation Kit for Imaging, Thermo Fisher). Images were
acquired with a Zeiss LSM700 or LSM880 confocal microscope.
809

Patient tissue samples and counting

Patient material was collected from the pathology archives of Dutch hospitals (UMC). In total five PJS
patients were included in this study. Five normal controls were randomly selected. In 10 microscopically
normal looking crypts MUC2$^+$ cells were counted from the bottom toward to the top of the crypt. Only
longitudinally oriented crypts that were completely visible on the slide, were counted. Cell positions were
defined as the bottom of the crypt is cell position one (1), going upward to cell position +2, +3, and so forth till
cell position 30. Within the part of the crypt between cell position 1 and 30, the number of positively labelled
cells were counted. Also the total cell position was determined by adding up each cell position of a labelled
cell and average cell position of the labelled cells was determined by adding up each cell position of a
labelled cell and dividing the sum by the total number of labelled cells counted.
820

Immunohistochemistry

Fresh mouse small intestinal organoids were fixed in 4% Formaldehyde for 1 hour followed by dehydration in
25%, 50% and 70% ethanol and embedding in paraffin. Briefly, 4µm FFPE organoid or PJS patient tissue
sections were obtained, deparaffinised, blocked in 0.3% H2O2 in methanol for 20 minutes. Antigen retrieval
was performed in Tris/EDTA buffer (10 mM/1 mM; pH 9.0) or Sodium-citrate buffer (0,01 M/pH6.0) for 20
minutes at 100°C. Nonspecific binding sites were blocked using Protein Block Serum free (DAKO) for 10
minutes, followed by primary antibody incubation. Antibody binding was visualized using the using
BrightVision Goat anti Rabbit / Poly-AP and BrightVision Goat anti Mouse / Poly-HRP (VWR/ImmunoLogic),
with DAB as chromogen (Sigma-Aldrich). Sections were counterstained with Haematoxylin or neutral red and
mounted using Pertex. Primary antibodies are described in **Table S8**.
831

Electron microscopy

Organoids in Matrigel were fixed in a solution of 2% paraformaldehyde and 2.5% glutaraldehyde in 0.1 M
phosphate buffer (pH 7.4) overnight at 4°C. On the following day, organoids were washed twice in 0.1 M

phosphate buffer (pH 7.4) and subsequently post-fixed with 1% Osmium tetroxide for 2 hours at 4°C.

Organoids were then stained with 0.5% uranyl acetate 1 hour at 4°C in the dark. Matrigel droplets were

transferred into glass vials for dehydration with increasing acetone concentrations according to the following

protocol: 70% overnight at 4°C, 90% for 15 minutes at RT, 96% for 15 minutes at RT, and 100% 3 times for

30 minutes each at RT. After dehydration, samples were embedded in EPON resin. Ultrathin sections of 65

nm were obtained with a Leica UC7 microtome and stained with uranyl and lead citrate. Electron microscopy

was performed in the Cell Microscopy Core (CMC) of the UMC Utrecht. Sections were imaged with a JEOL

JEM-1010 transmission electron microscope (TEM).

Cell viability assay

For the organoid reconstitution assays, organoids were dissociated to single cells after incubation with

TrypLE (Life Technologies) for 5 min at 37°C. Dissociated organoids were passed through a 40 µm strainer

and counted. Cells were seeded at a density of 5000 cells/10µl Matrigel and cultured in ENR medium with 10

µM Y-27632 (Sigma-Aldrich) for 7 days. Organoids were then lysed with CellTiter-Glo® (Promega) according

to the manufacturer's protocol. ATP produced by viable cells was measured by a Berthold luminometer

Centro LB960.

Bulk RNA sequencing and data analysis

Organoids were lysed in RLT lysis buffer (Qiagen) 3-5 days after passaging. Biological triplicates were

employed. RNA was obtained using the Qiagen QiaSymphony SP (Qiagen) according to the manufacturer's

protocol. Single-end RNASeq reads were processed using the UMCU RNASeq pipeline

(https://github.com/UMCUGenetics/RNASeq, v2.3.0) with default settings. Read quality was assessed by

FastQC (0.11.4) followed by splice-aware alignment against the mouse reference genome (GRCm38) with

STAR (2.4.2a). RNA expression quantification was performed with htseq-count (0.6.0) in reverse-stranded

mode. Differential gene expression analysis was performed with the DESeq2 (1.22.2) R package. In each

comparison, genes were selected with an absolute fold-change > 2 and alpha P > 0.05. Data clustering and

generation of heatmaps was performed with pheatmap. Gene Set Enrichment Analysis was performed using

GSEA (v4.2.0, Windows App) and visualized using ggplot2. Intestinal cell-type specific gene sets for GSEA

were obtained from Extended Figure Table 3 of Haber et al., 2017 (Haber *et al.*, 2017). The other datasets

employed in GSEA were derived from previously published studies (Mustata *et al.*, 2013; Gregorieff *et al.*,

2015; Nusse *et al.*, 2018; Yui, Azzolin, Maimets, Pedersen, Robert P. Fordham, *et al.*, 2018; Ayyaz *et al.*,

2019; Chen *et al.*, 2021; Qu *et al.*, 2021b). Human gene sets were converted to their respective mouse

homologs using biomaRt (2.50.3) in R where necessary.

Single cell RNA sequencing

Sample preparation and sequencing: Single cells from organoid were harvested using TrypLE incubation at

37 °C, four days after passaging. Cells were filtered through a 40 µm nylon cell strainer (Falcon). scRNA-seq

was performed according to the Sort-seq protocol (Muraro *et al.*, 2016). Libraries were sequenced on an

Illumina NextSeq500 at paired-end 60- and 26-bp read length and 75,000 reads per cell.

Mapping and filtering: The Sharq pipeline was used to process the sequencing data (Candelli *et al.*, 2018).

Mapping was performed using STAR (version 2.6.1), on the Genome Reference Consortium Mouse built 38

(GRCm38). Read assignment was with feature Counts version 1.5.2 using a gene annotation based on

GENCODE version M14.

External RNA controls (ERCCS) and transcripts mapping to the mitochondrial genome were removed from

all cells. Low quality cell barcodes having either less than 800 unique transcripts, 200 genes 33.3%

mitochondrial transcripts or over 4-fold non-exonic to exonic reads were excluded, as well as barcodes with

over 100,000 unique transcripts.

scRNA-seq analysis: Unique transcript counts were normalized using sctransform (Hafemeister and Satija,

2019) and analyzed using the Seurat R package (version 4.1.0) (Butler *et al.*, 2018). From the top 3000

variable genes, genes associated with cell cycle phase, dissociation stress (heat shock and chaperone

proteins according to GO:0006986), sex (Xist, Tsix, and Y chromosome-specific genes), and activity

(ribosomal protein genes according to GO:0022626) were removed from the list of variable genes to avoid

biases in cell clustering, as described before (Hanemaaijer *et al.*, 2021). The first 20 principal components

were used both to calculate dimensionality reduction using UMAP and to perform clustering with a resolution

of 1, k.param = 10 and the Louvain algorithm.

Supplementary information

Gene	Sequence
Lkb1 gRNA1	AGCTTGGCGCGTTTGCGGCG
Lkb1 gRNA2	TGAGGATCTTGACCGCCCTG
Apc	TGTCTGGCTCCGGTAAGTGA
Kras	CTTGTGGTGGTTGGAGCTGG

892 **Table S4: gRNA sequences**

894 **Table S5: Genotyping primers**

Gene	Forward primer	Reverse Primer
Lkb1 gRNA1	GGGGAAAATCAAAAGTGAAGAA	CCCTCTAGCTGCGTAAACAAAC
Lkb1 gRNA2	CTCCACCGAGGTAATCTACCAGCCG	TCTAACCGTTTTCTCTTCTCATTTCC
Apc	ATAGAAACAGCACTGACCCAAATTTCA	AGGCCTCTTTGCTTAGAGCTTTCATAA
Kras	TGGCTCCAACACAGATGTTC	GGATGGCATCTTGGACCTTA

896 **Table S6: Western Blot antibodies**

Antibody	Dilution	Supplier	Cat. number
Anti-LKB1	1:1000	Santa Cruz	sc-32245
Anti-GAPDH	1:8000	Millipore	CB1001
Anti-mouse Alexa 800	1:5000	LI-COR	926-32211
Anti-rabbit Alexa 680	1:5000	LI-COR	925-68070

898 **Table S7: Immunostaining antibodies**

Antibody	Dilution	Supplier	Cat. number
Anti-Muc2	1:200 (IF) 1:1000 (IHC)	Santa Cruz	sc-515032
Anti-Lysozyme1	1:200 (IF) 1:2000 (IHC)	Dako	A0099
Anti-ANXA1	1:200 (IF) 1:1000 (IHC)	Atlas Antibodies	HPA011271
Anti-YAP	1:200 (IF)	Santa Cruz	sc-101199
Anti-Cleasved-Caspase-3	1:200 (IF)	Cell Signalling	9661
Anti-rabbit Alexa Fluor 488	1:300 (IF)	Invitrogen	A11034
Anti-mouse Alexa 488	1:300 (IF)	Invitrogen	A11029
DAPI	1:1000 (IF)	Sigma	D9542
Click-iT™ EdU Alexa Fluor™	n.a.	Invitrogen	C10424

647			
Alexa Fluor™ 647 Phalloidin	1:500 (IF)	Invitrogen	A22287

Data availability

901 The scRNA-seq and bulk RNA-seq data generated in this study will be uploaded to Gene Expression
902 Omnibus (GEO) upon publication. If needed earlier, data can be requested via the corresponding author.
903

Author contributions

905 S.F.P, H.M., A.M. and M.M.M., conceived and designed the experiments. S.F.P, H.M., A.M., D.X., F.H.M.M.,
906 C.J. and A.R.K performed the experiments. S.F.P, H.M., A.M., J.Y.V., J.S., T.M., T.C., R.S., J.R., F.C.P.H.,
907 G.J.A.O. and M.M.M. analyzed the data. W.W.J.L. and G.J.A.O. provided patient material. S.F.P, H.M., A.M.,
908 J.Y.V. and M.M.M. wrote the manuscript, which was reviewed by all authors.
909

Disclosure and competing interest statement

911 M.M.M. is an inventor on patents related to membrane protein degradation; she is co-founder and
912 shareholder of Laigo Bio. J.R. is co-founder of Cyclomics BV., a genomics company.
913

Acknowledgements

The authors thank members of the Maurice laboratory for helpful discussions and suggestions. PiggyBac
Transposase and pPB-CAG-rtTA_Hygromycin were a kind gift from B.K. Koo. This work is part of the
Oncode Institute, which is partly financed by the Dutch Cancer Society. This work was supported by the
China Scholarship Council (to H.M.), the Dutch Cancer Society KWF Grant 13112 (to M.M.M.), Zon-MW VICI
grant 91815604 (to M.M.M.), Zon-MW TOP grant 91218050 (to M.M.M.) and NWO Gravitation project
IMAGINE! (to M.M.M.).

Figure 1

Figure 2

Figure 3

Figure 4

A

	WT	Lkb1[+/-]	Lkb1[bud -/-]	Lkb1[cys -/-]
DAPI				
Yap				
Merge				

B

Figure 5

Figure 6

Figure 7

Figure S1

Figure S2

Figure S3

| WT | Lkb1[+/-] | Lkb1[bud -/-] | Lkb1[cys -/-] |

Yap-responsive genes

Amotl1
Ctgf
Clu
Anxa1
Cyr61
Ereg
Areg

2
1
0
−1
−2

Figure S4

Figure S5

Figure S6

A

| Apc⁻ᐟ⁻ | Lkb1⁺ᐟ⁻ Apc⁻ᐟ⁻ | Lkb1ᵇᵘᵈ⁻ᐟ⁻ Apc⁻ᐟ⁻ | Lkb1ᶜʸˢ⁻ᐟ⁻ Apc⁻ᐟ⁻ |

B

TGGCACCCTCA-CTTACCGGAGCCAGACAAATAC
ACCGTGGGAGT-GAATGGCCTCGGTCTGTTTATG

Apc⁻ᐟ⁻
(p.Y620LfsX630)

TGGCACCCTCACCTTACCGGAGCCAGACAAATAC

Lkb1⁺ᐟ⁻ Apc⁻ᐟ⁻
(p.Y620LfsX630)

TGGCACCCTCACCTTACCGGAGCCAGACAAATAC

Lkb1ᵇᵘᵈ⁻ᐟ⁻ Apc⁻ᐟ⁻
(p.T619PfsX628)

TGGCAC-----CTTACCGGAGCCAGACAAATAC

Lkb1ᶜʸˢ⁻ᐟ⁻ Apc⁻ᐟ⁻
(p.Y620GfsX625)

TGGCACC-----CTTACCGGAGCCAGACAAATAC

Lkb1ᶜʸˢ⁻ᐟ⁻ Apc⁻ᐟ⁻
(p.T619LfsX629)

TGGCACCCT---CTTACCGGAGCCAGACAAATAC

C

| KrasG12D | Lkb1⁺ᐟ⁻ KrasG12D | Lkb1ᵇᵘᵈ⁻ᐟ⁻ KrasG12D | Lkb1ᶜʸˢ⁻ᐟ⁻ KrasG12D |

D

AAACTTGTGGTGGTTGGAGCTGGTGGCGTAGGC
TTTGAACACCACCAACCTCGACCACCGCATCCG

KrasG12D
(p.G12D)

AAACTTGTGGTGGTTGGAGCAGATGGCGTAGGC

Lkb1⁺ᐟ⁻ KrasG12D
(p.G12D)

AAACTTGTGGTGGTTGGAGCAGATGGCGTAGGC

Lkb1ᵇᵘᵈ⁻ᐟ⁻ KrasG12D
(p.G12D)

AAACTTGTGGTGGTTGGAGCAGATGGCGTAGGC

Lkb1ᵇᵘᵈ⁻ᐟ⁻ KrasG12D
(p.A11WfsX29)

AAACTTGTGGTGGTTGGA--TGGTGGCGTAGGC

Lkb1ᶜʸˢ⁻ᐟ⁻ KrasG12D
(p.G12D)

AAACTTGTGGTGGTTGGAGCAGATGGCGTAGGC

E

WT   Apc⁻ᐟ⁻   KrasG12D

Stem / Fetal

Ascl2, Znf3, Axin2, Rnf32, Lrp4, Ly6d, Anxa1, Clu, Anxa3, Anxa8, Msln

Lkb1⁺ᐟ⁻
Lkb1ᵇᵘᵈ⁻ᐟ⁻
Lkb1ᶜʸˢ⁻ᐟ⁻

F

iCMS2

Lkb1⁺ᐟ⁻ vs. WT
Lkb1ᵇᵘᵈ⁻ᐟ⁻ vs. WT
Lkb1ᶜʸˢ⁻ᐟ⁻ vs. WT

NES

iCMS3

Lkb1⁺ᐟ⁻ vs. WT
Lkb1ᵇᵘᵈ⁻ᐟ⁻ vs. WT
Lkb1ᶜʸˢ⁻ᐟ⁻ vs. WT

NES

FDR
0.0
0.1
0.2
0.3
0.4

Enriched in
WT
Lkb1-deficient

Significant
⊕ No
● Yes