# Intestinal LKB1 loss drives a pre-malignant program along the serrated cancer pathway

Plugge S.F.[#,1], Ma H.[#,1,2], van der Vaart J.Y.[1], Sprangers J.[1], Morsink F.H.M.[2], Xanthakis D.[1], Jamieson C.[1], Keijzer A.R.[1], Margaritis T[3]., Candelli T.[3], Straver R.[1], de Ridder J.[1], Holstege F.C.P.[3, 4], de Leng W.W.J.[2], Offerhaus G.J.A.[2], Merenda A.*[,1], Maurice M.M.*[,1]

1. Center for Molecular Medicine and Oncode Institute, University Medical Center Utrecht, Utrecht, The Netherlands
2. Department of Pathology, University Medical Center Utrecht, Utrecht, The Netherlands
3. Princess Maxima Center for Pediatric Oncology, Utrecht, Utrecht, The Netherlands
4. Center for Molecular Medicine, University Medical Center Utrecht, Utrecht, The Netherlands
[#]These authors contributed equally
*Corresponding authors

## Abstract

Peutz-Jeghers syndrome (PJS) is a familial disorder caused by heterozygous inactivating Liver Kinase B1 (*LKB1/STK11)* mutations that induce hamartomatous polyp formation in the gastrointestinal tract. Additionally, PJS patients carry an elevated risk of developing malignancies in multiple tissues, including the intestine. How *LKB1* deficiency alters the hierarchical organization and phenotypical landscape of epithelial tissues to mediate an increased cancer risk remains poorly understood. Here, we investigate these issues using small intestinal organoids that carry heterozygous or homozygous *Lkb1* mutations. We show that *Lkb1* loss causes allele dosage-dependent activation of transcriptional programs for tissue repair, particularly within stem and transit amplifying cell lineages. Furthermore, enhanced EGFR ligand expression promotes niche-independent growth of *Lkb1*-deficient organoids. Strikingly, we uncover that mono-allelic loss of *Lkb1* is sufficient to shift the epithelium into the serrated premalignant program for colorectal carcinogenesis, which is strongly amplified upon loss-of-heterozygosity (LOH) or activating *Kras* mutations. We conclude that heterozygous mutations in LKB1 predispose the intestinal epithelium to uncontrolled growth along the serrated pathway, providing an explanation for the increased cancer risk in PJS patients.

## Introduction

The Peutz-Jeghers syndrome (PJS) is a hereditary genetic disorder that occurs in approximately 1:50.000 - 1:200.000 individuals worldwide (Beggs *et al.*, 2010). PJS patients suffer from a life-long development of benign hamartomatous polyps throughout their gastrointestinal tract for which they need periodical surgical resection. Moreover, patients carry an increased risk of developing early-onset epithelial tumors, particularly within the gastrointestinal (GI) tract (up to 66%), breast (up to 54%), pancreas (up to 36%) and gonads (Giardiello *et al.*, 2000; Hearle *et al.*, 2006; Beggs *et al.*, 2010; Van Lier *et al.*, 2010), resulting in a cumulative lifetime risk of 93% (Giardiello *et al.*, 2000). Of note, PJS patients are predisposed to malignancies of the small intestine (frequency 13%), a condition that is otherwise very rare (0.03%) (Van Lier *et al.*, 2010; Tomasetti and Vogelstein, 2015).

Heterozygous inactivating mutations in the tumor suppressor gene Liver Kinase B1 (*LKB1/STK11*)
were identified as the main cause of PJS (Hemminki *et al.*, 1998; Jenne *et al.*, 1998). Furthermore, *LKB1*
mutations are causally linked to a broad range of sporadic cancer cases, including lung carcinoma,
melanoma, pancreatic and cervical cancers (Giardiello and Trimbath, 2006; Sanchez-Cespedes, 2007).
*LKB1* encodes a serine/threonine kinase implicated in the regulation of a variety of critical cellular processes,
including energy metabolism, cell adhesion, DNA methylation and apicobasal polarity (Shackelford and
Shaw, 2009; Kline *et al.*, 2013; Pierce *et al.*, 2021). Loss of heterozygosity (LOH) in this gene is regarded as
a driving force of cancer development, as over 64% of PJS-derived carcinomas present with inactivation of
the second copy of the gene (Entius *et al.*, 2001; De Leng *et al.*, 2003).
Over recent years, modelling of PJS in murine models revealed that *Lkb1* loss in either
mesenchymal or T cell lineages induces development of gastro-intestinal polyps with strong resemblance to
those observed in PJS patients. (Katajisto *et al.*, 2008; Ollila *et al.*, 2018; Poffenberger *et al.*, 2018; Cotton *et*
*al.*, 2022). These results have shifted focus to non-epithelial roles of *Lkb1* in driving PJS disease
development, with an emphasis on processes underlying polyposis. If and how polyp formation in PJS
patients links to cancer predisposition however remains a heavily debated issue, as LOH and dysplastic
transformation of PJS polyps were reported to be very rare (Shepherd, Bussey and Jass, 1987; Entius *et al.*,
2001; De Leng *et al.*, 2003; Korsse *et al.*, 2013). These observations differ from other polyposis syndromes,
such as familial adenomatous polyposis (FAP), where epithelial loss of the underlying germline mutation in
the adenomatous polyposis coli gene APC and its role in cancer development has been well described
(Fodde, 2002).
Since *LKB1* loss mediates predisposition to carcinoma development in multiple epithelial tissues
irrespective of polyp formation, we hypothesized that epithelial-intrinsic alterations may sensitize these
tissues to cancer formation. Previous studies indicate that intestinal epithelial loss of *LKB1* does not lead to
polyp formation while alterations in secretory cell fate specification were identified (Shorning *et al.*, 2009;
Gao *et al.*, 2020; Cotton *et al.*, 2022; Kim *et al.*, 2022). Despite these advancements, how epithelial *LKB1*
loss changes the phenotypic landscape of the intestinal epithelium and how these alterations may link to
cancer predisposition in PJS patients remains poorly understood.
Here, we model *mono-* and *bi*-allelic inactivation of *Lkb1* in small intestinal organoids to investigate
how *Lkb1* loss alters the dynamics of cellular identities and lineage relationships of the intestinal epithelium.
We uncover that *Lkb1* loss promotes features of epithelial transformation linked to the serrated pathway for
intestinal carcinogenesis. These features include loss of classical adult stem cell populations, activation of
transcriptional programs related to regeneration and intestinal-to-gastric cellular transdifferentiation, as well
as the acquirement of niche-independent growth properties. Our findings argue that *Lkb1* loss mediates
epithelium-wide activation of a pre-cancer program akin to sporadic serrated precursor lesions, providing an
explanation for the increased cancer risk observed in PJS patients. By linking *Lkb1* loss to a major pathway
for colorectal cancer development, these findings pave a path for the development of new strategies for
prevention, surveillance and precision treatment of PJS tumors as well as other malignancies harboring
*LKB1* inactivating mutations.

**Results**

***Lkb1* bi-allelic loss results in phenotypic heterogeneity and increased self-renewal potential of small**

**intestinal organoids.**

To study the role of *Lkb1* loss in the intestinal epithelium, we employed CRISPR/Cas9 technology to

generate *Lkb1*-deficient mouse small intestinal organoid lines. To obtain homogenous organoid cultures that

are either heterozygous or homozygous for *Lkb1* loss, we isolated single organoids from bulk cultures and

expanded these as individual clones. After genotyping, we selected one *Lkb1*$^{+/-}$ and two *Lkb1*$^{-/-}$ clonal lines

for further characterization (Figure S1A, Figure 1A-B). *Lkb1*$^{+/-}$ organoids were morphologically similar to wild

type (WT) organoids, with typical budding structures that reflect a crypt-villus organization (Sato *et al.*, 2009)

(Figure 1A). Remarkably, the introduction of homozygous *Lkb1* loss resulted in phenotypic bifurcation, with

clonal lines displaying either a budding or a cystic growth morphology (Figure 1A). Notably, from day 3 after

passaging, budding *Lkb1*$^{-/-}$ (*Lkb1*$^{bud\,-/-}$) organoids accumulated clusters of dead cells in their periphery (Figure

1A), as indicated by their positive staining for cleaved caspase 3 (Figure S1B). These observations are

different from *Lkb1*$^{+/-}$ and WT organoids that mainly accumulate dead cells in their lumen (Figure S1B) (Sato

*et al.*, 2009).

Next, we tested whether *Lkb1* loss alters the ability of intestinal organoids to reconstitute organoids

from single cells, as a proxy for their self-renewal capacity. While budding WT and *Lkb1*-mutant organoids

lines displayed similar outgrowth rates, cystic *Lkb1*$^{-/-}$ (*Lkb1*$^{cys\,-/-}$) organoids carried a significantly improved

capacity to form organoids upon seeding of single cells (Figure 1C). Regardless of these observations, both

homozygous mutant morphologies displayed a similar increase in the number of proliferative cells in

comparison to heterozygous and wild-type organoids, as shown by EdU incorporation (Figure 1D, E). To rule

out that changes in organoid morphology of the homozygous mutants are the result of CRISPR off-target

effects, we used an alternative gRNA to introduce *Lkb1* mutations, which resulted in an identical phenotypic

heterogeneity of *Lkb1*$^{-/-}$ organoid lines (Figure S1C-E). Moreover, these newly generated *Lkb1*$^{cys\,-/-}$ clones

also displayed superior reconstitution potential compared to their homozygous budding counterparts (Figure

S1F). These findings reveal that biallelic *Lkb1* loss results in phenotypic heterogeneity within the intestinal

epithelium, and that homozygous *Lkb1*-mutant organoids with cystic morphology display an increased

recovery potential.

***Lkb1* loss alters the number, morphology and positioning of secretory cells.**

To investigate whether our *Lkb1*-mutant organoid lines serve as representative models for intestinal

epithelial loss of *Lkb1 in vivo*, we examined alterations in localization and morphology of Paneth (Lyz$^+$) and

goblet (Muc2$^+$) cells that were previously reported for the *Lkb1*-mutant mouse intestine (Shorning *et al.*,

2009; Gao *et al.*, 2020). Compared to WT organoids, the localization of Lyz1$^+$ cells in *Lkb1*$^{+/-}$ and *Lkb1*$^{bud\,-/-}$

organoids was not restricted to the bottom of the crypt but displayed a more scattered distribution that

reached into the villus domain (Figure 2A). In addition, while Muc2$^+$ cells in WT and *Lkb1*$^{+/-}$ organoids were

present in modest numbers and always located to the villus region, Muc2$^+$ cell numbers were increased in

*Lkb1*$^{bud\,-/-}$ organoids and found scattered throughout the crypt-villus axis (Figure 2B). These observations are

highly reminiscent of the reported aberrancies in Paneth and Goblet cell numbers and localization in the

*Lkb1*-deficient mouse intestine *in vivo* (Shorning *et al.*, 2009). By contrast, *Lkb1*$^{cys\,-/-}$ organoids were devoid

of Lyz1$^+$ cells but did allow for formation of Muc2$^+$ cells that are usually lacking in cystic organoids (Sato, Es,

*et al.*, 2011), indicating that secretory lineage specification is also altered within these organoids (Figure 2A-

B).

We next investigated how these findings relate to intestinal tissue organization of human PJS

patients. While the average number and location of MUC2$^+$ cells per intestinal crypt was similar for healthy

donors and the non-transformed epithelium of PJS patients (Figure 2C, Figure S2A-B), LYZ1$^+$ cells were

clearly mislocalized within the PJS mucosa (Figure 2D). In the healthy intestine, LYZ1$^+$cells localized strictly

to the crypt base in the healthy intestine, as expected (Barker *et al.*, 2007). In the non-transformed

epithelium of PJS patients, however, LYZ1$^+$ cells frequently scattered towards higher regions of the crypt

(Figure 2D). Thus, heterozygous loss of *LKB1* in PJS patient-derived intestines leads to mislocalization of

LYZ1$^+$cells but does not alter MUC2$^+$ cell distribution, similar to our observations in *Lkb1$^{+/-}$* organoids.

To examine *Lkb1* loss-induced alterations in secretory cell morphology at the ultrastructural level, we

employed electron microscopy. In WT intestinal organoids, Paneth cells displayed a typical morphology with

prominent apical accumulation of spherically-shaped secretory granules that are uniformly filled with

electron-dense content (Figure 2E) (Satoh *et al.*, 1990; Sato *et al.*, 2009). By contrast, secretory granules of

*Lkb1$^{+/-}$* Paneth cells commonly contained an abnormally 'sausage-shaped' electron-dense core (Figure 2E).

In *Lkb1$^{bud-/-}$* organoids, the content of Paneth cell-secretory granules was even more aberrant, displaying a

small electron-dense core surrounded by a peripheral electron-lucent halo (Figure 2E). In WT intestinal

organoids, columnar-shaped goblet cells were identified with large translucent granules filled with mucus

(Figure 2F) (Troughton and Trier, 1969; Bansil and Turner, 2018). Compared to WT goblet cells, *Lkb1$^{+/-}$*

mucus-secreting cells displayed a rounded and more flattened morphology while *Lkb1$^{bud\ -/-}$* goblet-like cells

display granules with an unusual electron dense core (Figure 2F). In addition, in *Lkb1$^{cys-/-}$* organoids the

content of these mucus-like granules was frequently lost during EM preparation, suggesting an altered mucin

composition, which might also explain the limited number of identified Muc2$^+$ cells (Figure 2F). The

morphology of these Paneth-like and goblet-like cells in *Lkb1*-deficient organoids is reminiscent of a

previously described population of 'intermediate' cells that were proposed to represent transition states

between undifferentiated and more mature secretory cells (Troughton and Trier, 1969). Together, these

observations suggest that *Lkb1* loss may interfere with the terminal differentiation of secretory cells.

Furthermore, previous studies identified roles of *LKB1* in apical brush border formation, using 2D human

(cancer) cell lines (Baas *et al.*, 2004). Inspection of our EM images however did not reveal any evidence for

loss of these subcellular structures in *Lkb1*-deficient organoids (Figure S2C).

Taken together, our findings reveal that engineering *Lkb1* loss in organoids induces highly similar

alterations in secretory cell number, localization and morphology as those observed in *in vivo* mouse models

and the intestine of PJS patients. These results thus confirm the suitability and relevance of organoids for

modelling epithelial alterations in PJS. Furthermore, our results indicate that *Lkb1* loss-induced alterations in

cell lineage specification and positioning are intrinsic to the epithelium and do not rely on interactions with

the mesenchyme.

***Lkb1* mono- and bi-allelic loss induces incremental expression of a regenerative gene signature.**

To examine how *Lkb1* loss affects epithelial organization, we determined transcriptional changes induced by

*Lkb1* loss in organoids using bulk mRNA sequencing. Compared to WT organoids, we observed differential

expression of 1298, 2312 and 7816 genes in *Lkb1*$^{+/-}$, *Lkb1*$^{bud\ -/-}$ and *Lkb1*$^{cys\ -/-}$ organoids, respectively

(Supplementary Table 1). Using gene set enrichment analysis (GSEA) we first examined the consequences

of *Lkb1* loss on well-defined gene signatures derived from individual intestinal cell types (Haber *et al.*, 2017).

In agreement with our earlier results, all *Lkb1*-deficient clones showed significant downregulation of gene

sets linked to mature Paneth cells, whereas goblet cell gene sets were enriched within both budding and

cystic *Lkb1*$^{-/-}$ organoids in comparison to WT organoids, again suggesting alterations in secretory cell lineage

specification (Figure 3A).

Noticeably, the *Lgr5*$^{+}$ adult stem cell signature (Haber *et al.*, 2017) was downregulated in all *Lkb1*-

deficient clones (Figure 3A). These data are reminiscent of the previously reported growth state of fetal

intestinal spheroids that relies on proliferation of *Lgr5*-negative progenitors (Mustata *et al.*, 2013). Indeed, we

uncovered a gradual enrichment of an embryonic spheroid gene signature in *Lkb1*-deficient organoids

(Mustata *et al.*, 2013), that correlated with the number of lost *Lkb1* alleles (Figure 3A-B). The transition from

adult stem cell to a fetal spheroid signature was most striking for *Lkb1*$^{cys\ -/-}$ organoids compared to *Lkb1*$^{bud\ -/-}$

organoids (Figure 3B). Previous studies indicated that a transient upregulation of a fetal-like intestinal gene

signature associates with epithelial remodeling during injury repair and wound healing (Fordham *et al.*, 2013;

Nusse *et al.*, 2018; Yui, Azzolin, Maimets, Pedersen, Fordham, Hansen, Larsen, Guiu, M. R. P. Alves, *et al.*,

2018; Wang *et al.*, 2019). In line with these observations, *Lkb1*-deficient organoid clones also exhibited

enrichment of gene signatures associated with intestinal regeneration response following damage or

infection (Figure 3A). To rule out that organoid morphology rather than *Lkb1*-deficiency mediates expression

of this regenerative program, we analyzed the transcriptomes of *Apc*$^{-/-}$ and Wnt-treated organoids that also

exhibit a cystic shape but are WT for *Lkb1* (Merenda, Fenderico and Maurice, 2020). In comparison to both

*Apc*$^{-/-}$ and Wnt-treated organoids, *Lkb1*$^{cys\ -/-}$ organoids were significantly enriched for expression of these

regenerative gene sets, confirming that *Lkb1* loss itself promotes a regenerative state (Figure S3A).

Furthermore, *Lkb1*-deficient clones, as well as control fetal intestinal organoids (E16.5), displayed increased

expression of the fetal marker Anxa1, while this marker was not detected in WT adult intestinal organoids

(Figure 3C) (Yui, Azzolin, Maimets, Pedersen, Fordham, Hansen, Larsen, Guiu, M. R. P. Alves, *et al.*, 2018).

In addition, a dysplastic region of human PJS patient-derived duodenal epithelial tissue showed patches of

ANXA1-positive epithelial cells, while non-transformed PJS and healthy donor duodenum epithelial cells

remained negative (Figure 3D). Together, these findings indicate that *Lkb1* loss converts the epithelium into

a permanent regenerative state. Strikingly, this conversion towards a tissue repair growth state is already

apparent upon monoallelic *Lkb1* loss, a representative condition for the PJS intestinal epithelium.

**Lkb1-deficient organoids show a Yap-induced regenerative state and display intra-epithelial self-**

**sufficiency for Egf.**

Over recent years, Yap signaling was placed central to intestinal regeneration (Gregorieff *et al.*, 2015).

Additionally, *LKB1* was uncovered previously as a negatively regulator of the Hippo-Yap pathway in an

siRNA kinome screen (Mohseni *et al.*, 2014). We therefore wondered if *Lkb1* loss may activate Yap signaling

as part of the induced regenerative program in intestinal epithelial organoids. Indeed, we identified enhanced

expression of a Yap-regulated gene signature linked to intestinal regeneration (Gregorieff *et al.*, 2015) in

*Lkb1*-mutant organoids (Figure 3A). Notably, increased Yap target gene expression was already identified in

heterozygous *Lkb1*-mutant conditions and progressively increased with loss of both *Lkb1* alleles (Figure

S3B). Again, *Lkb1*$^{cys\ -/-}$ organoids showed the strongest induction of Yap target genes, in line with their robust

expression of a regenerative transcriptional program (Figure S3B). Immunofluorescence staining further

confirmed an increase in Yap protein expression in both budding and cystic *Lkb1*$^{-/-}$ organoids, and a

prominent nuclear localization of Yap, particularly within *Lkb1*$^{cys\ -/-}$ organoids (Figure 4A-B).

YAP pathway activation is well-known to induce enhanced expression of EGF family ligands,

particularly in the context of regeneration (Zhang *et al.*, 2009; Ren *et al.*, 2010; Yang *et al.*, 2012; Gregorieff

*et al.*, 2015). We therefore wondered whether *Lkb1*-deficient organoids display altered state of EGF pathway

dependency. Noticeably, we found that a combined set of the *Egf* ligand family was upregulated in *Lkb1*-

deficient organoids, in a gradual manner along with *Lkb1* allelic loss and alterations in organoid morphology

(Figure 5A). Furthermore, these changes were accompanied with an elevated expression of *Egf* receptor

family members (Figure 5B). As survival and growth of healthy homeostatic adult intestinal organoids

depends on supplementation with Egf, we investigated whether the enhanced expression of *Egf* family

ligands and its receptors in *Lkb1*-deficient organoids mediates niche factor self-sufficiency, by omitting Egf

from the culture medium. In contrast to WT organoids, all *Lkb1*-deficient organoids were capable of growing

without Egf supplementation (Figure 5C). In support of these findings, we confirmed that organoid survival

still depended on EGFR signaling, since *Lkb1*-deficient organoids were sensitive to Gefitinib, a potent EGFR

inhibitor (Figure 5D).

Together, these results demonstrate that *Lkb1* loss induces a constitutive Yap-induced regenerative

state and drives epithelial self-sufficiency for EGF ligand production. As niche-independent growth properties

are considered a hallmark of cancer progression (Fujii *et al.*, 2016), these results provide a potential

explanation for a growth advantage of *Lkb1*-deficient epithelial cells and the initiation of tumorigenesis.

**_Lkb1_-deficiency mediates reprogramming of intestinal stem cells into a regenerative state.**

To perform an in-depth analysis of how *Lkb1* loss alters intestinal lineage specification and cellular

hierarchies, we subjected WT and *Lkb1*-deficient organoids to single cell RNA sequencing. We included an

analysis of mouse fetal intestinal organoids, to explore a potential overlap with fetal gene expression.

Unsupervised clustering of single cells identified 19 distinct clusters (Figure S4A). A clear separation of

cellular profiles derived from *Lkb1*$^{cys\ -/-}$ organoids, fetal and WT cells was observed, while *Lkb1*$^{bud\ -/-}$ and

*Lkb1*$^{+/-}$ organoids displayed closer resemblance to WT cells (Figure 6A). The separation of *Lkb1*$^{cys\ -/-}$ and fetal

cell clusters in UMAP indicates that *Lkb1*$^{cys\ -/-}$ organoids do not fully convert to a fetal state but only show

partial expression of a fetal-like gene program (Figure 6A). Furthermore, *Lkb1*$^{cys\ -/-}$ organoids are composed

of a rather homogenous population of cells that are different from all budding organoid clusters and do not

display a known intestinal cellular identity. We therefore restricted annotation of intestinal cell types to

budding organoid lines, using publicly available intestinal cell type gene signatures (Figure 6C, Figure S4B-I)

(Haber *et al.*, 2017). As the gene signatures for Paneth and goblet cells mapped to the same cluster (Figure

S4E, S4H), we marked this cluster as secretory progenitors, similar to previous reports (He *et al.*, 2022). Cell

type fraction analysis revealed that *Lkb1*$^{bud\ -/-}$ organoids comprise an increased fraction of secretory

progenitor cells compared to WT organoids, confirming alterations in secretory cell differentiation (Figure

6B). Interestingly, the increase in secretory progenitors was accompanied with a decrease in enterocytes,

suggesting that the preferred differentiation trajectory in *Lkb1*-deficient budding organoids is shifted from

absorptive towards secretory lineages (Figure 6C). The previously observed morphological aberrancies in

secretory lineage specification were confirmed at the transcriptional level, as *Lkb1*-deficient organoids

displayed the occurrence of 'intermediate' cells that co-express features of both Paneth and goblet cells

(Figure 6D). Moreover, expression of Egf ligands was specifically increased in *Lkb1*-deficient

enteroendocrine and secretory progenitor cell populations (Figure 6E) suggesting that these cells might be

responsible for the observed niche-independency of *Lkb1*-mutant organoids.

As expected, expression of the fetal and regenerative marker *Anxa1* is highly abundant in both fetal

and *Lkb1$^{cys\,-/-}$* organoids (Figure 6F). Notably, we found that *Anxa1* expression in *Lkb1$^{bud\,-/-}$* organoids is

induced mainly within stem and transit amplifying (stem/TA) cell clusters, indicating that this population is the

main contributor to the observed epithelial conversion into a regenerative state (Figure 6A, F). To further test

this assumption, we performed reclustering of stem/TA cell populations including all *Lkb1$^{cys\,-/-}$* cells.

Unsupervised clustering indicates distinct clusters for WT, *Lkb1$^{+/-}$* and *Lkb1$^{bud\,-/-}$* stem/TA cells (Figure S4J-

K). To examine the effect of *Lkb1* loss on intestinal stem cells, we performed GSEA on genes differentially

expressed between WT stem cells and *Lkb1*-deficient stem cells (Supplementary Table 2). We observed

enrichment of goblet cell gene expression within *Lkb1*-deficient stem cell populations compared to WT

organoids (Figure 6G-H). This suggests that the capacity for preferential differentiation towards secretory

cells is already present in *Lkb1*-deficient stem cells. Furthermore, we identified an enrichment of four out of

six previously reported fetal and regenerative gene signatures, a depletion of the *Lgr5$^+$* stem cell signature

and an increased expression of Anxa1 in *Lkb1*-deficient stem cells (Figure 6G, I). These findings strengthen

a model in which induction of a regenerative response by *Lkb1* loss originates from the stem cell

compartment of the intestinal epithelium.

In conclusion, our results argue that *Lkb1*-deficient stem/TA cells reside in a regenerative state and

preferentially differentiate towards secretory rather than absorptive cell lineages. As a result, secretory cells

start secreting EGF ligands to drive niche-independent growth of *Lkb1*-deficient epithelium.

***Lkb1* loss drives a pre-malignant program along the serrated pathway which is amplified by**

**additional *Kras* mutations.**

Next, we aimed to obtain insight in how transcriptional reprogramming induced by *Lkb1* loss may promote

cancer predisposition. Colorectal cancer (CRC) develops mainly via two pathways that involve distinct

originating lesions and mutational events (Chen *et al.*, 2021). The classical adenoma-carcinoma sequence is

initiated by bi-allelic mutations in the tumor suppressor APC which drives formation of stem cell-enriched

adenomatous polyps that collect additional mutations (e.g. in *KRAS* and *TP53*) to develop into microsatellite-

stable CRC characterized by chromosomal instability (CIN) (Fearon and Vogelstein, 1990). Recently, an

alternative pathway emerged, in which colorectal cancer arises from serrated precursor polyps that are

initiated by *BRAF* or *KRAS* mutations and display combined features of fetal gene expression, differentiated

cells and gastric metaplasia (Patai *et al.*, 2013; Chen *et al.*, 2021). Whether and how *Lkb1* loss drives entry

into one of these pathways for CRC malignant progression is not known.

To investigate this issue, we examined expression of gene sets linked to the conventional and

serrated pathways of CRC development in *Lkb1*-deficient organoids (Chen *et al.*, 2021). Strikingly, all *Lkb1*-

deficient organoids display significant enrichment of genes that were previously found to mark neoplastic

populations of serrated-specific cells (SSC) (Figure 7A, Figure S5A-B). These effects gradually increased

with the number of lost *Lkb1* alleles, along with a simultaneous decrease in the expression of adenoma-

specific genes (ASC) (Figure 7B). Single cell analysis further revealed that the serrated gene program was
selectively increased in *Lkb1*-deficient stem cell populations (Figure S5C-E). Furthermore, *Lkb1*$^{cys\ -/-}$
organoids lose expression of the intestinal transcription factor *Cdx2* and upregulate expression of common
gastric markers like *Aqp5*, *Tacstd2* [*Trop2*], *Tff2*, and *Msln* (Figure 7B, Supplementary Table 1), indicating
that *Lkb1* loss promotes intestinal-to-gastric cellular transdifferentiation, often referred to as gastric
metaplasia (Balbinot *et al.*, 2018; Chen *et al.*, 2021).

To investigate how an *Lkb1*-deficient epithelial state cooperates with common cancer mutations in
CRC, we introduced *Apc* or *Kras* mutations in *Lkb1*-mutant organoids. Truncating *Apc* mutations are linked
to the classical adenoma sequence, while *Kras* mutations are observed in both pathways (Fearon and
Vogelstein, 1990; Patai *et al.*, 2013). As expected, based on earlier reports, introduction of a truncating *Apc*
mutation induced cystic morphologies of all organoids (Figure S6A-B) (Sato, Stange, *et al.*, 2011; Dow *et al.*,
2015). Surprisingly, however, introduction of a G12D *Kras* mutation in *Lkb1*-deficient organoids also
consistently mediated transition to a cystic morphology, while single *Kras*-mutant organoids maintained their
crypt-villus organization (Figure S6C-D). Functionally, *Kras* mutations conveyed synergistic effects on the
ability of single *Lkb1*$^{-/-}$ cells to reconstitute organoid growth (Figure 7C). By contrast, *Apc* mutations did not
add any significant synergy to the reconstitution capacity of *Lkb1*$^{-/-}$ organoids (Figure 7D).

To investigate how *Apc* or *Kras* mutations alter the phenotypic landscape of intestinal epithelial
organoids that already lost one or two copies of *Lkb1*, we examined transcriptional expression profiles of
double mutant organoid lines using bulk mRNA sequencing and compared them to their *Lkb1*-mutant
counterparts (Supplementary Table 3). With the introduction of mutant *Apc,* a slight increase was observed
in fetal and SSC gene programs in both *Lkb1*$^{+/-}$ and *Lkb1*$^{-/-}$ organoids, although the expression of stem cell
and ASC genes remained high for this mutational combination (Figure 7E, Figure S6E). By contrast,
introduction of mutant *Kras* in *Lkb1*$^{+/-}$ or *Lkb1*$^{bud\ -/-}$ backgrounds induced a strong shift towards expression of
fetal and SSC genes, while stem and ASC genes were downregulated, to a similar extent as observed for
*Lkb1*$^{cys\ -/-}$ organoids (Figure 7E, Figure S6E). In *Lkb1*$^{cys\ -/-}$ organoids, introduction of mutant *Kras* did not
further enhance the already strong serrated features, while introduction of truncated *Apc* in these organoids
clearly suppressed the serrated state and mediated reactivation of stem and ASC gene expression.
Quantification of these findings by GSEA confirmed a significant amplification of serrated signature by the
introduction of *Kras* mutations in both *Lkb1*$^{+/-}$ and *Lkb1*$^{bud\ -/-}$ organoids, and downregulation of the serrated
pathway by introduction of mutant Apc (Figure 7F). We conclude that *Lkb1* mutations synergize with mutant
*Kras,* but not *Apc,* to drive expression of the pre-malignant serrated gene program.

To investigate the link between *Lkb1* loss and advanced serrated colorectal tumors, we examined
the TCGA colorectal cancer database. First, we observed that *LKB1*-mutant tumors are significantly enriched
for mutations associated with serrated cancers, including *BRAF*, *RNF43* and *PIK3CA* mutations, while *APC*
mutations were significantly enriched in *LKB1* WT tumors (Figure 7G) (Day *et al.*, 2013; Robertson *et al.*,
2019). Mutations found in both classical and serrated adenocarcinomas, such as mutant *KRAS* and *TP53*,
were not significantly altered between *LKB1* WT and mutant tumors (Figure 7G). We next scored TCGA
colorectal samples for expression of a gene set that was upregulated in all *Lkb1*-deficient organoids.
Strikingly, expression of this *Lkb1*-deficient gene signature correlated with a mucinous histological CRC
subtype that is often found in serrated adenocarcinomas (Figure 7H) (Tuppurainen *et al.*, 2005). In addition,
among the consensus molecular subtypes (CMS), the serrated subtypes CMS1 and CMS4 carry significantly

higher expression of the *Lkb1*-deficient gene signature compared to the classical subtype CMS2 (Figure 7I)

(Nguyen, Goel and Chung, 2020). A recent study by Joanito *et al.* redefined the CMS classification and

found two epithelial intrinsic cell states in CRC samples (Joanito *et al.*, 2022). The iCMS2 state shows more

characteristics of classical CRCs, while the iCMS3 state correlates with features of serrated CRCs. Using

GSEA, we also found enrichment of iCMS3 transcriptional features in our *Lkb1*-deficient organoids, while the

iCMS2 state was downregulated (Figure S6F).

These results indicate that *Lkb1* loss drives cancer predisposition along the serrated pathway of

CRC development. *Kras* activating mutations cooperate with *Lkb1* loss by amplifying the serrated state.

Furthermore, upon homozygous *Lkb1* loss, a fraction of mutant cells grows out into *Lkb1*$^{cys\,-/-}$ organoids that

achieve an advanced serrated state without additional oncogenic mutations. Together, these findings

suggest that loss of *LKB1* converts intestinal epithelial cells in a pre-malignant state that primes for

progression along the serrated pathway.

**Discussion**

Over recent years, a variety of *Lkb1*-deficient mouse models have faithfully mimicked intestinal

hamartomatous polyp formation as a hallmark for PJS disease. The combined findings have promoted a

view in which non-epithelial functions of *LKB1* contribute to PJS disease development (Katajisto *et al.*, 2008;

Ollila *et al.*, 2018; Poffenberger *et al.*, 2018; Cotton *et al.*, 2022). Despite this progress, the relationship

between *LKB1* loss-induced hamartomatous polyposis and epithelial cancer predisposition has remained

highly debated and, moreover, does not explain cancer development in extra-intestinal epithelia (Shepherd,

Bussey and Jass, 1987; Entius *et al.*, 2001; De Leng *et al.*, 2003; Van Lier *et al.*, 2010; Korsse *et al.*, 2013).

Furthermore, how heterozygous and homozygous loss of *LKB1* mediates dose-dependent alterations in PJS

epithelial tissue organization at the single cell level has not been addressed, leaving the underlying cancer

driver mechanisms and pathways obscure.

Here, we employed mouse small intestinal organoids, an epithelium-only culture system, to obtain an

in-depth understanding of how *Lkb1* loss affects intestinal epithelial homeostatic cell identities and examine

how these alterations drive the epithelium into a pre-malignant state for colorectal cancer formation. By

generating both heterozygous and homozygous knockouts of *Lkb1*, we aimed to model aspects of cancer

predisposition and development in PJS patients, as over 64% of PJS-derived carcinomas present

inactivation of the second copy of the gene (Entius *et al.*, 2001; De Leng *et al.*, 2003). Our findings argue

that heterozygous *Lkb1* loss in the intestinal epithelium is sufficient to push cells into a premalignant program

for tumor development along the serrated colorectal cancer pathway, by a mechanism independent of the

micro-environment. Furthermore, expression of the serrated premalignant program is amplified upon loss of

the second allele. These results thus reveal that heterozygous *Lkb1*-mutant PJS epithelia already have

undergone alterations in phenotypic cellular identities in comparison to healthy individuals and supports the

view that loss of the second allele is a driver of cancer formation.

Our organoid-based model further confirms the previously described link between *Lkb1* loss and

aberrant secretory cell specification *in vivo* in mice, indicating that our epithelium-only model faithfully

reproduces the *in vivo* epithelium. We observed an increase in goblet cell numbers and mucus formation and

identified secretory cells that display 'intermediate' features, based both on morphology and transcriptome

analysis. In line with these findings, we show that expression of our newly identified *Lkb1*-deficient gene

signature correlates with a mucinous histological subtype of sporadic colorectal cancer cases. Together,

these findings indicate a common link between early premalignant alterations in secretory cell type

specification and colorectal cancer subtype progression. Similar *Lkb1* loss-induced alterations in secretory

cell identities may play a role in other epithelial tissues, as mucinous carcinomas in different organs were

frequently reported in PJS patients (Sato *et al.*, 2001; Osoegawa *et al.*, 2011; Collet *et al.*, 2020). If and how

aberrant secretory cell specification may facilitate cancer development however remains an issue for further

investigation.

Besides the observed alteration in secretory cell identities, transcriptomic analysis revealed that *Lkb1*

loss induces an allele dosage-dependent expression of a regenerative epithelial program. Notably, activation

of this tissue repair program occurred in regular culture conditions where no intestinal epithelial damage was

induced. We conclude that the *Lkb1*-deficient epithelium resides in a persistent regenerative state, a

condition that is worsened by loss-of-heterozygosity. A chronic state of regeneration may contribute to an

increase in tumor development as exemplified by the increased incidence of colorectal cancer formation in

patients suffering from inflammatory bowel disease (Lutgens *et al.*, 2013). As part of the induced

regenerative/YAP signaling-dependent program, the *Lkb1*-deficient epithelium showed an increase in

transcriptional expression of the *Egf* receptor family and its ligands, which allowed them to grow in the

absence of Egf supplementation, a condition that is not tolerated by healthy small intestinal organoids. These

findings thus reveal that the regenerative growth state induced by *Lkb1* loss endows the intestinal epithelium

with niche-independent properties, which is regarded as a hallmark of cancer (Hanahan and Weinberg,

2011). Furthermore, these results provide an explanation of the previously reported expanded proliferative

zone observed in the intestines of PJS patients (De Leng *et al.*, 2007)

Importantly, we uncovered that *Lkb1* loss predisposes the intestinal epithelium to express a gene

program related to the serrated pathway for colorectal cancer formation. In support of these observations,

*Lkb1* loss induced common features of early serrated lesions such as gastric metaplasia, induction of a

regenerative gene program and depletion of classical adult stem cells (Chen *et al.*, 2021). Furthermore,

inspection of a large set of sporadic colorectal cancer cases revealed that *LKB1* mutations frequently co-

occur with driver mutations linked to the serrated colorectal cancer pathway. Additionally, we observed

elevated expression of our *Lkb1*-deficient gene signature in the serrated colorectal cancer subtypes CMS1

and CMS4. Together, these findings thus suggest that the pathways employed for colorectal cancer

development in PJS patients are strikingly different from the well-studied cancer driver events in familial

adenomatous polyposis (FAP) patients that develop conventional *APC*-mutant adenocarcinomas (Li *et al.*,

2020).

Based on the work of Chen and colleagues (Chen *et al.*, 2021) a model was proposed in which

different environmental triggers, such as microbial dysbiosis and a subsequent MAPK pathway activation

drive the formation of sessile serrated lesions (IJspeert and Dekker, 2022; Shields *et al.*, 2022). In line with

this model, we observe that introduction of oncogenic mutant *Kras* in *Lkb1*-deficient organoids amplifies

serrated features and synergistically cooperates with *Lkb1* loss for organoid-forming capacity. Furthermore,

our results reveal that *Lkb1* loss-of-heterozygosity induces two distinct organoid phenotypes with either a

crypt-villus organization or a cystic morphology accompanied with strongly induced serrated features. Since

both *Lkb1*$^{-/-}$ organoid types carried similar *Lkb1* mutations (Figure S1C), we speculate that an external

stressor, such as the CRISPR procedure or clonal outgrowth, may promote this switch to a highly serrated

cystic phenotype. Although the responsible molecular triggers remain to be identified, our findings indicate

that *Lkb1*$^{-/-}$ cells are sensitized to undergo such a phenotypic switch and adopt a pre-malignant growth state.

Based on our findings, we conclude that the *Lkb1*-mutant epithelium is primed for the sessile serrated cancer

pathway and not the conventional adenoma-to-carcinoma sequence. In line with this reasoning, the

introduction of *Apc* mutations in *Lkb1*-deficient organoids did not display synergistic effects on pre-malignant

transcriptional features or organoid reconstitution capacity. Together, our findings suggest that the PJS

epithelium is sensitized to events that synergistically drive cells into a premalignant serrated state, such as

intestinal damage or activating mutations in MAPK pathway components.

PJS patients undergo extensive and life-long monitoring in the clinic, to resect obstructive polyps and

screen for (pre)cancerous lesions. Our results argue that monitoring of PJS patients will require specific

screening for the occurrence of serrated lesions that potentially may occur in non-polyp regions as well.

Serrated lesions are known to be frequently overlooked as their diagnosis relies on the existence of an

architecturally distorted serrated crypt that are difficult to identify with regular screening methods (Rosty *et*

*al.*, 2012). Diagnosed serrated lesions may be confirmed by various biomarkers including an increased

MUC5AC expression linked to the absence of CDX2 (Chen *et al.*, 2021), or the usage of additional markers

identified in this study such as ANXA1.

Taken together, our study reveals that intestinal organoids comprise a valuable model for *LKB1*

mutation-induced alterations in epithelial organization and growth in PJS patients. Our findings further

establish that induction of a premalignant sessile serrated program is a hallmark of PJS disease and

provides an explanation for increased cancer incidence in PJS patients. The presented models may provide

a starting point to uncover which and how additional mutations and environmental stressors synergistically

cooperate with *LKB1* mutations, and how interactions with stromal and immune components contribute to the

initiation of malignant growth of the PJS epithelium. Ultimately, in-depth knowledge on the pathways and

processes involved in driving PJS epithelia into a pre-malignant state may help to design strategies that re-

establish epithelial homeostasis and reduce the incidence of cancer development in PJS patients.

**References**

Ayyaz, A. *et al.* (2019) 'Single-cell transcriptomes of the regenerating intestine reveal a revival stem cell',

*Nature.* Springer US, 569(7754), pp. 121–125. doi: 10.1038/s41586-019-1154-y.

Baas, A. F. *et al.* (2004) 'Complete Polarization of Single Intestinal Epithelial Cells upon Activation of LKB1

by STRAD', *Cell.* Cell Press, 116(3), pp. 457–466. doi: 10.1016/S0092-8674(04)00114-X.

Balbinot, C. *et al.* (2018) 'The Cdx2 homeobox gene suppresses intestinal tumorigenesis through non-cell-

autonomous mechanisms', *Journal of Experimental Medicine*, 215(3), pp. 911–926. doi:

10.1084/jem.20170934.

Bansil, R. and Turner, B. S. (2018) 'The biology of mucus: Composition, synthesis and organization',

*Advanced Drug Delivery Reviews.* Elsevier, 124, pp. 3–15. doi: 10.1016/J.ADDR.2017.09.023.

Barker, N. *et al.* (2007) 'Identification of stem cells in small intestine and colon by marker gene Lgr5', *Nature*

*2007 449:7165.* Nature Publishing Group, 449(7165), pp. 1003–1007. doi: 10.1038/nature06196.

Beggs, A. D. *et al.* (2010) 'Peutz-Jeghers syndrome: a systematic review and recommendations for

management.', *Gut.* England, 59(7), pp. 975–986. doi: 10.1136/gut.2009.198499.

Butler, A. *et al.* (2018) 'Integrating single-cell transcriptomic data across different conditions, technologies,

and species', *Nature Biotechnology*, 36(5), pp. 411–420. doi: 10.1038/nbt.4096.

Candelli, T. *et al.* (2018) 'Sharq, A versatile preprocessing and QC pipeline for Single Cell RNA-seq',

*bioRxiv*, p. 250811. doi: 10.1101/250811.

Chen, B. *et al.* (2021) 'Differential pre-malignant programs and microenvironment chart distinct paths to

malignancy in human colorectal polyps', *Cell.* The Authors, 184(26), pp. 6262-6280.e26. doi:

10.1016/j.cell.2021.11.031.

Collet, L. *et al.* (2020) 'Kras and Lkb1 mutations synergistically induce intraductal papillary mucinous

neoplasm derived from pancreatic duct cells', *Gut.* BMJ Publishing Group, 69(4), pp. 704–714. doi:

10.1136/GUTJNL-2018-318059.

Cotton, J. L. *et al.* (2022) 'PTEN and LKB1 are differentially required in Gli1-expressing mesenchymal cells

to suppress gastrointestinal polyposis', *Cell Reports*, 40(3), p. 111125. doi:

10.1016/J.CELREP.2022.111125.

Day, F. L. *et al.* (2013) 'PIK3CA and PTEN gene and exon mutation-specific clinicopathologic and molecular

associations in Colorectal cancer', *Clinical Cancer Research.* American Association for Cancer Research,

19(12), pp. 3285–3296. doi: 10.1158/1078-0432.CCR-12-3614/85568/AM/PIK3CA-AND-PTEN-GENE-AND-

EXON-MUTATION-SPECIFIC.

Dow, L. E. *et al.* (2015) 'Apc Restoration Promotes Cellular Differentiation and Reestablishes Crypt

Homeostasis in Colorectal Cancer', *Cell.* Cell Press, 161(7), pp. 1539–1552. doi: 10.1016/j.cell.2015.05.033.

Entius, M. M. *et al.* (2001) 'Molecular genetic alterations in hamartomatous polyps and carcinomas of

patients with Peutz-Jeghers syndrome', *Journal of clinical pathology*, 54(2), pp. 126–131. doi:

10.1136/jcp.54.2.126.

Fearon, E. R. and Vogelstein, B. (1990) 'A genetic model for colorectal tumorigenesis.', *Cell*, 61(5), pp. 759–
67. doi: 10.1016/0092-8674(90)90186-i.

Fodde, R. (2002) 'The APC gene in colorectal cancer', *European Journal of Cancer*. Pergamon, 38(7), pp.
867–871. doi: 10.1016/S0959-8049(02)00040-0.

Fordham, R. P. *et al.* (2013) 'Transplantation of expanded fetal intestinal progenitors contributes to colon
regeneration after injury', *Cell Stem Cell*. The Authors, 13(6), pp. 734–744. doi: 10.1016/j.stem.2013.09.015.

Fujii, M. *et al.* (2015) 'Efficient genetic engineering of human intestinal organoids using electroporation',
*Nature Protocols 2015 10:10*. Nature Publishing Group, 10(10), pp. 1474–1485. doi:
10.1038/nprot.2015.088.

Fujii, M. *et al.* (2016) 'A Colorectal Tumor Organoid Library Demonstrates Progressive Loss of Niche Factor
Requirements during Tumorigenesis', *Cell Stem Cell*. Cell Press, 18(6), pp. 827–838. doi:
10.1016/J.STEM.2016.04.003.

Gao, Y. *et al.* (2020) 'LKB1 Represses ATOH1 via PDK4 and Energy Metabolism and Regulates Intestinal
Stem Cell Fate', *Gastroenterology*. W.B. Saunders, 158(5), p. 1401. doi: 10.1053/j.gastro.2019.12.033.

Giardiello, F. M. *et al.* (2000) 'Very high risk of cancer in familial Peutz-Jeghers syndrome.',
*Gastroenterology*. United States, 119(6), pp. 1447–1453. doi: 10.1053/gast.2000.20228.

Giardiello, F. M. and Trimbath, J. D. (2006) 'Peutz-Jeghers syndrome and management recommendations.',
*Clinical gastroenterology and hepatology : the official clinical practice journal of the American*
*Gastroenterological Association*. United States, 4(4), pp. 408–415. doi: 10.1016/j.cgh.2005.11.005.

Gregorieff, A. *et al.* (2015) 'Yap-dependent reprogramming of Lgr5+ stem cells drives intestinal regeneration
and cancer', *Nature*, 526(7575), pp. 715–718. doi: 10.1038/nature15382.

Haber, A. L. *et al.* (2017) 'A single-cell survey of the small intestinal epithelium', *Nature*. Nature Publishing
Group, 551(7680), pp. 333–339. doi: 10.1038/nature24489.

Hafemeister, C. and Satija, R. (2019) 'Normalization and variance stabilization of single-cell RNA-seq data
using regularized negative binomial regression', *Genome Biology*. Genome Biology, 20(1), pp. 1–15. doi:
10.1186/s13059-019-1874-1.

Hanahan, D. and Weinberg, R. A. (2011) 'Hallmarks of Cancer: The Next Generation', *Cell*. Cell Press,
144(5), pp. 646–674. doi: 10.1016/J.CELL.2011.02.013.

Hanemaaijer, E. S. *et al.* (2021) 'Single-cell atlas of developing murine adrenal gland reveals relation of
Schwann cell precursor signature to neuroblastoma phenotype', *Proceedings of the National Academy of*
*Sciences of the United States of America*, 118(5). doi: 10.1073/pnas.2022350118.

He, G.-W. *et al.* (2022) 'Optimized human intestinal organoid model reveals interleukin-22-dependency of
paneth cell formation', *Cell Stem Cell*, 29. doi: 10.1016/j.stem.2022.08.002.

Hearle, N. *et al.* (2006) 'Frequency and spectrum of cancers in the Peutz-Jeghers syndrome.', *Clinical cancer research : an official journal of the American Association for Cancer Research*. United States, 12(10), pp. 3209–3215. doi: 10.1158/1078-0432.CCR-06-0083.

Hemminki, A. *et al.* (1998) 'A serine/threonine kinase gene defective in Peutz–Jeghers syndrome', *Nature 1998 391:6663*. Nature Publishing Group, 391(6663), pp. 184–187. doi: 10.1038/34432.

IJspeert, J. E. G. and Dekker, E. (2022) 'Gastric metaplasia could initiate the serrated neoplasia pathway in CRC', *Nature Reviews Gastroenterology and Hepatology*, 19(4), pp. 217–218. doi: 10.1038/s41575-022-00592-z.

Jenne, D. E. *et al.* (1998) 'Peutz-Jeghers syndrome is caused by mutations in a novel serine threoninekinase', *Nature Genetics 1998 18:1*. Nature Publishing Group, 18(1), pp. 38–43. doi: 10.1038/ng0198-38.

Joanito, I. *et al.* (2022) 'Single-cell and bulk transcriptome sequencing identifies two epithelial tumor cell states and refines the consensus molecular classification of colorectal cancer', *Nature Genetics*, 54(7), pp. 963–975. doi: 10.1038/s41588-022-01100-4.

Katajisto, P. *et al.* (2008) 'LKB1 signaling in mesenchymal cells required for suppression of gastrointestinal polyposis', *Nature genetics*. Nat Genet, 40(4), pp. 455–459. doi: 10.1038/NG.98.

Kim, Y. *et al.* (2022) 'LKB1 in Intestinal Epithelial Cells Regulates Bile Acid Metabolism by Modulating FGF15/19 Production', *Cmgh*, 13(4), pp. 1121–1139. doi: 10.1016/j.jcmgh.2021.12.017.

Kline, E. R. *et al.* (2013) 'LKB1 Represses Focal Adhesion Kinase (FAK) Signaling via a FAK-LKB1 Complex to Regulate FAK Site Maturation and Directional Persistence', *The Journal of Biological Chemistry*. American Society for Biochemistry and Molecular Biology, 288(24), p. 17663. doi: 10.1074/JBC.M112.444620.

Korsse, S. E. *et al.* (2013) 'Identification of molecular alterations in gastrointestinal carcinomas and dysplastic hamartomas in Peutz-Jeghers syndrome', *Carcinogenesis*. Oxford Academic, 34(7), pp. 1611–1619. doi: 10.1093/CARCIN/BGT068.

De Leng, W. W. J. *et al.* (2003) 'Cyclooxygenase 2 expression and molecular alterations in Peutz-Jeghers hamartomas and carcinomas.', *Clinical cancer research : an official journal of the American Association for Cancer Research*. United States, 9(8), pp. 3065–72. Available at: http://www.ncbi.nlm.nih.gov/pubmed/12912958.

De Leng, W. W. J. *et al.* (2007) 'Peutz–Jeghers syndrome polyps are polyclonal with expanded progenitor cell compartment', *Gut*. BMJ Publishing Group, 56(10), pp. 1475–1476. doi: 10.1136/GUT.2007.128132.

Li, J. *et al.* (2020) 'Genomic and transcriptomic profiling of carcinogenesis in patients with familial adenomatous polyposis', *Gut*, 69, pp. 1283–1293. doi: 10.1136/gutjnl-2019-319438.

Van Lier, M. G. F. *et al.* (2010) 'High cancer risk in peutz-jeghers syndrome: A systematic review and surveillance recommendations', *American Journal of Gastroenterology*. Nature Publishing Group, 105(6), pp. 1258–1264. doi: 10.1038/ajg.2009.725.

Lutgens, M. W. M. D. *et al.* (2013) 'Declining Risk of Colorectal Cancer in Inflammatory Bowel DiseaseAn
Updated Meta-analysis of Population-based Cohort Studies', *Inflammatory Bowel Diseases*. Oxford
Academic, 19(4), pp. 789–799. doi: 10.1097/MIB.0B013E31828029C0.

Merenda, A., Fenderico, N. and Maurice, M. M. (2020) 'Wnt Signaling in 3D: Recent Advances in the
Applications of Intestinal Organoids Trends in Cell Biology', *Trends in Cell Biology*, 30(1). doi:
10.1016/j.tcb.2019.10.003.

Mohseni, M. *et al.* (2014) 'A genetic screen identifies an LKB1-MARK signalling axis controlling the Hippo-
YAP pathway', *Nature Cell Biology*. Nature Publishing Group, 16(1), pp. 108–117. doi: 10.1038/ncb2884.

Muraro, M. J. *et al.* (2016) 'A Single-Cell Transcriptome Atlas of the Human Pancreas', *Cell Systems*.
Elsevier, 3(4), p. 385. doi: 10.1016/J.CELS.2016.09.002.

Mustata, R. C. *et al.* (2013) 'Identification of Lgr5-Independent Spheroid-Generating Progenitors of the
Mouse Fetal Intestinal Epithelium', *Cell Reports*, 5(2), pp. 421–432. doi: 10.1016/j.celrep.2013.09.005.

Nguyen, L. H., Goel, A. and Chung, D. C. (2020) 'Pathways of Colorectal Carcinogenesis', *Gastroenterology*.
Elsevier, Inc, 158(2), pp. 291–302. doi: 10.1053/j.gastro.2019.08.059.

Nusse, Y. M. *et al.* (2018) 'Parasitic helminths induce fetal-like reversion in the intestinal stem cell niche',
*Nature*. Springer US, 559(7712), pp. 109–113. doi: 10.1038/s41586-018-0257-1.

Ollila, S. *et al.* (2018) 'Stromal Lkb1 deficiency leads to gastrointestinal tumorigenesis involving the IL-11-
JAK/STAT3 pathway', *The Journal of clinical investigation*. J Clin Invest, 128(1), pp. 402–414. doi:
10.1172/JCI93597.

Osoegawa, A. *et al.* (2011) 'LKB1 Mutations Frequently Detected in Mucinous Bronchioloalveolar
Carcinoma', *Japanese Journal of Clinical Oncology*. Oxford Academic, 41(9), pp. 1132–1137. doi:
10.1093/JJCO/HYR102.

Patai, Á. V. *et al.* (2013) 'Serrated pathway: Alternative route to colorectal cancer', *World Journal of
Gastroenterology*, 19(5), pp. 607–615. doi: 10.3748/wjg.v19.i5.607.

Pierce, S. E. *et al.* (2021) 'LKB1 inactivation modulates chromatin accessibility to drive metastatic
progression', *Nature Cell Biology*, 23(8), pp. 915–924. doi: 10.1038/s41556-021-00728-4.

Poffenberger, M. C. *et al.* (2018) 'LKB1 deficiency in T cells promotes the development of gastrointestinal
polyposis', *Science*. American Association for the Advancement of Science, 361(6400), pp. 406–411. doi:
10.1126/SCIENCE.AAN3975.

Qu, M. *et al.* (2021) 'Establishment of intestinal organoid cultures modeling injury-associated epithelial
regeneration', *Cell Research*. Springer US, 31(3), pp. 259–271. doi: 10.1038/s41422-020-00453-x.

Ran, F. A. *et al.* (2013) 'Genome engineering using the CRISPR-Cas9 system', *Nature Protocols*, 8(11), pp.
2281–2308. doi: 10.1038/nprot.2013.143.

Ren, F. *et al.* (2010) 'Hippo signaling regulates Drosophila intestine stem cell proliferation through multiple
pathways', *Proceedings of the National Academy of Sciences of the United States of America*. National

Academy of Sciences, 107(49), pp. 21064–21069. doi:
10.1073/PNAS.1012759107/SUPPL_FILE/PNAS.201012759SI.PDF.

Robertson, D. J. *et al.* (2019) 'Terminology, Molecular Features, Epidemiology, and Management of Serrated
Colorectal Neoplasia', *Gastroenterology*. Elsevier, 157(4), pp. 949-966.e4. doi:
10.1053/J.GASTRO.2019.06.041.

Rosty, C. *et al.* (2012) 'Serrated polyps of the large intestine: current understanding of diagnosis,
pathogenesis, and clinical management', *Journal of Gastroenterology 2012 48:3*. Springer, 48(3), pp. 287–
302. doi: 10.1007/S00535-012-0720-Y.

Sanchez-Cespedes, M. (2007) 'A role for LKB1 gene in human cancer beyond the Peutz-Jeghers
syndrome.', *Oncogene*. England, 26(57), pp. 7825–7832. doi: 10.1038/sj.onc.1210594.

Sato, N. *et al.* (2001) 'STK11/LKB1 Peutz-Jeghers Gene Inactivation in Intraductal Papillary-Mucinous
Neoplasms of the Pancreas', *The American Journal of Pathology*. Elsevier, 159(6), pp. 2017–2022. doi:
10.1016/S0002-9440(10)63053-2.

Sato, T. *et al.* (2009) 'Single Lgr5 stem cells build crypt-villus structures in vitro without a mesenchymal
niche', *Nature*. Macmillan Publishers Limited. All rights reserved, 459(7244), pp. 262–265. doi:
10.1038/nature07935.

Sato, T., Stange, D. E., *et al.* (2011) 'Long-term Expansion of Epithelial Organoids From Human Colon,
Adenoma, Adenocarcinoma, and Barrett's Epithelium', *Gastroenterology*. W.B. Saunders, 141(5), pp. 1762–
1772. doi: 10.1053/J.GASTRO.2011.07.050.

Sato, T., Es, J. H. Van, *et al.* (2011) 'Paneth cells constitute the niche for Lgr5 stem cells in intestinal crypts',
*Nature*, 469(7330), pp. 415–418. doi: 10.1038/nature09637.Paneth.

Satoh, Y. *et al.* (1990) 'Ultrastructure of Paneth cells in the intestine of various mammals', *Journal of Electron
Microscopy Technique*. John Wiley & Sons, Ltd, 16(1), pp. 69–80. doi: 10.1002/JEMT.1060160109.

Shackelford, D. B. and Shaw, R. J. (2009) 'The LKB1-AMPK pathway: Metabolism and growth control in
tumour suppression', *Nature Reviews Cancer*. Nature Publishing Group, 9(8), pp. 563–575. doi:
10.1038/nrc2676.

Shepherd, N. A., Bussey, H. J. and Jass, J. R. (1987) 'Epithelial misplacement in Peutz-Jeghers polyps. A
diagnostic pitfall.', *The American journal of surgical pathology*, 11(10), pp. 743–9. doi: 10.1097/00000478-
198710000-00001.

Shields, C. E. D. *et al.* (2022) 'Sensitive To Immune Checkpoint Therapy', 11(7), pp. 1792–1807. doi:
10.1158/2159-8290.CD-20-0770.Bacterial-driven.

Shorning, B. Y. *et al.* (2009) 'Lkb1 deficiency alters goblet and paneth cell differentiation in the small
intestine.', *PloS one*. Edited by R. E. Morty, 4(1), p. e4264. doi: 10.1371/journal.pone.0004264.

Tomasetti, C. and Vogelstein, B. (2015) 'Variation in cancer risk among tissues can be explained by the
number of stem cell divisions', *Science*. American Association for the Advancement of Science, 347(6217),

pp. 78–81. doi: 10.1126/SCIENCE.1260825.

Troughton, W. D. and Trier, J. S. (1969) 'Paneth and goblet cell renewal in mouse duodenal crypts.', *The*
*Journal of cell biology*, 41(1), pp. 251–268. doi: 10.1083/jcb.41.1.251.

Tuppurainen, K. *et al.* (2005) 'Morphology and microsatellite instability in sporadic serrated and non-serrated
colorectal cancer', *Journal of Pathology J Pathol*, 207, pp. 285–294. doi: 10.1002/path.1850.

Wang, Y. *et al.* (2019) 'Long-Term Culture Captures Injury-Repair Cycles of Colonic Stem Cells', *Cell*.
Elsevier Inc., 179(5), pp. 1144-1159.e15. doi: 10.1016/j.cell.2019.10.015.

Yang, N. *et al.* (2012) 'TAZ induces growth factor-independent proliferation through activation of EGFR
ligand amphiregulin', *Cell Cycle*. Taylor and Francis Inc., 11(15), pp. 2922–2930. doi: 10.4161/CC.21386.

Yui, S., Azzolin, L., Maimets, M., Pedersen, M. T., Fordham, R. P., Hansen, S. L., Larsen, H. L., Guiu, J.,
Alves, M. R. P., *et al.* (2018) 'YAP/TAZ-Dependent Reprogramming of Colonic Epithelium Links ECM
Remodeling to Tissue Regeneration', *Cell Stem Cell*. Cell Press, 22(1), pp. 35-49.e7. doi:
10.1016/j.stem.2017.11.001.

Zhang, J. *et al.* (2009) 'YAP-dependent induction of amphiregulin identifies a non-cell-autonomous
component of the Hippo pathway', *Nature Cell Biology*. doi: 10.1038/ncb1993.

**Figure legends**

**Figure 1. *Lkb1*-deficient organoids show heterogeneous morphology and viability.**
**A)** Representative brightfield images of mouse small intestine *Lkb1*-deficient organoid clones. Lower panel images show zoom-in images of boxed area in upper panel images. Scale = 100 µM **B)** Western blot analysis of Lkb1 protein levels in WT and *Lkb1*-deficient organoids. **C)** Organoid reconstitution assay from single cells performed at day 7. One-way ANOVA (Dunnett) was applied for analysis. * = *p < 0.05*. Each dot represents a biological replicate, *N* = 16. **D)** Representative images of an EdU incorporation (magenta) experiment marking the proliferative compartments in wild-type (WT) and *Lkb1*-deficient organoids. **E)** Quantification of EdU+ cells over the total number of cells (DAPI+ cells) of *Lkb1*-deficient organoids. For each condition, *N* = 1000 cells were analyzed derived from one independent experiment.

**Figure 2. *Lkb1* loss alters the number, location and granule composition of secretory cells. A)** Representative images of immunostaining for Lysozyme1 (Lyz1, green) to detect Lyz1+ cells in *Lkb1*-deficient clones. Cellular membranes are visualized using F-actin (white). Images are max projections of z-stack. Scale bar = 25 µm. White arrows indicate mislocalized Lyz1+ cells. **B)** Representative images of immunohistochemistry staining of Mucin 2 (Muc2) to visualize Muc2+ cells in *Lkb1*-deficient clones. Scale bar = 100 µm. **C-D)** Representative images of MUC2 **(C)** and LYZ1 **(D)** staining to visualize the localization of MUC2+ and LYZ1+ cells respectively in healthy intestinal epithelium and non-transformed PJS epithelium. Arrows indicate mislocalized LYZ1+ cells. Scalebar = 100 µm. **E)** Representative images of electron microscopy comparing the ultrastructure of Paneth cells and their secretory granules in wild-type organoids and *Lkb1*-deficient organoids. Lower panel images show zoom-in images of boxed area in upper panel images. Scale bar upper panels = 5 µm, scale bar lower panels = 1 µm. **F)** Representative images of electron microscopy comparing the ultrastructure of goblet cells and their secretory granules in wild-type organoids and *Lkb1*-deficient organoids. Lower panel images show zoom-in images of boxed area in upper panel images. Scale bar upper panels = 5 µm, scale bar lower panels = 1 µm.

**Figure 3. *Lkb1* loss upregulates a regenerative gene signature in the intestinal epithelium. A)** *Lkb1*-deficient versus wild-type (WT) organoid derived bulk mRNA-seq sequencing data was analyzed by gene set enrichment analysis (GSEA) for cell types (Haber *et al.*, 2017), colitis (Yui, Azzolin, Maimets, Pedersen, Fordham, Hansen, Larsen, Guiu, M. R. P. Alves, *et al.*, 2018), injury (Qu *et al.*, 2021), parasite (Nusse *et al.*, 2018), fetal spheroid (Mustata *et al.*, 2013), Yap regeneration (Gregorieff *et al.*, 2015) and revival stem cell (Ayyaz *et al.*, 2019) gene sets. The normalized enrichment score (NES) is depicted on the X-axis and the false discovery rates (FDRs) is indicated by the size of the dots. The color of the lines reflects whether a signature is enriched in WT (red) or *Lkb1*-deficient (blue) organoids. The shape of the dot (open/closed) indicates whether a signature is significantly enriched (FDR < 0.05). **B)** Heatmap of z-score transformed expression values of *Lgr5*+ stem cell genes (Stem) (Haber *et al.*, 2017) enriched in WT organoids (top cluster) and fetal genes (Fetal) (Mustata *et al.*, 2013) enriched in *Lkb1*-deficient organoids (bottom cluster). Key marker genes are highlighted. Each column represents a biological replicate of the indicated group, N = 3. **C)** Representative images of immunofluorescent staining to detect expression of the fetal marker Anxa1 (green) in *Lkb1*-deficient clones. Embryonic spheroids from E16.5 mice were used as positive control.

Cellular membranes are visualized by F-actin (red) and nuclei by DAPI (blue). **D)** Representative images of

immunohistochemistry staining of the fetal marker ANXA1 in healthy intestinal epithelium, PJS non-

transformed epithelium and PJS dysplastic tissue. High magnification of the area indicated by black arrows is

shown in the upper right corner. Scale bar = 100 μm

**Figure 4. *Lkb1*-deficient organoids show a Yap-induced regenerative state.**

**A)** Representative images of immunofluorescent staining of Yap (green) in wild-type (WT) and *Lkb1*-deficient

organoids. Nuclei are visualized by DAPI staining (blue). High magnification of the area indicated by white

box is shown in the lower three panels. Scale bar = 20 μm. **B)** Quantification of the nuclear fraction of Yap.

Dots represent individual organoids derived from two independent experiments. $N$ = 5 for WT organoids, $N$ =

7 for *Lkb1*$^{+/-}$ organoids, $N$ = 7 for *Lkb1*$^{bud\,-/-}$ organoids and $N$ = 7 for *Lkb1*$^{cys\,-/-}$. One-way ANOVA was applied

for analysis and statistical significance was defined as *\* = p < 0.05, \*\* = p < 0.01 and \*\*\*\* = p < 0.0001.*

**Figure 5. *Lkb1*-deficient organoids upregulate EGF family receptors and its ligands for niche**

**independency.**

**A-B)** Cumulative normalized counts of *Egfr* binding ligands **(A)** and *Egf* family receptors **(B)** in wild-type (WT)

and *Lkb1*-deficient organoids. One-way ANOVA was applied for analysis and statistical significance was

defined as *ns = not significant, \* = p < 0.05, \*\* = p < 0.01 and \*\*\*\* = p < 0.0001.* Each dot represents one

biological replicate, N = 3. **C)** Representative bright field images of WT and *Lkb1*-deficient organoids grown

in the presence (ENR) and absence (NR) of Egf. Organoids were incubated for 7 days in indicated medium

and split on day 4. Scale bar = 500 μm. **D)** Representative bright field images of WT and *Lkb1*-deficient

organoids cultured in NR and treated with DMSO or 100 nM Gefitinib. Organoids were incubated for 9 days

in indicated medium and split on day 6. Scale bar = 500 μm.

**Figure 6. *Lkb1*-deficiency redefines intestinal stem cell identity.**

**A-B)** UMAP visualization of scRNA-seq data from wild-type (WT (Adult)), fetal (WT(Fetal)) and *Lkb1*-deficient

organoids. Colors indicate the different organoids lines **(A)** and the different cell types **(B)**. **C)** Fraction of cell

types for each organoid line. Each color indicates a different cell type, related to figure B. **D)** Scatter plot

showing module scores for a Paneth gene signature (Haber *et al.*, 2017) on the y-axis and for a goblet cell

signature (Haber *et al.*, 2017) on the x-axis across all cells. Color/shape indicates the origin of the single cell.

**E)** Violin plot showing a module score for Egf family ligands within the goblet and enteroendocrine (EEC)

cluster. Colors reflect different organoid lines. **F)** Expression of the fetal marker *Anxa1* is overlaid onto the

UMAP plot for the different organoid lines. **G)** GSEA was performed on genes differentially expressed

between WT and *Lkb1*-deficient stem cells for cell types (Haber *et al.*, 2017), colitis (Yui, Azzolin, Maimets,

Pedersen, Fordham, Hansen, Larsen, Guiu, M. R. P. Alves, *et al.*, 2018), embryonic spheroid (Mustata *et al.*,

2013), Yap regeneration (Gregorieff *et al.*, 2015) and revival stem cell (Ayyaz *et al.*, 2019) gene sets. The

normalized enrichment score (NES) is depicted on the X-axis and the false discovery rates (FDRs) are

indicated by the size of the dots. The color of the lines reflects whether a signature is enriched in WT (red) or

*Lkb1*-deficient (blue) organoids. The shape of the dot (open/closed) indicates whether a signature is

significantly enriched (FDR < 0.05). **H)** Violin plot showing module scores for a goblet cell gene signature

(Haber *et al.*, 2017) across WT stem cells, *Lkb1*$^{+/-}$ stem cells, *Lkb1*$^{bud\,-/-}$ stem cells and all *Lkb1*$^{cys\,-/-}$ cells. **I)**

Violin plot showing expression of the regenerative marker *Anxa1* across WT stem cells, *Lkb1*^+/- stem cells,

*Lkb1*^bud -/- stem cells and all *Lkb1*^cys -/- cells.

**Figure 7. *Lkb1* loss drives a pre-malignant program along the serrated pathway which can be**

**amplified by additional *Kras* mutations. A)** GSEA was performed for a serrated-specific gene set (Chen *et*

*al.*, 2021). The normalized enrichment score (NES) is depicted on the X-axis and the false discovery rate

(FDR) is indicated by the size of the dots. The shape of the dot (open/closed) indicates whether a signature

is significantly enriched (FDR < 0.05). **B)** Heatmap of z-score transformed expression values of adenoma-

specific genes (ASC) (Chen *et al.*, 2021) enriched in WT organoids (top cluster) and serrated-specific genes

(SSC) (Chen *et al.*, 2021) enriched in *Lkb1*-deficient organoids (bottom cluster). Each column represents a

biological replicate of the indicated group, N = 3. **C)** Organoid reconstitution assay from single cells

performed at day 7 from trypsinization is shown for *Apc*-deficient and *Apc/Lkb1*-deficient organoids. Each dot

represents a biological replicate, *N* = 8 **D)** Cell reconstitution assay from single cells performed at day 7 from

trypsinization is shown for *Kras*-mutant and *Kras*-mutant/*Lkb1*-deficient organoids. Each dot represents a

biological replicate, *N* = 8 **E)** Heatmap of z-score transformed expression values of ASC genes enriched in

control organoids (top cluster) and SSC genes enriched in *Lkb1*-deficient organoids (bottom cluster) for *Lkb1*

single mutant (green) and *Apc/Lkb1* double mutant (blue) or *Kras*/*Lkb1* double mutant organoids (yellow).

Each column represents a biological replicate of the indicated group, N = 3. **F)** GSEA of a serrated-specific

gene set (Chen *et al.*, 2021) was performed on *Lkb1*-deficient and *Apc/Lkb1*-deficient or *Kras*-mutant/*Lkb1*-

deficient organoids. The normalized enrichment score (NES) is depicted on the X-axis and the FDR is

indicated by the size of the dots. The color of the lines reflects whether a signature is enriched in *Lkb1*-

deficient (red) or *Lkb1*-deficient/*Kras*-mutant (blue) organoids. **G)** Percentage of samples mutated for *BRAF,*

*RNF43, PIK3CA, APC, KRAS* or *TP53* in *LKB1* WT (*N* = 88) and *LKB1*-mutant (*N* = 7472) colorectal tumors

according to cBioPortal. The Benjamini-Hochberg procedure was applied for analysis and statistical

significance was determined as *\* = FDR < 0.05, \*\* = FDR < 0.01 and \*\*\*\* = FDR < 0.0001.* **H, I)** Box plots

showing log2 Z-scores for the *Lkb1*-deficient signature (genes significantly (*padj < 0.05*) upregulated in all

deficient clones compared to wild type organoids) over the colon adenocarcinoma TCGA bulk RNA

sequencing dataset grouped by histological classification as mucinous (*N* = 68) or non-mucinous (*N* = 437)

**(H)** or consensus molecular subtype (CMS1; *N* = 63, CMS2; *N* = 125, CMS3; *N* = 45, CMS 4; *N* = 73) **(I)**

Welch's unequal variances t-test was applied for analysis and statistical significance was determined as *\* = p*

*< 0.05, \*\* = p < 0.01 and \*\*\*\* = p < 0.0001.*

**Figure S1. *Lkb1* loss induced by different gRNAs results in similar heterogeneous phenotypes.**

**A)** Sequences of *Lkb1*-deficient organoids obtained with gRNA1 showing indels/deletions detected in exon 1

of the *Lkb1* gene. PAM sequences are underlined. **B)** Immunofluorescent images showing cleaved caspase

3 staining (green) marking apoptotic cells in *Lkb1*-deficient and wild type organoids. Cellular membranes are

visualized by F-actin (white). Scale bar = 33 μm. **C)** Sequences of *Lkb1*-deficient organoids obtained with

gRNA2 showing indels/deletions detected in exon 1 of the *Lkb1* gene. PAM sequences are underlined. **D)**

Representative brightfield images of *Lkb1*-deficient clones showing heterogeneous morphologies. **E)**

Western blot analysis of Lkb1 expression in each clone. **F)** Organoid reconstitution assay from single cells

for wild type and different morphologies of Lkb1$^{-/-}$ performed at day 7 from trypsinization. Friedman test was

applied for analysis. $* = p < 0.05$, $* = p < 0.01$. Each dot represents a biological replicate, $N = 5$.

**Figure S2. Goblet cell number and positioning are unaltered in PJS epithelia and *Lkb1*-deficient**

**organoids display proper brush border formation. A-B)** The average number **(A)** and position **(B)** of

goblet cells healthy and PJS epithelia. Each dot represents a single crypt. For healthy donors 40 crypts from

4 individuals were counted. For PJS untransformed epithelia 30 crypts from 3 individuals were counted.

Unpaired t-test was applied for analysis. *ns = not significant*. **C)** Representative electron microscopy images

of wild type (WT) and *Lkb1*-deficient organoids. Arrows indicate brush border formation in indicated organoid

lines. A zoom-in of the area indicated by the red arrows is shown in the bottom panel. Scale bar = 2 μm.

**Figure S3. Yap target gene expression progressively increases with loss of both *Lkb1* alleles.**

**A)** GSEA was performed for *Lkb1*$^{cys -/-}$ organoids versus WENR treated wild-type (WT (WENR)) organoids or

*Apc*$^{-/-}$ organoids. The following gene sets were tested: cell types (Haber *et al.*, 2017), colitis (Yui, Azzolin,

Maimets, Pedersen, Fordham, Hansen, Larsen, Guiu, M. R. P. Alves, *et al.*, 2018), injury (Qu *et al.*, 2021),

parasite (Nusse *et al.*, 2018), fetal spheroid (Mustata *et al.*, 2013), Yap regeneration (Gregorieff *et al.*, 2015)

and revival stem cell (Ayyaz *et al.*, 2019). The normalized enrichment score (NES) is depicted on the X-axis

and the false discovery rates (FDRs) is indicated by the size of the dots. The color of the lines reflects

whether a signature is enriched in WT (WENR) (blue), *Lkb1*$^{cys -/-}$ (green), or *Apc*$^{-/-}$ (red) organoids. The shape

of the dot (open/closed) indicates whether a signature is significantly enriched (FDR < 0.05). **B)** Heatmap of

z-score transformed expression values of Yap responsive genes (Gregorieff *et al.*, 2015) enriched in *Lkb1*-

deficient organoids. Key marker genes are highlighted. Each column represents a biological replicate of the

indicate group, N = 3.

**Figure S4. Identification of intestinal cell types of WT and *Lkb1*-deficient organoids.**

**A)** UMAP visualization of scRNA-seq data from wild-type (WT), fetal and *Lkb1*-deficient organoids. Colors

indicate different clusters as defined by Seurat analysis **B)** Expression of known markers for different cell

populations in WT and *Lkb1*-deficient organoids. Size of the dot represents percentage of cells expressing

marker gene and the color the average expression within that population. **C-I)** Module scores for enterocyte

**(C)**, enteroendocrine **(D)**, goblet **(E)**, stem **(F)**, transit amplifying (G1) **(G)**, Paneth **(H)** and tuft **(I)** cell gene

signatures (Haber *et al.*, 2017) overlaid on the UMAP plot of *Lkb1*-deficient, fetal and WT organoids. **J-K)**

Reclustering of stem cell populations from WT, *Lkb1*$^{+/-}$, *Lkb1*$^{-/- bud}$ and *Lkb1*$^{-/- cys}$ organoids showing clusters

as identified by Seurat analysis **(J)** and by organoid line **(K)**.

**Figure S5. Induction of serrated genes in *Lkb1*-deficient organoids is reflected at the single cell level.**

**A)** Module score for a serrated-specific (SSC) gene set (Chen *et al.*, 2021) on the UMAP plot of *Lkb1*-

deficient, fetal and WT organoids. **B)** Violin plot showing a module score for a serrated-specific gene set

across WT, fetal, *Lkb1*$^{+/-}$, *Lkb1*$^{bud -/-}$ and *Lkb1*$^{cys -/-}$ organoids. **C)** GSEA was performed for a serrated-specific

gene set (Chen *et al.*, 2021). The normalized enrichment score (NES) is depicted on the Y-axis and the false

discovery rate (FDR) is indicated by the size of the dots. Line colors reflects enrichment of the signature in

WT (blue) or *Lkb1*-deficient (red) stem cells. Dot shape (open/closed) indicates significant enrichment (FDR

< 0.05). **D)** Module score for a serrated-specific gene set (Chen *et al.*, 2021) on the UMAP plot of WT stem

cells, *Lkb1*$^{+/-}$ stem cells, *Lkb1*$^{bud-/-}$ stem cells and all *Lkb1*$^{cys-/-}$ cells. **E)** Violin plot showing module scores of a

serrated-specific gene set (Chen *et al.*, 2021) for WT stem cells, *Lkb1*$^{+/-}$ stem cells, *Lkb1*$^{bud-/-}$ stem cells and

all *Lkb1*$^{cys-/-}$ cells.

**Figure S6. *Lkb1* loss drives expression of a pre-malignant program along the serrated pathway which**

**can be amplified by additional *Kras* mutations. A)** Representative brightfield images of *Apc* and

*Apc/Lkb1*-deficient organoids. **B)** Representative brightfield images of *Kras*-mutant and *Kras*-mutant/*Lkb1*-

deficient organoids. **C)** Sequences of *Apc*$^{-/-}$ and *Lkb1/Apc* mutant organoids showing indels/deletions

detected in exon 15 of the *Apc* gene. PAM sequences are underlined. **D)** Sequences of *Kras*$^{G12D}$ and

*Lkb1/Kras* mutant organoids showing G12D mutations/deletions detected in exon 2 of the *Kras* gene. PAM

sequences are underlined. **E)** Heatmap of z-score transformed expression values of *Lgr5*$^{+}$ stem cell genes

(Stem) (Haber *et al.*, 2017) enriched in control organoids (top cluster) and fetal genes (Fetal) (Mustata *et al.*,

2013) enriched in *Lkb1*-deficient organoids (bottom cluster) for *Lkb1* single mutant (green) and *Apc/Lkb1*

double mutant (blue) or *Kras/Lkb1* double mutant organoids (yellow). Each column represents a biological

replicate of the indicate group, N = 3. **F)** GSEA of iCMS2 and iCMS3 gene signatures (Joanito *et al.*, 2022)

was performed on *Lkb1*-deficient organoids compared to control organoids. The normalized enrichment

score (NES) is depicted on the X-axis and the FDR is indicated by the size of the dots. Line color reflects

signature enrichment in *Lkb1*-deficient (red) or *Lkb1*-deficient/*Kras*-mutant (blue) organoids. Dot shape

(open/closed) indicates significant enrichment (FDR < 0.05).

## Materials and Methods

### Organoid culture

Small intestinal mouse organoids were established from crypts isolated from the small intestine of wild-type mice, as previously described (Sato *et al.*, 2009). Basal culture medium for mouse organoids contained advanced DMEM/F12 (Gibco) supplemented with 1% Penicillin-Streptomycin (Sigma-Aldrich), 10 mM HEPES (Gibco), 1% Ultraglutamine 1 (Lonza), 1% B27 (Sigma-Aldrich), 0.5% N2 (Sigma-Aldrich) and 1 mM N-acetylcysteine (Sigma- Aldrich). Basal culture medium was further supplemented with murine recombinant EGF (Peprotech), 2.5% R-spondin1 conditioned medium (homemade) and 10% Noggin conditioned medium (homemade) to obtain the complete ENR (EGF-Noggin-Rspondin) medium. Organoids were grown in Matrigel (Corning) or BME (Cultrex) and passaged every 6-8 days as previously described (Sato *et al.*, 2009); their medium was refreshed every 2-3 days. For growth factor withdrawal experiments, EGF was removed from organoid culture medium and organoids were monitored for a total of 16 days. Bright field images were acquired with the Invitrogen™ EVOS™ FL Color Imaging System (Thermo Fisher Scientific). To obtain wild type cystic organoids, wild type organoids were supplemented with 50% Wnt3a-conditioned medium (homemade) and 1mM Nicotinamide (Sigma-Aldrich)

### Vectors and sgRNAs

The pSpCas9(BB)-2A-Puro plasmid was obtained from Addgene (48139). Single guide RNAs (sgRNAs) and are listed in **Table S4**. Cloning of sgRNAs into the The pSpCas9(BB)-2A-Puro plasmid was performed according to previously published protocols (Ran *et al.*, 2013) and vector sequences conformed by Sanger sequencing.

### Organoid electroporation

For electroporation, mouse organoids were expanded in ENR medium without antibiotics and supplemented with 50% Wnt3a-conditioned medium (homemade) and 1mM Nicotinamide (Sigma-Aldrich). Electroporation of organoids was performed as previously described using a NEPA21 (NEPAGENE) electroporator (Fujii *et al.*, 2015). To obtain *Lkb1*-deficient organoids, a total of 20 µg of DNA (10 µg gRNA, 7.2 µg of pPB-CAG-rtTA_Hygromycin and 2.8 µg PiggyBac transposase) was used. To obtain *Apc* and *Kras* mutant organoids a total of 10 µg DNA (gRNA and gRNA with oligonucleotide respectively) was used. 100 µg/mL Hygromycin B was used to select *Lkb1*-deficient organoids and Noggin and Egf were omitted from the culture medium to select for *Apc* and *Kras* mutant organoids respectively. After selection, single organoids were manually picked for clonal expansion and then genotyped. Genomic DNA was isolated using the DNA micro kit (Qiagen) according to the manufacturer's instructions and PCR amplification was performed using the GoTaq Flexi DNA polymerase kit (Promega). Organoid genotype was confirmed by Sanger sequencing. Primers are listed in **Table S5.**

### Protein extraction and Western blotting

Organoids were harvested three days after splitting and washed in cold PBS and collected in cell recovery solution (Corning) according to the manufacturer's instructions. Whole cell lysates were obtained by direct lysis in RIPA buffer (50 mM Tris- HCl pH 8.0, 150 mM NaCl, 0.1% SDS, 0.5% Na-Deoxycholate, 1% NP-40)

containing complete protease inhibitors (Sigma-Aldrich) for 30 minutes on ice. Proteins were run on SDS–

PAGE gels and transferred to PVDF membranes, followed by blocking with Odyssey blocking buffer (LI-

COR). The antibodies used are described in **Table S6**. Blots were imaged with the Amersham Typhoon (GE

Healthcare Life Sciences).

**Immunofluorescence**

Organoids were seeded in 15-well Ibidi chambers after passaging and fixed 3-4 days later in 4%

Paraformaldehyde (Sigma-Aldrich) for 60 minutes at room temperature. After fixation organoids

were washed with cold PBS and permeabilized with PBD0.2T (48,5 mL PBS, 1 mL 10%Triton, 0,5mL DMSO

and 0,5 g BSA) for 20 minutes at room temperature, after which they were incubated with primary antibodies

diluted in PBD0.2T buffer at 4°C overnight. Washes were performed PBD0.2T buffer and secondary

antibodies incubated 2-3 hours at room temperature. All antibodies are described in **Table S7**. Quantification

of the nuclear fraction of Yap was analyzed using the JACoP Fiji plugin. EdU incorporation assays were

performed by culturing organoids in ENR medium containing 10 µM Edu for 6 hours to cells, after which

single organoids were harvested, fixed in 4% PFA and EdU-labeling was performed according to

manufacturer instructions (Click-iT EdU Cell Proliferation Kit for Imaging, Thermo Fisher). Images were

acquired with a Zeiss LSM700 or LSM880 confocal microscope.

**Patient tissue samples and counting**

Patient material was collected from the pathology archives of Dutch hospitals (UMC). In total five PJS

patients were included in this study. Five normal controls were randomly selected. In 10 microscopically

normal looking crypts MUC2$^+$ cells were counted from the bottom toward to the top of the crypt. Only

longitudinally oriented crypts that were completely visible on the slide, were counted. Cell positions were

defined as the bottom of the crypt is cell position one (1), going upward to cell position +2, +3, and so forth till

cell position 30. Within the part of the crypt between cell position 1 and 30, the number of positively labelled

cells were counted. Also, the total cell position was determined by adding up each cell position of a labelled

cell and average cell position of the labelled cells was determined by adding up each cell position of a

labelled cell and dividing the sum by the total number of labelled cells counted.

**Immunohistochemistry**

Fresh mouse small intestinal organoids were fixed in 4% Formaldehyde three days after splitting for 1 hour

followed by dehydration in 25%, 50% and 70% ethanol and embedding in paraffin. Briefly, 4µm FFPE

organoid or PJS patient tissue sections were obtained, deparaffinized, blocked in 0.3% H2O2 in methanol for

20 minutes. Antigen retrieval was performed in Tris/EDTA buffer (10 mM/1 mM; pH 9.0) or Sodium-citrate

buffer (0,01 M/pH6.0) for 20 minutes at 100°C. Nonspecific binding sites were blocked using Protein Block

Serum free (DAKO) for 10 minutes, followed by primary antibody incubation. Antibody binding was visualized

using the using BrightVision Goat anti Rabbit / Poly-AP and BrightVision Goat anti Mouse / Poly-HRP

(VWR/ImmunoLogic), with DAB as chromogen (Sigma-Aldrich). Sections were counterstained with

Haematoxylin or neutral red and mounted using Pertex. Primary antibodies are described in **Table S8**.

**Electron microscopy**

Three days after splitting, organoids were fixed in a solution of 2% paraformaldehyde and 2.5%

glutaraldehyde in 0.1 M phosphate buffer (pH 7.4) overnight at 4°C. On the following day, organoids were

washed twice in 0.1 M phosphate buffer (pH 7.4) and subsequently post-fixed with 1% Osmium tetroxide for

2 hours at 4°C. Organoids were then stained with 0.5% uranyl acetate 1 hour at 4°C in the dark. Matrigel

droplets were transferred into glass vials for dehydration with increasing acetone concentrations according to

the following protocol: 70% overnight at 4°C, 90% for 15 minutes at RT, 96% for 15 minutes at RT, and

100% 3 times for 30 minutes each at RT. After dehydration, samples were embedded in EPON resin.

Ultrathin sections of 65 nm were obtained with a Leica UC7 microtome and stained with uranyl and lead

citrate. Electron microscopy was performed in the Cell Microscopy Core (CMC) of the UMC Utrecht. Sections

were imaged with a JEOL JEM-1010 transmission electron microscope (TEM).

**Cell viability assay**

For the organoid reconstitution assays, organoids were dissociated to single cells after incubation with

TrypLE (Life Technologies) for 5 min at 37°C. Dissociated organoids were passed through a 40 μm strainer

and counted. Cells were seeded at a density of 5000 cells/10μl Matrigel and cultured in ENR medium with 10

μM Y-27632 (Sigma-Aldrich) for 7 days. Organoids were then lysed with CellTiter-Glo® (Promega) according

to the manufacturer's protocol. ATP produced by viable cells was measured by a Berthold luminometer

Centro LB960.

**Bulk RNA sequencing and data analysis**

Organoids were lysed in RLT lysis buffer (Qiagen) 3-5 days after passaging. Biological triplicates were

employed. RNA was obtained using the Qiagen QiaSymphony SP (Qiagen) according to the manufacturer's

protocol. Single-end RNASeq reads were processed using the UMCU RNASeq pipeline

([https://github.com/UMCUGenetics/RNASeq](https://github.com/UMCUGenetics/RNASeq), v2.3.0) with default settings. Read quality was assessed by

FastQC (0.11.4) followed by splice-aware alignment against the mouse reference genome (GRCm38) with

STAR (2.4.2a). RNA expression quantification was performed with htseq-count (0.6.0) in reverse-stranded

mode. Differential gene expression analysis was performed with the DESeq2 (1.22.2) R package. In each

comparison, genes were selected with an absolute fold-change > 2 and alpha P > 0.05. Data clustering and

generation of heatmaps was performed with pheatmap. Gene Set Enrichment Analysis was performed using

GSEA (v4.2.0, Windows App) and visualized using ggplot2. Intestinal cell-type specific gene sets for GSEA

were obtained from Extended Figure Table 3 of Haber et al., 2017 (Haber *et al.*, 2017). The other datasets

employed in GSEA were derived from previously published studies (Mustata *et al.*, 2013; Gregorieff *et al.*,

2015; Nusse *et al.*, 2018; Yui, Azzolin, Maimets, Pedersen, Fordham, Hansen, Larsen, Guiu, M. R. P. P.

Alves, *et al.*, 2018; Ayyaz *et al.*, 2019; Chen *et al.*, 2021; Qu *et al.*, 2021; Joanito *et al.*, 2022). Human gene

sets were converted to their respective mouse homologs using biomaRt (2.50.3) in R where necessary.

**Single cell RNA sequencing**

*Sample preparation and sequencing:* Single cells from organoid were harvested using TrypLE incubation at

37 °C, four days after passaging. Cells were filtered through a 40 μm nylon cell strainer (Falcon). scRNA-seq

was performed according to the Sort-seq protocol (Muraro *et al.*, 2016). Libraries were sequenced on an

Illumina NextSeq500 at paired-end 60- and 26-bp read length and 75,000 reads per cell.

*Mapping and filtering*: The Sharq pipeline was used to process the sequencing data (Candelli *et al.*, 2018).

Mapping was performed using STAR (version 2.6.1), on the Genome Reference Consortium Mouse built 38

(GRCm38). Read assignment was with feature Counts version 1.5.2 using a gene annotation based on

GENCODE version M14.

External RNA controls (ERCCS) and transcripts mapping to the mitochondrial genome were removed from

all cells. Low quality cell barcodes having either less than 800 unique transcripts, 200 genes 33.3%

mitochondrial transcripts or over 4-fold non-exonic to exonic reads were excluded, as well as barcodes with

over 100,000 unique transcripts.

*scRNA-seq analysis*: Unique transcript counts were normalized using sctransform (Hafemeister and Satija,

2019) and analyzed using the Seurat R package (version 4.1.0) (Butler *et al.*, 2018). From the top 3000

variable genes, genes associated with cell cycle phase, dissociation stress (heat shock and chaperone

proteins according to GO:0006986), sex (Xist, Tsix, and Y chromosome-specific genes), and activity

(ribosomal protein genes according to GO:0022626) were removed from the list of variable genes to avoid

biases in cell clustering, as described before (Hanemaaijer *et al.*, 2021). The first 20 principal components

were used both to calculate dimensionality reduction using UMAP and to perform clustering with a resolution

of 1, k.param = 10 and the Louvain algorithm.

**TCGA data analysis**

*STK11/LKB1* co-occurring mutations were determined using cBioPortal (https://www.cbioportal.org/). All

bowel datasets were selected and using the comparison tool co-occuring genomic alterations were identified

in *STK11/LKB1* WT and *STK11/LKB1* mutant samples. Signature scores for a *Lkb1*-deficient geneset were

calculated for each sample as log2 Z-score over the samples from the TCGA Colon Adenocarcinoma (2022

– v32) set using the R2 genomic analysis platform (http://r2.amc.nl). The *Lkb1*-deficient signature was

determined as genes that were significantly (padj < 0.05) upregulated in all comparisons (WT vs. *Lkb1*[+/-], WT

vs. *Lkb1*[bud -/-], WT vs. *Lkb1*[cys -/-]).

**Supplementary information**

**Table S4: gRNA sequences**

| Gene | Sequence |
|---|---|
| *Lkb1* gRNA1 | AGCTTGGCGCGTTTGCGGCG |
| *Lkb1* gRNA2 | TGAGGATCTTGACCGCCCTG |
| Apc | TGTCTGGCTCCGGTAAGTGA |
| Kras | CTTGTGGTGGTTGGAGCTGG |

957 **Table S5: Genotyping primers**

| Gene | Forward primer | Reverse Primer |
|---|---|---|
| *Lkb1* gRNA1 | GGGGAAAATCAAAAGTGAAGAA | CCCTCTAGCTGCGTAAACAAAC |
| *Lkb1* gRNA2 | CTCCACCGAGGTAATCTACCAGCCG | TCTAACCGTTTTCTCTTCTCATTTCC |
| *Apc* | ATAGAAACAGCACTGACCCAAATTTCA | AGGCCTCTTTGCTTAGAGCTTTCATAA |
| *Kras* | TGGCTCCAACACAGATGTTC | GGATGGCATCTTGGACCTTA |

959 **Table S6: Western Blot antibodies**

| Antibody | Dilution | Supplier | Cat. number |
|---|---|---|---|
| Anti-LKB1 | 1:1000 | Santa Cruz | sc-32245 |
| Anti-GAPDH | 1:8000 | Millipore | CB1001 |
| Anti-mouse Alexa 800 | 1:5000 | LI-COR | 926-32211 |
| Anti-rabbit Alexa 680 | 1:5000 | LI-COR | 925-68070 |

961 **Table S7: Immunostaining antibodies**

| Antibody | Dilution | Supplier | Cat. number |
|---|---|---|---|
| Anti-Muc2 | 1:200 (IF) 1:1000 (IHC) | Santa Cruz | sc-515032 |
| Anti-Lysozyme1 | 1:200 (IF) 1:2000 (IHC) | Dako | A0099 |
| Anti-ANXA1 | 1:200 (IF) 1:1000 (IHC) | Atlas Antibodies | HPA011271 |
| Anti-YAP | 1:200 (IF) | Santa Cruz | sc-101199 |
| Anti-Cleasved-Caspase-3 | 1:200 (IF) | Cell Signalling | 9661 |
| Anti-rabbit Alexa Fluor 488 | 1:300 (IF) | Invitrogen | A11034 |
| Anti-mouse Alexa 488 | 1:300 (IF) | Invitrogen | A11029 |
| DAPI | 1:1000 (IF) | Sigma | D9542 |
| Click-iT™ EdU Alexa Fluor™ 647 | n.a. | Invitrogen | C10424 |

| Alexa Fluor™ 647 Phalloidin | 1:500 (IF) | Invitrogen | A22287 |
|---|---|---|---|

## Data availability

964 The scRNA-seq and bulk RNA-seq data generated in this study will be uploaded to Gene Expression
965 Omnibus (GEO) upon publication. If needed earlier, data can be requested via the corresponding author.
966

## Author contributions

968 S.F.P, H.M., A.M. and M.M.M., conceived and designed the experiments. S.F.P, H.M., A.M., D.X., F.H.M.M.,
969 C.J. and A.R.K performed the experiments. S.F.P, H.M., A.M., J.Y.V., J.S., T.M., T.C., R.S., J.R., F.C.P.H.,
970 G.J.A.O. and M.M.M. analyzed the data. W.W.J.L. and G.J.A.O. provided patient material. S.F.P, H.M., A.M.,
971 J.Y.V. and M.M.M. wrote the manuscript, which was reviewed by all authors.
972

## Disclosure and competing interest statement

974 M.M.M. is co-founder and shareholder of Laigo Bio and inventor on patents related to membrane protein
975 degradation; J.R. is co-founder of Cyclomics BV., a genomics company.
976

## Acknowledgements

978 The authors thank members of the Maurice laboratory for helpful discussions and suggestions. PiggyBac
979 Transposase and pPB-CAG-rtTA_Hygromycin were a kind gift from B.K. Koo. This work is part of the
980 Oncode Institute, which is partly financed by the Dutch Cancer Society. This work was supported by the
981 China Scholarship Council (to H.M.), the Dutch Cancer Society KWF Grant 13112 (to M.M.M.), Zon-MW
982 TOP grant 91218050 (to M.M.M.) and NWO Gravitation project IMAGINE! (to M.M.M.).

**Figure 1**

**Figure 2**

# Figure 3

**A**

| Lkb1+/− vs. WT | Lkb1bud−/− vs. WT | Lkb1cys−/− vs. WT |

Stem, Paneth, Goblet, Colitis, Injury, Parasite, Fetal spheroid, Yap regeneration, Revival stem cell

NES

Enriched in: WT, Lkb1-deficient

FDR: 0.00, 0.02, 0.04, 0.06, 0.08

Significant: No, Yes

**B**

WT, Lkb1+/−, Lkb1bud −/−, Lkb1cys −/−

Stem, Fetal

Ascl2, Znf3, Axin2, Rnf32, Lrp4, Ly6d, Anxa1, Clu, Anxa3, Anxa8, Msln

**C**

Anxa1 | Merge

WT Adult, WT Fetal, Lkb1+/−, Lkb1bud −/−, Lkb1cys −/−

Anxa1, DAPI, Actin

**D**

Healthy donor | PJS non-transformed | PJS dysplasia

ANXA1

# Figure 4

**Figure 5**

# Figure 6

# Figure 7

# Figure S1

**Figure S2**

**Figure S3**

**A**

**B**

**Figure S4**

**Figure S5**

# Figure S6