1 **Possibilities of skin coat color-dependent risks and risk factors of squamous cell**
2 **carcinoma and deafness of domestic cats inferred via RNA-seq data**
3

4 Akinori Awazu[1,2*], Daigo Takemoto[1], Kaichi Watanabe[1], Naoaki Sakamoto[1,2]
5
6 1 Graduate School of Integrated Sciences for Life, Hiroshima University,
7 Higashihiroshima, Hiroshima 739-8526, Japan
8 2 Research Center for the Mathematics on Chromatin Live Dynamics, Hiroshima
9 University, Higashihiroshima, Hiroshima 739-8526, Japan
10
11 *Correspondence:
12 Akinori Awazu: awa@hiroshima-u.ac.jp

14 **Keywords:** coat color, deafness, domestic cats, *Felis catus,* insulator, squamous cell
15 carcinoma, transcriptome, white locus

Abstract

The transcriptome data of skin cells from domestic cats with brown, orange, and white coats were analyzed using a public database to investigate the possible relationship between coat color-related gene expression and squamous cell carcinoma risk, as well as the mechanism of deafness in white cats. We found that the ratio of the expression level of genes suppressing squamous cell carcinoma to that of genes promoting squamous cell carcinoma might be considerably lower than the theoretical estimation in skin cells with orange and white coats in white-spotted cat. We also found the possibility of the frequent production of KIT lacking the first exon (d1KIT) in skin cells with white coats, and d1KIT production exhibited a substantial negative correlation with the expression of SOX10, which is essential for melanocyte formation and adjustment of hearing function. Additionally, the production of d1KIT was expected to be due to the insulating activity of the feline endogenous retrovirus 1 (FERV1) LTR in the 1st intron of *KIT* and its CTCF binding sequence repeat. These results contribute to basic veterinary research to understand the relationship between cat skin coat and disease risk, as well as the underlying mechanism.

Introduction

Domestic cats are among the most important companion animals, and feline health occupies an important position in human society. It has also recently been suggested that dementia and breast cancer in cats share various similarities with Alzheimer's disease (Chambers et al., 2015; Sordo et al., 2021) and triple-negative breast cancer in humans (Wiese et al., 2013; Caliari et al., 2014; Cannon, 2015; Thomas, 2015; Adega et al., 2016; Howard et al., 2020; Sommerville et al., 2022), respectively. Therefore, advances in feline medicine are expected to make important contributions to advances in human medicine.

In recent years, research on various feline diseases has progressed, leading to the development of vaccines for various lethal infectious diseases caused by virus such as Feline leukemia virus (FeLV) (Hofmann-Lehmann et al., 2020) and Feline immunodeficiency virus (FIV) (Westman et al., 2019; Andrade et al., 2022), therapeutic drugs for Feline infectious peritonitis (FIP) (Pedersen et al., 2019; Cook et al., 2022), and drugs for kidney disease (Sugisawa et al., 2016; Arai et al., 2016) that have been put into practical use, consequently extending the lifespan of domestic cats. However, these methods have only solved some diseases in cats. Basic medical research focusing

on the genetics and physiology of cats is essential for establishing preventive and
therapeutic methods for various feline diseases.
Domestic cats have different colors and patterns on their skin and coat, which
vary from individual to individual depending on the combination of genes related to
color (white, orange, brown, black, etc.) and patterns (agouti, tabby, dilute, etc.) (Lyons
et al., 2005; Schmidt-Küntzel et al., 2005; Imes et al., 2006; Ishida et al., 2006;
Schmidt-Küntzel et al., 2009; Eizirik et al., 2010; David et al., 2014; Lyons, 2021), and
the pattern of X-chromosome inactivation in female. In addition, certain skin colors in
cats have been suggested to be associated with specific diseases. For example,
light-skinned individuals have been suggested to have a higher risk of squamous cell
carcinoma than dark-skinned individuals (Dorn et al., 1971; Thomson, 2007; Murphy,
2013). This tendency has been attributed to the fact that light-colored individuals with
less melanin have a weak resistance to ultraviolet (UV) rays. However, squamous cell
carcinoma also develops in the lining of the oral cavity, tongue, and tonsils, which are
less likely to be directly exposed to UV light. Therefore, there may be other marked
factors besides UV light (Baer et al., 1993; Bertone et al., 2003; Sabattini et al., 2010;
Munday et al., 2011; Munday et al., 2012; Dermatology et al., 2016, Sequeira et al.,
2022).
Individuals with a white coat all over their bodies develop deafness at a high
rate (Bergsma et al., 1971; Strain, 1996; Geigy et al., 2007; Ryugo et al., 2012; Strain,
2015; Kral et al., 2015; Strain, 2017; Mari et al., 2019; Bianchi et al., 2020). This
disease is thought to result from the inhibition of melanin synthesis and the
differentiation of melanocytes form the neural crest cells when *White* allele is present at
white locus (Geigy et al., 2007; David et al., 2014; Strain, 2017). In a cat with white
coat, a sequence derived from feline endogenous retrovirus 1 (*FERV1*) is inserted at the
white locus in the first intron of the tyrosine protein kinase (*KIT*) gene (David et al.,
2014). The *white spotting* allele of an individual with white-coated skin on a part of the
body surface (white-spotted cat) contains the full-length sequence of *FERV1* (7125 bp).
In contrast, the *White* allele of an individual whose whole body is completely covered
with a white coat (dominant white cat) contains only the long terminal repeat (LTR) of
FERV1 (622 bp) and not the full length. This suggests that the presence of LTR caused
a decrease in melanin synthesis. However, the role of LTR and the mechanism through
which melanin synthesis is suppressed remains unclear. White coat skin formation with
decreased melanin production is also associated with deafness in other animals, such as
dogs, miniature pigs, and yaks (Strain, 2015; Strain, 2017; Qi et al., 2022). However,
the underlying mechanisms have not been elucidated.

Thus, the relationship between the mechanisms that determine skin coat color
and disease risk remains poorly understood. Therefore, in this study, we analyzed the
RNA-seq data obtained from differently colored domestic cat skin cells published in a
public database. Through this analysis, the possible relationship between skin color and
squamous cell carcinoma risk and deafness as well as the mechanism by which genes
that determine skin coat color is involved in these diseases were inferred.

This study had two further aims. First was to report the relationship between
coat color and disease risk in domestic cats and its mechanism. The second was to
promote basic veterinary research on feline diseases and data accumulation that serves
as backbone for such research. Unfortunately, the availability of experimental data for
domestic cats is currently substantially limited compared to that for humans and canines.
In this study, only one replicate per skin cell color was available for RNA-seq analysis,
which limited the application of conventional methods to assess differentially expressed
genes (DEGs). Therefore, only forecast scenarios based on limited analysis could be
determined. However, this study is expected to achieve both purposes by providing the
motivation for various experimental studies and additional data analysis to validate the
present hypothesis.
126

Materials and Methods
Quantification of transcriptome data in 3 types of skin cells of *Felis catus*
From the Ensembl public database, specifically the file transfer protocol (ftp)
site, mapped RNA-seq data on the *Felis catus* genome, Fca126_mat1.0, were obtained
in bam and bam.bai formats. Data were retrieved from the orange skin cells of a cat with
orange and white coats (referred to as cell O), white skin cells of the cat with orange and
white coats (referred to as cell W), and skin cells of a cat ticked with white torties
(referred to as cell T). Each cell line had one replicate and the original Sequence Read
Archive (SRA) identifiers were SRS1335428, SRS1319391, and SRS1319719. Here, O
and W cells were obtained from the same cat, whereas T cells were obtained from a
different cat (https://www.ebi.ac.uk/biosamples/samples/SAMN04498546;
https://www.ebi.ac.uk/biosamples/samples/SAMN04498547;
https://www.ebi.ac.uk/biosamples/samples/SAMN04498539).
We should note the following facts for the samples:
I) Both cats belonged to the white-spotted cat classification, which means that the cell
genomes contained the full-length sequence derived from *FERV1* in the first intron of
the *KIT* gene.

II) Most of the body surface of cats with white tortie-ticked tabby is covered with a

dark-colored coat, such as brown or black. Therefore, for the analysis, it was assumed

that cell T represented a typical dark-colored skin cell. As mentioned below, this

assumption was supported by the expression levels of *tyrosinase* (*TYR*) and *tyrosinase*

related protein 1 (*TYRP1*) genes, which are located in the "color locus" and "brown

locus" loci.

III) Cats with O and W cells are carriers of lipoprotein lipase deficiency.

151    The annotation file of the *F. catus* genome (Fca126_mat1.0) in the gene

152 transfer format (GTF) was obtained from the National Centre for Biotechnology

153 Information (NCBI) database (Bethesda, MD, USA). In the file, the chromosome

154 numbers were converted from NC_058368.1, NC_058369.1, etc. to A1, A2, etc.

155    Using FeatureCounts with the options -M -O --fraction (Liao et al., 2014), the

number of reads for each gene described in the GTF file was obtained from the bam and

bam.bai formatted files for cells T, O, and W. From these counts, the transcript level of

each gene was calculated by dividing the read count by the total exon length. The total

transcript level was defined as the sum of all the transcript levels of the genes. Based on

this result, the transcripts per million (TPM) of each gene were calculated as 1000000 *

[transcript level] / [total transcript level] (see Table S1).

162    It is worth noting that the TPM values of the *TYR* and *TYRP1* genes in T cells

163 were more than ten times higher than those in W cells (Table S1). This observation

164 supports the previous assumption that T cells were sampled from dark-colored skin

165 rather than white skin in cats with a ticked tabby and white torties.

167 **Size factor of RNA-seq data**

168    As an example of size factors of RNA-seq data, the median of the ratios

(MOR) defined by the recent literature for DESeq (Anders, et al, 2010) was estimated

using read count data. For cells T, O, and W, the MORs were obtained 0.952545606,

1.201469023, and 0.868500362, respectively, and [standard deviation of MOR over

three cells] / [average MOR over three cells] was obtained 0.140326793. Based on these

results, it was expected that the size factor does not considerably affect the comparison

of TPM values.

Quantification of exon transcription levels of *KIT* in 3 types of skin cells of *F. catus*

177    A GTF file specifying each exon region of *KIT* was obtained from the GTF file

Fca126_mat1.0. The read count in each exon region of *KIT* was obtained from bam and

bam.bai formatted files with FeatureCounts (using the options: -t feature -M -O

--fraction) (Liao et al., 2014) for T, O, and W cells. The transcript level of each region
was calculated as the ratio of read count to region length. From the results, the TPM in
each exon region was obtained as 1000000 * [transcript level] / [total transcript level]
(Table S2), where the total transcript level uses the value calculated in the previous
section.
185
186 **Extraction of skin coat color-dependent differentially expressed genes (SCDEGs)**
187 The median TPM values (Table S1) obtained for all genes in all the cells (T, O,
188 and W) were approximately 0.987. Based on this fact, we considered genes that met the
189 following conditions as skin coat color-dependent differentially expressed genes
190 (SCDEGs):

- The maximum TPM values obtained for T, O, and W were more than u-times of
  the minimum value.
- The maximum TPM values obtained for T, O, and W were v or higher.

The following arguments, $u = 2$ and $v = 2$ (approximately equal to twice the median
value of TPM values) were assumed to estimate SCDEGs. Additionally, the SRY-Box
Transcription Factor 10 (*SOX10*) was included as an SCDEG. Although the maximum
expression level obtained from T, O, and W was 1.9 times the minimum value, which
does not satisfy the above requirement, *SOX10* is one of the predominant factors in
determining skin coat color (Mort et al., 2015; Strain, 2015; Qi et al., 2022).
Furthermore, the maximum expression of *SOX10* was nearly twice the minimum value.
Therefore, *SOX10* was specifically considered an SCDEG (Table S3).
202
Enrichment analysis
Pathway enrichment analysis of the SCDEGs was conducted using g:Profiler
(https://biit.cs.ut.ee/gprofiler/gost) (Raudvere et al., 2019). For this analysis, "felis
catus" was chosen as the target organism, and "only annotated genes" were selected as
the target gene set. The focus of the analysis was to identify significantly enriched
Kyoto Encyclopedia of Genes and Genomes (KEGG) pathways with a false discovery
rate (FDR) threshold of less than 0.1.
210
Classification into squamous cell carcinoma-promoting and suppressing genes
First, SCDEGs associated with the KEGG pathway were identified from the
results obtained through the aforementioned pathway enrichment analysis. Next, using
databases and prior literature, the extracted SCDEGs were categorized into the
following groups:

1. Carcinoma-promoting genes: These genes exhibited an excessive increase in
expression leading to promotion of squamous cell carcinoma.
2. Carcinoma suppressor genes: These were expressed at decreased levels or were
mutated; these genes contributed to the development of squamous cell
carcinoma.
3. Other genes: SCDEGs that did not fall into the above categories.
Categorization was based on the information obtained from databases and previous
studies (Table S5).
224
Theoretical ratio of the number of "tumor-promoting genes" and
"tumor-suppressor genes"
A file named cancermine_collated.tsv, which contained cancer gene roles and
citations, was obtained from the data download page
(https://zenodo.org/record/7689627) of the public database CancerMine
(http://bionlp.bcgsc.ca/cancermine/) (Lever et al., 2019). This file provided a list of
oncogenes and tumor suppressor genes for each of the 526 cancer types covered in the
database. The total number of oncogenes and tumor suppressor genes in all carcinomas
was 18,994 and 11,295, respectively. Therefore, we assumed that the "theoretical ratio
for all carcinomas" between the number of "carcinoma-promoting genes" and the
number of "carcinoma-suppressor genes" could be estimated based on these data.
Additionally, CancerMine contains data of 17 types of squamous cell
carcinomas (Table S4), where the total number of oncogenes and tumor suppressor
genes in all squamous cell carcinomas was 904 and 584, respectively. Therefore, we
also assumed that the "theoretical ratio for all squamous cell carcinomas" between the
number of "carcinoma-promoting genes" and the number of "carcinoma-suppressor
genes" could be estimated.
Although the database data used for estimating this theoretical ratio are specific
to human cancer, it is reasonable to assume that the ratio of the number of genes would
be similar between the two species because humans and cats are both mammals.
245
Evaluation of skin coat color dependent squamous cell carcinoma risk
To assess the cancer risk, the expression levels of carcinoma-promoting and
carcinoma-suppressor genes within the SCDEGs, specifically those contributing to
pathways related to squamous cell carcinoma, were compared among T, O, and W cells
using the following criteria:

1. The number of carcinoma-promoting genes whose expression level in cell X (X
= T, O, or W) was at least twice the minimum expression level among cells T, O,
or W was denoted as P_X.
2. The number of carcinoma-suppressor genes whose expression level in cell X (X
= T, O, or W) was at least twice the minimum expression level in cells T, O, or
W was denoted as S_X.
3. If P_X was greater than S_X, a comparison was made with the theoretical ratio
obtained earlier using an exact test of goodness-of-fit. If the resulting *p*-value is
less than 0.05, it indicates that cell X is in a state where squamous cell
carcinoma is more likely to develop and progress than in the other cells.

Now, it should be noted that whether the decreases in the expression levels of
tumor suppressor genes in normal cells could be useful for predicting carcinogenesis is
not trivial. Generally, mutations in some genes trigger tumorigenesis. For some
carcinomas, however, recent studies suggest that candidate tumor suppressor genes,
such as CDH1, PTEN, and KiSS1, expression levels decreased by epigenetic silencing,
rather than by deletion, cause carcinogenesis (Wang, et al., 2019). Thus, the ratio of
excessively expressed carcinoma promoting and suppressor genes in normal cells were
focused because such ratio might provide nonnegligible contributions to squamous cell
carcinoma risk.
270

Evaluations of motif sequence of CTCF binding and promoter activity in white
locus LTR
T, O, and W cells were derived from individual cats partially surrounded by
white skin and fur. In the genomes of these white-spotted cats, an LTR sequence
derived from *FERV1* was located in the first intron of *KIT*. To determine the specificity
of this LTR sequence, we searched for binding motif sequences of the CCCTC-binding
factor (CTCF) and motif sequence for core promoter elements within the LTR.
For this purpose, we utilized the CTCFBS Prediction Tool available in
CTCFBSDB 2.0 (https://insulatordb.uthsc.edu/) (Ziebarth et al., 2013) and Eukaryotic
Core Promoter Predictor (YAPP: http://www.bioinformatics.org/yapp/cgi-bin/yapp.cgi).
The former tool is based on a database containing the CTCF-binding motif sequences
from humans and mice, where the given sequence is considered to involve
CTCF-binding activity if the score of this tool is larger than 3. The latter tool is based
on the statistical analysis of data sets of eukaryotic core promoter elements (Cartharius
et al., 2005; Gershenzon et al., 2005; Jin et al., 2006), where the sequence with a score

larger than 0.8 (0 < score < 1) is considered to exhibit promoter activity; the sequences
with a score equal to or larger than 0.9 were focused in this study.
Feline CTCF has more than 99% of the same amino acid sequence as human
CTCF, and all of the DNA-binding motif amino acid sequences are the same (Fig. S1).
Therefore, although the CTCFBSDB database was primarily constructed based on
research conducted in humans and mice CTCF, the use of this database is reasonable for
the CTCF analysis of domestic cats.
293
294 **Statistical analysis**
295   *P*-values for various statistical analyses were calculated using R (version
296 4.2.3).
297
298
Result
SCDEGs and results of KEGG pathway enrichment analysis
To investigate the relationship between SCDEGs (Table S3) and the risk of
squamous cell carcinoma, KEGG pathway enrichment analysis was conducted using
g:Profiler (Table S5a). The analysis identified the following pathways related to
squamous cell carcinoma in which significantly larger number of SCDEGs were
involved.
1. Pathways in cancer (FDR ~ 3.58×10^{-3})
2. The p53 signaling pathway (FDR ~ 0.013×10^{-2})
3. Human papillomavirus (HPV) infection (FDR ~ 8.33×10^{-2})
HPV infection is implicated in the development of human squamous cell carcinoma and
cervical cancer (Hennessey et al., 2009; Husain et al., 2017; Augustin et al., 2020; Bao
et al., 2021; Chrysovergis et al., 2021; The et al., 2021; Karina, 2022; Kwon et al.,
2022). Feline papillomavirus (FPV) type-2 has been proposed as the primary cause of
squamous cell carcinoma in domestic cats (Munday et al., 2008; Munday et al., 2011;
O'Neill et al., 2011; Munday et al., 2012; Thomson et al., 2016; Munday et al., 2017;
Carrai et al., 2020; Yamashita-Kawanishi et al., 2021; Teh et al., 2021; Altamura et al.,
2022). HPV and FPV are distinct viruses; however, their *E5*, *E6*, and *E7* oncogenes
exhibit similar features (Jiang et al., 2014; Altamura et al., 2016; Karina, 2022).
Additionally, the pathogenesis of FPV-2 infection share pathways with HPV-induced
pathogenesis and carcinogenesis (Teh et al., 2021; Karina, 2022). Consequently, genes
involved in processes driven by HPV infection should also be considered genes
associated with squamous cell carcinoma.

322   KEGG pathway enrichment analysis might change depending on the chosen
323   SCDEGs determined by u and v values used. Here, to clarify the limitation of the above
324   result, the cases with u and v larger than 2 were exceptionally considered. Consequently,
325   the abovementioned three KEGG pathways related to squamous cell carcinoma
326   contained a substantial large number of SCDEGs (FDR < 0.1) when $u \leq 2.5$ and $v \leq 4.5$
327   (Table S5b).
328   Notably, the same analysis using RNA-seq read data of mesenteric lymph node
cells from 4 different domestic short hair cats did not exhibited similar results, where
"pathways in cancer" and "Human papillomavirus (HPV) infection" did not contain
significantly larger number of SCDEGs (Supporting information 1, Table S6a, and S6b).
This fact may support that the SCDEGs among T, O, and W cells and their squamous
cell carcinoma-related characteristics were rather specific than emerged due to
variations among individuals.
335

336   **Assessment of squamous cell carcinoma risk by comparing the expression of**
337   **carcinoma-promoting and carcinoma-suppressing genes**
338   SCDEGs that contributed to cancer pathways, the p53 signaling pathway, and
human papillomavirus infection were classified into squamous cell
carcinoma-promoting and squamous cell carcinoma-suppressing genes based on
previous studies (Table 1, Table S7). For each of the T, O, and W cells, the P_X and S_X
that exhibited stronger expression than the others were counted. The relationship
between P_X and S_X (X = T, O, W) was compared to the theoretical ratio of the number
of carcinoma-promoting and carcinoma-suppressor genes using the exact goodness of
fit test (Table 2).
346   The analysis revealed that the P_O in O cell was significantly higher than the
number of carcinoma-suppressor genes (S_O) ($P = 0.02246$ compared with the theoretical
ratio for all carcinomas, and 0.01421 compared with all squamous cell carcinomas). The
relationship between P_W and S_W displayed similar features, although the P-value
obtained was slightly larger ($P = 0.05194$ compared with the theoretical ratio for all
carcinomas, and 0.03445 compared with all squamous cell carcinomas). The values for
P_T, P_O, and P_W were similar. In contrast, S_O and S_W values were lower than that of S_T (P
value = 0.1296 using the Fisher's exact test), indicating that the expression levels of
carcinoma-suppressing genes were considerably lower in O and W cells than in T cells.
A similar result was also confirmed when $u = 2.5$ and $v = 4.5$ (Table S5c).
356

CTCF binding motif repeat in LTR at white locus in 1st intron of *KIT* gene

White-coated cats possessed a DNA sequence derived from *FERV1* located in

the first intron of the *KIT* gene. Furthermore, both *White* and *white spotting* alleles

includes a 5' LTR region, which is expected to inhibit melanocyte differentiation (David

et al., 2014). In this study, we examined the characteristics of the LTR sequence by

evaluating the presence of CTCF-binding motifs in this region using CTCFBS

Prediction Tool. The analysis identified putative seven CTCF-binding motifs within the

LTR that showed similarity to known CTCF-binding sites (Fig. 1a). Furthermore, these

motif sequences were arranged in a repeat pattern with a spacing of 46 base pairs

between them.

1st exon-lacking *KIT* (*d1KIT*) expression level increased in white skin cell

The repeat region of the putative CTCF-binding motif within the LTR of the

first *KIT* intron suggested a potential role of the LTR by binding of CTCF and

functioning as an insulator, thereby acting as a boundary for topologically associated

domains (TADs). *CTCF* has been found to be enriched at the edges of chromatin loops

that form TADs and cooperates with cohesins to regulate TAD boundary formation

(Parelho et al., 2008; Wendt et al., 2008; Merkenschlager et al., 2016). Genes located

within neighboring TADs are generally known to be independently regulated in their

expression.

Retroviral LTRs typically function as transcription promoters (Klaver et al.,

1994; Pereira et al., 2000; Dunn et al., 2003; Cohen et al., 2009). The *FERV1* LTR may

also possess promoter activity because the prediction tool YAPP found an initiator

motif (INR), which is a typical consensus motif sequence of core promoter region, at

the downstream of the repeat region of the putative CTCF-binding motif in LTR (Fig.

1a). Additionally, a downstream promoter element (DPE), which is also typical

consensus motif sequence of core promoter region, was also found at the downstream of

the INR (Fig. 1a). The DPE and the INR can function as TFIID-binding site

cooperatively, leading to accurate and efficient transcription (Burke and Kadonaga,

1996, 1997). Therefore, if CTCF-binding motifs exhibit insulating activity, it is possible

that the second exon and downstream regions of the *KIT* gene can be transcribed

independently of the native promoter, utilizing the region regulated by above pair of

INR and DPE as a core promoter (Fig. 1b). Now, we tentatively named the *KIT*

transcribed in such a manner and lacking the first exon *d1KIT* and quantified its

expression level in each cell type in the next section.

Based on the estimated TPM values provided in Tables S1 and S2, the ratio of

[TPM of the 1st exon] / [TPM of KIT] in T and O cells was in the range of 0.8 and 0.84,

respectively. Additionally, the ratio of [Average TPM of 2nd-22nd exons] / [TPM of
KIT] in T, O, and W cells was close to 1. However, in W cell, the ratio of [TPM of 1st
exon] / [TPM of KIT] was approximately 0.48, which was close to half of the values
observed in the other cells (Fig. 2). This suggests that approximately half of the *KIT*
mRNAs generated by W cells lacked the first exon.

The approximate transcript levels (TPM) of normal *KIT* and *d1KIT* in T, O, and
W cells were estimated as follows. First, since the amount of normal *KIT* mRNA was
expected to consist of those of the first exon and the second and subsequent exons,
[TPM of normal KIT] was assumed to be the minimum value between [TPM of 1st
exon] and [TPM of KIT]. Second, the TPM of d1KIT was estimated as follows: [TPM
of d1KIT] = [TPM of KIT] – [TPM of normal KIT]. The TPM of normal *KIT* in T, O,
and W cells was approximately 9.35, 20.50, and 9.87, respectively, whereas that of
d1KIT in T, O, and W cells was approximately 1.67, 5.00, and 10.49, respectively (Fig.
2). Here, the ratio [TPM of d1KIT]/[TPM of normal KIT] in T, O, and W cells was
0.179929164, 0.244270964, and 1.063338179, respectively.

Notably, the transcript levels of normal KIT and d1KIT could not be obtained
directly from short read RNA-seq data as analyzed in this study; however, they could be
estimated using various methods. Thus, to confirm the validity of the abovementioned
results, [d1KIT expression]/[Normal KIT expression] was estimated using the
maximum likelihood estimate (MLE) method, using the number of read of each exon of
KIT. Here, the [d1KIT expression]/[Normal KIT expression] for T, O, and W cells was
0.191814285, 0.259627163, and 1.112849284, respectively, which seemed similar to
the above ratio [TPM of d1KIT]/[TPM of normal KIT] (Supporting information 2).
Therefore, [TPM of normal KIT] and [TPM of d1KIT] were used as transcript levels of
normal KIT and d1KIT in the following arguments.

***d1KIT* expression correlated negatively to gene expressions for melanocyte**
formation

To determine how normal KIT and d1KIT are involved in melanocyte
formation, we examined the correlation between their expression levels and those of
typical genes involved in melanoblast and melanocyte formation (Table S8). The
expression level of *d1KIT* showed a significant negative correlation with the expression
level of *SOX10*, with a correlation coefficient of approximately -0.99957 (*P*-value ~
0.01876) (Fig. 3). Furthermore, strong negative correlations were observed with the
expression levels of melanocyte-related *DKK4* and *TYRP1*, with correlation coefficients
of approximately -0.98777 (*P*-value ~ 0.09968) and -0.98000 (*P*-value ~ 0.12763),

respectively. In contrast, the correlation coefficient between the expression levels of
normal *KIT* and melanocyte protein *PMEL* ~ 0.94548 (*P*-value ~ 0.20302) that
exhibited the largest absolute value of correlation coefficient (Fig. 3). This fact
suggested the expression level of normal KIT did not strongly correlate with the
expression of genes involved in melanoblast and melanocyte formation.
435
Discussion
Squamous cell carcinoma tends to occur more frequently in light-coated cats,
which may be due to a lack of melanin production, resulting in poor UV resistance. If
this is the case, it would be expected that white-coated skin, without melanin production,
would be most susceptible to carcinoma. However, in our analysis, we found the
possibility that, in terms of gene expression, the expression level of a factor that
suppresses squamous cell carcinoma was significantly lower in cells with orange-coated
skin with *white spotting* allele. In other words, when considering gene expression, cells
with not only white- but also orange-coated skin may have higher risks of cancer than
cells with other skin colors.
We also observed that the ratio of the expression level of genes suppressing
squamous cell carcinoma to that of genes promoting squamous cell carcinoma was
lower than the theoretical estimation in cells with white coated skin with *white spotting*
allele. It is worth noting that the orange- and white-coated skin cells analyzed in this
study were sampled from the same individuals and were derived from carriers of
lipoprotein lipase deficiency. These findings suggest two possible mechanisms
underlying the risk of squamous cell carcinoma.
1. Cells for light-colored skin, such as white- and orange-coated skin, tend to
454   exhibit low expression of carcinoma suppressor genes.
455   2. Cells derived from carriers with lipoprotein lipase deficiency tended to exhibit
456   low expression of carcinoma-suppressor genes.
Validation of these possibilities through analysis of more accumulated data and
additional experiments should be conducted in future.
Here, squamous cell carcinoma develops from cells in the epidermis. Thus, the
ideal samples would be squamous cells or at least epidermal cells rather than skin cells
as used in the present study. Feline squamous cell carcinoma is highly malignant. For
example, oral squamous cell carcinoma has a poor prognosis with a one-year survival
rate of less than 10% (Martin et al., 2011). However, there are no public RNA-seq data
on squamous cells or epidermal cells derived from cats with different coat colors.
Therefore, in this study, the analysis of the data obtained from skin cells was performed

to see the possibilities expected from it, knowing that there were several limitations.
However, more accurate prediction of squamous cell carcinoma risk becomes possible
by obtaining and analyzing the data through experiments of squamous cells and
epidermal cells from cats with different coat colors. Such studies should be progressed
in the future.
The contributions of each gene to carcinogenesis seemed to be different. For
example, change in expression level of the *TP53* should be critical, while those in others
might have moderate effects. However, there were no appropriate way to quantify the
contribution weight of each gene to the carcinoma risks. Thus, in the present study, we
assumed that such weight was approximately uniform as first trial analysis. However,
the estimation of such weight is crucial for a more accurate carcinoma risk assessment.
In white-coated skin cells, a large amount of *d1KIT* mRNA, in which the 1st
exon of the *KIT* gene is deleted, might be produced. Furthermore, we found the
possibility that the LTR of *FERV1*, which was present in *White* and *white spotting*
alleles at white locus in the first intron of the *KIT* gene in the genome of a domestic cat
with white-coated skin, contains a repeated CTCF-binding site. This finding suggests
the possibility that the transcription from the 1st exon and that from the 2nd exon are
separately regulated by the presence of the promoter and insulator activities of this LTR,
which provided the mechanism for d1KIT production.
Additionally, the amount of *d1KIT* transcript was found to exhibit a significant
negative correlation with the expression levels of *SOX10*, as well as strong negative
correlations with genes that induce differentiation into melanocytes. These results
suggest the possibility that the insulating activity at the white locus, specifically the
LTR of *FERV1*, resulted in the production of d1KIT and subsequent inhibition of the
SOX10 expression, which was crucial for melanocyte formation.
Notably, Sox10 mutations cause deafness in miniature pigs, leading to
degeneration and a considerable decrease in the density of utricular hair cells, cochlear
abnormalities, and saccular hypofunction (Qi et al., 2022). Therefore, when the
aforementioned LTR activities occur throughout the body, it is conceivable that the
entire skin becomes covered with white fur and deafness is induced. We hypothesized
that this is the mechanism underlying dominant white cat deafness.
However, in white-spotted cats, from which the cells sampled in this study
were derived, LTR activity and its influences on the KIT gene expression might be
limited depending on cell lineages or cell types, since the full-length of FERV1 is
present at *white* locus. For example, between O and W cells differences in 3D genomic
structures of FERV1 other than LTR or those in DNA methylation status of the LTR

promoter region might appear, and contributed to the around twice difference of
expression of the d1KIT appeared. However, the precise mechanism remains unclear,
and its elucidation is an issue for future research through the epigenetic measurements
such as Hi-C and 4C-seq, methylation, and nucleosome binding around FERV1 in the
first intron of KIT.

Furthermore, for both dominant white and white spotted cats, it was difficult to
show a causal relationship between d1KIT production and *SOX10* expression only in
the present study. Such relationship should be confirmed by future experiments.

As mentioned in the last part of introduction section, the availability of
experimental data for *F. catus* (domestic cats) is currently significantly limited
compared to that for humans and canines, which are also common companion animals.
For instance, the number of *F. catus* RNA-seq samples registered in the public database
COXPRESDB (https://coxpresdb.jp/RNA-seq) (Obayashi et al., 2023), which allows
the search of co-expressed genes, is only 267, whereas the numbers for humans and
canines are much larger (235,187 and 1,361, respectively). Additionally, the specific
roles of each gene in the KEGG pathways obtained through enrichment analysis have
rarely been directly confirmed in domestic cats, with most predictions being
extrapolated from results in humans and mice.

To advance the field of basic veterinary medicine and support clinical
veterinary practices not only for domestic cats, but also for various domestic and
companion animals, it is crucial to accumulate a substantial amount of diverse basic
data.

524
525

Author Contributions
AA, KW and NS conceived and designed the study. AA and DK performed the analyses.
AA and NS drafted the manuscript.

Acknowledgements
This work was supported by a Grant-in-Aid for Scientific Research (KAKENHI) [Grant
Number 21K06124] from the Japan Society for the Promotion of Science.
Computations were partially performed on the NIG supercomputer at ROIS National
Institute of Genetics.

535
536

537 **References**

538 Adega, F., Borges, A., & Chaves, R. (2016). Cat Mammary Tumors: Genetic Models
539 for the Human Counterpart. *Veterinary Sciences*, *3*(3), 17.
540 https://doi.org/10.3390/vetsci3030017
541
542 Altamura, G., Corteggio, A., Pacini, L., Conte, A., Pierantoni, G. M., Tommasino, M.,
543 Accardi, R., & Borzacchiello, G. (2016). Transforming properties of Felis catus
544 papillomavirus type 2 E6 and E7 putative oncogenes in vitro and their transcriptional
545 activity in feline squamous cell carcinoma in vivo. *Virology*, *496*, 1-8.
546 https://doi.org/10.1016/j.virol.2016.05.017
547
548 Altamura, G., Cuccaro, B., Eleni, C., Strohmayer, C., Brandt, S., & Borzacchiello, G.
549 (2022). Investigation of multiple Felis catus papillomavirus types (-1/-2/-3/-4/-5/-6)
550 DNAs in feline oral squamous cell carcinoma: A multicentric study. *The Journal of*
551 *Veterinary Medical Science*, *84*(6), 881-884. https://doi.org/10.1292/jvms.22-0060
552
553 Anders, S., & Huber, W. (2010). Differential expression analysis for sequence count
554 data. *Genome Biol.*, *11*, R106.
555 https://doi.org/10.1186/gb-2010-11-10-r106
556
557 Andrade, L. A., Versiani, A. F., F., E., dos Reis, J. K., dos Reis, J. G., & da Fonseca, F.
558 G. (2022). Developing a Feline Immunodeficiency Virus Subtype B Vaccine Prototype
559 Using a Recombinant MVA Vector. *Vaccines*, *10*(10), 1717.
560 https://doi.org/10.3390/vaccines10101717
561
562 Arai, S., Kitada, K., Yamazaki, T., Takai, R., Zhang, X., Tsugawa, Y., ... & Miyazaki, T.
563 (2016). Apoptosis inhibitor of macrophage protein enhances intraluminal debris
564 clearance and ameliorates acute kidney injury in mice. *Nature Medicine*, *22*(2), 183-193.
565 https://doi.org/10.1038/nm.4012
566
567 Augustin, J. G., Lepine, C., Morini, A., Brunet, A., Veyer, D., Brochard, C., Mirghani,
568 H., Péré, H., & Badoual, C. (2020). HPV Detection in Head and Neck Squamous Cell
569 Carcinomas: What Is the Issue? *Frontiers in Oncology*, *10*, 553220.
570 https://doi.org/10.3389/fonc.2020.01751
571

572   Baer, K. E., & Helton, K. (1993). Multicentric squamous cell carcinoma in situ
573   resembling Bowen's disease in cats. *Veterinary Pathology*, *30*(6), 535-543.
574   https://doi.org/10.1177/030098589303000607
575

Bao, Y., Gabrielpillai, J., Dietrich, J., Zarbl, R., Strieth, S., Schröck, F., & Dietrich, D.
(2021). Fibroblast growth factor (FGF), FGF receptor (FGFR), and cyclin D1 (CCND1)
DNA methylation in head and neck squamous cell carcinomas is associated with
transcriptional activity, gene amplification, human papillomavirus (HPV) status, and
sensitivity to tyrosine kinase inhibitors. *Clinical Epigenetics 13*, 228.
https://doi.org/10.1186/s13148-021-01212-4
582

583   Bergsma, D. R., & Brown, K. S. (1971). White Fur, Blue Eyes, and Deafness in the
584   Domestic Cat. *Journal of Heredity*, *62*(3), 171-183.
585   https://doi.org/10.1093/jhered/62.3.171
586

587   Bertone, E. R., Snyder, L. A., & Moore, A. S. (2003). Environmental and Lifestyle Risk
588   Factors for Oral Squamous Cell Carcinoma in Domestic Cats. *Journal of Veterinary*
589   *Internal Medicine*, *17*(4), 557-562. https://doi.org/10.1111/j.1939-1676.2003.tb02478.x
590

591   Bianchi, E., Cantoni, A. M., & Poncelet, L. (2020). The acoustically evoked short
592   latency negative response (ASNR) in a unilaterally deaf cat with
593   histologically-confirmed cochleosaccular degeneration. *BMC Veterinary Research*, *16*,
594   1-5. https://doi.org/10.1186/s12917-020-02426-z
595

596   Bredemeyer, K. R., Harris, A. J., Li, G., Zhao, L., Foley, N. M., J, S., Lyons, L. A.,
597   Warren, W. C., & Murphy, W. J. (2021). Ultracontinuous Single Haplotype Genome
598   Assemblies for the Domestic Cat (Felis catus) and Asian Leopard Cat (Prionailurus
599   bengalensis). *Journal of Heredity*, *112*(2), 165-173.
600   https://doi.org/10.1093/jhered/esaa057
601

602   Burke, T.W., & Kadonaga, J.T. (1996). Drosophila TFIID binds to a conserved
603   downstream basal promoter element that is present in many TATA-box-deficient
604   promoters. *Genes & Development*, 10(6), 711-724. https://doi.org/10.1101/gad.10.6.711
605

606   Burke, T.W., & Kadonaga, J.T. (1997). The downstream core promoter element, DPE,
607   is conserved from Drosophila to humans and is recognized by TAFII60 of Drosophila.
608   Genes & Development, 11(22), 3020-3031. https://doi.org/10.1101/gad.11.22.3020
609
610   Caliari, D., Zappulli, V., Rasotto, R., Cardazzo, B., Frassineti, F., Goldschmidt, M. H.,
611   & Castagnaro, M. (2014). Triple-negative vimentin-positive heterogeneous feline
612   mammary carcinomas as a potential comparative model for breast cancer. *BMC*
613   *Veterinary Research*, *10*(1), 185. https://doi.org/10.1186/s12917-014-0185-8
614
615   Cannon, C. M. (2015). Cats, Cancer and Comparative Oncology. *Veterinary Sciences*,
616   *2*(3), 111-126. https://doi.org/10.3390/vetsci2030111
617
618   Carrai, M., Van Brussel, K., Shi, M., Li, C. X., Chang, W. S., Munday, J. S., ... &
619   Beatty, J. A. (2020). Identification of a novel papillomavirus associated with squamous
620   cell carcinoma in a domestic cat. *Viruses*, *12*(1), 124.
621   https://doi.org/10.3390/v12010124
622
623   Cartharius, K., Frech, K., Grote, K., Klocke, B., Haltmeier, M., Klingenhoff, A., Frisch,
624   M., Bayerlein, M., & Werner, T. (2005). MatInspector and beyond: promoter analysis
625   based on transcription factor binding sites. *Bioinformatics*, 21(13), 2933-2942.
626   https://doi.org/10.1093/bioinformatics/bti473
627
628   Chambers, J. K., Tokuda, T., Uchida, K., Ishii, R., Tatebe, H., Takahashi, E., Tomiyama,
629   T., Une, Y., & Nakayama, H. (2015). The domestic cat as a natural animal model of
630   Alzheimer's disease. *Acta Neuropathologica Communications*, *3*, 78.
631   https://doi.org/10.1186/s40478-015-0258-3
632
633   Chrysovergis, A., Papanikolaou, V., Mastronikolis, N., Spyropoulou, D., Adamopoulou,
634   M., Tsiambas, E., ... & Kyrodimos, E. (2021). C-Fos Digital Expression Analysis in
635   Human Papillomavirus Related Oral Squamous Cell Carcinoma. *Asian Pacific*
636   *Journal of Cancer Biology*, *6*(2), 117-121. 10.31557/APJCB.2021.6.2.117-121
637
638   Cohen, C. J., Lock, W. M., & Mager, D. L. (2009). Endogenous retroviral LTRs as
639   promoters for human genes: A critical assessment. *Gene*, *448*(2), 105-114.
640   https://doi.org/10.1016/j.gene.2009.06.020
641

Cook, S., Wittenburg, L., Yan, V. C., Theil, J. H., Castillo, D., Reagan, K. L., ... &
Murphy, B. G. (2022). An Optimized Bioassay for Screening Combined
Anticoronaviral Compounds for Efficacy against Feline Infectious Peritonitis Virus with
Pharmacokinetic Analyses of GS-441524, Remdesivir, and Molnupiravir in Cats.
Viruses, *14*(11), 2429.   https://doi.org/10.3390/v14112429

Cunningham, F., Allen, J. E., Allen, J., Amode, M. R., Armean, I. M., Azov, A. G., …
Flicek, P. (2022). Ensembl 2022. *Nucleic Acids Research*, *50*(D1), D988-D995.
https://doi.org/10.1093/nar/gkab1049

David, V. A., Wallace, A. C., Roelke, M., Kehler, J., Leighty, R., Eizirik, E., … &
Ryugo, D. K. (2014). Endogenous Retrovirus Insertion in the KIT Oncogene
Determines White and White spotting in Domestic Cats. *G3 Genes|Genomes|Genetics*,
4(10), 1881-1891. https://doi.org/10.1534/g3.114.013425

Dorn, C. R., Taylor, D. O., & Schneider, R. (1971). Sunlight Exposure and Risk of
Developing Cutaneous and Oral Squamous Cell Carcinomas in White Cats. *JNCI:*
Journal of the National Cancer Institute, *46*(5), 1073-1078.
https://doi.org/10.1093/jnci/46.5.1073

Dunn, C. A., Medstrand, P., & Mager, D. L. (2003). An endogenous retroviral long
terminal repeat is the dominant promoter for human β1,3-galactosyltransferase 5 in the
colon. *Proceedings of the National Academy of Sciences of the United States of America*,
100(22), 12841-12846. https://doi.org/10.1073/pnas.2134464100

Eizirik, E., David, V. A., Buckley-Beason, V., Roelke, M. E., Schäffer, A. A., Hannah,
S. S., … & Reed, K. (2010). Defining and mapping mammalian coat pattern genes:
multiple genomic regions implicated in domestic cat stripes and spots. *Genetics*. *184*(1),
267–75.   https://doi.org/10.1534/genetics.109.109629

Geigy, C. A., Heid, S., Steffen, F., Danielson, K., Jaggy, A., & Gaillard, C. (2007).
Does a pleiotropic gene explain deafness and blue irises in white cats? *The Veterinary*
Journal, *173*(3), 548-553. https://doi.org/10.1016/j.tvjl.2006.07.021
675

Gershenzon, N., & Ioshikhes, I. (2005). Synergy of human Pol II core promoter

elements revealed by statistical sequence analysis. *Bioinformatics*, 21(8), 1295-300.

https://doi.org/10.1093/bioinformatics/bti172

Hennessey, P. T., Westra, W. H., & Califano, J. A. (2009). Human Papillomavirus and

Head and Neck Squamous Cell Carcinoma: Recent Evidence and Clinical Implications.

Journal of Dental Research, *88*(4), 300-306.

https://doi.org/10.1177/0022034509333371

Hofmann-Lehmann, R., & Hartmann, K. (2020). Feline leukaemia virus infection: A

practical approach to diagnosis. *Journal of Feline Medicine and Surgery*, *22*(9),

831-846. https://doi.org/10.1177/1098612X20941785

Howard, J., Wyse, C., Argyle, D., Quinn, C., Kelly, P., & McCann, A. (2020).

Exosomes as biomarkers of human and feline mammary tumours; a comparative

medicine approach to unravelling the aggressiveness of TNBC. *Biochimica et*

Biophysica Acta (BBA)-Reviews on Cancer, *1874*(2), 188431.

10.1016/j.bbcan.2020.188431

Husain, N., & Neyaz, A. (2017). Human papillomavirus associated head and neck

squamous cell carcinoma: Controversies and new concepts. *Journal of Oral Biology*

and Craniofacial Research, *7*(3), 198-205. https://doi.org/10.1016/j.jobcr.2017.08.003

Imes, D. L., Geary, L. A., Grahn, R. A., & Lyons, L. A. (2006). Albinism in the

domestic cat (Felis catus) is associated with a tyrosinase (TYR) mutation. *Animal*

Genetics, *37*(2), 175-178. https://doi.org/10.1111/j.1365-2052.2005.01409.x

Ishida, Y., David, V. A., Eizirik, E., Schäffer, A. A., Neelam, B. A., Roelke, M. E., …

& Menotti-Raymond, M. (2006). A homozygous single-base deletion in MLPH causes

the dilute coat color phenotype in the domestic cat. *Genomics*, *88*, 698–705.

10.1016/j.ygeno.2006.06.006

Jiang, P., & Jiang, P. (2014). Human papillomavirus oncoproteins and apoptosis

(Review). *Experimental and Therapeutic Medicine*, *7*, 3-7.

https://doi.org/10.3892/etm.2013.1374

Jin, V.X., Singer, G.A., Agosto-Perez, F.J., Liyanarachchi, S., & Davuluri, RV. (2006).

Genome-wide analysis of core promoter elements from conserved human and mouse

orthologous pairs. *BMC Bioinformatics*, 7, 114.

https://doi.org/10.1186/1471-2105-7-114

Karina, A. (2022). Pathological similarities in the development of

Papillomavirus-associated cancer in humans, dogs, and cats. *Animals*, *12*(18), 2390.

https://doi.org/10.3390/ani12182390

Klaver, B., & Berkhout, B. (1994). Comparison of 5' and 3' long terminal repeat

promoter function in human immunodeficiency virus. *Journal of Virology*, *68*(6),

3830-3840. https://doi.org/10.1128/jvi.68.6.3830-3840.1994

Kral, A., & Lomber, S. G. (2015). Deaf white cats. *Current Biology*, *25*(9), R351-R353.

https://doi.org/10.1016/j.cub.2015.02.040

Kwon, E. J., Lee, H. R., Lee, J. H., Seo, C., Ha, M., Roh, J., ... & Jang, J. Y. (2022).

Identification of differentially expressed genes and pathways for risk stratification in

HPV-associated cancers governing different anatomical sites. *Frontiers in*

Bioscience-Landmark, *27*(1), 2. https://doi.org/10.31083/j.fbl2701002

Lever, J., Zhao, E. Y., Grewal, J., Jones, M. R., & Jones, S. J. (2019). CancerMine: A

literature-mined resource for drivers, oncogenes and tumor suppressors in cancer.

Nature Methods, *16*(6), 505-507. https://doi.org/10.1038/s41592-019-0422-y

Liao, Y., Smyth, G. K., & Shi, W. (2014). FeatureCounts: An efficient general purpose

program for assigning sequence reads to genomic features. *Bioinformatics*, *30*(7),

923-930. https://doi.org/10.1093/bioinformatics/btt656

Lyons, L. A., Foe, I. T., Rah, H. C., & Grahn, R. A. (2005). Chocolate coated cats:

TYRP1 mutations for brown color in domestic cats. *Mammalian Genome*, *16*(5),

356–66. https://doi.org/10.1007/s00335-004-2455-4.

Lyons, L. A., Buckley, R. M., Harvey, R. J., & Genome Consortium, L. C. (2021).

Mining the 99 Lives Cat Genome Sequencing Consortium database implicates genes

and variants for the Ticked locus in domestic cats (Felis catus). *Animal Genetics*, *52*(3),

321-332. https://doi.org/10.1111/age.13059

Mari, L., Freeman, J., Dijk, J. V., & Risio, L. D. (2019). Prevalence of congenital

sensorineural deafness in a population of client-owned purebred kittens in the United

Kingdom. *Journal of Veterinary Internal Medicine*, *33*(4), 1707-1713.

https://doi.org/10.1111/jvim.15515

Martin, C. K., Tannehill-Gregg, S. H., Wolfe, T. D., & Rosol, T. J. (2011).

Bone-Invasive Oral Squamous Cell Carcinoma in Cats: Pathology and Expression of

Parathyroid Hormone-Related Protein. Veterinary Pathology, 48(1), 302-312.

https://doi.org/10.1177/0300985810384414

Merkenschlager, M., & Nora, E. P. (2016). CTCF and Cohesin in Genome Folding and

Transcriptional Gene Regulation. *Annual Review of Genomics and Human Genetics*, *17*,

17–43.https://doi.org/10.1146/annurev-genom-083115-022339

Millanta, F., Andreani, G., Rocchigiani, G., Lorenzi, D., & Poli, A. (2016). Correlation

Between Cyclo-oxygenase-2 and Vascular Endothelial Growth Factor Expression in

Canine and Feline Squamous Cell Carcinomas. *Journal of Comparative Pathology*,

154(4), 297-303. https://doi.org/10.1016/j.jcpa.2016.02.005

Munday, J. S., & Aberdein, D. (2012). Loss of retinoblastoma protein, but not p53, is

associated with the presence of papillomaviral DNA in feline viral plaques, Bowenoid

in situ carcinomas, and squamous cell carcinomas. *Veterinary Pathology*, *49*(3),

538-545.https://doi.org/10.1177/0300985811419534

Munday, J. S., Gibson, I., & French, A. F. (2011). Papillomaviral DNA and increased

p16CDKN2A protein are frequently present within feline cutaneous squamous cell

carcinomas in ultraviolet-protected skin. *Veterinary Dermatology*, *22*(4), 360-366.

https://doi.org/10.1111/j.1365-3164.2011.00958.x

Munday, J. S., Kiupel, M., French, A. F., & Howe, L. (2008). Amplification of

papillomaviral DNA sequences from a high proportion of feline cutaneous in situ and

invasive squamous cell carcinomas using a nested polymerase chain reaction.

Veterinary Dermatology, *19*(5),

259-263.https://doi.org/10.1111/j.1365-3164.2008.00685.x

Munday, J. S., Thomson, N. A., & Luff, J. A. (2017). Papillomaviruses in dogs and cats.

The Veterinary Journal, *225*, 23-31. https://doi.org/10.1016/j.tvjl.2017.04.018

Murphy, S. (2013). Cutaneous squamous cell carcinoma in the cat: current

understanding and treatment approaches. *Journal of Feline Medicine and Surgery*, *15*(5),

401-407. https://doi.org/10.1177/1098612X1348323

O'Neill, S. H., Newkirk, K. M., Anis, E. A., Brahmbhatt, R., Frank, L. A., & Kania, S.

A. (2011). Detection of human papillomavirus DNA in feline premalignant and invasive

squamous cell carcinoma. *Veterinary Dermatology*, *22*(1), 68-74.

https://doi.org/10.1111/j.1365-3164.2010.00912.x

Obayashi, T., Kodate, S., Hibara, H., Kagaya, Y., & Kinoshita, K. (2023). COXPRESdb

v8: An animal gene coexpression database navigating from a global view to detailed

investigations. *Nucleic Acids Research*, *51*(D1), D80-D87.

https://doi.org/10.1093/nar/gkac983

Parelho, V., Hadjur, S., Spivakov, M., Leleu, M., Sauer, S., Gregson, H. C., … &

Merkenschlager, M. (2008). Cohesins Functionally Associate with CTCF on

Mammalian Chromosome Arms. *Cell*, *132*(3),

422–433.https://doi.org/10.1016/j.cell.2008.01.011

Pedersen, N. C., Perron, M., Bannasch, M., Montgomery, E., Murakami, E., Liepnieks,

M., & Liu, H. (2019). Efficacy and safety of the nucleoside analog GS-441524 for

treatment of cats with naturally occurring feline infectious peritonitis. *Journal of Feline*

Medicine and Surgery, *21*(4), 271-281. https://doi.org/10.1177/1098612X19825701

Pereira, L. A., Bentley, K., Peeters, A., Churchill, M. J., & Deacon, N. J. (2000).

SURVEY AND SUMMARY A compilation of cellular transcription factor interactions

with the HIV-1 LTR promoter. *Nucleic Acids Research*, *28*(3), 663-668.

https://doi.org/10.1093/nar/28.3.663

Qi, J. C., Jiang, Q. Q., Ma, L., Yuan, S. L., Sun, W., Yu, L. S., ... & Yang, S. M. (2022).
Sox10 gene is required for the survival of saccular and utricular hair cells in a porcine
model. *Molecular Neurobiology*, *59*(6), 3323-3335.
https://doi.org/10.1007/s12035-021-02691-5

Raudvere, U., Kolberg, L., Kuzmin, I., Arak, T., Adler, P., Peterson, H., & Vilo, J.
(2019). G:Profiler: a web server for functional enrichment analysis and conversions of
gene lists (2019 update). *Nucleic Acids Research*, *47*(W1), W191-W198.
https://doi.org/10.1093/nar/gkz369

Ryugo, D. K., & Menotti-Raymond, M. (2012). Feline Deafness. *Veterinary Clinics of*
North America: Small Animal Practice, *42*(6), 1179.
https://doi.org/10.1016/j.cvsm.2012.08.008

Sabattini, S., Marconato, L., Zoff, A., Morini, M., Scarpa, F., Capitani, O., & Bettini, G.
(2010). Epidermal growth factor receptor expression is predictive of poor prognosis in
feline cutaneous squamous cell carcinoma. *Journal of Feline Medicine and Surgery*,
12(10), 760-768. https://doi.org/10.1016/j.jfms.2010.04.010

Schmidt-Küntzel, A., Nelson, G., David, V. A., Schäffer, A. A., Eizirik, E., Roelke, M.
E., … & Menotti-Raymond, M. (2009). A domestic cat X chromosome linkage map and
the sex-linked orange locus: mapping of orange, multiple origins and epistasis over
nonagouti. *Genetics*, *181*(4), 1415–25. https://doi.org/10.1534/genetics.108.095240

Schmidt-Küntzel, A., Eizirik, E., O'Brien, S. J., & Menotti-Raymond, M. (2005).
Tyrosinase and tyrosinase related protein 1 alleles specify domestic cat coat color
phenotypes of the albino and brown loci. *Journal of Heredity*, *96*(4), 289-301.
https://doi.org/10.1093/jhered/esi066

Sequeira, I., Pires, M. D., Leitão, J., Henriques, J., Viegas, C., & Requicha, J. (2022).
Feline Oral Squamous Cell Carcinoma: A Critical Review of Etiologic Factors.
Veterinary Sciences, *9*(10), 558. https://doi.org/10.3390/vetsci9100558

Sommerville, L., Howard, J., Evans, S., Kelly, P., & McCann, A. (2022). Comparative
gene expression study highlights molecular similarities between triple negative breast

cancer tumours and feline mammary carcinomas. *Veterinary and Comparative*
Oncology, *20*(2), 535-538. https://doi.org/10.1111/vco.12800
854
855 Sordo, L., & Gunn-Moore, D. A. (2021). Cognitive Dysfunction in Cats: Update on
856 Neuropathological and Behavioural Changes Plus Clinical Management. *Veterinary*
857 *Record*, *188*(1), N/A. https://doi.org/10.1002/vetr.3
858
859 Strain, G. M. (1996). Aetiology, prevalence and diagnosis of deafness in dogs and cats.
860 *British Veterinary Journal*, *152*(1), 17-36.
861 https://doi.org/10.1016/S0007-1935(96)80083-2
862
863 Strain, G. M. (2015). The Genetics of Deafness in Domestic Animals. *Frontiers in*
864 *Veterinary Science*, *2*, 155695. https://doi.org/10.3389/fvets.2015.00029
865
866 Strain, G. M. (2017). Hearing disorders in cats: Classification, pathology and diagnosis.
867 *Journal of Feline Medicine and Surgery*, *19*(3), 276-287.
868 https://doi.org/10.1177/1098612X17695062
869
870 Sugisawa, R., Hiramoto, E., Matsuoka, S., Iwai, S., Takai, R., Yamazaki, T., ... &
871 Miyazaki, T. (2016). Impact of feline AIM on the susceptibility of cats to renal disease.
872 *Scientific Reports*, *6*(1), 35251. https://doi.org/10.1038/srep35251
873
874 Teh, A., & Krockenberger, M. (2021). Do papillomaviruses cause feline cutaneous
875 squamous cell carcinoma? *Veterinary Evidence*, *6*(3).
876 https://doi.org/10.18849/ve.v6i3.402
877
878 Thomas, R. (2015). Cytogenomics of Feline Cancers: Advances and Opportunities.
879 *Veterinary Sciences*, *2*(3), 246-258. https://doi.org/10.3390/vetsci2030246
880
881 Thomson, M. (2007). Squamous Cell Carcinoma of the Nasal Planum in Cats and Dogs.
882 *Clinical Techniques in Small Animal Practice*, *22*(2), 42-45.
883 https://doi.org/10.1053/j.ctsap.2007.03.002
884
Thomson, N. A., Munday, J. S., & Dittmer, K. E. (2016). Frequent detection of
transcriptionally active Felis catus papillomavirus 2 in feline cutaneous squamous cell

carcinomas. *Journal of General Virology*, *97*(5), 1189-1197.

https://doi.org/10.1099/jgv.0.000416

Wang, L. H., Wu, C. F., Rajasekaran, N., & Shin, Y. K. (2019). Loss of tumor

suppressor gene function in human cancer: an overview. Cellular Physiology and

Biochemistry, 51(6), 2647-2693. https://doi.org/10.1159/000495956.

Wendt, K. S., Yoshida, K., Itoh, T., Bando, M., Koch, B., Schirghuber, E., … & Peters,

J. M. (2008). Cohesin mediates transcriptional insulation by CCCTC-binding factor.

Nature, *451*(7180), 796–801. https://doi.org/10.1038/nature06634

Westman, M. E., Malik, R., & Norris, J. M. (2019). Diagnosing feline

immunodeficiency virus (FIV) and feline leukaemia virus (FeLV) infection: an update

for clinicians. *Australian Veterinary Journal*, *97*(3), 47-55.

https://doi.org/10.1111/avj.12781

Wiese, D. A., Thaiwong, T., Yuzbasiyan-Gurkan, V., & Kiupel, M. (2013). Feline

mammary basal-like adenocarcinomas: a potential model for human triple-negative

breast cancer (TNBC) with basal-like subtype. *BMC Cancer*, *13*(1), 403.

https://doi.org/10.1186/1471-2407-13-403

Yamashita-Kawanishi, N., Gushino, Y., Chang, C. Y., Chang, H. W., Chambers, J. K.,

Uchida, K., & Haga, T. (2021). Full-genome characterization of a novel Felis catus

papillomavirus 4 subtype identified in a cutaneous squamous cell carcinoma of a

domestic cat. *Virus Genes*, *57*(4), 380-384. https://doi.org/10.1007/s11262-021-01843-y

Ziebarth, J. D., Bhattacharya, A., & Cui, Y. (2013). CTCFBSDB 2.0: a database for

CTCF-binding sites and genome organization. *Nucleic Acids Research*, *41*(D1):

D188-D194. https://doi.org/10.1093/nar/gks1165

Table 1: List of squamous cell carcinoma promoting and suppressor SCDEGs.

Carcinoma promoting					Carcinoma suppressor	
AKT3	CRB3	GNB4	MGST1	VEGFC	AIFM2	LAMA2
ATP6V1C2	CXCL8	GNG11	MMP9	WNT10A	BBC3	MFNG

CAMK2A	E2F1	HEY1	PDGFA	WNT10B	CD82	MGST2
CCNA1	ERBB2	IBSP	PDGFB	WNT16	CDKN1A	PARD6G
CCNA2	FGF1	IGF1	PGF	WNT2	CDKN2A	PERP
CCNB2	FGF2	IL7R	PIM1	WNT2B	CHEK1	PIK3R1
CDK1	FGFR1	ITGA5	PLCB1	WNT3A	EDNRB	RASGRP2
CDK2	FGFR2	ITGA7	PRKCI	WNT4	ESR1	RASSF5
CEBPA	FGFR3	ITGB4	PTGS2	WNT6	FLT3LG	RXRG
CKS2	FLAI-K	ITGB6	ROCK2	WNT7B	FOXO1	SESN1
COL1A1	FLT3	JUN	RRM2		GADD45G	TGFBR2
COL4A1	FLT4	JUP	SERPINE1		GSTM3	TP53
COL4A5	FOS	KIT	SPI1		ITGA8	ZBTB16
COL4A6	FZD10	LAMA4	VEGFB		ITGA9	

Table 2: Number of squamous cell carcinoma promoting and suppressor genes, which are more strongly expressed, for cells T (P_T and S_T), O (P_O and S_O), and W (P_W and S_W), and P-values of exact test of goodness-of-fit between promoting-suppressor genes of each case and the theoretical ratio, where ALL indicate the theoretical ratio for all carcinomas and SCC indicate squamous cell carcinomas classified in CancerMine (Table S4). P-value of Fisher's exact test for a 2×3 table with P_T, S_T, P_O, S_O, P_W, and S_W was obtained ~ 0.1296.

	T	O	W	ALL	SCC
Carcinoma promoting genes	$P_T = 27$	$P_O = 32$	$P_W = 31$	18994	904
Carcinoma suppressor genes	$S_T = 17$	$S_o = 8$	$S_W = 9$	11295	584
P-value (for ALL)	0.8768	0.02246	0.05194		
P-value (for SCC)	1	0.01421	0.03445		

(a)

TGTTATGCCCAAAAATTCGTGATCCCCAAAGACCACCAGGGAGCCGAGT
CCGATGCAAAAGCAAAGAGCCTTTATTCGAGCTAGCTCGAGCTCAATC
CCCTACCTGCACCGACGCAGCGGTGAGATACCAGGGAAAGAGCACGA
GTTTCAAAAAGGACAAAGGTTTTATTGGGGCCTGGGGGCAGTTGGTG
AGGTAATGGCTGTGGCCTCAGCTGATTGGCTGGGGAGGGGTCCTGGG
GAAGGGTCTGGCAGGTGAGGGAGGGTTTACTCAAGGGGAGGAGGTG
TGGTCAAGGTGAAGGACACAGAACAAGATGGCGAGGGGAGGAGGTG
TGGTCAAGGTGAAGGACACAGAACAAGATGGCGAGGGGAGGAGGTG
TGGTCAAGGTGAAGGACACAGAACAAGATGGCGAGGGGAGGAGGTG
TGGTCAAGGTGAAGGACACAGAACAAGATGGCGAGGGGAGGAGGTG
TGGTCAAGGTGAAGGACACAGAACAAGATGGCGAGGGGAGGAGGTG
TGGTCAAGGTGAAGGACACAGAACAAGATGGCGAGGGGAGGAGGTG
TGGTCAAGGTGAAGGTCACAGAACAAGATGGCGACGGCTGGCGTAGG
CCCGCCCTTTCACTTTctgcaaaatcttacttggatcctaaagctgtagtgaaaatccggttta
tctgcgcgggaacttaggtctgaaggtggaggagga

	Motif PWM	Motif Sequence	Orientation	Score
CTCF-binding motif	EMBL_M1	CGCCATCTTGTTCT	–	11.349
	EMBL_M2	GGCATAACA	–	4.0407
	EMBL_M2	CAATCAGCTGAGG CCACAGC	–	4.0407
Promoter motif	Initiator	TCACTTT	+	0.95
	DPE	ggatc	+	0.93

(b)

Figure 1: Positions of motif sequences with higher scores of CTCF-binding and core promoter activity in *FERV1* LTR in the first intron of the *KIT* gene. Detailing position and sequence of each CTCF-binding motif (cyan, green, and yellow shadowed) and each core promoter motif (magenta and red texts) (a), and illustration of inferred promoter and insulator activities of *FERV1* LTR (b). Capital letters indicated the sequence in LTR and lower case letters indicated outside of LTR.

 Figure 2: TPM of the following: entire KIT, 1^{st} exon of KIT, averages over $2^{nd} \sim 22^{nd}$
943   exon, normal KIT, and d1KIT of T, O, and W cells. Values indicate TPM and length of
944   each green bar is proportional to TPM.
945
946

947
948   **Figure 3:** Correlation coefficients between TPMs of typical SCDEGs that are involved
949   in melanoblast and melanocyte formation and TPM of normal KIT (Blue) or d1KIT
950   (Orange). P-values of correlation coefficient between normal *KIT* and *PMEL* ~ 0.20302,
951   that between *d1KIT* and *SOX10* ~ 0.01876, that between *d1KIT* and *Dkk4* ~ 0.09968,
952   and that between *d1KIT* and *TYRP1* ~ 0.12763.