Trabid patient mutations impede the axonal trafficking of adenomatous polyposis coli to disrupt neurite growth

Daniel Frank[1,7,14], Maria Bergamasco[2,7,14], Michael Mlodzianoski[3,7,14], Andrew Kueh[4,7,8,9], Ellen Tsui[5,7], Cathrine Hall[6,7], Georgios Kastrappis[10], Anne Kathrin Voss[2,7], Catriona McLean[11], Maree Faux[12], Kelly Rogers[3,7], Bang Tran[10], Elizabeth Vincan[10,13], David Komander[1,7], Grant Dewson[1,7,15], and Hoanh Tran[1,7,10,15]

[1]Ubiquitin Signalling Division, [2]Epigenetics and Development Division, [3]Centre for Dynamic Imaging, [4]Melbourne Advanced Genome Editing Centre, [5]Histology Facility, [6]Inflammation Division, Walter and Eliza Hall Institute of Medical Research, Parkville, VIC 3052, Australia
[7]Department of Medical Biology, The University of Melbourne, Parkville, VIC 3052, Australia
[8]Olivia Newton-John Cancer Research Institute, Heidelberg, VIC 3084, Australia
[9]School of Cancer Medicine, La Trobe University, Heidelberg, VIC 3084, Australia
[10]Department of Infectious Diseases, The University of Melbourne at The Peter Doherty Institute for Infection and Immunity, Melbourne, VIC 3000, Australia
[11]Department of Anatomical Pathology, The Alfred Hospital, Melbourne, VIC 3004, Australia
[12]Neuro-Oncology Group, Murdoch Children's Research Institute, Parkville, VIC 3052, Australia
[13]The Victorian Infectious Diseases Reference Laboratory, Royal Melbourne Hospital at The Peter Doherty Institute for Infection and Immunity, Melbourne, VIC 3000, Australia
[14]These authors contributed equally to this work
[15]Corresponding authors dewson@wehi.edu.au, tran.h@wehi.edu.au

Running Title: Deubiquitylating enzyme Trabid controls polarised axon elongation

Abstract

Trabid/ZRANB1 missense mutations have been identified in children diagnosed with a range of congenital disorders including reduced brain size, but how Trabid regulates neurodevelopment is not understood. We have characterised these patient mutations in cells and mice to identify a key role for Trabid in the regulation of neurite growth. One of the patient mutations flanked the catalytic cysteine of Trabid and its deubiquitylating (DUB) activity was abrogated. The second variant retained DUB activity, but failed to bind STRIPAK, a large multiprotein assembly implicated in cytoskeleton organisation and neural development. *Trabid/ZRANB1* knock-in mice harbouring either of these patient mutations exhibited reduced neuronal and glial cell densities in the brain and a motor deficit consistent with fewer dopaminergic neurons and projections. Mechanistically, both DUB-impaired and STRIPAK-binding-deficient Trabid variants impeded the trafficking of adenomatous polyposis coli (APC) to microtubule plus-ends. Consequently, the formation of neuronal growth cones and the trajectory of neurite outgrowth from mutant midbrain progenitors were severely compromised. We propose that STRIPAK recruits Trabid to deubiquitylate APC, and that in cells with mutant Trabid, APC becomes hyperubiquitylated and mislocalised causing impaired organisation of the cytoskeleton that underlie the neuronal and developmental phenotypes.

Introduction

Deubiquitylating enzymes (DUBs) are integral components of the ubiquitin system that control protein functions essential for healthy development and aging (Clague et al. 2019). DUBs catalyze the removal of ubiquitin from protein substrates to regulate protein stability, activity, interaction, or localisation (Komander et al. 2009). Ubiquitin can be assembled into polymers linked through one of eight internal ubiquitin residues – Met1, Lys6, Lys11, Lys27, Lys29, Lys33, Lys48 and Lys63 – and impairment of DUB function can lead to the accumulation of one or more of these ubiquitin chain linkages on a protein substrate with deleterious consequences. For example, LINKage-specific deubiquitylation deficiency-induced Embryonic Defects (LINKED), a human syndrome caused by loss-of-function mutations in the DUB OTUD5, is associated with increased Lys48-linked ubiquitylation and turnover of chromatin remodeling and transcription factors that are critical for normal embryonic development (Beck et al. 2021).

Consequently, LINKED syndrome patients manifest severe congenital malformations and die in
early infancy (Beck et al. 2021). OTUD5 belongs to a 17-member family of human DUBs that
possess an OTU (*o*varian *tu*mour) catalytic domain and include the closely related OTU DUBs
A20, Cezanne and Trabid. Trabid missense mutations have been identified in children diagnosed
with a range of developmental disorders including microcephaly (Deciphering Developmental
Disorders 2015).
69
Trabid has two defining biochemical features. First, it exhibits strong DUB specificity for Lys29-
and Lys33-linked ubiquitin chains, cleaving these chain types 40-fold more efficiently than
Lys63-linked chains (Licchesi et al. 2012). Lys29-linked polyubiquitin exists mainly as
heterotypic chains containing Lys48 linkages that target substrates for degradation (Kristariyanto
et al. 2015; Leto et al. 2019; Harris et al. 2021). Trabid is thought to modulate the levels of
Lys29/Lys48 mixed chains to regulate proteostasis, autophagy and cell division (Chen et al.
2021; Yu et al. 2021; Vaughan et al. 2022). Lys33-linked ubiquitin polymers have been
implicated in post-Golgi protein trafficking (Yuan et al. 2014), and Lys63-linked chains are
abundant cellular adducts with established roles in the sorting of endosomal/lysosomal cargo
(Erpapazoglou et al. 2014) and mediating the formation of protein assemblies (Tran and Polakis
2012). The second defining feature of Trabid is that it binds the Striatin-interacting phosphatase
and kinase (STRIPAK) complex (Sowa et al. 2009; Tran et al. 2013; Harris et al. 2021), a large
multiprotein assembly implicated in cytoskeleton organisation, cell migration and neural
development (Hwang and Pallas 2014; Sakuma et al. 2014; Madsen et al. 2015; Bazzi et al.
2017; Kuck et al. 2019).
85
We have previously identified the adenomatous polyposis coli (APC) protein as a candidate
Trabid substrate (Tran et al. 2008). Trabid knockdown in HEK293 cells caused APC to become
modified with polyubiquitin and aggregate in the cytoplasm, whereas high Trabid levels
correlated with *hypo*ubiquitylated APC that accumulated in the membrane protrusions of long
cell processes. These observations led us to propose that Trabid regulates the ubiquitylation and
subcellular localisation of APC (Tran et al. 2013). In neurons, APC organises the cortical
cytoskeleton to promote the formation of growth cones and the steering of growing axons that
are essential for embryonic brain development (Yokota et al. 2009; Preitner et al. 2014;

Dogterom and Koenderink 2019; Efimova et al. 2020). High expression of Trabid, APC and

STRIPAK proteins in neural stem cells (Castets et al. 2000; Blanpain et al. 2004; Yokota et al.

2009) indicate that Trabid's association with STRIPAK and its ability to regulate APC

polyubiquitylation may be important for neurogenesis. We now show that two Trabid mutants

found in children with developmental deficits are impaired in two distinct biochemical activities

that culminate in the perturbed trafficking of APC to neurite tips. We propose that the neuronal

and developmental phenotypes associated with these Trabid loss-of-function mutations are

primarily caused by the mislocalisation of APC that leads to defective cytoskeleton organisation

and aberrant cell locomotion.

Results

Trabid patient variants are impaired in DUB activity and STRIPAK binding

Trabid/ZRANB1 patient missense mutations, R438W and A451V, are linked to developmental

microcephaly (Deciphering Developmental Disorders 2015). Mapping these residues onto the

domain structure of Trabid show that they flank either side of Trabid's catalytic cysteine C443

(Fig. 1A). Residue R438 projects prominently into the catalytic cleft formed by the ankyrin

repeats and the OTU core of Trabid's catalytic domain (Licchesi et al. 2012), whereas residue

A451 resides at the back of the active site, opposite to C443 on alpha helix 4 (Fig. 1B). To

determine if patient mutations R438W and A451V influence substrate catalysis, we purified the

mutant AnkOTU domains and tested their ability to hydrolyse synthetic ubiquitin chains *in vitro*

(Licchesi et al. 2012). Wild-type Trabid AnkOTU generated appreciable amounts of mono-

ubiquitin within 15 min incubation with either Lys29- or Lys63-linked di-ubiquitin chains (Fig

1C and D). In contrast, Trabid R438W showed near total loss of DUB activity and cleaved

ubiquitin product was only readily detected after a prolonged 120 min incubation (Fig. 1C and

D). A catalytically inactive C443S mutant AnkOTU domain failed to hydrolyse ubiquitin chains

even after 2 h incubation (Fig. 1C and D). Interestingly, the Trabid A451V AnkOTU domain

cleaved Lys29- or Lys63-linked di-ubiquitin with comparable efficiency to wild-type Trabid,

despite it being slightly less stable as a recombinant protein based on thermal stability assay

(Supplementary Fig. 1A). A ubiquitin suicide probe assay revealed that R438W and A451V

Trabid AnkOTU proteins retained a functional catalytic interaction with ubiquitin

(Supplementary Fig. 1B).

In cells, the inability of the DUB-inactive Trabid C443S to cleave polyubiquitin led to stable

interaction of this mutant with polyubiquitylated substrates (Tran et al. 2008; Licchesi et al.

2012). We therefore asked whether the DUB-impaired R438W Trabid likewise exhibited

increased binding to polyubiquitin in cells. Polyubiquitin chains readily co-precipitated with both

FLAG-tagged Trabid R438W and FLAG-Trabid C443S expressed in HEK293T cells, consistent

with the compromised DUB activity of these mutants (Fig. 1E). The Trabid A451V variant did

not co-precipitate abundant polyubiquitylated substrates, consistent with it retaining full DUB

activity (Fig 1E).

We have previously identified an interaction between Trabid and STRIPAK that is important for

the deubiquitylation of substrate APC (Tran et al. 2013). Whilst A451V retained full DUB

activity, strikingly, in contrast to FLAG-Trabid wild-type, R438W and C443S, the FLAG-Trabid

A451V mutant failed to efficiently co-precipitate several STRIPAK components, including

Striatin3 and STRIP1 (Fig. 1E). All FLAG-Trabid proteins co-precipitated similar levels of the

substrate E3 ubiquitin ligase HECTD1 (Tran et al. 2013; Harris et al. 2021). Together, these

results suggest that the Trabid patient mutations produce hypomorphic variants impaired in two

distinct biochemical activities: polyubiquitin hydrolysis and STRIPAK-binding.

Decreased cell density in the brains of mice harbouring Trabid patient mutations

Given the microcephaly observed in children with Trabid mutations, and to understand the

consequence of Trabid hypomorphic variants *in vivo*, we examined knock-in mice carrying

Trabid patient mutations for evidence of brain development abnormalities (Fig. 2A). Mice

heterozygous or homozygous for the R438W or A451V mutation were viable, fertile, and born at

expected Mendelian ratios (Fig. 2B). Because germline Trabid knockout or C443S homozygous

knock-in mice exhibit perinatal lethality (Dickinson et al. 2016) (our unpublished data), these

new R438W and A451V mutant mouse strains allow us to interrogate Trabid function *in vivo*.

Homozygous mutant mice from the R438W colony weighed on average ~15% less than wild-

type littermates, while R438W heterozygous and A451V mutant mice did not exhibit significant

weight loss (Fig. 2C). Immunohistochemical analysis of brain sections from weaned littermate

mice of both mutant strains revealed normal structure and laminar organisation of the cerebral

cortex, but intriguingly a reduction in cell number was apparent in different brain regions of

R438W and A451V homozygous mice compared to wild-type littermate controls (Fig. 2D). The

number of Ctip2+ medium spiny neurons in the striatum of R438W and A451V mutant mice

were reduced compared with wild-type littermates, and homozygous mice showed a greater cell

number loss than heterozygous mice, suggesting a gene dosage effect (Fig. 2E). Moreover,

R438W and A451V mutant mice had reduced number of Olig2+ oligodendrocytes in the

forebrain and midbrain compared to wild-type littermate mice (Fig. 2F). Olig2+ cells in

homozygous brain sections were consistently strongly reduced (>30%) compared to wild-type

mice, independent of age or sex. Of a combined 16 sets of littermate mice from both R438W and

A451V colonies, reduced neuronal or glial cell numbers were conspicuous in homozygous brain

sections of 12 littermate sets (Fig. 2E, F; see also Fig. 3), indicating incomplete penetrance of the

mutant phenotype. Collectively, these results suggest that Trabid's DUB and STRIPAK-binding

activities are required to produce the correct numbers of neuronal and glial cells in the

developing brain.

Trabid mutant mice exhibit a motor deficit consistent with reduced numbers of

dopaminergic neurons and projections

Given the cell number deficits in the midbrain of Trabid mutant mice and the reported midbrain

neurodegeneration and locomotor defects in Trabid mutant *Drosophila* (Kounatidis et al. 2017),

we asked if the numbers of dopaminergic neurons that control motor function might be affected

in Trabid R438W and A451V mutant mice. Midbrain and striatal coronal sections were

immunostained for Tyrosine Hydroxylase (TH), the rate-limiting enzyme for dopamine synthesis

and a marker of dopaminergic neurons. A reduction in the numbers of TH+ neurons was

observed in the substantia nigra pars compacta (SNc) of Trabid mutant mice from both R438W

and A451V colonies, independent of age or sex (Fig. 3A). SNc neurons project axons to the

striatum to produce an extensive network of axonal terminals that communicate with striatal

neurons (Matsuda et al. 2009). Consistent with the lack of prominent TH+ neuronal processes

emanating from the SNc neurons that remain in the mutant midbrain, TH immunoreactivity was

reduced in the striatum of homozygous R438W and A451V mutant mice (Fig. 3B, C). Also in

mutant brain sections, the reduced abundance and intensity of TH+ neuronal processes were
readily apparent in cortical regions including the motor cortex and claustrum that regulate motor
responses (Fig. 3B, C). Differences in TH staining in the mutant striatum compared to wild-type
were modest or not observed in all littermates examined (Fig. 3D), indicating incomplete
penetrance of the mutant Trabid alleles.
191
Given the altered abundance of TH+ cells and projections in motor-related brain regions of
Trabid mutant mice, we evaluated the motor function of these mice using a rotarod assay.
Homozygous R438W mice showed a significantly reduced latency to fall at 20, 25, 30 and 35
RPM, compared to controls (Fig. 3E), as did homozygous A451V animals at 30 and 35 RPM
(Fig. 3E). This indicates that Trabid patient mutations impair motor coordination, consistent with
a deficiency in dopaminergic neurons. The smaller size of R438W homozygous mice compared
to wild-type littermates (Fig. 2C) could be a contributing factor to the compromised rotarod
performance of these mutants. However, given that A451V homozygous mice on average are not
smaller than wild-type littermates, the motor deficit exhibited by both R438W and A451V
homozygous mice is likely caused by the abnormal cell numbers and reduced dopaminergic
neurons in their brains (Figs. 2 and 3). Taken together, these results suggest that Trabid regulates
the brain cellular architecture and neuronal circuitry required for normal motor function.
204
Trabid patient mutants fail to efficiently limit APC ubiquitylation in cells
To understand the molecular basis for the neuronal and behavioral phenotypes of Trabid mutant
mice, we focused on APC— an established regulator of mammalian brain development (Yokota
et al. 2009; Preitner et al. 2014) and a protein that we have shown becomes strongly
ubiquitylated upon loss of Trabid or Striatin (Tran et al. 2008; Tran et al. 2013). Moreover, given
that Striatin binds directly to APC (Breitman et al. 2008), we hypothesized that the Trabid
patient mutants impaired in DUB activity or Striatin-binding would impact APC ubiquitylation
and function in cells. To test this, we generated cells with doxycycline (dox)-inducible
expression of FLAG-Trabid and examined the ubiquitylation status of endogenous APC in dox-
treated cells. The levels of ubiquitin-modified APC were strongly suppressed in cells expressing
wild-type FLAG-Trabid (Fig. 4). In contrast, induction of the FLAG-Trabid R438W mutant, like
the catalytically dead FLAG-Trabid C443S, did not repress APC ubiquitylation, and induction of

FLAG-Trabid A451V expression only partially reduced APC ubiquitylation compared to control

cells (Fig. 4). These results suggest that both DUB and STRIPAK-binding activities of Trabid

are required for efficient deubiquitylation of APC in cells.

Trabid patient mutants impede EGFP-APC transport to the leading edge of migrating cells

APC decorates the membrane protrusions at the leading edge of migrating cells (Nathke et al.

1996) and we have shown that Trabid is a key regulator of APC localisation to these cortical

structures (Tran et al. 2013). Consistent with this insight, FLAG-Trabid localised prominently

with EGFP-APC in the tips of long cell processes (Fig. 5A). Analysis of several of these tip

clusters by super resolution microscopy revealed ~30% overlap between the FLAG-Trabid and

EGFP-APC signals, indicating that a significant fraction of these proteins co-localise in cortical

cell protrusions (Fig. 5B). In sub-confluent cultures, EGFP-APC localised strongly to the

lamellipodial leading edge of the majority of cells co-transfected wild-type FLAG-Trabid (Fig.

5C, D). By contrast, in cells transfected with the FLAG-Trabid mutants R438W and A451V,

EGFP-APC formed irregular puncta at sub-cortical regions close to the leading edge and EGFP-

APC aggregates are often sequestered to puncta formed by the R438W FLAG-Trabid mutant

(Fig. 5C, D). Such sites may contain ubiquitylated substrates bound to DUB-defective Trabid

mutants, as we have previously demonstrated with Trabid C443S (Tran et al. 2008; Licchesi et

al. 2012). Consistently, FLAG-Trabid C443S formed puncta that contained EGFP-APC near the

tips of long cell processes (Fig. 5E). The Trabid substrate HECTD1 was sequestered to C443S

puncta in the cytosol, but intriguingly not to puncta residing in proximity of the leading edge

membrane (Fig. 5E). In near-confluent cultures following prolonged co-transfection of FLAG-

Trabid mutants and EGFP-APC, we observed the striking phenomena of strong EGFP-APC

aggregation on or near the plasma membrane and thin tube-like processes extending between

neighbouring cells decorated with abundant EGFP-APC puncta/aggregates of various sizes (Fig.

5F). In cells co-transfected with wild-type FLAG-Trabid, EGFP-APC concentrated in cell tips

(Fig. 5F). Collectively, these results indicate that Trabid's DUB and STRIPAK-binding activities

are required for the efficient localisation of APC to plasma membrane sites involved in polarised

cell migration.

Trabid patient mutations perturb the axonal trafficking of APC-tdTomato and impair the
trajectory of neurite outgrowth

APC regulates the dynamic interactions of the microtubule and actin cytoskeletons at cortical
membranes to direct the formation of neuronal growth cones and the trajectory of axon growth
(Dogterom and Koenderink 2019). To investigate the effect of patient Trabid mutations on APC
localisation in primary neurons, we generated mice expressing an APC-tdTomato fluorescent
protein under the control of the endogenous *Apc* gene promoter (Fig. 6A). APC-tdTomato mice
were bred with knock-in mice carrying either the Trabid R438W or A451V mutant allele (Fig. 2)
and neural progenitors were isolated at embryonic day E11.5 from compound heterozygotes. We
then performed live cell imaging to analyse the trafficking of APC-tdTomato in neurites
extending from progenitors undergoing differentiation (Fig. 6A). Endogenous Trabid protein
levels were comparable in neural progenitor cultures derived from embryos bearing wild-type or
R438W and A451V mutant Trabid alleles (Fig. 6B). In neurons with wild-type Trabid, APC-
tdTomato accumulated strongly in the tips of growing neurites, marking prominent growth cones
that drive axon elongation (Fig. 6C; *Zranb1*$^{+/+}$). In neurons with heterozygous Trabid R438W or
A451V mutation, APC-tdTomato exhibited broad distribution along the length of the neurite and
the formation of growth cones were severely compromised (Fig. 6C, *Zranb1*$^{R438W/+}$). Tracking
analysis of the turn angles of APC-tdTomato intensities revealed that neurites with wild-type
Trabid elongated in a polarised manner, whereas neurites with R438W or A451V mutant Trabid
turned back or retracted at a greater frequency (Fig. 6D and E). These data suggest that Trabid's
DUB and STRIPAK-binding activities are required for efficient transport of APC to neurite tips
to promote the formation growth cones required for polarised axon elongation.

Discussion

We describe a novel function for the deubiquitylating enzyme Trabid as a key regulator of
axonal growth and guidance that likely underpins the neurodevelopmental defects observed in
children with Trabid mutation. Our data suggests that Trabid's mechanism of action is to
suppress the ubiquitylation of APC to regulate its intracellular trafficking. A *hypo*ubiquitylated
APC pool is efficiently localised to the cortical cytoskeleton where it directs neuronal growth
cone formation and polarised axon elongation. The identification of human Trabid variants that

disrupt the distribution of APC to the leading edge of migrating cells provides a plausible

explanation for the associated patient neuro-developmental disorders. Our work identifies

Trabid's DUB- and STRIPAK-binding activities, and the control of APC localisation, as crucial

events during embryonic and neural development.

The two patients carrying Trabid missense mutations R438W and A451V were diagnosed with a

range of distinct congenital disorders including craniofacial abnormalities, seizures,

developmental delay, autism—and both patients presented with microcephaly and constipation

(Deciphering Developmental Disorders 2015). We propose that the underlying cause of these

seemingly unrelated conditions, broadly classified as neurocristopathies, is the abnormal

specification or migration of neural crest cells in the developing embryo (Vega-Lopez et al.

2018). Discrete neural crest populations contribute to the development of craniofacial structures,

the forebrain and midbrain, and the enteric nervous system (Anderson et al. 2006; Creuzet et al.

2006). Therefore, errors in the formation or polarised migration of neural crest cells could

account for all the patient phenotypes, including the reduction in brain volume (microcephaly).

Presently, we can only speculate that the irregular cellular architecture and reduced cell numbers

in the brains of mice harbouring Trabid patient mutations (Figs. 2 and 3) reflect the errant

migration of neural crest or neural progenitor cell populations in early embryogenesis (Silva et

al. 2019). While Trabid mutant mice did not exhibit microcephaly, they showed a motor deficit

(Fig. 3) consistent with the locomotor defects reported for Trabid loss-of-function in *Drosophila*

(Kounatidis et al. 2017). Intriguingly, the fly Striatin homolog CKA is involved in axonal

transport and motor coordination (Neisch et al. 2017), and *Drosophila* Strip1 regulates

endosomal trafficking and axon elongation (Sakuma et al. 2014). Endosomal trafficking defects

have been linked to reduced proliferation of neural progenitors and microcephaly (Carpentieri et

al. 2022). Three independent groups have identified Trabid's association with STRIPAK (Sowa

et al. 2009; Tran et al. 2013; Harris et al. 2021). We have now established the functional

significance of this interaction. The requirement of Trabid binding to STRIPAK for efficient

protein trafficking, polarised axon growth, and motor coordination (Figs. 3, 5, 6), supports the

view that Trabid-STRIPAK regulates an evolutionarily conserved mechanism of cell movement

required for normal brain development and establishment of the correct neuronal circuitry in the

motor system.

The impairment of two distinct Trabid functions—polyubiquitin hydrolysis and STRIPAK-

binding—led to common cellular, developmental, and behavioral phenotypes (Figs. 2-6)

(Deciphering Developmental Disorders 2015), strongly suggesting that these activities act in the

same pathway. We propose that APC is the primary molecular target of Trabid action in

cytoskeleton organisation and polarised axon growth (Fig. 7). APC is an established regulator of

cell adhesion and migration, and it governs the cortical actin and microtubule cytoskeleton

dynamics required to form and steer axonal growth cones (Dogterom and Koenderink 2019;

Efimova et al. 2020). In a yeast two hybrid screen, we have found that Trabid and Striatin

interacted with the armadillo repeat domain (ARD) of APC, but not to an APC ARD mutant that

caused cell-cell adhesion defects (Hamada and Bienz 2002; Tran et al. 2008). This implies that a

functional interaction between Trabid, Striatin and APC promotes the fidelity of cell-cell or cell-

substratum contacts. Our model of Trabid action in Figure 7 integrates published data showing

that (1) Trabid complexes with Striatin/STRIPAK in human cell lines (Sowa et al. 2009; Tran et

al. 2013; Harris et al. 2021), and (2) Striatin binds directly to the ARD domain of APC (Breitman

et al. 2008). We propose that STRIPAK recruits Trabid to deubiquitylate APC. This allows APC

to accumulate at the leading edge of migrating cells to promote efficient, polarised locomotion

(Fig. 7A). The Trabid R438W mutant still binds STRIPAK and is recruited to APC, but it is

impaired in its ability to cleave ubiquitin chains from APC. And although the Trabid A451V

mutant retains full DUB activity, it cannot be recruited to APC via STRIPAK. Both mutant

scenarios cause APC to become persistently modified with ubiquitin chains that lead to APC

delocalisation from cell tips and defective cell movement (Fig. 7B). Consistent with this model,

Trabid and/or Striatin deficiency caused APC hyperubiquitylation and aggregation, perturbed

actin assembly and microtubule stability, and inhibited the migration of mouse and human cell

lines (Bai et al. 2011; Tran et al. 2013) (our unpublished data).

We have shown that ubiquitin-modified APC correlated with its binding to Axin in the β-catenin

destruction complex (Tran and Polakis 2012), whereas non-ubiquitin-modified APC accumulates

in membrane protrusions (Tran et al. 2013). Thus, reversible modification with polyubiquitin

could be the long-hypothesised molecular switch that regulates the distribution of APC between

its many functional pools in cells (Dikovskaya et al. 2001; Bienz 2002). Chronic APC

ubiquitylation in Trabid deficient/mutant cells might result in increased APC sequestration into

Axin destruction complexes or promote spurious interactions with ubiquitin binding proteins that

retard APC trafficking along microtubules. Rescue of the APC transport defect could

hypothetically be achieved by inhibition of glycogen synthase kinase 3 (GSK3), which we have

shown abolishes APC ubiquitylation (Tran and Polakis 2012). In support of this idea is the

finding that GSK3 inactivation promotes the association of APC with microtubule plus ends to

drive polarised cell migration (Etienne-Manneville and Hall 2003).

Optimal cell migration and adhesion requires stable, acetylated microtubules (Aguilar et al.

2014; Bance et al. 2019). Of note, loss of APC and the STRIPAK proteins STRIP1 and

CTTNBP2 strongly reduced the acetylated microtubule network in neurons (Yokota et al. 2009;

Shih et al. 2014; Sakuma et al. 2015). Furthermore, Striatin depletion perturbed cell-cell

adhesion (Breitman et al. 2008; Lahav-Ariel et al. 2019) and axon elongation defects caused by

STRIP1 mutation are linked to the dysregulation of neuronal adhesion (Sakuma et al. 2014). It

would be interesting to investigate if chronic APC ubiquitylation and dysfunction underlie all

Trabid and STRIPAK mutant phenotypes.

Materials and Methods

Protein purification and characterisation

Wild-type (WT), R438W, A451V, and C443S Trabid AnkOTU domains (245-697) were

expressed in BL21 competent *E. coli* and purified as described previously (Licchesi et al. 2012)

with minor modifications. *E. coli* cells were induced with 300 mM IPTG and grown overnight at

16°C. Cells were lysed by sonication in 50 mL lysis buffer (20 mM Tris pH 8.5, 200 mM NaCl,

10% glycerol, 1 mM PMSF, 2x Roche protease inhibitor tablets, 3 mM $MgCl_2$, 0.1 mg/mL

DNAse, 150 μL lysozyme, 10 mM β-mercaptoethanol) per 2 L culture. Anion-exchange

chromatography (MonoQ 5/50) was performed using the ÄKTA pure system at 4°C. See

Supplementary Figure 1 for the characterisation of the thermal stability and ubiquitin reactivity

of purified Trabid AnkOTU proteins. For DUB assays, a total of 20 μL reaction consisting of

0.25 μM purified AnkOTU domain was incubated with 1 μM of K29, K63 di-ubiquitin chains, or

K33-linked Tetra-Ub chains (R&D Systems) in DUB reaction buffer (50 mM NaCl, 50 mM Tris

pH 7.4, and 5 mM DTT) at 37°C. Reactions were stopped at the indicated times by addition of

SDS sample buffer and visualized by SDS-PAGE and silver staining (Silver Stain Plus; Biorad).

Cells, plasmids, and antibodies

An inducible lentiviral CRISPR/Cas9 system (Aubrey et al. 2015) was used to overexpress

Trabid in Figure 4. PCR amplified DNA encoding human wild-type or mutant Trabid with an N-

terminal FLAG tag was cloned into the EcoRI/NheI sites of a doxycycline-inducible pFTREtight

MCS rtTAadvanced puro lentiviral vector (Brumatti et al. 2013). Transduced HEK293 cells were

selected for puromycin resistance and pooled clones were used for experiments. Transient

transfection of plasmids pcDNA3.1-FLAG-Trabid (Tran et al. 2013), FLAG-Trabid mutants

generated by site-directed mutagenesis (QuikChange, Agilent), pEGFP-C1-APC (Rosin-

Arbesfeld et al. 2001), and pCMV-HA-HECTD1 (Sarkar and Zohn 2012) in HEK293T cells was

performed using Lipofectamine 2000 (Invitrogen). HEK293T cells (ATCC) and HEK293 cells

(Cellbank Australia) were authenticated by STR profiling and frequently tested for mycoplasma

contamination. Cells were cultured in DMEM supplemented with 10% FBS and 2 mM L-

Glutamine in a humidified incubator at 37°C with 5% CO_2. Mouse monoclonal Trabid antibodies

and rabbit polyclonal HectD1 antibodies have been described (Tran et al. 2013). Commercial

antibodies were purchased from a variety of vendors: anti-APC ALi 12-28 (Santa Cruz), anti-

Striatin (BD Biosciences), anti-Striatin3 (SG2NA, Novus Biologicals), anti-Strip1 (Abcepta)

anti-ubiquitin P4D1 (Cell Signaling), anti-FLAG M2, anti-β-actin-HRP AC-15, anti-β-tubulin

(Sigma), anti-HA 3F10 (Roche), anti-Tyrosine Hydroxylase (Millipore, AB152), anti-Ctip2

(Abcam, ab18465), and anti-Olig2 (Abcam, ab109186).

Protein enrichment from cell lysates

Confluent cells in 10 cm dishes were lysed on ice in 1 mL lysis buffer containing 1% IGEPAL

CA-630, 50 mM Tris-HCl, pH 7.5, 120 mM NaCl, 1 mM EDTA, and protease inhibitor tablets

(Roche). Lysates were centrifuged at 14,000 rpm for 10 min to remove cell debris. The

supernatant was assayed for protein (BCA kit, Pierce), then 1 mg total protein was

immunoprecipitated with anti-FLAG M2 antibody and protein G sepharose in a total volume of 1

mL lysis buffer. Enrichment of polyubiquitin chains was performed using purified GST-TUBE

(Tandem Ubiquitin Binding Entity) protein (Hjerpe et al. 2009). A volume of cell supernatant

containing 1 mg total protein was diluted 1:1 with 4M urea then incubated with 10 µg GST-

TUBE and Glutathione sepharose 4B. Immunoprecipitations and GST-TUBE pulldowns were

performed overnight with constant rotation at 4°C. Protein complexes were washed twice with

lysis buffer and once with PBS, then eluted at 95°C for 5 min in Laemmli buffer for Western

blotting analysis.

Mice

All mouse studies complied with relevant ethical regulations and were approved by the Walter

and Eliza Hall Institute Animal Ethics Committee. The *Zranb1^{R438W}*, *Zranb1^{A451V}*, and *Apc^{tdTomato}*

knock-in mice were generated on a C57BL/6J background using CRISPR–Cas9-mediated gene

editing by the Melbourne Advanced Genome Editing Centre (MAGEC) at the Walter and Eliza

Hall Institute. To generate a R438W mutation within the *Zranb1* gene on mouse chromosome 7,

a single guide (sg) RNA of the sequence GAC TAT ATG CAC TTT GGA AC was used to

create double stranded breaks within the Zranb1 locus to stimulate homologous recombination

and an oligo donor of the sequence TAT AAA CTG GTC TTT GGA GTT GGC TAC ACG TCT

GGA CAG TAG ACT ATA TGC ACT TTG AAC TGA CTG CGG AGA TTG TTT ACT

TGA CTC AGT ACT ACA AGC TAC ATG GGG CAT TTA TGA CAA A was used to

introduce the R438W mutation. The sgRNA and donor sequence along with Cas9 mRNA were

injected into the cytoplasm of fertilized one-cell stage embryos generated from wild-type

C57BL/6J breeders. To generate the A451V mutation, a sgRNA of the sequence ACT CAG TAC

TAC AAG CTA CA and an oligo donor of the sequence ACA GTA GAC TAT ATG CAC TTT

GGA ACC GGA CTG CCG GAG ATT GTT TAC TTG ACT CAG TAC TAC AAG TCA CAT

GGG GCA TTT ATG ACA AAG ACT CGG TGC TTC GGA AAG CCC TGC ATG ACA GCC

TG CAT was used. Twenty-four hours later, two-cell stage embryos were transferred into the

uteri of pseudo-pregnant female mice. Viable offspring were genotyped by next-generation

sequencing. Targeted animals were backcrossed twice to wild-type C57BL/6J to eliminate off-

target mutations. Generation of C57BL/6J mice expressing the APC-tdTomato fusion protein

under the control of the endogenous *Apc* promoter (B6J.*Apc*^{tdTom}) were based on methods

previously described (Ng et al. 2020). Briefly, a sgRNA of the sequence AGA CGT CAC GAG

GTA AGA CC was used to create double stranded breaks within the *Apc* locus to stimulate

homologous recombination. A targeting vector containing homology arms of ~1.4 kilobases was

used to introduce the tdTomato coding sequence after the last *Apc* coding exon. Forward (ACC

TGT TCC TGT ACG GCA TG) and reverse (GCC TCC CAA AAT GAC CAG TG) primers to

detect the tdTomato sequence were used to screen viable pups for integration of the targeting

vector by PCR.

Neurite outgrowth from cultured neural progenitors

To generate mouse embryos expressing Apc-tdTomato with wild-type or mutant Trabid, mice

heterozygous for the Apc-tdTomato allele (*Apc*$^{tdTom/+}$) were outcrossed with mice heterozygous

for the Trabid R438W or A451V mutant allele (*Zranb1*$^{R438W/+}$ or *Zranb1*$^{A451V/+}$). Embryos at

E11.5 were harvested from pregnant females and ventral midbrains were dissected as previously

described (Thompson and Parish 2013). The isolated midbrain tissue was enzymatically

dissociated in Hank's Balanced Salt Solution containing 0.05% trypsin and 0.1% DNase I for 12

minutes at 37°C. Cells in the tissue were separated by mechanical dissociation, counted, then

plated in serum-free N2 medium consisting of a 1:1 mixture of Ham's F12 and Minimum

Essential Medium supplemented with 15 mM HEPES buffer, 1 mM glutamine, 6 mg/ml glucose,

1 mg/ml bovine serum albumin and N2 100X supplement. Cells were seeded at a density of

250,000 cells per well in a 24-well plate at 37°C, 3% CO_2. An aliquot of the resultant

neurosphere cultures were sequenced to ascertain genotypes. To differentiate midbrain

progenitors towards the dopaminergic lineage, neurospheres were resuspended in N2 medium

containing BDNF and GDNF (30 ng/ml each) then seeded in glass bottom ibidi chamber slides

pre-coated with poly-D-lysine and laminin (10 µg/ml each) for live imaging of neurite outgrowth

as described below.

Histology and IHC analysis

Mice were euthanized by CO_2 inhalation and intracardial perfusion was performed to fix the

tissues. Perfusion was initiated with Dulbecco's phosphate buffered saline followed by 10%

neutral buffered formalin (NBF). After perfusion, brain and tissues were dissected and post-fixed

in 10% NBF for 24h before paraffin-embedding with the Tissue-Tek VIP 6 automated tissue

processor (Sakura Finetek USA). Formalin-fixed paraffin-embedded tissues were sectioned

into 7 µm slices using a microtome and mounted on positively charged slides.

Immunohistochemistry (IHC) was performed with the Omnis Auto-immunostaining platform

using Agilent EnVision Target retrieval solution and the optimal dilution of the primary antibody

against the target protein. Secondary antibody detection was performed with the

Dako EnVision+ Single Reagents HRP and FLEX DAB+ Substrate Chromogen System. Slides

were counterstained with haematoxylin. Sections from wild-type, heterozygous and homozygous

littermate mice were mounted and stained on the same slide to ensure IHC signals can be directly

compared. Cell counts and IHC signal intensities were quantified using Fiji software (NIH). The

workflow for counting cells from IHC images is described in Supplementary Figure 2. IHC

signal intensities were measured as the optical density proportional to the concentration of the

stain. Briefly, the colour spectra of DAB- and haematoxylin-stained image was separated by

colour deconvolution, converted to greyscale, thresholding was applied, and the mean pixel

intensity of the region of interest was measured.

Rotarod performance test

The motor coordination of 3- to 4-month-old littermate mice from Trabid R438W and A451V

colonies were measured using a rotating rod (Rotamex-5, Columbus Instruments). Mice were

lowered onto a 3 cm diameter rod rotating at 12, 16, 20, 25, 30, 35, 40 or 45 revolutions per min

(RPM) for 2 min or until they fell. Animals were given a 5 min rest between RPM sessions. Two

trials were performed with a 1 h break between trials. The latency to fall was recorded for each

RPM. A 1 sec penalty was added if an animal failed to walk in time with the rod but rather

gripped the rod and rotated with it (cartwheel). Animals underwent 2 trials per day across 3 days.

Microscopy and image analysis

Confocal microscopy – Cells grown in Lab-Tek II chamber slides were fixed with 4%

paraformaldehyde for 10 min at room temperature, then permeabilized with 0.2% Triton-X100

and blocked with 5% normal goat serum (NGS). Cells were incubated with primary antibodies

diluted in 5% NGS overnight at 4°C, followed by Alexa Fluor-conjugated secondary antibodies

for 1h at room temperature. ProLong Gold antifade reagent with DAPI was used to mount

coverslips to microscope slides. Confocal images were acquired on a Zeiss LSM 880 Airyscan

microscope using a 63x/1.4 N.A. oil objective, and 405, 488 and 594nm lasers. Maximum

intensity projections of raw images comprising 10-15 *z* sections were created using Fiji.

3D structured illumination microscopy – Super-resolution three-dimensional structured

illumination microscopy (3D-SIM) was performed on the DeltaVision OMX-SR system using

a 60x/1.42 N.A. PlanApo oil immersion objective, sCMOS cameras, and 488, 568 and 640 nm

lasers. 3D-SIM images consisted of 15 raw images per focal plane per colour channel with 125

nm between each z-step. Images were reconstructed and the colour channels aligned using the

reconstruction and alignment algorithms in softWoRx 7.0. Fraction of overlap between FLAG-

Trabid and EGFP-APC were measured in Imaris 9.2.

Lattice lightsheet time-lapse imaging and analysis – Live cell time-lapse imaging of APC-

tdTomato fluorescence in growing neurites was acquired using the Zeiss Lattice Light Sheet 7

microscope. A 561 nm laser formed a light sheet of length 30 μm with a thickness of 1 μm at the

sample plane via a 13.3 x, 0.44 NA objective. tdTomato fluorescence was collected via a 44.83

X, 1 NA detection objective lens. Data was collected with a frame rate of 60 ms and a z-step of

300 nm. Each region was imaged at 15-min intervals for 4-6 hours. Fluorescence was collected

via a multi-band stop, LBF 405/488/561/633, filter. Images were subsequently deskewed using

Zeiss's Zen 3.4 software. Samples were measured at 37°C and 5% CO_2. Maximum intensity

projections were created from the deskewed data in Fiji. Neurite tips were tracked using

Trackmate (v6.0.3) and LoG detector with 12 μm diameter, 0.08 threshold, no filters, and

subpixel localisation turned on. A minimum of 5 frames were set for track inclusion. Turn angles

were calculated using Matlab R2019b. Three points were used to calculate the turn. The first two

points determine the direction of neurite growth. The angle between this direction to the third

point determines the turn angle of the neurite. The turn angle was normalized such that 0º

constitutes a step straight forward with no turn and 180º is a turn backwards.

Acknowledgements

We are indebted to Prof David L. Vaux for his generous support and mentorship in the early

stages of this study. We thank Drs S. Wilcox, S. Scutts, J. Heath, D. Newgreen, J. Vince, C.

Parish, K. Scicluna, M. Herold, V. Wimmer, T. Thomas, K. Newton and V. Dixit for discussion,

reagents and technical support; and WEHI's Bioservices staff for outstanding animal care. Part

of the funding for this study was provided by the Australian National Health and Medical

Research Council (Ideas Grant #1181580 to E.V., H.T., and B.M.T). G.D. was supported by a

fellowship from the Bodhi Education Fund. Work in the authors' laboratory is made possible by

operational infrastructure grants through the Australian Government Independent Research

Institutes Infrastructure Support (IRISS) and the Victorian State Government OIS. The

generation of all mutant and transgenic mice used in this study was supported by Phenomics

Australia and the Australian Government through the National Collaborative Research

Infrastructure Strategy (NCRIS) program.

Author Contributions

D.F. designed and performed the protein biochemistry experiments, quantified cell numbers from

histological sections, and assisted with the generation and maintenance of neurosphere cultures.

M.B. designed and performed the rotarod experiments and assisted with analysis of the rotarod

and histology data. M.M. performed 3D-SIM microscopy, live cell imaging of Apc-tdTomato

trafficking, and analysed the data. A.K. designed the targeting vectors for generation of knock-in

mice. E.T. assisted with the design of histology experiments. C.H. performed the genotyping.

G.K. assisted with statistical analysis of the rotarod data and quantified IHC intensities. A.V.,

C.M., M.F., K.R., B.T., and E.V. analysed data. D.K. analysed data and provided study feedback

and advice. G.D. supervised the study, analysed and conceptualized data, and edited the

manuscript. H.T. conceived the project, designed, performed and supervised experiments,

analysed and conceptualized data, and wrote the manuscript. All authors have read and

commented on the manuscript.

Competing Interest Statement

551 D.K. is founder, shareholder and serves on the SAB of Entact Bio.

References

Aguilar A, Becker L, Tedeschi T, Heller S, Iomini C, Nachury MV. 2014. Alpha-tubulin K40 acetylation is required for contact inhibition of proliferation and cell-substrate adhesion. *Mol Biol Cell* **25**: 1854-1866.

Anderson RB, Newgreen DF, Young HM. 2006. Neural crest and the development of the enteric nervous system. *Adv Exp Med Biol* **589**: 181-196.

Aubrey BJ, Kelly GL, Kueh AJ, Brennan MS, O'Connor L, Milla L, Wilcox S, Tai L, Strasser A, Herold MJ. 2015. An inducible lentiviral guide RNA platform enables the identification of tumor-essential genes and tumor-promoting mutations in vivo. *Cell Rep* **10**: 1422-1432.

Bai SW, Herrera-Abreu MT, Rohn JL, Racine V, Tajadura V, Suryavanshi N, Bechtel S, Wiemann S, Baum B, Ridley AJ. 2011. Identification and characterization of a set of conserved and new regulators of cytoskeletal organization, cell morphology and migration. *BMC Biol* **9**: 54.

Bance B, Seetharaman S, Leduc C, Boeda B, Etienne-Manneville S. 2019. Microtubule acetylation but not detyrosination promotes focal adhesion dynamics and astrocyte migration. *J Cell Sci* **132**.

Bazzi H, Soroka E, Alcorn HL, Anderson KV. 2017. STRIP1, a core component of STRIPAK complexes, is essential for normal mesoderm migration in the mouse embryo. *Proc Natl Acad Sci U S A* **114**: E10928-E10936.

Beck DB, Basar MA, Asmar AJ, Thompson JJ, Oda H, Uehara DT, Saida K, Pajusalu S, Talvik I, D'Souza P et al. 2021. Linkage-specific deubiquitylation by OTUD5 defines an embryonic pathway intolerant to genomic variation. *Sci Adv* **7**.

Bienz M. 2002. The subcellular destinations of APC proteins. *Nat Rev Mol Cell Biol* **3**: 328-338.

Blanpain C, Lowry WE, Geoghegan A, Polak L, Fuchs E. 2004. Self-renewal, multipotency, and the existence of two cell populations within an epithelial stem cell niche. *Cell* **118**: 635-648.

Breitman M, Zilberberg A, Caspi M, Rosin-Arbesfeld R. 2008. The armadillo repeat domain of the APC tumor suppressor protein interacts with Striatin family members. *Biochim Biophys Acta* **1783**: 1792-1802.

Brumatti G, Salmanidis M, Kok CH, Bilardi RA, Sandow JJ, Silke N, Mason K, Visser J, Jabbour AM, Glaser SP et al. 2013. HoxA9 regulated Bcl-2 expression mediates survival of

myeloid progenitors and the severity of HoxA9-dependent leukemia. *Oncotarget* **4**: 1933-1947.

Carpentieri JA, Di Cicco A, Lampic M, Andreau D, Del Maestro L, El Marjou F, Coquand L, Bahi-Buisson N, Brault JB, Baffet AD. 2022. Endosomal trafficking defects alter neural progenitor proliferation and cause microcephaly. *Nat Commun* **13**: 16.

Castets F, Rakitina T, Gaillard S, Moqrich A, Mattei MG, Monneron A. 2000. Zinedin, SG2NA, and striatin are calmodulin-binding, WD repeat proteins principally expressed in the brain. *J Biol Chem* **275**: 19970-19977.

Chen YH, Huang TY, Lin YT, Lin SY, Li WH, Hsiao HJ, Yan RL, Tang HW, Shen ZQ, Chen GC et al. 2021. VPS34 K29/K48 branched ubiquitination governed by UBE3C and TRABID regulates autophagy, proteostasis and liver metabolism. *Nat Commun* **12**: 1322.

Clague MJ, Urbe S, Komander D. 2019. Breaking the chains: deubiquitylating enzyme specificity begets function. *Nat Rev Mol Cell Biol*.

Creuzet SE, Martinez S, Le Douarin NM. 2006. The cephalic neural crest exerts a critical effect on forebrain and midbrain development. *Proc Natl Acad Sci U S A* **103**: 14033-14038.

Deciphering Developmental Disorders S. 2015. Large-scale discovery of novel genetic causes of developmental disorders. *Nature* **519**: 223-228.

Dickinson ME, Flenniken AM, Ji X, Teboul L, Wong MD, White JK, Meehan TF, Weninger WJ, Westerberg H, Adissu H et al. 2016. High-throughput discovery of novel developmental phenotypes. *Nature* **537**: 508-514.

Dikovskaya D, Zumbrunn J, Penman GA, Nathke IS. 2001. The adenomatous polyposis coli protein: in the limelight out at the edge. *Trends Cell Biol* **11**: 378-384.

Dogterom M, Koenderink GH. 2019. Actin-microtubule crosstalk in cell biology. *Nat Rev Mol Cell Biol* **20**: 38-54.

Efimova N, Yang C, Chia JX, Li N, Lengner CJ, Neufeld KL, Svitkina TM. 2020. Branched actin networks are assembled on microtubules by adenomatous polyposis coli for targeted membrane protrusion. *J Cell Biol* **219**.

Erpapazoglou Z, Walker O, Haguenauer-Tsapis R. 2014. Versatile roles of k63-linked ubiquitin chains in trafficking. *Cells* **3**: 1027-1088.

Etienne-Manneville S, Hall A. 2003. Cdc42 regulates GSK-3beta and adenomatous polyposis coli to control cell polarity. *Nature* **421**: 753-756.

618   Hamada F, Bienz M. 2002. A Drosophila APC tumour suppressor homologue functions in cellular
619   adhesion. *Nat Cell Biol* **4**: 208-213.

620   Harris LD, Le Pen J, Scholz N, Mieszczanek J, Vaughan N, Davis S, Berridge G, Kessler BM,
621   Bienz M, Licchesi JDF. 2021. The deubiquitinase TRABID stabilizes the K29/K48-
622   specific E3 ubiquitin ligase HECTD1. *J Biol Chem* **296**: 100246.

623   Hjerpe R, Aillet F, Lopitz-Otsoa F, Lang V, England P, Rodriguez MS. 2009. Efficient protection
624   and isolation of ubiquitylated proteins using tandem ubiquitin-binding entities. *EMBO Rep*
625   **10**: 1250-1258.

626   Hwang J, Pallas DC. 2014. STRIPAK complexes: structure, biological function, and involvement
627   in human diseases. *Int J Biochem Cell Biol* **47**: 118-148.

628   Komander D, Clague MJ, Urbe S. 2009. Breaking the chains: structure and function of the
629   deubiquitinases. *Nat Rev Mol Cell Biol* **10**: 550-563.

630   Kounatidis I, Chtarbanova S, Cao Y, Hayne M, Jayanth D, Ganetzky B, Ligoxygakis P. 2017. NF-
631   kappaB Immunity in the Brain Determines Fly Lifespan in Healthy Aging and Age-Related
632   Neurodegeneration. *Cell Rep* **19**: 836-848.

633   Kristariyanto YA, Abdul Rehman SA, Campbell DG, Morrice NA, Johnson C, Toth R, Kulathu
634   Y. 2015. K29-selective ubiquitin binding domain reveals structural basis of specificity and
635   heterotypic nature of k29 polyubiquitin. *Mol Cell* **58**: 83-94.

636   Kuck U, Radchenko D, Teichert I. 2019. STRIPAK, a highly conserved signaling complex,
637   controls multiple eukaryotic cellular and developmental processes and is linked with
638   human diseases. *Biol Chem* **400**: 1005-1022.

639   Lahav-Ariel L, Caspi M, Nadar-Ponniah PT, Zelikson N, Hofmann I, Hanson KK, Franke WW,
640   Sklan EH, Avraham KB, Rosin-Arbesfeld R. 2019. Striatin is a novel modulator of cell
641   adhesion. *FASEB J* **33**: 4729-4740.

642   Leto DE, Morgens DW, Zhang L, Walczak CP, Elias JE, Bassik MC, Kopito RR. 2019. Genome-
643   wide CRISPR Analysis Identifies Substrate-Specific Conjugation Modules in ER-
644   Associated Degradation. *Mol Cell* **73**: 377-389 e311.

645   Licchesi JD, Mieszczanek J, Mevissen TE, Rutherford TJ, Akutsu M, Virdee S, El Oualid F, Chin
646   JW, Ovaa H, Bienz M et al. 2012. An ankyrin-repeat ubiquitin-binding domain determines
647   TRABID's specificity for atypical ubiquitin chains. *Nature structural & molecular biology*
648   **19**: 62-71.

Madsen CD, Hooper S, Tozluoglu M, Bruckbauer A, Fletcher G, Erler JT, Bates PA, Thompson
B, Sahai E. 2015. STRIPAK components determine mode of cancer cell migration and
metastasis. *Nat Cell Biol* **17**: 68-80.

Matsuda W, Furuta T, Nakamura KC, Hioki H, Fujiyama F, Arai R, Kaneko T. 2009. Single
nigrostriatal dopaminergic neurons form widely spread and highly dense axonal
arborizations in the neostriatum. *J Neurosci* **29**: 444-453.

Nathke IS, Adams CL, Polakis P, Sellin JH, Nelson WJ. 1996. The adenomatous polyposis coli
tumor suppressor protein localizes to plasma membrane sites involved in active cell
migration. *J Cell Biol* **134**: 165-179.

Neisch AL, Neufeld TP, Hays TS. 2017. A STRIPAK complex mediates axonal transport of
autophagosomes and dense core vesicles through PP2A regulation. *J Cell Biol* **216**: 441-
461.

Ng SS, De Labastida Rivera F, Yan J, Corvino D, Das I, Zhang P, Kuns R, Chauhan SB, Hou J,
Li XY et al. 2020. The NK cell granule protein NKG7 regulates cytotoxic granule
exocytosis and inflammation. *Nat Immunol* **21**: 1205-1218.

Preitner N, Quan J, Nowakowski DW, Hancock ML, Shi J, Tcherkezian J, Young-Pearse TL,
Flanagan JG. 2014. APC is an RNA-binding protein, and its interactome provides a link to
neural development and microtubule assembly. *Cell* **158**: 368-382.

Rosin-Arbesfeld R, Ihrke G, Bienz M. 2001. Actin-dependent membrane association of the APC
tumour suppressor in polarized mammalian epithelial cells. *EMBO J* **20**: 5929-5939.

Sakuma C, Kawauchi T, Haraguchi S, Shikanai M, Yamaguchi Y, Gelfand VI, Luo L, Miura M,
Chihara T. 2014. Drosophila Strip serves as a platform for early endosome organization
during axon elongation. *Nat Commun* **5**: 5180.

Sakuma C, Okumura M, Umehara T, Miura M, Chihara T. 2015. A STRIPAK component Strip
regulates neuronal morphogenesis by affecting microtubule stability. *Sci Rep* **5**: 17769.

Sarkar AA, Zohn IE. 2012. Hectd1 regulates intracellular localization and secretion of Hsp90 to
control cellular behavior of the cranial mesenchyme. *J Cell Biol* **196**: 789-800.

Shih PY, Lee SP, Chen YK, Hsueh YP. 2014. Cortactin-binding protein 2 increases microtubule
stability and regulates dendritic arborization. *J Cell Sci* **127**: 3521-3534.

Silva CG, Peyre E, Nguyen L. 2019. Cell migration promotes dynamic cellular interactions to
control cerebral cortex morphogenesis. *Nat Rev Neurosci* **20**: 318-329.

680 Sowa ME, Bennett EJ, Gygi SP, Harper JW. 2009. Defining the human deubiquitinating enzyme
681   interaction landscape. *Cell* **138**: 389-403.

682 Thompson LH, Parish CL. 2013. Transplantation of fetal midbrain dopamine progenitors into a
683   rodent model of Parkinson's disease. *Methods Mol Biol* **1059**: 169-180.

684 Tran H, Bustos D, Yeh R, Rubinfeld B, Lam C, Shriver S, Zilberleyb I, Lee MW, Phu L, Sarkar
685   AA et al. 2013. HectD1 E3 ligase modifies adenomatous polyposis coli (APC) with
686   polyubiquitin to promote the APC-axin interaction. *J Biol Chem* **288**: 3753-3767.

687 Tran H, Hamada F, Schwarz-Romond T, Bienz M. 2008. Trabid, a new positive regulator of Wnt-
688   induced transcription with preference for binding and cleaving K63-linked ubiquitin
689   chains. *Genes Dev* **22**: 528-542.

690 Tran H, Polakis P. 2012. Reversible modification of adenomatous polyposis coli (APC) with K63-
691   linked polyubiquitin regulates the assembly and activity of the beta-catenin destruction
692   complex. *J Biol Chem* **287**: 28552-28563.

693 Vaughan N, Scholz N, Lindon C, Licchesi JDF. 2022. The E3 ubiquitin ligase HECTD1
694   contributes to cell proliferation through an effect on mitosis. *Sci Rep* **12**: 13160.

695 Vega-Lopez GA, Cerrizuela S, Tribulo C, Aybar MJ. 2018. Neurocristopathies: New insights 150
696   years after the neural crest discovery. *Dev Biol* **444 Suppl 1**: S110-S143.

697 Yokota Y, Kim WY, Chen Y, Wang X, Stanco A, Komuro Y, Snider W, Anton ES. 2009. The
698   adenomatous polyposis coli protein is an essential regulator of radial glial polarity and
699   construction of the cerebral cortex. *Neuron* **61**: 42-56.

700 Yu Y, Zheng Q, Erramilli SK, Pan M, Park S, Xie Y, Li J, Fei J, Kossiakoff AA, Liu L et al. 2021.
701   K29-linked ubiquitin signaling regulates proteotoxic stress response and cell cycle. *Nat*
702   *Chem Biol* **17**: 896-905.

703 Yuan WC, Lee YR, Lin SY, Chang LY, Tan YP, Hung CC, Kuo JC, Liu CH, Lin MY, Xu M et
704   al. 2014. K33-Linked Polyubiquitination of Coronin 7 by Cul3-KLHL20 Ubiquitin E3
705   Ligase Regulates Protein Trafficking. *Mol Cell* **54**: 586-600.

 Figures

Figure 1

Figure 1. Trabid patient mutants are impaired in DUB activity and STRIPAK binding.

A. Trabid patient mutations R438W and A451V flank either side of the catalytic cysteine C443

in Trabid's AnkOTU domain.

B. Residue R438 project prominently into the catalytic cleft formed by the Ank and OTU

domain, whereas residue A451 reside at the back of the active site. The positions of the catalytic

triad residues C443, H585 and D404 are shown. Crystal structure of AnkOTU domain: PDB

3ZRH.

C and **D**. *In vitro* DUB assays. Hydrolysis of Lys29-linked (C) and Lys63-linked (D) di-

ubiquitin chains by purified wild-type and mutant Trabid AnkOTU proteins. The Trabid R438W

AnkOTU domain exhibit strongly reduced DUB activity M, molecular weight markers in

kilodaltons.

E. Immunoprecipitation of FLAG-Trabid from the lysates of transfected HEK293T cells showed

that FLAG-Trabid R438W and C443S mutants co-precipitated high amounts of endogenous

polyubiquitin (PolyUb(n) smear), indicating impaired DUB activity. FLAG-Trabid A451V failed

to efficiently co-precipitate Striatin3 and STRIP1, indicating loss of binding to the STRIPAK

complex. All FLAG-Trabid proteins co-precipitated similar amounts of the E3 ubiquitin ligase

HECTD1. EV, empty vector control.

Figure 2

Figure 2. Decreased cell density in the brains of mice harbouring Trabid patient mutations.

A. Schematic of knock-in mouse strains harbouring *Zranb1/Trabid* patient mutations generated by CRISPR/Cas9 genome editing.

B. Numbers of mice of all genotypes produced from heterozygous intercrosses in the R438W or A451V colony, respectively.

C. Normalized weight of 10 sets of weaned littermate mice (5 males, 5 females; age range P26-P222) from each Trabid mutant colony. For each set of littermates, the weight of the wild-type (+/+) mouse was set at 1 and the weight of KI/+ or KI/KI mutant mice expressed as a ratio of the wild-type mouse weight.

D. Coronal brain sections (Bregma ±0.3mm) of weaned (postnatal day P26) littermate males from the R438W and A451V mouse colonies were stained with Cresyl Violet to assess general anatomy and cellular organisation in the cerebral cortex and striatum. LV, lateral ventricle; SVZ, subventricular zone. An overall decrease in Cresyl Violet staining intensity was evident in mutant homozygous (KI/KI) sections from both mouse strains compared to similar sections from the respective wild-type (+/+) littermate.

E. Ctip2 IHC revealed reduced numbers of medium spiny neurons in the striatum of mutant mice from the R438W and A451V colonies compared to wild-type littermate mice. Representative images shown are coronal brain sections of P26 male littermates from the R438W colony. Each data point represents the cell count in a randomly selected, non-overlapping 1 mm square area in the striatum from both brain hemispheres. The age and brain region of the cell count for 3 sets of littermate mice from the indicated mutant strain are specified. Cell numbers were quantified blinded to genotype using images processed in Fiji software. See Supplementary Figure 2.

F. Olig2 IHC revealed reduced numbers of oligodendrocytes in the striatum and midbrain of homozygous mice from the R438W and A451V colonies compared to respective wild-type littermates. Representative images shown are coronal sections from the ventral midbrain of P301 female littermates from the A451V colony (AQ, cerebral aqueduct; PAQ, periaqueductal gray). Each data point represents the cell count in a randomly selected, non-overlapping 1 mm square area in the striatum or midbrain from both brain hemispheres. The age and brain region of the cell count for 3 sets of littermate mice from the indicated mutant strain are specified. Cell numbers were quantified blinded to genotype using images processed in Fiji. See Supplementary Figure 2.

Figure 3

Figure 3. Trabid mutant mice exhibit a motor deficit consistent with reduced numbers of dopaminergic neurons and projections.

A. Tyrosine Hydroxylase (TH) IHC of coronal midbrain sections revealed reduced numbers of TH+ dopaminergic neurons in the substantia nigra pars compacta (SNc) of homozygous mutant mice from the Trabid R438W and A451V colonies compared to similar midbrain sections of the respective wild-type littermates. Representative images shown are midbrain sections of P26 female littermates from the R438W colony. VTA, ventral tegmental area. Numbers of TH+ SNc neurons were quantified using IHC images processed in Fiji. See Supplementary Figure 2.

B. Tyrosine Hydroxylase IHC of coronal brain sections (Bregma ±0.3mm) of P26 littermate mice from the Trabid R438W colony revealed reduced TH staining intensity in the striatum and fewer TH+ projections in the adjacent cortex of homozygous mutant mice compared to similar regions in the wild-type littermate (magnified area).

C. Tyrosine Hydroxylase IHC of coronal brain sections (Bregma ±0.3mm) of P133 littermate mice from the Trabid A451V colony revealed reduced abundance and intensity of TH+ projections in several cortical regions including the primary/secondary motor cortex of homozygous mutant mice compared to similar regions in the wild-type littermate (magnified area).

D. Tyrosine Hydroxylase staining intensity in the striatum of mutant mice from the R438W and A451V colonies normalized to the staining intensity of wild-type littermate sections. IHC signals were measured as the optical density of the region of interest demarcated manually using Fiji. Each symbol represents one mouse of the indicated genotype belonging to a set of littermate mice from the R438W colony (n=4 sets: P26 males, P93 females, P307 males, P480 females) or the A451V colony (n=5 sets: P26 males, P41 females, P133 males, P188 females, P303 females).

E. Rotarod performance of 3 to 4-month-old littermate mice from the R438W and A451V colonies. Each data point represents the average latencies of 8 mice (4 males, 4 females), where each mouse was subjected to 6 trials over 3 days. Wild-type (+/+), heterozygous (KI/+) and homozygous (KI/KI) littermate mice were tested together. The experimenter was blinded to genotype. A repeated measures two-way ANOVA and Dunnett's multiple comparisons test was applied to the data, using wild-type mice as the control group. Asterisks denote statistically significant differences between wild-type (+/+) and homozygous (KI/KI) mutant mice. *$p<0.05$, **$p<0.01$, ***$p<0.001$. Error bars, ±SEM.

Figure 4

Figure 4. Trabid patient mutants fail to efficiently suppress APC ubiquitylation in cells.
HEK293 cells stably transfected with doxycycline inducible constructs for the expression of
wild-type or mutant FLAG-Trabid were untreated (-) or treated with dox (+; 100 ng/ml) for
24h. Cell lysates were subjected to a GST-TUBE pulldown to enrich the ubiquitylated proteome.
Precipitated material and input lysates were processed for Western blotting using the indicated
antibodies. Endogenous APC protein levels remained unchanged irrespective of FLAG-Trabid
expression (input lysates), but ubiquitylated APC species were only efficiently suppressed by
wild-type FLAG-Trabid expression ($APC^{Ub(n)}$; WT +). EV, empty vector control.

Figure 5

Figure 5. Trabid patient mutants impede the transport of EGFP-APC to the leading edge

of migrating cells.

A. Combined differential interference contrast and immunofluorescence images of HEK293T

cells extending long cell processes that contain FLAG-Trabid and EGFP-APC in the cortical

protrusions of co-transfected cells (arrowhead in merged image). Scale bar, 10 μm.

B. 3D-SIM super resolution microscopy analysis of HEK293T cell tip clusters (n=5) revealed

~30% overlap of FLAG-Trabid and EGFP-APC signals at resolutions of ~100 nm in *xy* and 320

nm in *z*. Scale bar, 100 nm.

C. EGFP-APC membrane clusters at the leading edge of migrating HEK293T cells (green

arrowheads) are readily detected with FLAG-Trabid wild-type co-expression. In cells expressing

FLAG-Trabid patient mutants, EGFP-APC aggregate in punctate structures near the leading edge

(white arrowheads), that are often sequestered together with the FLAG-Trabid R438W mutant

(yellow arrowheads). Scale bar, 10 μm.

D. Quantification of strong EGFP-APC clustering at or away from the leading edge membrane of

co-transfected HEK293T cells, as shown in C. Numbers of cells analysed from 3 independent

transfections are shown.

E. In co-transfected HEK293T cells, DUB-dead FLAG-Trabid C443S form puncta that

sequesters EGFP-APC near the tips of long cell protrusions or sequesters HA-HECTD1 in the

cytoplasm (yellow arrowheads). Notably, HA-HECTD1 was not detected in FLAG-Trabid

C443S puncta that form near the leading edge membrane of migrating cells (red arrowheads).

Scale bar, 10 μm.

F. In near-confluent HEK293T cells 48h post transfection, FLAG-Trabid patient mutants induce

long thin tubes that connect neigbouring cells and extensive EGFP-APC aggregations are

conspicuous along these tubular structures. This phenomenon was not observed in cells

transfected with wild-type FLAG-Trabid. Scale bar, 10 μm.

Figure 6

Figure 6. Trabid patient mutations perturb the trafficking of APC-tdTomato and the trajectory of neurite outgrowth.

A. Schematic of outcross between knock-in mice harbouring *Zranb1/Trabid* patient mutations and *Apc^{tdTomato}* mice, where the APC-tdTomato fluorescent protein is expressed under the control of the endogenous *Apc* gene promoter. Midbrain neural progenitors derived from E11.5 embryos of these outcrosses were used for live cell imaging of APC-tdTomato trafficking in growing neurites.

B. Western blot analysis of Trabid protein expression in E11.5 midbrain neural progenitors derived from heterozygous intercrosses in the *Zranb1/Trabid* R438W or A451V mouse colonies. Mutant Trabid protein levels are expressed at comparable levels to wild-type Trabid protein (compare KI/KI to +/+).

C. Live cell imaging of APC-tdTomato in neurites extending from E11.5 midbrain neural progenitors of a *Zranb1$^{R438W/+}$;Apc$^{tdTom/+}$* outcross. The frames shown are from a 4h time lapse portraying the movement of APC-tdTomato. APC-tdTomato accumulated in the tips of neurites with wild-type Trabid (*Zranb1$^{+/+}$*) to generate growth cones that drive axon elongation (white arrows). In neurons with mutant Trabid (*Zranb1*$^{R438W/+}$; *Zranb1*$^{A451V/+}$ data not shown), APC-tdTomato intensities were broadly distributed along neurites (purple arrowheads) and the formation of growth cones was abrogated. Scale bar, 10 μm.

D and **E**. The movement of Apc-tdTomato in individual neurite tips was tracked to visualize growth trajectory. Neurites extending from E11.5 midbrain neural progenitors of *Zranb1$^{R438W/+}$;Apc$^{tdTom/+}$* (D) or *Zranb1$^{A451V/+}$;Apc$^{tdTom/+}$* (E) outcrosses were analysed. Each data point represents the location of APC-tdTomato at 15-min intervals of a 4h time lapse. APC-tdTomato fluorescence synchronised with the movement of a membrane dye during neurite extension/retraction (data not shown) and was therefore used as a proxy for neurite growth along an arbitrary X and Y plane. Neurite tracking data were compiled from the imaging of neurite outgrowth in 3 independent neurosphere cultures established from E11.5 embryos of 3 independent *Zranb1$^{R438W/+}$;Apc$^{tdTom/+}$* or *Zranb1$^{A451V/+}$;Apc$^{tdTom/+}$* outcrosses. All image acquisition, processing and analyses were performed blinded to genotype.

Figure 7

A

B

885
886
Figure 7. Model of Trabid's mechanism of action in axon growth and guidance.

A. STRIPAK recruits Trabid to deubiquitylate APC (and possibly other substrates). This

promotes efficient APC accumulation at microtubule plus ends to coordinate the actin and

microtubule cytoskeleton dynamics that drive directional cell migration and polarised axon

growth.

B. In cells expressing Trabid mutants that are DUB-impaired (R438W) or STRIPAK-binding-

deficient (A451V), APC becomes persistently modified with ubiquitin chains that retard its

transport to cell tips. This impairs cytoskeleton organisation which leads to defective axon

elongation and cell migration. Future studies will aim to identify the mechanism of Trabid

recruitment to STRIPAK, the ubiquitin-modified residue(s) on APC (and possibly other

substrates), and the composition of polyubiquitin linkages on substrates including the Lys29- and

Lys33-linked atypical chain types preferentially targeted by Trabid.