Structural insights into complex I deficiency and assembly from the disease-related *ndufs4*[-/-] mouse

Zhan Yin[†], Ahmed-Noor A. Agip[‡], Hannah R. Bridges[*], and Judy Hirst[*]

The Medical Research Council Mitochondrial Biology Unit, University of Cambridge, Keith Peters Building, Cambridge Biomedical Campus, Cambridge, United Kingdom

† Current address: Department of Biochemistry, University of Cambridge, Tennis Court Road, Cambridge, CB2 1GA, UK

‡ Current Address: Max-Planck-Institute of Biophysics, Frankfurt 60438, Germany

* e-mail: hrb@mrc-mbu.cam.ac.uk; jh480@cam.ac.uk

Character count: 56,796 not including spaces

Abstract

Respiratory complex I (NADH:ubiquinone oxidoreductase) is essential for cellular energy production and NAD^+ homeostasis. Complex I mutations cause neuromuscular, mitochondrial diseases, such as Leigh Syndrome, but their molecular-level consequences remain poorly understood. Here, we use a popular complex I-linked mitochondrial disease model, the $ndufs4^{-/-}$ mouse, to define the structural, biochemical and functional consequences of the absence of subunit NDUFS4. Cryo-EM analyses of mouse-heart $ndufs4^{-/-}$ complex I revealed a loose association of the NADH-dehydrogenase module, and discrete classes containing either assembly factor NDUFAF2 or subunit NDUFS6. Subunit NDUFA12 (that replaces its paralogue NDUFAF2 in mature complex I) is absent from all classes, compounding the deletion of NDUFS4 and preventing maturation of an NDUFS4-free but otherwise complete enzyme. We propose NDUFAF2 as the recruiter of the NADH-dehydrogenase module during assembly of the complex. Our results provide new molecular-level understanding of the $ndufs4^{-/-}$ mouse model and complex I-linked mitochondrial disease.

Introduction

Mammalian complex I (NADH:ubiquinone oxidoreductase) is central to cellular metabolism, with key roles in NAD^+ homeostasis, respiration and oxidative phosphorylation. It couples NADH oxidation in the mitochondrial matrix to ubiquinone reduction in the inner membrane, reoxidizing the NAD^+ pool to sustain metabolism, and captures the energy from the redox reaction to transfer protons across the inner membrane and create the proton-motive force that drives ATP synthesis. Together with the substantial contribution of complex I to cellular reactive oxygen species production, these key metabolic roles make complex I dysfunctions some of the most frequent causes of primary mitochondrial diseases, including Leber's hereditary optic neuropathy (LHON) and Leigh syndrome (Koene *et al*, 2012; Fiedorczuk & Sazanov, 2018; Padavannil *et al*, 2022; Fassone & Rahman, 2012).

Complex I is a 'boot-shaped' enzyme, with a hydrophilic domain in the mitochondrial matrix that contains the cofactors for redox catalysis, and a membrane-bound domain that transports the protons. Mammalian complex I comprises 45 subunits: 14 conserved core subunits that are essential for catalysis (seven in the hydrophilic domain and seven in the membrane domain) and 31 supernumerary subunits (Brandt, 2006; Zhu *et al*, 2016; Padavannil *et al*, 2022; Hirst *et al*, 2003). Single-particle electron cryo-microscopy (cryo-EM) has

revolutionized our structural knowledge of complex I, and high-resolution structures have now been determined from several mammalian species [*Bos taurus* (cow, bovine complex I), *Ovis aries* (sheep, ovine complex I), *Mus musculus* (mouse, murine complex I) and *Sus scrofa* (pig, porcine complex I)] (Chung *et al*, 2022b; Gu *et al*, 2022; Kravchuk *et al*, 2022; Bridges *et al*, 2023) as well as from the complex I-model eukaryote, the yeast *Yarrowia lipolytica* (Parey *et al*, 2019). NADH is oxidized by a flavin mononucleotide in subunit NDUFV1 at the top of the hydrophilic domain, within a subdomain called the N-module (subunits NDUFV1, NUDUFV2, NDUFS1, NDUFA2 and NDUFV3). Then, electrons are transferred along a chain of seven iron-sulphur (FeS) clusters to bound ubiquinone, which enters 'head first' from the membrane and transits along a long, narrow channel to approach the final cluster in the chain (cluster N2). The mechanism by which ubiquinone reduction then drives proton transfer across the membrane at distant sites remains controversial (Kampjut & Sazanov, 2022; Chung *et al*, 2022a).

NDUFS4 (also known as 18 kDa, ND-18, AQDQ or NUYM) is a supernumerary subunit in the hydrophilic domain of complex I, located where the N-module is bound to the rest of the complex (Figure 1A-B). In the mammalian enzyme, it comprises N- and C-terminal coils flanking a central 4-strand β-sheet and α-helix. The N-terminus reaches down to NDUFA6 on the lower section of the hydrophilic domain (also known as the Q-module), while the C-terminus runs up over the N-module, and the central β-sheet and α-helix domain is sandwiched between the N-module (NDUFS1) and the Q-module (predominantly NDUFS3) (Figure 1B), anchoring them together. In *Escherichia coli* complex I, which lacks NDUFS4, the region occupied by the central domain is occupied instead by a C-terminal extension of NDUFS1 (Padavannil *et al*, 2022; Kolata & Efremov, 2021). These elements of NDUFS4 are all on the front of the hydrophilic domain (we use the boot analogy to identify different parts of the enzyme), while the β2-β3 loop runs through the domain, under NDUFS1, to form interactions with subunits located on the heel, including NDUFS6 and NDUFA12 (Figure 1C). Thus, with structural elements interacting with both the N-module and the Q-module, a central domain in the 'crevice' between them, and a loop inserted through the enzyme to act like a rivet, NDUFS4 is key for complex I stability. In this location, NDUFS4 is within 12 Å of five iron-sulphur clusters (Figure 1D). Finally, mammalian NDUFS4 contains a canonical protein kinase A site at Ser131, near its C-terminus, which has been proposed to be important for both catalysis and homeostasis (Signorile *et al*, 2002; Rasmo *et al*, 2010), including resistance of the N-module to degradation in mouse-brain astroglial cells (Jimenez-Blasco *et*

al, 2020). However, phosphorylation of this buried residue has not yet been observed in any
structure.

Mammalian complex I is assembled from a set of assembly intermediates, comprising
different segments of the enzyme, that are built stepwise from subunits and assembly factors
then joined together to form the mature complex as the assembly factors are released
(Guerrero-Castillo *et al*, 2017; Stroud *et al*, 2016). In the model of Nijtmans and coworkers
(Guerrero-Castillo *et al*, 2017), the N-module intermediate (core subunits NDUFV1,
NDUFV2, NDUFS1 and supernumerary subunit NDUFA2) is added to the nascent complex
in the final stage of assembly, along with four supernumerary subunits (NDUFS4, NDUFS6,
NDUFA12 and NDUFA7) that are the focus of the work presented here, and supernumerary
subunits NDUFV3 (N-module), NDUFA6 and NDUFAB1 (Q-module), and NDUFA11
(proximal membrane domain). Subunits NDUFS4, NDUFS6, NDUFA12 all form contacts
between the N-module and the Q-module, and are all conserved in complex I from the α-
proteobacterium *Paracoccus denitrificans* (Yip *et al*, 2011), an ancient relative of the
mitochondrion. As the N-module is added to the nascent complex, assembly factor
NDUFAF2 (also known as B17.2L) is released and replaced by its paralogous subunit
NDUFA12 (also known as B17.2) (Lazarou *et al*, 2007).

Many clinical studies have identified *ndufs4* mutations in severe paediatric neurological
disorders such as Leigh or Leigh-like Syndrome, with symptoms including hypotonia,
abnormal ocular movement and visual impairment, psychomotor arrest or regression and
respiratory failure (Budde *et al*, 2003; Ortigoza-Escobar *et al*, 2016; van de Wal *et al*, 2022;
Koene *et al*, 2012). Analyses of patient fibroblasts and muscle biopsies uncovered substantial
loss of complex I activity, along with characteristic features in blue-native PAGE (BN-PAGE)
analyses: lower abundance of intact complex I plus appearance of an ~830 kDa subcomplex
that lacks in-gel NADH-oxidation activity, indicating the absence of the N-module (Scacco *et*
al, 2003; Assouline *et al*, 2012; Ugalde, 2004). Underlining their functional relationship,
mutations in *ndufs6*, *ndufa12* and *ndufaf2* have all also been detected in cases of Leigh
Syndrome (Spiegel *et al*, 2009; Ostergaard *et al*, 2011; Hoefs *et al*, 2009; Kahlhöfer *et al*,
2021).

The *ndufs4* gene is highly conserved between humans and mice, and most known pathogenic
mutations lead to absence of the whole NDUFS4 protein (van de Wal *et al*, 2022), so *ndufs4*[-/-]
mice have proved a powerful and representative model for studying NDUFS4-linked

complex I deficiency, mechanisms of pathogenicity, and intervention testing (Kruse *et al*,
2008; Breuer *et al*, 2013; van de Wal *et al*, 2022). Both whole-body and neuron-specific
ndufs4[-/-] mice develop early-onset Leigh-like phenotypes (van de Wal *et al*, 2022), and
complex I activity and stability decreases substantially in all the tissues studied in whole-
body *ndufs4*[-/-] mice (Calvaruso *et al*, 2012). Intriguingly, *ndufs4*[-/-] mice exposed to prolonged
hypoxia show marked improvements in their condition, and extended lifespan compared to
mice kept under normoxia (Jain *et al*, 2016). Detailed proteomic studies in *ndufs4*[-/-] mice
observed the expected ~830 kDa subcomplex in BN-PAGE analyses, substantially lower
complex I subunit levels in brain than in liver, heart, kidney and skeletal muscle, the near-
complete absence of NDUFA12, and increased complex I assembly factors, most notably
NDUFAF2; matching results were obtained from *ndufs4*[−/−] mouse embryonic fibroblasts and
human patient cells, and it was proposed that absence of NDUFS4 induces absence of
NDUFA12, and that NDUFAF2 stabilizes the *ndufs4*[-/-] enzyme (Adjobo-Hermans *et al*,
2020).

Here, we use single-particle cryo-EM to determine the structure of complex I from *ndufs4*[-/-]
mouse-heart mitochondria, building on our previous structures of the *wild-type* enzyme (Agip
et al, 2018; Bridges *et al*, 2020) and the ND6-P25L variant (Yin *et al*, 2021). The ND6-P25L
mouse is also a complex I-linked mitochondrial disease variant (though with milder
phenotype than *ndufs4*[-/-]) that acts through a subtle 'deactivating' change in the membrane
domain to produce an enzyme incapable of reverse electron transfer. In the absence of
substrates, mammalian complex I converts gradually from an 'active' ready-to-go resting
state to the 'deactive' pronounced resting state, which requires substrates to reactivate it
(Kotlyar & Vinogradov, 1990; Chung *et al*, 2022a). Deactivation also occurs in the *wild-type*
mouse enzyme, but much more slowly, whereas the *ndufs4*[-/-] complex is observed completely
in the active state. It is also incompletely assembled, its N-module is not stably attached, and
its composition is heterogenous. By considering the structural determinants of catalytic
activity and stability in *ndufs4*[-/-] mouse complex I we advance understanding of its molecular
phenotype and the basis of complex I-linked mitochondrial dysfunctions.

Results

Deletion of NDUFS4 decreases complex I stability and its activity in membranes

To assess the structural integrity of complex I without the NDUFS4 subunit, we first investigated the complexes present in mitochondrial membranes from *ndufs4*[-/-] (Kruse, et al., 2008) and *wild-type* mouse heart mitochondria by BN-PAGE (Figure 2A). The gels were probed with nitroblue tetrazolium (NBT), which reacts at the NADH-reduced complex I flavin site: *ndufs4*[-/-] membranes exhibited a markedly less intense NADH:NBT-active complex I band at ~1 MDa than *wild-type* membranes, plus a new NADH:NBT-active band at much lower mass not evident in *wild-type* samples. Unusually strong Coomassie-staining was also observed at the position corresponding to the migration of complex V (ATP synthase).

BN-PAGE analyses of *ndufs4*[-/-] and *wild-type* mouse heart mitochondrial membranes were then subjected to complexomic analyses to characterize the subunit compositions of complex I-related species, and to confirm the identity of *ndufs4*[-/-]-specific bands (Figures 2B and EV1). For the *wild-type* membranes, 40 of the 44 complex I proteins were detected in the ~1 MDa complex I band. Only subunits ND3, ND4L and ND6, which are hydrophobic proteins with few tryptic peptides, were not detected. For the *ndufs4*[-/-] membranes, two new *ndufs4*[-/-]-specific bands were observed at ~800 kDa and ~200 kDa (by including species with known masses as standards in the BN-PAGE analyses their masses were estimated as 812 and 199 kDa, respectively, Figure EV1). Mapping the 32 complex I subunits detected in the ~800 kDa band to the structure of the *wild-type* enzyme showed that they represent a substantial subcomplex containing the whole of the membrane domain and the lower (Q-module) section of the hydrophilic domain, but lacking the top section of the hydrophilic domain that contains the active site for NADH oxidation (the NADH-dehydrogenase module or N-module) (Figure 2B and EV1). The N-module subunits (NDUFV1, NDUFV2, NDUFV3, NDUFS1 and NDUFA2) were identified in the NADH:NBT-active ~200 kDa band. The ~800 kDa subcomplex contains the subunits that catalyze ubiquinone reduction and proton pumping, so we refer to it as the QP subcomplex. Two complex I assembly factors, ACAD9 and NDUFAF2, also comigrate with the QP subcomplex, although NDUFAF2 is present in a similar position in the *wild-type* membrane analyses, one gel slice below the intact complex. An additional three subunits, NDUFS6, NDUFA12 and NDUFA7, usually located at the

interface between the N-module and QP subcomplex, were detected elsewhere in the analyses, not associated with either. The complexome analyses thus indicate that the absence of NDUFS4 weakens or prevents the attachment of the N-module (and adjacent subunits) to the complex.

Kinetic measurements of the specific NADH:O_2 oxidoreductase activity (turnover of the CI-III-IV pathway, Figure 2C) show that the rate of complex I-linked respiration by *ndufs4*$^{-/-}$ mouse heart mitochondrial membranes is only ~10% of that of *wild-type* membranes. No corresponding difference was observed for the specific succinate:O_2 oxidoreductase activity (CII-III-IV, Figure 2D), confirming that the loss of NADH:O_2 activity stems from a complex I defect. It could indicate a lower complex I content, and/or a lower intrinsic activity of the *ndufs4*$^{-/-}$ complex relative to the *wild-type* complex. The amount of intact complex I present (irrespective of its capability to reduce ubiquinone) is typically assessed by i) semi-quantitative evaluation of the intensities of Coomassie or NBT-stained bands on BN-PAGE (precluded here by enzyme instability, see below), or ii) quantitative determination of the rates of flavin-localized reactions that couple NADH oxidation to reduction of artificial electron acceptors such as hexaammineruthenium III (HAR) or hexacyanoferrate (FeCN). Here, we found that the kinetics of the flavin-localized reactions were drastically altered in *ndufs4*$^{-/-}$ membranes (Figure EV2): the K_M value for HAR is decreased 120-fold relative to in the *wild-type* enzyme, and the K_M values for NADH are increased 3-fold for both the NADH:HAR and NADH:FeCN reactions. The altered values are consistent with the proximity of NDUFS4 to the flavin site, and the drastically decreased K_M value for HAR also suggests the possibility of a new HAR binding site opening up in the *ndufs4*$^{-/-}$ variant. They also preclude flavin-site kinetics being used to determine the complex I content. Instead, we turned to mass spectrometry of whole membrane digests and performed unlabeled quantification relative to the sum of peptides from VDAC1, 2 and 3: the relative abundance of *ndufs4*$^{-/-}$ complex I peptides was 66 ± 2% of that observed for *wild-type* membranes (Figure 2E). This experiment was carried out using membranes from kidney to minimize animal usage and, in the samples studied, the NADH:O_2 rates were 1.36 ± 0.38 and 0.38 ± 0.03 µmol NADH mg^{-1} min^{-1} (S.D. n = 8) for *wild-type* and *ndufs4*$^{-/-}$ membranes, respectively (28 ± 2%, matching the earlier value of 25 ± 11% from kidney tissue (Calvaruso *et al*, 2012)). Our data thus suggest that the lower NADH:O_2 activity in kidney membranes is due to both decreased content and decreased intrinsic catalytic activity of the *ndufs4*$^{-/-}$ enzyme, which we estimate to catalyze at ~40% of the *wild-type* rate.

The N-module is hydrophilic, so if it dissociates from the complex then it would be expected
to be lost in the supernatant following collection of the membranes by centrifugation – yet we
see it in mass spectrometry analyses of our membrane preparations. This observation
suggests the N-module remains attached to the QP subcomplex *in vivo* and in membrane
preparations, but dissociates during BN-PAGE. To test this proposal, *ndufs4*[-/-] membranes
were treated with the cross-linker bis(sulfosuccinimidyl)suberate (BS[3], which links nearby
lysine residues to one another) before BN-PAGE, substantially increasing the amount of
intact complex I observed (Figure 2F). Note, however, that BS[3] treatment was deleterious for
catalysis, decreasing the rate of NADH:decylubiquinone oxidoreduction to $9.1 \pm 1.2\%$ of the
control rate when added at 2.5 mM to DDM-solubilized bovine heart membranes, which we
attribute to effects on catalytic enzyme motions and/or substrate access.

Together, our results suggest that lack of NDUFS4 destabilizes the association of the N-
module to the *ndufs4*[-/-] complex, but that the N-module remains bound *in vivo* and is lost
during BN-PAGE analyses.

Purification of *ndufs4*[-/-] complex I and initial cryo-EM study of the kidney enzyme

Complex I from *ndufs4*[-/-] mice was isolated using the same method as for the *wild-type*
complex (Agip *et al*, 2018), by solubilization with DDM, ion-exchange and size-exclusion
chromatography. As expected, the yield was lower, but the complex was purified successfully
from both heart and kidney membranes with the same apparent molecular mass as for the
wild-type enzyme (Figure 3A-B). Kinetic measurements were then compared for the *ndufs4*[-/-]
and *wild-type* enzymes, using both heart membranes (Figure 3C) and purified heart enzymes
(Figure 3D). The rates of NADH:ubiquinone oxidoreduction (NADH:O_2 and NADH:DQ
reactions, respectively) were much lower for the *ndufs4*[-/-] variant (<10% of the *wild-type*
activity). Flavin-site assays using artificial electron acceptors (NADH:FeCN, HAR and
APAD[+]) also showed decreased activities for *ndufs4*[-/-] relative to *wild-type*, although the
relative rates of each reaction varied. The rate of H_2O_2 (from reaction of the NADH-reduced
flavin with O_2) also decreased for *ndufs4*[-/-] relative to *wild-type* complex I. As the purified
ndufs4[-/-] complex I was determined to contain 0.93 ± 0.02 molecules of FMN per 1 MDa of
protein (close to the expected 1 FMN per MDa), and the thermal stability of the bound FMN
was not altered substantially (Figure 3E), it appears that the decreased flavin-site activities
reflect a change in the catalytic properties of the site itself (rather than a decreased content of
FMN). Finally, mass spectrometry analyses on the enzyme from *ndufs4*[-/-] heart membranes

showed that two additional complex I-linked proteins, ACAD9 and NDUFAF2, were co-
purified with it (Figure EV3, Table S1), with a substantial band visible on SDS-PAGE where
both ACAD9 and a related protein, ACADVL, were detected. ACAD9 and ACADVL both
contain FAD, and a substoichiometric level of 0.35 ± 0.01 FAD was detected per MDa of
protein. Mass spectrometry analyses of the *wild-type* enzyme only detected NDUFAF2 with
poor quality peptide data and did not detect ACAD9, but a matching band on SDS PAGE was
identified as ACADVL (Figure EV3, Table S2).

For preliminary structure evaluation, complex I from *ndufs4*$^{-/-}$ mouse kidney was subjected to
cryo-EM imaging (by working with the enzyme from kidney as well as from heart we were
able to decrease the number of mice required). No cross-linker treatment was used. The
enzyme was frozen onto PEGylated UltrAuFoil R0.6/1 gold grids (Meyerson *et al*, 2014;
Blaza *et al*, 2018) and imaged with an FEI Falcon III detector in integrating mode. Around
150,000 particles were picked, but 2D and 3D classification using cryoSPARC (Punjani *et al*,
2017) eliminated most of them because they did not resemble intact complex I, leading
eventually to a reconstruction from only 7,563 particles with an estimated resolution of 6.2 Å
(Figure S1, Table S3) [Electron Microscopy Data (EMD): 16514]. Despite the low yield of
intact complex I particles and low final resolution, particularly for the N-module, the map
was sufficient to observe a mature complex I structure, which appeared to lack density for the
NDUFS4 subunit, as well as for subunit NDUFS6 (Figure EV4). As the protein eluting from
the size exclusion column has a mass consistent with intact complex I (Figure 3A), damage at
the air-water interface during grid freezing probably causes the more fragile *ndufs4*$^{-/-}$
complex to fragment, much like during BN-PAGE, leading to loss of most of the intact
particles.

Cryo-EM structures of *ndufs4*$^{-/-}$ complex I from mouse heart mitochondria

To stabilize the complex during grid freezing and improve the yield of intact particles, the
BS^3 cross-linking procedure tested on membranes was extended to the purified enzyme. BS^3
was applied twice, once during membrane solubilization and once immediately before the
enzyme was added to the grids, a procedure that enabled a substantial proportion of the
complex to survive BN-PAGE analysis (Figure 3F). The cross-linked complex was frozen
onto PEGylated UltrAuFoil R0.6/1 gold grids and 7,309 micrographs from a single grid were
collected on an FEI Titan Krios microscope fitted with a Gatan Quantum K3 Summit detector
with energy filter (Figure S2, Table S3). A low (7.1 Å)-resolution class, lacking clear density

for the N-module, was found that corresponds to the shape predicted for the ~800 kDa QP

subcomplex identified in complexome analyses, and the reconstruction of the intact complex

I achieved a global resolution of 2.9 Å (Figure S2). A survey of lysine densities failed to

identify any continuous features from Lys-Lys cross-links, suggesting the enzyme is

stabilized by a mixture of low-occupancy links. As the map for the intact enzyme showed

clear signs of heterogeneity, the intact particles were then subjected to focused classification

using cryoSPARC, resulting in three major classes (Figure S3, S4). The class 1 map exhibited

a large low-resolution protein density adjacent to subunit ND1, the class 2 map (70% of the

particles) resembled the consensus map, and the class 3 map exhibited an additional protein

density on the heel of the complex, where subunit NDUFA12 usually resides. All three

classes were compared by map-to-map correlation to the mouse *wild-type* active and deactive

resting states (EMD: 11377 (Bridges *et al*, 2020) and EMD: 11810 (Yin *et al*, 2021)) and all

three were found to resemble the active state more closely than the deactive state (Table S4).

Furthermore, all the specific hallmarks of the active state are present in all three maps,

including the relative orientation of NDUFA5 and NDUFA10, the lack of π-bulge in ND6-

TMH3, and the ordered states of loops in NDUFS2, NDUFA9, ND3 and ND1 (Agip *et al*,

2018; Chung *et al*, 2022a).

Deletion of NDUFS4 weakens the attachment of the N-module

Comparing the class 2 model of *ndufs4$^{-/-}$* complex I to the model of the active state of *wild-*

type mouse complex I (we use PDB: 6ZR2 (Bridges *et al*, 2020) as our reference structure

throughout) shows that deletion of the NDUFS4 subunit leaves a deep groove on the surface

of the hydrophilic domain (Figure 4A). In the consensus class 2 map, the density for the N-

module is substantially lower in resolution than the density for the rest of the enzyme but

when the N-module is locally refined its resolution improves to the same level (Figure 4B,

S4A-B). The resultant N-module model (subunits NDUFV1, NDUFV2, NDUFV3, NDUFS1

and NDUFA2) has an all-atom RMSD of 1.20 Å against the equivalent N-module of the

wild-type (NDUFS4-containing) reference model, and the density for the FMN is clear, in its

expected location. Thus, the structure of the N-module itself is closely conserved in structure

between the *wild-type* and *ndufs4$^{-/-}$* enzymes, but in *ndufs4$^{-/-}$* the module is not rigidly fixed to

the rest of the enzyme and adopts a distribution of positions relative to it (Figure 4C and

Supplementary Movie 1). Finally, the NDUFA9 subunit drops slightly towards the membrane,

relative to its position in the *wild-type* reference structure (Figure 4D and Supplementary

Movie 1), due to loss of stabilizing contacts from the NDUFA9 N-terminus to subunits
NDUFS4, NDUFS6 and NDUFA12, at a crucial junction between all four subunits (Figure
1C).

Within the *ndufs4*[-/-] N-module, residues 61-68 of subunit NDUFV3 that are adjacent to
NDUFS4 and resolved in the *wild-type* enzyme, remain disordered even after local
refinement. Subunits NDUFS2, NDUFS3, NDUFS8, and NDUFA5, which are adjacent to the
N-module in the QP subcomplex and usually interact with NDUSF4, are not changed by its
absence. The C-terminus of NDUFA6 is unchanged in its position on the N-module, but
shifted relative to the QP subcomplex. There is no density observed for subunit NDUFA12,
or for the N-terminal regions of NDUFS6 (residues 1-39), NDUFA7 (residues 1-35), and
NDUFA9 (residues 1-17) that are well ordered in the *wild-type* reference structure (Figure
EV5). The proximity of these features in the *wild-type* enzyme (Figure 4E) suggests a
cascade of effects, in which the absence of NDUFS4 affects the N-terminus of NDUFA9 and
the C-terminal loop of NDUFA12, which is normally overlaid by the N-terminal domain of
NDUFS6, and loss of NDUFA12 propagates to the adjacent N-terminus of NDUFA7. Loss of
NDUFA12 is consistent with our mass spectrometry analyses (Figures EV1, EV3), as well as
with previous proteomic studies of *ndufs4*[-/-] mouse tissues, *ndufs4*[-/-] mouse embryonic
fibroblasts, and NDUFS4-mutated human cells (Adjobo-Hermans *et al*, 2020; Stroud *et al*,
2016). The C-terminal domain of NDUFS6 and the rest of NDUFA7 appear otherwise
unaffected structurally, with the Zn cofactor of NDUFS6 in its expected location. However,
like the N-module, these two subunits were not found to co-migrate with the rest of the
ndufs4[-/-] complex in complexome analyses (Figures 2B, EV1) suggesting they are also only
weakly bound.

Association of assembly factor NDUFAF2 with *ndufs4*[-/-] complex I

The structure of the class 3 *ndufs4*[-/-] complex I matches that from class 2 closely, except that
assembly factor NDUFAF2 is bound in the place occupied by subunit NDUFA12 in the *wild-*
type complex (Figure 5A), and the whole of the adjacent NDUFS6 subunit is absent,
including the Zn-containing C-terminal domain present in class 2 (see Table 1 for a
comparison of complex I compositions). The presence of NDUFAF2 is consistent with our
complexomic analyses and mass spectrometry on the isolated enzyme, although the particle
classification data suggest that only ~14% of the population contains NDUFAF2, while the
other ~86% (classes 1 & 2) contain NDUFS6 instead: NDUFS6 and NDUFAF2 were not

observed together in any class. The location of NDUFAF2 overlaps with that of NDUFA12
(in the *wild-type* enzyme) through a common three-strand β-sheet and adjacent short helix;
however, NDUFAF2 also contains a long 'diagonal' helix that is not present in NDUFA12,
which crosses the region occupied by the small central helix of NDUFS6 in the *wild-type*
enzyme (Figures 5B and 1C). A 35-residue unresolved stretch of NDUFAF2 is followed by a
C-terminal β-hairpin, which interacts with subunit NDUFS1 (Figure 5B-C). The location and
fold of NDUFAF2 matches well to that observed in the model of the ΔNDUFS6-complex I
from *Y. lipolytica* (Parey *et al*, 2019) (Figure 5C) except that the *Y. lipolytica* model lacks the
β-hairpin. Finally, the low resolution map from *ndufs4$^{-/-}$* kidney matches best to the heart
class 3 map, with density for the NDUFAF2 diagonal helix clearly visible, and no density
observed for NDUFS6 (Figure EV5).

ACAD association with *ndufs4$^{-/-}$* complex I

The 12% of particles in class 1 of the *ndufs4$^{-/-}$* complex were found to have an additional
large low-resolution density on the heel of the complex. Particle subtraction of this region
followed by local refinement with C2 symmetry (Figures 6A, S3) produced a 4.2-Å
resolution map for the additional density, which was visually identified as a dimer of an
ACAD-family protein. Mass spectrometry on purified heart *ndufs4$^{-/-}$* complex I (prepared
without cross-linking) identified two candidates, ACAD9 and ACADVL, both with mass ~65
kDa (Figure EV3). Although our complexome analyses of DDM-solubilized *ndufs4$^{-/-}$* heart
membranes (Figures 2B, EV1) revealed co-migration of ACAD9 and the ~800 kDa QP
subcomplex (but not co-migration of ACADVL), our MALDI-TOF and LC/MS analyses
both contained substantially more peptides (and greater sequence coverage) for ACADVL
(Table S1). Considering the sequence similarity between the two proteins, we expect a
similar number of peptides from each if they are present in equimolar quantities, so these data
suggest ACADVL is present in greater abundance than ACAD9. We thus chose to model the
density as a dimer of ACADVL (Figure 6B), and a survey of positions where bulky residues
(Tyr, Trp and Phe) vary between the two proteins subsequently confirmed that ACADVL is
the major species present.

The resolution achieved was sufficient to model the two FAD cofactors in the ACAD dimer
(Figure 6B), accounting for the ~0.3 FAD observed per 1 MDa of protein in purified complex
I (two FAD per dimer bound to 12% of particles). Overall, the density corresponds closely to

the model of human ACADVL solved by X-ray crystallography (PDB: 3B96) (McAndrew *et*

al, 2008), but with an extra α-helical region (residues 490-518) resolved. This region is

inserted into the micelle of our cryo-EM density and braces the dimeric ACAD against the

hydrophobic heel of the complex (Figure 6C), with one helix embedded in the micelle,

interacting with core subunit ND1, and the other less deeply embedded, approaching subunit

NDUFA3. However, the position of the ACAD dimer is flexible relative to complex I and

their interface is not well resolved. Fitting both models into a composite map showed the

closest proximity residues to be ACADVL-Thr514 (monomer B) with NDUFA3-Ile34, and

ACADVL-Arg513 (monomer A) with ND1-His171. Comparison of the local structures of

ndufs4$^{-/-}$ complex I subunits (ND1, NDUFA3, NDUFA8 and NDUFA13) with the *wild-type*

reference model revealed no differences, including in the closest interacting sidechains, and

no substantive differences were observed between the structures from class 1 (ACAD-

containing) and class 2 (no ACAD) either. Finally, as the ACAD bound to class 1 does not

overlap spatially with the NDUFAF2 bound to class 3 we do not expect them to be mutually

exclusive, so it is likely that the expected fourth class that contains both has simply been

missed due to its low population (predicted ~1250 particles or 1.8%).

The location of ACAD binding in our map (Figure 6A) is surprising: ACAD proteins are

known to be membrane-associated and located in the mitochondrial matrix (McAndrew *et al*,

2008), but in our structure the ACAD dimer appears to have 'slipped' around, into a position

that would be occupied by the lipid tails of a membrane bilayer. No density to suggest the

presence of a similarly-bound ACAD could be identified in the maps from the *ndufs4*$^{-/-}$

kidney enzyme, which was not treated with a cross-linker, or in any of our structural data on

wild-type mouse complex I (Agip *et al*, 2018; Bridges *et al*, 2020; Grba *et al*, 2022), even

though the abundance of ACAD proteins in *ndufs4*$^{-/-}$ and *wild-type* mitochondria are similar

(Figure 6D). Therefore, we conclude the ACAD dimer is artefactually bound and stabilized in

the *ndufs4*$^{-/-}$ enzyme by the cross-linking procedure that was applied to stabilize the N-

module, and that its observed binding mode is not physiologically relevant. Although we do

not observe connecting density, the nitrogens of the Lys sidechains on ACADVL (monomer

A, Lys556 and Lys557) and NDUFA8 (Lys105) are modelled ~4 and ~14 Å apart,

respectively, consistent with reaction with the BS[3] cross-linker (Figure 6E).

Structure of complex I from *ndufs4*^{-/-} mouse contains bound ubiquinone

Continuous density resembling that expected for ubiquinone-9 was identified along the length of the ubiquinone-binding channel in the class 2 *ndufs4*^{-/-} enzyme (Figure S5A). The density feature observed is substantially longer than the BS³ cross-linker molecule, and so we conclude it originates from bound ubiquinone, even though the density for the ubiquinone headgroup is poorly defined, precluding a confident model being built. Similar but less contiguous densities were also observed in class 3, whereas the densities observed in class 1, as well as in the reference map from the *wild-type* complex (EMD: 11377 (Bridges *et al*, 2020)), were fragmented and could not be interpreted. Although the class 2 *ndufs4*^{-/-} ubiquinone headgroup is not resolved, the residues around its binding site match closely to the residues in the *wild-type* reference model (Figure S5B). Comparison to the models for mouse complex with inhibitors bound (Grba *et al*, 2022; Bridges *et al*, 2020), and to other mammalian complexes with ubiquinone-10 bound in the active site (Chung *et al*, 2022b; Gu *et al*, 2022) (Figure S5C), show that the isoprenoid chain overlays well with the location of other inhibitors and substrates modelled in the channel. Finally, our model provides no explanation for why our class 2 *ndufs4*^{-/-} map exhibits better density for ubiquinone-9 than the *wild-type* enzyme (Agip *et al*, 2018; Bridges *et al*, 2020), even at higher resolution. The difference may be due to an indirect effect of the absence of NDUFS4 or the BS³ cross-linker, perhaps restricting an enzyme motion required for ubiquinone exchange, consistent with the loss of catalytic activity in the *ndufs4*^{-/-} complex, and with further loss upon cross-linker treatment.

Discussion

Is *ndufs4*^{-/-} complex I unstable or incompletely assembled?

Purification and cryo-EM reconstruction of complex I from *ndufs4*^{-/-} mouse kidney and heart tissue yielded maps of an intact complex I, with the N-module bound, which contains almost all the subunits of the enzyme (including all the core-subunits) and exhibits a *wild-type*-like mass on BN-PAGE. The ~800 kDa subcomplex identified on BN-PAGE both here and previously (Adjobo-Hermans *et al*, 2020; Calvaruso *et al*, 2012) likely results from fragmentation of the intact complex under BN-PAGE conditions, which we mitigated by adding a cross-linker (Figure 2F). The presence of an intact complex in mouse *in vivo* explains why the *ndufs4*^{-/-} variant is not embryonically lethal, although the amount of enzyme

present is substantially diminished (by ~50%), as well as less stable than the *wild-type*

enzyme. A similar conclusion, that the *ndufs4^-/-* enzyme is intact *in vivo*, was drawn for *Y.*

lipolytica complex I (Parey *et al*, 2019; Kahlhöfer *et al*, 2017), and has also been suggested

previously for the mammalian complex (van de Wal *et al*, 2022; Adjobo-Hermans *et al*,

2020). The conclusion that the ~800 kDa subcomplex observed on BN-PAGE is thus an

artefact of the analysis urges caution when using BN-PAGE to assess clinically identified

complex I defects, where genetic variants that result in a less stable enzyme may be conflated

with those that genuinely result in interrupted-assembly intermediates. However, as well as

being unstable, *ndufs4^-/-* complex I is also incompletely assembled: although it contains the

N-module, it lacks NDUFA12 as well as NDUFS4, nearby structural elements are disordered,

and it exists in different stages of completion, with some complexes still containing

NDUFAF2, and some with NDUFS6 missing or incompletely associated (Figures 4, 5).

Loss of catalytic activity in the absence of NDUFS4

Our kinetic assays revealed substantially lower NADH:ubiquinone oxidoreductase activity

for *ndufs4^-/-* relative to *wild-type* complex I (Figure 3C-D), even when the decreased amount

of enzyme present in membranes is taken into account. The reason for the lower activity,

which was also observed for the *Y. lipolytica ndufs4^-/-* enzyme (Kahlhöfer *et al*, 2017), is

obscure. The static structure of the flavin NADH-oxidizing site does not appear to be

changed by the absence of NDUFS4, but the kinetics of reactions normally localized only at

the flavin site are altered (Figure 3C-D, EV2) and therefore the dynamics of the site may be

different. However, NADH oxidation is not rate limiting during turnover (Birrell 2010) so the

moderate decrease in NADH oxidation rate by the *ndufs4^-/-* complex is unlikely to affect the

overall rate substantially. The NDUFS4 C-terminus lies within 8 Å of the cluster 1 in the FeS

chain (the 4Fe-4S cluster in NDUFV2 that accepts the electrons from the flavin), the $\beta1$-$\beta2$

loop within 12 Å of clusters 2, 3 and 4 (NDUFS1), and the $\beta2$-$\beta3$ loop within 12 Å of cluster

6 (the penultimate, 4Fe-4S cluster in NDUFS8) (Figure 1D). Furthermore, the small drop in

NDUFA9 towards the membrane in the *ndufs4^-/-* variant expands a solvent-filled cavity that

extends towards clusters 5 and 6, the two [4Fe-4S] clusters in NDUFS8, although neither

becomes directly solvent accessible. It is possible that the altered kinetics (particularly the

decreased K_M value) for the NADH:HAR oxidoreduction reaction result from interaction of

HAR at a new *ndufs4^-/-*-specific site in this region. In *Y. lipolytica* complex I, deletion of

NDUFS4 perturbed the electron paramagnetic resonance spectra of reduced FeS cluster 1

(signal N3), 2 (signal N1b, [2Fe-2S] cluster in NDUFS1), and 6 (signal N4), consistent with
their changed dielectric environment (Kahlhöfer *et al*, 2017). However, as for NADH
oxidation, the rate of intramolecular electron transfer is considered fast (the FeS chain is
reduced during steady-state catalysis in *wild-type* complex I) and so it is unlikely to become
rate limiting. For *Y. lipolytica ndufs4*[-/-] the increased solvent accessibility of the clusters was
proposed to increase electron leakage to O_2, then further oxidative damage to the enzyme in a
vicious cycle (Kahlhöfer *et al*, 2017). Increased cellular ROS production, relative to *wild-type*,
was reported in *ndufs4*[-/-] mouse fibroblasts (Valsecchi *et al*, 2013), but not in *ndufs4*[-/-] mouse
heart (Chouchani *et al*, 2014), and here we did not observe any substantial increase in ROS
production in either *ndufs4*[-/-] mitochondrial membranes or complex I (Figure 3C-D), with the
rate of H_2O_2 production remaining two orders of magnitude slower than the rate of turnover.
Increased complex I-linked ROS in mitochondria may result from changes to the
environment, not the complex I molecule itself, as a result of increased reduction of the
NAD^+ pool (Kussmaul & Hirst, 2006).

The decreased catalytic activity may also result from the altered dynamics and
conformational landscape that likely results from loss of the structural junction between
NDUFS4, NDUFS6 and NDUFA12, which pins N-module subunit NDUFS1 to NDUFA9.
The NDUFA9 C-terminal domain, buttressed by the NDUFS7 C-terminus at the base of the
hydrophilic domain, makes tight interactions with specific phospholipids at the membrane
interface, and may affect the membrane topology near the ubiquinone-binding site. A further
effect on the local membrane structure may arise from the absence of the NDUFA12
amphipathic helix, which is not present in NDUFAF2 (Padavannil *et al*, 2022), and together
these may change the ubiquinone-9/10 binding dynamics. It is possible that the same effect
explains the unexpected retention of ubiquinone-9 in the *ndufs4*[-/-] enzyme (Figure S5).
Finally, conformational changes in the NDUFA9 C-terminal domain have been observed in
the complex I active-to-deactive transition (Zhu *et al*, 2016), and the C-terminal peptide
contacts the start of the ND3-TMH1-2 loop, a structural element central to deactivation
(Chung *et al*, 2022a) that may also change its conformation during catalysis (Cabrera-Orefice
et al, 2018). We were, unusually, unable to find a class of particles that resembled the
deactive state in our *ndufs4*[-/-] complex I, so these subtle changes may also alter the active-to-
deactive transition in the *ndufs4*[-/-] enzyme. More complete understanding of the structural
changes that occur during catalysis is required to further evaluate these proposals.

Comparison with *Y. lipolytica ndufs4*[-/-] complex I

In the *Y. lipolytica ndufs4*[-/-] strain, complex I levels decreased to ~40% of *wild type*, similar to here, but the subunits of the N-module plus NDUFS6, NDUFA12 and NDUFAF2 were all found to stay associated with the intact complex on BN-PAGE (Kahlhöfer *et al*, 2017). Subsequently, a 4 Å-resolution cryo-EM map (PDB: 6RFS) showed that NDUFS6 and NDUFA12 were bound as normal, but no density for assembly factor NDUFAF2 was observed (Parey *et al*, 2019) (see Table 1). Thus, unlike the mouse *ndufs4*[-/-] enzyme, the *Y. lipolytica ndufs4*[-/-] complex has completed assembly of its hydrophilic domain without NDUFS4. Conversely, the mouse *ndufs4*[-/-] enzyme contains a mature membrane domain, but the *Y. lipolytica ndufs4*[-/-] complex does not: it lacks density for the N-terminal half of NDUFA11 and the C-terminus of ND5 (part of the transverse helix and TMH16), and has lost structural integrity for ND6-TMH4 and adjacent loops, as well as for the N-terminus of NDUFS2. NDUFA11 has been reported to associate in the final stages of complex I assembly (Andrews *et al*, 2013; Guerrero-Castillo *et al*, 2017) when the N-module joins the complex, so it is possible that assembly of NDUFA11 onto the *Y. lipolytica ndufs4*[-/-] complex stalls without NDUFS4. We note that the structural changes at/around *Y. lipolytica* NDUFA11 resemble those observed in the 'slack' state of mammalian complex I (Zhu *et al*, 2016; Chung *et al*, 2022b), so may also be caused by instability of the *ndufs4*[-/-] complex, either during purification or cryo-EM grid freezing. Finally, like our mouse *ndufs4*[-/-] class 3 structure, the structure of *Y. lipolytica ndufs6*[-/-] complex I (PDB: 6RFQ) (Parey *et al*, 2019) contains NDUFAF2 instead of NDUFA12, consistent with a helix in NDUFAF2 occupying the same space as the loop in NDUFS6 that connects its two domains together, so that both cannot bind together.

Assembly pathway

Work on the assembly of both mammalian and yeast complexes I has led to a model in which the assembly factor NDUFAF2 recruits the N-module onto the nascent complex; NDUFS4 (and perhaps NDUFS6) may also be required to stabilise the N-module, before NDUFAF2 is displaced by NDUFS6 and NDUFA12 (perhaps as a single unit) to complete the enzyme (Lazarou *et al*, 2007; Stroud *et al*, 2016; Parey *et al*, 2019; Adjobo-Hermans *et al*, 2020; Pereira *et al*, 2013; Guerrero-Castillo *et al*, 2017). NDUFAF2, NDUFS4, NDUFS6 and NDUFA12 have also been considered an 'assembly checkpoint', blocking completion of the hydrophilic arm until the rest of the complex is complete (Padavannil *et al*, 2022).

Our structures of mouse *ndufs4*[-/-] complex I show that the N-module can be stabilized on the
nascent complex by NDUFAF2 (without NDUFS4, NDUFS6 or NDUFA12, in class 3), or by
the N-terminal domain of NDUFS6 (without NDUFS4, NDUFA12 or NDUFAF2, in class 2).
We do not observe the N-module bound alone. However, while the N-terminal domain of
NDUFS6 binds between N-module subunits NDUFS1 and NDUFV2 and QP-subcomplex
subunits NDUFS2 and NDUFS8 (Figure 5B), clearly stabilizing attachment of the N-module,
the stabilizing effects of NDUFAF2 are unclear. Its N-terminal domain (modelled from
Arg20 to Tyr106) is bound (predominantly) to subunits NDUFS7, ND1 and NDUFS8 on the
heel of the complex, whereas its C-terminal domain (modelled from Glu141 to Glu161) is
located on NDUFS1 in the N-module (Figure 5B). There is no sturdy connection between
them, since residues 107 to 140 are disordered – and indeed this stretch of the sequence is
unstructured in the alphafold prediction AF-Q59J78 (Jumper *et al*, 2021). Therefore,
NDUFAF2 does not form a stabilizing bridge to attach the N-module. Instead, we propose
that it acts as a 'lasso' to capture it and guide it onto the complex: with its N-terminal domain
firmly attached to the heel, its intrinsically disordered region allows the C-terminus to search
the surrounding space, capture the N-module, and then restrict its freedom to promote
docking (Figure 7). Our proposal explains the importance of NDUFAF2 for N-module
attachment, and why it cannot be substituted by NDUFA12, which subsequently replaces it
on the mature *wild-type* complex, but lacks the flexibly-linked β-hairpin anchor to NDUFS1.
Thus, our class 3 NDUFAF2-bound structure likely represents the first N-module-containing
species on the assembly pathway.

Our structures show that the three subunits required to complete the mature protein (NDUFS4,
NDUFS6 and NDUFA12) all overlap with NDUFAF2 and therefore likely contribute to
displacing it, suggesting they all bind only after the N-module has been recruited. The β2-β3
hairpin-loop (residues 62-77) of NDUFS4, which is probably inserted from the front of the
hydrophilic domain, to pass under the N-module and connect to NDUFS6, NDUFA12 and
NDUFA9 (Figure 1C), displaces a short section of the C-terminus of NDUFAF2. This section
is ordered in our class 3 *ndufs4*[-/-] structure, but disordered in the structure of *Y. lipolytica*
ndufs6[-/-] complex I that contains both NDUFS4 and NDUFAF2 (Parey *et al*, 2019). The
NDUFA12 N-terminal domain replaces the NDUFAF2 N-terminal domain on the heel, and
its C-terminus replaces the NDUFAF2 C-terminus on NDUFS1 (Figure 5C). The connecting
loop, which is ordered in NDUFA12 and interacts with the NDUFS4 β2-β3 loop and
NDUFS6, is trapped under the NDUFS6 C-terminal domain in the mature complex (Figure

1C), which must thus bind either after or along with it. Our class 2 *ndufs4$^{-/-}$* structure,

representing the majority of the particles in the heart complex, and containing the N-terminal

domain of NDUFS6 but not NDUFAF2, is thus surprising: first, because it suggests that

NDUFS6 alone is sufficient to dislodge NDUFAF2, and second, because NDUFA12 is

absent despite the availability of its binding site. Whereas in *Y. lipolytica*, *ndufs4$^{-/-}$* complex I

binds both NDUFS6 and NDUFA12 to complete assembly without NDUFS4 (Parey *et al*,

2019), assembly of the mouse *ndufs4$^{-/-}$* complex stalls as NDUFA12 and the C-terminal

domain of NDUFS6 fail to associate. However, the difference in the molecular phenotypes

can be assigned, not to structural effects, but to the near-complete absence of NDUFA12 in

ndufs4$^{-/-}$ mouse tissues, embryonic fibroblasts and NDUFS4-mutated Leigh syndrome patient

cells (Adjobo-Hermans *et al*, 2020); similar observations were also made for *ndufs4$^{-/-}$* mouse

glutamatergic vestibular neurons, that are affected in Leigh syndrome (Gella *et al*, 2020).

Why NDUFA12 is also absent, as well as NDUFS4, is currently unclear, as well as

unfortunate, since it compounds the *ndufs4$^{-/-}$* defect and its pathological consequences by

preventing the final stages in assembly.

Implications of the NDUFS4 mouse model for complex I-linked disease

The *ndufs4$^{-/-}$* mouse displays a severe Leigh-like phenotype (Kruse *et al*, 2008), consistent

with decreased complex I catalysis (which results in decreased ATP production) causing

increased steady-state NADH and O_2 levels, which together may result in increased ROS

production. The discovery that a hypoxic environment mitigates the *ndufs4$^{-/-}$* mouse

phenotype (Jain *et al*, 2016) suggested oxidative damage as a substantial contributor to

disease, with emphasis later placed on locally increased O_2 in the *ndufs4$^{-/-}$* mouse brain (Jain

et al, 2019). These data are consistent with our assays on isolated *ndufs4$^{-/-}$* mitochondrial

membranes and complex I (Figure 3), which support the concept that increased *ndufs4$^{-/-}$* ROS

production in mitochondria stems from the altered mitochondrial conditions (Kussmaul &

Hirst, 2006), rather than from an increased molecular-level propensity of the *ndufs4$^{-/-}$*

complex I to produce ROS. However, this mechanism suggests that hypoxia should benefit

all complex I-linked mitochondrial diseases that similarly compromise the rate of catalysis,

not be specific to a particular subset (including the *ndufs4$^{-/-}$* family) or a specific genetic

background (or species). It is thus essential to determine how general an effect hypoxia-

rescue is. Should it be specific to *ndufs4$^{-/-}$* variants, then one possible explanation is increased

susceptibility to O_2-induced FeS cluster degradation, based on the proximity of NDUFS4 to

clusters along the chain (Figure 1D). The cluster chain extends beyond the N-module, such
that repair or replacement may require either complete regeneration of the complex, or
disassembly and replacement of a larger proportion of the hydrophilic arm than proposed for
either an N-module-specific salvage pathway (Szczepanowska *et al*, 2020), or enhanced
turnover of subunits in the N-module and adjacent hydrophilic arm (Dieteren *et al*, 2012;
Krishna *et al*, 2021). Differences between our *ndufs4*[−/−] and *wild-type* mouse complex I
structures have also provided some intriguing hints for additional subtle molecular changes,
including the presence of bound ubiquinone, changes to the active-deactive transition, and the
substantially decreased catalytic activity, which we cannot currently explain through
incomplete assembly of the enzyme, or instability of the N-module. Wider knowledge of the
effects of hypoxia on different complex I defects with varying molecular phenotypes will
thus be crucial for elucidating both the origins of individual defects, the mechanism of
hypoxia-rescue, and the balance of enzyme synthesis, assembly and degradation that occurs
through complex feedback loops in mitochondrial homeostasis.

Materials and methods

Animals

All procedures were carried out in accordance with the UK Animals (Scientific Procedures)
Act, 1986 (PPL: P6C97520A) and the University of Cambridge Animal Welfare Policy.
Animals were maintained in a temperature- and humidity-controlled animal care facility with
a 12-h light/12-h dark cycle and free access to water and food and sacrificed by cervical
dislocation. To produce constitutive *ndufs4* knockout (KO) mice, mice heterozygous for the
ndufs4 knockout (*ndufs4*[+/−]) were bred together, and *wild-type* littermates used as controls
(Kruse *et al*, 2008). The genotype of each mouse was confirmed by PCR.

Preparation of mitochondrial membranes from mouse heart and kidney

Heart and kidney tissues were excised from *ndufs4*[−/−] and wild type mice, and mitochondria
isolated as described previously (Fernández-Vizarra *et al*, 2010). Heart tissues were
immersed immediately in ice-cold buffer AT (10 mM Tris-HCl, 75 mM sucrose, 225 mM
sorbitol, 1 mM ethylene glycol-bis (β-aminoethyl ether)-N, N,N′,N′-tetraacetic acid (EGTA)
and 0.1% (w/v) fatty acid-free bovine serum albumin, pH 7.4 at 4 °C). Kidney tissues were

stored in ice-cold buffer A (0.32 M sucrose, 1 mM ethylenediaminetetraacetic acid (EDTA),

and 10 mM Tris–HCl, pH 7.4 at 4 °C).

The hearts were diced, washed, then homogenized in 10 mL buffer AT per gram of tissue by

seven to ten strokes of a Potter–Elvehjem homogenizer fitted with a Teflon pestle at ~1,000

rpm. The homogenate was centrifuged (1,000 x g, 5 min), then the supernatant was

centrifuged (9,000 x g, 10 min) to collect the crude mitochondria. The kidneys were diced,

washed, then homogenized in 5 mL buffer A per gram of tissue by five to eight strokes of the

homogenizer at ~600 rpm. The homogenate was centrifuged (1,000 x g, 5 min), then the

supernatant centrifuged (15,000 x g, 10 min) to collect the crude mitochondria. The pellets

were suspended in resuspension buffer (20 mM Tris-HCl, 1 mM EDTA, 10% glycerol, pH

7.4 at 4 °C) to a protein concentration of ~10 mg mL^{-1}. Pierce protease inhibitor tablets were

added (1 tablet to 50 mL suspension) and aliquots were frozen and stored at –80 °C.

Membranes were prepared as described previously (Agip et al, 2018), where mitochondrial

suspensions were thawed on ice, sonicated using a Q700 Sonicator (Qsonica, USA), at 65%

amplitude with three 5-s bursts of sonication interspersed by 30-s intervals on ice and

centrifuged at 75,000 × g for 1 h. The pellets containing the mitochondrial membranes were

homogenized in resuspension buffer to ca. 5 mg mL^{-1} and stored at −80 °C.

Blue-native (BN) PAGE

For BN PAGE (Schägger & von Jagow, 1991), mitochondrial membranes were solubilized

using 1% dodecyl-β-D-maltoside (DDM) (Glycon, Germany) for 30 min, centrifuged to

remove insoluble materials (32,000 × g, 30 min), and run on Native PAGE™ Novex 3–12%

Bis-Tris gels (ThermoFisher Scientific) according to the manufacturer's instructions, except

that the gel was run for 30 min. at 100 V, then the inner buffer was exchanged for one

containing 1/10th of the standard cathode buffer and run for a further 1.5 h at 180 V. Proteins

were visualized using colloidal Coomassie R250, or the gels de-stained with MilliQ water for

in-gel complex I activity assays using 100 μM NADH and 1 mg mL^{-1} nitroblue tetrazolium

(NBT) (Wittig et al, 2007).

Catalytic activity measurements

Assays on membranes were carried out in 10 mM Tris-SO$_4$ (pH 7.4) and 250 mM sucrose at

32 °C in a SPECTRAmax PLUS384 spectrophotometer (Molecular Devices, UK) using 10

µg-protein mL^{-1} membranes supplemented by 1.5 µM horse heart cytochrome c and 15 µg

mL^{-1} alamethicin. NADH:O$_2$ oxidoreduction was initiated by addition of 200 µM NADH,

monitored at 340-380 nm (ε = 4.81 mM^{-1} cm^{-1}) and its inhibitor sensitivity checked by

addition of 1 µM piericidin A (Sigma). Succinate:O$_2$ oxidoreduction was initiated by addition

of 5 mM succinate, monitored using the reduction of 2 mM NADP$^+$ at 340-380 nm (ε = 4.81

mM^{-1} cm^{-1}) by a coupled assay system comprising 90 µg mL^{-1} fumarate hydratase (FumC)

and 500 µg mL^{-1} oxaloacetate decarboxylating malic dehydrogenase (MaeB) with 2 mM

MgSO$_4$ and 2 mM K$_2$SO$_4$ (Jones & Hirst, 2013), and its inhibitor sensitivity checked by

addition of 2 µM atpenin A5 (Santa Cruz Biotechnology, USA). Flavin-site activity

measurements were carried out in the presence of 15 µg mL^{-1} alamethicin and 2 µM

piericidin A as follows: NADH:FeCN oxidoreduction used standard concentrations of 200

µM NADH and 1 mM FeCN and was monitored using ε = 4.81 mM^{-1} cm^{-1} at 340-380;

NADH:APAD$^+$ oxidoreduction used 200 µM NADH and 1 mM APAD$^+$ (ε = 3.16 mM^{-1} cm^{-1}

4 450-500 nm) and NADH:HAR oxidoreduction used 100 µM NADH and 3.5 mM HAR (ε

= 4.81 mM^{-1} cm^{-1} at 340-380 nm).

H$_2$O$_2$ production was measured using 10 U mL^{-1} horseradish peroxidase, 10 µM Amplex red

and 30 µM NADH with the background rates in the presence of catalase (5000 Units mL^{-1})

and in the absence of protein subtracted (Kussmaul & Hirst, 2006). Assays using purified

complex I were carried out similarly at 32 °C in 20 mM Tris-Cl pH 7.5 (in the absence of

alamethicin, cytochrome c or piericidin A). NADH:DQ oxidoreduction was measured using

100 µM DQ and 100 µM NADH in the presence of 0.15% soy bean asolectin and 0.15% 3-

[(3-cholamidopropyl)dimethylammonio]-1-propanesulfonate (CHAPS).

Preparation of complex I from mitochondrial membranes

Ndufs4$^{-/-}$ complex I was prepared as described previously with minor alterations (Agip *et al*,

2018; Yin *et al*, 2021). 3–4 mL of membrane suspension were solubilized by addition of 1%

DDM, along with 0.005% phenylmethane sulfonyl fluoride (PMSF, Sigma-Aldrich, UK) for

30 min. 0.25 mM of the bis(sulfosuccinimidyl)suberate) (BS^3) cross-linker was added 5 min.

after the start of the solubilization, and the cross-linking reaction quenched after 25 min. (at

the end of the solubilization) by addition of 50 mM Tris-HCl from a 1 M stock solution (pH

7.5, 4 °C). Following centrifugation (32,000 × g, 30 min), the solubilized enzymes were

loaded onto a 1 mL Q-sepharose HP column (GE Healthcare, UK) pre-equilibrated in buffer

containing 20 mM Tris-HCl (pH 7.14 at 20 °C), 1 mM EDTA, 0.1% DDM, 10% (v/v)

ethylene glycol, 0.005% asolectin (Avanti Polar Lipid, USA) and 0.005% CHAPS

(Calbiochem, Merck, Germany), and eluted by an increasing proportion of the same buffer

plus 1 M NaCl. Complex I eluted in ~35% buffer B. The complex I-containing fractions were

collected, concentrated using a 100 kDa MWCO Vivaspin 500 concentrator (Sartorius,

Germany), and eluted from a Superose 6 Increase 5/150GL size exclusion column (GE

Healthcare, UK) in 20 mM Tris-HCl (pH 7.14 at 20 °C), 150 mM NaCl and 0.05% DDM.

Protein concentrations were determined using the BCA assay (Fisher Scientific UK). For the

heart samples for cryo-EM analyses, 2.5 mM BS^3 was added for 30 min. on ice immediately

before cryo-EM grid preparation.

Cryo-EM data collection

UltrAuFoil gold grids (R0.6/1, Quantifoil Micro Tools GmbH, Germany) were prepared as

described previously (Blaza *et al*, 2018). Briefly, they were glow discharged (20 mA, 90 s),

incubated in a solution of 5 mM 11-mercaptoundecyl hexaethyleneglycol (SensoPath

Technologies, USA) in ethanol (Meyerson *et al*, 2014) for 48 hours in an anaerobic glovebox,

then washed with ethanol and air-dried just before use. Using an FEI Vitrobot Mark IV, 2.5

µL of complex I solution (2.55 mg mL^{-1} for kidney CI, 3.82 mg mL^{-1} for heart CI) was

applied to each grid at 4 °C in 100% humidity and blotted for 9–10s at blotting force setting -

10, before the grid was frozen by plunging it into liquid ethane. The highest quality cryo-

frozen grids were identified using a 200 keV Talos Arctica microscope. Datasets for

reconstruction were then collected using a Titan Krios (300 keV) microscope at the UK

National Electron Bio-Imaging Centre at the Diamond Light Source.

A total of 3,373 micrographs from the *ndufs4$^{-/-}$* mouse kidney complex I sample were

collected with a Falcon III detector in integrating mode, each with 59 movie frames, using the

FEI EPU software. The calibrated pixel size was 1.46 Å, the defocus range was -2.1 to -3.3

µm and the total electron dose was 49.24 electrons Å^{-2} s^{-1} over a total exposure time of 1.5 s

for each micrograph. Following inspection, 1,982 micrographs were retained for analysis. A

total of 7,309 micrographs from the *ndufs4$^{-/-}$* mouse heart complex I sample were collected

with a Gatan K3 camera in super-resolution mode, each with 25 movie frames, using the FEI

EPU software with aberration-free image shift. The calibrated pixel size was 1.352 Å, the

defocus range was -1.5 to -2.9 µm and the total electron dose was 45 electrons Å^{-2} s^{-1} over a

total exposure time of 5.4 s for each micrograph.

Cryo-EM data processing for complex I from *ndufs4$^{-/-}$* mouse kidney

Cryo-EM data processing was carried out using RELION 3.0 (Scheres, 2012). Micrographs were motion corrected using Motioncor2 with dose-weighting (Zheng *et al*, 2017) followed by CTF estimation using Gctf (Zhang, 2016). 149,430 particles were automatically picked and extracted with a box size of 360 pixels. The particles were imported to cryoSPARC 3.3.2 (Punjani *et al*, 2017) and first 2D classified to remove non-protein junk. The resulting 115,461 particles went through rounds of *ab-initio* and heterogeneous refinements to produce 3D volumes that were used to re-classify them into 6 classes. The class resembling intact complex I contained 19,260 particles and was further subclassified through *ab-initio* and heterogeneous refinements to remove particles that lacked the N-module of the complex. This approach yielded 7,563 particles that were refined using non-uniform refinement, minimizing per-particle scales, and optimizing defocus, beam tilt and trefoil. The final mask for post processing was created in RELION using a map generated from the model of the *ndufs4$^{-/-}$* mouse heart class 3 using the molmap command in Chimera (Pettersen *et al*, 2004) with a binary extension of 4 pixels and a soft edge of 8 pixels. Postprocessing was performed on the final map using RELION. The global resolution of the map was 6.2 Å based on the FSC = 0.143 criterion.

Cryo-EM data processing for complex I from *ndufs4$^{-/-}$* mouse heart

Micrographs were motion corrected using Motioncor2 in RELION 3.1 and imported into cryoSPARC v3 (Punjani *et al*, 2017) followed by CTF estimation using Gctf (Zhang, 2016). The initial auto-picking process obtained 1,141,407 particles, which were cleaned by two rounds of 2D classifications resulting in 84,141 complex I-like particles. The particles were further cleaned by a reference-free *ab-initio* 3D classification and 71,761 were selected. These particles were exported into RELION 3.1 (Zivanov *et al*, 2020) and subjected to CTF refinement and Bayesian polishing using all frames. The polished particles were exported to cryoSPARC and subjected to rounds of 3D classification; first the particles were homogeneously refined, and then classified into two classes using a very generous mask including the entire micelle, and a target resolution of 20 Å. The two classes were then each subclassified; one subclass contained additional density to the side of the micelle, and another contained density in the region where NDUFA12 is usually found. Both classes were subclassified into two classes using a mask around the areas of interest and target resolutions of 10 and 15 Å, respectively. Two of the resulting maps were found to be essentially identical

and were pooled together, and all three maps were refined using non-uniform refinement. The

two maps with additional densities still suffered poorer clarity of these areas than the bulk of

the protein, and so they were subclassified once more using masks around the respective

additional density areas and using the refined volumes (with and without the additional

density) as input models. This resulted in a further class that appeared essentially identical to

the class without additional density, and so all these particles were pooled together. A class of

3035 particles that appeared incompletely classified in the region of additional density were

excluded from further analyses. The three classes identified were refined by non-uniform

refinement using per particle scale factors, and further postprocessed in RELION using a

mask generated with a soft edge of 8 pixels, from a volume made in UCSF Chimera

(Pettersen *et al*, 2004) from a preliminary PDB model based on 6ZR2 (Bridges *et al*, 2020),

with the addition of alphafold models AF-P50544 for ACADVL and AF-Q59J78 for

NDUFAF2 (Jumper *et al*, 2021) as appropriate. Global resolutions are reported at FSC =

0.143. Particle subtraction was performed in cryoSPARC on the ACAD-containing map in

the region of the ACAD dimer using a mask generated with a soft edge of 8 pixels based on a

crudely placed volume made from a dimer of ACADVL in UCSF Chimera. Subtracted

particles were reconstructed, and the volume aligned to find the symmetry axis, for local

refinement with C2 symmetry enforced.

Model building and refinement for complex I from *ndufs4^-/-* mouse heart

Model building and refinement were performed using Coot 0.9.3 (Emsley *et al*, 2010) and

Phenix-1.18.2-3874 (Afonine *et al*, 2018). The active mouse complex I (PDB: 6ZR2) with

the deletion of NDUFS4 and NDUFA12 subunits was rigid body–fitted into the map using

UCSF ChimeraX as the initial model. Where appropriate, alphafold models AF-P50544 and

AF-Q59J78 (Jumper *et al*, 2021) were placed in the density using the fit-in-map function of

ChimeraX, and fitted in initially using ISOLDE (Croll, 2018) in ChimeraX using default

parameters. The models were then subject to cycles of manual adjustment in Coot 0.9.3 using

the globally sharpened map and refined with secondary structure restraints by

phenix.realspace. The ubiquinone-9 molecule was manually fitted into the density and real

space refined by Phenix. Model-to-map FSC curves were generated using the Phenix

Comprehensive validation (cryo-EM) tools and final model statistics were produced by the

Phenix implementations of MolProbity 4.4, and EMRinger.

Complexome profiling experiments

Blue native PAGE/complexome profiling was performed as described previously (Bridges *et al*, 2017), based on the method developed by Heide et al. (Heide *et al*, 2012). The tryptic peptides from excised 1 mm slices (62 from *ndufs4*[-/-] and 64 from *wild-type*) from BN-PAGE gel lanes were analyzed by LC/MS/MS using nanoscale reverse-phase HPLC and an LTQ Orbitrap XL mass spectrometer (Thermo Scientific). Proteins were identified with the MASCOT algorithm (v2.4, Matrix Sciences) and the results from each gel slice compiled into a consensus report using Proteome Discoverer (Thermo Scientific). The MS data were compared with the Uniprot sequence database (August 2020 version containing 563,000 sequences) with peptide mass and fragment mass tolerances of 10 ppm and 0.5 Da, respectively. Variable protein modifications included methionine oxidation, cysteine propionamide, and protein N-terminal acetylation or formylation, with the potential for one missed tryptic cleavage site. Protein quantities in each gel slice were estimated from the average of the intensity values of the three highest intensity peptides.

Protein identification

Proteins from 10 μg of purified complex I from *ndufs4*[-/-] mouse heart (not treated with cross-linker) were resolved by SDS PAGE. The gel was stained with Coomassie R-250, and blue stained bands were excised, digested with trypsin, and the peptides extracted as described previously (Shevchenko *et al*, 1996). Tryptic peptides were analyzed using either a MALDI-TOF-TOF mass spectrometer (Sciex Model 4800 Plus) using α-cyano-4-hydroxycinnamic acid as the matrix compound, or by LC-MS-MS in an Orbitrap XL mass spectrometer as described previously (Andrews *et al*, 2013). Peptide masses and peptide fragment data were identified as described above, except that the tolerances for peptide mass and fragment data were 70 ppm and 0.8 Da for MALDI-TOF-TOF data, or 5 ppm and 0.5 Da for Orbitrap data. Either 1 or 4 missed tryptic cleavages were permitted, respectively.

Relative protein quantification

Tryptic digests were produced from 20 μg samples of mouse kidney mitochondrial membranes. Proteolysis was maximized by the presence of 0.5% (w/v) sodium deoxycholate in 100 mM ammonium bicarbonate and three additions of trypsin (1:100 w/w) over an incubation period of 48 h at 37 °C (Masuda *et al*, 2008). Deoxycholate was removed from the digest mixture by phase extraction with acidified ethyl acetate. Portions of the digests were

analyzed using a Q-Exactive plus mass spectrometer (Thermo Scientific) as described

previously (Bridges *et al*, 2017), and the data compared with the UNIPROT database

comprising 17,107 mouse sequences as described above. The comparisons were performed

with mass tolerances of 5 ppm and 0.01 Da for peptide masses and fragment ion masses,

respectively, an allowance of two missed tryptic cleavages and the potential for methionine

oxidation. These analyses were performed in triplicate.

Common peptides amongst the replicates from *ndufs4*[-/-] and *wild-type* samples were identified

for each protein of interest (complexes I, II, III, IV, ACAD9 and ACAD-VL) and the

VDAC1, VDAC2 and VDAC3 proteins used for normalization. The sum of the peptide areas

for these common peptides were used to produce an abundance value for each protein or

protein complex. The values were internally normalized for each technical replicate to the

sum of common VDAC1+VDAC2+VDAC3 peptides and the average and standard deviation

of the *ndufs4*[-/-] and *wild-type* samples were calculated for each protein or protein complex of

interest. Relative abundance of proteins in the *ndufs4*[-/-] sample were calculated and expressed

as a fraction of the *wild-type* value, with appropriate error propagation.

Data Availability

Cryo-EM maps and models have been deposited in the Electron Microscopy Data Bank

(https://www.ebi.ac.uk/pdbe/emdb/) and Protein Data Bank (https://pdbj.org/) with accession

numbers EMD-16398 and PDB:8C2S (*ndufs4*[-/-] heart class 1); EMD-16515 and PDB:8CA1

(class 1 ACADVL); EMD-16516 and PDB:8CA3 (*ndufs4*[-/-] heart class 2); EMD-16517 and

PDB:8CA4 (*ndufs4*[-/-] heart class 2 N-module); EMD-16518 and PDB:8CA5 (*ndufs4*[-/-] heart

class 3); and EMD-16514 (*ndufs4*[-/-] kidney). Raw cryo-EM images have been deposited into

EMPIAR (https://www.ebi.ac.uk/pdbe/emdb/empiar/) with IDs: EMPIAR-11356 for complex

1554

Acknowledgements

This research was funded by the Medical Research Council (MC_UU_00015/2 and

MC_UU_00028/1 to JH). We thank Shujing Ding, Ian M. Fearnley and Michael E. Harbour

(MBU proteomics facility) for mass spectrometry analyses, James N. Blaza (MBU, now

University of York) for assistance with preliminary cryo-EM experiments, Diamond for

access and support of the cryo-EM facilities at the UK national electron Bio-Imaging Centre

(eBIC), proposals EM17057-2 and BI22238-4, funded by the Wellcome Trust, MRC, and

BBSRC, as well as staff at University of Cambridge cryo-EM facility for grid screening.

Author Contributions

ZY – Data curation, Formal analysis, Investigation, Methodology, Writing – original draft,

Writing – review and editing; ANAA – Formal analysis, Investigation, Methodology, Writing

– review & editing; HRB – Data curation, Formal analysis, Investigation, Methodology,

Supervision, Validation, Visualisation, Writing – original draft, Writing – review and editing;

JH – Conceptualization, Funding acquisition, Methodology, Project administration,

Supervision, Writing – review and editing.

Disclosure and competing interests statement

1571 The authors declare that they have no conflict of interest.

References

Adjobo-Hermans MJW, de Haas R, Willems PHGM, Wojtala A, van Emst-de Vries SE, Wagenaars JA, van den Brand M, Rodenburg RJ, Smeitink JAM, Nijtmans LG, *et al* (2020) NDUFS4 deletion triggers loss of NDUFA12 in Ndufs4 mice and Leigh syndrome patients: A stabilizing role for NDUFAF2. *Biochim Biophys Acta - Bioenerg* 1861: 148213

Afonine P V., Poon BK, Read RJ, Sobolev O V., Terwilliger TC, Urzhumtsev A & Adams PD (2018) Real-space refinement in PHENIX for cryo-EM and crystallography. *Acta Crystallogr Sect D Struct Biol* 74: 531–544

Agip ANA, Blaza JN, Bridges HR, Viscomi C, Rawson S, Muench SP & Hirst J (2018) Cryo-em structures of complex i from mouse heart mitochondria in two biochemically defined states. *Nat Struct Mol Biol* 25: 548–556

Andrews B, Carroll J, Ding S, Fearnley IM & Walker JE (2013) Assembly factors for the membrane arm of human complex I. *PNAS* 110: 18934–18939

Assouline Z, Jambou M, Rio M, Bole-Feysot C, de Lonlay P, Barnerias C, Desguerre I, Bonnemains C, Guillermet C, Steffann J, *et al* (2012) A constant and similar assembly defect of mitochondrial respiratory chain complex I allows rapid identification of NDUFS4 mutations in patients with Leigh syndrome. *Biochim Biophys Acta - Mol Basis Dis* 1822: 1062–1069

Blaza JN, Vinothkumar KR & Hirst J (2018) Structure of the Deactive State of Mammalian Respiratory Complex I. *Structure* 26: 312-319.e3

Brandt U (2006) Energy converting NADH:quinone oxidoreductase (complex I). *Annu Rev Biochem* 75: 69–92

Breuer ME, Willems PHGM, Smeitink JAM, Koopman WJH & Nooteboom M (2013) Cellular and animal models for mitochondrial complex I deficiency: A focus on the NDUFS4 subunit. *IUBMB Life* 65: 202–208

Bridges HR, Blaza JN, Yin Z, Chung I, Pollak MN & Hirst J (2023) Structural basis of mammalian respiratory complex I inhibition by medicinal biguanides. *Science (80-)* 379:

351–357

Bridges HR, Fedor JG, Blaza JN, Di Luca A, Jussupow A, Jarman OD, Wright JJ, Agip ANA,
Gamiz-Hernandez AP, Roessler MM, *et al* (2020) Structure of inhibitor-bound
mammalian complex I. *Nat Commun* 11: 5261

Bridges HR, Mohammed K, Harbour ME & Hirst J (2017) Subunit NDUFV3 is present in
two distinct isoforms in mammalian complex I. *Biochim Biophys Acta - Bioenerg* 1858:
197–207

Budde SMS, van den Heuvel LPWJ, Smeets RJP, Skladal D, Mayr JA, Boelen C, Petruzzella
V, Papa S & Smeitink JAM (2003) Clinical heterogeneity in patients with mutations in
the NDUFS4 gene of mitochondrial complex I. *J Inherit Metab Dis* 26: 813–815

Cabrera-Orefice A, Yoga EG, Wirth C, Siegmund K, Zwicker K, Guerrero-Castillo S,
Zickermann V, Hunte C & Brandt U (2018) Locking loop movement in the ubiquinone
pocket of complex I disengages the proton pumps. *Nat Commun* 9: 4500

Calvaruso MA, Willems P, van den Brand M, Valsecchi F, Kruse S, Palmiter R, Smeitink J
& Nijtmans L (2012) Mitochondrial complex III stabilizes complex I in the absence of
NDUFS4 to provide partial activity. *Hum Mol Genet* 21: 115–120

Chouchani ET, Methner C, Buonincontri G, Hu CH, Logan A, Sawiak SJ, Murphy MP &
Krieg T (2014) Complex I deficiency due to selective loss of Ndufs4 in the mouse heart
results in severe hypertrophic cardiomyopathy. *PLoS One*

Chung I, Grba DN, Wright JJ & Hirst J (2022a) Making the leap from structure to
mechanism: are the open states of mammalian complex I identified by cryoEM resting
states or catalytic intermediates? *Curr Opin Struct Biol* 77: 102447

Chung I, Wright JJ, Bridges HR, Ivanov BS, Biner O, Pereira CS, Arantes GM & Hirst J
(2022b) Cryo-EM structures define ubiquinone-10 binding to mitochondrial complex I
and conformational transitions accompanying Q-site occupancy. *Nat Commun* 13: 2758

Croll TI (2018) ISOLDE : a physically realistic environment for model building into low-
resolution electron-density maps. *Acta Crystallogr Sect D Struct Biol* 74: 519–530

Dieteren CEJ, Koopman WJH, Swarts HG, Peters JGP, Maczuga P, van Gemst JJ,
Masereeuw R, Smeitink JAM, Nijtmans LGJ & Willems PHGM (2012) Subunit-specific
Incorporation Efficiency and Kinetics in Mitochondrial Complex I Homeostasis. *J Biol
Chem* 287: 41851–41860

Emsley P, Lohkamp B, Scott WG & Cowtan K (2010) Features and development of Coot.
Acta Crystallogr Sect D Biol Crystallogr 66: 486–501

Fassone E & Rahman S (2012) Complex I deficiency: clinical features, biochemistry and
molecular genetics. *J Med Genet* 49: 578–590

Fernández-Vizarra E, Ferr\'\in G, Pérez-Martos A, Fernández-Silva P, Zeviani M &
Enr\'\iquez JA (2010) Isolation of mitochondria for biogenetical studies: An update.
Mitochondrion 10: 253–262

Fiedorczuk K & Sazanov LA (2018) Mammalian Mitochondrial Complex I Structure and
Disease-Causing Mutations. *Trends Cell Biol* 28: 835–867

Gella A, Prada-Dacasa P, Carrascal M, Urpi A, González-Torres M, Abian J, Sanz E &
Quintana A (2020) Mitochondrial Proteome of Affected Glutamatergic Neurons in a
Mouse Model of Leigh Syndrome. *Front Cell Dev Biol*

Grba DN, Blaza JN, Bridges HR, Agip ANA, Yin Z, Murai M, Miyoshi H & Hirst J (2022)
Cryo-electron microscopy reveals how acetogenins inhibit mitochondrial respiratory
complex I. *J Biol Chem* 298: 101602

Gu J, Liu T, Guo R, Zhang L & Yang M (2022) The coupling mechanism of mammalian
mitochondrial complex I. *Nat Struct Mol Biol* 29: 172–182

Guerrero-Castillo S, Baertling F, Kownatzki D, Wessels HJ, Arnold S, Brandt U & Nijtmans
L (2017) The Assembly Pathway of Mitochondrial Respiratory Chain Complex I. *Cell*

Metab 25: 128–139

Heide H, Bleier L, Steger M, Ackermann J, Dröse S, Schwamb B, Zörnig M, Reichert AS, Koch I, Wittig I, *et al* (2012) Complexome profiling identifies TMEM126B as a component of the mitochondrial complex I assembly complex. *Cell Metab* 16: 538–549

Hirst J, Carroll J, Fearnley IM, Shannon RJ & Walker JE (2003) The nuclear encoded subunits of complex I from bovine heart mitochondria. *Biochim Biophys Acta - Bioenerg* 1604: 135–150

Hoefs SJG, Dieteren CEJ, Rodenburg RJ, Naess K, Bruhn H, Wibom R, Wagena E, Willems PH, Smeitink JAM, Nijtmans LG, *et al* (2009) Baculovirus complementation restores a novel NDUFAF2 mutation causing complex I deficiency. *Hum Mutat*

Jain IH, Zazzeron L, Goldberger O, Marutani E, Wojtkiewicz GR, Ast T, Wang H, Schleifer G, Stepanova A, Brepoels K, *et al* (2019) Leigh Syndrome Mouse Model Can Be Rescued by Interventions that Normalize Brain Hyperoxia, but Not HIF Activation. *Cell Metab* 30: 824-832.e3

Jain IH, Zazzeron L, Goli R, Alexa K, Schatzman-Bone S, Dhillon H, Goldberger O, Peng J, Shalem O, Sanjana NE, *et al* (2016) Hypoxia as a therapy for mitochondrial disease. *Science (80-)* 352: 54–61

Jimenez-Blasco D, Busquets-Garcia A, Hebert-Chatelain E, Serrat R, Vicente-Gutierrez C, Ioannidou C, Gómez-Sotres P, Lopez-Fabuel I, Resch-Beusher M, Resel E, *et al* (2020) Glucose metabolism links astroglial mitochondria to cannabinoid effects. *Nature* 583: 603–608

Jones AJY & Hirst J (2013) A spectrophotometric coupled enzyme assay to measure the activity of succinate dehydrogenase. *Anal Biochem* 442: 19–23

Jumper J, Evans R, Pritzel A, Green T, Figurnov M, Ronneberger O, Tunyasuvunakool K, Bates R, Žídek A, Potapenko A, *et al* (2021) Highly accurate protein structure prediction with AlphaFold. *Nature* 596: 583–589

Kahlhöfer F, Gansen M & Zickermann V (2021) Accessory Subunits of the Matrix Arm of Mitochondrial Complex I with a Focus on Subunit NDUFS4 and Its Role in Complex I Function and Assembly. *Life* 11: 455

Kahlhöfer F, Kmita K, Wittig I, Zwicker K & Zickermann V (2017) Accessory subunit NUYM (NDUFS4) is required for stability of the electron input module and activity of mitochondrial complex I. *Biochim Biophys Acta - Bioenerg* 1858: 175–181

Kampjut D & Sazanov LA (2022) Structure of respiratory complex I – An emerging blueprint for the mechanism. *Curr Opin Struct Biol* 74: 102350

Koene S, Rodenburg RJ, van der Knaap MS, Willemsen MAAP, Sperl W, Laugel V, Ostergaard E, Tarnopolsky M, Martin MA, Nesbitt V, *et al* (2012) Natural disease course and genotype-phenotype correlations in Complex I deficiency caused by nuclear gene defects: what we learned from 130 cases. *J Inherit Metab Dis* 35: 737–747

Kolata P & Efremov RG (2021) Structure of Escherichia coli respiratory complex I reconstituted into lipid nanodiscs reveals an uncoupled conformation. *Elife* 10

Kotlyar AB & Vinogradov AD (1990) Slow active/inactive transition of the mitochondrial NADH-ubiquinone reductase. *Biochim Biophys Acta - Bioenerg* 1019: 151–158

Kravchuk V, Petrova O, Kampjut D, Wojciechowska-Bason A, Breese Z & Sazanov L (2022) A universal coupling mechanism of respiratory complex I. *Nature* 609: 808–814

Krishna S, Arrojo e Drigo R, Capitanio JS, Ramachandra R, Ellisman M & Hetzer MW (2021) Identification of long-lived proteins in the mitochondria reveals increased stability of the electron transport chain. *Dev Cell* 56: 2952-2965.e9

Kruse SE, Watt WC, Marcinek DJ, Kapur RP, Schenkman KA & Palmiter RD (2008) Mice with Mitochondrial Complex I Deficiency Develop a Fatal Encephalomyopathy. *Cell Metab* 7: 312–320

Kussmaul L & Hirst J (2006) The mechanism of superoxide production by NADH:ubiquinone oxidoreductase (complex I) from bovine heart mitochondria. *Proc Natl Acad Sci U S A* 103: 7607–7612

Lazarou M, McKenzie M, Ohtake A, Thorburn DR & Ryan MT (2007) Analysis of the Assembly Profiles for Mitochondrial- and Nuclear-DNA-Encoded Subunits into Complex I. *Mol Cell Biol* 27: 4228–4237

Masuda T, Tomita M & Ishihama Y (2008) Phase Transfer Surfactant-Aided Trypsin Digestion for Membrane Proteome Analysis. *J Proteome Res* 7: 731–740

McAndrew RP, Wang Y, Mohsen A-W, He M, Vockley J & Kim J-JP (2008) Structural Basis for Substrate Fatty Acyl Chain Specificity. *J Biol Chem* 283: 9435–9443

Meyerson JR, Rao P, Kumar J, Chittori S, Banerjee S, Pierson J, Mayer ML & Subramaniam S (2014) Self-assembled monolayers improve protein distribution on holey carbon cryo-EM supports. *Sci Rep* 4: 7084

Ortigoza-Escobar JD, Oyarzabal A, Montero R, Artuch R, Jou C, Jiménez C, Gort L, Briones P, Muchart J, López-Gallardo E, *et al* (2016) Ndufs4 related Leigh syndrome: A case report and review of the literature. *Mitochondrion* 28: 73–78

Ostergaard E, Rodenburg RJ, van den Brand M, Thomsen LL, Duno M, Batbayli M, Wibrand F & Nijtmans L (2011) Respiratory chain complex I deficiency due to NDUFA12 mutations as a new cause of leigh syndrome. *J Med Genet*

Padavannil A, Ayala-Hernandez MG, Castellanos-Silva EA & Letts JA (2022) The Mysterious Multitude: Structural Perspective on the Accessory Subunits of Respiratory Complex I. *Front Mol Biosci* 8: 798353

Parey K, Haapanen O, Sharma V, Köfeler H, Züllig T, Prinz S, Siegmund K, Wittig I, Mills DJ, Vonck J, *et al* (2019) High-resolution cryo-EM structures of respiratory complex I: Mechanism, assembly, and disease. *Sci Adv* 5: eaax9484

Pereira B, Videira A & Duarte M (2013) Novel Insights into the Role of Neurospora crassa NDUFAF2, an Evolutionarily Conserved Mitochondrial Complex I Assembly Factor. *Mol Cell Biol* 33: 2623–2634

Pettersen EF, Goddard TD, Huang CC, Couch GS, Greenblatt DM, Meng EC & Ferrin TE (2004) UCSF Chimera - A visualization system for exploratory research and analysis. *J Comput Chem* 25: 1605–1612

Punjani A, Rubinstein JL, Fleet DJ & Brubaker MA (2017) cryoSPARC: algorithms for rapid unsupervised cryo-EM structure determination. *Nat Methods* 14: 290–296

Rasmo D De, Palmisano G, Scacco S, Technikova-Dobrova Z, Panelli D, Cocco T, Sardanelli AM, Gnoni A, Micelli L, Trani A, *et al* (2010) Phosphorylation pattern of the NDUFS4 subunit of complex I of the mammalian respiratory chain. *Mitochondrion* 10: 464–471

Scacco S, Petruzzella V, Budde S, Vergari R, Tamborra R, Panelli D, van den Heuvel LP, Smeitink JA & Papa S (2003) Pathological Mutations of the Human NDUFS4 Gene of the 18-kDa (AQDQ) Subunit of Complex I Affect the Expression of the Protein and the Assembly and Function of the Complex. *J Biol Chem* 278: 44161–44167

Schägger H & von Jagow G (1991) Blue native electrophoresis for isolation of membrane protein complexes in enzymatically active form. *Anal Biochem* 199: 223–231

Scheres SHW (2012) RELION: Implementation of a Bayesian approach to cryo-EM structure determination. *J Struct Biol* 180: 519–530

Shevchenko A, Wilm M, Vorm O & Mann M (1996) Mass Spectrometric Sequencing of Proteins from Silver-Stained Polyacrylamide Gels. *Anal Chem* 68: 850–858

Signorile A, Sardanelli AM, Nuzzi R & Papa S (2002) Serine (threonine) phosphatase(s) acting on cAMP-dependent phosphoproteins in mammalian mitochondria. *FEBS Lett* 512: 91–94

Spiegel R, Shaag A, Mandel H, Reich D, Penyakov M, Hujeirat Y, Saada A, Elpeleg O &

Shalev SA (2009) Mutated NDUFS6 is the cause of fatal neonatal lactic acidemia in Caucasus Jews. *Eur J Hum Genet*

Stroud DA, Surgenor EE, Formosa LE, Reljic B, Frazier AE, Dibley MG, Osellame LD, Stait T, Beilharz TH, Thorburn DR, *et al* (2016) Accessory subunits are integral for assembly and function of human mitochondrial complex I. *Nature* 538: 123–126

Szczepanowska K, Senft K, Heidler J, Herholz M, Kukat A, Höhne MN, Hofsetz E, Becker C, Kaspar S, Giese H, *et al* (2020) A salvage pathway maintains highly functional respiratory complex I. *Nat Commun* 11: 1643

Ugalde C (2004) Differences in assembly or stability of complex I and other mitochondrial OXPHOS complexes in inherited complex I deficiency. *Hum Mol Genet* 13: 659–667

Valsecchi F, Grefte S, Roestenberg P, Joosten-Wagenaars J, Smeitink JAM, Willems PHGM & Koopman WJH (2013) Primary fibroblasts of NDUFS4−/− mice display increased ROS levels and aberrant mitochondrial morphology. *Mitochondrion* 13: 436–443

van de Wal MAE, Adjobo-Hermans MJW, Keijer J, Schirris TJJ, Homberg JR, Wieckowski MR, Grefte S, van Schothorst EM, van Karnebeek C, Quintana A, *et al* (2022) Ndufs4 knockout mouse models of Leigh syndrome: pathophysiology and intervention. *Brain* 145: 45–63

Wittig I, Karas M & Schägger H (2007) High Resolution Clear Native Electrophoresis for In-gel Functional Assays and Fluorescence Studies of Membrane Protein Complexes. *Mol Cell Proteomics* 6: 1215–1225

Yin Z, Burger N, Kula-Alwar D, Aksentijević D, Bridges HR, Prag HA, Grba DN, Viscomi C, James AM, Mottahedin A, *et al* (2021) Structural basis for a complex I mutation that blocks pathological ROS production. *Nat Commun* 12: 707

Yip CY, Harbour ME, Jayawardena K, Fearnley IM & Sazanov LA (2011) Evolution of respiratory complex I 'Supernumerary' subunits are present in the α-proteobacterial enzyme. *J Biol Chem* 286: 5023–5033

Zhang K (2016) Gctf: Real-time CTF determination and correction. *J Struct Biol* 193: 1–12

Zheng SQ, Palovcak E, Armache J-P, Verba KA, Cheng Y & Agard DA (2017) MotionCor2: anisotropic correction of beam-induced motion for improved cryo-electron microscopy. *Nat Methods* 14: 331–332

Zhu J, Vinothkumar KR & Hirst J (2016) Structure of mammalian respiratory complex i. *Nature* 536: 354–358

Zivanov J, Nakane T & Scheres SHW (2020) Estimation of high-order aberrations and anisotropic magnification from cryo-EM data sets in RELION-3.1. *IUCrJ* 7: 253–267

		PDB code	Subunits					
			N-module	NDUFS4	NDUFS6	NDUFA12	NDUFAF2	ACADVL
mouse heart	*ndufs4*[-/-] class 1	8C2S	+		+			+
	ndufs4[-/-] class 2	8CA3	+		+			
	ndufs4[-/-] class 3	8CA5	+				+	
	wild-type	6ZR2	+	+	+	+		
Y. lipolytica	*ndufs4*[-/-]	6RFS	+		+	+		
	ndufs6[-/-]	6RFQ	+	+			+	
	wild-type	6YJ4	+	+	+	+		

Table 1. Comparison of the subunit compositions of mouse and *Y. lipolytica* complex I models.

 Figure Legends

Figure 1. The NDUFS4 subunit in *wild-type* complex I from mouse-heart mitochondria. A) Overview
structures of complex I viewed from the membrane and from the heel showing the locations of NDUFS4 (red),
the N-module (blue) and subunits NDUFS6 (cyan), NDUFA12 (purple) and NDUFA7 (gold). B) View of the
front of the hydrophilic domain showing NDUFS4 (red) and the surrounding subunits. C) Subunits surrounding
NDUFA12 on the heel of the complex, and structural elements interacting with the β2-β3 loop of NDUFS4 that
passes underneath NDUFS1 to form a junction of NDUSF4, NDUFA12, NDUFA6 and NDUF9 (right, close-up
of the boxed region from a different viewpoint to focus on the loop). D) NDUFS4 approaches four of the seven
FeS clusters in the FeS chain between the FMN and the ubiquinone-binding site.

Figure 2. Characterization of *ndufs4*[-/-] mouse mitochondrial membranes. A) BN-PAGE analyses of heart
mitochondrial membranes solubilized by 1% DDM and stained with the in-gel NADH-linked NBT assay with
30 μg of protein loaded on each lane. B) Summary of the complexomic analyses of complex I subunits and
assembly factors in *wild-type* and *ndufs4*[-/-] heart mitochondrial membranes. The complete datasets are shown in
Figure EV1. C) The piericidin A-sensitive rates of NADH oxidation by *wild-type* and *ndufs4*[-/-] heart
mitochondrial membranes using 200 μM NADH. Data are mean averages ± SEM (n = 4 or 6) analyzed with the
Student's t-test (**** $p < 0.0001$). D) The atpenin A-sensitive rates of succinate oxidation using 5 mM
succinate. Data are mean averages ± SEM (n = 6) analyzed with the Student's t-test. In C and D, each point
represents an individual membrane preparation from more than four mice. E) Relative quantification of
respiratory complexes I to IV in kidney mitochondrial membranes of *ndufs4*[-/-] mice compared to *wild-type* mice,
calculated by normalizing the sum of MS peptide areas for the proteins in each complex to the sum of those for
VDAC1, VDAC2 and VDAC3. Data are mean averages ± SEM (n = 3 or 4). F) BN-PAGE analyses of *ndufs4*[-/-]
heart mitochondrial membranes solubilized by 1% DDM with and without cross-linking by BS[3] (0.1 mM for 25
min). then stained with NADH-linked NBT. N, N-module of complex I (~200 kDa); QP, QP subcomplex of
complex I (~800 kDa); SC, supercomplexes.

Figure 3. Characterization of purified *ndufs4*[-/-] complex I. Size exclusion chromatography of complex I from
A) kidney and B) heart mitochondrial membranes. The absorbance at 280 nm is in dark blue and at 420 nm in
red. The peak elution volume of *wild-type* complex I is marked with a in dashed line. The peak volumes for
complex I elution vary (~1.35 in A *vs.* ~1.6 mL in B) due to use of different columns (of the same type). C)
Rates of kinetic reactions for *ndufs4*[-/-] heart mitochondrial membranes as a proportion of the *wild-type* rates
(mean ± S.E.M., n ≥ 3). D) Rates of kinetic reactions for purified *ndufs4*[-/-] heart complex I as a proportion of the
wild-type rates (mean ± S.E.M., n ≥ 3). E) Loss of FMN from complex I purified from *wild-type* and *ndufs4*[-/-]
heart as the temperature is increased, measured as the fluorescence of the free flavin. F) The effect of BS[3] cross-
linking on BN-PAGE of the enzymes purified from heart; *wild-type* enzyme was not cross-linked (BS[3] 0);
ndufs4[-/-] complex I was cross-linked with 0.25 mM BS[3] during solubilization from the membranes, and then run
as prepared (BS[3] 1); *ndufs4*[-/-] complex I was reacted again with 2.5 mM BS[3] after size exclusion
chromatography (BS[3] 2). The second reaction further increased the stability of the enzyme (compare to the
untreated sample in Figure 2F).

Figure 4. Structural differences between *ndufs4⁻/⁻* heart complex I (class 2) and the *wild-type* enzyme. A) View of the active *wild-type* reference model (PDB: 6ZR2) and the *ndufs4⁻/⁻* heart class 2 model (PDB: 8CA3) showing the crevice left by the absence of NDUFS4. Red, NDUFS4; gold, NDUFA7; white, NDUFA9. B) Local resolutions of the consensus map of *ndufs4⁻/⁻* heart complex I (class 2, EMD-16516) and the locally refined N-module map (class 2, EMD-16517). C) Overlay of the PDB models for the *ndufs4⁻/⁻* (PDB: 8CA3) and *wild-type* (PDB: 6ZR2) enzymes. The *wild-type* model is shown in semi-transparent, and the rotation of the colored subunits of the N-module between the two enzymes is observed upon global model alignment. Green, NDUFV1; yellow, NDUFV3; purple, NDUFS1; blue, NDUFA2. D) Overlay of the PDB models for the *ndufs4⁻/⁻* (PDB: 8CA3) and *wild-type* (PDB: 6ZR2) enzymes focusing on the NDUFA9 subunit. The *wild-type* model is shown in white, and subunits of the *ndufs4⁻/⁻* enzyme are in color. With the models aligned on core subunit ND1, NDUFA9 drops slightly in the *ndufs4⁻/⁻* enzyme (see arrow). E) The regions of the active *wild-type* enzyme which appear disordered in *ndufs4⁻/⁻* maps.

Figure 5. NDUFAF2 binding in *ndufs4⁻/⁻* class 3 and comparison of the NDUFAF2- and NDUFS6-bound *ndufs4⁻/⁻* structures. A) Complex I models for the *wild-type* (left, PDB: 6ZR2) and *ndufs4⁻/⁻* classes 1 (PDB: 8C2S), 2 (PDB: 8CA3) and 3 (PDB: 8CA5) shown in surface view for the majority of the complex. Cartoon view is shown for subunits NDUFS6 (teal), NDUFA7 (gold), NDUFA9 (grey), NDUFA12 (purple), and NDUFAF2 (green). Protein compositions are also summarized in Table 1. B) Relationships between NDUFAF2 and surrounding subunits in class 3, and the equivalent view for class 2. An equivalent view for the wild-type complex is shown in Figure 1C. C) Comparison of the structures for NDUFA12 (from *wild-type* PDB: 6ZR2) and NDUFAF2 (from PDB: 8C2S (mouse) and PDB: 7RFQ (*Y. lipolytica*)) aligned to one another.

Figure 6. ACADVL association with purified *ndufs4⁻/⁻* heart complex I. A) Cryo-EM density map showing the location of the ACAD dimer (blue) attached to the detergent micelle (yellow) and next to subunit ND1 (purple). B) The model for the ACADVL dimer and cryoEM density (at a threshold of 1.74) is shown with the location of the FAD cofactors highlighted in yellow; the FAD density is shown separately also (right, at a threshold of 1.74). C) The dimeric model of ACADVL aligned to its position relative to complex I, within the detergent micelle. D) Relative quantification of ACAD9 and ACADVL. The sum of peptide areas for each protein were normalized to the sum of peptide areas for VDAC1, VDAC 2 and VDAC3, and displayed as the ratio between wild type and *ndufs4⁻/⁻* kidney mitochondrial membranes, mean and S.E.M (n = 3-4). E) The short Lys-Lys distances between ACADVL and NDUFA8 that may be responsible for stabilization of ACAD binding by crosslinking.

Figure 7. Schematic diagram showing our proposed assembly pathway for mouse *ndufs4⁻/⁻* complex I. NDUFAF2 contains two domains, connected by a flexible linker. NDUFS6 also contains two domains linked together; in the class 2 complex I structure only the N-terminal domain is bound and observed in the density map.

Expanded view Figure legends

Figure EV1. Complexome profiles from BN-PAGE analyses of solubilized mitochondrial membranes from A) *wild-type* and B) *ndufs4*-/- mouse heart. The heatmap represents the protein abundance, based on the relative peak areas of the three top scoring peptides in each gel slice for each protein from LC-MS analyses. The N-module subunits (red) are in the order NDUFV1, NDUFV2, NDUFV3, NDUFS1, NDUFS4, NDUFA2, NDUFS6, NDUFA12; subunits from the lower section of the hydrophilic domain (purple) are in the order NDUFS2, NDUFS3, NDUFS7, NDUFS8, NDUFA6, NDUFA5, NDUFA7, NDUFA9; subunits from the membrane domain (blue) are in the order ND1, ND2, ND4, ND5, NDUFS5, NDUFA1, NDUFA3, NDUFA8, NDUFA10, NDUFA11, NDUFA13, NDUFAB1, NDUFB1, NDUFB2, NDUFB3, NDUFB4, NDUFB5, NDUFB6, NDUFB7, NDUFB8, NDUFB9, NDUFB10, NDUFB11, NDUFC1, NDUFC2; assembly factors are in the order NDUFA2 and ACAD9 (green); NDUFAF1, NDUFAF3, NDUFAF4, NDUFAF5, ECSIT, TMEM126A and TMEM126B (yellow); and ACADVL is in orange. Where a protein was not detected in one sample, the column is completely black. Subunits ND3, ND4L and ND6 were not detected in either sample. The masses of complex I-related bands specific to *ndufs4*-/- membranes of ~800 (QP subcomplex) and ~200 kDa (N module) were estimated using the migration and masses of known complexes (inset).

Figure EV2. Flavin-site kinetics in *ndufs4*-/- mitochondrial membranes compared to *wild-type*. A) Kinetic evaluation of the Michaelis constant (K_M) for HAR in the NADH:HAR oxidoreductase reaction catalyzed by mouse heart membranes in the presence of 100 μM NADH. b) Kinetic evaluation of the Michaelis constant (K_M) for NADH in the presence of 1 mM HAR. C) Kinetic evaluation of the Michaelis constant (K_M) for NADH in the NADH:FeCN oxidoreductase reaction catalyzed by mouse kidney membranes in the presence of 100 μM FeCN. Due to the scarcity of materials, the assays in panels A and B were performed on different membrane samples, so the absolute rates are not directly comparable between them. Data points are mean averages ± SEM (n = 3).

Figure EV3. SDS PAGE of complex I purified from *ndufs4*-/- and *wild-type* heart mitochondrial membranes. Bands were manually excised, and proteins identified by MALDI-TOF-TOF analyses of tryptic digests. Complex I subunits are labelled in black, known assembly factors in blue, and other proteins detected in red. For proteins labelled in brackets individual peptide scores were below the 95% threshold but their protein scores were above the peptide threshold. Information on protein identification is in Tables S2 and S3.

Figure EV4. Views of the cryo-EM density map of complex I from *ndufs4*-/- kidney and comparison to the model for wild-type heart complex I. The wild-type complex I model (PDB:6ZR2) is docked into the *ndufs4*-/- complex I density map (grey, EMD:16514); no density is present in the locations of NDUFS4 (red model) and NDUFS6 (purple model).

Figure EV5. The cryo-EM densities for NDUFS6, NDUFA7, NDUFAF2 and NDUFA9. The models for *wild-type* heart complex I (PDB: 6ZR2, for NDUFS6, NDUFA7 and NDUFA9) and the class 3 model (for NDUFAF2) from were fitted into each density map using the fit-in-map function in Chimera X. The density within 3 Å of each fitted model is shown for A) subunit NDUFS6; B) subunit NDUFA7; C) assembly factor NDUFAF2; and D) subunit NDUFA9. They are shown for the map of *wild-type* heart complex I (EMD-11377);

1901   the *ndufs4*^{-/-} heart complex I class 1, 2 and 3 maps; and the *ndufs4*^{-/-} kidney complex I map, with map thresholds

1902   of 0.027, 1.76, 2.18, 1.52 and 1.12, respectively.

1903   **Suplementary Movie 1. Relative movement between the *wild-type* and *ndufs4*^{-/-} heart class 2 models, and**

1904   **motion within the cryo-EM density of *ndufs4*^{-/-} heart class 2 from 3DVA analyses.**

1905

1906

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Ndufs4$^{-/-}$ 800 kDa intermediate QP subcomplex

N-terminal domain of NDUFAF2 binds to QP subcomplex

NDUFAF2

C-terminal domain of NDUFAF2 (attached by flexible linker) captures the N-module

N-module

NDUFAF2 docks the N-module onto the QP subcomplex

Ndufs4$^{-/-}$ heart complex I class 3

NDUFAF2

N-terminal domain of NDUFS6 displaces NDUFAF2 and binds to the nascent complex

NDUFS6

Ndufs4$^{-/-}$ heart complex I class 2

Absence of NDUFA12 precludes binding of the C-terminal domain of NDUFS6 and completion of assembly

Figure 7

A

B

Figure EV1

Figure EV2

Figure EV3

Figure EV4

Figure EV5