Title: Natural disaster alters the adaptive benefits of sociality

Authors: C. Testard[1]*, C. Shergold[2], A. Acevedo-Ithier[1], J. Hart[2], JE. Negron-Del Valle[3,4], D. Philips[3,4], MM. Watowich[3,4,5], JI. Sanguinetti-Scheck[6], MJ. Montague[1], N. Snyder-Mackler[3,4], JP. Higham[7,8], ML. Platt[1,9,10], LJN. Brent[2]
*Corresponding author. Email: camille.testard@pennmedicine.upenn.edu

Affiliations:
1. Department of Neuroscience, University of Pennsylvania, Philadelphia, PA, USA
2. Centre for Research in Animal Behaviour, University of Exeter, United Kingdom
3. Center for Evolution and Medicine, Arizona State University, Tempe, AZ, USA.
4. School of Life Sciences, Arizona State University, Tempe, AZ, USA.
5. Department of Biological Sciences, Vanderbilt University, Nashville, TN, USA.
6. Department of Organismic and Evolutionary Biology, Harvard University, MA, USA.
7. Department of Anthropology, New York University, New York, NY, USA.
8. New York Consortium in Evolutionary Primatology, NYCEP, New York, NY, USA.
9. Department of Psychology, University of Pennsylvania, Philadelphia, PA, USA
10. Marketing Department, University of Pennsylvania, Philadelphia, PA, USA

Abstract:
Global warming intensifies weather-related disasters that radically alter ecosystems. How animals respond to these abrupt and lasting ecological changes remains poorly understood. Here, we leveraged an unprecedented study of rhesus macaques from 5 years before to 5 years after a category 4 hurricane, which destroyed 63% of the island's vegetation and exacerbated monkeys' exposure to intense heat. In response, macaques dramatically and persistently increased tolerance for other monkeys. Social tolerance predicted individual survival after the hurricane–but not before it–by facilitating access to now scarce shade critical for lowering body temperature. We demonstrate that an extreme climatic event altered the adaptive benefits of sociality and triggered substantial and persistent changes in social structure of group-living animals–with consequences for the resilience of populations living in degraded ecosystems.

One-Sentence Summary: *Testard et al.* show that natural disasters alter the adaptive benefits of sociality triggering persistent behavioral changes in animals.

MAIN TEXT

With the intensifying climate crisis, extreme weather events are expected to become less predictable and increase in both frequency and force (*1*, *2*). Weather-related natural disasters can cause widespread and sustained destruction of the natural landscape, deplete resources, and degrade infrastructure upon which humans and other animals depend (*3*). Disaster-induced ecosystem modifications may then alter the selection pressures exerted on individuals living

within them, enforcing changes in their behavior (*4*). Flexibility in social behavior has been proposed as an adaptive trait for animals living in unstable environments such as those impacted by extreme climatic events (*5, 6*). This idea has remained largely untested due to the difficulty of studying behavioral responses to environmental instability, and thus the flexibility and adaptive value of sociality in these contexts, remain open questions (*7*). Moreover, due to the highly unpredictable nature of extreme weather events and the impossibility of replicating these conditions in the laboratory, how and to what extent animals respond to abrupt and lasting ecosystem degradation remains largely unknown (*4, 7*).

On September 20th 2017, Hurricane Maria made landfall as a Category 4 hurricane on Puerto Rico, causing devastating physical destruction and a human death toll reaching the thousands, one of the deadliest hurricanes in Caribbean history (*8*). A population of rhesus macaques living on a small island off the coast of Puerto Rico, Cayo Santiago, was also hit by this storm (**Figure 1A**). Hurricane Maria destroyed 63% of Cayo Santiago's vegetation, along with nearly all existing research infrastructure (**Figure 1B-C**). Five years later (Spring 2022), the canopy cover remains far below pre-hurricane levels (**Figure 1B, S1A**), and surface temperatures on the island regularly exceed 40ºC outside of shaded areas (**Figure S1D**). Hurricane-induced deforestation resulted in the majority of the island being exposed to these high temperatures. From 2018 to 2021, 48% of sample points from temperature loggers deployed on exposed (i.e., non-shaded) parts of the island detected temperatures above 40°C during the day (10am - 5pm) (N=190,240 sample points from 25 temperature loggers, see **Methods**, **Figure S1D**). Such extreme temperatures compromise the body's ability to regulate its internal temperature, and can result in heat cramps, heat exhaustion, heatstroke, and hyperthermia (*9, 10*). Primates have limited capacity for physiological thermoregulation (*11, 12*) and therefore often cope with extreme temperatures by changing their behavior (*13–15*). Cayo Santiago's macaques address thermoregulatory challenges posed by the hot Caribbean sun primarily by accessing shade provided by trees (on average 7°C cooler than exposed areas, **Figure 1D**).

Previously, we reported that the macaques on Cayo Santiago dramatically increased their tolerance for other monkeys in the year following Hurricane Maria (*16*). We hypothesized social tolerance helped monkeys cope with the drastic hurricane-induced devegetation which exacerbated heat stress. Here, we test the persistence and adaptive function of this social response in a dramatically altered landscape, using the richest behavioral and demographic dataset of a non-human animal population available to date, from before and after a natural disaster.

Figure 1. Hurricane Maria led to a persistent drop in vegetation and a concomitant increase in surface temperature. (**A**) Left: Path of Hurricane Maria making landfall on Puerto Rico, including Cayo Santiago, on September 20th 2017. VIIRS image captured by the National Oceanic and Atmospheric Administration's Suomi NPP satellite. (**B**) Google Earth Images taken at three time points: 7 months before Maria (left), 2 weeks after Maria (middle) and 4.5 years after (right). Tree census conducted in April 2023 revealed only 167 trees were left on the upper part of "Big Cayo" (see **Figure S1** for satellite images from more time points and tree census on the rest of Cayo Santiago). (C) Foliage cover from Cayo Santiago Island, as measured by greenness from satellite imagery (NDVI), decreased by 63% in the 3 months following Hurricane Maria (thick gray bar). Note that NDVI index captures all types of vegetation, including grass and small bushes. Although satellite NDVI increases steadily up until 2022, this largely reflects grass (**Figure S1**) which does not provide shade. Even when including all vegetation types, vegetation cover still remains far below pre-storm levels. (**D**) Distribution of temperatures post-hurricane (2018-2022) in three types of terrain: *Exposed* - areas without vegetation cover both before and after Hurricane Maria; *De-vegetated* - areas which lost their vegetation cover after Hurricane Maria; and *Vegetated* - areas with vegetation cover. (**E**) Social network data used in this study were collected from 5 years before Maria to 5 years after, across 7 social groups.

Social tolerance is higher for 5 years after the hurricane

To assess the long-term effects of Hurricane Maria on the Cayo Santiago macaques' rate of social interaction, we used behavioral data collected during a 10 year period from 5 years before Hurricane Maria through 5 years after (pre-hurricane years: 2013-2017; post-hurricane: 2018-2022, excluding 2020 due to the Covid19 pandemic, **Figure 1E**) in 7 distinct social groups. A total of 790 unique adult individuals with behavioral data before and/or after Hurricane Maria were included in this analysis (i.e. individuals did not need to have data collected both before and

after Maria to be considered, as in (*16*)). We focused our analyses on social tolerance, operationalized as the propensity to be in close proximity to conspecifics (i.e., within 2m of another adult monkey (*17*)), but see **Figure S2** for results on grooming and aggressive interactions.

To establish if changes in proximity interactions persisted over time, we constructed social networks for each group in each year from 2013 to 2022 and compared individual propensity to be in proximity to a conspecific before and after the hurricane (details in **Methods**). We used a linear mixed model to compare proximity in each post-hurricane year to pre-hurricane years, controlling for potential confounding variables such as group size, individual age, and sex (see **Methods**). Our models accounted for uncertainty in edge weights due to unequal sampling across individuals, groups and years (see **Methods,** (*18*)). We first report changes in proximity in the year immediately following Maria compared to before Maria, which measures the acute response, and next quantify changes 2-5 years after the storm, which measures the persistence of the response.

In the year following Hurricane Maria, a macaque's propensity to be in proximity to another macaque was on average 1.71 standard deviations higher–more than triple–compared to pre-hurricane years (95% CI = [1.60 - 1.83], **Figure 2C**; mean pre-hurricane was 0.18 compared to 0.58 in 2018). Two-to-five years after the storm, proximity amongst macaques remained between 0.47 and 0.84 standard deviations higher than pre-hurricane levels ($P<0.001$ for all years post-hurricane, **Figure 2C**). Thus, increased social tolerance levels persisted for 5 years after Hurricane Maria, peaking in the year immediately following the disaster ($P<0.001$, **Figure 2**, **S2**). Population density remained relatively stable over the 10 years analyzed and therefore cannot explain these results (mean total population size = 1498 in pre-hurricane years; mean = 1491.2 in post-hurricane years). Evidence of higher tolerance is corroborated by reduced aggression in the two to five years after the disaster (**Figure S2**). Higher tolerance and lower aggression were not associated with consistently higher frequencies of grooming in the five years following the hurricane (**Figure S2**).

Social tolerance after Hurricane Maria, but not before, predicts survival
Next, we investigated whether macaques' higher tolerance predicted their survival following Hurricane Maria (study period: September 2017 - August 2022). We tested the effect of two individual-level network metrics from the proximity network: degree (i.e., the number of proximity partners) and strength (i.e., the propensity to be in proximity to others). This analysis included 430 unique adult individuals from 5 social groups, of which 56 died during the study period. Because the values of network metrics changed over the study period, we used time-varying mixed effect cox-proportional hazard models to evaluate whether proximity predicted survival from one year to the next during our study period (**Figure 2D**). All models

implicitly accounted for age (see **Methods**), sex was included as a fixed effect and social group, individual ID and year as random effects.

We found that both proximity degree and strength predicted survival following the hurricane, suggesting the behavioral responses were adaptive (**Figure 2E**). Each additional standard deviation in an individual's number of proximity partners (+12.7 partners) resulted in a 42.69% decrease in mortality risk (Model estimate = -0.54, Hazard Ratio = 0.58, 95%CI = [0.40 0.86]) and an additional standard deviation in the propensity to be in proximity to another macaque (i.e., proximity strength) led to 37.3% decrease in mortality risk (Model estimate = -0.48, Hazard Ratio = 0.62, 95%CI = [0.41 0.95]).

To test whether the adaptive benefits of proximity after the hurricane were specifically due to the storm and its aftermath, we tested whether proximity also predicted survival before Hurricane Maria. Using the same modeling approach but focused on *pre*-hurricane years (study period: Jan. 2013 - Sept. 2017; N = 616 individuals, 65 deaths, see **Methods**), we found no evidence that proximity strength or degree predicted survival (**Figure 2E**). These findings indicate that, by drastically modifying Cayo Santiago's ecosystem, Hurricane Maria altered the adaptive benefits of tolerating conspecifics in close proximity.

Figure 2. Cayo Santiago macaques exhibit a persistent increase in social tolerance after Hurricane Maria, which predicts survival. (**A**) Proximity networks for social group V from 2013 to 2022, excluding 2020 due to the COVID19 crisis are shown as example networks. Each node represents an individual and edges represent the probability for two individuals to be in proximity to one another. The thicker the line the higher the probability of proximity. Red dotted line marks Hurricane Maria. (**B**) Probability of being observed in proximity to another monkey from 2013 to 2022, across 790 individuals in 6 social groups, 24 group-year combinations. (**C**) Linear mixed model estimates for the long-term change in standardized proximity network strength compared to pre-hurricane levels (for grooming and aggression results see **Figure S2**; **Table S1**). Error bars represent the 95% confidence interval (CI), pooled across 100 network iterations (distribution of possible networks given the observed interaction data, see **Methods**). Red: increase relative to pre-hurricane years. (**D**) Graphical explanation of cox-proportional hazard models used to test the effect of sociality before and after the hurricane on survival. HR: Hazard ratio. (**E**) Estimates from survival models for the impact of degree and strength in the proximity networks on mortality risk. Estimates <1: lower mortality risk, >1: higher mortality. Hazard ratio = e^estimate. Error bars represent the 95% CI pooled across bayesian-generated social networks (see **Methods**). Green: pre-hurricane. Brown: Post-hurricane. n.s.: not significant.

Stronger thermoregulatory pressure post-hurricane may drive the adaptive benefits of social tolerance

The dramatic drop in shade brought by Hurricane Maria elevated thermoregulatory burdens on Cayo Santigo's macaques. Temperatures in non-shaded areas of the island regularly exceed 40ºC (**Figure S1D**) and persistent exposure to these temperatures increases the risk of severe heat shock and generally degrades health and physical condition (*9*). Anecdotal reports from observers on the ground suggested that macaques concentrated in shady areas afforded by the remaining vegetation following Maria (**Figure 3A**). From these initial observations we hypothesized that individuals' increased proximity to conspecifics after the hurricane reflected, in part, attempts to access scarce shade to deal with the intensified thermoregulatory burden. If increased proximity to conspecifics following Hurricane Maria reflects adaptation to stronger thermoregulatory pressure we predicted, first, that we would find higher proximity between macaques during hotter hours of the day compared to cooler hours and, second, a stronger relationship between proximity to others and survival during during hotter hours of the day than during cooler daytime hours. We found that differences in proximity pre-to-post hurricane were almost twice as strong in the afternoon, when temperatures were highest (**Figure 3B**), compared to cooler morning hours (0.92 SD higher proximity strength post-hurricane for afternoon hours compared to 0.58 SD higher in morning hours, **Figure 3C**). We also found that, after Maria, proximity to conspecifics in the afternoon better predicted survival than proximity in the morning (**Figure 3D**), consistent with the hypothesis that increased tolerance after the hurricane reflects an adaptive behavioral response to increased thermoregulatory stress.

Figure 3. Thermoregulatory needs drive macaques to become more tolerant after Hurricane Maria. (A) Photo taken two years after the storm showcasing Cayo Santiago macaques concentrated in a slim shady area behind a dead tree. (B) Temperature (°C) in exposed vs. shaded areas in the morning (left) vs. afternoon (right). (C) Model estimates comparing differences in standardized strength in the proximity network pre-to-post hurricane, considering morning (6am-10am) or afternoon hours (10am-3pm when data collection stops). Error bars represent 95% confidence interval (CI). (D) Proximity network measures from data collected after the hurricane in the afternoon (orange) better predict survival than from data collected in the morning (green). N.s.: not significant. ***: *P*<0.001. Bars represent 95% CI.

DISCUSSION

We found that monkeys were persistently more tolerant of others in their vicinity for 5 years after Hurricane Maria. Social tolerance predicted individual survival after the storm, especially during the hottest hours of the day, supporting our hypothesis that the adaptive benefits of social tolerance are linked to behavioral thermoregulation in response to increased heat-stress. Importantly, social tolerance did not predict survival before the hurricane, demonstrating that hurricane-induced changes in ecological pressures altered the benefits individuals gain from social relationships, leading to flexible adjustments in how they interact with others.

Social tolerance following the hurricane was associated with a 42% decrease in mortality risk, an effect comparable to the striking survival impacts of social support in humans (*19*). If fitness-enhancing tolerance is heritable and the environmental conditions favoring this phenotype do not change, hurricane-induced ecological degradation may select for more tolerant individuals, leading to an overall more tolerant macaque society on Cayo Santiago in the future. However, the adaptive benefits of increased tolerance may change dynamically over time as new challenges emerge. For example, persistent increases in proximity or close contact may increase disease transmission (*20*), changing the balance of the ultimate costs and benefits of social tolerance. The ongoing behavioral and demographic study of this population will provide unprecedented opportunities to investigate these possibilities.

Our findings, in conjunction with previous work on this population (*16*), demonstrate that rhesus monkeys display considerable social flexibility in response to environmental disruption and that social connections, rather than being static, are dynamically renegotiated to adjust to current demands. The sudden and persistent vegetation drop induced by Hurricane Maria increased competition for shade, which could have provoked higher aggression and greater mortality, but instead elicited higher social tolerance – a remarkable change in social behavior given that rhesus macaques are widely considered to be aggressive and strongly hierarchical primates (*21–23*). Previous research shows how group-living animals use their social connections to cope with environmental challenges (*24–28*), but whether they could alter their patterns of social interactions when the challenges they face change abruptly was less clear (*29, 30*). Our study provides rare evidence of social flexibility in an animal population, which promoted resilience to an unpredictable, extreme, and persistently impactful climatic event. By increasing the frequency of such events, climate change may ultimately select for species with flexible social systems capable of facultative adjustments to rapidly-changing environments (*6, 31*).

Increased social tolerance peaked in the year immediately following the hurricane (**Figure 2C**), despite the lack of recovery in vegetative cover in the ensuing years (only a few hundred trees were left on the main part of the island in 2023, **Figure 1B-S1** & **Methods** for tree census details). The social response to the disaster thus included both an acute and a persistent

component, indicating partial decoupling from changes in the environment. The acute component might reflect stronger social cohesion, defined here as a willingness to be in proximity to others and cooperate rather than being defensive or aggressive, following a traumatic event, which is often observed in humans following both natural disasters (*32*) and terrorist attacks (*33*). When a highly unpredictable shock to the system has occurred, animals might hedge their bets in terms of which social connections might end up being useful. Since who will be helpful and when is uncertain, individuals may be motivated to be tolerant of more individuals. As the environment stabilizes around a "new normal", less useful social connections might be pruned as a result. Yet another possibility is that the effect on sociality in the year immediately following Maria may have been stronger because, after this point, macaques gradually habituated to the greater temperature through physiological changes. However, this explanation seems unlikely given the limited capacity for physiological thermoregulation in primates (*11*, *12*).

As temperatures rise due to global warming, heat waves are becoming more frequent and intense (*34*), increasing thermoregulatory burdens on animal populations worldwide, including humans. These observations foreshadow future changes in social behavior to deal with heat stress, particularly in species with a limited capacity for physiological thermoregulation (*35*). These social changes can lead to increased epidemic risk, representing compound threats to animal and human health, even years after the disaster itself (*36*). Understanding how animals cope with increased heat and other environmental consequences of climate change, and how these responses shape both intra- and inter-specific interactions, will be important for predicting, and potentially mitigating, the impacts of climate change on ecosystems.

Using the richest behavioral dataset of a non-human animal population affected by a natural disaster, we found that extreme climatic events can alter the adaptive benefits of sociality and trigger persistent changes in social structure. Furthermore, we provide rare evidence that social flexibility promotes resilience in the aftermath of a category 4 hurricane. Our findings provide an unprecedented window into how behavioral responses to heat, which is intensifying with global warming, influence social interactions–with potential consequences for social transmission of information, genes and disease.

Acknowledgments: We thank the Caribbean Primate Research Center staff for their important roles in data collection and in the maintenance of Cayo Santiago. We also thank Nacho Sanguinetti, Greg Albery, Erin Siracusa, and Delphine de Moor for insightful comments that enriched our analyses and discussion of the results.

Funding: Support for this research was provided by the National Institutes of Health (R01MH118203, U01MH121260, R01MH096875, P40OD012217, R01AG060931, R00AG051764, R56AG071023), a RAPID award from the National Science Foundation (1800558), a Royal Society Research Grant (RGS/R1/191182) and a European Research Council Consolidator Grant (864461 - FriendOrigins). Cayo Santiago is supported by grant 8-P40

OD012217-25 from the National Center for Research Resources (NCRR) and the Office of Research Infrastructure Programs (ORIP) of the National Institutes of Health.

Author contributions: CT and LJNB conceptualized the study; JAH developed methodology specific to this study; CT, AA and CS analyzed the data with input from LJNB; JN, DP and MW collected the data; NSM, JH, MM, MLP, LJNB secured funding for this work; CT and LJNB drafted the manuscript with input from all authors; LJNB supervised this work. All authors read and approved the manuscript.

Competing interests: MLP is a scientific advisory board member, consultant, and/or co-founder of Blue Horizons International, NeuroFlow, Amplio, Cogwear Technologies, Burgeon Labs, and Glassview, and receives research funding from AIIR Consulting, the SEB Group, Mars Inc, Slalom Inc, the Lefkort Family Research Foundation, Sisu Capital, and Benjamin Franklin Technology Partners. All other authors declare no competing interests.

Data and materials availability: Upon publication, data will be available on Open Science Framework platform and code GitHub at https://github.com/camilletestard/Cayo-Maria-Survival

Supplementary materials
Materials and Methods
Supplementary text
Figs. S1 to S2
Table S1
References (37-45)

REFERENCES

1. K. Balaguru, G. R. Foltz, L. R. Leung, K. A. Emanuel, Global warming-induced upper-ocean freshening and the intensification of super typhoons. *Nat. Commun.* **7**, 13670 (2016).

2. K. Emanuel, Will Global Warming Make Hurricane Forecasting More Difficult? *Bull. Am. Meteorol. Soc.* **98**, 495–501 (2017).

3. S. R. Weiskopf, M. A. Rubenstein, L. G. Crozier, S. Gaichas, R. Griffis, J. E. Halofsky, K. J. W. Hyde, T. L. Morelli, J. T. Morisette, R. C. Muñoz, A. J. Pershing, D. L. Peterson, R. Poudel, M. D. Staudinger, A. E. Sutton-Grier, L. Thompson, J. Vose, J. F. Weltzin, K. P. Whyte, Climate change effects on biodiversity, ecosystems, ecosystem services, and natural resource management in the United States. *Sci. Total Environ.* **733**, 137782 (2020).

4. D. N. Fisher, R. J. Kilgour, E. R. Siracusa, J. R. Foote, E. A. Hobson, P.-O. Montiglio, J. B. Saltz, T. W. Wey, E. W. Wice, Anticipated effects of abiotic environmental change on intraspecific social interactions. *Biol. Rev. Camb. Philos. Soc.* **96**, 2661–2693 (2021).

5. C. Schradin, N. Pillay, C. Bertelsmeier, Social flexibility and environmental unpredictability in African striped mice. *Behav. Ecol. Sociobiol.* **73**, 1–12 (2019).

6. E. A. Beever, L. E. Hall, J. Varner, A. E. Loosen, J. B. Dunham, M. K. Gahl, F. A. Smith, J. J. Lawler, Behavioral flexibility as a mechanism for coping with climate change. *Front. Ecol. Environ.* **15**, 299–308 (2017).

7. D. T. Blumstein, L. D. Hayes, N. Pinter-Wollman, Social consequences of rapid environmental change. *Trends Ecol. Evol.* **38**, 337–345 (2023).

8. D. D. Kane, Mortality in Puerto Rico after Hurricane Maria. *N. Engl. J. Med.* **379** (2018), p. e30.

9. W. G. Whitford, R. E. Sherman, Sweating responses in the rhesus monkey (Macaca mulatta). *Comp. Biochem. Physiol. A Physiol.* **53**, 337–341 (1976).

10. K. L. Ebi, A. Capon, P. Berry, C. Broderick, R. de Dear, G. Havenith, Y. Honda, R. S. Kovats, W. Ma, A. Malik, N. B. Morris, L. Nybo, S. I. Seneviratne, J. Vanos, O. Jay, Hot weather and heat extremes: health risks. *Lancet.* **398**, 698–708 (2021).

11. Y. G. Kamberov, S. M. Guhan, A. DeMarchis, J. Jiang, S. S. Wright, B. A. Morgan, P. C. Sabeti, C. J. Tabin, D. E. Lieberman, Comparative evidence for the independent evolution of hair and sweat gland traits in primates. *J. Hum. Evol.* **125**, 99–105 (2018).

12. A. Best, J. M. Kamilar, The evolution of eccrine sweat glands in human and nonhuman primates. *J. Hum. Evol.* **117**, 33–43 (2018).

13. R. McFarland, L. Barrett, M.-A. Costello, A. Fuller, R. S. Hetem, S. K. Maloney, D. Mitchell, P. S. Henzi, Keeping cool in the heat: Behavioral thermoregulation and body temperature patterns in wild vervet monkeys. *Am. J. Phys. Anthropol.* **171**, 407–418 (2020).

14. R. McFarland, B. Majolo, Coping with the cold: predictors of survival in wild Barbary macaques, Macaca sylvanus. *Biol. Lett.* **9**, 20130428 (2013).

15. G. Schino, A. Troisi, Behavioral thermoregulation in long-tailed macaques: effect on social preference. *Physiol. Behav.* **47**, 1125–1128 (1990).

16. C. Testard, S. M. Larson, M. M. Watowich, C. H. Kaplinsky, A. Bernau, M. Faulder, H. H. Marshall, J. Lehmann, A. Ruiz-Lambides, J. P. Higham, M. J. Montague, N. Snyder-Mackler, M. L. Platt, L. J. N. Brent, Rhesus macaques build new social connections after a natural disaster. *Curr. Biol.* **31**, 2299–2309.e7 (2021).

17. S. E. DeTroy, D. B. M. Haun, E. J. C. van Leeuwen, What isn't social tolerance? The past, present, and possible future of an overused term in the field of primatology. *Evol. Anthropol.* **31**, 30–44 (2022).

18. J. D. A. Hart, M. N. Weiss, D. W. Franks, L. J. N. Brent, BISoN: A Bayesian Framework for Inference of Social Networks. *bioRxiv* (2022), p. 2021.12.20.473541.

19. J. Holt-Lunstad, T. B. Smith, J. B. Layton, Social Relationships and Mortality Risk: A Meta-analytic Review. *PLoS Med.* **7**, e1000316 (2010).

20. C. J. Carlson, G. F. Albery, C. Merow, C. H. Trisos, C. M. Zipfel, E. A. Eskew, K. J. Olival, N. Ross, S. Bansal, Climate change increases cross-species viral transmission risk. *Nature*. **607**, 555–562 (2022).

21. B. Thierry, A. N. Iwaniuk, S. M. Pellis, The influence of phylogeny on the social behaviour of macaques (primates: Cercopithecidae, genus Macaca). *Ethology*. **106**, 713–728 (2000).

22. D. Maestripieri, C. L. Hoffman, "Behavior and social dynamics of rhesus macaques on Cayo Santiago" in *Bones, Genetics, and Behavior of Rhesus Macaques: Macaca Mulatta of Cayo Santiago and Beyond*, Q. Wang, Ed. (Springer, New York, 2012), pp. 247–262.

23. E. B. Cooper, L. J. N. Brent, N. Snyder-Mackler, M. Singh, A. Sengupta, S. Khatiwada, S. Malaivijitnond, Z. Qi Hai, J. P. Higham, The rhesus macaque as a success story of the Anthropocene. *Elife*. **11**, e78169 (2022).

24. S. Cohen, T. A. Wills, Stress, social support, and the buffering hypothesis. *Psychol. Bull.* **98**, 310–357 (1985).

25. J. B. Silk, The adaptive value of sociality in mammalian groups. *Philos. Trans. R. Soc. Lond. B Biol. Sci.* **362**, 539–559 (2007).

26. R. McFarland, D. Murphy, D. Lusseau, S. P. Henzi, J. L. Parker, T. V. Pollet, L. Barrett, The "strength of weak ties" among female baboons: fitness-related benefits of social bonds. *Anim. Behav.* **126**, 101–106 (2017).

27. S. Ellis, N. Snyder-Mackler, A. Ruiz-Lambides, M. L. Platt, L. J. N. Brent, Deconstructing sociality: the types of social connections that predict longevity in a group-living primate. *Proc. Biol. Sci.* **286**, 20191991 (2019).

28. N. Snyder-Mackler, J. R. Burger, L. Gaydosh, D. W. Belsky, G. A. Noppert, F. A. Campos, A. Bartolomucci, Y. C. Yang, A. E. Aiello, A. O'Rand, K. M. Harris, C. A. Shively, S. C. Alberts, J. Tung, Social determinants of health and survival in humans and other animals. *Science* (2020), doi:10.1126/science.aax9553.

29. P. M. Kappeler, L. Barrett, D. T. Blumstein, T. H. Clutton-Brock, Constraints and flexibility in mammalian social behaviour: introduction and synthesis. *Philos. Trans. R. Soc. Lond. B Biol. Sci.* **368**, 20120337 (2013).

30. B. Thierry, Where do we stand with the covariation framework in primate societies? *Am. J. Biol. Anthropol.* **178 Suppl 74**, 5–25 (2022).

31. C. Parmesan, Ecological and Evolutionary Responses to Recent Climate Change. *Annu. Rev. Ecol. Evol. Syst.* **37**, 637–669 (2006).

32. S. Sweet, The Effect of a Natural Disaster on Social Cohesion: A Longitudinal Study. *Int. J. Mass Emerg. Disasters*. **16**, 321–331 (1998).

33. G. Dezecache, J.-R. Martin, C. Tessier, L. Safra, V. Pitron, P. Nuss, J. Grèzes, Nature and

determinants of social actions during a mass shooting. *PLoS One*. **16**, e0260392 (2021).

34. J. H. Stillman, Heat Waves, the New Normal: Summertime Temperature Extremes Will Impact Animals, Ecosystems, and Human Communities. *Physiology* . **34**, 86–100 (2019).

35. J. M. Alston, M. J. Joyce, J. A. Merkle, R. A. Moen, Temperature shapes movement and habitat selection by a heat-sensitive ungulate. *Landsc. Ecol.* **35**, 1961–1973 (2020).

36. Alba Motes-Rodrigo, Gregory F. Albery, Josue E. Negron-Del Valle, Daniel Philips, Cayo Biobank Research Unit, Michael L. Platt, Lauren J. N. Brent, Camille Testard, Social responses to a hurricane drive greater epidemic risk. *Submitted* (2023).

37. J. Altmann, Observational study of behavior: sampling methods. *Behaviour*. **49**, 227–267 (1974).

38. N. P. Robinson, B. W. Allred, M. O. Jones, A. Moreno, J. S. Kimball, D. E. Naugle, T. A. Erickson, A. D. Richardson, A Dynamic Landsat Derived Normalized Difference Vegetation Index (NDVI) Product for the Conterminous United States. *Remote Sensing*. **9**, 863 (2017).

39. J. M. Setchell, *Field and Laboratory Methods in Primatology: A Practical Guide* (Cambridge University Press, Cambridge, 2 Edition., 2011).

40. H. Whitehead, *Analyzing Animal Societies: Quantitative Methods for Vertebrate Social Analysis* (University of Chicago Press, 2008).

41. J. Krause, R. James, D. W. Franks, D. P. Croft, *Animal Social Networks* (Oxford University Press, 2015).

42. D. Bates, M. Mächler, B. Bolker, S. Walker, Fitting Linear Mixed-Effects Models Using lme4. *Journal of Statistical Software*. **67** (2015), pp. 1–48.

43. D. Lüdecke, M. S. Ben-Shachar, I. Patil, P. Waggoner, D. Makowski, performance: An R Package for Assessment, Comparison and Testing of Statistical Models. *Journal of Open Source Software*. **6** (2021), p. 3139.

44. T. M. Therneau, coxme: mixed effects Cox models. *R package version*.

45. T. M. Therneau, "A Package for Survival Analysis in R" (2023), (available at https://CRAN.R-project.org/package=survival).

Supplementary Materials

Materials and Methods:

1. Experimental model and subject details

We studied a population of rhesus macaques living in a semi free-ranging colony on Cayo Santiago, Puerto Rico (1809 N, 65 44 W). The colony has been almost continuously monitored since it was established in 1938 following the release of 409 animals originally captured in India. Cayo Santiago is managed by the Caribbean Primate Research Center (CPRC), which supplies food to the population daily and water *ad libitum*. Animals are fed once a day in the morning between 7 and 9am. There is no contraceptive use and no medical intervention aside from tetanus inoculation when animals are weaned yearlings. Animals are free to aggregate into social units as they would in the wild. There are no natural predators on the island.

Subjects for this study were adult males and females (at least 6 years old), individually recognizable by tattoos, ear notches, and facial features. We used 7 groups, TT, F, KK, V, R, HH and S, for which there was behavioral data either before or after Hurricane Maria, between 2013 and 2022. Our dataset included 790 unique adult individuals (see **Table S1** for more details per group and year). These groups had home ranges on different parts of the island. Group V ranged alone on ''small Cayo''; an area whose land bridge was severed from the main part of island (''big Cayo'') after the hurricane. In contrast, Group TT, KK, R, HH, S and F ranged on big Cayo with all other groups. We used multiple available years of observational data (HH: 2014, 2016; R: 2015, 2016; F: 2013, 2014, 2015, 2016, 2017; KK: 2013, 2015, 2017, 2018; V: 2015, 2016, 2017) to characterize social behavior before the hurricane (''pre-hurricane''). Post-hurricane, our sample includes observational data from 2018, 2019, 2021 and 2022 (S: 2019; TT:2022; F:2021,2022; V:2018,2019,2021,2022; KK: 2018).

2. Behavioral data collection

In all years of data collection except 2018 (year immediately following Hurricane Maria) and three social groups in 2016, 2017 and 2019 (HH, KK and S respectively), behavioral data were collected with 10-min focal animal samples, where the duration and partner identity of positive (e.g., grooming) and negative (e.g., aggression, threats, submissions, and displacements) social interactions with adults were recorded. At the 0-, 5-, and 10-min marks of the focal follow, we collected instantaneous scan samples during which we recorded the state behavior of the subject (grooming, feeding, resting, and traveling) and the identity of all adults within two meters (i.e., in proximity). For groups HH in 2016, KK in 2017 and S in 2019, we collected 5-min focal animal samples with instantaneous scan samples at the beginning and end of the focal follow, the data collection protocol was otherwise identical to other non-hurricane years. Importantly for this study, grooming, and proximity were mutually exclusive: grooming took precedence over proximity such that whenever two individuals were grooming they were not recorded as being in

proximity as well. We balanced the collection of focal samples on individuals across time of day and across months to account for temporal variation in behaviors. Data-collection hours were from 6am to 2:30pm due to CPRC operating hours. To collect behavioral data before 2020 we used Teklogic Psion WorkAbout Pro handheld computers with Noldus Observer software. After 2020, we used the Animal Observer App on Apple iPads.

In the year following Hurricane Maria (from November 2017 to September 2018), damage resulting in inconsistent access to electricity in Puerto Rico imposed the adoption of a downgraded means of recording data using basic tablets. We recorded group-wide instantaneous scan samples at 10-min intervals. For all animals in view of an observer, we recorded the state behavior of the subject, the identity of their adult social partner when relevant (i.e., if they were grooming) and the identity of all adults within two meters (i.e., in proximity)–similarly to instantaneous scans recorded during focal follows prior to the hurricane. Observers were given 15 mins to complete a group-wide scan session, were required to stand a minimum of 4 m from monkeys and, because of very good visibility of these terrestrial animals, were able to identify them at distances upward of 30 m. While aggressive interactions were recorded during scans in 2018 they were only recorded during focal samples in all other years. Scan samples and focal samples can provide different estimates of brief behaviors like aggression that are not extended in time (*37*). Given this limitation, we excluded 2018 in our comparison of aggressive rates pre-to-post hurricane in this study.

Our subjects were observed over a mean (SD) of 2.61 (1.70) years, and did not have to be observed both before and after the hurricane to be included in analyses. In other words we did not conduct within-individual comparisons as in (*16*). In non-Hurricane Maria years, we included on average 4.64 (2.16) h of focal follows and 84.32 (36.96) scan observations embedded within focal follows per individual per year, and 533.36 (127.41) scans per individual post-hurricane (November 2017 - September 2018). Note that because of the impacts of Hurricane Maria and other storms, data collection stopped on August 31, 2017 and didn't resume until November 2nd, 2017.

Dominance ranks for individuals were determined separately for each group and year. Rank was also determined separately for males and females. For males, the direction and outcome of win-loss agonistic interactions recorded during focal animal samples or during *ad libitum* observations of a given year was used to determine rank for that year. For females, rank was determined using both outcomes of win-loss agonistic interactions and matriline rank. Female macaques inherit their rank from their mothers, and female ranks are linear and relatively stable over time. In order to account for group size, dominance rank was defined by the percentage of same sex individuals outranked, and ranged between 0 to 100 (0 = lowest rank, outranks 0% of same sex individuals; 100 = highest rank, outranks 100%).

3. <u>Changes in vegetation after Hurricane Maria</u>

We quantified changes to vegetation cover as a result of Hurricane Maria, and in particular tree coverage which provides shade, using two quantitative approaches and a qualitative one.

a. Greenery index for short-term vegetation drop pre-to-post hurricane

From satellite images, we extracted the ''Normalized Difference Vegetation Index'' (NDVI) which is the most widely used remote sensing index for assessing vegetation cover (*38*). NDVI is measured using the near-infrared radiation from photosynthetic pigments to assess the photosynthetic activity of vegetation. Importantly, NDVI is sensitive to all types of vegetation, including grass and short bushes which do not provide shade for mid-sized to large vertebrates as tree coverage would. Starting about one year after the hurricane, grass and other ground vegetation started to emerge and are captured by the NDVI index as greenery, even if tree coverage has not recovered (**Figure S1**). We used NDVI to measure the drop in all vegetation types.

NDVI was extracted from satellite images from Sentinel-hub EO-Browser. We used images from Sentinel 2 L2A, a satellite operated by the US Geological Survey that has a 16-day repeat cycle (i.e., visiting Cayo Santiago every approximately 16 days). Images from Sentinel 2 can be viewed in many formats, including NDVI image format. We created a geojson shapefile of coordinates outlining the entirety of Cayo Santiago, including both the large and small islands, which allowed us to specifically search for images in which there was very little cloud cover over the island. We compiled NDVI scores (0 representing no vegetation and 1 representing full vegetation cover) from satellite images with < 10% cloud cover over Cayo Santiago from January 8th 2017 to August 6th 2018 (approximately 1 year before and after Hurricane Maria). In total, we used 45 images, 19 from before Hurricane Maria (01/08/2017–08/31/2017) and 38 from after (10/26/2017–08/06/2018). Following Hurricane Maria, vegetation on the island decreased by 63% (*t*-test, $P < 0.000001$).

b. Tree census

In March 2023, we undertook a tree census of Cayo Santiago, splitting the island into three areas: Upper Cayo, Mangroves and Small Cayo (see **Figure S1C**). We censused trees that were ≥ 20cm in diameter at breast height and with leaves covering a minimum of 20% of their branches in an attempt to count only trees that might provide substantial shade (*39*). On Upper Cayo, we performed a total tree count due to the small number of trees remaining and found a total of 167 trees. In the Mangroves and Small Cayo, where tree density was greater, we conducted a random sampling of tree coverage using a circular plot approach. In brief, we counted all trees of the characteristics described above in randomly selected circular plots of 5m² radius. From a starting point, observers walked 30 meters in a compass direction determined via random number generator. That point was the center of the circle of 5m² radius. This process was repeated, without repeated sampling of areas, until approximately 20% of the total area was covered. In

Sentinel Hub EO browser we used the area measuring tool to quantify the total area of Small Cayo and the Mangroves, which were 20,000m^2 and 30,000m^2 respectively. On Small Cayo, we found a median of 1 tree per 78m^2. Extrapolating to the total area of Small Cayo (excluding cliff areas) we estimated the total tree coverage to be 20,000/78x1 = 256. For Mangroves, we found a median of 0 trees per 78m^2 which, using the same approach, yields 0 trees. If using the mean instead of median, we estimated a total of 30,000/78*0.7 = 269 trees in the Mangroves. Upon visual inspection of the google Earth images, the complete tree count likely lies between these two estimates.

c. Qualitative assessment of Google Earth images

Google Earth images allowed us to visualize the state of Cayo Santiago island at multiple time points from September 2016 until February 2022 (**Figure S1A**). Although the quality and viewpoints of the images prevented us from performing a quantitative assessment of the tree coverage, they were sufficient for a qualitative assessment and provided valuable historical information about tree coverage pre-hurricane that we do not otherwise have access to. We concluded from this timeline of images and on-the-ground experience that the vegetation cover is far from pre-hurricane levels.

4. Temperature logging after Maria

Surface temperature of the island was quantified using 39 temperature loggers (IButton DS1922L Thermochron Data Logger) dispatched across Cayo Santiago in 2018, i.e., after Hurricane Maria. Loggers were dispatched in three types of terrain: exposed, de-vegetated and vegetated. 12 temperature loggers were located in 'exposed' terrain, which did not have tree coverage either before or after the hurricane. 13 loggers were located in 'de-vegetated' terrain which used to have tree coverage but did not anymore after the hurricane. Finally, 14 loggers were located in 'vegetated' terrain with tree coverage (i.e., in the shade). We compared temperatures from all 39 loggers in the three types of terrain from July 7th 2018 to October 10th 2021, only including temperatures from 6am to 5pm, amounting to a total of 463,510 samples. We find a mean difference of 7ºC between shaded areas ('vegetated' terrain) and non-shaded areas (exposed and de-vegetated, $P < 0.00001$).

When plotting the mean temperature detected across all loggers in non-shaded areas (i.e., 'exposed' or 'de-vegetated') by time of day (6am-5pm, considering all months of the year), we find that temperatures exceed 38ºC, dangerously intense heat for macaques (9), beginning at 10am (**Figure S1D**). To assess the effect of temperature on sociality and survival patterns (see below for more details), we split our data into cooler morning hours, from 6-10am, and hotter afternoon hours, 10am-2:30pm.

5. Estimating social networks and associated metrics using a Bayesian approach (BISoN)

For each group and year separately, we estimated three types of social networks based on three different social interactions: proximity, aggression and grooming. For each social interaction type, we quantified two individual-level network metrics that allowed us to measure individuals' sociality: (1) the number of social partners an individual interacted with, or number of connections in the network (i.e., degree); and (2) the propensity at which individuals interacted with their partners, or the sum of probabilities of interaction with all individuals in the network (i.e., strength)(*27*).

Networks are usually constructed by taking a normalized measure of sociality, such as the proportion of sampling periods each pair spends engaged in a social behavior (e.g. the Simple Ratio Index; SRI). These normalized measures ensure that pairs that are observed for longer are not erroneously determined to be more social (*40*). This is important because observing social events can be challenging and uniform sampling over all pairs is not always possible (*41*). Though normalized measures of sociality will not be biased by sampling time, they will be accompanied by varying levels of certainty. For example, edge weights will be treated as a certain 0.5 for both a case where individuals have been seen together once and apart once, and equally where individuals have been together 100 times and apart 100 times. Despite there being considerably more certainty around the values of the latter example than the former, methods to estimate and propagate uncertainty through social network analyses remain largely unused.

In this study, we generated weighted Bayesian networks using the BISoN framework and bisonR package (*18*). This framework allowed us to account for uncertainty in the edges connecting individuals in the network based on how often they were sampled and, more importantly, propagate this uncertainty to subsequent analyses. In the next paragraphs we describe how we modeled each network type in BISoN:

Proximity networks were estimated based on data collected through focal scan sampling (*37*) across all groups and years. Edges in the networks represented the number of times a pair of individuals were observed in proximity relative to the total observation effort for the dyad (i.e., total scans individual A + total scans individual B), or the probability of interaction. Given that the observed interaction data was binary (i.e. for every observation sample, was the focal animal in proximity to another monkey? Y/N) we modeled the uncertainty around the edges using as weak prior a beta distribution with alpha = 0.1 and beta = 0.1. Proximity networks were undirected.

Aggressive networks, on the other hand, were estimated based on data collected through continuous focal sampling across all groups and years except the year immediately following Hurricane Maria (2018). In 2018 aggressive interactions were collected through scan sampling which makes this year hardly comparable to all other years. As such, 2018 were excluded from our analyses for aggression data. Edges in the network represented the number of aggressive

interactions between members of a dyad over the total observation effort for the dyad (i.e., total number of hours observed individual A + total number of hours observed individual B), or the rate of interaction. Given that the interaction data consisted of counts of aggressive events over total time observed, we modeled the uncertainty around the edges using a gamma distribution with alpha = 0.1 and beta = 0.1 as weak prior. Aggression networks were directed.

Grooming interactions were captured both through scan sampling and continuous focal sampling such that we quantified grooming networks using both approaches described above. Scan-based grooming networks were modeled with a beta distribution (as for proximity) and continuous sampling-based grooming networks were modeled with a gamma distribution (as for aggression). We did not collect continuous focal sampling in 2018 therefore continuous sampling-based grooming networks do not include 2018. Grooming networks were directed.

The BISoN bayesian framework computes a posterior distribution of possible networks from the observed data, from which we sampled 100 grooming, proximity and aggression networks for each group and year. These networks sampled from posterior distributions were used for downstream analyses.

Networks generated with BISoN include edges between all dyads by default, as it assumes non-zero probability for all potential interactions, even if that probability is exceedingly small. To compute degree (or number of connections in the network for each individual), we therefore defined a threshold of 0.0001, 1 interaction in 10,000 observations, to differentiate dyads that did interact versus those that did not. After this thresholding, we computed two individual-level network metrics for each possible BISoN network, which were used for downstream analyses: (1) nodal degree using the 'degree' function from *igraph* (i.e., number of edges for each individual or node in the network); and (2) relationship strength to all partners in the network using "strength" function in *igraph* (summing all the weights of an individual's edges in the network). Thresholds of 0.001 or 0.00001 for defining an edge in networks for degree calculations yielded very similar results.

6. Methodological improvements since previous work

Note that we used different analytical methods to conduct network analyses in this study compared to our previous work (*16*). Critically, methods differ in how they account for uncertainty around edge weight estimates, especially for edge weight = 0. In our previous work, we used a subsampling bootstrap approach to control for observation effort differences pre- vs. post-hurricane. However, if no interaction was observed between two nodes in the network, the edge weight would always stay equal to 0 (i.e., no interaction can ever be picked up by sub-sampling) - biasing networks with lower sampling effort toward appearing sparser (with less connections) compared to better-sampled networks, even if observation effort is kept the same through sub-sampling. In this current study, to address this issue, which incidentally pervades the

field of animal social network analysis, we used a new Bayesian network method, BISoN, which adds uncertainty around edge weights based on observation effort, such that 0 observed interactions can be modeled as more or less likely to be 0 depending on the observation effort. In pre-hurricane networks, which were based on less sampling effort compared to 2018 ones (~5-10 times lower), this new method avoids 0-inflation in edge-weights due to sparse sampling. The methodological improvement in this study explains discrepancies with our previous results regarding changes in grooming networks, which were previously reported to be denser in 2018 compared to before the hurricane, but which we now found to be equally dense or sparser in 2018 compared to pre-hurricane years (**Figure S2**). Note that grooming results differ if we use focal sampling or scan sampling to collect our behavioral data (**Figure S2**), and as a result patterns of grooming post-hurricane compared to pre-hurricane remain unclear.

7. Long-term change in sociality

To assess long-term changes in the propensity to be in proximity pre- to post-hurricane, we used linear mixed models using the R package lme4 (*42*). We included all individuals for which we had behavioral data in any year from 2013 to 2022 across 7 social groups, yielding a total of 790 unique individuals. In all our models our response variable was standardized individual network 'strength' (i.e., the sum of weights of an individual's connections in the network) and our main predictor of interest was the year post-hurricane compared to pre-hurricane years. We included this 'year' predictor as a factor in our model, where all pre-hurricane years were combined into a "pre-hurricane" category and post-hurricane years were split into 2018, 2019, 2021 and 2022 (we did not collect data in 2020 due to the Covid19 crisis). We also included covariates age and sex as fixed effects, and social group and individual ID as random effects to account for repeated measures. The same analysis pipeline was also done for grooming and aggression data. Finally, to compare changes in proximity in the cooler hours of the day vs. the hotter hours of the day (after 10am when temperatures reached 38C, **Figure S1**), we repeated the same analytical approach but simply considered data collected in earlier in the day (7-10am) vs later in the days (10am-2:30pm) to construct social networks.

For all models we ran the following assumption checks to ensure model estimates were accurately estimated using the function 'check_model' from R package *performance (43)*: posterior predictive checks, homogeneity of variance, colinearity of fixed effects, normality of residuals, normality of random effects. We did not find violations of the assumptions checked.

8. Mixed effect cox-proportional hazard survival models

We investigated whether social tolerance, as measured by the propensity to be in proximity to other monkeys and the number of proximity partners, predicted survival in the years before (2013-2017) and after the hurricane (2018-2022). We used time-varying mixed-effect Cox proportional hazard models, through the *coxme* R package (*44*), divvying up our analysis by year pre- and post-hurricane. This is because the value of social covariates changed from year-to-year.

We used behavioral data from each year to predict survival to the next year. For 2020 in which we did not collect behavioral data, we used social covariates calculated from the previous year (i.e. 2019). Our results remain qualitatively the same if we exclude 2020 from our analysis.

For all models, the response variable was a 'Surv' object with age in days at the start of the study period, age in days at the time of event (which could be death, removal or end of study), and the event outcome (0: the animal is censored, 1: the animal is dead). Culled individuals during the study period were censored to their cull date and all animals still alive at the end of the study period were censored to the end date. With this approach, age was accounted for implicitly in all models.

Importantly, study start and end dates varied with the follow-up period considered. In explicit terms: when measuring the survival effects of proximity in 2013, we only considered individuals for which we had behavioral data in 2013. The start of the study was January 1st 2013 and the study end date was December 31st 2013. Individuals who died after 2013 were considered alive when quantifying the effect of 2013 proximity on survival (i.e., event outcome = 0, the animal is censored). All other years were treated the same except 2017 and 2018. For 2017, study end date was set to Sept 20th to exclude post-hurricane deaths. For 2018, study start date was set to Sept 21st to include deaths immediately following Maria. This means that individuals who died immediately of the hurricane were excluded from our analysis (since we do not have behavioral data for them for the Oct. 2017- December 2018 observation year).

For pre-hurricane (2013-2017) and post-hurricane (2018-2022) periods separately, we ran two survival models with either nodal strength or nodal degree in proximity network as social covariates (i.e., a total of four models). All four models also included sex and social status as a fixed effects and social group, individual ID and year considered (or follow-up period) as random effects. All adult monkeys with behavioral data in any year from 2013 to 2022 were considered for this analysis. Pre-hurricane, we considered a total of 616 unique individuals across 5 social groups (HH, R, F, V and KK), of which 65 died in the study period. Post-hurricane, we considered 430 unique individuals across 4 social groups (S, KK, V and F) of which 56 died.

To compare how proximity predicted individuals' survival earlier in the day compared to later in the day (when hotter), we repeated the same analysis but only considering proximity data collected from 7-10 am or from 10am-2:30pm.

For all cox-proportional hazard models we ran the following assumption checks to ensure model estimates were accurately estimated: proportional hazard over time using the R function 'cox.zph' (*45*) and (2) linear relationship between the log hazard and each covariate using R function 'residuals'.

9. <u>Pooling model results across sampled network metrics</u>

In order to propagate the uncertainty around social networks constructed from the observed interaction data to downstream analyses, all models described above were computed across 100 possible social networks for each group-year-behavior combination generated in BISoN (i.e., extracted from the networks' posterior distribution). We used the function 'pool' from the R package *mice* to combine estimates and the uncertainty around those estimates from the 100 repeated models. As an example, to compute the effect of pre-hurricane proximity on survival, we ran 100 survival models using network metrics extracted from 100 different possible proximity networks per group-year and pooled estimated across those 100 models.

Supplementary Text:

<u>Changes in feeding patterns after Maria do not explain survival results</u>

Before the hurricane, 33.6% of Cayo macaques' diet was vegetation-based, the rest being 'monkey chow' supplemented by the Caribbean primate research center (CPRC) in centralized feeders. Hurricane-induced vegetation loss led to vegetation-based diet dropping to 21.5%, the rest being CPRC-distributed monkey chow. Moreover, starting in April 2017 (6 months before Maria), the CPRC began distributing food across the island resulting in *less* centralized feeding. These feeding changes are unlikely to explain both higher social tolerance post-hurricane and its effect on survival for the following reasons. First, being in proximity to conspecifics did not correlate with the amount of time an individual spent feeding post-hurricane (r=0.06, $P = 0.10$) suggesting one does not enable the other. Second, feeding only occurred once a day in the morning, and afternoon proximity was both highest compared to pre-hurricane and better predicted survival. Reduced aggression post-hurricane, however, may be partly explained by less centralized feeding post-hurricane.

Figure S1. Vegetation and temperature after Hurricane Maria. (A) Timeline of Google Earth Satellite images of Cayo Santiago. (B) Map of the sections of Cayo Santiago for tree census. (C) Average day temperatures from 2018 to 2023 in exposed temperature loggers.

Figure S2. Long term effect of Maria on sociality. Results for linear mixed models for grooming, proximity and aggression.

	2013	2014	2015	2016	2017	2018
F	100	106	103	123	113	0
HH	0	45	0	69	0	0
KK	30	0	54	0	61	63
R	0	0	94	125	0	0
S	0	0	0	0	0	0
TT	0	0	0	0	0	0
V	0	0	80	90	102	67

Table S1. Sample sizes by group and year. Note that an individual can be sampled in multiple years.