Title: Inflammasomes primarily restrict cytosolic *Salmonella* replication within human

macrophages

Authors: Marisa S. Egan[1], Emily A. O'Rourke[1], Shrawan Kumar Mageswaran[2], Biao

Zuo[3], Inna Martynyuk[3], Tabitha Demissie[1], Emma N. Hunter[1], Antonia R. Bass[1†], Yi-Wei

Chang[2], Igor E. Brodsky[4], and Sunny Shin[1#]

[1]Department of Microbiology, Perelman School of Medicine, University of Pennsylvania,

Philadelphia, PA

[2]Department of Biochemistry and Biophysics, Perelman School of Medicine, University

of Pennsylvania, Philadelphia, PA

[3]Electron Microscopy Resource Laboratory, Department of Biochemistry & Biophysics,

Perelman School of Medicine, University of Pennsylvania, Philadelphia, PA

[4]Department of Pathobiology, University of Pennsylvania School of Veterinary Medicine,

Philadelphia, PA

Current Address:

[†]MRL, Merck & Co., Inc., Kenilworth, NJ

#Address correspondence to Sunny Shin, sunshin@pennmedicine.upenn.edu.

Abstract

Salmonella enterica serovar Typhimurium is a facultative intracellular pathogen

that utilizes its type III secretion systems (T3SSs) to inject virulence factors into the host

cell and colonize the host. In turn, a subset of cytosolic immune receptors respond to

T3SS ligands by forming multimeric signaling complexes called inflammasomes, which

activate caspases that induce interleukin-1 (IL-1) family cytokine release and an

inflammatory form of cell death called pyroptosis. Human macrophages mount a

multifaceted inflammasome response to *Salmonella* infection that ultimately restricts

intracellular bacterial replication. However, how inflammasomes restrict *Salmonella*

replication remains unknown. We find that caspase-1 is essential for mediating

inflammasome responses to *Salmonella* and subsequent restriction of bacterial

replication within human macrophages, with caspase-4 contributing as well. We also

demonstrate that the downstream pore-forming protein gasdermin D (GSDMD) and

ninjurin-1 (NINJ1), a mediator of terminal cell lysis, play a role in controlling *Salmonella*

replication in human macrophages. Notably, in the absence of inflammasome

responses, we observed hyperreplication of *Salmonella* within the cytosol of infected

cells, and we also observed increased bacterial replication within vacuoles, suggesting

that inflammasomes control *Salmonella* replication primarily within the cytosol and also

within vacuoles. These findings reveal that inflammatory caspases and pyroptotic

factors mediate inflammasome responses that restrict the subcellular localization of

intracellular *Salmonella* replication within human macrophages.

<u>Introduction</u>

Intracellular bacterial pathogens create and maintain replicative niches within host cells. To survive inside the host, these pathogens must remodel the host cellular landscape and overcome immune defenses. *Salmonella enterica* serovar Typhimurium (*Salmonella*) is a facultative intracellular pathogen and a major cause of food-borne illness worldwide (Majowicz et al., 2010). Following ingestion, *Salmonella* colonizes the intestinal tract, where it can invade, replicate, and survive in host cells, including macrophages (Crowley et al., 2016; R. L. Santos & Bäumler, 2004). *Salmonella* employs type III secretion systems (T3SSs) that act as molecular syringes to inject virulence factors, or effectors, into the host cell cytosol (Agbor & McCormick, 2011; Crowley et al., 2016). Specifically, *Salmonella* relies on two distinct T3SSs encoded on *Salmonella* Pathogenicity Islands 1 and 2 (SPI-1 and SPI-2) to invade and replicate within host cells, respectively (Mills et al., 1995; Shea et al., 1996; Hensel et al., 1998; Galan & Zhou, 2000; Galán, 1999; Galán & Collmer, 1999; Galan & Curtiss, 1989; Ochman et al., 1996; Cirillo et al., 1998). The SPI-2 T3SS translocates effectors to facilitate biogenesis and maintenance of the *Salmonella*-containing vacuole (SCV), wherein *Salmonella* resides and replicates (Cirillo et al., 1998; Takeuchi, 1967; Takeuchi & Sprinz, 1967; Kihlström & Latkovic, 1978; Garcia-del Portillo et al., 1993; Garcia-del Portillo & Finlay, 1995; Steele-Mortimer et al., 1999; Hansen-Wester et al., 2002; Steele-Mortimer, 2008; Jennings et al., 2017; Brumell, Goosney, et al., 2002; Brumell, Tang, et al., 2002; Beuzón et al., 2000; Hensel et al., 1998). In epithelial cells, *Salmonella* escapes its vacuolar niche and hyperreplicates in the host cell cytosol (Knodler, Crowley, et al., 2014; Knodler et al., 2010; Knodler, Nair, et al., 2014; Malik-

Kale et al., 2012). While T3SSs are essential for *Salmonella* to infect host cells, they

also inject flagellin (Sun et al., 2007) and structural components of the T3SS into the

host cell cytosol, which conversely leads to cytosolic immune detection of *Salmonella*.

The mammalian innate immune system detects violations of cytosolic sanctity,

such as the presence of intracellular bacterial pathogens, through cytosolic pattern

recognition receptors (PRRs) (Janeway, 1989; Medzhitov & Janeway, 2002). A subset

of these PRRs includes the nucleotide-binding domain, leucine-rich repeat (NLR) family

proteins. NLRs respond to their cognate stimuli by oligomerizing and inducing the

assembly of multiprotein complexes termed inflammasomes, which activate

inflammatory caspases (Broz & Dixit, 2016; Lamkanfi & Dixit, 2009, 2014; Martinon et

al., 2002). Canonical inflammasomes recruit and activate caspase-1, which cleaves and

activates pro-inflammatory interleukin-1 (IL-1) family cytokines and the pore-forming

protein gasdermin D (GSDMD) (Kuida et al., 1995; Li et al., 1995; Agard et al., 2010;

Thornberry et al., 1992; Shi et al., 2015; Kayagaki et al., 2015). Alternatively,

noncanonical inflammasomes are formed by caspase-11 in mice and two orthologs in

humans, caspase-4 and caspase-5, in response to cytosolic lipopolysaccharide (LPS)

(Casson et al., 2015; Hagar et al., 2013; Kayagaki et al., 2011, 2013; Lagrange et al.,

2018; Schmid-Burgk et al., 2015; Shi et al., 2014). These caspases directly process and

activate GSDMD (Agard et al., 2010; Kayagaki et al., 2015; Shi et al., 2015). Liberated

GSDMD N-terminal fragments oligomerize to create pores in the host plasma

membrane, through which IL-1 family cytokines and other alarmins are released to

promote a lytic form of inflammatory cell death termed pyroptosis (Agard et al., 2010;

Ding et al., 2016; Kayagaki et al., 2015; Shi et al., 2015).

Salmonella activates several inflammasomes in human macrophages. One such

inflammasome is the NLR family, apoptosis inhibitory protein (NAIP)/NLR family, CARD

domain-containing protein 4 (NLRC4) inflammasome (Bierschenk et al., 2019; Gram et

al., 2021; Naseer, Egan, et al., 2022). NAIP detects the cytosolic presence of T3SS

structural components and flagellin (Grandjean et al., 2017; Kofoed & Vance, 2011;

Kortmann et al., 2015; Miao, Mao, et al., 2010; Molofsky et al., 2006; Rauch et al., 2016;

Rayamajhi et al., 2013; Ren et al., 2006; Reyes Ruiz et al., 2017; Sun et al., 2007; Yang

et al., 2013; Zhao et al., 2011, 2016). Upon ligand recognition, NAIP recruits NLRC4,

which oligomerizes to form the active NAIP/NLRC4 inflammasome (Diebolder et al.,

2015; Z. Hu et al., 2015; Zhang et al., 2015). *Salmonella* also activates the NLR pyrin

domain-containing protein 3 (NLRP3) inflammasome and the noncanonical

inflammasome in human macrophages (Bierschenk et al., 2019; Casson et al., 2015;

Gram et al., 2021; Naseer, Egan, et al., 2022). NLRP3 responds to diverse stimuli,

including ionic fluxes as a result of host plasma membrane damage (Franchi et al.,

2007; Hornung et al., 2008; Mariathasan et al., 2006; Muñoz-Planillo et al., 2013;

Perregaux & Gabel, 1994), and can be secondarily activated by the noncanonical

inflammasome (Baker et al., 2015; Casson et al., 2015; Kayagaki et al., 2013; Pilla et

al., 2014; Rathinam et al., 2012; Rühl & Broz, 2015; Schmid-Burgk et al., 2015; Shi et

al., 2014).

Inflammasome activation is critical for host defense against *Salmonella*. In mice,

the NAIP/NLRC4 inflammasome is required to control *Salmonella* infection (Carvalho et

al., 2012; Franchi et al., 2012; Hausmann et al., 2020; Miao et al., 2006; Miao, Mao, et

al., 2010; Rauch et al., 2016, 2017; Sellin et al., 2014; Zhao et al., 2016). In murine

intestinal epithelial cells (IECs), NAIP/NLRC4 inflammasome activation triggers

pyroptosis and expulsion of infected cells, and is both necessary and sufficient in IECs

to restrict *Salmonella* replication and prevent bacterial dissemination to distal organ

sites (Hausmann et al., 2020; Rauch et al., 2017; Sellin et al., 2014). Unlike murine

IECs, human epithelial cells do not rely on NAIP/NLRC4 or caspase-1, but instead rely

on caspase-4 to control *Salmonella* replication, in part through pyroptosis and cell

extrusion (Holly et al., 2020; Knodler, Crowley, et al., 2014; Knodler et al., 2010;

Naseer, Zhang, et al., 2022). Moreover, in murine macrophages, inflammasome

activation controls the replication of a mutant strain of *Salmonella* that aberrantly

invades the cytosol, but only slightly limits replication of wild-type (WT) *Salmonella*

(Thurston et al., 2016). In contrast, we found that human macrophages rely on

NAIP/NLRC4- and NLRP3-dependent inflammasome responses to control intracellular

WT *Salmonella* replication (Naseer, Egan, et al., 2022). However, how inflammasome

signaling restricts *Salmonella* replication in human macrophages remains unknown.

In this study, we show that caspase-1 is required for inflammasome responses

and the restriction of intracellular *Salmonella* replication early during infection in human

macrophages, and that caspase-4 contributes to the restriction of *Salmonella* later

during infection. We find that the cell lysis mediators GSDMD and ninjurin-1 (NINJ1)

contribute to bacterial restriction. Importantly, we observed that while human

macrophages unable to undergo inflammasome responses showed slightly elevated

bacterial replication within SCVs, they became permissive to hyperreplication of

Salmonella within the cytosolic compartment. Thus, inflammasome activation appears

to preferentially restrict cytosolic *Salmonella* replication. Our results offer insight into

how inflammasomes control *Salmonella* in human macrophages and restrict the distinct

intracellular spatial niches that *Salmonella* occupy in these cells.

<u>Results</u>

<u>Caspase-1 promotes the control of *Salmonella* replication within human macrophages</u>

Human macrophages undergo NAIP/NLRC4- and NLRP3-dependent

inflammasome activation during *Salmonella* infection (Bierschenk et al., 2019; Gram et

al., 2021; Naseer, Egan, et al., 2022), which restricts intracellular *Salmonella* replication

(Naseer, Egan, et al., 2022). Nonetheless, how inflammasome activation controls

Salmonella replication in human macrophages remains unclear. Inflammasomes recruit

and activate caspase-1 in both murine and human cells (Franchi et al., 2006; Man,

Hopkins, et al., 2014; Mariathasan et al., 2004; Miao et al., 2006; Ross et al., 2022;

Zamboni et al., 2006), and caspase-1 promotes the control of *Salmonella* both *in vivo*

and *in vitro* (Broz et al., 2010, 2012; Crowley et al., 2020; Hausmann et al., 2020; Holly

et al., 2020; Lara-Tejero et al., 2006; Miao, Leaf, et al., 2010; Rauch et al., 2017;

Raupach et al., 2006; Sellin et al., 2014; Thurston et al., 2016). Thus, we sought to test

whether caspase-1 restricts *Salmonella* replication in human macrophages.

To first interrogate the impact of caspase activity on *Salmonella* replication in

human macrophages, we pretreated macrophages derived from the human monocytic

cell line, THP-1, with Ac-YVAD-cmk (YVAD), a chemical inhibitor of caspase-1 activity,

or Z-VAD-FMK (ZVAD), a pan-caspase inhibitor. Upon infection with wild-type (WT)

Salmonella, WT THP-1 cells pretreated with either YVAD or ZVAD had significantly

reduced levels of IL-1β secretion and cell death compared to infected WT cells

pretreated with DMSO, the vehicle control (Figure 1 – figure supplement 1A-B). Since

pretreatment with YVAD largely phenocopies ZVAD, these data suggest that caspase-1

is the primary caspase that responds to *Salmonella* infection in THP-1 cells. We next

assessed intracellular *Salmonella* burdens by determining bacterial colony forming units

(CFUs). The increase in bacterial CFUs was significantly higher in WT cells pretreated

either YVAD or ZVAD compared to cells pretreated with DMSO (Figure 1 – figure

supplement 1C). Microscopic analysis revealed that WT cells treated with either YVAD

or ZVAD prior to infection harbored significantly higher intracellular *Salmonella* burdens

compared to cells pretreated with DMSO (Figure 1 – figure supplement 1D). Overall,

these results suggest that caspase activity, primarily caspase-1 activity, controls

Salmonella burdens in human macrophages.

Next, to genetically test the requirement of caspase-1, we used two independent

CASP1$^{-/-}$ THP-1 single cell clones generated through CRISPR/Cas9-mediated deletion

(Okondo et al., 2017). In agreement with previous reports (Bierschenk et al., 2019;

Gram et al., 2021; Naseer, Egan, et al., 2022), WT THP-1 cells infected with WT

Salmonella exhibited high levels of IL-1β, IL-18, and IL-1α release as well as cell death

at 6 hpi (Figure 1A-B; Figure 1 – figure supplement 2A-B). In contrast, *CASP1$^{-/-}$* THP-1

cells released negligible levels of inflammasome-dependent cytokines and did not

undergo substantial cell death upon infection (Figure 1A-B; Figure 1 – figure

supplement 2A-B). Infected WT and *CASP1$^{-/-}$* THP-1 cells released similar levels of the

inflammasome-independent cytokine TNF-α (Figure 1 – figure supplement 2C). These

results indicate that caspase-1 is required for inflammasome responses to *Salmonella*

infection in human macrophages.

We next tested whether caspase-1 is required to restrict *Salmonella* replication in

human macrophages. At 1 hour post-infection (hpi), we did not observe any significant

differences in bacterial uptake between WT and *CASP1*[-/-] THP-1 cells (Figure 1 – figure

supplement 2D). However, at 6 hpi, *CASP1*[-/-] cells harbored significantly higher bacterial

burdens and a significant fold-increase in bacterial CFUs compared to WT cells (Figure

1C; Figure 1 – figure supplement 2E). To assess intracellular bacterial burdens on a

single-cell level, we quantified the amount of WT *Salmonella* per cell by microscopy. WT

cells contained relatively low numbers of *Salmonella* on average (~5 bacteria per cell),

while *CASP1*[-/-] cells harbored significantly higher numbers of *Salmonella* on average

(~30 bacteria per cell) at 6 hpi (Figure 1D-E). Overall, these data indicate that caspase-

1 limits intracellular *Salmonella* replication in human macrophages.

<u>Caspase-4 contributes to the control of *Salmonella* replication within human</u>

<u>macrophages later during infection</u>

Caspase-11 is another member of the inflammatory caspase family, and plays a

role in host defense against *Salmonella* in mice (Aachoui et al., 2013; Crowley et al.,

2020; Knodler, Crowley, et al., 2014; Sellin et al., 2014). Humans harbor two orthologs

of murine caspase-11: caspase-4 and caspase-5 (Shi et al., 2014). Caspases-11/4/5

recognize cytosolic LPS and form the noncanonical inflammasome to mediate

downstream inflammatory responses (Casson et al., 2015; Hagar et al., 2013; Kayagaki

et al., 2011, 2013; Lagrange et al., 2018; Schmid-Burgk et al., 2015; Shi et al., 2014).

Caspase-4 contributes to inflammasome responses during *Salmonella* infection of THP-

1 macrophages and primary human macrophages (Casson et al., 2015; Naseer, Egan,

et al., 2022), and caspases-4/5 are required for inflammasome responses to *Salmonella*

in THP-1 monocytes (Baker et al., 2015). In human intestinal epithelial cells (IECs),

caspase-4 drives inflammasome responses during *Salmonella* infection and limits

intracellular bacterial replication (Holly et al., 2020; Knodler, Crowley, et al., 2014;

Naseer, Zhang, et al., 2022). However, whether caspase-4 contributes to restriction of

intracellular *Salmonella* replication within human macrophages is unclear.

To genetically test the contribution of caspase-4 during *Salmonella* infection in

THP-1 macrophages, we used CRISPR/Cas9-mediated deletion to disrupt the *CASP4*

gene. We selected and sequence-validated two independent *CASP4*$^{-/-}$ THP-1 single cell

clones (Figure 2 – figure supplement 1). We observed a slight decrease in secreted IL-

1β levels in *CASP4*$^{-/-}$ THP-1 macrophages infected with WT *Salmonella* compared to

infected WT THP-1 macrophages at 6 hpi (Figure 2 – figure supplement 2A). We also

observed a slight decrease in cell death at 6 hpi in infected *CASP4*$^{-/-}$ clone #2 compared

to infected WT cells, whereas cytotoxicity levels were unaffected in infected *CASP4*$^{-/-}$

clone #6 (Figure 2 – figure supplement 2B). However, at 24 hpi, we observed a

significant decrease in IL-1 release and cell death in both *CASP4*$^{-/-}$ clones infected with

WT *Salmonella* compared to infected WT cells (Figure 2A-B; Figure 2 – figure

supplement 3A-B). WT and *CASP4*$^{-/-}$ cells also released similar levels of the

inflammasome-independent cytokine TNF-α upon infection with WT *Salmonella* at 24

hpi (Figure 2 – figure supplement 3C). Overall, these data indicate that caspase-4

contributes to inflammasome responses in THP-1 macrophages later during *Salmonella*

infection.

Since caspase-4 restricts *Salmonella* replication in human epithelial cells (Holly

et al., 2020; Knodler, Crowley, et al., 2014; Naseer, Zhang, et al., 2022), we then asked

whether caspase-4 contributes to the control of *Salmonella* replication in human

macrophages. Upon examination of the fold-change in bacterial CFUs at 6 hpi, we did

not observe any significant differences between WT and *CASP4*$^{-/-}$ THP-1 cells (Figure 2

– figure supplement 2C). However, at 24 hpi, *CASP4*$^{-/-}$ cells harbored higher bacterial

burdens and a significant fold-increase in bacterial CFUs compared to WT cells (Figure

2C; Figure 2 – figure supplement 3D). To confirm these findings, we enumerated the

amount of WT *Salmonella* per cell by microscopy. At 6 hpi, WT and *CASP4*$^{-/-}$ THP-1

cells contained comparable numbers of *Salmonella* per cell (Figure 2 – figure

supplement 2D-E). In contrast, at 24 hpi, *CASP4*$^{-/-}$ cells harbored significantly higher

burdens of *Salmonella* per cell on average (~20 bacteria per cell) compared to WT cells

(~10 bacteria per cell) (Figure 2D-E). Altogether, these results suggest that caspase-4

plays a larger role in controlling *Salmonella* burdens later during infection.

<u>GSDMD promotes the control of *Salmonella* replication within human macrophages</u>

Inflammasome activation triggers a lytic form of cell death known as pyroptosis

(Lamkanfi & Dixit, 2014). Death of the infected host cell eliminates *Salmonella*'s

intracellular replicative niche and thus, may contribute to restricting intracellular bacterial

replication. Upon its cleavage by inflammatory caspases, GSDMD forms pores in the

host plasma membrane, resulting in pyroptosis (Agard et al., 2010; Ding et al., 2016;

Kayagaki et al., 2015; Shi et al., 2015). Whether GSDMD contributes to the control of

Salmonella replication in human macrophages is unknown.

First, we asked whether GSDMD pore formation plays a role in controlling

Salmonella replication in human macrophages. To do this, we pretreated THP-1

macrophages and primary human monocyte-derived macrophages (hMDMs) with the

chemical inhibitor disulfiram. Disulfiram prevents cleaved GSDMD from inserting into

the host plasma membrane, thereby limiting GSDMD-mediated pore formation (J. J. Hu

et al., 2020). Disulfiram treatment led to the loss of IL-1β release and cytotoxicity in

macrophages infected with WT *Salmonella*, compared to treatment with the vehicle

control, DMSO (Figure 3 – figure supplement 1A-B, D-E). Next, we examined the effect

of GSDMD-mediated pore formation on intracellular *Salmonella* burdens. The fold-

increase in bacterial CFUs at 6 hpi was significantly higher in cells treated with

disulfiram compared to cells treated with DMSO (Figure 3 – figure supplement 1C, F).

Collectively, these results suggest that GSDMD-mediated pore formation promotes the

restriction of intracellular *Salmonella* replication.

Consistent with our findings with disulfiram, we observed a significant decrease

in IL-1 cytokine release in *GSDMD*[-/-] THP-1 macrophages (Okondo et al., 2017;

Taabazuing et al., 2017) infected with WT *Salmonella* compared to infected WT THP-1

macrophages (Figure 3A; Figure 3 – figure supplement 2A-B). These results suggest

that human macrophages undergo GSDMD-dependent IL-1 release during infection.

Interestingly, loss of GSDMD did not completely abrogate the release of IL-1,

suggesting that there is also GSDMD-independent IL-1 release (Figure 3A; Figure 3 –

figure supplement 2A-B). Infected WT and *GSDMD*[-/-] THP-1 cells secreted similar levels

of the inflammasome-independent cytokine TNF-α (Figure 3 – figure supplement 2C).

Importantly, infected *GSDMD*[-/-] cells failed to undergo substantial cell death, in contrast

to infected WT cells, suggesting that GSDMD facilitates cell death during *Salmonella*

infection (Figure 3B). Overall, these results indicate that GSDMD plays an important

role in mediating inflammasome responses during *Salmonella* infection of human

macrophages.

We next examined whether GSDMD controls intracellular *Salmonella* replication

in human macrophages. While we did not observe any differences in bacterial uptake

between WT and *GSDMD*[-/-] cells at 1 hpi (Figure 3 – figure supplement 2D), we did

observe significantly higher bacterial CFUs in *GSDMD*[-/-] cells compared to WT cells at 6

hpi (Figure 3 – figure supplement 2E). Moreover, the increase in bacterial CFUs at 6 hpi

was significantly higher in *GSDMD*[-/-] cells than in WT cells (Figure 3C). Next, we used

microscopy to further interrogate intracellular *Salmonella* burdens. As expected, WT

cells contained low numbers of *Salmonella* per cell (~5 bacteria per cell) (Figure 3D-E).

However, *GSDMD*[-/-] cells harbored significantly higher numbers of *Salmonella* per cell

on average (~20 bacteria per cell) at 6 hpi (Figure 3D-E), suggesting that GSDMD

restricts intracellular *Salmonella* burdens. Altogether, these data indicate that GSDMD

promotes the control of *Salmonella* replication in human macrophages.

NINJ1 contributes to the control of *Salmonella* replication within human macrophages

Terminal cell lysis is regulated downstream of GSDMD cleavage and pore

formation (Bjanes et al., 2021; Kayagaki et al., 2021). While GSDMD pores release a

subset of molecules, including IL-1 family cytokines, the extensive release of cellular

contents is mediated by plasma membrane rupture (Ding et al., 2016; Ruan et al.,

2018). Recently, NINJ1 was identified as an executioner of terminal cell lysis (Bjanes et

al., 2021; Borges et al., 2022; Degen et al., 2023; Kayagaki et al., 2021, 2023). NINJ1 is

a transmembrane protein that oligomerizes in the host plasma membrane to induce lytic

rupture of the cell after the initiation of pyroptosis and other forms of regulated cell death

(Degen et al., 2023; Kayagaki et al., 2021). Cells can be protected from plasma

membrane rupture through treatment with the amino acid glycine (Fink & Cookson,

2006; Frank et al., 2000; Heilig et al., 2018; Verhoef et al., 2005), which interferes with

the clustering of NINJ1 (Borges et al., 2022). Thus, upon incubation with glycine, cells

can still form GSDMD pores but cannot undergo terminal cell lysis (Fink & Cookson,

2006; Heilig et al., 2018; Tsuchiya et al., 2021; Verhoef et al., 2005). Whether NINJ1-

dependent cell lysis might contribute to control of *Salmonella* replication in human

macrophages remains unknown.

First, to determine whether glycine exerts a cytoprotective effect on THP-1

macrophages, we pretreated WT and *NAIP*[-/-] THP-1 cells with glycine prior to infection

with WT *Salmonella* and assayed for downstream inflammasome responses. Notably,

WT and *NAIP*[-/-] cells pretreated with glycine exhibited significantly decreased cell death

and a small defect in IL-1β release following infection compared to infected cells treated

with the vehicle control (Figure 4 – figure supplement 1A-B). Release of the

inflammasome-independent cytokine TNF-α was unaffected by glycine treatment

(Figure 4 – figure supplement 1C). Altogether, these results indicate that glycine

prevents cell lysis and limits IL-1 release in THP-1 macrophages upon *Salmonella*

infection.

Given glycine's cytoprotective effect on infected THP-1 macrophages, we next

sought to determine whether glycine impacts intracellular *Salmonella* burdens in human

macrophages. THP-1 macrophages pretreated with glycine retained significantly higher

intracellular bacterial burdens at 6 hpi compared to cells given the vehicle control

(Figure 4 – figure supplement 1D). We did not detect any significant differences in

bacterial uptake (Figure 4 – figure supplement 1E). Moreover, WT and $NAIP^{-/-}$ THP-1

cells pretreated with glycine exhibited a greater increase in Salmonella CFUs compared

to cells treated with the vehicle control (Figure 4 – figure supplement 1F), suggesting

that glycine treatment impedes intracellular control of Salmonella. Therefore,

cytoprotection by glycine appears to interfere with the restriction of Salmonella burdens

in human macrophages.

Next, to genetically test the role of NINJ1, we transfected WT THP-1

macrophages with small interfering RNA (siRNA) targeting NINJ1 or control scrambled

siRNA. WT cells treated with control siRNA exhibited robust IL-1β release and

cytotoxicity upon infection with WT Salmonella (Figure 4A-B). However, knockdown of

NINJ1 resulted in a significant decrease, yet not complete abrogation, of cytotoxicity in

infected cells (Figure 4B). Knockdown of NINJ1 also led to defect in IL-1β secretion

(Figure 4A). In these experiments, we observed efficient siRNA-mediated knockdown of

NINJ1 ranging from 77% to 85%. Collectively, these data suggest a critical role for

NINJ1 in contributing to IL-1 release and cell death during Salmonella infection in

human macrophages.

We then enumerated bacterial CFUs to assess whether NINJ1 impacts

intracellular Salmonella replication. At 1 hpi, we did not observe any differences in

bacterial uptake between WT THP-1 cells treated with control siRNA or NINJ1 siRNA

(Figure 4C). However, at 6 hpi, we found that cells treated with NINJ1 siRNA contained

higher intracellular bacterial burdens than cells treated with control siRNA (Figure 4D).

Furthermore, there was a greater increase in bacterial CFUs at 6 hpi in *NINJ1* siRNA-

treated cells compared to control siRNA-treated cells (Figure 4E). Together, these

findings suggest that NINJ1 contributes to intracellular bacterial control.

<u>Inflammasome activation primarily controls cytosolic *Salmonella* replication in human</u>

<u>macrophages</u>

In epithelial cells, WT *Salmonella* can replicate in both vacuolar and cytosolic

compartments, specifically hyperreplicating in the cytosol (Knodler, Crowley, et al.,

2014; Knodler et al., 2010; Knodler, Nair, et al., 2014; Malik-Kale et al., 2012). However,

in murine macrophages, WT *Salmonella* appears to replicate exclusively in vacuoles

(Beuzón et al., 2002; Thurston et al., 2016). *Salmonella* lacking the SPI-2 effector SifA

(Δ*sifA*), which is required for SCV membrane stability, frequently enter the cytosol and

cannot replicate efficiently in murine macrophages (Beuzón et al., 2000, 2002; Thurston

et al., 2016). Whether *Salmonella* replicates within vacuoles or the cytosol of human

macrophages remains largely unknown. One study suggested that a small but

significant proportion of *Salmonella* are exposed to the cytosol in THP-1 macrophages

(Fisch et al., 2020). Our recently published data and current findings indicate that when

human macrophages fail to undergo robust inflammasome responses, *Salmonella*

hyperreplicates to large numbers within these cells (Naseer, Egan, et al., 2022). The

hyperreplication that we observed in human macrophages was reminiscent of previous

reports describing *Salmonella* hyperreplication within the cytosol of IECs (Knodler,

Crowley, et al., 2014; Knodler et al., 2010; Knodler, Nair, et al., 2014; Malik-Kale et al.,

2012). Thus, we hypothesized that inflammasome activation limits *Salmonella*

replication in both SCVs and the host cell cytosol, and that in the absence of

inflammasome responses, *Salmonella* hyperreplicates to large numbers within the host

cell cytosol and also exhibits increased replication within SCVs.

First, to test whether inflammasome activation restricts the ability of *Salmonella*

to replicate in vacuolar and cytosolic compartments of human macrophages, we used a

chloroquine (CHQ) resistance assay. As a weak base, CHQ accumulates in endosomal

compartments, including the SCV, without entering the cytosol of host cells (Klein,

Powers, et al., 2017; Knodler, Nair, et al., 2014; Steinberg, 1994). Thus, vacuolar

bacteria are CHQ-sensitive, while cytosolic bacteria are CHQ-resistant (Knodler, Nair, et

al., 2014). We infected THP-1 macrophages with WT *Salmonella* and treated the

infected cells with CHQ. Then, we determined the bacterial CFUs at 6 hpi, quantifying

the numbers of vacuolar bacteria (CHQ-sensitive) and cytosolic bacteria (CHQ-

resistant). We found that WT THP-1 cells contained mostly vacuolar *Salmonella* while

also retaining some cytosolic *Salmonella* (Figure 5A), suggesting that a subset of

Salmonella dwells in the cytosol of THP-1 macrophages, in agreement with a previous

study (Fisch et al., 2020). In *NAIP*[-/-] and *CASP1*[-/-] THP-1 cells, we observed slightly

higher burdens of vacuolar *Salmonella* compared to WT cells (Figure 5A). Strikingly, we

also observed significantly larger numbers of cytosolic *Salmonella* in *NAIP*[-/-] and

CASP1[-/-] cells compared to WT cells (Figure 5A). These data indicate that

inflammasome signaling restricts *Salmonella* primarily within the host cell cytosol but

also within SCVs in human macrophages.

The SPI-1 T3SS can damage the SCV (Roy et al., 2004), and the SPI-1 T3SS

and its effectors contribute to *Salmonella* replication in the cytosol of epithelial cells

(Chong et al., 2019; Klein, Grenz, et al., 2017; Knodler, Nair, et al., 2014). So, we next

assessed whether the SPI-1 T3SS was required for the cytosolic exposure of

Salmonella in human macrophages. We infected WT and *CASP1*$^{-/-}$ THP-1 macrophages

with *Salmonella* lacking the SPI-1 T3SS translocon protein SipB (Δ*sipB*), thus

preventing SPI-1 T3SS effector translocation into host cells. While we observed a

significant increase in the amount of vacuolar and cytosolic Δ*sipB Salmonella* in

CASP1$^{-/-}$ cells compared to WT cells at 6 hpi, the numbers of cytosolic Δ*sipB* were

significantly lower than vacuolar Δ*sipB* in both WT and *CASP1*$^{-/-}$ cells (Figure 5B). We

also observed lower cytosolic Δ*sipB Salmonella* burdens compared to WT *Salmonella* in

both WT and *CASP1*$^{-/-}$ THP-1 cells (Figure 5A-B). Therefore, these results suggest that

there is a partial dependence on the SPI-1 T3SS for *Salmonella*'s cytosolic access in

human macrophages.

Since CFU assays are population-based, we next used single-cell methods to

interrogate the subcellular localization of WT *Salmonella* in human macrophages. We

relied on a strain of WT *Salmonella* that constitutively expresses mCherry and maintains

a reporter plasmid, pNF101, that expresses *gfp-ova* under the control of a promoter

responsive to the host cytosolic metabolite glucose-6-phosphate (Lau et al., 2019). We

scored the number of GFP-positive/mCherry-positive bacteria (cytosolic) and GFP-

negative/mCherry-positive bacteria (vacuolar) in WT, *NAIP*$^{-/-}$ and *CASP1*$^{-/-}$ THP-1

macrophages at 8 hpi by microscopy. We observed low numbers of both vacuolar and

cytosolic populations of *Salmonella* in WT THP-1 cells (Figure 5C-D). In contrast, *NAIP*$^{-}$

⁠/- and *CASP1⁻/⁻* THP-1 cells maintained significantly larger numbers of vacuolar

Salmonella, with a substantial increase in the numbers of cytosolic *Salmonella* (Figure

5C-D). Taken together, these results suggest that inflammasome responses primarily

restrict *Salmonella* replication within the host cell cytosol and also control bacterial

replication within SCVs in human macrophages.

We next asked whether inflammasome responses also restrict the replication of

Salmonella within the cytosol and SCV in primary human macrophages. We pretreated

hMDMs with either ZVAD, to inhibit inflammasome responses mediated by caspase

activity, or the vehicle control DMSO, and then infected the cells with WT *Salmonella*

constitutively expressing mCherry and harboring pNF101. We observed that hMDMs

treated with ZVAD contained significantly more cytosolic *Salmonella* than hMDMs

treated with DMSO (Figure 5 – figure supplement 1A-B). We observed no significant

differences between the vacuolar burdens of *Salmonella* in hMDMs pretreated with

ZVAD or DMSO (Figure 5 – figure supplement 1A-B). Thus, inflammasome responses

appear to primarily control *Salmonella* replication within the cytosol of primary human

macrophages.

<u>Inflammasome activation modulates the cytosolic exposure of *Salmonella* in human</u>

<u>macrophages</u>

Finally, to characterize the effect of inflammasomes on the subcellular niches of

Salmonella in human macrophages at higher resolution, we used transmission electron

microscopy (TEM). In WT THP-1 cells, TEM analysis revealed that the majority of

bacteria resided in a membrane-bound compartment (vacuolar, white arrows) (Figure

6A; Figure 6 – figure supplement 1B). In *CASP1*$^{-/-}$ THP-1 cells, we observed a more

mixed population of vacuolar bacteria and bacteria that were exposed to the host cell

cytosol to varying extents (Figure 6B; Figure 6 – figure supplement 1B). We observed

Salmonella free-living in the cytosol without a vacuolar membrane (fully cytosolic, black

asterisk) as well as *Salmonella* exposed to the cytosol within discontinuous vacuolar

membranes (partially cytosolic, cyan arrows) (Figure 6B; Figure 6 – figure supplement

1B). We observed a greater proportion of cytosol-exposed *Salmonella* in *CASP1*$^{-/-}$ cells

compared to WT cells (Figure 6 – figure supplement 1A). Strikingly, these TEM results

revealed distinct intracellular populations of *Salmonella* in human macrophages,

providing further evidence that inflammasome signaling controls the replicative niches

that *Salmonella* occupies in human macrophages.

447   We next sought to further characterize the cytosolic exposure of *Salmonella* in

CASP1$^{-/-}$ THP-1 cells using electron tomography (ET), a technique that can reveal three-

dimensional (3D) detail about SCV membranes. Indeed, we confirmed the presence of

Salmonella in SCVs with varying degrees of vacuolar membrane discontinuities,

yielding various extents of cytosolic exposure (Figure 6C; Figure 6 – figure supplement

1C; Video 1). Altogether, our TEM and ET data reveal the full spectrum of cytosolic

exposure for *Salmonella* in *CASP1*$^{-/-}$ cells as a result of vacuolar membrane

discontinuities. Overall, our findings indicate that inflammasome responses play a role

in modulating the subcellular populations of *Salmonella*, thereby controlling the number

of *Salmonella* able to replicate within the cytosol and SCV in human macrophages.

Discussion

Our data reveal that inflammatory caspases and downstream cell lysis mediators

are required to restrict *Salmonella* replication in human macrophages. Furthermore, our

findings indicate that inflammasomes restrict *Salmonella* hyperreplication within the

cytosol of human macrophages. Caspase-1 is required for inflammasome responses

and control of intracellular *Salmonella* replication early during infection. In contrast,

caspase-4 contributed minimally early during infection and instead played a larger role

in inflammasome responses and restriction of *Salmonella* replication at later timepoints.

We also found that GSDMD and NINJ1 were required for cell death, IL-1 cytokine

release, and control of *Salmonella* replication. Finally, in the absence of these

inflammasome components and effectors in human macrophages, we observed a

hyperreplicating population of *Salmonella* within the cytosol, as well as increased

bacterial loads within the SCV, suggesting that inflammasome activation prevents

hyperreplication of *Salmonella* within the cytosol and increased replication within SCVs.

There are multiple downstream consequences of inflammasome activation,

including pyroptosis, cytokine secretion, phagolysosomal fusion, the formation of pore-

induced intracellular traps (PITs), and the generation of reactive oxygen species (ROS),

each of which could be responsible for the control of intracellular *Salmonella* replication

in human macrophages. Previous studies have suggested that inflammatory caspases

limit the intracellular replication of WT *Salmonella* in human epithelial cells and murine

macrophages through host cell death (Aachoui et al., 2013; Holly et al., 2020; Knodler,

Crowley, et al., 2014). Interestingly, in murine macrophages, the restriction of a mutant

strain of *Salmonella* that frequently enters the cytosol is dependent on caspase-1/11 but

independent of IL-1 signaling and host cell death (Thurston et al., 2016). Moreover,

caspase-1/11-dependent production of mitochondrial ROS and hydrogen peroxide

contributes to the control of WT *Salmonella* replication in SCVs in murine macrophages

(Man, Ekpenyong, et al., 2014). Future studies investigating the downstream

consequences of inflammasome activation will help elucidate the mechanisms by which

inflammatory caspases and cell lysis mediators control intracellular *Salmonella*

replication in human macrophages.

Whereas caspase-1 is the primary caspase that mediates inflammasome

responses and control of *Salmonella* burdens in human macrophages, our data indicate

that caspase-4 also plays a role at a later stage of infection. Whether caspase-4 is

activated by vacuolar *Salmonella* or hyperreplicating cytosolic *Salmonella* in human

macrophages remains an open question. Other host immune factors may also

contribute to caspase-4-dependent inflammasome responses to *Salmonella*. Guanylate

binding proteins (GBPs) can modulate inflammasome responses to intracellular LPS

and impact intracellular bacterial replication (Degrandi et al., 2007; Fisch et al., 2020;

Kim et al., 2011; Kutsch et al., 2020; Meunier et al., 2014; Pilla et al., 2014; J. C. Santos

et al., 2020; Tietzel et al., 2009; Wandel et al., 2020).

The impact of GSDMD on the restriction of intracellular *Salmonella* replication

may be due to downstream responses, such as IL-1 release and pyroptosis. Moreover,

pyroptosis can induce PIT formation, which traps and damages intracellular bacteria,

rendering them more susceptible to immune defenses, such as neutrophil-mediated

killing (Jorgensen et al., 2016). It is possible that pyroptosis leads to the development of

PITs in human macrophages during *Salmonella* infection, thereby limiting *Salmonella*'s

intracellular replication. In addition, the cleaved N-terminal fragment of GSDMD can

directly kill bacteria (Ding et al., 2016; Liu et al., 2016; Wang et al., 2019), so perhaps

GSDMD directly targets *Salmonella* to mediate restriction. It remains unknown whether

other cell lysis factors, like NINJ1, together with GSDMD, directly bind the SCV,

affecting vacuolar integrity and thus influencing intracellular *Salmonella* replication.

Our data indicate that inflammasome responses primarily control *Salmonella*

replication within the cytosol of human macrophages and also within SCVs. How

inflammasome activation inhibits *Salmonella* replication in the cytosol of human

macrophages remains unknown. *Salmonella* hyperreplicates in the cytosol of human

epithelial cells, which fail to undergo caspase-1-dependent inflammasome responses

(Holly et al., 2020; Knodler, Crowley, et al., 2014; Knodler et al., 2010; Knodler, Nair, et

al., 2014; Naseer, Zhang, et al., 2022). Our study suggests that *CASP1*$^{-/-}$ THP-1

macrophages behave similarly to human IECs, as they support *Salmonella*

hyperreplication in the cytosol in the absence of caspase-1. Inflammasome activation

could curtail cytosolic replication of *Salmonella* through host cell death, direct GSDMD

targeting of cytosolic *Salmonella,* and/or another effector mechanism. Alternatively,

inflammasome activation could regulate the cytosolic access of *Salmonella* through host

factors that directly damage the SCV, such as pore-forming proteins like GSDMD.

The bacterial factors that facilitate *Salmonella*'s cytosolic lifestyle in human

macrophages remain largely unknown. Our findings indicate that the SPI-1 T3SS is

partially required for *Salmonella* to access the cytosol in THP-1 macrophages. In

epithelial cells, the SPI-1 T3SS SopB and SipA enable *Salmonella* to efficiently colonize

and replicate in the cytosol (Chong et al., 2019; Klein, Grenz, et al., 2017). Future

studies are needed to elucidate whether these same effectors play a role in supporting

Salmonella's cytosolic replication in human macrophages.

Of note, our data indicate that there is considerable single-cell heterogeneity in

terms of total intracellular bacterial burdens as well as proportions of vacuolar and

cytosolic *Salmonella* in human macrophages using single-cell microscopic analysis.

Several factors could account for this phenotypic heterogeneity, including expression

levels of inflammasome components, the amount of translocated *Salmonella* ligands, or

the extent of inflammasome activation, all of which might differ at a single cell level and

are masked in bulk-population assays. Notably, heterogeneity in *Salmonella* gene

expression, intracellular *Salmonella* populations, and intracellular *Salmonella*

proliferation and viability has been previously observed in human epithelial cells and

murine macrophages (Helaine et al., 2010, 2014; Knodler, Nair, et al., 2014; Malik-Kale

et al., 2012; Powers et al., 2021).

Collectively, our results reveal that inflammasome responses restrict intracellular

Salmonella replication, particularly within the cytosol of human macrophages. These

findings are in contrast to mouse macrophages, where inflammasomes are activated

but only marginally restrict the intracellular replication of WT *Salmonella*. Our findings

provide insight into how human macrophages leverage inflammasomes to restrict

Salmonella intracellular replication. Moreover, our work offers a basis for future studies

to investigate how inflammasome activation modulates the subcellular localization of

bacterial replicative niches within host cells.

<u>Materials and Methods</u>

Ethics statement

All experiments on primary human monocyte-derived macrophages (hMDMs) were

performed in compliance with the requirements of the US Department of Health and

Human Services and the principles expressed in the Declaration of Helsinki. hMDMs

were derived from samples obtained from the University of Pennsylvania Human

Immunology Core, and they are considered to be a secondary use of deidentified

human specimens and are exempt via Title 55 Part 46, Subpart A of 46.101 (b) of the

Code of Federal Regulations.

Cell culture of THP-1 cells

THP-1 cells (TIB-202; American Type Culture Collection) were maintained in RPMI

supplemented with 10% (vol/vol) heat-inactivated FBS, 0.05 nM β-mercaptoethanol, 100

IU/mL penicillin, and 100 µg/mL streptomycin at 37°C in a humidified incubator. Two

days prior to experimentation, the cells were replated in media without antibiotics in a

48-well plate at a concentration of 2×10^5 cells/well or in a 24-well plate at a

concentration of 3.5×10^5 cells per well and incubated with phorbol 12-myristate 13-

acetate (PMA) for 24 hours to allow differentiation into macrophages. Then,

macrophages were primed with 100 ng/mL Pam3CSK4 (Invivogen) for 16 to 20 hours

prior to bacterial infections. For fluorescence microscopy experiments, cells were plated

on glass coverslips in a 24-well plate.

Cell culture of primary human monocyte-derived macrophages (hMDMs)

Purified human monocytes from deidentified healthy human donors were obtained from

the University of Pennsylvania Human Immunology Core. The monocytes were cultured

in RPMI supplemented with 10% (vol/vol) heat-inactivated FBS, 2 mM L-glutamine, 100

IU/mL penicillin, 100 µg/ml streptomycin, and 50 ng/ml recombinant human M-CSF

(Gemini Bio-Products). Cells were cultured for 4 days in 10 mL of media in at a

concentration of 4-5 × 10^5 cells/mL in 10 cm-dishes. Then, 10 mL of fresh growth media

was added for an additional 2 days to promote complete differentiation into

macrophages. One day prior to infection, the cells were rinsed with cold PBS, gently

detached with trypsin-EDTA (0.05%) and replated in media without antibiotics and with

25 ng/mL human M-CSF in a 48-well plate at a concentration of 1 × 10^5 cells per well or

in a 24-well plate at a concentration of 2 × 10^5 cells per well. For fluorescence

microscopy, cells were replated on glass coverslips in a 24-well plate. For experiments

involving LPS, cells were primed with 500 ng/mL LPS (Sigma-Aldrich) for 3 hours prior

to bacterial infections.

<u>Inhibitor experiments</u>

Human macrophages were treated 1 hour prior to infection at the indicated

concentrations with the following inhibitors: 20 µM of the pan-caspase inhibitor Z-

VAD(OMe)-FMK (SM Biochemicals; SMFMK001), 25 µM of the caspase-1 inhibitor Ac-

YVAD-cmk (Sigma-Aldrich; SML0429), and 40 µM of the GSDMD inhibitor disulfiram

(Sigma). DMSO treatment was used as a vehicle control with these inhibitors. To

prevent cell lysis, cells were treated with 20 mM glycine (Fisher Scientific) for 30

minutes prior to infection. Distilled water was used as a vehicle control.

Bacterial strains and growth conditions

Salmonella enterica serovar Typhimurium SL1344 WT was routinely grown shaking

overnight at 37°C in Luria-Bertani (LB) broth with streptomycin (100 μg/mL). For

infection of cultured cells, the overnight culture was diluted in LB with streptomycin

(100 μg/mL) containing 300 mM NaCl and grown standing for 3 hours at 37°C to induce

SPI-1 expression (Lee & Falkow, 1990).

SL1344 WT *glmS::Ptrc-mCherryST::FRT* pNF101 (Lau et al., 2019) was kindly provided

by Dr. Leigh Knodler. This strain constitutively expresses *mCherry* that is

chromosomally encoded. It also harbors the *PuhpT-gfpova* plasmid, pNF101, where

expression of GFP is under the control of the glucose-6-phosphate responsive *uhpT*

promoter derived from *Shigella flexneri*. SL1344 WT *glmS::Ptrc-mCherryST::FRT*

pNF101 was grown shaking overnight at 37°C in Luria-Bertani (LB) broth with

streptomycin (100 μg/mL) and ampicillin (100 μg/mL). For SPI-1 induction prior to

infection, the overnight culture was diluted in LB with streptomycin (100 μg/mL) and

ampicillin (100 μg/mL) that also contained 300 mM NaCl and then grown standing for 3

hours at 37°C (Lee & Falkow, 1990).

Bacterial infections

Overnight cultures of *Salmonella* were diluted into LB broth with streptomycin

(100 μg/mL) containing 300 mM NaCl and grown for 3 hours standing at 37°C to induce

SPI-1 gene expression (Lee & Falkow, 1990). Bacterial cultures were then pelleted at

6,010 × g for 3 minutes, washed once with PBS, and resuspended in PBS. Human

macrophages were infected with *Salmonella* at a multiplicity of infection (MOI) of 20.

Infected cells were centrifuged at 290 × g for 10 minutes and incubated at 37°C. At 30

minutes post-infection, the cells were treated with 100 ng/mL of gentamicin to kill any

extracellular *Salmonella*. Then, the infection proceeded at 37°C for 6 to 8 hours, as

indicated. For all experiments, control cells were mock-infected with PBS.

<u>Bacterial intracellular burden assay</u>

Cells were infected with WT *Salmonella* as described above at an MOI of 20. Then, 30

minutes post-infection, cells were treated with 100 μg/ml of gentamicin to kill any

extracellular bacteria. 1 hour post-infection, the media was replaced with fresh media

containing 10 μg/ml of gentamicin. At the indicated time points, the infected cells were

lysed with PBS containing 0.5% Triton to collect all intracellular *Salmonella*. Harvested

bacteria were serially diluted in PBS and plated on LB agar plates containing

streptomycin (100 μg/ml) to enumerate colony forming units (CFUs). Plates were

incubated overnight at 37°C and CFUs were subsequently counted.

<u>Chloroquine (CHQ) Resistance Assay</u>

THP-1 macrophages were infected in 48-well plates as described above. For each

timepoint, triplicate wells were incubated in the presence of CHQ (500 μM) and

gentamicin (100 ng/mL) for 1 hour to quantify the CHQ-resistant bacteria (Bârzu et al.,

1997; Fernandez et al., 2001; Klein, Powers, et al., 2017; Knodler, Nair, et al., 2014;

Zychlinsky et al., 1994). Another triplicate wells were incubated with gentamicin (100

ng/mL) only to quantify the total intracellular bacteria. At the indicated time points, the

infected cells were lysed with PBS containing 0.5% Triton to collect intracellular

Salmonella. Harvested bacteria were serially diluted in PBS and plated on LB agar

plates containing streptomycin (100 µg/ml) to enumerate colony forming units (CFUs).

Plates were incubated overnight at 37°C and CFUs were subsequently counted.

ELISAs

Harvested supernatants from infected human macrophages were assayed for cytokine

levels using ELISA kits for human IL-1α (R&D Systems), IL-18 (R&D Systems), IL-1β

(BD Biosciences), and TNF-α (R&D Systems).

LDH cytotoxicity assays

Harvested supernatants from infected human macrophages were assayed for

cytotoxicity by quantifying the loss of cellular membrane integrity via lactate

dehydrogenase (LDH) activity. LDH release was measured using an LDH Cytotoxicity

Detection Kit (Clontech) according to the manufacturer's instructions and normalized to

mock-infected cells.

siRNA-mediated knockdown of genes

All Silencer Select siRNA oligos were purchased from Ambion (Life Technologies).

Individual siRNA targeting *NINJ1* (ID# s9556) was used. The two Silencer Select

negative control siRNAs (Silencer Select Negative Control No. 1 siRNA and Silencer

Select Negative Control No. 2 siRNA) were used as a control. Three days prior to the

infection, 30 nM of siRNA was transfected into the human macrophages using

Lipofectamine RNAiMAX transfection reagent (Thermo Fisher Scientific) following the

manufacturer's protocol. Then, 24 hours after transfection, the media was replaced with

fresh media containing antibiotics. Finally, 16 hours before infection, the media was

replaced with fresh antibiotic-free media containing 100 ng/ml Pam3CSK4.

Quantitative RT-PCR Analysis

RNA was isolated using the RNeasy Plus Mini Kit (Qiagen) following the manufacturer's

protocol. Human macrophages were lysed in 350 µL RLT buffer with β-mercaptoethanol

and centrifuged through a QIAshredder spin column (Qiagen). cDNA was synthesized

from isolated RNA using SuperScript II Reverse Transcriptase (Invitrogen) following the

manufacturer's protocol. Quantitative PCR was conducted with the CFX96 real-time

system from Bio-Rad using the SsoFast EvaGreen Supermix with Low ROX (Bio-Rad).

To calculate knockdown efficiency, mRNA levels of siRNA-treated cells were

normalized to the housekeeping gene *HPRT* and control siRNA-treated cells using the

$2^{-\Delta\Delta CT}$ (cycle threshold) method (Livak & Schmittgen, 2001). The following primers from

PrimerBank were used. The PrimerBank identifications are *NINJ1* (148922910c1),

CASP4 (73622124c2) and *HPRT* (164518913c1); all primers listed as 5′–3′:

NINJ1 forward: TCAAGTACGACCTTAACAACCCG

NINJ1 reverse: TGAAGATGTTGACTACCACGATG

CASP4 forward: TCTGCGGAACTGTGCATGATG

CASP4 reverse: TGTGTGATGAAGATAGAGCCCAT

HPRT forward: CCTGGCGTCGTGATTAGTGAT

HPRT reverse: AGACGTTCAGTCCTGTCCATAA

<u>Immunoblot analysis</u>

Cell lysates were harvested for immunoblot analysis by adding 1X SDS/PAGE sample

buffer to cells. All protein samples (lysates) were boiled for 5 minutes. Samples were

separated by SDS/PAGE on a 12% (vol/vol) acrylamide gel and transferred to PVDF

Immobilon-P membranes (Millipore). Primary antibodies specific for caspase-4 (4450S;

Cell Signaling) and β-actin (4967L; Cell Signaling) and HRP-conjugated secondary

antibodies anti-mouse IgG (F00011; Cell Signaling) and anti-rabbit IgG (7074S; Cell

Signaling) were used. ECL Western Blotting Substrate (Pierce Thermo Scientific) was

used as the HRP substrate for detection.

<u>Fluorescent microscopy of intracellular *Salmonella*</u>

Primary hMDMs or THP-1 cells were plated on glass coverslips in a 24-well plate as

described above. Cells were either infected with WT *Salmonella* constitutively

expressing GFP [Sl1344 harboring pFPV25.1 (Valdivia & Falkow, 1996)], or WT

Salmonella constitutively expressing mCherry with a cytosolic GFP reporter [SL1344

glmS::Ptrc-mCherryST::FRT pNF101 (Lau et al., 2019)] at an MOI of 20 as described

above. At the indicated timepoints following infection, cells were washed 2 times with

PBS and then fixed with 4% paraformaldehyde for 10 minutes. Following fixation, cells

were mounted on glass slides with DAPI mounting medium (Sigma Fluoroshield).

Coverslips were imaged on an inverted fluorescence microscope (IX81; Olympus), and

the images were collected using a high-resolution charge-coupled-device camera

(FAST1394; QImaging) at a magnification of 100×. All images were analyzed and

presented using SlideBook (version 5.0) software (Intelligent Imaging Innovations, Inc.)

and ImageJ software. For experiments with WT *Salmonella* constitutively expressing

GFP, the proportion of infected cells containing GFP-expressing *Salmonella* (green)

were scored by counting 50 infected cells per coverslip. 150 total infected cells were

scored for each condition. For experiments with WT *Salmonella* constitutively

expressing mCherry with a cytosolic GFP reporter, the proportion of infected cells

containing GFP-positive *Salmonella* (cytosolic) and GFP-negative, mCherry-positive

Salmonella (vacuolar) were scored by counting 50 infected cells per coverslip. 150 total

infected cells were scored for each condition.

<u>Transmission electron microscopy</u>

Two days prior to experimentation, 2×10^6 cells THP-1 cells were replated in 10-cm

dishes in media without antibiotics and incubated with phorbol 12-myristate 13-acetate

(PMA) for 24 hours to allow differentiation into macrophages. Then, macrophages were

primed with 100 ng/mL Pam3CSK4 (Invivogen) for 16 hours prior to bacterial infection.

Cells were infected with WT *Salmonella* at an MOI of 20 as described above. At 8 hpi,

the media was aspirated, and the cells were fixed with 2.5% glutaraldehyde, 2.0%

paraformaldehyde in 0.1M sodium cacodylate buffer, pH 7.4. Then, at the Electron

Microscopy Resource Laboratory in the Perelman School of Medicine, after subsequent

buffer washes, the samples were post-fixed in 2.0% osmium tetroxide with 1.5%

K3Fe(CN)6 for 1 hour at room temperature, and rinsed in dH2O. After dehydration

through a graded ethanol series, the tissue was infiltrated and embedded in EMbed-812

(Electron Microscopy Sciences, Fort Washington, PA). Thin sections were stained with

uranyl acetate and SATO lead and examined with a JEOL 1010 electron microscope

fitted with a Hamamatsu digital camera and AMT Advantage NanoSprint500 software.

<u>Electron tomography</u>

Electron tomography was performed at room temperature on a ThermoFisher Krios G3i

TEM equipped with a 300 keV field emission gun. Imaging was performed using the

SerialEM software (Mastronarde, 2005) on a K3 direct electron detector (Xuong et al.,

2007) (Gatan Inc., Pleasanton, CA, USA) operated in the electron-counted mode. We

additionally used the Gatan Imaging Filter (Gatan Inc., Pleasanton, CA, USA) with a slit

width of 20 eV to increase contrast by removing inelastically scattered electrons

(Krivanek et al., 1995). After initially assessing the infected macrophages at lower

magnifications, tilt series were collected at a magnification of 26,000X (with a

corresponding pixel size of 3.38 Å) and a defocus of −6 µm. Precooking the target areas

with a total dosage of 1000-1500 e/Å^2 was required to minimize sample shrinking and

drifting during tilt series collection. A bi-directional tilt scheme was employed with 2°

increments and 120° span (−60° to +60°). The cumulative dose of each tilt-series was in

the order of ~500 e−/Å2. Once acquired, tilt series were aligned (using the patch

tracking function) and reconstructed into tomograms, both using the IMOD software

package (Kremer et al., 1996). After careful assessment of the 3-dimensional (3-D)

tomograms, optimal 2-D slices were chosen for figure presentations. Color overlays

provide additional assistance with interpretation.

Statistical analysis

Prism 9.5.0 (GraphPad Software) was utilized for the graphing of data and all statistical

analyses. Statistical significance for experiments with THP-1 macrophages and hMDMs

was determined using the appropriate test and are indicated in each figure legend.

Differences were considered statistically significant if the *p* value was <0.05.

Acknowledgements

We thank members of Igor Brodsky's and Sunny Shin's laboratories for their scientific

discussions and continuous support. We thank Leigh Knodler for providing SL1344

glmS::Ptrc-mCherryST::FRT pNF101 and Cornelius Taabazuing for providing *CASP1*[-/-]

and *GSDMD*[-/-] THP-1 cells. We thank the Human Immunology Core of the Penn Center

for AIDS Research and Abramson Cancer Center for providing purified primary human

monocytes. We also thank the Electron Microscopy Resource Lab (EMRL) at the

Perelman School of Medicine, University of Pennsylvania for TEM specimen

processing, sectioning, and staining along with microscopy training. Additionally, we

thank the EMRL for allowing us access and usage of the transmission electron

microscope, JEOL JEM-1010. Lastly, we thank Gordon Ruthel at the Penn Vet Imaging

Core and Ronit Schwartz for providing helpful insights on fluorescence microscopy.

This work was supported by National Institutes of Health (NIH)/National Institute of

Allergy and Infectious Diseases (NIAID) grants: AI151476, AI118861, and AI123243

(S.S.), AI128630, AI163596, and AI139102 (I.E.B). This work was also supported by the

Burroughs-Wellcome Fund Investigators in the Pathogenesis of Infectious Disease

Award (S.S. and I.E.B.), the National Science Foundation Graduate Fellowships DGE-

1321851 (M.S.E.) and DGE-1845298 (A.R.B.), the NIH/NIGMS grant T32GM07229

(E.A.O.), the David and Lucile Packard Fellowship for Science and Engineering 2019–

69645 (Y.-W.C.), and the Pennsylvania Department of Health FY19 Health Research

Formula Fund (Y.-W.C.). The funders had no role in study design, data collection and

analysis, decision to publish, or preparation of the manuscript.

786
<u>References</u>

Aachoui, Y., Leaf, I. A., Hagar, J. A., Fontana, M. F., Campos, C. G., Zak, D. E., Tan, M.
H., Cotter, P. A., Vance, R. E., Aderem, A., & Miao, E. A. (2013). Caspase-11
Protects Against Bacteria That Escape the Vacuole. *Science*, *339*(6122), 975–
978. https://doi.org/10.1126/science.1230751
Agard, N. J., Maltby, D., & Wells, J. A. (2010). Inflammatory Stimuli Regulate Caspase
Substrate Profiles. *Molecular & Cellular Proteomics*, *9*(5), 880–893.
https://doi.org/10.1074/mcp.M900528-MCP200
Agbor, T. A., & McCormick, B. A. (2011). Salmonella effectors: Important players
modulating host cell function during infection: Salmonella effectors. *Cellular*
Microbiology, *13*(12), 1858–1869. https://doi.org/10.1111/j.1462-
5822.2011.01701.x
Baker, P. J., Boucher, D., Bierschenk, D., Tebartz, C., Whitney, P. G., D'Silva, D. B.,
Tanzer, M. C., Monteleone, M., Robertson, A. A. B., Cooper, M. A., Alvarez-Diaz,
S., Herold, M. J., Bedoui, S., Schroder, K., & Masters, S. L. (2015). NLRP3
inflammasome activation downstream of cytoplasmic LPS recognition by both
caspase-4 and caspase-5: Innate immunity. *European Journal of Immunology*,
45(10), 2918–2926. https://doi.org/10.1002/eji.201545655
Bârzu, S., Benjelloun-Touimi, Z., Phalipon, A., Sansonetti, P., & Parsot, C. (1997).
Functional analysis of the Shigella flexneri IpaC invasin by insertional
mutagenesis. *Infection and Immunity*, *65*(5), 1599–1605.
https://doi.org/10.1128/iai.65.5.1599-1605.1997
Beuzón, C. R., Méresse, S., Unsworth, K. E., Ruíz-Albert, J., Garvis, S., Waterman, S.
R., Ryder, T. A., Boucrot, E., & Holden, D. W. (2000). Salmonella maintains the
integrity of its intracellular vacuole through the action of SifA. *The EMBO Journal*,
19(13), 3235–3249. https://doi.org/10.1093/emboj/19.13.3235
Beuzón, C. R., Salcedo, S. P., & Holden, D. W. (2002). Growth and killing of a
Salmonella enterica serovar Typhimurium sifA mutant strain in the cytosol of
different host cell lines. *Microbiology*, *148*(9), 2705–2715.
https://doi.org/10.1099/00221287-148-9-2705

Bierschenk, D., Monteleone, M., Moghaddas, F., Baker, P. J., Masters, S. L., Boucher, D., & Schroder, K. (2019). The *Salmonella* pathogenicity island-2 subverts human NLRP3 and NLRC4 inflammasome responses. *Journal of Leukocyte Biology*, *105*(2), 401–410. https://doi.org/10.1002/JLB.MA0318-112RR

Bjanes, E., Sillas, R. G., Matsuda, R., Demarco, B., Fettrelet, T., DeLaney, A. A., Kornfeld, O. S., Lee, B. L., Rodríguez López, E. M., Grubaugh, D., Wynosky-Dolfi, M. A., Philip, N. H., Krespan, E., Tovar, D., Joannas, L., Beiting, D. P., Henao-Mejia, J., Schaefer, B. C., Chen, K. W., … Brodsky, I. E. (2021). Genetic targeting of Card19 is linked to disrupted NINJ1 expression, impaired cell lysis, and increased susceptibility to Yersinia infection. *PLOS Pathogens*, *17*(10), e1009967. https://doi.org/10.1371/journal.ppat.1009967

Borges, J. P., Sætra, R. S., Volchuk, A., Bugge, M., Devant, P., Sporsheim, B., Kilburn, B. R., Evavold, C. L., Kagan, J. C., Goldenberg, N. M., Flo, T. H., & Steinberg, B. E. (2022). Glycine inhibits NINJ1 membrane clustering to suppress plasma membrane rupture in cell death. *ELife*, *11*, e78609. https://doi.org/10.7554/eLife.78609

Broz, P., & Dixit, V. M. (2016). Inflammasomes: Mechanism of assembly, regulation and signalling. *Nature Reviews Immunology*, *16*(7), 407–420. https://doi.org/10.1038/nri.2016.58

Broz, P., Newton, K., Lamkanfi, M., Mariathasan, S., Dixit, V. M., & Monack, D. M. (2010). Redundant roles for inflammasome receptors NLRP3 and NLRC4 in host defense against Salmonella. *Journal of Experimental Medicine*, *207*(8), 1745–1755. https://doi.org/10.1084/jem.20100257

Broz, P., Ruby, T., Belhocine, K., Bouley, D. M., Kayagaki, N., Dixit, V. M., & Monack, D. M. (2012). Caspase-11 increases susceptibility to Salmonella infection in the absence of caspase-1. *Nature*, *490*(7419), 288–291. https://doi.org/10.1038/nature11419

Brumell, J. H., Goosney, D. L., & Finlay, B. B. (2002). SifA, a Type III Secreted Effector of *Salmonella typhimurium* , Directs *Salmonella* -Induced Filament (Sif) Formation Along Microtubules: **Sif Formation Along Microtubules**. *Traffic*, *3*(6), 407–415. https://doi.org/10.1034/j.1600-0854.2002.30604.x

Brumell, J. H., Tang, P., Zaharik, M. L., & Finlay, B. B. (2002). Disruption of the Salmonella-containing vacuole leads to increased replication of Salmonella enterica serovar typhimurium in the cytosol of epithelial cells. *Infection and Immunity*, *70*(6), 3264–3270. https://doi.org/10.1128/iai.70.6.3264-3270.2002

Carvalho, F. A., Nalbantoglu, I., Aitken, J. D., Uchiyama, R., Su, Y., Doho, G. H., Vijay-Kumar, M., & Gewirtz, A. T. (2012). Cytosolic flagellin receptor NLRC4 protects mice against mucosal and systemic challenges. *Mucosal Immunology*, *5*(3), 288–298. https://doi.org/10.1038/mi.2012.8

Casson, C. N., Yu, J., Reyes, V. M., Taschuk, F. O., Yadav, A., Copenhaver, A. M., Nguyen, H. T., Collman, R. G., & Shin, S. (2015). Human caspase-4 mediates noncanonical inflammasome activation against gram-negative bacterial pathogens. *Proceedings of the National Academy of Sciences*, *112*(21), 6688–6693. https://doi.org/10.1073/pnas.1421699112

Chong, A., Starr, T., Finn, C. E., & Steele-Mortimer, O. (2019). A role for the *Salmonella* Type III Secretion System 1 in bacterial adaptation to the cytosol of epithelial

cells. *Molecular Microbiology*, *112*(4), 1270–1283.
https://doi.org/10.1111/mmi.14361

Cirillo, D. M., Valdivia, R. H., Monack, D. M., & Falkow, S. (1998). Macrophage-dependent induction of the *Salmonella* pathogenicity island 2 type III secretion system and its role in intracellular survival. *Molecular Microbiology*, *30*(1), 175–188. https://doi.org/10.1046/j.1365-2958.1998.01048.x

Crowley, S. M., Han, X., Allaire, J. M., Stahl, M., Rauch, I., Knodler, L. A., & Vallance, B. A. (2020). Intestinal restriction of Salmonella Typhimurium requires caspase-1 and caspase-11 epithelial intrinsic inflammasomes. *PLOS Pathogens*, *16*(4), e1008498. https://doi.org/10.1371/journal.ppat.1008498

Crowley, S. M., Knodler, L. A., & Vallance, B. A. (2016). Salmonella and the Inflammasome: Battle for Intracellular Dominance. In S. Backert (Ed.), *Inflammasome Signaling and Bacterial Infections* (Vol. 397, pp. 43–67). Springer International Publishing. https://doi.org/10.1007/978-3-319-41171-2_3

Degen, M., Santos, J. C., Pluhackova, K., Cebrero, G., Ramos, S., Jankevicius, G., Hartenian, E., Guillerm, U., Mari, S. A., Kohl, B., Müller, D. J., Schanda, P., Maier, T., Perez, C., Sieben, C., Broz, P., & Hiller, S. (2023). Structural basis of NINJ1-mediated plasma membrane rupture in cell death. *Nature*. https://doi.org/10.1038/s41586-023-05991-z

Degrandi, D., Konermann, C., Beuter-Gunia, C., Kresse, A., Würthner, J., Kurig, S., Beer, S., & Pfeffer, K. (2007). Extensive characterization of IFN-induced GTPases mGBP1 to mGBP10 involved in host defense. *Journal of Immunology (Baltimore, Md.: 1950)*, *179*(11), 7729–7740. https://doi.org/10.4049/jimmunol.179.11.7729

Diebolder, C. A., Halff, E. F., Koster, A. J., Huizinga, E. G., & Koning, R. I. (2015). Cryoelectron Tomography of the NAIP5/NLRC4 Inflammasome: Implications for NLR Activation. *Structure*, *23*(12), 2349–2357. https://doi.org/10.1016/j.str.2015.10.001

Ding, J., Wang, K., Liu, W., She, Y., Sun, Q., Shi, J., Sun, H., Wang, D.-C., & Shao, F. (2016). Pore-forming activity and structural autoinhibition of the gasdermin family. *Nature*, *535*(7610), 111–116. https://doi.org/10.1038/nature18590

Fernandez, I. M., Silva, M., Schuch, R., Walker, W. A., Siber, A. M., Maurelli, A. T., & McCormick, B. A. (2001). Cadaverine Prevents the Escape of *Shigella flexneri* from the Phagolysosome: A Connection between Bacterial Dissemination and Neutrophil Transepithelial Signaling. *The Journal of Infectious Diseases*, *184*(6), 743–753. https://doi.org/10.1086/323035

Fink, S. L., & Cookson, B. T. (2006). Caspase-1-dependent pore formation during pyroptosis leads to osmotic lysis of infected host macrophages. *Cellular Microbiology*, *8*(11), 1812–1825. https://doi.org/10.1111/j.1462-5822.2006.00751.x

Fisch, D., Clough, B., Domart, M.-C., Encheva, V., Bando, H., Snijders, A. P., Collinson, L. M., Yamamoto, M., Shenoy, A. R., & Frickel, E.-M. (2020). Human GBP1 Differentially Targets Salmonella and Toxoplasma to License Recognition of Microbial Ligands and Caspase-Mediated Death. *Cell Reports*, *32*(6), 108008. https://doi.org/10.1016/j.celrep.2020.108008

Franchi, L., Amer, A., Body-Malapel, M., Kanneganti, T.-D., Özören, N., Jagirdar, R., Inohara, N., Vandenabeele, P., Bertin, J., Coyle, A., Grant, E. P., & Núñez, G. (2006). Cytosolic flagellin requires Ipaf for activation of caspase-1 and interleukin 1β in salmonella-infected macrophages. *Nature Immunology*, *7*(6), 576–582. https://doi.org/10.1038/ni1346

Franchi, L., Kamada, N., Nakamura, Y., Burberry, A., Kuffa, P., Suzuki, S., Shaw, M. H., Kim, Y.-G., & Núñez, G. (2012). NLRC4-driven production of IL-1β discriminates between pathogenic and commensal bacteria and promotes host intestinal defense. *Nature Immunology*, *13*(5), 449–456. https://doi.org/10.1038/ni.2263

Franchi, L., Kanneganti, T.-D., Dubyak, G. R., & Núñez, G. (2007). Differential Requirement of P2X7 Receptor and Intracellular K+ for Caspase-1 Activation Induced by Intracellular and Extracellular Bacteria. *Journal of Biological Chemistry*, *282*(26), 18810–18818. https://doi.org/10.1074/jbc.M610762200

Frank, A., Rauen, U., & Groot, H. de. (2000). Protection by glycine against hypoxic injury of rat hepatocytes: Inhibition of ion fluxes through nonspecific leaks. *Journal of Hepatology*, *32*(1), 58–66. https://doi.org/10.1016/S0168-8278(00)80190-7

Galán, J. E. (1999). Interaction of Salmonella with host cells through the centisome 63 type III secretion system. *Current Opinion in Microbiology*, *2*(1), 46–50. https://doi.org/10.1016/S1369-5274(99)80008-3

Galán, J. E., & Collmer, A. (1999). Type III Secretion Machines: Bacterial Devices for Protein Delivery into Host Cells. *Science*, *284*(5418), 1322–1328. https://doi.org/10.1126/science.284.5418.1322

Galan, J. E., & Curtiss, R. (1989). Cloning and molecular characterization of genes whose products allow Salmonella typhimurium to penetrate tissue culture cells. *Proceedings of the National Academy of Sciences*, *86*(16), 6383–6387. https://doi.org/10.1073/pnas.86.16.6383

Galan, J. E., & Zhou, D. (2000). Striking a balance: Modulation of the actin cytoskeleton by Salmonella. *Proceedings of the National Academy of Sciences*, *97*(16), 8754–8761. https://doi.org/10.1073/pnas.97.16.8754

Garcia-del Portillo, F., & Finlay, B. B. (1995). Targeting of Salmonella typhimurium to vesicles containing lysosomal membrane glycoproteins bypasses compartments with mannose 6-phosphate receptors. *Journal of Cell Biology*, *129*(1), 81–97. https://doi.org/10.1083/jcb.129.1.81

Garcia-del Portillo, F., Zwick, M. B., Leung, K. Y., & Finlay, B. B. (1993). Salmonella induces the formation of filamentous structures containing lysosomal membrane glycoproteins in epithelial cells. *Proceedings of the National Academy of Sciences of the United States of America*, *90*(22), 10544–10548. https://doi.org/10.1073/pnas.90.22.10544

Gram, A. M., Wright, J. A., Pickering, R. J., Lam, N. L., Booty, L. M., Webster, S. J., & Bryant, C. E. (2021). *Salmonella* Flagellin Activates NAIP/NLRC4 and Canonical NLRP3 Inflammasomes in Human Macrophages. *The Journal of Immunology*, *206*(3), 631–640. https://doi.org/10.4049/jimmunol.2000382

Grandjean, T., Boucher, A., Thepaut, M., Monlezun, L., Guery, B., Faudry, E., Kipnis, E., & Dessein, R. (2017). The human NAIP-NLRC4-inflammasome senses the

Pseudomonas aeruginosa T3SS inner-rod protein. *International Immunology*, *29*(8), 377–384. https://doi.org/10.1093/intimm/dxx047

Hagar, J. A., Powell, D. A., Aachoui, Y., Ernst, R. K., & Miao, E. A. (2013). Cytoplasmic LPS Activates Caspase-11: Implications in TLR4-Independent Endotoxic Shock. *Science*, *341*(6151), 1250–1253. https://doi.org/10.1126/science.1240988

Hansen-Wester, I., Stecher, B., & Hensel, M. (2002). Type III Secretion of *Salmonella enterica* Serovar Typhimurium Translocated Effectors and SseFG. *Infection and Immunity*, *70*(3), 1403–1409. https://doi.org/10.1128/IAI.70.3.1403-1409.2002

Hausmann, A., Böck, D., Geiser, P., Berthold, D. L., Fattinger, S. A., Furter, M., Bouman, J. A., Barthel-Scherrer, M., Lang, C. M., Bakkeren, E., Kolinko, I., Diard, M., Bumann, D., Slack, E., Regoes, R. R., Pilhofer, M., Sellin, M. E., & Hardt, W.-D. (2020). Intestinal epithelial NAIP/NLRC4 restricts systemic dissemination of the adapted pathogen Salmonella Typhimurium due to site-specific bacterial PAMP expression. *Mucosal Immunology*, *13*(3), 530–544. https://doi.org/10.1038/s41385-019-0247-0

Heilig, R., Dick, M. S., Sborgi, L., Meunier, E., Hiller, S., & Broz, P. (2018). The Gasdermin-D pore acts as a conduit for IL-1β secretion in mice. *European Journal of Immunology*, *48*(4), 584–592. https://doi.org/10.1002/eji.201747404

Helaine, S., Cheverton, A. M., Watson, K. G., Faure, L. M., Matthews, S. A., & Holden, D. W. (2014). Internalization of Salmonella by macrophages induces formation of nonreplicating persisters. *Science (New York, N.Y.)*, *343*(6167), 204–208. https://doi.org/10.1126/science.1244705

Helaine, S., Thompson, J. A., Watson, K. G., Liu, M., Boyle, C., & Holden, D. W. (2010). Dynamics of intracellular bacterial replication at the single cell level. *Proceedings of the National Academy of Sciences*, *107*(8), 3746–3751. https://doi.org/10.1073/pnas.1000041107

Hensel, M., Shea, J. E., Waterman, S. R., Mundy, R., Nikolaus, T., Banks, G., Vazquez-Torres, A., Gleeson, C., Fang, F. C., & Holden, D. W. (1998). Genes encoding putative effector proteins of the type III secretion system of *Salmonella* pathogenicity island 2 are required for bacterial virulence and proliferation in macrophages. *Molecular Microbiology*, *30*(1), 163–174. https://doi.org/10.1046/j.1365-2958.1998.01047.x

Holly, M. K., Han, X., Zhao, E. J., Crowley, S. M., Allaire, J. M., Knodler, L. A., Vallance, B. A., & Smith, J. G. (2020). Salmonella enterica Infection of Murine and Human Enteroid-Derived Monolayers Elicits Differential Activation of Epithelium-Intrinsic Inflammasomes. *Infection and Immunity*, *88*(7), e00017-20. https://doi.org/10.1128/IAI.00017-20

Hornung, V., Bauernfeind, F., Halle, A., Samstad, E. O., Kono, H., Rock, K. L., Fitzgerald, K. A., & Latz, E. (2008). Silica crystals and aluminum salts activate the NALP3 inflammasome through phagosomal destabilization. *Nature Immunology*, *9*(8), 847–856. https://doi.org/10.1038/ni.1631

Hu, J. J., Liu, X., Xia, S., Zhang, Z., Zhang, Y., Zhao, J., Ruan, J., Luo, X., Lou, X., Bai, Y., Wang, J., Hollingsworth, L. R., Magupalli, V. G., Zhao, L., Luo, H. R., Kim, J., Lieberman, J., & Wu, H. (2020). FDA-approved disulfiram inhibits pyroptosis by blocking gasdermin D pore formation. *Nature Immunology*, *21*(7), 736–745. https://doi.org/10.1038/s41590-020-0669-6

Hu, Z., Zhou, Q., Zhang, C., Fan, S., Cheng, W., Zhao, Y., Shao, F., Wang, H.-W., Sui, S.-F., & Chai, J. (2015). Structural and biochemical basis for induced self-propagation of NLRC4. *Science*, *350*(6259), 399–404. https://doi.org/10.1126/science.aac5489

Janeway, C. A. (1989). Approaching the asymptote? Evolution and revolution in immunology. *Cold Spring Harbor Symposia on Quantitative Biology*, *54 Pt 1*, 1–13. https://doi.org/10.1101/sqb.1989.054.01.003

Jennings, E., Thurston, T. L. M., & Holden, D. W. (2017). Salmonella SPI-2 Type III Secretion System Effectors: Molecular Mechanisms And Physiological Consequences. *Cell Host & Microbe*, *22*(2), 217–231. https://doi.org/10.1016/j.chom.2017.07.009

Jorgensen, I., Zhang, Y., Krantz, B. A., & Miao, E. A. (2016). Pyroptosis triggers pore-induced intracellular traps (PITs) that capture bacteria and lead to their clearance by efferocytosis. *Journal of Experimental Medicine*, *213*(10), 2113–2128. https://doi.org/10.1084/jem.20151613

Kayagaki, N., Kornfeld, O. S., Lee, B. L., Stowe, I. B., O'Rourke, K., Li, Q., Sandoval, W., Yan, D., Kang, J., Xu, M., Zhang, J., Lee, W. P., McKenzie, B. S., Ulas, G., Payandeh, J., Roose-Girma, M., Modrusan, Z., Reja, R., Sagolla, M., … Dixit, V. M. (2021). NINJ1 mediates plasma membrane rupture during lytic cell death. *Nature*, *591*(7848), 131–136. https://doi.org/10.1038/s41586-021-03218-7

Kayagaki, N., Stowe, I. B., Alegre, K., Deshpande, I., Wu, S., Lin, Z., Kornfeld, O. S., Lee, B. L., Zhang, J., Liu, J., Suto, E., Lee, W. P., Schneider, K., Lin, W., Seshasayee, D., Bhangale, T., Chalouni, C., Johnson, M. C., Joshi, P., … Dixit, V. M. (2023). Inhibiting membrane rupture with NINJ1 antibodies limits tissue injury. *Nature*. https://doi.org/10.1038/s41586-023-06191-5

Kayagaki, N., Stowe, I. B., Lee, B. L., O'Rourke, K., Anderson, K., Warming, S., Cuellar, T., Haley, B., Roose-Girma, M., Phung, Q. T., Liu, P. S., Lill, J. R., Li, H., Wu, J., Kummerfeld, S., Zhang, J., Lee, W. P., Snipas, S. J., Salvesen, G. S., … Dixit, V. M. (2015). Caspase-11 cleaves gasdermin D for non-canonical inflammasome signalling. *Nature*, *526*(7575), 666–671. https://doi.org/10.1038/nature15541

Kayagaki, N., Warming, S., Lamkanfi, M., Walle, L. V., Louie, S., Dong, J., Newton, K., Qu, Y., Liu, J., Heldens, S., Zhang, J., Lee, W. P., Roose-Girma, M., & Dixit, V. M. (2011). Non-canonical inflammasome activation targets caspase-11. *Nature*, *479*(7371), 117–121. https://doi.org/10.1038/nature10558

Kayagaki, N., Wong, M. T., Stowe, I. B., Ramani, S. R., Gonzalez, L. C., Akashi-Takamura, S., Miyake, K., Zhang, J., Lee, W. P., Muszynski, A., Forsberg, L. S., Carlson, R. W., & Dixit, V. M. (2013). Noncanonical Inflammasome Activation by Intracellular LPS Independent of TLR4. *Science*, *341*(6151), 1246–1249. https://doi.org/10.1126/science.1240248

Kihlström, E., & Latkovic, S. (1978). Ultrastructural studies on the interaction between Salmonella typhimurium 395 M and HeLa cells. *Infection and Immunity*, *22*(3), 804–809. https://doi.org/10.1128/iai.22.3.804-809.1978

Kim, B.-H., Shenoy, A. R., Kumar, P., Das, R., Tiwari, S., & MacMicking, J. D. (2011). A family of IFN-γ-inducible 65-kD GTPases protects against bacterial infection. *Science (New York, N.Y.)*, *332*(6030), 717–721. https://doi.org/10.1126/science.1201711

Klein, J. A., Grenz, J. R., Slauch, J. M., & Knodler, L. A. (2017). Controlled Activity of the *Salmonella* Invasion-Associated Injectisome Reveals Its Intracellular Role in the Cytosolic Population. *MBio*, *8*(6). https://doi.org/10.1128/mBio.01931-17

Klein, J. A., Powers, T. R., & Knodler, L. A. (2017). Measurement of Salmonella enterica Internalization and Vacuole Lysis in Epithelial Cells. In R. Botelho (Ed.), *Phagocytosis and Phagosomes* (Vol. 1519, pp. 285–296). Springer New York. https://doi.org/10.1007/978-1-4939-6581-6_19

Knodler, L. A., Crowley, S. M., Sham, H. P., Yang, H., Wrande, M., Ma, C., Ernst, R. K., Steele-Mortimer, O., Celli, J., & Vallance, B. A. (2014). Noncanonical Inflammasome Activation of Caspase-4/Caspase-11 Mediates Epithelial Defenses against Enteric Bacterial Pathogens. *Cell Host & Microbe*, *16*(2), 249–256. https://doi.org/10.1016/j.chom.2014.07.002

Knodler, L. A., Nair, V., & Steele-Mortimer, O. (2014). Quantitative Assessment of Cytosolic Salmonella in Epithelial Cells. *PLoS ONE*, *9*(1), e84681. https://doi.org/10.1371/journal.pone.0084681

Knodler, L. A., Vallance, B. A., Celli, J., Winfree, S., Hansen, B., Montero, M., & Steele-Mortimer, O. (2010). Dissemination of invasive *Salmonella* via bacterial-induced extrusion of mucosal epithelia. *Proceedings of the National Academy of Sciences*, *107*(41), 17733–17738. https://doi.org/10.1073/pnas.1006098107

Kofoed, E. M., & Vance, R. E. (2011). Innate immune recognition of bacterial ligands by NAIPs determines inflammasome specificity. *Nature*, *477*(7366), 592–595. https://doi.org/10.1038/nature10394

Kortmann, J., Brubaker, S. W., & Monack, D. M. (2015). Cutting Edge: Inflammasome Activation in Primary Human Macrophages Is Dependent on Flagellin. *Journal of Immunology (Baltimore, Md.: 1950)*, *195*(3), 815–819. https://doi.org/10.4049/jimmunol.1403100

Kremer, J. R., Mastronarde, D. N., & McIntosh, J. R. (1996). Computer Visualization of Three-Dimensional Image Data Using IMOD. *Journal of Structural Biology*, *116*(1), 71–76. https://doi.org/10.1006/jsbi.1996.0013

Krivanek, O. L., Friedman, S. L., Gubbens, A. J., & Kraus, B. (1995). An imaging filter for biological applications. *Ultramicroscopy*, *59*(1–4), 267–282. https://doi.org/10.1016/0304-3991(95)00034-X

Kuida, K., Lippke, J. A., Ku, G., Harding, M. W., Livingston, D. J., Su, M. S.-S., & Flavell, R. A. (1995). Altered Cytokine Export and Apoptosis in Mice Deficient in Interleukin-1β Converting Enzyme. *Science*, *267*(5206), 2000–2003. https://doi.org/10.1126/science.7535475

Kutsch, M., Sistemich, L., Lesser, C. F., Goldberg, M. B., Herrmann, C., & Coers, J. (2020). Direct binding of polymeric GBP1 to LPS disrupts bacterial cell envelope functions. *The EMBO Journal*, *39*(13), e104926. https://doi.org/10.15252/embj.2020104926

Lagrange, B., Benaoudia, S., Wallet, P., Magnotti, F., Provost, A., Michal, F., Martin, A., Di Lorenzo, F., Py, B. F., Molinaro, A., & Henry, T. (2018). Human caspase-4 detects tetra-acylated LPS and cytosolic Francisella and functions differently from murine caspase-11. *Nature Communications*, *9*(1), 242. https://doi.org/10.1038/s41467-017-02682-y

Lamkanfi, M., & Dixit, V. M. (2009). Inflammasomes: Guardians of cytosolic sanctity. *Immunological Reviews*, *227*(1), 95–105. https://doi.org/10.1111/j.1600-065X.2008.00730.x

Lamkanfi, M., & Dixit, V. M. (2014). Mechanisms and Functions of Inflammasomes. *Cell*, *157*(5), 1013–1022. https://doi.org/10.1016/j.cell.2014.04.007

Lara-Tejero, M., Sutterwala, F. S., Ogura, Y., Grant, E. P., Bertin, J., Coyle, A. J., Flavell, R. A., & Galán, J. E. (2006). Role of the caspase-1 inflammasome in Salmonella typhimurium pathogenesis. *The Journal of Experimental Medicine*, *203*(6), 1407–1412. https://doi.org/10.1084/jem.20060206

Lau, N., Haeberle, A. L., O'Keeffe, B. J., Latomanski, E. A., Celli, J., Newton, H. J., & Knodler, L. A. (2019). SopF, a phosphoinositide binding effector, promotes the stability of the nascent Salmonella-containing vacuole. *PLOS Pathogens*, *15*(7), e1007959. https://doi.org/10.1371/journal.ppat.1007959

Lee, C. A., & Falkow, S. (1990). The ability of Salmonella to enter mammalian cells is affected by bacterial growth state. *Proceedings of the National Academy of Sciences*, *87*(11), 4304–4308. https://doi.org/10.1073/pnas.87.11.4304

Li, P., Allen, H., Banerjee, S., Franklin, S., Herzog, L., Johnston, C., McDowell, J., Paskind, M., Rodman, L., Salfeld, J., Towne, E., Tracey, D., Wardwell, S., Wei, F.-Y., Wong, W., Kamen, R., & Seshadri, T. (1995). Mice deficient in IL-1β-converting enzyme are defective in production of mature IL-1β and resistant to endotoxic shock. *Cell*, *80*(3), 401–411. https://doi.org/10.1016/0092-8674(95)90490-5

Liu, X., Zhang, Z., Ruan, J., Pan, Y., Magupalli, V. G., Wu, H., & Lieberman, J. (2016). Inflammasome-activated gasdermin D causes pyroptosis by forming membrane pores. *Nature*, *535*(7610), 153–158. https://doi.org/10.1038/nature18629

Livak, K. J., & Schmittgen, T. D. (2001). Analysis of relative gene expression data using real-time quantitative PCR and the 2(-Delta Delta C(T)) Method. *Methods (San Diego, Calif.)*, *25*(4), 402–408. https://doi.org/10.1006/meth.2001.1262

Majowicz, S. E., Musto, J., Scallan, E., Angulo, F. J., Kirk, M., O'Brien, S. J., Jones, T. F., Fazil, A., & Hoekstra, R. M. (2010). The Global Burden of Nontyphoidal *Salmonella* Gastroenteritis. *Clinical Infectious Diseases*, *50*(6), 882–889. https://doi.org/10.1086/650733

Malik-Kale, P., Winfree, S., & Steele-Mortimer, O. (2012). The Bimodal Lifestyle of Intracellular Salmonella in Epithelial Cells: Replication in the Cytosol Obscures Defects in Vacuolar Replication. *PLoS ONE*, *7*(6), e38732. https://doi.org/10.1371/journal.pone.0038732

Man, S. M., Ekpenyong, A., Tourlomousis, P., Achouri, S., Cammarota, E., Hughes, K., Rizzo, A., Ng, G., Wright, J. A., Cicuta, P., Guck, J. R., & Bryant, C. E. (2014). Actin polymerization as a key innate immune effector mechanism to control *Salmonella* infection. *Proceedings of the National Academy of Sciences*, *111*(49), 17588–17593. https://doi.org/10.1073/pnas.1419925111

Man, S. M., Hopkins, L. J., Nugent, E., Cox, S., Gluck, I. M., Tourlomousis, P., Wright, J. A., Cicuta, P., Monie, T. P., & Bryant, C. E. (2014). Inflammasome activation causes dual recruitment of NLRC4 and NLRP3 to the same macromolecular complex. *Proceedings of the National Academy of Sciences*, *111*(20), 7403–7408. https://doi.org/10.1073/pnas.1402911111

Mariathasan, S., Newton, K., Monack, D. M., Vucic, D., French, D. M., Lee, W. P., Roose-Girma, M., Erickson, S., & Dixit, V. M. (2004). Differential activation of the inflammasome by caspase-1 adaptors ASC and Ipaf. *Nature*, *430*(6996), 213–218. https://doi.org/10.1038/nature02664

Mariathasan, S., Weiss, D. S., Newton, K., McBride, J., O'Rourke, K., Roose-Girma, M., Lee, W. P., Weinrauch, Y., Monack, D. M., & Dixit, V. M. (2006). Cryopyrin activates the inflammasome in response to toxins and ATP. *Nature*, *440*(7081), 228–232. https://doi.org/10.1038/nature04515

Martinon, F., Burns, K., & Tschopp, J. (2002). The Inflammasome. *Molecular Cell*, *10*(2), 417–426. https://doi.org/10.1016/S1097-2765(02)00599-3

Mastronarde, D. N. (2005). Automated electron microscope tomography using robust prediction of specimen movements. *Journal of Structural Biology*, *152*(1), 36–51. https://doi.org/10.1016/j.jsb.2005.07.007

Medzhitov, R., & Janeway, C. A. (2002). Decoding the patterns of self and nonself by the innate immune system. *Science (New York, N.Y.)*, *296*(5566), 298–300. https://doi.org/10.1126/science.1068883

Meunier, E., Dick, M. S., Dreier, R. F., Schürmann, N., Broz, D. K., Warming, S., Roose-Girma, M., Bumann, D., Kayagaki, N., Takeda, K., Yamamoto, M., & Broz, P. (2014). Caspase-11 activation requires lysis of pathogen-containing vacuoles by IFN-induced GTPases. *Nature*, *509*(7500), 366–370. https://doi.org/10.1038/nature13157

Miao, E. A., Alpuche-Aranda, C. M., Dors, M., Clark, A. E., Bader, M. W., Miller, S. I., & Aderem, A. (2006). Cytoplasmic flagellin activates caspase-1 and secretion of interleukin 1β via Ipaf. *Nature Immunology*, *7*(6), 569–575. https://doi.org/10.1038/ni1344

Miao, E. A., Leaf, I. A., Treuting, P. M., Mao, D. P., Dors, M., Sarkar, A., Warren, S. E., Wewers, M. D., & Aderem, A. (2010). Caspase-1-induced pyroptosis is an innate immune effector mechanism against intracellular bacteria. *Nature Immunology*, *11*(12), 1136–1142. https://doi.org/10.1038/ni.1960

Miao, E. A., Mao, D. P., Yudkovsky, N., Bonneau, R., Lorang, C. G., Warren, S. E., Leaf, I. A., & Aderem, A. (2010). Innate immune detection of the type III secretion apparatus through the NLRC4 inflammasome. *Proceedings of the National Academy of Sciences*, *107*(7), 3076–3080. https://doi.org/10.1073/pnas.0913087107

Mills, D. M., Bajaj, V., & Lee, C. A. (1995). A 40 kb chromosomal fragment encoding Salmonella typhimurium invasion genes is absent from the corresponding region of the Escherichia coli K-12 chromosome. *Molecular Microbiology*, *15*(4), 749–759. https://doi.org/10.1111/j.1365-2958.1995.tb02382.x

Molofsky, A. B., Byrne, B. G., Whitfield, N. N., Madigan, C. A., Fuse, E. T., Tateda, K., & Swanson, M. S. (2006). Cytosolic recognition of flagellin by mouse macrophages restricts *Legionella pneumophila* infection. *The Journal of Experimental Medicine*, *203*(4), 1093–1104. https://doi.org/10.1084/jem.20051659

Muñoz-Planillo, R., Kuffa, P., Martínez-Colón, G., Smith, B. L., Rajendiran, T. M., & Núñez, G. (2013). K$^+$ efflux is the common trigger of NLRP3 inflammasome activation by bacterial toxins and particulate matter. *Immunity*, *38*(6), 1142–1153. https://doi.org/10.1016/j.immuni.2013.05.016

Naseer, N., Egan, M. S., Reyes Ruiz, V. M., Scott, W. P., Hunter, E. N., Demissie, T., Rauch, I., Brodsky, I. E., & Shin, S. (2022). Human NAIP/NLRC4 and NLRP3 inflammasomes detect Salmonella type III secretion system activities to restrict intracellular bacterial replication. *PLOS Pathogens*, *18*(1), e1009718. https://doi.org/10.1371/journal.ppat.1009718

Naseer, N., Zhang, J., Bauer, R., Constant, D. A., Nice, T. J., Brodsky, I. E., Rauch, I., & Shin, S. (2022). Salmonella enterica Serovar Typhimurium Induces NAIP/NLRC4- and NLRP3/ASC-Independent, Caspase-4-Dependent Inflammasome Activation in Human Intestinal Epithelial Cells. *Infection and Immunity*, *90*(7), e00663-21. https://doi.org/10.1128/iai.00663-21

Ochman, H., Soncini, F. C., Solomon, F., & Groisman, E. A. (1996). Identification of a pathogenicity island required for Salmonella survival in host cells. *Proceedings of the National Academy of Sciences*, *93*(15), 7800–7804. https://doi.org/10.1073/pnas.93.15.7800

Okondo, M. C., Johnson, D. C., Sridharan, R., Go, E. B., Chui, A. J., Wang, M. S., Poplawski, S. E., Wu, W., Liu, Y., Lai, J. H., Sanford, D. G., Arciprete, M. O., Golub, T. R., Bachovchin, W. W., & Bachovchin, D. A. (2017). DPP8 and DPP9 inhibition induces pro-caspase-1-dependent monocyte and macrophage pyroptosis. *Nature Chemical Biology*, *13*(1), 46–53. https://doi.org/10.1038/nchembio.2229

Perregaux, D., & Gabel, C. A. (1994). Interleukin-1 beta maturation and release in response to ATP and nigericin. Evidence that potassium depletion mediated by these agents is a necessary and common feature of their activity. *The Journal of Biological Chemistry*, *269*(21), 15195–15203.

Pilla, D. M., Hagar, J. A., Haldar, A. K., Mason, A. K., Degrandi, D., Pfeffer, K., Ernst, R. K., Yamamoto, M., Miao, E. A., & Coers, J. (2014). Guanylate binding proteins promote caspase-11-dependent pyroptosis in response to cytoplasmic LPS. *Proceedings of the National Academy of Sciences*, *111*(16), 6046–6051. https://doi.org/10.1073/pnas.1321700111

Powers, T. R., Haeberle, A. L., Predeus, A. V., Hammarlöf, D. L., Cundiff, J. A., Saldaña-Ahuactzi, Z., Hokamp, K., Hinton, J. C. D., & Knodler, L. A. (2021). Intracellular niche-specific profiling reveals transcriptional adaptations required for the cytosolic lifestyle of Salmonella enterica. *PLOS Pathogens*, *17*(8), e1009280. https://doi.org/10.1371/journal.ppat.1009280

Rathinam, V. A. K., Vanaja, S. K., Waggoner, L., Sokolovska, A., Becker, C., Stuart, L. M., Leong, J. M., & Fitzgerald, K. A. (2012). TRIF Licenses Caspase-11-Dependent NLRP3 Inflammasome Activation by Gram-Negative Bacteria. *Cell*, *150*(3), 606–619. https://doi.org/10.1016/j.cell.2012.07.007

Rauch, I., Deets, K. A., Ji, D. X., von Moltke, J., Tenthorey, J. L., Lee, A. Y., Philip, N. H., Ayres, J. S., Brodsky, I. E., Gronert, K., & Vance, R. E. (2017). NAIP-NLRC4 Inflammasomes Coordinate Intestinal Epithelial Cell Expulsion with Eicosanoid and IL-18 Release via Activation of Caspase-1 and -8. *Immunity*, *46*(4), 649–659. https://doi.org/10.1016/j.immuni.2017.03.016

Rauch, I., Tenthorey, J. L., Nichols, R. D., Al Moussawi, K., Kang, J. J., Kang, C., Kazmierczak, B. I., & Vance, R. E. (2016). NAIP proteins are required for

cytosolic detection of specific bacterial ligands in vivo. *The Journal of Experimental Medicine*, *213*(5), 657–665. https://doi.org/10.1084/jem.20151809

Raupach, B., Peuschel, S.-K., Monack, D. M., & Zychlinsky, A. (2006). Caspase-1-mediated activation of interleukin-1beta (IL-1beta) and IL-18 contributes to innate immune defenses against Salmonella enterica serovar Typhimurium infection. *Infection and Immunity*, *74*(8), 4922–4926. https://doi.org/10.1128/IAI.00417-06

Rayamajhi, M., Zak, D. E., Chavarria-Smith, J., Vance, R. E., & Miao, E. A. (2013). Cutting edge: Mouse NAIP1 detects the type III secretion system needle protein. *Journal of Immunology (Baltimore, Md.: 1950), 191*(8), 3986–3989. https://doi.org/10.4049/jimmunol.1301549

Ren, T., Zamboni, D. S., Roy, C. R., Dietrich, W. F., & Vance, R. E. (2006). Flagellin-Deficient Legionella Mutants Evade Caspase-1- and Naip5-Mediated Macrophage Immunity. *PLoS Pathogens*, *2*(3), e18. https://doi.org/10.1371/journal.ppat.0020018

Reyes Ruiz, V. M., Ramirez, J., Naseer, N., Palacio, N. M., Siddarthan, I. J., Yan, B. M., Boyer, M. A., Pensinger, D. A., Sauer, J.-D., & Shin, S. (2017). Broad detection of bacterial type III secretion system and flagellin proteins by the human NAIP/NLRC4 inflammasome. *Proceedings of the National Academy of Sciences*, *114*(50), 13242–13247. https://doi.org/10.1073/pnas.1710433114

Ross, C., Chan, A. H., Von Pein, J. B., Maddugoda, M. P., Boucher, D., & Schroder, K. (2022). Inflammatory Caspases: Toward a Unified Model for Caspase Activation by Inflammasomes. *Annual Review of Immunology*, *40*(1), 249–269. https://doi.org/10.1146/annurev-immunol-101220-030653

Roy, D., Liston, D. R., Idone, V. J., Di, A., Nelson, D. J., Pujol, C., Bliska, J. B., Chakrabarti, S., & Andrews, N. W. (2004). A Process for Controlling Intracellular Bacterial Infections Induced by Membrane Injury. *Science*, *304*(5676), 1515–1518. https://doi.org/10.1126/science.1098371

Ruan, J., Xia, S., Liu, X., Lieberman, J., & Wu, H. (2018). Cryo-EM structure of the gasdermin A3 membrane pore. *Nature*, *557*(7703), 62–67. https://doi.org/10.1038/s41586-018-0058-6

Rühl, S., & Broz, P. (2015). Caspase-11 activates a canonical NLRP3 inflammasome by promoting K $^+$ efflux. *European Journal of Immunology*, *45*(10), 2927–2936. https://doi.org/10.1002/eji.201545772

Santos, J. C., Boucher, D., Schneider, L. K., Demarco, B., Dilucca, M., Shkarina, K., Heilig, R., Chen, K. W., Lim, R. Y. H., & Broz, P. (2020). Human GBP1 binds LPS to initiate assembly of a caspase-4 activating platform on cytosolic bacteria. *Nature Communications*, *11*(1), 3276. https://doi.org/10.1038/s41467-020-16889-z

Santos, R. L., & Bäumler, A. J. (2004). Cell tropism of Salmonella enterica. *International Journal of Medical Microbiology*, *294*(4), 225–233. https://doi.org/10.1016/j.ijmm.2004.06.029

Schmid-Burgk, J. L., Gaidt, M. M., Schmidt, T., Ebert, T. S., Bartok, E., & Hornung, V. (2015). Caspase-4 mediates non-canonical activation of the NLRP3 inflammasome in human myeloid cells: Innate immunity. *European Journal of Immunology*, *45*(10), 2911–2917. https://doi.org/10.1002/eji.201545523

Sellin, M. E., Müller, A. A., Felmy, B., Dolowschiak, T., Diard, M., Tardivel, A., Maslowski, K. M., & Hardt, W.-D. (2014). Epithelium-Intrinsic NAIP/NLRC4 Inflammasome Drives Infected Enterocyte Expulsion to Restrict Salmonella Replication in the Intestinal Mucosa. *Cell Host & Microbe*, *16*(2), 237–248. https://doi.org/10.1016/j.chom.2014.07.001

Shea, J. E., Hensel, M., Gleeson, C., & Holden, D. W. (1996). Identification of a virulence locus encoding a second type III secretion system in Salmonella typhimurium. *Proceedings of the National Academy of Sciences*, *93*(6), 2593–2597. https://doi.org/10.1073/pnas.93.6.2593

Shi, J., Zhao, Y., Wang, K., Shi, X., Wang, Y., Huang, H., Zhuang, Y., Cai, T., Wang, F., & Shao, F. (2015). Cleavage of GSDMD by inflammatory caspases determines pyroptotic cell death. *Nature*, *526*(7575), 660–665. https://doi.org/10.1038/nature15514

Shi, J., Zhao, Y., Wang, Y., Gao, W., Ding, J., Li, P., Hu, L., & Shao, F. (2014). Inflammatory caspases are innate immune receptors for intracellular LPS. *Nature*, *514*(7521), 187–192. https://doi.org/10.1038/nature13683

Steele-Mortimer, O. (2008). The Salmonella-containing vacuole—Moving with the times. *Current Opinion in Microbiology*, *11*(1), 38–45. https://doi.org/10.1016/j.mib.2008.01.002

Steele-Mortimer, O., Meresse, S., Gorvel, J.-P., Toh, B.-H., & Finlay, B. B. (1999). Biogenesis of Salmonella typhimurium-containing vacuoles in epithelial cells involves interactions with the early endocytic pathway. *Cellular Microbiology*, *1*(1), 33–49. https://doi.org/10.1046/j.1462-5822.1999.00003.x

Steinberg, T. H. (1994). Cellular Transport of Drugs. *Clinical Infectious Diseases*, *19*(5), 916–921. https://doi.org/10.1093/clinids/19.5.916

Sun, Y.-H., Rolan, H. G., & Tsolis, R. M. (2007). Injection of Flagellin into the Host Cell Cytosol by Salmonella enterica Serotype Typhimurium. *Journal of Biological Chemistry*, *282*(47), 33897–33901. https://doi.org/10.1074/jbc.C700181200

Taabazuing, C. Y., Okondo, M. C., & Bachovchin, D. A. (2017). Pyroptosis and Apoptosis Pathways Engage in Bidirectional Crosstalk in Monocytes and Macrophages. *Cell Chemical Biology*, *24*(4), 507-514.e4. https://doi.org/10.1016/j.chembiol.2017.03.009

Takeuchi, A. (1967). Electron microscope studies of experimental Salmonella infection. I. Penetration into the intestinal epithelium by Salmonella typhimurium. *The American Journal of Pathology*, *50*(1), 109–136.

Takeuchi, A., & Sprinz, H. (1967). Electron-Microscope Studies of Experimental Salmonella Infection in the Preconditioned Guinea Pig: II. Response of the Intestinal Mucosa to the Invasion by Salmonella typhimurium. *The American Journal of Pathology*, *51*(1), 137–161.

Thornberry, N. A., Bull, H. G., Calaycay, J. R., Chapman, K. T., Howard, A. D., Kostura, M. J., Miller, D. K., Molineaux, S. M., Weidner, J. R., Aunins, J., Elliston, K. O., Ayala, J. M., Casano, F. J., Chin, J., Ding, G. J.-F., Egger, L. A., Gaffney, E. P., Limjuco, G., Palyha, O. C., … Tocci, M. J. (1992). A novel heterodimeric cysteine protease is required for interleukin-1βprocessing in monocytes. *Nature*, *356*(6372), 768–774. https://doi.org/10.1038/356768a0

Thurston, T. L. M., Matthews, S. A., Jennings, E., Alix, E., Shao, F., Shenoy, A. R., Birrell, M. A., & Holden, D. W. (2016). Growth inhibition of cytosolic Salmonella by caspase-1 and caspase-11 precedes host cell death. *Nature Communications*, *7*(1), 13292. https://doi.org/10.1038/ncomms13292

Tietzel, I., El-Haibi, C., & Carabeo, R. A. (2009). Human guanylate binding proteins potentiate the anti-chlamydia effects of interferon-gamma. *PloS One*, *4*(8), e6499. https://doi.org/10.1371/journal.pone.0006499

Tsuchiya, K., Hosojima, S., Hara, H., Kushiyama, H., Mahib, M. R., Kinoshita, T., & Suda, T. (2021). Gasdermin D mediates the maturation and release of IL-1α downstream of inflammasomes. *Cell Reports*, *34*(12), 108887. https://doi.org/10.1016/j.celrep.2021.108887

Valdivia, R. H., & Falkow, S. (1996). Bacterial genetics by flow cytometry: Rapid isolation of *Salmonella typhimurium* acid-inducible promoters by differential fluorescence induction . *Molecular Microbiology*, *22*(2), 367–378. https://doi.org/10.1046/j.1365-2958.1996.00120.x

Verhoef, P. A., Kertesy, S. B., Lundberg, K., Kahlenberg, J. M., & Dubyak, G. R. (2005). Inhibitory effects of chloride on the activation of caspase-1, IL-1beta secretion, and cytolysis by the P2X7 receptor. *Journal of Immunology (Baltimore, Md.: 1950)*, *175*(11), 7623–7634. https://doi.org/10.4049/jimmunol.175.11.7623

Wandel, M. P., Kim, B.-H., Park, E.-S., Boyle, K. B., Nayak, K., Lagrange, B., Herod, A., Henry, T., Zilbauer, M., Rohde, J., MacMicking, J. D., & Randow, F. (2020). Guanylate-binding proteins convert cytosolic bacteria into caspase-4 signaling platforms. *Nature Immunology*, *21*(8), 880–891. https://doi.org/10.1038/s41590-020-0697-2

Wang, J., Deobald, K., & Re, F. (2019). Gasdermin D Protects from Melioidosis through Pyroptosis and Direct Killing of Bacteria. *The Journal of Immunology*, *202*(12), 3468–3473. https://doi.org/10.4049/jimmunol.1900045

Xuong, N., Jin, L., Kleinfelder, S., Li, S., Leblanc, P., Duttweiler, F., Bouwer, J. C., Peltier, S. T., Milazzo, A., & Ellisman, M. (2007). Future Directions for Camera Systems in Electron Microscopy. In *Methods in Cell Biology* (Vol. 79, pp. 721–739). Elsevier. https://doi.org/10.1016/S0091-679X(06)79028-8

Yang, J., Zhao, Y., Shi, J., & Shao, F. (2013). Human NAIP and mouse NAIP1 recognize bacterial type III secretion needle protein for inflammasome activation. *Proceedings of the National Academy of Sciences of the United States of America*, *110*(35), 14408–14413. https://doi.org/10.1073/pnas.1306376110

Zamboni, D. S., Kobayashi, K. S., Kohlsdorf, T., Ogura, Y., Long, E. M., Vance, R. E., Kuida, K., Mariathasan, S., Dixit, V. M., Flavell, R. A., Dietrich, W. F., & Roy, C. R. (2006). The Birc1e cytosolic pattern-recognition receptor contributes to the detection and control of Legionella pneumophila infection. *Nature Immunology*, *7*(3), 318–325. https://doi.org/10.1038/ni1305

Zhang, L., Chen, S., Ruan, J., Wu, J., Tong, A. B., Yin, Q., Li, Y., David, L., Lu, A., Wang, W. L., Marks, C., Ouyang, Q., Zhang, X., Mao, Y., & Wu, H. (2015). Cryo-EM structure of the activated NAIP2-NLRC4 inflammasome reveals nucleated polymerization. *Science*, *350*(6259), 404–409. https://doi.org/10.1126/science.aac5789

Zhao, Y., Shi, J., Shi, X., Wang, Y., Wang, F., & Shao, F. (2016). Genetic functions of
the NAIP family of inflammasome receptors for bacterial ligands in mice. *The*
Journal of Experimental Medicine, *213*(5), 647–656.
https://doi.org/10.1084/jem.20160006
Zhao, Y., Yang, J., Shi, J., Gong, Y.-N., Lu, Q., Xu, H., Liu, L., & Shao, F. (2011). The
NLRC4 inflammasome receptors for bacterial flagellin and type III secretion
apparatus. *Nature*, *477*(7366), 596–600. https://doi.org/10.1038/nature10510
Zychlinsky, A., Kenny, B., Ménard, R., Prévost, M.-C., Holland, I. B., & Sansonetti, P. J.
(1994). IpaB mediates macrophage apoptosis induced by Shigella flexneri.
Molecular Microbiology, *11*(4), 619–627. https://doi.org/10.1111/j.1365-
2958.1994.tb00341.x

1373
1374

Figure Legends

Figure 1. Caspase-1 promotes the control of *Salmonella* replication within human

macrophages. WT and two independent clones of *CASP1*$^{-/-}$ THP-1 monocyte-derived

macrophages were primed with 100 ng/mL Pam3CSK4 for 16 hours. Cells were then

infected with PBS (Mock), WT *S*. Typhimurium (A, B, C), or WT *S*. Typhimurium

constitutively expressing GFP (D,E) at an MOI = 20. (A) Release of IL-1β into the

supernatant was measured by ELISA at 6 hpi. (B) Cell death (percentage cytotoxicity)

was measured by lactate dehydrogenase release assay and normalized to mock-

infected cells at 6 hpi. (C) Cells were lysed at 1 hpi and 6 hpi, and bacteria were

subsequently plated to calculate CFU. Fold-change in CFU/well was calculated. (D, E)

Cells were fixed at 6 hpi and stained for DAPI to label DNA (blue). (D) The number of

bacteria per cell at 6 hpi was scored by fluorescence microscopy. Each small dot

represents one infected cell. 150 infected cells were scored for each genotype. (E)

Representative images are shown. Scale bar represents 10 μm. Bars represent the

mean for each genotype, and error bars represent the standard deviation of triplicate

wells from one experiment (A, B, C, D). ***p < 0.001, ****p < 0.0001 by Dunnett's

multiple comparisons test (A, B, C, D). Data shown are representative of at least three

independent experiments.

1393

Figure 2. Caspase-4 contributes to the control of *Salmonella* replication within

human macrophages later during infection. WT and two independent clones of

CASP4^{-/-} THP-1 monocyte-derived macrophages were primed with 100 ng/mL

Pam3CSK4 for 16 hours. Cells were then infected with PBS (Mock), WT *S.*

Typhimurium (A, B, C), or WT *S.* Typhimurium constitutively expressing GFP (D, E) at

an MOI = 20. (A) Release of IL-1β into the supernatant was measured by ELISA at 24

hpi. (B) Cell death (percentage cytotoxicity) was measured by lactate dehydrogenase

release assay and normalized to mock-infected cells at 24 hpi. (C) Cells were lysed at 1

hpi and 24 hpi, and bacteria were subsequently plated to calculate CFU. Fold-change in

CFU/well was calculated. (D, E) Cells were fixed at 24 hpi and stained for DAPI to label

DNA (blue). (D) The number of bacteria per cell at 24 hpi was scored by fluorescence

microscopy. Each small dot represents one infected cell. 150 infected cells were scored

for each genotype. (E) Representative images are shown. Scale bar represents 10 μm.

Bars represent the mean for each genotype, and error bars represent the standard

deviation of triplicate wells from one experiment (A, B, C, D). ns – not significant, *p <

0.05 by Dunnett's multiple comparisons test (A, B, C, D). Data shown are representative

of at least three independent experiments.

1411

Figure 3. GSDMD promotes the control of *Salmonella* replication within human

macrophages. WT and *GSDMD^{-/-}* THP-1 monocyte-derived macrophages were primed

with 100 ng/mL Pam3CSK4 for 16 hours. Cells were then infected with PBS (Mock), WT

S. Typhimurium (A, B, C), or WT *S.* Typhimurium constitutively expressing GFP (D,E) at

an MOI = 20. (A) Release of IL-1β into the supernatant was measured by ELISA at 6

hpi. (B) Cell death (percentage cytotoxicity) was measured by lactate dehydrogenase

release assay and normalized to mock-infected cells at 6 hpi. (C) Cells were lysed at 1

hpi and 6 hpi, and bacteria were subsequently plated to calculate CFU. Fold-change in

CFU/well was calculated. (D, E) Cells were fixed at 6 hpi and stained for DAPI to label

DNA (blue). (D) The number of bacteria per cell at 6 hpi was scored by fluorescence

microscopy. Each small dot represents one infected cell. 150 infected cells were scored

for each genotype. (E) Representative images are shown. Scale bar represents 10 µm.

Bars represent the mean for each genotype, and error bars represent the standard

deviation of triplicate wells from one experiment (A, B, C, D). **p< 0.01, ***p < 0.001,

****p < 0.0001 by Šídák's multiple comparisons test (A) or by unpaired t-test (B, C, D).

Data shown are representative of at least three independent experiments.

1428

Figure 4. NINJ1 contributes to the control of *Salmonella* replication within human

macrophages. WT THP-1 monocyte-derived macrophages were treated with siRNA

targeting a control scrambled siRNA or siRNA targeting *NINJ1* for 72 hours prior to

infection. Cells were primed with 100 ng/mL Pam3CSK4 for 16 hours. Cells were then

infected with PBS (Mock) or WT *S*. Typhimurium at an MOI = 20. (A) Release of IL-1β

into the supernatant was measured by ELISA at 6 hpi. (B) Cell death (percentage

cytotoxicity) was measured by lactate dehydrogenase release assay and normalized to

mock-infected cells at 6 hpi. (C, D, E) Cells were lysed at 1 hpi and 6 hpi, and bacteria

were subsequently plated to calculate CFU. (C) CFU/well at 1 hpi. (D) CFU/well at 6 hpi.

(E) Fold-change in CFU/well was calculated. Bars represent the mean for each

condition. Error bars represent the standard deviation of triplicate wells from one

experiment. ns – not significant, *p< 0.05, **p< 0.01 by unpaired t-test. Data shown are

representative of at least three independent experiments.

1442

Figure 5. Inflammasome activation primarily controls cytosolic *Salmonella*

replication in human macrophages. WT, *NAIP*[-/-] (A), and *CASP1*[-/-] (A, B) THP-1

monocyte-derived macrophages were primed with 100 ng/mL Pam3CSK4 for 16 hours.

Cells were then infected with PBS (Mock) or WT *S*. Typhimurium (A) or Δ*sipB S*.

Typhimurium (B) at an MOI = 20. At 5 hpi, cells were left untreated or treated with 500

μM CHQ for 1 hour. Then, at 6 hpi, cells were lysed, and bacteria were subsequently

plated to calculate CFU. CFU/well of vacuolar (CHQ-sensitive) and cytosolic (CHQ-

resistant) bacteria were calculated from the total bacteria. (C,D) WT, *NAIP*[-/-], and

CASP1[-/-] THP-1 monocyte-derived macrophages were primed with 100 ng/mL

Pam3CSK4 for 16 hours. Cells were then infected with PBS (Mock) or WT *S.*

Typhimurium constitutively expressing mCherry and harboring the GFP cytosolic

reporter plasmid, pNF101, at an MOI = 20. Cells were fixed at 8 hpi and stained for

DAPI to label DNA (blue). (C) The number of GFP-positive, mCherry-positive bacteria

(cytosolic) per cell and the number of GFP-negative, mCherry-positive bacteria

(vacuolar) per cell were scored by fluorescence microscopy. Each small dot represents

one infected cell. 150 total infected cells were scored for each genotype. (D)

Representative images from 8 hpi are shown. Scale bar represents 10 μm. White

arrows indicate cytosolic bacteria (GFP-positive, mCherry-positive). Bars represent the

mean for each genotype (A, B). Error bars represent the standard deviation of triplicate

wells from one experiment (A, B). ns – not significant, *p< 0.05, ****p < 0.0001 by

Dunnett's multiple comparisons test (A) or by Tukey's multiple comparisons test (B).

Data shown are representative of at least three independent experiments (A, B, C).

1465

Figure 6. Inflammasome activation modulates the cytosolic exposure of

***Salmonella* in human macrophages.** (A, B) WT and *CASP1$^{-/-}$* THP-1 monocyte-

derived macrophages were primed with 100 ng/mL Pam3CSK4 for 16 hours. Cells were

then infected with WT *S*. Typhimurium at an MOI = 20. At 8 hpi, cells were fixed and

collected to be processed for transmission electron microscopy. Representative

transmission electron micrographs are shown. *Salmonella* (green) in an SCV (maroon)

with cytosolic or vacuolar clearance (yellow). Scale bar represents 1 µm [A(i), B(i)] or

400 nm [A(ii), B(ii)]. (A) WT THP-1s, (ii) is an inset from (i). White arrows indicate

vacuolar bacteria. (B) *CASP1$^{-/-}$* THP-1s, (ii) is an inset from (i). White arrows indicate

vacuolar bacteria, cyan arrows indicate partially cytosolic bacteria, and black asterisk

indicates a fully cytosolic bacterium. (C) *CASP1$^{-/-}$* THP-1 monocyte-derived

macrophages were primed with 100 ng/mL Pam3CSK4 for 16 hours. Cells were then

infected with WT *S*. Typhimurium at an MOI = 20. At 8 hpi, cells were fixed and

collected to be processed for transmission electron microscopy. Representative

tomogram slices are shown, depicting *Salmonella* (green) in an SCV with a

discontinuous vacuolar membrane (maroon).

1482

Video 1. Characterization of the cytosolic exposure of *Salmonella* in *CASP1*^{-/-}

human macrophages. *CASP1*^{-/-} THP-1 monocyte-derived macrophages were primed

with 100 ng/mL Pam3CSK4 for 16 hours. Cells were then infected with WT *S.*

Typhimurium at an MOI = 20. At 8 hpi, cells were fixed and collected to be processed

for transmission electron microscopy. Representative tomogram shown, depicting

Salmonella in an SCV with a discontinuous vacuolar membrane.

Fig. 1—Fig. Sup. 1

Fig. 1

Fig. 1—Fig. Sup. 2

Fig. 2—Fig. Sup. 1

Fig. 2—Fig. Sup. 2

Fig. 2

Fig. 2—Fig. Sup. 3

Fig. 3—Fig. Sup. 1

Fig. 3

Fig. 3—Fig. Sup. 2

Fig. 4—Fig. Sup. 1

Fig. 4

Fig. 5

A.

B.

A. WT

B. *CASP1*⁻/⁻

C.

Salmonella

Cytosolic/vacuolar clearance

SCV/SCV-like membranes

Fig. 6

A.

Cytosol-exposed bacteria

B.

WT

CASP1-/-

C.

Fig. 6—Fig. Sup. 1