Genomic insights into hybrid zone formation: the role of climate, landscape, and demography in the emergence of a novel hybrid lineage between *Populus trichocarpa* and *P. balsamifera*

Authors: Constance E. Bolte, Tommy Phannareth, Michelle Zavala-Paez, Brianna N. Sutara, Muhammed F. Can, Matthew C. Fitzpatrick, Jason A. Holliday, Stephen R. Keller, Jill A. Hamilton

Corresponding author:
Jill A. Hamilton, jvh6349@psu.edu

Abstract

Population demographic changes, alongside landscape, geographic, and climate heterogeneity, can influence the timing, stability, and extent of introgression where species hybridize. Thus, quantifying interactions across diverged lineages, and the relative contributions of interspecific genetic exchange and selection to divergence at the genome-wide level is needed to better understand the drivers of hybrid zone formation and maintenance. We used seven latitudinally arrayed transects to quantify the contributions of climate, geography, and landscape features to broad patterns of genetic structure across the hybrid zone of *Populus trichocarpa* and *P. balsamifera* and evaluated the demographic context of hybridization over time. We found genetic structure differed among the seven transects. While ancestry was structured by climate, landscape features influenced gene flow dynamics. Demographic models indicated a secondary contact event may have influenced contemporary hybrid zone formation with the origin of a putative hybrid lineage that inhabits regions with higher aridity than either of the ancestral groups. Phylogenetic relationships based on chloroplast genomes support the origin of this hybrid lineage inferred from demographic models based on nuclear data. Our results point towards the importance of climate and landscape patterns in structuring the contact zones between *P. trichocarpa* and *P. balsamifera* and emphasize the value whole genome sequencing can have to advancing our understanding of how neutral processes influence divergence across space and time.

Keywords: Natural hybridization, speciation, forest trees, gene flow*,* climate, *Populus trichocarpa, Populus balsamifera*

Introduction

Understanding the processes influencing the formation and maintenance of species is a central goal of evolutionary biology and is crucial to the management and conservation of biodiversity in a rapidly changing environment (Frankham, 2010; Hoffmann et al., 2015; Payseur & Rieseberg, 2016). As such, hybridization and introgression have long been of interest (Dobzhansky, 1936; Heiser, 1949; Jeffrey, 1916; Stebbins, 1959), and recent advances in population-scale genome sequencing have revolutionized our understanding of the frequency and pervasiveness of hybridization in nature (Hamilton & Miller, 2016; Janes & Hamilton, 2017; VanWallendael et al., 2022). Multiple taxa are known to form natural hybrid zones; however, they are particularly prevalent among forest tree species (Abbott, 2017; Suarez-Gonzalez et al., 2018; Swenson & Howard, 2004). Climatic oscillations during glacial and interglacial periods have influenced the distributions of many temperate and boreal trees, facilitating opportunities for interspecific gene flow during times of secondary contact (Hamilton et al., 2015; Hewitt, 2004; Jump & Peñuelas, 2005; Soltis et al., 2006; Swenson & Howard, 2004). However, despite widespread observations of hybridization across forest tree species, gaps remain in our understanding of how demographic, neutral and nonneutral evolutionary processes influence the formation and maintenance of natural hybrid zones in forested landscapes across space and time. Teasing apart these processes requires an assessment of the historical, landscape, and climatic factors that underlay hybrid zone formation. Ultimately, understanding how population demographic changes, alongside landscape heterogeneity and climate adaptation contribute to the timing, stability, and extent of hybridization will be critical to managing species in a rapidly changing climate.

Population demographics, including expansion and contraction of a species' range can have substantial influence on standing genetic variation, impacting the evolutionary trajectory of populations across space and time. Hybrid zones often form when previously isolated lineages come into contact and the impacts of these processes, including genetic drift and gene flow, likely play an important role shaping the genetic structure of a contact zone (Abbott, 2017; N. H. Barton & Hewitt, 1989). Shifts between isolation and contact can coincide with changes in effective population size (N_e) influencing the direction and extent of genetic exchange, while fine-scale variance at the population or genome-level can further influence elimination or fixation of alleles. In forest trees, genetic exchange often appears to be largely asymmetrical favoring the genomic background of one species due solely or in part to differential dispersal capacity, wind-patterns, and unidirectional reproductive incompatibilities (El Mujtar et al., 2017; Hamilton et al., 2013a; Lepais et al., 2009; Lexer et al., 2005, 2006). Given that an influx of genetic variants via interspecific gene flow not only buffers N_e, but can generate novel genetic recombinants upon which natural selection can act, characterizing genetic structure and the impact genetic exchange may have across space and time is needed to predict evolution.

Demographic inference can be particularly useful for characterizing processes underlying patterns of genetic exchange that can influence species evolutionary relationships, distributional shifts, or adaptive potential (Bacilieri et al., 1996; Petit et al., 2004). Several forest tree hybrid zones have characterized patterns of introgression, including oaks (Cannon & Petit, 2020; Eaton et al., 2015; McVay et al., 2017), poplars (Chhatre et al., 2018; Christe et al., 2017; Lexer et al., 2005; Suarez-Gonzalez et al., 2016; Suarez-Gonzalez et al., 2018), and spruce (Hamilton et al., 2013a, 2013b). However, few studies have leveraged whole genome sequencing to tease apart the temporal and spatial dynamics underlying the evolutionary history of contact zones using both biparentally and uniparentally inherited genomes. Viewing distinct evolutionary trajectories associated with biparentally and uniparentally inherited genomes through the lens of demographic change and landscape heterogeneity enables assessment of the role these evolutionary processes may play to the formation and maintenance of natural hybrid zones.

97 Here, we use *Populus*, an ecologically and economically important model system in forest
98 trees that naturally hybridize with congeners where geographical ranges overlap. *Populus*
99 *trichocarpa,* the first tree to have its genome fully sequenced (Tuskan et al., 2006), has
100 played an extensive role in our understanding of tree genome biology, comparative
101 genomics, and adaptive introgression (e.g., Jansson & Douglas, 2007; Shang et al., 2020;
102 Suarez-Gonzalez et al., 2016). Natural hybrid zones exist between *P. trichocarpa* and *P.*
103 *balsamifera*, largely associated with geographically and climatically steep transitions from
104 maritime to continental climates west and east of the Rocky Mountains. However,
105 divergence has likely involved a combination of intrinsic and extrinsic factors, including a
106 dynamic history of population size change throughout glacial and interglacial periods,
107 landscape-related barriers to gene flow, and climatically-related responses to selection
108 (Geraldes et al., 2014; Keller et al., 2010; Levsen et al., 2012; Slavov et al., 2012). We
109 compare whole-genome sequence data for biparentally (nuclear) and uniparentally
110 (chloroplast) inherited genomes to quantify divergence and the evolutionary relationship
111 between species across space and time. This will provide an understanding of the role
112 neutral and non-neutral processes play in the formation of these long-lived hybrid zones
113 (Petit et al. 2005).

115 Using a sampling design of latitudinally arrayed transects, we sequenced whole genomes
116 for 576 trees to capture parental-types and hybrids across repeated zones of contact
117 between *P. trichocarpa* and *P. balsamifera* distributed across their entire range of overlap.
118 With these data, we observed genetic structure within and across each contact zone and
119 asked (1) How has climate, geography, and landscape-level barriers influenced genetic
120 structure, and (2) What is the history and extent of interspecific gene flow? These data
121 provide new insights into how different evolutionary processes contribute to the formation
122 and persistence of hybrid zones over varying spatial and temporal scales.

Materials and Methods

Sampling of plant material
130

In January 2020, vegetative branch cuttings were collected from 576 poplar trees spanning repeated natural contact zones between *Populus trichocarpa* and *Populus balsamifera* (Figure 1a). These represent individual collections from seven latitudinally distributed transects (hereafter referred to as Alaska_1, Alaska_2, Cassiar, Chilcotin, Jasper, Crowsnest, and Wyoming) spanning much of the species' overlapping distributions, from 40°N to 65°N and -100°W to -150°W longitude. On average the distance between samples was approximately 43 kilometers (range: <1 - 962 km). Vegetative branch cuttings were transported back to a greenhouse at Virginia Tech (Blacksburg, VA, USA) for propagation. Details associated with propagation and greenhouse conditions are provided in Appendix 1 of Supplementary.S1. Young leaf tissue was sampled from each of the 576 propagated branch cuttings, placed immediately on dry ice, and then transferred to a -80°C freezer for long-term preservation.

Genomic Library Preparation and Bioinformatics
145

For each of the 576 samples, DNA was extracted from approximately 100mg of leaf tissue using the Qiagen plant DNeasy kit, with one modification (details available in Supplementary.S1; Appendix 2). Genomic DNA libraries were constructed at the Duke University Center for Genomic and Computational Biology, using an Illumina DNA Prep kit (Illumina Inc., San Diego, USA). Genomic libraries were sequenced on an S4 flow cell in 2x150bp format on an IlluminaNovaSeq 6000 instrument with 64 samples per lane. De-indexing, QC, trimming adapter sequences, and sequence preprocessing were completed by the sequencing facility.

Subsequent bioinformatic analysis was performed on Virginia Tech's Advanced Research Computing System (ARC) using Burrows-Wheeler Aligner (BWA) to map reads to the *P. trichocarpa* reference genome (v4.0), and the resulting SAM files were

converted to BAM format with SAMtools (Li & Durbin, 2010). The Genome Analysis

Toolkit (v3.7) Haplotype Caller algorithm was then used to generate individual gVCF

files, which were merged in a single VCF file with the GATKGenotype GVCFs function.

This raw VCF had ~82 million variant calls and was quality-filtered for variants that had

poor map quality (MQ < 40.00), elevated strand bias (FS > 40.000, SOR > 3.0),

differential map quality between reference and alternative alleles (MQRankSum < -

12.500), positional bias between reference and alternate alleles (ReadPosRankSum < -

8.000), or low coverage depth (QD < 2.0).

After quality filtering, ~34 million single nucleotide polymorphisms (SNPs) and ~4.8

million indels remained in the variant call set. Filtering removed all of the ~28 million

multiallelic sites (defined as having more than two allelic states), of which ~5 million were

SNP sites (i.e., did not include indels). For 575 of the 576 samples, missing data was on

average 0.7 % (SD: 0.1 - 1.3 %). One sample, GPR-14_S50_L001, had 99.9% missing

data and was therefore not retained for further analysis.

Analysis of population structure

To account for potential sampling of non-focal poplar species and remove those

samples, we merged the genetic data for our 575 samples with genetic data available

for 48 samples of *P. angustifolia*, which overlaps in portions of the southern distribution

of our sampling region (Chhatre et al., 2018). Methods in Supplementary.S1 (Appendix

3) detail the analyses used. While we found no need to remove samples due to

admixture with *P. angustifolia*, we removed 29 of the 575 samples with unknown

ancestry (Supplementary.S1; Figure S1 and S2), leaving 546 samples for subsequent

genetic analyses.

To assess the genetic structure of the contact zones between *P. balsamifera* and *P.*

trichocarpa, we first thinned variant sites in high linkage disequilibrium using *Plink* (v1.9)

based on 10,000 bp windows sliding in increments of 1000 bp along each chromosome,

flagging site pairs with an r^2 > 0.1. One site for each pair exceeding the r^2 threshold (i.e.,

'prune.out' file), along with singletons (--mac 2) and those with missing data (--max-

missing 1.0), were removed using *vcftools*, version 0.1.16. The resulting 299,335 low LD

variant sites were used to visualize the scale and spatial extent of hybridization.

Genomic structure was assessed using a principal component analysis (PCA) with the

'prcomp' function of the *stats* package in R, version 4.2.1. We then performed an

ADMIXTURE analysis (Alexander et al., 2009) using cluster assignments ranging from

K=2 to K=6. The cluster assignment (K) with the lowest cross validation error (k-folds =

10) was determined to best fit the data (Figure 1c).

Assessing evolutionary relationships from nuclear data

To evaluate the evolutionary relationships between genetic clusters, we used the three

genetic clusters identified from ADMIXTURE (K = 3; see Results) to determine the

topology with the greatest support. We identified parental-type samples for each genetic

cluster, hereafter referred to as *P. balsamifera*, *coastal P. trichocarpa*, and *interior P.*

trichocarpa, based on K=3 using an ancestry coefficient threshold of Q > 0.90. Of the

546 samples included in the analysis, 185 parental-types were identified (112 *P.*

balsamifera, 51 *coastal P. trichocarpa*, and 22 *interior P. trichocarpa*). Using parental-

types, we evaluated three different topologies based on the proposed divergence history

between *interior P. trichocarpa* and the other two parental-type genetic lineages (Table

1). The first topology evaluated expectations based on a recent divergence from *coastal*

*P. trichocarp*a, while the second tested for a recent divergence from *P. balsamifera*, and

the third evaluated whether *interior P. trichocarpa* was a hybrid lineage formed from an

admixture event between *coastal P. trichocarpa* and *P. balsamifera*.

The evolutionary relationship of the three genetic clusters was evaluated using the

genotypes from the 185 parental-types (112 *P. balsamifera*, 51 *coastal P. trichocarpa*,

and 22 *interior P. trichocarpa*) and Diffusion Approximation for Demographic Inference

($\partial a \partial i$ v.2.1.1;(Gutenkunst et al., 2009). We used the unfolded joint site frequency

spectrum (jSFS) of derived allele counts following polarization of genotypes following the

approach of (Luqman et al., 2023). Details pertaining to data sources and program

implementation are provided in Supplementary_S1, Appendix 4. We assigned allelic

states as either ancestral or derived based on the consensus ancestral state: 6495 sites

(2.260%) had the reference allele assigned to the derived state and 275,751 sites
(95.959%) had the reference allele assigned to the ancestral state. The remaining 1.781%
of sites had undefined ancestral alleles. To increase computational efficiency, we used
the derived allele frequencies for a random sample of 10 diploid individuals per parental-
type cluster to build an unfolded 3-dimensional joint site frequency spectrum (jSFS) using
the function 'GenerateFs' in the dadi-cli program (X. Huang et al., 2023). We performed
ten replicate runs of each model in $\partial a \partial i$ with a 40 x 50 x 60 grid space and the nonlinear
Broyden-Fletcher-Goldfarb-Shannon (BFGS) optimization routine. Akaike information
Criterion (AIC; Akaike 1974) was employed for model selection, with the best replicate
run (highest log composite likelihood) for each model used to calculate ΔAIC ($AIC_{model\ i}$ −
$AIC_{best\ model}$) scores (Burnham & Anderson, 2002).

To estimate the timing of divergence and quantify the extent and directionality of gene
flow between genetic clusters, we tested four hypotheses, or models, of connectivity and
divergence based on the best supported topology (Table 1). The first model assumed a
scenario of recurrent gene flow, while the other three models assumed scenarios of
isolation that varied based on the timing and frequency of secondary contact
(Supplementary.S1, Figure S4). We performed ten replicate runs of each model in $\partial a \partial i$
with a 40 x 50 x 60 grid space and the nonlinear Broyden-Fletcher-Goldfarb-Shannon
(BFGS) optimization routine. Akaike information Criterion (AIC; Akaike 1974) was
employed for model selection, and Fisher Information Matrix (FIM)-based uncertainty
analysis was then conducted for the best supported model to obtain upper and lower 95%
confidence intervals (CIs) for the inferred parameter estimates. To unscale parameter
estimates and their 95% CIs we used a mutation rate of 1.33 x 10^{-10} substitutions/site/year
rate for *Populus trichocarpa* (Hofmeister et al., 2020), a generation time of 15 years, and
a calculated effective genome length (Supplementary.S3). To calculate effective genome
length, we multiplied the proportion of variant sites used in the modeling (count of filtered
sites in each pairwise inference with no missing data and singletons removed / count of
variant sites prior to filtering) by the number of base pairs from the *P. trichocarpa*
reference genome considered in variant calling (389,204,664 bp).

Assessing evolutionary relationships from chloroplast data

To gain insight into the evolutionary relationships among the three genetic clusters identified from nuclear data, we used the 185 parental-types to examine differentiation across uni-parentally inherited chloroplast genomes. Chloroplast (cp) genomes were assembled using NOVOPlasty (Dierckxsens et al., 2017). The published Nisqaully-1 *P. trichocarpa* chloroplast genome (NC_009143.1) was used as the reference sequence for genome assembly. The ribulose-bisphosphate carboxylase (*rbc*L) gene was manually extracted and used as a seed, serving as an initial reference point for the chloroplast genome assembly. Assembled chloroplast genomes were annotated using Geneious v7.1.4 (Dierckxsens et al. 2016). Following annotation, inverted repeats and coding regions were manually checked for each annotated individual. In total, 184 out of the 185 parental-types were successfully assembled, with one individual (genotype 831) excluded due to incomplete assembly.

A maximum likelihood (ML) phylogenetic tree was built to assess the relationships among the chloroplast genomes for each parental type. The chloroplast genomes from two *P. angustifolia* samples (NC_037413 and MW376761) were downloaded from NCBI and selected as an outgroup. Chloroplast genomes for all 184 samples and outgroups were aligned using MAFFT V.7.453 (Katoh & Standley, 2013) to perform a ML phylogeny in IQ-tree 1.6.11 (Trifinopoulos et al., 2016). The best fit evolutionary model for the sequence alignment was calculated by ModelFinder (Kalyaanamoorthy et al., 2017), with the selected model being the transversion model with empirical base frequencies and FreeRate model (TVM+F+R2). A ML tree was generated following 1,000 bootstrap replications and a consensus phylogenetic tree was created based on all parental-type samples using the FigTree software v1.4.13.

Associating genetic structure with climate and geography

To quantify the multivariate relationships between genetic structure, climate, and geographic variation across the hybrid zone, a redundancy analysis (RDA) was conducted within the *vegan* package, version 2.6-2 (Oksanen et al., 2020) in R version 4.2.1. Climate data at 30 second resolution averaged across the years 1961-1990 for

twenty-five variables associated with geographic origin were extracted using *ClimateNA* (version 5.2; (Wang et al., 2016) for each sampled tree (Supplementary.S2) and then tested for collinearity using variance inflation factor (VIF) analysis implemented in the *usdm* R package (version 1.1-18; (Naimi et al., 2014). Six climatic variables with a VIF less than 10 were retained for subsequent analysis, including continentality (TD), mean annual temperature (MAT), mean annual precipitation (MAP), climate moisture deficit (CMD), relative humidity (RH), and precipitation as snow (PAS). Geographic variables included for subsequent analysis were latitude, longitude, and elevation.

Genotype data (299,335 nuclear SNPs and 2,828 chloroplast SNPs) for all 546 samples was converted to minor allele counts per sample using vcftools (version 0.1.16). A permutation-based analysis of variance (ANOVA) procedure with 999 permutations (Legendre & Legendre, 2012) was used to test the overall statistical significance of the RDA model (α = 0.05) as well as the significance of each RDA axis separately. Variance partitioning was then performed using the 'varpart' function of the *vegan* package to quantify the influence of predictor variables and their confounding effects to observed genetic variation. This analysis was repeated without hybrid samples (0.10 < Q < 0.90) to compare the multivariate relationships among the 185 parental-type samples.

Visualizing barriers to gene flow

To evaluate genetic connectivity across the contact zones between *P. trichocarpa* and *P. balsamifera,* we used EEMS (Estimated Effective Migration Surfaces; (Petkova et al., 2016) and observed correspondence between landscape features, ancestry, and patterns of genetic connectivity. A genetic dissimilarity matrix was generated using the 'bed2diffs script' included within the EEMS program informed by the 299,335 genome-wide nuclear SNPs across 546 individuals. Two additional geographic files were created using geographic coordinates; including the location data of each individual (.coord extension) and a large general habitat polygon (.outer extension) which was manually obtained using Google Maps API v3 Tool. The geographical coordinates for each sample alongside the parameterized grid space (500 demes with each grid cell ~21,000 km^2) assigned individuals to one of 49 demes, defined as pseudo populations of close geographical

proximity to a vertex of the defined grid space (Supplementary.S1, Figure S3). We then

estimated the effective migration rate using a stepping-stone model. Model parameters

included a burn-in of 25,000,000 with an MCMC length of 50,000,000 and thinning to

sample 1 in every 100,000 iterations. The results of three replicate runs were averaged

to visualize effective diversity (q) and effective migration rate (m) surfaces (in $\log_{10}$ scale)

across the contact zone using the R package *rEEMSplots* v 0.0.1 (Petkova et al., 2016).

Results

Genomic structure of hybrid zone identifies three distinct genetic clusters

Using all 546 individuals, the nuclear genomic PCA ascribed genetic structure along PC1

(1.764% variance explained) to species-specific differences between *P. trichocarpa* (far

left of PC1) and *P. balsamifera* (far right of PC1) and PC2 (0.925% variance explained)

to genetic structure within *P. trichocarpa* (Figure 1b). Admixture analysis best supported

the presence of three genetic clusters (*K* = 3), with admixed individuals present across all

transects (Figure 1c). Two of the three genetic groups clustered geographically within the

range of *P. trichocarpa* (Little 1971) – this included a *coastal P. trichocarpa* cluster which

had a northern, coastal distribution, and an *interior P. trichocarpa* cluster*,* which had a

southern, interior distribution (Figure 1d). The third genetic cluster was *P. balsamifera* in

the K = 3 admixture analysis. Pairwise estimates of genome-wide genetic differentiation

(F_{ST}) were similar across the three genetic clusters, ranging from 0.053 for *coastal -*

interior P. trichocarpa, 0.054 for *interior P. trichocarpa - P. balsamifera*, and 0.056 for

coastal P. trichocarpa - P. balsamifera comparisons.

Of the 546 samples included in the analysis, 185 parental-types associated with *K* = 3

genetic clusters were identified based on Q > 0.90 (112 individuals identified as *P.*

balsamifera, 51 individuals identified as *coastal P. trichocarpa*, and 22 individuals

identified as *interior P. trichocarpa). Coastal P. trichocarpa* and *P. balsamifera* genetic

ancestry was observed in Alaska and Cassiar transects, although the Alaska transect did

not appear to have any *coastal P. trichocarpa* parental-types. All three genetic clusters
were observed in Chilcotin, Jasper, Crowsnest, and Wyoming transects (Figure 1c).
Shared ancestry associated with *P. balsamifera* along the Crowsnest transect was
restricted to the eastern edge of the *P. trichocarpa* distribution. The Wyoming transect
was dominated by *interior P. trichocarpa* parental-type ancestry (Q > 0.90 at *K* = 3;
Supplementary.S2), whereas individuals sampled near the western edge of this transect
exhibited mixed ancestry between the coastal and interior genetic clusters of *P.*
trichocarpa. Consistent with the Crowsnest transect, *P. balsamifera* ancestry was limited
to the far eastern distribution of *P. trichocarpa,* with *P. balsamifera* parental-types
restricted to fragmented populations geographically distant from the main distribution of
either species.

Nuclear and chloroplast genomes identify the persistence of three genetic lineages
Given the similar estimates of genome-wide nuclear differentiation (F_{ST}), we performed a
topology test to better understand the relationships among the three genetic clusters. The
best fit topology as determined by AIC (Table 1a) supported *interior P. trichocarpa* as an
ancient lineage of admixed origin. The second-best fit topology placed *interior P.*
trichocarpa as a split from *coastal P. trichocarpa*, but delta AIC was 4,943 units higher.

To complement assessment of nuclear topologies, evolutionary relationships were also
assessed using ML analysis of chloroplast genomes. The alignment of the 184 individuals
and outgroup genomes produced a single matrix with 164,183 nucleotide sites, including
1,020 polymorphic sites. Three distinct clades were detected using the ML tree, including
coastal P. trichocarpa, *interior P. trichocarpa*, and *P. balsamifera* respectively. The
coastal P. trichocarpa clade appeared ancestral to both *interior P. trichocarpa* and *P.*
balsamifera, with all three exhibiting robust bootstrap support. Lineage assignments were
congruent between nuclear and chloroplast genomes, except for individual 757, which
clustered with *interior P. trichocarpa* in the nuclear admixture analysis but in the
chloroplast ML tree belonged to the *P. balsamifera* clade.

Climate strongly associates with genetic structure

Climate and geography explained 3.79% (r^2) of the genetic variance across the 546

sampled trees and 299,335 SNP dataset. Most of the explainable genetic variance (PVE)

was accounted for by the first RDA axis (38.26%, Figure 2a) with continentality (TD) and

mean annual temperature (MAT) having the highest predictor loadings (eigenvectors).

Individuals identified as *P. balsamifera* were associated with higher TD and lower MAT

on average. The second RDA axis captured 19.23% (PVE) of the explainable genetic

variance, with the highest predictor loadings on the second RDA axis being relative

humidity (RH), elevation, and longitude. These variables were associated with

differentiation within *P. trichocarpa* across the seven transects. Associations between

climate, geography and genetic variation were not observed along RDA2 for *P.*

balsamifera (Figure 2). Combining loadings for RDA1 and RDA2, we found latitude, TD,

and climate moisture deficit (CMD) had the strongest associations with genetic variation

within and among sampled transects.

Because geographic and climatic predictor variables are often correlated, we performed

variance partitioning to tease apart the respective contributions of climate, geography,

and their interactions to genomic variance. We observed a large, confounded effect

(58.69%) between the climatic and geographic variables chosen for analysis. Removing

confounded effects through conditioning revealed climate was a stronger predictor of

genetic variance. Climate, conditioned on geography, accounted for 27.28% of

explainable genetic variance, whereas geography, conditioned on climate, accounted for

just 14.03% of explainable genetic variance (Supplementary.S1; Table S2).

Using the same suite of analyses to investigate associations between environment and

ancestry for the 185 parental-types revealed higher contributions of climate and

geography to explainable genetic structure. From side-by-side comparisons of nuclear

and chloroplast genetic structure in the PCA, we observed similar clustering (Figure 2).

Considering the differentiation along PC1 axis, *coastal P. trichocarpa* and *interior P.*

trichocarpa appear less diverged based on nuclear data when compared to chloroplast.

Similar clustering and environmental associations were also observed in the RDA (Figure

2) and aligned with the associations presented for the 546 samples (Figure 2a). However,

with hybrids removed from the analysis, the explanatory variance increased (adj. r^2 =
7.8%; PVE = 51.65%). The explanatory variance based on chloroplast structure was the
highest observed (adj. r^2 = 30.1%; PVE = 66.26%). Variance partitioning was also
conducted with the hybrids excluded, and with the confounding effects of geography
removed, climate explained approximately 40-42% of the genetic variance in both the
chloroplast and nuclear data. Conversely, after conditioning on climate, geography
explained only 9.46% of the nuclear genetic variance and 2.87% of the chloroplast genetic
variance (Supplementary.S1, Table S3).

Gene flow dynamics shaped by physical barriers
Using departures from isolation-by-distance (IBD) to indicate barriers to gene flow, EEMS
analysis identified the continental divide of the Rocky Mountains as a large barrier
between *P. trichocarpa* and *P. balsamifera*. Additional barriers to gene flow were
identified between transects associated with elevational gradients (Figure 3). For
example, *interior P. trichocarpa* within the most southern transect was observed in a
region of high local connectivity but was isolated from neighboring individuals identified
as *coastal P. trichocarpa* and *P. balsamifera* in other analyses. The EEMS results
indicated both Alaska transects were more isolated from the other northern transect,
Cassiar, than expected based on IBD. In contrast, individuals spanning *coastal P.*
trichocarpa parental-type and hybrids across the Chilcotin, Jasper, and Crowsnest
transects were more similar than expected under IBD, suggesting substantial intraspecific
migration across latitudes within the southern range of this hybrid zone. This pattern was
not observed for *P. balsamifera,* as only individuals located east of the Rocky Mountains
exhibited evidence of greater effective migration than expected.

Demographic inference reveals periods of interspecific vicariance, gene flow, and
the emergence of a hybrid lineage
Of the four demographic models of connectivity and divergence evaluated (Table 1b), the
model with the greatest support inferred an initial divergence between *coastal P.*
trichocarpa - P. balsamifera approximately 2.12 million years ago (95% CI: 2.11 – 2.13
million years ago). Following initial divergence, limited gene flow (mA ~ 1.90e-04)

continued until approximately 860,000 years ago when gene flow ceased, during which
time the two lineages were isolated (Figure 4). Following approximately 96,000 years of
isolation, the two lineages experienced secondary contact and 764,000 years ago formed
the *interior P. trichocarpa* lineage via admixture. During the initial formation of *interior P.*
trichocarpa the proportions of each progenitor lineage were inferred to be approximately
34.8% *coastal P. trichocarpa* (parameter *f*; Table 2) and 65.2% *P. balsamifera* (1 - *f*).
Following secondary contact, asymmetrical gene flow was inferred until present, with the
highest rates of gene flow observed between *coastal P. trichocarpa* and *interior P.*
trichocarpa (*m13* ~ 5.08e-05 and *m31* ~ 4.58e-05; Table 2), with a slightly greater rate of
genetic exchange from *interior* into *coastal* (*m13*). Gene flow appeared to be strongly
asymmetrical, with more movement from *P. balsamifera* into *coastal* and *interior P.*
trichocarpa (*m12* ~ 3.64e-05, *m32* ~ 3.13e-05, respectively) and less *coastal* or *interior*
P. trichocarpa movement into *P. balsamifera* (*m21* ~ 1.30e-05 and *m23* ~ 9.08e-06; Table
3). The highest effective population size inferred was for *P. balsamifera* (N_2 ~ 100,000)
followed by *interior P. trichocarpa* (N_3 ~ 53,800) and *coastal P. trichocarpa* (N_1 ~ 36,500).
The residuals between the model and data fit were small and evenly distributed,
suggesting good model fit (Supplementary.S1, Figure S5). In addition, confidence
intervals for each parameter were tightly bound to the estimations inferred (Table 2).

Discussion

This study provides valuable insights into how complex spatial and temporal dynamics
influence hybridization, highlighting the role neutral processes can play in shaping the
genetic structure of hybrid zones. A dynamic history between glacial and interglacial
cycles suggests that secondary contact over time has had a substantial influence on
hybrid zone structure. We identified three distinct genetic clusters geographically
distributed across our seven repeated zones of species overlap. Differentiation within and
across nuclear and chloroplast genomes supported the presence of three genetic

lineages, with tests of topology suggesting one genetic lineage reflects an ancient
admixture event between *coastal P. trichocarpa* and *P. balsamifera* leading to the origin
of a new hybrid lineage, *interior P. trichocarpa*. While physical geographic barriers have
played a significant role influencing historical and contemporary asymmetrical genetic
exchange, our results highlight the important role climate gradients alongside geography
have played influencing hybrid zone structure. Together, this suggests that genetic
patterns observed could be sensitive to climatic shifts, which has implications for
understanding the future of these contact zones in a changing climate.

Demographic shifts associate with shifts in historical climate
According to our best fit 3D demographic model, initial divergence between *P.*
balsamifera and coastal P. trichocarpa (~ 2.12 million years ago) aligns with a shift from
the Pliocene to the Pleistocene Climate Era (Tan et al., 2018). This initial divergence was
inferred to occur amid recurrent gene flow for approximately 1.25 million years, followed
by a period of isolation lasting ~ 96,000 years. Interestingly, this period of isolation not
only aligns with the Mid-Pleistocene Transition, when glacial cycles became longer (from
~40,000 to ~100,000 years long) and shifts to interglacial climates become abrupt and
extreme, but it also coincides in timing with a glacial maximum (~800,000 years ago;
(Chalk et al., 2017; Ellis & Palmer, 2016; Willeit et al., 2019). Secondary contact between
P. balsamifera and *coastal P. trichocarpa*, along with the origin of *interior P. trichocarpa,*
was estimated to have occurred ~764,000 years ago and coincides with a transition to
interglacial climate (Ellis & Palmer, 2016). This suggests opportunities for genetic
exchange were regained following expansion from isolated refugia.

While climate has continued to oscillate since the inferred secondary contact event, which
might be expected to reduce opportunities for gene flow given the different climatic
affinities of each lineage, our demographic model suggested continued genetic exchange.
One explanation may be that species distributions have moved in tandem with climate
such that populations were proximal enough to maintain gene flow, resulting in a stable
hybrid zone. Indeed, the long-distance dispersal of seed and pollen in *Populus* and other
forest trees blurs definitions for how geographically isolated species need to be to disrupt

gene flow. An alternative explanation may be that periods of strict isolation did occur

during extreme climatic shifts but were not sustained long enough for genomic patterns

of divergence associated with allopatry to develop, or patterns were broken down

following subsequent contact. This latter possibility may characterize hybridizing species

with porous genomes whose species boundaries vary temporally with repeated cycles of

reinforcement and homogenization (e.g., Mix-Isolation-Mix hypothesis; He et al., 2019) or

due to changes in the strength and direction of selection over historically varying climatic

optima (e.g., Shang et al., 2020). Among the divergence histories inferred for forest tree

species, there is growing evidence for divergence with recurrent gene flow despite abrupt

shifts in climate (e.g., Bolte et al., 2022; He et al., 2019; Lexer et al., 2006; Menon et al.,

2018).

Gene flow asymmetries in the context of future climate change

Gene flow was nearly three times higher from *P. balsamifera* into *coastal P. trichocarpa*

than from *coastal P. trichocarpa* into *P. balsamifera.* This observation is corroborated by

previous estimates of genetic exchange in this hybrid zone by Suarez-Gonzalez et al.

(2018), who noted asymmetrical introgression from *P. balsamifera* into *P. trichocarpa*.

This asymmetry may suggest *P. balsamifera* has a demographic advantage within the

ecotonal transition (e.g., dispersal down an elevational gradient or with river flow from

inland to coast). On the other hand, lower rates of introgression between *P. balsamifera*

and *interior P. trichocarpa* may be indicative of stronger geographical isolation, a lack of

hybrids geographically positioned between these genetic clusters to facilitate

backcrossing, or even more likely, asynchronous reproductive phenology. According to

descriptions within (Haeussler et al., 1990) flowering times for interior populations of *P.*

trichocarpa can be nearly one month later than those reported for populations of *P.*

trichocarpa (coastal) and *P. balsamifera* in British Columbia, Canada.

Gene flow between *coastal P. trichocarpa* and *interior P. trichocarpa* were greatest

relative to all inferred between-lineage contemporary estimates (Figure 4b). Indeed,

interior P. trichocarpa is genetically more similar to *coastal P. trichocarpa* based on

patterns of nuclear differentiation. However, despite this, the demographic model that

received the greatest support suggested that in the formation of *interior P. trichocarpa*
(~764,000 years ago) only 35% of the hybrid genome reflected ancestry from *coastal P.*
trichocarpa. Taken together, these results may indicate that higher rates of contemporary
gene flow between *coastal* and *interior P. trichocarpa* homogenized population
differences following the initial divergence of the *interior P. trichocarpa* lineage.
Regardless, the influx of new genetic recombinants through introgression is evident
across all three parental-types and may impact adaptation to a changing climate (Aitken
et al., 2008; N. Barton & Bengtsson, 1986; Martinsen et al., 2001).

Climate and landscape heterogeneity contribute to genetic structure
Landscape-induced restrictions to gene flow have contributed to patterns of differentiation
across the hybrid zone. Estimated effective migration surfaces (EEMS) suggest gene flow
has likely been suppressed by elevational barriers through the Rocky Mountains of
northwestern North America. West of the Rocky Mountains, we observed variation in
effective migration rates within the *P. trichocarpa* distribution described by (EL Little Jr.,
1971). For example, effective migration was greater within the northern transects of
Alaska and Cassiar but not between these two transects. Similarly, there appeared to be
substantial gene flow between hybrids within the 'mid-latitudinal' transects (Chilcotin,
Jasper, and Crowsnest), but migration was restricted along the coastal regions of these
three transects where parental-types of *coastal P. trichocarpa* were identified. This
suggests gene flow between *P. trichocarpa* populations is lower than expected based on
expectations of isolation-by-distance. Thus, landscape barriers to gene flow likely
contribute to strong population structure observed within the range of *P. trichocarpa*
(Geraldes et al., 2014; Slavov et al., 2012).

Redundancy analysis (RDA) and variance partitioning revealed genetic structure and
climate are significantly associated across the hybrid zone. The Rocky Mountains present
environmental transitions from drier and colder conditions on the eastern side (favoring
P. balsamifera) to more maritime climates on the western side of the Rockies (favoring
coastal P. trichocarpa). The distributions and climate associations we observed between
P. balsamifera, coastal P. trichocarpa, and *interior P. trichocarpa* (Figure 2a) are

comparable to other hybrid zones within the region, including *Picea glauca, P. sitchensis*,

and *P. engelmannii* hybrid zones (De La Torre et al., 2015; Hamilton et al., 2013b, 2015).

In both systems, maritime to continental climate transitions were associated with genomic

ancestry. Because the frequency of hybrid genotypes is largely associated with ecotonal

transitions, this suggests these hybrid zones likely reflect a model of bounded hybrid

superiority. However, quantifying fitness in reciprocal transplant experiments is needed

to test the bounded hybrid superiority model.

The admixed lineage, *interior P. trichocarpa*, appears to be bounded to a drier and warmer

climate than either of its progenitors. The genetic structure of *interior P. trichocarpa* is

associated with high climate moisture deficit. If selection maintains genetic structure

among the lineages observed with our study, we may find trait divergence, especially

those underlying water-use, across genetic lineages. Interestingly, *Populus* hybrid

lineages of the Qinghai-Tibet Plateau (QTP) also inhabit landscapes with higher aridity

relative to their progenitors. However, survival and growth measurements within common

garden experiments suggested phenotypic plasticity may have contributed to adaptive

divergence and persistence in environments beyond the parental species' range (Shi et

al., 2024). A similar experimental design could be employed for *coastal P. trichocarpa,*

interior P. trichocarpa, and *P. balsamifera* to test if the admixed lineage (*interior P.*

trichocarpa) exhibit transgressive traits relative to parents while also evaluating the role

of plasticity to success in novel environments. In the absence of trait data, we can

leverage genomic data to differentiate between, and functionally assess, genomic regions

associated with restricted or excess introgression that may underlay adaptive divergence.

Nuclear and chloroplast genomes provide complementary insights into hybrid

zone dynamics

The estimates observed herein suggest 2.11 - 2.13 million years ago *P. trichocarpa* and

P. balsamifera diverged and are congruent with observations for this species-pair from

(Sanderson et al., 2023). The differentiation we observed in the chloroplast not only aligns

with patterns of nuclear differentiation among the three assigned parental-types but also

supports the observations and interpretations of previous chloroplast assessments of

divergence in *Populus*. (D. I. Huang et al., 2014) ascribed the deep chloroplast divergence
of *P. trichocarpa* from *P. balsamifera* to chloroplast capture of an ancient, now extinct,
lineage. While the genome-wide patterns of nuclear differentiation (e.g., PCA, F_{ST})
suggests *coastal P. trichocarpa* and *interior P. trichocarpa* are more closely related, our
phylogenetic inference from the chloroplast placed *coastal P. trichocarpa* outside the
clade of *interior P. trichocarpa* and *P. balsamifera*. Only one individual out of 185 did not
have a matching chloroplast - nuclear parental-type assignment. Considering our
demographic modeling and the fact that our focal taxa exhibit dioecy with maternal
inheritance of the chloroplast genome, the origin of *interior P. trichocarpa* likely resulted
from admixture between *P. balsamifera* seed parents and *coastal P. trichocarpa* pollen
parents. Given the inferred length of time that has passed since the admixed origin
(~764,000 years), the *interior P. trichocarpa* chloroplast genome has likely evolved
following divergence. Co-evolution and co-adaptation between the chloroplast and
nuclear genomes may have contributed to chloroplast divergence of *interior P.*
trichocarpa from the original seed parent (Greiner & Bock, 2013) and would be an
interesting future avenue for research.
607

Conclusions
Here, we provide insight into the evolutionary processes shaping divergence and
hybridization across space and time in a *Populus* system of western North America.
Demographic models support the origin of a hybrid lineage following secondary contact
of ancestral lineages and a history of asymmetrical gene flow between focal taxa. Our
results highlight the influence of climate and geography to genetic structure and
divergence, as well as the influence of landscape-level attributes to gene flow dynamics.
Overall, our study sheds light on the role demographic processes can play within a long-
lived hybrid zone across space and time and highlight the need for fine-scale assessment
of regions of the genome important to introgression and adaptation for species
management in a changing climate.
619
620
621

Acknowledgements:

We thank Joe Braasch, Lionel Di Santo, Sonia DeYoung, Sara Klopf, Jack Woods, Ben Woods, Raju Soolanayakanahally for assisting with field sample collection. We also thank Kyle Peer, Clay Sawyers, and Deborah Bird from the Virginia Tech Reynold's Homestead Forestry Research Station for their assistance with plant propagation. We thank Ryan Gutenkunst and Andrew Eckert for advice and communications related to demographic inference. We acknowledge the valuable questions from and conversations with our team members during project calls and meetings: Cigdem Kansu, Susanne Lachmuth, Baxter Worthing, and Alayna Mead.

Competing interests: All authors claim no competing interests.

Author contributions:

MCF, JH, SRK, and JAH conceived the study and collected the samples. MC extracted DNA. TP processed genomic data. CEB analyzed genomic data, conducted demographic inference, and performed genetic-environment associations. MZ and BS assembled chloroplast genomes and performed chloroplast analyses. SRK, JH and JAH advised analyses. CEB and JAH led the writing of the manuscript. SRK, JH, and MCF provided valuable feedback.

Data availability: Sequence reads for the 546 samples analyzed in this manuscript have been archived on NCBI (PRJNA996882). Chloroplast genomes are archived under BankIt2793276 (Accession numbers PP274128 - PP274690). Both nuclear and chloroplast sequences will remain under embargo until publication. Scripts for variance partitioning, RDA, and demographic inference can be found at https://github.com/boltece/Poplar_divergence.

Funding: National Science Foundation grant IOS-1856450, USDA National Institute of Food and Agriculture project and Hatch Appropriations (PEN04809 and Accession 7003639), NIFA project VA-136641, Schatz Center for Tree Molecular Genetics

Supporting Information:

Supplementary.S1 - Supplementary methods (appendices) and figures

Supplementary.S2 - Sample summaries for climate and geographic data, admixture coefficients, and PCA and RDA coordinates

Supplementary.S3 - Summarized results from demographic inference including AIC calculations, unscaling of $\partial a \partial i$ parameter estimates, and confidence interval calculations

References:

Abbott, R. J. (2017). Plant speciation across environmental gradients and the

    occurrence and nature of hybrid zones. *Journal of Systematics and Evolution*,

    55(4), 238–258. https://doi.org/10.1111/JSE.12267

Aitken, S. N., Yeaman, S., Holliday, J. A., Wang, T., & Curtis-McLane, S. (2008).

    Adaptation, migration or extirpation: Climate change outcomes for tree

    populations. *Evolutionary Applications*, *1*(1), 95–111.

    https://doi.org/10.1111/J.1752-4571.2007.00013.X

Alexander, D. H., Novembre, J., & Lange, K. (2009). Fast model-based estimation of

    ancestry in unrelated individuals. *Genome Research*, *19*(9).

    https://doi.org/10.1101/gr.094052.109

Bacilieri, R., Ducousso, A., & Kremer, A. (1996). Comparison of morphological

    characters and molecular markers for the analysis of hybridization in sessile and

    pedunculate oak. *Annales Des Sciences Forestières*, *53*(1), 79–91.

    https://doi.org/10.1051/forest:19960106

Barton, N., & Bengtsson, B. O. (1986). The barrier to genetic exchange between

hybridising populations. *Heredity*, *57*(3), 357–376.

https://doi.org/10.1038/hdy.1986.135

Barton, N. H., & Hewitt, G. M. (1989). Adaptation, speciation and hybrid zones. *Nature*,

341(6242), 497–503. https://doi.org/10.1038/341497a0

Bolte, C. E., Faske, T. M., Friedline, C. J., & Eckert, A. J. (2022). Divergence amid

recurring gene flow: Complex demographic histories for two North American pine

species (Pinus pungens and P. rigida) fit growing expectations among forest

trees. *Tree Genetics & Genomes*, *18*(5), 35. https://doi.org/10.1007/s11295-022-

01565-8

Burnham, K. P., & Anderson, D. R. (2002). *Model selection and multimodel inference: A*

practical information-theoretic approach. Springer.

Cannon, C. H., & Petit, R. J. (2020). The oak syngameon: More than the sum of its

parts. *New Phytologist*, *226*(4), 978–983. https://doi.org/10.1111/nph.16091

Chalk, T. B., Hain, M. P., Foster, G. L., Rohling, E. J., Sexton, P. F., Badger, M. P. S.,

Cherry, S. G., Hasenfratz, A. P., Haug, G. H., Jaccard, S. L., Martínez-García,

A., Pälike, H., Pancost, R. D., & Wilson, P. A. (2017). Causes of ice age

intensification across the Mid-Pleistocene Transition. *Proceedings of the National*

Academy of Sciences, *114*(50), 13114–13119.

https://doi.org/10.1073/pnas.1702143114

Chhatre, V. E., Evans, L. M., DiFazio, S. P., & Keller, S. R. (2018). Adaptive

introgression and maintenance of a trispecies hybrid complex in range-edge

populations of Populus. *Molecular Ecology*, *27*(23), 4820–4838.

https://doi.org/10.1111/mec.14820

Christe, C., Stölting, K. N., Paris, M., Fraïsse, C., Bierne, N., & Lexer, C. (2017).

Adaptive evolution and segregating load contribute to the genomic landscape of

divergence in two tree species connected by episodic gene flow. *Molecular

Ecology*, *26*(1), 59–76. https://doi.org/10.1111/mec.13765

De La Torre, A., Ingvarsson, P. K., & Aitken, S. N. (2015). Genetic architecture and

genomic patterns of gene flow between hybridizing species of picea. *Heredity*,

115(2), 153–164. https://doi.org/10.1038/hdy.2015.19

Dierckxsens, N., Mardulyn, P., & Smits, G. (2017). NOVOPlasty: *De novo* assembly of

organelle genomes from whole genome data. *Nucleic Acids Research*, gkw955.

https://doi.org/10.1093/nar/gkw955

Dobzhansky, T. H. (1936). STUDIES ON HYBRID STERILITY. 11. LOCALIZATION OF

STERILITY FACTORS I N DROSOPHILA PSEUDOOBSCURA HYBRIDS.

Genetics, *21*(113).

Eaton, D. A. R., Hipp, A. L., González-Rodríguez, A., & Cavender-Bares, J. (2015).

Historical introgression among the American live oaks and the comparative

nature of tests for introgression: INTROGRESSION IN THE AMERICAN LIVE

OAKS. *Evolution*, *69*(10), 2587–2601. https://doi.org/10.1111/evo.12758

EL Little Jr. (1971). *Atlas of United States trees. Vol. 1. Conifers and important

hardwoods*. USDA Misc. Publ. 1146 (200 maps).

El Mujtar, V., Sola, G., Aparicio, A., & Gallo, L. (2017). Pattern of natural introgression in

a Nothofagus hybrid zone from South American temperate forests. *Tree Genetics*

& Genomes, *13*(2), 49. https://doi.org/10.1007/s11295-017-1132-1

Ellis, R., & Palmer, M. (2016). Modulation of ice ages via precession and dust-albedo

feedbacks. *Geoscience Frontiers*, *7*(6), 891–909.

https://doi.org/10.1016/j.gsf.2016.04.004

Frankham, R. (2010). Challenges and opportunities of genetic approaches to biological

conservation. *Biological Conservation*, *143*(9), 1919–1927.

https://doi.org/10.1016/j.biocon.2010.05.011

Geraldes, A., Farzaneh, N., Grassa, C. J., Mckown, A. D., Guy, R. D., Mansfield, S. D.,

Douglas, C. J., & Cronk, Q. C. B. (2014). Landscape genomics of populus

trichocarpa: The role of hybridization, limited gene flow, and natural selection in

shaping patterns of population structure. *Evolution*, *68*(11), 3260–3280.

https://doi.org/10.1111/evo.12497

Greiner, S., & Bock, R. (2013). Tuning a ménage à trois: Co-evolution and co-

adaptation of nuclear and organellar genomes in plants. *BioEssays*, *35*(4), 354–

365. https://doi.org/10.1002/bies.201200137

Gutenkunst, R. N., Hernandez, R. D., Williamson, S. H., & Bustamante, C. D. (2009).

Inferring the joint demographic history of multiple populations from

multidimensional SNP frequency data. *PLoS Genetics*, *5*(10).

https://doi.org/10.1371/journal.pgen.1000695

Haeussler, S., Coates, D., & Mather, J. (1990). *Autecology of Common Plants in British*

Columbia: A Literature Review. Forestry Canada, Pacific Forestry Center; British

Columbia Ministry of Forests, Research Branch.

Hamilton, J. A., De la Torre, A. R., & Aitken, S. N. (2015). Fine-scale environmental

variation contributes to introgression in a three-species spruce hybrid complex.

Tree Genetics and Genomes, *11*(1). https://doi.org/10.1007/s11295-014-0817-y

Hamilton, J. A., Lexer, C., & Aitken, S. N. (2013a). Differential introgression reveals

candidate genes for selection across a spruce (Picea sitchensis × P. glauca)

hybrid zone. *New Phytologist*, *197*(3), 927–938.

https://doi.org/10.1111/nph.12055

Hamilton, J. A., Lexer, C., & Aitken, S. N. (2013b). Genomic and phenotypic

architecture of a spruce hybrid zone (Picea sitchensis × P. glauca). *Molecular*

Ecology, *22*(3), 827–841. https://doi.org/10.1111/mec.12007

Hamilton, J. A., & Miller, J. M. (2016). Adaptive introgression as a resource for

management and genetic conservation in a changing climate. *Conservation*

Biology, *30*(1), 33–41. https://doi.org/10.1111/cobi.12574

He, Z., Li, X., Yang, M., Wang, X., Zhong, C., Duke, N. C., Wu, C. I., & Shi, S. (2019).

Speciation with gene flow via cycles of isolation and migration: Insights from

multiple mangrove taxa. *National Science Review*, *6*(2), 275–288.

https://doi.org/10.1093/nsr/nwy078

Heiser, C. (1949). Natural hybridization with particular reference to introgression. *The*

Botanical Review, *15*, 645–687.

Hewitt, G. M. (2004). Genetic consequences of climatic oscillations in the Quaternary.

Philosophical Transactions of the Royal Society B: Biological Sciences,

359(1442), 183–195. https://doi.org/10.1098/rstb.2003.1388

Hoffmann, A., Griffin, P., Dillon, S., Catullo, R., Rane, R., Byrne, M., Jordan, R.,

Oakeshott, J., Weeks, A., Joseph, L., Lockhart, P., Borevitz, J., & Sgrò, C.

(2015). A framework for incorporating evolutionary genomics into biodiversity

conservation and management. *Climate Change Responses*, *2*(1).

https://doi.org/10.1186/s40665-014-0009-x

Hofmeister, B. T., Denkena, J., Colomé-Tatché, M., Shahryary, Y., Hazarika, R.,

Grimwood, J., Mamidi, S., Jenkins, J., Grabowski, P. P., Sreedasyam, A., Shu,

S., Barry, K., Lail, K., Adam, C., Lipzen, A., Sorek, R., Kudrna, D., Talag, J.,

Wing, R., … Schmitz, R. J. (2020). A genome assembly and the somatic genetic

and epigenetic mutation rate in a wild long-lived perennial Populus trichocarpa.

Genome Biology, *21*(1). https://doi.org/10.1186/s13059-020-02162-5

Huang, D. I., Hefer, C. A., Kolosova, N., Douglas, C. J., & Cronk, Q. C. B. (2014).

Whole plastome sequencing reveals deep plastid divergence and cytonuclear

discordance between closely related balsam poplars, Populus balsamifera and P.

trichocarpa (Salicaceae). *New Phytologist*, *204*(3), 693–703.

https://doi.org/10.1111/nph.12956

Huang, X., Struck, T. J., Davey, S. W., & Gutenkunst, R. N. (2023). *dadi-cli: Automated*

and distributed population genetic model inference from allele frequency spectra

[Preprint]. Bioinformatics. https://doi.org/10.1101/2023.06.15.545182

Janes, J. K., & Hamilton, J. A. (2017). Mixing it up: The role of hybridization in forest

management and conservation under climate change. *Forests*, *8*(7).

https://doi.org/10.3390/f8070237

Jansson, S., & Douglas, C. J. (2007). Populus: A Model System for Plant Biology.

Annual Review of Plant Biology, *58*(1), 435–458.

https://doi.org/10.1146/annurev.arplant.58.032806.103956

Jeffrey, E. C. (1916). Hybridism and the rate of evolution in angiosperms. *The American*

Naturalist.

Jump, A. S., & Peñuelas, J. (2005). Running to stand still: Adaptation and the response

of plants to rapid climate change. *Ecology Letters*, *8*(9), 1010–1020.

https://doi.org/10.1111/j.1461-0248.2005.00796.x

Kalyaanamoorthy, S., Minh, B. Q., Wong, T. K. F., Von Haeseler, A., & Jermiin, L. S.

(2017). ModelFinder: Fast model selection for accurate phylogenetic estimates.

Nature Methods, *14*(6), 587–589. https://doi.org/10.1038/nmeth.4285

Katoh, K., & Standley, D. M. (2013). MAFFT Multiple Sequence Alignment Software

Version 7: Improvements in Performance and Usability. *Molecular Biology and*

Evolution, *30*(4), 772–780. https://doi.org/10.1093/molbev/mst010

Kearse, M., Moir, R., Wilson, A., Stones-Havas, S., Cheung, M., Sturrock, S., Buxton,

S., Cooper, A., Markowitz, S., Duran, C., Thierer, T., Ashton, B., Meintjes, P., &

Drummond, A. (2012). Geneious Basic: An integrated and extendable desktop

software platform for the organization and analysis of sequence data.

Bioinformatics, *28*(12), 1647–1649. https://doi.org/10.1093/bioinformatics/bts199

Keller, S. R., Olson, M. S., Salim, S., William, S. A., & Peter, T. (2010). Genomic

809   diversity, population structure, and migration following rapid range expansion in

810   the Balsam Poplar, Populus balsamifera. *Molecular Ecology*, *19*(6), 1212–1226.

811   https://doi.org/10.1111/j.1365-294X.2010.04546.x

812   Legendre, P., & Legendre, L. (2012). *Numerical ecology*. Elsevier.

813   Lepais, O., Petit, R. J., Guichoux, E., Lavabre, J. E., Alberto, F., Kremer, A., & Gerber,

814   S. (2009). Species relative abundance and direction of introgression in oaks.

815   *Molecular Ecology*, *18*(10), 2228–2242. https://doi.org/10.1111/j.1365-

816   294X.2009.04137.x

817   Levsen, N. D., Tiffin, P., & Olson, M. S. (2012). Pleistocene speciation in the genus

818   populus (salicaceae). *Systematic Biology*, *61*(3), 401–412.

819   https://doi.org/10.1093/sysbio/syr120

820   Lexer, C., Fay, M. F., Joseph, J. A., Nica, M. S., & Heinze, B. (2005). Barrier to gene

821   flow between two ecologically divergent Populus species, P. alba (white poplar)

822   and P. tremula (European aspen): The role of ecology and life history in gene

823   introgression. *Molecular Ecology*, *14*(4), 1045–1057.

824   https://doi.org/10.1111/j.1365-294X.2005.02469.x

825   Lexer, C., Kremer, A., & Petit, R. J. (2006). Shared alleles in sympatric oaks: Recurrent

826   gene flow is a more parsimonious explanation than ancestral polymorphism.

827   *Molecular Ecology*, *15*(7), 2007–2012. https://doi.org/10.1111/j.1365-

828   294X.2006.02896.x

Li, H., & Durbin, R. (2010). Fast and accurate long-read alignment with Burrows-

Wheeler transform. *Bioinformatics*, *26*(5), 589–595.

https://doi.org/10.1093/bioinformatics/btp698

Luqman, H., Wegmann, D., Fior, S., & Widmer, A. (2023). Climate-induced range shifts

drive adaptive response via spatio-temporal sieving of alleles. *Nature*

Communications, *14*(1), 1080. https://doi.org/10.1038/s41467-023-36631-9

Martinsen, G. D., Whitham, T. G., Turek, R. J., & Keim, P. (2001). HYBRID

POPULATIONS SELECTIVELY FILTER GENE INTROGRESSION BETWEEN

SPECIES. In *Evolution* (Vol. 55, Issue 7, pp. 1325–1335).

McVay, J. D., Hipp, A. L., & Manos, P. S. (2017). A genetic legacy of introgression

confounds phylogeny and biogeography in oaks. *Proceedings of the Royal*

Society B: Biological Sciences, *284*(1854).

https://doi.org/10.1098/rspb.2017.0300

Menon, M., Bagley, J. C., Friedline, C. J., Whipple, A. V., Schoettle, A. W., Leal-Sàenz,

A., Wehenkel, C., Molina-Freaner, F., Flores-Rentería, L., Gonzalez-Elizondo, M.

S., Sniezko, R. A., Cushman, S. A., Waring, K. M., & Eckert, A. J. (2018). The

role of hybridization during ecological divergence of southwestern white pine

(Pinus strobiformis) and limber pine (P. flexilis). *Molecular Ecology*, *27*(5), 1245–

1260. https://doi.org/10.1111/MEC.14505

Naimi, B., Hamm, N. A. S., Groen, T. A., Skidmore, A. K., & Toxopeus, A. G. (2014).

Where is positional uncertainty a problem for species distribution modelling?

Ecography, *37*(2), 191–203. https://doi.org/10.1111/j.1600-0587.2013.00205.x

Oksanen, J., Blanchet, F., Friendly, M., Kindt, R., & Legendre, P. (2020). Package

"vegan"—Community Ecology Package Version 2.5-6. *Cran. Ism. Ac. Jp.*

Payseur, B. A., & Rieseberg, L. H. (2016). A genomic perspective on hybridization and

speciation. *Molecular Ecology*, *25*(11), 2337–2360.

https://doi.org/10.1111/mec.13557

Petit, R. J., Bodénès, C., Ducousso, A., Roussel, G., & Kremer, A. (2004). Hybridization

as a mechanism of invasion in oaks. *New Phytologist*, *161*(1), 151–164.

https://doi.org/10.1046/j.1469-8137.2003.00944.x

Petkova, D., Novembre, J., & Stephens, M. (2016). Visualizing spatial population

structure with estimated effective migration surfaces. *Nature Genetics*, *48*(1), 94–

100. https://doi.org/10.1038/ng.3464

Sanderson, B. J., Ghambir, D., Feng, G., Hu, N., Cronk, Q. C., Percy, M., Molina

Freaner, F., Johnson, M. G., Smart, L. B., Yin, T., Ma, T., DiFazio, S. P., & Liu, J.

(2023). Phylogenomics reveals patterns of ancient hybridization and differential

diversification that contribute to phylogenetic conflict in willows, poplars, and

close relatives. *Systematic Biology*, *syad042*.

https://doi.org/10.1093/sysbio/syad042

Shang, H., Hess, J., Pickup, M., Field, D. L., Ingvarsson, P. K., Liu, J., & Lexer, C.

(2020). Evolution of strong reproductive isolation in plants: Broad-scale patterns

and lessons from a perennial model group: Evolution of strong barriers in plants.

Philosophical Transactions of the Royal Society B: Biological Sciences,

375(1806). https://doi.org/10.1098/rstb.2019.0544

Shi, Y.-J., Huang, J.-L., Mi, J.-X., Li, J., Meng, F.-Y., Zhong, Y., He, F., Tian, F.-F.,

Zhang, F., Chen, L.-H., Yang, H.-B., Hu, H.-L., & Wan, X.-Q. (2024). A model of

hybrid speciation process drawn from three new poplar species originating from

distant hybridization between sections. *Molecular Phylogenetics and Evolution*,

190, 107966. https://doi.org/10.1016/j.ympev.2023.107966

Slavov, G. T., Difazio, S. P., Martin, J., Schackwitz, W., Muchero, W., Rodgers-Melnick,

E., Lipphardt, M. F., Pennacchio, C. P., Hellsten, U., Pennacchio, L. A., Gunter,

L. E., Ranjan, P., Vining, K., Pomraning, K. R., Wilhelm, L. J., Pellegrini, M.,

Mockler, T. C., Freitag, M., Geraldes, A., … Tuskan, G. A. (2012). Genome

resequencing reveals multiscale geographic structure and extensive linkage

disequilibrium in the forest tree Populus trichocarpa. *New Phytologist*, *196*(3),

713–725. https://doi.org/10.1111/j.1469-8137.2012.04258.x

Soltis, D. E., Morris, A. B., McLachlan, J. S., Manos, P. S., & Soltis, P. S. (2006).

Comparative phylogeography of unglaciated eastern North America. *Molecular*

Ecology, *15*(14), 4261–4293. https://doi.org/10.1111/j.1365-294X.2006.03061.x

Stebbins, G. L. (1959). The Role of Hybridization in Evolution. *Proceeding of the*

American Philosophical Society, *103*, 231–251.

Suarez-Gonzalez, A., Hefer, C. A., Christe, C., Corea, O., Lexer, C., Cronk, Q. C. B., &

Douglas, C. J. (2016). Genomic and functional approaches reveal a case of

adaptive introgression from Populus balsamifera (balsam poplar) in

P. trichocarpa (black cottonwood). *Molecular Ecology*, *25*(11), 2427–2442.

https://doi.org/10.1111/mec.13539

Suarez-Gonzalez, A., Hefer, C. A., Lexer, C., Cronk, Q. C. B., & Douglas, C. J. (2018).

Scale and direction of adaptive introgression between black cottonwood (

Populus trichocarpa) and balsam poplar (*P. balsamifera*). *Molecular Ecology*,

27(7), 1667–1680. https://doi.org/10.1111/mec.14561

Suarez-Gonzalez, A., Lexer, C., & Cronk, Q. C. B. (2018). Adaptive introgression: A

plant perspective. *Biology Letters*, *14*(3). https://doi.org/10.1098/rsbl.2017.0688

Swenson, N. G., & Howard, D. J. (2004). Do suture zones exist? *Evolution*, *58*(11),

2391–2397. https://doi.org/10.1111/j.0014-3820.2004.tb00869.x

Tan, N., Ladant, J.-B., Ramstein, G., Dumas, C., Bachem, P., & Jansen, E. (2018).

Dynamic Greenland ice sheet driven by pCO2 variations across the Pliocene

Pleistocene transition. *Nature Communications*, *9*(1), 4755.

https://doi.org/10.1038/s41467-018-07206-w

Trifinopoulos, J., Nguyen, L.-T., von Haeseler, A., & Minh, B. Q. (2016). W-IQ-TREE: A

fast online phylogenetic tool for maximum likelihood analysis. *Nucleic Acids*

Research, *44*(W1), W232–W235. https://doi.org/10.1093/nar/gkw256

Tuskan, G. A., DiFazio, S., Jansson, S., & Bohlmann, J. (2006). The genome of black

cottonwood, Populus trichocarpa (Torr. & Gray). *Science*, *313*, 1596–1605.

VanWallendael, A., Lowry, D. B., & Hamilton, J. A. (2022). One hundred years into the

study of ecotypes, new advances are being made through large-scale field

experiments in perennial plant systems. *Current Opinion in Plant Biology*, *66*,

102152. https://doi.org/10.1016/j.pbi.2021.102152

Wang, T., Hamann, A., Spittlehouse, D., & Carroll, C. (2016). Locally downscaled and

spatially customizable climate data for historical and future periods for North

America. *PLoS ONE, 11*(6). https://doi.org/10.1371/journal.pone.0156720

Willeit, M., Ganopolski, A., Calov, V., & Brovkin, V. (2019). Mid-Pleistocene transition in

glacial cycles explained by declining CO2 and regolith removal. *SCIENCE*

ADVANCES, *5*(4), 1–8. https://doi.org/DOI: 10.1126/sciadv.aav7337

Table 1. Demographic inference AIC comparisons across 3 population models. Model
abbreviations are defined in the footnote. The lower the AIC and delta AIC the better the
model fit to the data. The first three rows (Table 1a) are results from the topology test to
identify the evolutionary relationship of *interior P. trichocarpa* (I) to *P. balsamifera* (B)
and *coastal P. trichocarpa* (C). The following four rows (Table 1b) are results of four
models with gene flow parameters tested in various scenarios under the best fit
topology (Topology 3; Figure S6) of admixed origin. Bolded models in each of the two
inference routines signify best fit with delta AIC = 0.00.

a) Model	k	AIC	delta AIC
Topology 1: SI	6	68,945.17	4,942.96
Topology 2: SI	6	75,772.18	11,769.97
Topology 3: SI	6	64,002.21	**0.00**

b) Model	k	AIC	delta AIC
Topology 3: T2_SC	12	34,935.00	136.26
Topology 3: T2_IM	12	43,391.55	8,592.80
Topology 3: T3_IM_SC	14	34,798.75	**0.00**
Topology 3: T4_SC_SC	14	35,426.09	627.34

958 *Models; SI (strict isolation), IM (isolation-with-migration), SC (secondary contact), T#
959 (number of time intervals considered)

Table 2. Parameter estimates and 95% confidence intervals (CI) associated with the best fit 3 population demographic model (Topology 3 - T3_IM_SC_IM). Calculations used to unscaling parameter estimates (using theta, mutation rate per generation, and genome length) to meaningful units are provided in Supplementary S3. Each value for T (time) is an interval. Interpretation of each migration (gene flow) parameter is provided in the footnote of this table. The parameter *f* is an estimate of the proportion of *coastal P. trichocarpa* in the formation of *interior P. trichocarpa* at the start of T3. The proportion of *P. balsamifera* is thus 1 - *f.*

Parameter	$\partial a \partial i$ scaled	$\partial a \partial i$ unscaled	95% CI: (lower)	95% CI: (upper)
N_ref (ancestral, theta)	4938	14,031	13,650	14,413
N_1 (*coastal trichocarpa*)	2.60	36,501	35,851	37,150
N_2 (*balsamifera*)	7.12	99,958	96,125	103,792
N_3 (*interior trichocarpa*)	3.83	53,809	51,106	56,512
T1 (ancient gene flow)	2.99	1,259,951	1,254,996	1,264,907
T2 (no gene flow)	0.23	96,645	93,272	100,018
T3 (admixture)	1.82	764,327	763,623	765,031
$M_A \rightarrow mA$	5.34	1.90E-04	1.58E-04	2.22E-04
$M_{12} \rightarrow m12$	1.02	3.64E-05	3.56E-05	3.71E-05
$M_{21} \rightarrow m21$	0.36	1.30E-05	1.15E-05	1.45E-05
$M_{23} \rightarrow m23$	0.25	9.08E-06	7.48E-06	1.07E-05
$M_{32} \rightarrow m32$	0.88	3.13E-05	2.89E-05	3.37E-05
$M_{13} \rightarrow m13$	1.43	5.08E-05	5.01E-05	5.14E-05
$M_{31} \rightarrow m31$	1.28	4.58E-05	4.28E-05	4.88E-05
f	0.348	34.8%	0.346	0.350

**M is a scaled estimate of the number of migrants per generation. *m* is an estimate of gene flow rate. After M (or *m*) the notation of A is ancestral migration, 12 is migrations from *balsamifera* into *coastal trichocarpa*, 21 is migration from *coastal trichocarpa* into *balsamifera*, 23 is migration from *interior trichocarpa* into *balsamifera*, 32 is migration from *balsamifera* into *interior trichocarpa*, 13 is migrations from *interior trichocarpa* into *coastal trichocarpa*, and 31 is migration from *coastal trichocarpa* into *interior trichocarpa.*

Figure 1. Spatial patterns of genetic structure and ancestry across 546 trees sampled from a) seven transects of the *Populus balsamifera* and *P. trichocarpa* hybrid zone, labeled by transect; b) transect-specific genomic structure based on Principal Component Analysis for PC1 and PC2 c) results of admixture analysis for each sample at *K* = 3, the best supported cluster assignment on the basis on lowest cross validation (CV) error; d) geographic distribution of *K* = 3 ancestry proportions illustrated as pie charts; e) ML phylogeny of chloroplast evolutionary relationships among parental-type samples from *K* = 3 admixture results (Q > 0.90 for each cluster). Bootstrap confidence is provided on each branch.

Figure 2. Genetic structure (PCA) and Redundancy analysis (RDA) based on associations between 6 climate, 3 geographic variables for a) genome-wide nuclear data of all 546 samples analyzed, versus b) genome-wide nuclear and c) chloroplast differentiation across parental-types. In each RDA plot, the direction and length of each arrow corresponds with the strength of prediction (i.e. loading) along each RDA axis. Climate and geographic variables include MAT (mean annual temperature), TD (continentality), PAS (precipitation as snow), RH (relative humidity), MAP (mean annual precipitation), CMD (climate moisture deficit), latitude, longitude, and elevation. Patterns between chloroplast and nuclear DNA are congruent with the exception of one sample (ID= 757_S202_L004) which has *interior P. trichocarpa* nuclear genetic background with *P. balsamifera* chloroplast identity.

1012
1013
1014
1015

1016
1017
Figure 3. Estimated effective migration surfaces (EEMS) analysis of the relationship between genetic distance and geographical distance for 546 sampled trees across seven transects of the *P. trichocarpa* and *P. balsamifera* hybrid zone to visualize potential barriers to gene flow. Panel a) is a raster map of elevational differences (in meters) across the studied region, and panel b) illustrates putative barriers to gene flow within and among sampled transects. The black dots represent the geographic location of each of the 546 samples. Regions with negative migration rate predictions (brown colors on the log 10 scale) indicate lower genetic connectivity (gene flow) than expected. Regions with positive migration rate predictions (blue colors on the log 10 scale) indicate higher genetic connectivity than expected.

1028

1029

1030

1031

1032

1033

1034

1035

1036
1037
1038 **Figure 4.** Distribution of samples used in three population demographic inference (a)
1039 and the best fit model of divergence among *coastal P. trichocarpa* (in yellow), *P.*
1040 *balsamifera* (in dark blue), and *interior P. trichocarpa* (in light blue). Parameter
1041 estimates and confidence intervals for the 3-population model are provided in Table 2.
1042 Note that T1, T2, and T3 are time intervals in millions of years (my).
1043

1044

1045
1046