Alternative splicing and its regulation in the malaria vector *Anopheles gambiae*

Bárbara Díaz-Terenti[1], Elena Gómez-Díaz[1*]

[1]Instituto de Parasitología y Biomedicina López-Neyra, Consejo Superior de Investigaciones Científicas (IPBLN, CSIC), Granada, Spain

* Correspondence:

Elena Gómez-Díaz

elena.gomez@csic.es

Acknowledgements

We are grateful to Dr Lisa Ranford-Cartwright and Dr José L. Ruiz for sharing data. We thank the bioinformatics unit of the IPBLN-CSIC for computing facilities and advice on the analysis.

Abstract

Alternative splicing (AS) is a highly conserved mechanism that allows to expand the coding capacity of the genome, by modifying how multiple isoforms are expressed or used to generate different phenotypes. Despite its importance in physiology and disease, genome-wide studies of AS are lacking in most insects, including mosquitoes. Even for model organisms, chromatin associated processes involved in the regulation AS are poorly known. In this study, we investigated AS in the mosquito *Anopheles gambiae* in the context of tissue-specific gene expression and mosquito responses to a *Plasmodium falciparum* infection, as well as the relationship between patterns of differential isoform expression and usage with chromatin

accessibility changes. For this, we combined RNA-seq and ATAC-seq data from *A. gambiae*

midguts and salivary glands, and from infected and non-infected midguts. We report differences

between tissues in the expression of 456 isoforms and in the use of 211 isoforms. Secondly, we

find a clear and significant association between chromatin accessibility states and tissue-specific

patterns of AS. The analysis of differential accessible regions located at splicing sites permitted

the identification of several motifs resembling the binding sites of *Drosophila* transcription

factors. Finally, the genome-wide analysis of tissue-dependent enhancer activity revealed that

approximately 20% of *A. gambiae* transcriptional enhancers annotate to a differentially

expressed or used isoform and that their activation status is linked to AS differences between

tissues. This research illuminates the role of AS in gene expression in vector mosquitoes, and

identifies regulatory regions potentially involved in AS regulation, which could reveal novel

strategies for vector control.

Keywords: Mosquitoes, Plasmodium, Isoform expression, Isoform usage,

Chromatin accessibility

Abbreviations [1]

1   Introduction

42 Alternative splicing (AS from here) is an important source of phenotypic variation in eukaryotes

43 that allows the generation of multiple transcripts from a single gene, expanding the coding

44 capacity and functionality of the genome. AS regulates gene expression by acting on mRNA

---

[1] **Abbreviations**: DEMG, Differential isoform expression of multisoform genes; DUI, Differential isoform usage of multisoform genes; SS, splicing sites.

transcripts at a post-transcriptional level and is critical for diverse cellular processes, including

cell differentiation and development as well as cell reprogramming and tissue remodeling.

AS is mediated by four types of basic events: alternative 5′ splice-site, alternative 3′ splice-site,

cassette-exon inclusion or skipping, and intron retention. These processes can be combined to

obtain dozens of isoforms from a single gene. On the other hand, AS regulation include different

mechanisms as the temporal and spatial differential expression of the isoforms, the activation of

the RNA Pol II or the on-off regulation by nonsense-mediated decay (Wang and Cooper, 2007,

Nilsen and Graveley, 2010, Kelemen et al., 2013, Shenasa and Hertel, 2019, Ule and Blencowe,

2019, Wright, Smith and Jiggins, 2022). Another process associated to the AS is regulating

isoform usage. Isoform usage is defined as, the participation of each isoform in the gene

expression, so when these proportions switch between two conditions, some isoforms are

differentially used (Vitting-Seerup and Sandelin, 2017, Merino, Conesa and Fernández, 2019,

De La Fuente et al., 2020, Li et al., 2020a, Chen et al., 2022). As a result, the function of gene

might change because different variant proteins are produced, while the total gene expression

can remain unaltered.

The type of events and mechanisms of AS vary significantly amongst species. Despite of this,

AS is highly conserved in eukaryotes playing fundamental functions in health and disease,

adaptation to the environment, and defence against pathogens (Nilsen and Graveley, 2010,

Kelemen et al., 2013, Martinez and Lynch, 2013, Bush et al., 2017, Vitting-Seerup and

Sandelin, 2017, Ule and Blencowe, 2019, Zhao, 2019, Li et al., 2020a, Song et al., 2020, Verta

and Jacobs, 2022, Wright, Smith and Jiggins, 2022).

A simple way to study AS is to quantify and compare the expression of isoforms that are

generated in different tissues or experimental conditions (Katz et al., 2010, Glaus, Honkela and

Rattray, 2012). Transcriptomic data obtained by RNA-seq is mostly used for gene level analysis

(Vitting-Seerup and Sandelin, 2017, Zhao, 2019), but offers the possibility to study AS to

analyze and count known transcripts or to detect new splicing events (Zhao, 2019).

Regulation of AS is intrinsically related to chromatin dynamics. To obtain diverse transcripts,

cis and *trans*-regulatory elements participate both in the generation of transcripts and in marking

splice sites (Bush et al., 2017, Shenasa and Hertel, 2019, Zhao, 2019, Verta and Jacobs, 2022).

In addition, AS regulation can be affected by posttranslational histone modifications,

nucleosome positioning, and chromatin accessibility (Davuluri et al., 2008, Luco et al., 2011,

Montes et al., 2012, Klemm, Shipony and Greenleaf, 2019, Nussinov et al., 2021). However,

how these different epigenetic layers interact, and which are the mechanisms that underlie the

great diversification of gene functions through AS, has been poorly investigated (Davuluri et al.,

2008, Luco et al., 2011, Klemm, Shipony and Greenleaf, 2019). The integrated analysis of

transcriptomic and epigenetic data could bring new insights into how AS is regulated and how it

changes under different conditions.

AS has been widely studied in model organisms like *Drosophila melanogaster*, *Danio renio* or

Caenorhabditis elegans (Zahler, 2005, Venables, Tazi and Juge, 2012, Lin et al., 2022). In

Drosophila melanogaster AS plays important roles in sex-determination, muscle, neural

development among other biological processes (Venables, Tazi and Juge, 2012). More precisely

the relationship between AS and chromatin has been investigated in *Drosophila* gonads. That

study revealed that the splicing factors expressed in the undifferentiated cells act coordinately

with chromatin remodeling factors and histone modifying enzymes to maintain the transcription

network through post-transcriptional mechanisms (Gan et al., 2010). Despite of this, there is a

lack of the knowledge in how the AS interacts with chromatin to regulate gene function under

different physiological or cellular conditions (Nilsen and Graveley, 2010, Luco et al., 2011,

Kelemen et al., 2013).

In mosquitoes, little is known about AS and its regulation. Given that AS is evolutionarily

conserved, one would expect that functions and mechanisms are maintained across insects.

There is, however, marked inter-species variation in alternative splicing that probably resulted

through selective pressures exerted by environment and lifestyle (Kurtz and Armitage, 2006,

Malko et al., 2006). In *Aedes aegypti*, previous studies have focused on sex differentiation

(Salvemini et al., 2011, 2013, Calkins et al., 2015). In the human malaria mosquito *Anopheles*

gambiae, the study of AS is limited to a few genes, such as *doublesex* and *Dscam* (Tsujimoto et

al., 2013, Krzywinska et al., 2016, Sinkins, 2016, Djihinto et al., 2022), tissues like salivary

glands (Dixit et al., 2009) or signalling pathways (Meister et al., 2005). These studies suggest

that AS could play a crucial role in vector competence and mosquito immunity. However, a

more systematic analysis onto the role of AS and its regulation in mosquitoes, is lacking.

In this study, we combined transcriptomic and chromatin accessibility data of infected and non-

infected *A. gambiae* salivary glands and midguts to interrogate about AS regulation. In

particular, which isoforms belonging to multisoform genes are differentially expressed and used

in different mosquito tissues infected with the malaria parasite *Plasmodium falciparum*, and

whether chromatin-associated mechanisms underlie these events (Love, Huber and Anders,

2014, Vitting-Seerup, Sandelin and Berger, 2019). We contend that a deeper understanding of

the mechanisms of AS regulation can reveal key *cis* and *trans* regulatory elements, like splicing

transcription factors and enhancers, involved in mosquito responses to *P. falciparum* infection

and/or vector competence.

113 2   Results

114 2.1   Alternative splicing (AS) regulates tissue-specific gene expression in

115   infected *A. gambiae*

Considering the 1,263 multisoform genes detected in *A. gambiae*, we aimed to identify

alternative splicing events and their role in gene expression regulation in different tissues and

infection conditions. For this, we performed two types of analysis: one to identify differentially

expressed isoforms belonging to multisoform genes (DEMG), and another to detect isoforms

differentially used (DUI); isoforms that are altered in their relative proportion to the total

expression of the gene.

In the comparison between *P. falciparum* infected and non-infected (control) mosquito midguts,

32 isoforms appeared differentially regulated out of the 848 multisoform genes present in this

dataset (with counts > 10 as the detectability threshold). Of these, 19 (2%) correspond to

differentially expressed and 13 (1.5%) to differentially used isoforms. In salivary glands, we

detected 977 expressed multisoform genes of which 91 appear differentially regulated; 56 (6%)

were differentially expressed and 35 (3.5%) differentially used.

Then, we examined differences in AS regulation between tissues: midguts and salivary glands

infected with *P. falciparum*. We detect 920 multisoform genes: 456 (50%) DEMG and 211

(23%) DUI.

We also report multisoform genes in which, one or more isoforms were differentially expressed

and used (Fig 1a). In our results, the overlap between AS mechanisms represented 22% (105) of

the multisoform genes that display AS when comparing midguts and salivary glands (Fig 1b),

and less than 1% (0 for the midguts and 1 for the salivary glands) when comparing infected and

non-infected mosquito tissues (Supplementary table S1 and S2).

To test for differential functionality of DEMG and DUI isoforms between tissues, we performed

GO enrichment analysis. For the differentially expressed isoforms that appear more expressed in

midguts, the 10 first GO terms (ordered by less p-value) include metabolic process, regulation

of response to stimulus and signaling. Contrary, differentially expressed isoform genes that are

more expressed in salivary glands correspond to functions related to fiber and cytoskeleton

organization, ion transport, and muscle contraction (Supplementary table S3).

In the case of differentially used isoforms that appear overused in midguts, the 10 first GO

terms correspond to functions related to muscle, contraction, development and differentiation,

proton transmembrane transport and metabolic processes. In salivary glands, overused isoforms

displayed functions like metabolic process of nucleotides and carboxylic acids, and muscle

contraction (Supplementary table S3).

Next, we described the types of events that lead to differential isoform expression (DEMG)

and/or isoform usage (DUI) in our set of *A.gambiae* multisoform genes. Our results show that

Exon Skipping (ES) and Alternative Start Site (ATSS) are the two most frequent types of events.

For differentially expressed isoforms, we report differences in the frequency of these events

between conditions for both ES and ATSS: 68 ES events in *Inf MG vs.* 102 ES events in *Inf SG*

and in the case of ATSS*,* 89 ATSS events in *Inf MG vs.* 105 ATSS events in *Inf SG.* For the

differentially used isoforms, only the frequency of ATSS is moderately altered when comparing

Inf MG vs. Inf SG (*43 ATSS events in SG compared to 51 events in MG*).

Finally, we interrogated whether our differentially expressed and used isoforms coincided with

genes known to be involved in mosquito immunity (Table S4) (Sreenivasamurthy et al., 2013).

We found 8 genes among our differentially expressed and used isoforms, that encode for

proteins involved in anti-*Plasmodium* mosquito responses (STAT, PGRPLC, OXR1). One

example are the AGAP005203-RA and AGAP005203-RE isoforms of the PGRPLC gene,

identified in the present work as differentially expressed (DEMG). The RA isoform is

overexpressed during midgut infection, while RE isoform, is overrepresented in infected

salivary glands compared to control. RB and RC appear similarly expressed, and the rest of

isoforms have values below the expression threshold, so they were considered as non-expressed

(Fig 1c).

2.2   Alternative Splicing is positively correlated with chromatin

accessibility in malaria-infected *A. gambiae* tissues

2.2.1   Tissue-specific isoform expression is positively associated to differential

chromatin accessibility

First, we tested if higher expression of the isoforms was associated with higher levels of

chromatin accessibility in the isoform promoter (1kb upstream of the TSS/ATG of the isoform)

or the isoform body. The results for the differentially expressed group show a positive and

significant correlation between isoform expression and chromatin accessibility for the isoform

body, both in salivary glands (*rho* = 0,328, *p-value* = 8,095e-07) and midguts (*rho* = 0,369, *p-*

value = 3.009e-07), but not for the promoter region (Table S5A) (Fig 2).

Next, we examined whether the relationship is quantitative, that is, if the levels of expression of

the isoform differentially expressed (categorized as high, medium, or low) were correlated with

the degree of chromatin accessibility in the isoform promoter. This relationship appears not

significant either in midguts or salivary glands (*rho* < 0,02, *p-value* > 0.1 for all levels and

tissues, Table S5B) (Fig S2).

For the analysis of chromatin accessibility in relation to differential usage, we used processed

data of differential accessibility peaks previously published by us (Ruiz, Ranford-Cartwright

and Gómez-Díaz, 2021). We then selected DiffBind peaks that were located at the splicing site

or in the region nearby (+/-100 Kb), and that annotated to differentially used isoforms. We

report a clear association between differential usage and accessibility with 98 differentially used

isoforms, 54,4% of the total number of isoforms differentially used, displaying a DiffBind peak

located at or nearby splicing sites.

2.2.2   Motif analysis

We performed a motif analysis using HOMER on the set of DiffBind peaks that annotated to

differentially used isoforms and located at or nearby splicing sites. The purpose of this analysis

was to detect overrepresented motifs that point to transcription factors involved in mosquito AS

regulation.

Using the hierarchical clustering of MEME *de novo motif discovery* option, several motif hits

with distinct similarities with *Drosophila* sequences appear overrepresented (summary in Table

1) (Table S6B). For the subset of differentially used isoforms, we found *de novo* motifs related

with brachyentron (byn), implicated in midgut development, snail (sna), which participates in

epithelial transition, grainy head (grh) involved in epithelial cell fate or chromatin remodellers

like the Chromatin-linked adaptor for MSL proteins (CLAMP).

Now analyzing the set of "known" motifs, we found only two motifs overrepresented. The

motif CATCMCTA matches a *Drosophila* promoter element of unknown function, and the motif

GGYCATAAAW corresponds to the caudal homeobox of *Drosophila* (Table S6A).

2.3   Mosquito enhancers involved in AS

We aimed to study if *A. gambiae* previously identified transcriptional enhancers which

participated in the regulation of AS. Such a relationship has been shown in human and mouse in

which 5% of the enhancers appear associated to AS changes (Shiau et al., 2021). For this

analysis, we used *A. gambiae* enhancers described by our lab in a previous study (Ruiz,

Ranford-Cartwright and Gómez-Díaz, 2021). This dataset included 4,268 enhancers identified

computationally by orthology with *Drosophila* enhancers, or experimentally discovered by

others in mosquitoes, and that were validated by ours using the combined analysis of differential

chromatin accessibility and histone modifications marking, as well as gene expression changes

between tissues (Ruiz, Ranford-Cartwright and Gómez-Díaz, 2021). Of these mosquito

enhancers, 811 appear active in our dataset.

In the differentially expressed isoforms we found a total of 125 accessible enhancers that

annotate to 76 genes (21,1% of the 360 differentially expressed isoforms belonging to

multisoform genes) (Supplementary Table S7A). Of these 125 accessible (active) enhancers, 26

of them coincide with a Diffbind peak, that is, they are differentially accessible between

infected midguts and salivary glands.

For the group of differentially used isoforms, we identified 39 accessible enhancer regions that

annotate to 23 genes (19,8% of the 116 differentially used isoforms belonging to multisoform

genes) (Supplementary Table S7B), and 11 exhibit Diffbind peaks, indicating differential

accessibility between infected midguts and salivary glands.

Lastly, we examined the location of the enhancers annotated to differentially expressed and used

isoforms. For both, the DEMG and DUI groups, the number of intragenic enhancers almost

double of the number of intergenic (DEMG: 44 intergenic and 81 intragenic enhancers; DUI: 14

intergenic and 25 intragenic enhancers).

One example is the gene AGAP005234 that encodes for a copper-zinc superoxide dismutase.

This protein has functions involved in mosquito immunity, including anti-Plasmodium

responses, as well as response to oxygen radicals. (Marikovsky et al., 2003, Molina-Cruz et al.,

2008). The gene contains two different isoforms (AGAP005234-RA and AGAP005234-RB),

RB isoform is differentially expressed and used in salivary glands and RA is differentially used

in midguts. The enhancer associated to this gene is intergenic, located 2 Kbp upstream of the

start of the gene, and its accessibility in higher in SG coinciding with the expression and use

that is also higher in SG (Fig 3).

3   Discussion

Alternative splicing has been relatively little studied in mosquitoes, and the relationship of this

process with chromatin dynamics remains mostly unknown. In this study we investigated

alternative splicing in *A. gambiae* and the role of chromatin structure in its regulation. We

compared patterns of AS in different conditions, first between two mosquito tissues: midguts

and salivary glands, and second between *P. falciparum* infected and non-infected (control)

mosquito midguts and salivary glands.

The first part of this work aimed at identifying how prevalent is AS in mosquitoes and how AS

patterns change between tissues and infection status. From a total of 1,263 *A. gambiae* genes

susceptible of AS, we detect around a 67-77% being differentially expressed or used at the level

of isoforms.

These differences were largely due AS events at the tissue-level. Indeed, when comparing the

two infected tissues, the 49% of isoforms (456) appeared differentially expressed (DEMG) and

22.9% (211) differentially used (DUI), suggesting that AS plays an important role in regulating

tissue-specific expression in *A. gambiae*. The role of AS in tissue specific processes has been

reported in different organisms like in humans, *D. melanogaster*, *C. elegans* (Wang et al., 2008,

Telonis-Scott et al., 2009, Venables, Tazi and Juge, 2012, Ragle et al., 2015) or even in

Anopheles stephensi (Sreenivasamurthy et al., 2017).

However, while in this study the AS patterns vary significantly between tissues, differences

were minimal when comparing infected and uninfected tissues for both groups, differentially

expressed, and used isoforms (only 32 DEMG and DUI isoforms detected in case of midgut

infection and 91 DEMG and DUI isoforms detected in case of salivary glands infection). These

results can be explained by the inter-sample variability in these comparisons as revealed the

PCA analysis. That may be reducing the statistical power of the expression and differential use

tests, resulting in low numbers of differentially expressed and used isoforms.

Among the tissue-specific multisoform genes, we report several genes with immune functions

that have a specific isoform differentially expressed in midguts and another isoform

differentially expressed in salivary glands, meaning that each has a different function depending

on the tissue in which it is expressed. An example of this is the AGAP005203 gene, that encodes

for a Peptidoglycan recognition protein (PGRPLC) and is involved in the Imd pathway (Meister

et al., 2009). This protein has been reported to be implicated in anti-*Plasmodium* responses at

the level of the mosquito midgut, but without information about the functionality of their

isoforms (Sreenivasamurthy et al., 2013). This gene consists of six isoforms corresponding to

three different variants. PGRPLC gene produces three primary protein isoforms each with a

distinct PGRP domain and an optional 75-nucleotide cassette located at the 3' end of the shared

exon 3. Consequently, each isoform is present in two forms, the longer version is 25 amino

acids longer than the shorter version. AGAP005203 has 2 isoforms differentially expressed in

our data all of them corresponding to the shorter version of the variants. RA form has been

reported to be overexpressed during midgut infection and has been proposed to be involved in

the mosquito immune response against *Plasmodium* (Meister et al., 2009). On the other side, the

RE isoform, was overexpressed in salivary glands, and although its function in this tissue is still

unknown. Previous work reported that the RD and RE isoforms are positively regulated in

response to stress and antimicrobial responses (Lin et al., 2007, Chen, Ling and Weng, 2009,

Meister et al., 2009).

Chromatin dynamics and transcriptional regulation are intimately related, but its role in the

regulation of alternative splicing remains little known. There are very few studies that

investigated how epigenetic mechanisms such as histone modifications or nucleosome

positioning, would contribute to changes in the AS pattern. This is despite the functional

consequences that AS events can have during development and disease (Simon et al., 2014,

Petrova, Song, Consortium, NordströmNordstr, et al., 2022). Previous evidence in model

organisms reported local changes in nucleosome positioning histone modifications and

chromatin accessibility near the exon that undergoes splicing (Naftelberg et al., n.d.,

Chodavarapu et al., 2010, Li et al., 2020b).

Here, we proposed a model for *A. gambiae* to describe the association between isoform

expression and/or usage and chromatin structure dynamics (Fig. 4). For differentially expressed

isoforms we expected increased promoter accessibility in the tissue where the isoform was more

expressed (Li et al., 2020b, Ruiz, Ranford-Cartwright and Gómez-Díaz, 2021). For the

differentially used isoforms, we expected accessibility changes associates to the differential use,

but localized at or nearby AS regulatory regions (splicing sites). In agreement with this model,

our results indicate that there is a relationship between the differential expression and the use of

isoforms with changes in chromatin accessibility enrichment suggesting that chromatin structure

would play a role in AS-mediated gene expression regulation in *A. gambiae*. However, this

relationship was only significant in the region of the isoform body. This is supported by

previous studies in mammals and plants, reporting higher chromatin accessibility coinciding

with the region of the intron retention (Ullah et al., 2018, Petrova, Song, Consortium,

Nordström, et al., 2022).

In addition, changes in accessibility along the isoform body have also been shown to alter the

speed of RNA polymerase II (RNA pol II) (Saldi et al., 2016) or, alternatively, impact the

recruitment of splicing factors through direct or indirect recognition of the different chromatin

marks (Schor, Gómez Acuña and Kornblihtt, 2013). Indeed, an increase in chromatin

accessibility in histone-depleted genes was shown to alter the elongation rate of Pol II, resulting

in several transcripts with increased intron retention and changes in alternative splicing (Jimeno-

González et al., 2015).

For differentially used isoforms, we already predicted that changes at the level of splicing sites

would be more important. Our results confirm the existence of differential accessibility peaks at

splicing sites for around 54% of the isoforms. What we cannot predict from our results,

however, are the precise mechanism and the functional consequences of this change, that is,

whether a local increase in chromatin accessibility at the splicing site would result in one or

another alternatively spliced isoform due the selective binding of splicing repressor versus an

inducer. Indeed, according to a recent study, the type of event and the choice of a particular

isoform would be multifactorial and depend on a trade-off between chromatin accessibility at

regulatory sequences (such as splicing sites), and variable RNA pol II elongation rates (Pai and

Luca, 2019).

Motif analysis can give some insights on the underlying mechanisms of AS regulation. Looking

at the *de novo* motifs identified in the differentially expressed and used isoforms, we found

overrepresented motifs that resemble several transcription and slicing factors of *Drosophila*

such as snail (sna), grainy head (grh) and chromatin-linked adaptor for MSL proteins (CLAMP).

In the case of CLAMP, this is a chromatin remodeling protein with many different functions in

regulating RNA Pol II dynamics, chromatin architecture, and alternative splicing

(Schuettengruber et al., 2009) (Urban et al., 2017). It has been described for example, that the

interaction of CLAMP with the proteins Squid (Sqd) and Syncrip (Syp), members of the

heterogeneous nuclear ribonucleoproteins (hnRNPs), has a role in the regulation of the AS in

Drosophila (Blanchette et al., 2009).

Another aspect that we approached in this study is the relationship between the enhancer

function, based on histone PTMs and accessibility patterns, and AS events in *A. gambiae*.

Previous evidence suggests that enhancers could mediate isoform-specific expression by

regulating the production and binding of antisense transcripts, as well as by regulating

messenger RNA 3′ end processing (Onodera et al., 2012, Shiau et al., 2021, Kwon et al., 2022).

In such cases, enhancers were expected to be located intragenically, near the exons or introns

that undergo AS (Lam and Hertel, 2002, Matlin, Clark and Smith, 2005).

When we look at the presence of enhancers in our AS susceptible isoforms (DEMG and DUI),

we discovered that approximately 19-23% of differentially expressed and used isoforms have an

enhancer annotated, and approximately 18-20% of these are active in our study, i.e. accessible.

This observation suggests a relationship of the enhancers with the expression and use of the

isoforms. Indeed, we also report that 20% of the enhancers active in differentially expressed

group and 28% in differentially used isoforms correspond to differentially accessible enhancers,

i.e. they have differential activation status, between tissues. Furthermore, most of our accessible

enhancers (64.6% of the total of both groups) appear to be intragenic. Although their function in

alternative splicing still needs to be experimentally confirmed, the fact that their activation

status changes according to changes in the expression and use of the isoforms to which they are

annotated, suggests a possible role in AS-mediated isoform expression regulation.

One example of a AS gene with an associated enhancer that is differentially accessible between

conditions is the immune gene AGAP005234. In *A. gambiae* the function of this gene is

associated the production of reactive oxygen species (ROS), that have been implicated in the

immune response to pathogens (Marikovsky et al., 2003), including a *Plasmodium* infection

(Molina-Cruz et al., 2008). Our data shows that the RB isoform is more used and expressed in

salivary glands, whereas the RA isoform is only used in midguts. There is an enhancer

annotated to the AGAP005234 gene that is more accessible in salivary glands, suggesting a

possible regulatory function of this enhancer in determining the choice of which of the isoforms

of this gene is used (RA or RB) or differentially expressed (RB).

To conclude, this is, to our knowledge, the first study that analyze comprehensively alternative

splicing and its regulation in *A. gambiae*. We report genome-wide alterations in isoform

expression and usage between different tissues and, less so, linked to the infection status. Also,

the study of the relationship between AS with chromatin, points to the importance of chromatin

structure dynamics at splicing sites in tissue-specific gene expression regulation. Finally, the

identification of active enhancers associated to differentially expressed and used isoforms

suggests a possible role of these regulatory elements on the splicing process. Our study lays the

foundation for further research on the mechanisms of alternative splicing in *A. gambiae*,

including the functional characterization of regulatory elements like enhancers and splicing

factors as potential candidates for vector control.

4 Experimental Procedures

4.1 External data

External RNA-seq data analysed in this study was downloaded from GEO accession number

GSE120076 and corresponds to samples infected and non-infected in midguts (Ruiz et al.,

2019). Unpublished RNA-seq data corresponds to samples of salivary glands infected and non-

infected from the same study (Ruiz et al., 2019), and has been assigned to GEO accession

number GSE243215. Previously published ATAC-seq data for infected tissues, both in midgut

and salivary glands, was downloaded from GEO accession number GSE152924 (Ruiz, Ranford-

Cartwright and Gómez-Díaz, 2021).

4.2 Genes susceptible of alternative splicing and experimental design

To study the role of Alternative Splicing in *A. gambiae* we considered genes that are susceptible to be regulated by this mechanism. Only 1,263 genes in the *A. gambiae* genome (13,094 genes) have more than one transcript and could be classified as multisoform genes.

We used 12 RNA-seq and 4 ATAC-seq datasets corresponding to 8 *A. gambiae* infected and 4 non-infected (control) samples from two mosquito tissues (midguts and salivary glands). Midgut datasets correspond to published data from two previous studies(Ruiz et al., 2019, Ruiz, Ranford-Cartwright and Gómez-Díaz, 2021). And there is additional data from two samples (infected and non-infected salivary glands) that are new to this study. Samples are distributed in three different comparisons: infection *vs.* control in midguts and salivary glands, and infected midguts *vs.* infected salivary glands, having each of them 2 replicates per condition.

4.3 RNA-seq data analysis

Quality control of RNA-seq data was performed with FastQC v0.11.9 (Babraham Bioinformatics - FastQC A Quality Control tool for High Throughput Sequence Data, 2022). A replicability test based on correlation analysis and principal component analysis (PCA), support the use of the different samples as replicates (Supplementary figure 1).

We used SortMeRNA v4.2.0 (Kopylova, Noé and Touzet, 2012) to determine and eliminate possible ribosomal RNA contamination. Filtered reads were mapped to the reference genome (Anopheles gambiae PEST genome, version P4.12) available in VectorBase (release 53) (Giraldo-Calderón et al., 2015), using Salmon v.0.11.2 (Patro et al., 2017). This software is specifically used for the alignment and quantification of isoforms (Zhang et al., 2016) and was run using the standard parameters *-l A --validateMappings --go 4 --mismatchSeedSkip 5*. For this analysis we applied a conservative filter, only isoforms with a count value greater than 10 were considered as being expressed. A total number of 1,561 (Inf MG vs. Inf SG comparison),

402 1,394 (Inf vs. Ctrl MG), 1,679 (Inf vs. Ctrl SG) isoforms passed this threshold and were

403 included in all subsequent analysis.

4.4   AS mechanisms and differential analyses

The counts generated by Salmon were used to perform various analyses. First, to study the most

prevailing alternative splicing mechanisms in each condition we used the R package

IsoformSwitchAnalyceR v1.12.0 (Vitting-Seerup, Sandelin and Berger, 2019). We applied the

function *analyzeAlternativeSplicing* with the argument *onlySwitchingGenes=F*. We then

compared the frequencies of the AS events between conditions (tissues and infection status) to

determine how are they altered.

The differential expression analysis was performed using the R package (v4.0.2) DESeq2

v1.30.0, (Love, Huber and Anders, 2014). Taking multisoform genes, we considered

differentially expressed isoforms (DEMG), those that have differential expression between

conditions (the isoform not the gene). To be differentially expressed, isoforms must have a

significant P-value of less than 0.05 and a LogFoldChange greater than 1 or less than -1

(corresponding to a 10% change). In the comparison between Inf MG vs. Inf SG a LogFC < 0

means overexpression in MG, whereas a LogFC > 0 means overexpression in SG. In the

comparison between Inf vs. Control a LogFC > 0 means more expression in the infected tissue.

The study of the differential usage of the isoforms was performed using

IsoformSwitchAnalyceR v1.12.0 (Vitting-Seerup, Sandelin and Berger, 2019). In this case we

used the parameter, difference in isoform fraction (dIF) to measure the magnitude of change.

The isoform fraction (IF) is a measurement that determines the proportion of the total

expression of a parent gene that comes from a particular isoform (isoform_exp / gene_exp).

Isoforms significantly differentially used (DUI) were those with a significance P-value below

0.05. In the comparison between Inf MG vs. Inf SG a dIF < 0 means more used in MG, and dIF

> 0 means more used in SG. In the comparison between Inf vs. Control a dIF < 0 means more

usage in the infected tissue.

We performed Gene Ontology (GO) terms overrepresentation tests for the sets of differentially

expressed or used multisoform genes between Infected midguts *vs.* Infected salivary glands

using VectorBase Overrepresentation Test and ImmunoDB database to know which isoforms are

immunologically relevant (Giraldo-Calderón et al., 2015, EZlab | Computational Evolutionary

Genomics group, 2022). We chose the p-value as a measure of the likelihood that a certain GO

term appear among the genes in your results more often than it appears in the set of all genes for

that organism (background) and we applied a threshold of p-value < 0.05.

4.5   ATAC-seq data analyses

For the analysis of ATAC-seq data in relation to AS, we used processed data from a recent

article previously published by our group (Ruiz, Ranford-Cartwright and Gómez-Díaz, 2021)

the nucleosome free reads, the ATAC peaks resulting from the analysis with MACS2 (Zhang et

al., 2008a) and the differentially accessible peaks resulting from the analysis with Diffbind

(Stark R. and Brown G., 2011).

In brief, raw reads were trimmed 20 bases from the 3′ end of each read (−3 20) and aligned to

the *A. gambiae* PEST genome with Bowtie2 (Langmead and Salzberg, 2012) (v2.4.1) using

default parameters, except for –no-unal –no-mixed -X 2000. We then applied a MAPQ score

threshold of 10 and sorted and deduplicated the reads using SAMtools (Li et al., 2009) (v1.10).

To adjust the known bias and ensure the mapping of Tn5 cutting sites, we shifted aligned reads

+4 bp for + strands and −5 bp for − strands with ATACSeqQC (Ou et al., 2018) (v1.10). We

removed not properly paired reads and extracted nucleosome-free fragments with a size

threshold of 130 bp (SAMtools). We performed peak-calling on nucleosome-free reads with

MACS2 (Zhang et al., 2008b) (v.2.1.2) using the callpeak module and the following parameters:

-f BAMPE -g 273109044 -q 0.01 -B –keep-dup all –nomodel –nolambda. We refer to ATAC-seq

peaks as Tn5 hypersensitive sites (THSs). To identify THSs unique or common across samples,

we used BEDTools intersect (Quinlan and Hall, 2010) requiring a minimum overlap of 51% (-

f/-F 0.51). We annotated THSs to genomic features combining HOMER (Heinz et al., 2010)

(v4.11).

From the nucleosome free reads, we quantified the ATAC-seq nucleosome-free signal

enrichment at our genomic regions of interest (promoter, isoform body) using BEDTools

v2.30.0 (Quinlan and Hall, 2010) with the parameter *intersect* and the option *-c*. Read counts

were normalized (RPKM).

We used the DiffBind R package (Stark R. and Brown G., 2011) (v2.14) to assess differential

chromatin accessibility at given locations between P. falciparum-infected mosquito MGs and

SGs. As input for DiffBind, we used the ATAC-seq nucleosome-free reads and the THSs.

Infection 1 and Infection 2 were used as biological replicates.

4.6   Relationship between gene expression and chromatin accessibility

4.6.1   Model and predictions

The relationship between the gene expression and chromatin structure has been examined in a

few model organisms (Luco et al., 2011). These previous studies showed that various epigenetic

mechanisms such as nucleosome positioning, post-transcriptional modifications (Luco et al.,

2011) or the modification of the chromatin accessibility (Li et al., 2020b, Nussinov et al., 2021),

alter or correlate with AS dynamics. In *A. gambiae*, an earlier work investigated the relationship

between chromatin accessibility and gene expression regulation at gene level (Ruiz, Ranford-

Cartwright and Gómez-Díaz, 2021), but here we focused on gene regulation occurring at the

isoform level via alternative splicing.

We made some predictions on how chromatin structure may be altered when comparing

different mosquito tissues in which AS would play a role in gene expression regulation (see

previous section), and the pattern expected considering DEMG and DUI genes. Our expectation

was to find a positive relationship between chromatin accessibility differences and the

differential expression and/or usage of the isoforms (Fig 4).

For the differentially expressed group, we expected to find an increased accessibility in the

promoter of one or more isoforms in the tissue where the expression of the isoform/s is higher

(Fig 4a). In the differentially used isoforms group, we expected to find changes in chromatin

accessibility at regulatory regions coinciding or near alternative splicing sites in the isoform

more used in each tissue, as is shown in Fig 4b. For example, for an exon skipping event we

expected more accessibility in the surroundings of the exon to be added to the transcript. In our

data the alternative splicing event more used is the ES and the majority of the MACS2 peaks

were in the isoform body. Alternatively, this association could also be evident in the promoter of

isoforms differentially used for AS events such as ES, ATTS and IR. (Li et al., 2020b).

4.6.2   Data analysis

To test the possible relationship between differential expression of isoforms and chromatin

accessibility, we discarded genes that overlap by more than 25% with other genes. We used the

Spearman's correlation test from R v4.2.0 to analyze the possible association between isoform

expression and chromatin accessibility at various regions where the AS mechanisms would

operate: The isoform promoter, which comprises 1Kbp from the TSS (end 5' UTR), and the

isoform body including all exons and introns. For those isoforms that lack a TSS annotated, we

considered the ATG as the reference point (supplementary figure 2).

We used a R script to classify the multisoform genes into three groups - high, medium, and low

- depending on the expression level of the isoforms. The threshold values for this classification

were determined by dividing the signal into three quantile groups based on their means, using

the Hmisc::cut2 R function. We then used the Spearman's correlation test to compare the

isoform expression and chromatin accessibility for each group separately. The software *ngsplot*

was used to visualize as Average profile plots the chromatin accessibility (nucleosome free

region signal) at the TSS in the different groups of isoforms classified by their expression level

(high, medium, low) (Shen et al., 2014) (Supplementary figure 3).

In the case of differentially used isoforms, we analyse the association between isoform usage

and chromatin accessibility focusing on alternative splicing sites. To do this, we first searched

for significantly differential peaks between conditions (previously obtained by DiffBind) that

coincide with *A. gambiae* known splicing sites (SS) adding a small distance to see what is

happening around these sites. We set +/- 100bp being restrictive, and subsequently we counted

how many of those peaks annotate to isoforms classified as differentially used. On those peaks,

we then we performed a motif analysis using HOMER software v4.11 (Homer Software and

Data Download, 2022). We used *findMotifGenome.pl* to determine known and *de novo* motifs

overrepresented in our sequences, and then used *annotatePeaks.pl* to link the motifs back to the

differentially used isoforms and the splicing sites. Only motifs with significance p-values less

than 1e-25 were considered.

4.7   Enhancers associated to AS in *A. gambiae*

Based on the integrative analysis of the ATAC-seq and RNA-seq data, we used the set of *A.*

gambiae enhancers characterized in a previous study (Ruiz, Ranford-Cartwright and Gómez-

Díaz, 2021), to interrogate whether these regulatory sequences could be involved in AS events.

The list included 1,402 *A. gambiae* enhancers predicted by homology with *Drosophila*

enhancers and, 2,866 *Anopheles* enhancers identified either computationally or experimentally

by different studies (Ruiz, Ranford-Cartwright and Gómez-Díaz, 2021). We selected only the

enhancers that appeared to be active in at least one of our conditions, that is, enhancers that have

a MACS2 chromatin accessibility peak and appear enriched in H3K27ac. As a result, we

obtained a list of 811 active enhancers which we overlapped with our differentially expressed

and used isoforms using BEDtools *intersect*. We also classified these enhancers as intragenic or

intergenic using BEDtools *intersect*.

Finally, we quantified which portion of the active enhancers that annotated to differentially

expressed and used isoforms, display a change in accessibility; that is, are differentially

accessible between tissues coinciding with the change in the expression of the isoform/s.

5   Funding sources

532 This work was supported by the Spanish Ministry of Science and Innovation [grant no.

533 PID2019-111109RB-I00]. BDT is funded by a doctoral training fellowship [grant no. PRE2020-

534 095076].

6   Conflict of interest

536 Authors declare no conflict of interest.

7   References

Babraham Bioinformatics - FastQC A Quality Control tool for High Throughput Sequence Data.
(2022) [Online]. Available: https://www.bioinformatics.babraham.ac.uk/projects/fastqc/
[Accessed 11 Nov 2022].

Blanchette, M., Green, R. E., MacArthur, S., Brooks, A. N., Brenner, S. E., Eisen, M. B. and
Rio, D. C. (2009) Genome-wide analysis of alternative pre-mRNA splicing and RNA
binding specificities of the Drosophila hnRNP A/B family members. Molecular cell.
[Online] Vol.33 (4), p.438. Available: /pmc/articles/PMC2674966/ [Accessed 5 May
2023].

Bush, S. J., Chen, L., Tovar-Corona, J. M. and Urrutia, A. O. (2017) Alternative splicing and the
evolution of phenotypic novelty. Philosophical Transactions of the Royal Society B:
Biological Sciences. [Online] Vol.372 (1713). Available:
https://royalsocietypublishing.org/doi/10.1098/rstb.2015.0474 [Accessed 8 Sep 2022].

Calkins, T. L., Woods-Acevedo, M. A., Hildebrandt, O. and Piermarini, P. M. (2015) The
molecular and immunochemical expression of innexins in the yellow fever mosquito,
Aedes aegypti: insights into putative life stage- and tissue-specific functions of gap
junctions. Comparative biochemistry and physiology. Part B, Biochemistry & molecular
biology. [Online] Vol.183, p.11. Available: /pmc/articles/PMC4380524/ [Accessed 28 Nov
2022].

Chen, L., Chen, K., Hong, Y., Xing, L., Zhang, J., Zhang, K. and Zhang, Z. (2022) The
landscape of isoform switches in sepsis: a multicenter cohort study. Scientific Reports
2022 12:1. [Online] Vol.12 (1), pp.1–16. Available:
https://www.nature.com/articles/s41598-022-14231-9 [Accessed 7 Sep 2022].

Chen, Y., Ling, E. and Weng, Z. (2009) Functional characterization of PGRP-LC1 of Anopheles
gambiae through deletion and RNA interference. Insect Science. [Online] Vol.16 (6),
pp.443–453. Available: https://onlinelibrary.wiley.com/doi/full/10.1111/j.1744-
7917.2009.01296.x [Accessed 1 Dec 2022].

Chodavarapu, R. K., Feng, S., Bernatavichute, Y. V., Chen, P. Y., Stroud, H., Yu, Y., Hetzel, J.
A., Kuo, F., Kim, J., Cokus, S. J., Casero, D., Bernal, M., Huijser, P., Clark, A. T., Krämer,
U., Merchant, S. S., Zhang, X., Jacobsen, S. E. and Pellegrini, M. (2010) Relationship
between nucleosome positioning and DNA methylation. Nature 2010 466:7304. [Online]
Vol.466 (7304), pp.388–392. Available: https://www.nature.com/articles/nature09147
[Accessed 14 Nov 2022].

Davuluri, R. V., Suzuki, Y., Sugano, S., Plass, C. and Huang, T. H. M. (2008) The functional
consequences of alternative promoter use in mammalian genomes. Trends in Genetics.
Vol.24 (4), pp.167–177.

Dixit, R., Sharma, A., Mourya, D. T., Kamaraju, R., Patole, M. S. and Shouche, Y. S. (2009)
Salivary gland transcriptome analysis during Plasmodium infection in malaria vector
Anopheles stephensi. International Journal of Infectious Diseases. Vol.13 (5), pp.636–646.

Djihinto, O., Saizonou, H. D. M., Djogbenou, L. S., Nolan, T. and Lee, Y. (2022) Single
nucleotide polymorphism (SNP) in the doublesex (dsx) gene splice sites and relevance for
its alternative splicing in the malaria vector Anopheles gambiae [version 1; peer review: 1
approved with reservations, 1 not approved]. [Online]. Available:
https://doi.org/10.12688/wellcomeopenres.17572.1 [Accessed 9 Sep 2022].

EZlab | Computational Evolutionary Genomics group. (2022) [Online]. Available:
https://www.ezlab.org/ [Accessed 14 Nov 2022].

Gan, Q., Chepelev, I., Wei, G., Tarayrah, L., Cui, K., Zhao, K. and Chen, X. (2010) Dynamic
regulation of alternative splicing and chromatin structure in Drosophila gonads revealed by
RNA-seq. Cell research. [Online] Vol.20 (7), p.763. Available:
/pmc/articles/PMC2919574/.

Giraldo-Calderón, G. I., Emrich, S. J., MacCallum, R. M., Maslen, G., Emrich, S., Collins, F.,
Dialynas, E., Topalis, P., Ho, N., Gesing, S., Madey, G., Collins, F. H., Lawson, D.,
Kersey, P., Allen, J., Christensen, M., Hughes, D., Koscielny, G., Langridge, N., Gallego,
E. L., Megy, K., Wilson, D., Gelbart, B., Emmert, D., Russo, S., Zhou, P., Christophides,
G., Brockman, A., Kirmitzoglou, I., MacCallum, B., Tiirikka, T., Louis, K., Dritsou, V.,
Mitraka, E., Werner-Washburn, M., Baker, P., Platero, H., Aguilar, A., Bogol, S.,
Campbell, D., Carmichael, R., Cieslak, D., Davis, G., Konopinski, N., Nabrzyski, J.,
Reinking, C., Sheehan, A., Szakonyi, S. and Wieck, R. (2015) VectorBase: an updated
bioinformatics resource for invertebrate vectors and other organisms related with human
diseases. Nucleic Acids Research. [Online] Vol.43 (D1), pp.D707–D713. Available:
https://academic.oup.com/nar/article/43/D1/D707/2437683 [Accessed 11 Nov 2022].

Glaus, P., Honkela, A. and Rattray, M. (2012) Identifying differentially expressed transcripts
from RNA-seq data with biological variation. Bioinformatics. [Online] Vol.28 (13),
pp.1721–1728. Available:

https://academic.oup.com/bioinformatics/article/28/13/1721/235585 [Accessed 25 Nov
2022].

Heinz, S., Benner, C., Spann, N., Bertolino, E., Lin, Y. C., Laslo, P., Cheng, J. X., Murre, C.,
Singh, H. and Glass, C. K. (2010) Simple combinations of lineage-determining
transcription factors prime cis-regulatory elements required for macrophage and B cell
identities. Molecular cell. [Online] Vol.38 (4), pp.576–589. Available:
https://pubmed.ncbi.nlm.nih.gov/20513432/ [Accessed 3 Sep 2023].

Homer Software and Data Download. (2022) [Online]. Available:
http://homer.ucsd.edu/homer/motif/ [Accessed 14 Nov 2022].

Jimeno-González, S., Payán-Bravo, L., Muñoz-Cabello, A. M., Guijo, M., Gutierrez, G., Prado,
F. and Reyes, J. C. (2015) Defective histone supply causes changes in RNA polymerase II
elongation rate and cotranscriptional pre-mRNA splicing. Proceedings of the National
Academy of Sciences of the United States of America. [Online] Vol.112 (48), pp.14840–
14845. Available: /pmc/articles/PMC4672771/ [Accessed 27 Apr 2023].

Katz, Y., Wang, E. T., Airoldi, E. M. and Burge, C. B. (2010) Analysis and design of RNA
sequencing experiments for identifying isoform regulation. Nature Methods 2010 7:12.
[Online] Vol.7 (12), pp.1009–1015. Available:
https://www.nature.com/articles/nmeth.1528 [Accessed 25 Nov 2022].

Kelemen, O., Convertini, P., Zhang, Z., Wen, Y., Shen, M., Falaleeva, M. and Stamm, S. (2013)
Function of alternative splicing. Gene. [Online] Vol.514 (1), p.1. Available:
/pmc/articles/PMC5632952/ [Accessed 8 Sep 2022].

Klemm, S. L., Shipony, Z. and Greenleaf, W. J. (2019) Chromatin accessibility and the
regulatory epigenome. Nature Reviews Genetics 2018 20:4. [Online] Vol.20 (4), pp.207–
220. Available: https://www.nature.com/articles/s41576-018-0089-8 [Accessed 8 Sep
2022].

Kopylova, E., Noé, L. and Touzet, H. (2012) SortMeRNA: fast and accurate filtering of
ribosomal RNAs in metatranscriptomic data. Bioinformatics. [Online] Vol.28 (24),
pp.3211–3217. Available:
https://academic.oup.com/bioinformatics/article/28/24/3211/246053 [Accessed 11 Nov
2022].

Krzywinska, E., Dennison, N. J., Lycett, G. J. and Krzywinski, J. (2016) A maleness gene in the
malaria mosquito Anopheles gambiae. Science. [Online] Vol.353 (6294), pp.67–69.
Available: https://www.science.org/doi/10.1126/science.aaf5605 [Accessed 28 Nov 2022].

Kurtz, J. and Armitage, S. A. O. (2006) Alternative adaptive immunity in invertebrates. Trends
in Immunology. Vol.27 (11), pp.493–496.

Kwon, B., Fansler, M. M., Patel, N. D., Lee, J., Ma, W. and Mayr, C. (2022) Enhancers regulate
3′ end processing activity to control expression of alternative 3′UTR isoforms. Nature
Communications. [Online] Vol.13 (1). Available: /pmc/articles/PMC9114392/ [Accessed
28 Apr 2023].

De La Fuente, L., Arzalluz-Luque, Á., Tardáguila, M., Del Risco, H., Martí, C., Tarazona, S.,
Salguero, P., Scott, R., Lerma, A., Alastrue-Agudo, A., Bonilla, P., Newman, J. R. B.,
Kosugi, S., McIntyre, L. M., Moreno-Manzano, V. and Conesa, A. (2020) TappAS: A
comprehensive computational framework for the analysis of the functional impact of
differential splicing. Genome Biology. [Online] Vol.21 (1), pp.1–32. Available:

646 https://genomebiology.biomedcentral.com/articles/10.1186/s13059-020-02028-w
647 [Accessed 7 Sep 2022].

648 Lam, B. J. and Hertel, K. J. (2002) A general role for splicing enhancers in exon definition.
649 RNA. [Online] Vol.8 (10), p.1233. Available: /pmc/articles/PMC1370333/?report=abstract
650 [Accessed 16 Jan 2023].

651 Langmead, B. and Salzberg, S. L. (2012) Fast gapped-read alignment with Bowtie 2. Nature
652 Methods 2012 9:4. [Online] Vol.9 (4), pp.357–359. Available:
653 https://www.nature.com/articles/nmeth.1923 [Accessed 3 Sep 2023].

654 Li, H., Handsaker, B., Wysoker, A., Fennell, T., Ruan, J., Homer, N., Marth, G., Abecasis, G.
655 and Durbin, R. (2009) The Sequence Alignment/Map format and SAMtools.
656 Bioinformatics (Oxford, England). [Online] Vol.25 (16), pp.2078–2079. Available:
657 https://pubmed.ncbi.nlm.nih.gov/19505943/ [Accessed 3 Sep 2023].

658 Li, Y., Liu, Y., Yang, H., Zhang, T., Naruse, K. and Tu, Q. (2020a) Dynamic transcriptional and
659 chromatin accessibility landscape of medaka embryogenesis. Genome Research. [Online]
660 Vol.30 (6), pp.924–937. Available: /pmc/articles/PMC7370878/ [Accessed 9 Aug 2022].

661 Li, Y., Liu, Y., Yang, H., Zhang, T., Naruse, K. and Tu, Q. (2020b) Dynamic transcriptional and
662 chromatin accessibility landscape of medaka embryogenesis. Genome Research. [Online]
663 Vol.30 (6), pp.924–937. Available: /pmc/articles/PMC7370878/.

664 Lin, H., Zhang, L., Luna, C., Hoa, N. T. and Zheng, L. (2007) A splice variant of PGRP-LC
665 required for expression of antimicrobial peptides in Anopheles gambiae. Insect Science.
666 [Online] Vol.14 (3), pp.185–192. Available:
667 https://onlinelibrary.wiley.com/doi/full/10.1111/j.1744-7917.2007.00142.x [Accessed 1
668 Dec 2022].

669 Lin, X., Liu, F., Meng, K., Liu, H., Zhao, Y., Chen, Y., Hu, W. and Luo, D. (2022)
670 Comprehensive Transcriptome Analysis Reveals Sex-Specific Alternative Splicing Events
671 in Zebrafish Gonads. Life. [Online] Vol.12 (9), p.1441. Available:
672 https://www.mdpi.com/2075-1729/12/9/1441/htm [Accessed 25 Nov 2022].

673 Love, M. I., Huber, W. and Anders, S. (2014) Moderated estimation of fold change and
674 dispersion for RNA-seq data with DESeq2. Genome Biology. [Online] Vol.15 (12), pp.1–
675 21. Available: https://genomebiology.biomedcentral.com/articles/10.1186/s13059-014-
676 0550-8 [Accessed 7 Sep 2022].

677 Luco, R. F., Allo, M., Schor, I. E., Kornblihtt, A. R. and Misteli, T. (2011) Epigenetics in
678 Alternative Pre-mRNA Splicing. Cell. Vol.144 (1), pp.16–26.

679 Malko, D. B., Makeev, V. J., Mironov, A. A. and Gelfand, M. S. (2006) Evolution of exon–
680 intron structure and alternative splicing in fruit flies and malarial mosquito genomes.
681 Genome Research. [Online] Vol.16 (4), pp.505–509. Available:
682 https://genome.cshlp.org/content/16/4/505.full [Accessed 28 Nov 2022].

683 Marikovsky, M., Ziv, V., Nevo, N., Harris-Cerruti, C. and Mahler, O. (2003) Cu/Zn Superoxide
684 Dismutase Plays Important Role in Immune Response. The Journal of Immunology.
685 [Online] Vol.170 (6), pp.2993–3001. Available:
686 https://journals.aai.org/jimmunol/article/170/6/2993/71379/Cu-Zn-Superoxide-Dismutase-
687 Plays-Important-Role-in [Accessed 19 Jan 2023].

688 Martinez, N. M. and Lynch, K. W. (2013) Control of alternative splicing in immune responses:
689 many regulators, many predictions, much still to learn. Immunological Reviews. [Online]

Vol.253 (1), pp.216–236. Available:
https://onlinelibrary.wiley.com/doi/full/10.1111/imr.12047 [Accessed 9 Aug 2022].

Matlin, A. J., Clark, F. and Smith, C. W. J. (2005) Understanding alternative splicing: towards a cellular code. Nature Reviews Molecular Cell Biology 2005 6:5. [Online] Vol.6 (5), pp.386–398. Available: https://www.nature.com/articles/nrm1645 [Accessed 16 Jan 2023].

Meister, S., Agianian, B., Turlure, F., Relógio, A., Morlais, I., Kafatos, F. C. and Christophides, G. K. (2009) Anopheles gambiae PGRPLC-Mediated Defense against Bacteria Modulates Infections with Malaria Parasites. PLoS Pathogens. [Online] Vol.5 (8). Available: /pmc/articles/PMC2715215/ [Accessed 9 Sep 2022].

Meister, S., Kanzok, S. M., Zheng, X. L., Luna, C., Li, T. R., Hoa, N. T., Clayton, J. R., White, K. P., Kafatos, F. C., Christophides, G. K. and Zheng, L. (2005) Immune signaling pathways regulating bacterial and malaria parasite infection of the mosquito Anopheles gambiae. Proceedings of the National Academy of Sciences of the United States of America. [Online] Vol.102 (32), pp.11420–11425. Available: https://www.pnas.org/doi/abs/10.1073/pnas.0504950102 [Accessed 9 Sep 2022].

Merino, G. A., Conesa, A. and Fernández, E. A. (2019) A benchmarking of workflows for detecting differential splicing and differential expression at isoform level in human RNA-seq studies. Briefings in Bioinformatics. [Online] Vol.20 (2), pp.471–481. Available: https://academic.oup.com/bib/article/20/2/471/4524048 [Accessed 7 Sep 2022].

Molina-Cruz, A., DeJong, R. J., Charles, B., Gupta, L., Kumar, S., Jaramillo-Gutierrez, G. and Barillas-Mury, C. (2008) Reactive oxygen species modulate Anopheles gambiae immunity against bacteria and Plasmodium. Journal of Biological Chemistry. [Online] Vol.283 (6), pp.3217–3223. Available: http://www.jbc.org/article/S0021925820697848/fulltext [Accessed 19 Jan 2023].

Montes, M., Becerra, S., Sánchez-Álvarez, M. and Suñé, C. (2012) Functional coupling of transcription and splicing. Gene. Vol.501 (2), pp.104–117.

Naftelberg, S., Schor, I. E., Ast, G. and Kornblihtt, A. R. (n.d.) Regulation of Alternative Splicing Through Coupling with Transcription and Chromatin Structure. [Online]. Available: www.annualreviews.org [Accessed 5 Dec 2022].

Nilsen, T. W. and Graveley, B. R. (2010) Expansion of the eukaryotic proteome by alternative splicing. Nature 2010 463:7280. [Online] Vol.463 (7280), pp.457–463. Available: https://www.nature.com/articles/nature08909 [Accessed 8 Sep 2022].

Nussinov, R., Zhang, M., Maloney, R. and Jang, H. (2021) Ras isoform-specific expression, chromatin accessibility, and signaling. Biophysical Reviews 2021 13:4. [Online] Vol.13 (4), pp.489–505. Available: https://link.springer.com/article/10.1007/s12551-021-00817-6 [Accessed 8 Sep 2022].

Onodera, C. S., Underwood, J. G., Katzman, S., Jacobs, F., Greenberg, D., Salama, S. R. and Haussler, D. (2012) Gene Isoform Specificity through Enhancer-Associated Antisense Transcription. PLoS ONE. [Online] Vol.7 (8), p.43511. Available: /pmc/articles/PMC3427357/ [Accessed 28 Apr 2023].

Ou, J., Liu, H., Yu, J., Kelliher, M. A., Castilla, L. H., Lawson, N. D. and Zhu, L. J. (2018) ATACseqQC: A Bioconductor package for post-alignment quality assessment of ATAC-seq data. BMC Genomics. [Online] Vol.19 (1), pp.1–13. Available:

https://bmcgenomics.biomedcentral.com/articles/10.1186/s12864-018-4559-3 [Accessed 3
Sep 2023].

Pai, A. A. and Luca, F. (2019) Environmental influences on RNA processing: biochemical,
molecular and genetic regulators of cellular response. Wiley interdisciplinary reviews.
RNA. [Online] Vol.10 (1), p.e1503. Available: /pmc/articles/PMC6294667/ [Accessed 28
Apr 2023].

Patro, R., Duggal, G., Love, M. I., Irizarry, R. A. and Kingsford, C. (2017) Salmon provides fast
and bias-aware quantification of transcript expression. Nature Methods 2017 14:4.
[Online] Vol.14 (4), pp.417–419. Available: https://www.nature.com/articles/nmeth.4197
[Accessed 11 Nov 2022].

Petrova, V., Song, R., Consortium, D., Nordström, K. J. V, Walter, J., Wong, J. J. L., Armstrong,
N. J., Rasko, J. E. J. and Schmitz, U. (2022) Increased chromatin accessibility facilitates
intron retention in specific cell differentiation states. Nucleic Acids Research. [Online]
Vol.50 (20), p.11563. Available: /pmc/articles/PMC9723627/ [Accessed 27 Apr 2023].

Petrova, V., Song, R., Consortium, D., NordströmNordstr, K. J., Walter, orn, J-L Wong, J.,
Armstrong, N. J., Rasko, J. E. and Schmitz, U. (2022) Increased chromatin accessibility
facilitates intron retention in specific cell differentiation states. Nucleic Acids Research.
[Online] (1). Available: https://academic.oup.com/nar/advance-
article/doi/10.1093/nar/gkac994/6819938 [Accessed 5 Dec 2022].

Quinlan, A. R. and Hall, I. M. (2010) BEDTools: a flexible suite of utilities for comparing
genomic features. Bioinformatics (Oxford, England). [Online] Vol.26 (6), pp.841–842.
Available: https://pubmed.ncbi.nlm.nih.gov/20110278/ [Accessed 3 Sep 2023].

Ragle, J. M., Katzman, S., Akers, T. F., Barberan-Soler, S. and Zahler, A. M. (2015)
Coordinated tissue-specific regulation of adjacent alternative 3′ splice sites in C. elegans.
Genome Research. [Online] Vol.25 (7), p.982. Available: /pmc/articles/PMC4484395/
[Accessed 29 Nov 2022].

Ruiz, J. L., Ranford-Cartwright, L. C. and Gómez-Díaz, E. (2021) The regulatory genome of the
malaria vector Anopheles gambiae: integrating chromatin accessibility and gene
expression. NAR Genomics and Bioinformatics. [Online] Vol.3 (1). Available:
https://academic.oup.com/nargab/article/3/1/lqaa113/6104947 [Accessed 11 Nov 2022].

Ruiz, J. L., Yerbanga, R. S., Lefèvre, T., Ouedraogo, J. B., Corces, V. G. and Gómez-Díaz, E.
(2019) Chromatin changes in Anopheles gambiae induced by Plasmodium falciparum
infection 06 Biological Sciences 0604 Genetics 11 Medical and Health Sciences 1108
Medical Microbiology. Epigenetics and Chromatin. [Online] Vol.12 (1), pp.1–18.
Available: https://link.springer.com/articles/10.1186/s13072-018-0250-9 [Accessed 11
Nov 2022].

Saldi, T., Cortazar, M. A., Sheridan, R. M. and Bentley, D. L. (2016) Coupling of RNA
polymerase II transcription elongation with pre-mRNA splicing. Journal of molecular
biology. [Online] Vol.428 (12), p.2623. Available: /pmc/articles/PMC4893998/ [Accessed
27 Apr 2023].

Salvemini, M., D'Amato, R., Petrella, V., Aceto, S., Nimmo, D., Neira, M., Alphey, L., Polito,
L. C. and Saccone, G. (2013) The Orthologue of the Fruitfly Sex Behaviour Gene Fruitless
in the Mosquito Aedes aegypti: Evolution of Genomic Organisation and Alternative
Splicing. PLoS ONE. [Online] Vol.8 (2), p.48554. Available: /pmc/articles/PMC3572092/
[Accessed 28 Nov 2022].

Salvemini, M., Mauro, U., Lombardo, F., Milano, A., Zazzaro, V., Arcà, B., Polito, L. C. and
Saccone, G. (2011) Genomic organization and splicing evolution of the doublesex gene, a
Drosophila regulator of sexual differentiation, in the dengue and yellow fever mosquito
Aedes aegypti. BMC Evolutionary Biology. [Online] Vol.11 (1), p.41. Available:
/pmc/articles/PMC3045327/ [Accessed 28 Nov 2022].

Schor, I. E., Gómez Acuña, L. I. and Kornblihtt, A. R. (2013) Coupling between transcription
and alternative splicing. Cancer Treatment and Research. [Online] pp.1–24. Available:
https://link.springer.com/chapter/10.1007/978-3-642-31659-3_1 [Accessed 27 Apr 2023].

Schuettengruber, B., Ganapathi, M., Leblanc, B., Portoso, M., Jaschek, R., Tolhuis, B., Van
Lohuizen, M., Tanay, A. and Cavalli, G. (2009) Functional Anatomy of Polycomb and
Trithorax Chromatin Landscapes in Drosophila Embryos. PLoS Biology. [Online] Vol.7
(1), p.1000013. Available: /pmc/articles/PMC2621266 [Accessed 5 Dec 2022].

Shen, L., Shao, N., Liu, X. and Nestler, E. (2014) Ngs.plot: Quick mining and visualization of
next-generation sequencing data by integrating genomic databases. BMC Genomics.
[Online] Vol.15 (1), pp.1–14. Available:
https://bmcgenomics.biomedcentral.com/articles/10.1186/1471-2164-15-284 [Accessed 1
Dec 2022].

Shenasa, H. and Hertel, K. J. (2019) Combinatorial regulation of alternative splicing.
Biochimica et Biophysica Acta (BBA) - Gene Regulatory Mechanisms. Vol.1862 (11–12),
p.194392.

Shiau, C. K., Huang, J. H., Liu, Y. T. and Tsai, H. K. (2021) Genome-wide identification of
associations between enhancer and alternative splicing in human and mouse. BMC
Genomics. [Online] Vol.22 (Suppl 5). Available: /pmc/articles/PMC9082955/ [Accessed
28 Apr 2023].

Simon, J. M., Hacker, K. E., Singh, D., Brannon, A. R., Parker, J. S., Weiser, M., Ho, T. H.,
Kuan, P. F., Jonasch, E., Furey, T. S., Prins, J. F., Lieb, J. D., Rathmell, W. K. and Davis, I.
J. (2014) Variation in chromatin accessibility in human kidney cancer links H3K36
methyltransferase loss with widespread RNA processing defects. Genome Research.
[Online] Vol.24 (2), pp.241–250. Available: https://genome.cshlp.org/content/24/2/241.full
[Accessed 5 Dec 2022].

Sinkins, S. P. (2016) Yob makes mosquitoes male. Science. [Online] Vol.353 (6294), pp.33–34.
Available: https://www.science.org/doi/10.1126/science.aag1012 [Accessed 28 Nov 2022].

Song, L., Pan, Z., Chen, L., Dai, Y., Wan, J., Ye, H., Nguyen, H. T., Zhang, G. and Chen, H.
(2020) Analysis of Whole Transcriptome RNA-seq Data Reveals Many Alternative
Splicing Events in Soybean Roots under Drought Stress Conditions. Genes 2020, Vol. 11,
Page 1520. [Online] Vol.11 (12), p.1520. Available: https://www.mdpi.com/2073-
4425/11/12/1520/htm [Accessed 6 Sep 2022].

Sreenivasamurthy, S. K., Dey, G., Ramu, M., Kumar, M., Gupta, M. K., Mohanty, A. K.,
Harsha, H., Sharma, P., Kumar, N., Pandey, A., Kumar, A. and Prasad, T. K. (2013) A
compendium of molecules involved in vector-pathogen interactions pertaining to malaria.
Malaria Journal. [Online] Vol.12 (1), p.216. Available: /pmc/articles/PMC3734095/
[Accessed 27 Apr 2023].

Sreenivasamurthy, S. K., Madugundu, A. K., Patil, A. H., Dey, G., Mohanty, A. K., Kumar, M.,
Patel, K., Wang, C., Kumar, A., Pandey, A. and Prasad, T. S. K. (2017) Mosquito-Borne
Diseases and Omics: Tissue-Restricted Expression and Alternative Splicing Revealed by

Transcriptome Profiling of Anopheles stephensi. OMICS A Journal of Integrative Biology.
Vol.21 (8), pp.488–497.

Stark R. and Brown G. (2011) DiffBind: differential binding analysis of ChIP-Seq peak data. R
Package Version. Vol.100, pp.4–3.

Telonis-Scott, M., Kopp, A., Wayne, M. L., Nuzhdin, S. V. and McIntyre, L. M. (2009) Sex-
Specific Splicing in Drosophila: Widespread Occurrence, Tissue Specificity and
Evolutionary Conservation. Genetics. [Online] Vol.181 (2), p.421. Available:
/pmc/articles/PMC2644937/ [Accessed 29 Nov 2022].

Tsujimoto, H., Liu, K., Linser, P. J., Agre, P. and Rasgon, J. L. (2013) Organ-Specific Splice
Variants of Aquaporin Water Channel AgAQP1 in the Malaria Vector Anopheles gambiae.
PLoS ONE. [Online] Vol.8 (9). Available: /pmc/articles/PMC3774814/ [Accessed 28 Nov
2022].

Ule, J. and Blencowe, B. J. (2019) Alternative Splicing Regulatory Networks: Functions,
Mechanisms, and Evolution. Molecular Cell. [Online] Vol.76 (2), pp.329–345. Available:
http://www.cell.com/article/S1097276519307026/fulltext [Accessed 5 Sep 2022].

Ullah, F., Hamilton, M., Reddy, A. S. N. and Ben-Hur, A. (2018) Exploring the relationship
between intron retention and chromatin accessibility in plants. BMC Genomics. [Online]
Vol.19 (1), p.21. Available: /pmc/articles/PMC5756433/ [Accessed 27 Apr 2023].

Urban, J. A., Urban, J. M., Kuzu, G. and Larschan, E. N. (2017) The Drosophila CLAMP
protein associates with diverse proteins on chromatin. PLoS ONE. [Online] Vol.12 (12).
Available: /pmc/articles/PMC5744976/ [Accessed 5 Dec 2022].

Venables, J. P., Tazi, J. and Juge, F. (2012) Regulated functional alternative splicing in
Drosophila. Nucleic Acids Research. [Online] Vol.40 (1), pp.1–10. Available:
https://academic.oup.com/nar/article/40/1/1/1272632 [Accessed 9 Sep 2022].

Verta, J. P. and Jacobs, A. (2022) The role of alternative splicing in adaptation and evolution.
Trends in Ecology & Evolution. Vol.37 (4), pp.299–308.

Vitting-Seerup, K. and Sandelin, A. (2017) The landscape of isoform switches in human
cancers. Molecular Cancer Research. [Online] Vol.15 (9), pp.1206–1220. Available:
https://aacrjournals.org/mcr/article/15/9/1206/268100/The-Landscape-of-Isoform-
Switches-in-Human [Accessed 7 Sep 2022].

Vitting-Seerup, K., Sandelin, A. and Berger, B. (2019) IsoformSwitchAnalyzeR: analysis of
changes in genome-wide patterns of alternative splicing and its functional consequences.
Bioinformatics. [Online] Vol.35 (21), pp.4469–4471. Available:
https://academic.oup.com/bioinformatics/article/35/21/4469/5466456 [Accessed 7 Sep
2022].

Wang, E. T., Sandberg, R., Luo, S., Khrebtukova, I., Zhang, L., Mayr, C., Kingsmore, S. F.,
Schroth, G. P. and Burge, C. B. (2008) Alternative isoform regulation in human tissue
transcriptomes. Nature 2008 456:7221. [Online] Vol.456 (7221), pp.470–476. Available:
https://www.nature.com/articles/nature07509 [Accessed 29 Nov 2022].

Wang, G. S. and Cooper, T. A. (2007) Splicing in disease: disruption of the splicing code and the
decoding machinery. Nature Reviews Genetics 2007 8:10. [Online] Vol.8 (10), pp.749–
761. Available: https://www.nature.com/articles/nrg2164 [Accessed 25 Nov 2022].

Wright, C. J., Smith, C. W. J. and Jiggins, C. D. (2022) Alternative splicing as a source of
phenotypic diversity. <u>Nature Reviews Genetics 2022</u>. [Online] pp.1–14. Available:
https://www.nature.com/articles/s41576-022-00514-4 [Accessed 5 Sep 2022].

Zahler, A. M. (2005) Alternative splicing in C. elegans. <u>WormBook□: the online review of C.</u>
<u>elegans biology</u>. [Online] pp.1–13. Available:
https://www.ncbi.nlm.nih.gov/books/NBK19730/ [Accessed 25 Nov 2022].

Zhang, C., Zhang, B., Vincent, M. S. and Zhao, S. (2016) Bioinformatics Tools for RNA-seq
Gene and Isoform Quantification. [Online]. Available: www.gencodegenes.org [Accessed
11 Nov 2022].

Zhang, Y., Liu, T., Meyer, C. A., Eeckhoute, J., Johnson, D. S., Bernstein, B. E., Nussbaum, C.,
Myers, R. M., Brown, M., Li, W. and Shirley, X. S. (2008a) Model-based Analysis of
ChIP-Seq (MACS). <u>Genome Biology</u>. [Online] Vol.9 (9), p.R137. Available:
/pmc/articles/PMC2592715/ [Accessed 13 Jun 2023].

Zhang, Y., Liu, T., Meyer, C. A., Eeckhoute, J., Johnson, D. S., Bernstein, B. E., Nussbaum, C.,
Myers, R. M., Brown, M., Li, W. and Shirley, X. S. (2008b) Model-based analysis of
ChIP-Seq (MACS). <u>Genome biology</u>. [Online] Vol.9 (9). Available:
https://pubmed.ncbi.nlm.nih.gov/18798982/ [Accessed 3 Sep 2023].

Zhao, S. (2019) Alternative splicing, RNA-seq and drug discovery. <u>Drug Discovery Today</u>.
Vol.24 (6), pp.1258–1267.

Tables

Motif	Name	P-value	Flybase ID	Flybase name	Nº of DUI isoforms with the motif
CACCWATT	sna/MA0086.2	1E-34	FBgn0003448	snail	22
CTTGCAAGGT	byn/dmmpmm	1E-31	FBgn0011723	brachyenteron	21
ACARAGAAAT	sd/dmmpmm	1E-28	FBgn0003345	scalloped	26
ACACAAAG	cad/dmmpmm	1E-27	FBgn0000251	caudal	26
GAGTCTTTAT	br-Z4/dmmpmm	1E-27	FBgn0283451	broad	20
GAAAACAGTT	grh/MA1457.1	1E-27	FBgn0259211	grainy head	21
AAACAGTAAAAG	br-Z2/dmmpmm	1E-27	FBgn0283451	broad	20
CAAGAAGA	Aef1/dmmpmm	1E-27	FBgn0005694	Adult enhancer factor 1	25
GAMCGAGH	Clamp/MA1700.1	1E-27	FBgn0032979	Chromatin-linked adaptor for MSL proteins	24
TTATAAYT	br-Z1/dmmpmm	1E-27	FBgn0283451	broad	22

900 **Table 1.** Ten first "de novo" motifs identified by Homer that appear overrepresented in the splicing sites of differentially expressed and used isoforms that

901 display differential accessibility between tissues. Additional information in Supplementary table S6.

Figure legends

Fig 1. Patterns of differentially expressed isoforms belonging to multisoform genes (DEMG)
and differentially used isoforms between tissues. **a**, Diagram illustrating possible DEMG and
DUI patterns expected in our data. The gene can be differentially expressed, and the isoform
differentially expressed but not differentially used, only one more of the isoforms differentially
used and the isoforms can be differentially expressed and used at the same time; **b**, Volcano plot
showing total usage change (i.e., expression redistribution between isoforms) versus log-
transformed values of isoform expression fold change between tissues. Isoforms that are DUI
and DEMG are shown in red whereas black dots correspond to differentially expressed
isoforms. Labels identify isoforms corresponding to *A. gambiae* genes with immune functions;
c, IGV representation of the AGAP005203 (PGRPLC) gene. Red lines highlight differential
expression and accessibility at RA and RE differential exons. The red box on the left points to a
higher expression of the RA isoform in midguts, while the box on the right points to a higher
expression in salivary glands.

Fig 2. Differential expression of the isoforms correlates with changes in chromatin accessibility
between tissues in the isoform body. Heatmap showing ATAC-seq nucleosome-free enrichment
at promoters or in the body of DEMG isoforms and their expression levels. Genes are ordered
by normalized RNA-seq isoform expression (RPKM). Changes tend to occur in the same
direction, and there is a positive and significant correlation between the magnitude of changes in
accessibility and expression at the isoform body in both tissues.

Fig 3. Enhancer function and alternative splicing of the gene AGAP005234. **a**, Scheme

summarizing gene and isoform expression between tissues; **b**, IGV screenshot showing the

enhancer annotated to the AGAP005234 gene which is differentially accessible between tissues

coinciding with the differential expression and usage of their isoforms. RB isoform is

differentially expressed and used in salivary glands and the enhancer is more accessible in this

tissue. All tracks are shown at equal scale.

Fig 4. Model for the relationship between isoform expression, isoform usage and chromatin

accessibility. The ATAC-seq peaks are shown in red. **a**, For the DEMG group the expectation is

that the accessibility of the promoter will increase in the tissue where the expression of the

isoform is higher i.e, The RA isoform in midguts display a higher ATAC-seq peak. For the RB

isoform, the expression is higher in salivary glands the accessibility peak is higher in this tissue;

b, For the DUI group, we expect to find chromatin accessibility changes near the splicing sites

coinciding with the different usage of the isoform/s. The ATAC-seq peaks are around the

differentially exons and are higher in the tissues where the isoforms are more used. The RA

isoform is less expressed in midguts and becomes more expressed in salivary glands, displaying

a higher ATAC-seq peak.

a

DEG + DEMG + no DUI

No DEG + no DEMG + DUI

DEG + DEMG + DUI

b

c

Midguts

Salivary glands

RNA and ATAC mean (log2)

2.8

−2.8

Expression

Accessibility in the
isoform body

Accessibility in the
isoform promoter

Expression

Accessibility in the
isoform body

Accessibility in the
isoform promoter

a

No DEG

Gene Expresión levels

615,57 (MG) 595 (SG)

DEMG + DUI

Isoform Expresión levels

158,29 RA
21,41 RB
87,48
79,74

MG   SG

b

Track	
RNA-seq MG	[0-200]
ATAC-seq MG	[0-200]
HCS MG	
RNA-seq SG	[0-200]
ATAC-seq SG	[0-200]
HCS SG	
Diffbind Peaks	
Promoters	AGAP005234 promoter
Enhancers	
GFF *A. gambiae*	AGAP005234 / AGAP005234-RA / AGAP005234-RB