Title

Multiple axes of visual system diversity in Ithomiini, an ecologically diverse tribe of mimetic

butterflies

Running title

Visual system diversity in Ithomiini

Authors and affiliations

J. Benito Wainwright[1], Corin Schofield[1], Max Conway[1], Daniel Phillips[1], Elizabeth Martin-

Silverstone[2], Emelie A. Brodrick[3], Francesco Cicconardi[1], Martin J. How[1], Nicholas W.

Roberts[1] and Stephen H. Montgomery[1]

[1]School of Biological Sciences, University of Bristol, 24 Tyndall Avenue, Bristol, BS8 1TQ,

UK

[2]Bristol Palaeobiology Group, School of Earth Sciences, University of Bristol, 24 Tyndall

Avenue, Bristol, BS8 1TQ

[3]Living Systems Institute, University of Exeter, Stocker Road, Exeter, EX4 4QD

Corresponding authors:

J. B. Wainwright: jw14637@bristol.ac.uk

S. H. Montgomery: s.montgomery@bristol.ac.uk

Key words

Allometric scaling, apposition compound eye, Lepidoptera, neuroecology, optic lobe, visual

ecology

Summary statement

Physiological, anatomical, and molecular evidence of extensive visual system variation within a diverse butterfly radiation, hinting at the lability of visual systems to evolve specialisations to distinct visual environments.

Abstract

The striking structural variation seen in arthropod visual systems can be explained by the overall quantity and spatio-temporal structure of light within habitats coupled with developmental and physiological constraints. However, little is currently known about how fine-scale variation in visual structures arise across shorter evolutionary and ecological scales. In this study, we characterise patterns of interspecific (between species), intraspecific (between sexes) and intraindividual (between eye regions) variation in the visual system of four ithomiine butterfly species. These species are part of a diverse 26-Myr-old Neotropical radiation where changes in mimetic colouration are associated with fine-scale shifts in ecology, such as microhabitat preference. By using a combination of selection analyses on visual opsin sequences, in-vivo ophthalmoscopy, micro-computed tomography (micro-CT), immunohistochemistry, confocal microscopy, and neural tracing, we quantify and describe physiological, anatomical, and molecular traits involved in visual processing. Using these data, we provide evidence of substantial variation within the visual systems of Ithomiini, including: i) relaxed selection on visual opsins, perhaps mediated by habitat preference, ii) interspecific shifts in visual system physiology and anatomy, and iii) extensive sexual dimorphism, including the complete absence of a butterfly-specific optic neuropil in the males of some species. We conclude that considerable visual system variation can exist within diverse insect radiations, hinting at the evolutionary lability of these systems to rapidly develop specialisations to distinct visual ecologies, with selection acting at both the perceptual, processing, and molecular level.

Introduction

The diversity of animal visual specialisations reflects their central role in key behaviours such as foraging, navigation, communication, and predator avoidance (Cronin *et al.* 2014; Endler *et al.* 2005). As the maintenance of complex visual adaptations incurs significant energetic

costs (Land & Nilsson, 2012; Laughlin *et al.* 1998; Moran *et al.* 2015; Niven *et al.* 2007;

Niven & Laughlin, 2008) divergence in specialization at any functional, physiological,

anatomical or molecular level is likely the result of ecological selection pressures (e.g.

Gonzalez-Bellido *et al.* 2011; Hofmann *et al.* 2009; Huber *et al.* 1997; Scales & Butler, 2016;

Sugawara *et al.* 2005; Zhao *et al.* 2009). However, such specializations are also limited by a

number of trade-offs including evolutionary history, body size and other allometric and

physiological constraints (Land, 1997; Land & Nilsson, 2012; Warrant & McIntyre, 1993).

Understanding how an animal's visual ecology interacts with these constraints to shape

variation in visual systems is a central question to sensory and neuroecology (Cronin *et al.*

2014; Lythgoe, 1979; Striedter, 2005).

Visual system diversity is greatest within the Arthropoda (Osorio *et al.* 1995; Stansbury &

Moczek, 2013; Yilmaz *et al.* 2022). This diversity is perhaps best reflected by their colour

vision which is acquired via the tuning of visual pigments found within photoreceptor cells.

These pigments consist of a retinal-based chromophore attached to an opsin protein, where

the sequence of critical amino acid residues within the chromophore binding pocket of the

opsin can shift the wavelength sensitivity which varies between species with different visual

ecologies (Fain *et al.* 2010; Feuda *et al.* 2016; Frentiu *et al.* 2007; Terakita, 2005). In

apposition compound eyes, each facet typically projects light along a single rhabdom within

an ommatidium, separated from adjacent ommatidia by light-absorbing pigments (Land,

1989). However, anatomical, and physiological parameters such as total eye size, shape, lens

diameter, acuity, sensitivity, number of ommatidia, and pupillary response vary, even

between closely related species (Bartholomée *et al.* 2023; Greiner, 2006; Greiner *et al.*

2004a; Land, 1989; Narendra *et al.* 2013; Scales & Butler, 2016; Somanathan *et al.* 2009;

Warrant, 2001). For example, shifts to a nocturnal lifestyle have led to larger ommatidial

facets and rhabdom diameters in the apposition eye morphology of the sweat bee, *Megalopta*

genalis, compared to its close relatives (Greiner *et al.* 2004a; Warrant, 2017). Microhabitat

partitioning within communities of Hawaiian damselflies and hemiboreal bumblebees has

also resulted in rapid shifts in eye sensitivity at a much finer ecological scale (Bartholomée *et*

al. 2023; Scales & Butler, 2016). Spatial variation in light abundance and spectral

composition within habitats has also promoted compound eye regionalisation whereby

different eye regions are fine-tuned to optimally receive ecologically relevant stimuli (e.g.

Arikawa *et al.* 2009; Lehrer, 1998; Labhart & Meyer, 1999; Meyer & Labhart, 1992; Nilsson

& Smolka, 2021; Nilsson *et al.* 2022; Stavenga, 1992; Stavenga *et al.* 2001; White *et al.*

2003; Zufall *et al.* 1989). For example, the dorsal rim area (DRA) of many insects is

specialised for polarization vision, useful for orientation and navigation (Dacke *et al.* 2002;

Mappes & Homberg, 2004; Reppert *et al.* 2004; Sauman *et al.* 2005; Stalleicken *et al.* 2006).

The fine structure and function of the primary insect visual neuropils can also differ

(Sinakevitch *et al.* 2003). While the general structure of optic lobes is largely conserved,

typically subdivided into four main synapse-dense regions (the lamina, medulla, lobula and

the lobula plate) (Fig. 1a; Strausfeld & Nässel, 1980), the way they have evolved to process

different forms of visual information again reflects their visual environment and evolutionary

history. For example, neural adaptations for spatial and temporal summation within the

lamina are repeatedly associated with shifts towards nocturnal diel patterns (Greiner *et al.*

2004b; Warrant *et al.* 2004, Stöckl *et al.* 2016b; 2020).

Understanding how visual systems evolve in response to novel habitats requires comparative

study systems of species with diverse and well documented ecologies. Lepidopterans are one

such group, occupying a wide range of light environments, relying heavily on the use of

visual cues and signals for a variety of behavioural tasks, and exhibiting considerable levels

of investment in specialized visual system physiology and anatomy (Bergman *et al.* 2021;

Briscoe & Chittka, 2001; Couto *et al.* 2020; Nilsson *et al.* 1988; Stavenga & Arikawa, 2006;

St☐ckl *et al.* 2016a,b). Like other arthropods, lepidopteran species that occupy high-light

intensity environments tend to have larger optic lobe neuropils than species found in low-

light habitats, or nocturnal conditions (Couto *et al.* 2020; Heinze & Reppert, 2012;

Montgomery & Ott, 2015; Montgomery & Merrill, 2017; Wainwright & Montgomery, 2022).

The opposite appears true for relative eye size, where nocturnal species have evolved larger

facets and wider and longer rhabdoms to maximise light collection and absorption

(Frederiksen & Warrant, 2008; Yack *et al.* 2007). However, in butterflies there is also some

evidence that visual systems vary across more subtle ecological differences, and readily adapt

over short evolutionary scales. For example, a comparison of two *Boloria* butterfly species

identified larger eyes and facets in the frontal eye region in *B. aquilonaris* whose habitat is

naturally fragmented relative to *B. eunomia*, suggesting that differences in eye morphology

can evolve remarkably rapidly to ecological challenges (Turlure *et al.* 2016). Consistent

patterns of neural divergence in response to ecological preference shifts have also been

observed between parapatric *Heliconius* butterfly species, which also display eye structural

variation, separated across continuous environmental gradients within tropical forests

(Hebberecht *et al.* 2023; Montgomery & Merrill, 2017; Montgomery *et al.* 2021; Seymoure *et al.* 2015).

To better explore how visual systems evolve in response to subtle environmental changes we present here a detailed study of interspecific and intraspecific differences in both eye and brain structure, across representatives of a diverse 26-Myr-old tribe of Neotropical butterflies, the Ithomiini (Nymphalidae: Danainae) (Chazot *et al.* 2019). Historically, ithomiines have been primarily studied for their Müllerian mimicry rings (also referred to as mimicry complexes) where multiple species have evolved convergent wing colours, patterns, and morphologies to amplify their aposematic signal to predators within sympatric communities (Bates, 1862; Beccaloni, 1997; Elias & Joron, 2015; Hill, 2021; Müller, 1879). Critically, for mimicry to be effective, co-mimics must signal to the same predators, whose distribution and abundance varies across forest habitats (Gompert *et al.* 2011; Willmott *et al.* 2017). As a result, data on forest structure suggest that mimicry rings are segregated across microhabitats, which likely expose closely related species from different mimicry rings to divergent visual environments (Beccaloni, 1997; Elias *et al.* 2008; Hill, 2010; Wainwright & Montgomery, 2022; Willmott *et al.* 2017). Typically, colourful ithomiine mimics are found in more open forest compared with cryptic, transparent-winged ithomiines which reside in shaded inner-forest (Elias *et al.* 2008; Pliske, 1975), and these habitat shifts have been linked to variation in investment in sensory neuropils (Montgomery & Ott 2015; Wainwright & Montgomery, 2022). In addition, while ithomiines are generally not sexually dichromatic, evidence of dimorphism in olfactory and visual processing centres might hint at the possibility of additional sex-specific patterns of visual investment (Montgomery & Ott 2015; Morris *et al.* 2021; Pliske, 1975; Wainwright & Montgomery, 2022).

Here, we focus on representatives of four subtribes within the ithomiine radiation: *Greta morgane* (subtribe: Godyridina), *Mechanitis polymnia* (subtribe: Mechanitina), *Methona confusa* (subtribe: Methonina) and *Tithorea harmonia* (subtribe: Tithoreina), hereafter referred to by their genus name for brevity (Fig. 1b; Chazot *et al.* 2019; de Silva *et al.* 2016). *Mechanitis, Methona* and *Tithorea* have converged on an open forest microhabitat, flying at high elevations (Elias *et al.* 2008), whereas the transparent *Greta* flies lower down in shaded forest understory (Fig. 1b; Elias & Joron, 2015; Willmott & Mallet, 2004). By combining results from selection analyses on opsin sequences from recent genomic data, ophthalmoscopy, micro-CT, immunohistochemistry, and neural tracing, we describe and

quantify variation for several aspects of the ithomiine visual system. Our data reveal multiple

examples of anatomical, physiological, and molecular variation within and between these

four ithomiine species, highlighting potential adaptations for optimising visual perception and

processing during microhabitat niche partitioning.

Methods

1. *Molecular analyses of visual opsins*

1. i. Opsin gene alignment and data mining

Opsin sequences for *Danaus plexippus* (Nymphalidae: Danainae), the most closely related

species to the Ithomiini for which all three visual opsins (ultraviolet, blue and long-

wavelength) are publicly available on GenBank, were used as query sequences. These

sequences were utilised for finding homologous opsins within recently assembled genomes of

each of four ithomiine species (Cicconardi *et al.* in prep) using Exonerate v2.2.0 with '-model

protein2genome -percent 50' options (Slater & Birney, 2005; Zhan & Reppert, 2013). To

increase the power of our comparative analyses, we used GenBank deposited ultraviolet

(UV), blue (B) and long-wavelength (LW) opsin sequences from 13 additional nymphalid

butterfly species with well-established ecologies (all sequences used in our analyses,

including accession IDs, can be found in Table S1).

To confirm functional opsin number for each ithomiine species, the transmembrane structure

of each opsin sequence was predicted using Phobius, implemented through Protter (Kall *et al.*

2007; Omasits *et al.* 2014), a webserver-based tool for making protein structural predictions.

Any opsin sequences predicted to have six or seven transmembrane domains were considered

putative functional opsins. Rh7 opsins were also surveyed, however this gene family was not

included in downstream analyses due to a poor abundance of other available nymphalid Rh7

sequences, as previously noted by Sondhi *et al.* (2021). The 3D protein structure for the

Greta and *Tithorea* UV opsins were also modelled using a homology-based approach in

Swiss-Model (Waterhouse *et al.* 2018). The jumping spider Rhodopsin-1 (9i9k.1.A) was used

as a template because this had the highest identity score and coverage (*Greta*: GMQE = 0.67,

identity = 37.39; *Tithorea*: GMQE = 0.68, identity = 38.34).

1. ii. Gene tree estimation

Gene sequences for each opsin family (UV, B, LW) were aligned separately using MACSE

v2 (Ranwez *et al.* 2018), a tool which accounts for underlying codon structure when aligning

protein-coding nucleotide sequences. Aligned sequences were manually cleaned and trimmed

in SeaView v5 (Gouy *et al.* 2021) to ensure the alignment contained no stop or incomplete

codons. The final alignments all contained approximately 1100 nucleotide bases with 370

amino acids which falls within the range of previously characterised lepidopteran opsin

sequences (White *et al.* 2003; Zhan & Reppert, 2012). IQ-TREE (multi-core v1.6.12) was

then used to build a maximum likelihood (ML) nucleotide gene tree for each opsin from these

alignments (iqtree -s alignment_name.fasta -st DNA -bb 10000 -nt AUTO -alrt 1000) using

ModelFinder and ultrafast bootstrap (Fig. 2a; Hoang *et al.* 2018; Kalyaanamoorthy *et al.*

2017; Trifinopoulos *et al.* 2016).

1. iii. Selection analyses

We tested whether an ecological shift from open to shaded forest was associated with relaxed

or intensifying selection on the visual opsins of *Greta* by applying RELAX implemented in

HyPhy, accessed via the Datamonkey webserver (Kosakovsky Pond *et al.* 2020; Muse *et al.*

2018; Wertheim *et al.* 2015). When given prespecified test (*T*) and reference (*R*) branches

from the gene tree, RELAX estimates the rate of nonsynonymous to synonymous nucleotide

base substitutions (d_N/d_S, or ω) among three rate categories using a BS-REL (branch site

random effects likelihood) model (Kosakovsky Pond *et al.* 2011) before raising these values

to the parameter k ($ω^k$) in *T*, which estimates the strength of selection at *T* relative to *R*. When

k = 1, as is the case in the null model, the ω distributions of *T* and *R* are the same. If k < 1, *T*

is under relaxed selection relative to *R*, whereas if k > 1, *T* is under intensified selection

relative to *R* (Wertheim *et al.* 2015). Null and alternative models are then compared with

likelihood ratio tests using a $χ^2$ distribution. In our analyses, the *Greta* branch tip was

specified as *T* for each opsin class with all other branches being assigned to *R,* with the

exception of the UV opsin analysis where duplicated opsins were included, with these

portions of the tree being left as unspecified branches. Additional tests were performed to

compare patterns of selection following gene duplications, where the branch stemming from

the duplicated node were specified as *T* and all other branches assigned as *R.*

2. Animals for physiology and anatomy

Butterflies used in physiological and anatomical comparisons were purchased as pupae or as

live adults from The Entomologist Ltd (*Mechanitis, Methona* and *Tithorea*) and Tropical

Butterflies UK (Edu-Sci Ltd) (*Greta*) respectively. Individuals were reared in 2.2 x 1.8 x

1.8m cages kept at 28-35°C and 80% relative humidity at the University of Bristol's Old Park

Hill Greenhouse facility and were regularly fed 30% sucrose solution. Adults were marked

and sexed immediately post-eclosion and matured for a minimum of three days before being

sampled.

3. Eye physiology comparisons

3.i. Ophthalmoscopy

Interspecific and intraspecific variation in eye physiology were explored using a custom-built

ophthalmoscope (Fig. S1), based on the system described by Brodrick *et al.* (2020). It

consisted of a UI-3590CP-C-HQ-R2 camera with a CMOS colour sensor (Imaging

Development Systems, Germany), 10x 0.25 NA objective lens (Plan N, Olympus, Tokyo,

Japan), beam-splitter (Thorlabs, Newton, MA) and a portable fibre-coupled broadband (470-

850 nm) light emitting diode (LED, MBB1F1 Thorlabs, Munich, Germany) which was

projected along an optical fibre, providing co-axial illumination of the eye. The camera was

connected to a computer with the uEye Cockpit program (part of the IDS Software Suite

4.95) installed for capturing and recording images and videos.

Twelve live individuals of each of the four species were used in the investigation, with

approximately equal numbers of males and females (Table S2). Intact butterflies were

mounted in slotted plastic tubes and further immobilised using plasticine before being

suspended at the centre of rotation of a Newport M-MGM80MS 90° rotation goniometric

cradle by attaching the plastic tube to the arm of a micromanipulator (Fig. S1). Butterflies

were then oriented to set the eye's direction of view. The ophthalmoscope was adjusted to

focus on a region of the cornea where the optical axes of several ommatidia are directly

facing the objective lens, therefore viewing the individual's pseudopupil. After dark adapting

for as briefly as 10 seconds, the LED can be switched on to reveal the eyeshine (or luminous

pseudopupil) created by unabsorbed light reflecting off a mirror-like tracheolar tapetum at the

proximal end of each rhabdom (Franceschini & Kirschfeld, 1971; Stavenga, 1979, 2002).

Unabsorbed reflected light leaves the eyes, referred to as eyeshine, the colour and

heterogeneity of which varies hugely between butterfly species (Belušič *et al.* 2021; Briscoe

& Bernard, 2005; Stavenga *et al.* 2001). As expected, the intensity of this eyeshine

diminished within seconds of illumination due to the intracellular migration of pigment

granules which move from the soma towards the rhabdomeres, preventing light from

reaching the tapetum (Qiu *et al.* 2002; Stavenga *et al.* 1977). This is known hereafter as the

pupillary response. We used the duration of the pupillary response to assay how rapidly eyes

physiologically respond to sudden changes in their light environment.

Video recordings of the eyeshine were taken in the dorsal, frontal, and ventral eye regions

(achieved by rotation of the goniometric cradle) after ten minutes of dark adaptation under

standard laboratory conditions (21℃). When video recordings were made in the frontal eye

region, individuals were also dark adapted for 0.5, 1, 5 and 20 minutes, in a randomised

order, to confirm whether any interspecific patterns are still observed for different lengths of

dark adaptation. We also recorded whether the rate of habituation to a flashing light stimulus

differed between species by switching the LED light source on and off at ten second intervals

for a period of five minutes. Here, our intention was to simulate changes in light conditions as

individuals transition between microhabitats. Videos were imported into FIJI/ImageJ

(Schindelin *et al.* 2012) where response time was visualised and estimated using the View5D

plugin (Fig. 3a). Reflecting facets were counted and categorized as being red or yellow,

where red reflecting ommatidia indicate the presence of long-wavelength shifted

photoreceptors. Using this, the ratio of yellow:red ommatidia was also calculated for the

frontal region of each individual.

Once eye physiological data had been collected, half of the individuals from the eyeshine

study were sampled for micro-computed tomography (micro-CT; see section 4). The second

half of individuals from the eyeshine experiment were sampled for immunohistochemistry

(see section 5).

3. ii. Statistical analysis

All statistical analyses were performed in R (R Core Team 2016). Following Shapiro-Wilk

tests for normality and Bartlett's test for homogeneity of variances, linear models were

constructed, testing the effect of species, sex, and their interaction, on response time in the

frontal region. Subsequent post-hoc comparisons were performed using the TukeyHSD

function. Similar analyses were also performed to test for species and sex differences in

response time in the habituation experiment. The ratios of yellow:red reflecting frontal

ommatidia did not follow a normal distribution, so the non-parametric Kruskal-Wallis test

was performed instead. Intraindividual differences in the number of reflecting ommatidia and

response time between eye regions were analysed by building linear models with the mean

fitted as a fixed effect, the null hypothesis being that the mean difference between eye regions

is zero. For response time, species, sex, and their interaction were included as additional

independent variables.

4. *Micro-computed tomography*

4.i Image acquisition

Heads were severed and placed directly into a chilled fixative (4% paraformaldehyde, 2%

glutaraldehyde, 2% glucose) in 0.1 M sodium phosphate buffer solution (pH 7.4) for 24 hours

under agitation. Heads were dehydrated in a graded ethanol series (10%, 30%, 50%, 70%,

90%, 95%, ten minutes each), before being stained with 1% iodine dissolved in 100% ethanol

for three days under agitation to enhance X-ray absorption contrast (Smith *et al.* 2016; Swart

et al. 2016). Samples were subsequently washed in 100% ethanol three times, each for 30

minutes and stored in ethanol at 4°C until ready for use. The body weight (g) of each

individual was also recorded.

Tomographic images of the samples were taken at the University of Bristol's X-ray

tomography (XTM) facility using a Nikon XTH225ST scanner with a 180kV transmission

target. Heads were typically imaged using x0.33 total magnification and a ~2.2 μm effective

pixel size (see Table S5 for full scanning parameters). 23 heads were scanned in total (n = 6

per species with the exception of *Tithorea harmonia* where one individual experienced

bacterial growth and was thus removed from any subsequent analyses; see Table S3). Scan

acquisition and reconstruction was achieved using Nikon CT Pro and the resulting TIF files

were exported using VG Studio MAX (Fig. 4a). The resolution of the micro-CT scans

allowed identification of the corneal lens of each ommatidium, as well as the area where the

crystalline cones, and light sensitive rhabdoms are located (Fig. 4b-d).

4.ii. Eye volumetric reconstructions

The resulting TIF stacks were compressed from 32 bit to 8 bit and cropped in FIJI/ImageJ (Schindelin *et al.* 2012). They were then loaded into Amira 3D 2021.2 (ThermoFisher Scientific, FEI Visualization Sciences Group). Label files were created for each individual using the *labelfield* module, and segmentation was performed by manually delineating the boundaries of the eye based on intensity contrasts (Fig. 4c,d). Features were labelled every 5-10 slices before being interpolated so all intervening slices could be assigned to the structure of interest. Reconstructions were edited and smoothed before total volumes were extracted using the *measure statistics* module. The volume of each bilaterally paired structure was then multiplied by two and $\log_{10}$ transformed before any analyses.

4. iii. Eye anatomical measurements

Further anatomical measurements were made with the *3D Measurement* tool in Amira to manually estimate ommatidial length (L, µm), interommatidial angle ($\Delta\Phi$, °), and facet diameter (D, µm) (Fig. 4d,e). 5-10 measurements of L, $\Delta\Phi$ and D were taken in each eye region with a mean calculated per region per individual. Eye regions were defined in the XY dimension with the upper, middle, and lower thirds of the eye comprising the dorsal, frontal, and ventral eye regions respectively. In practice, no measurements were taken in ambiguous regions and the distribution of dorsal and ventral measurements are skewed towards the extreme ends of those boundaries as a result. To minimise measurement error when estimating D, the width of five adjacent facets were measured before dividing by five. We also measured interocular distance (IOD, µm), defined as the minimum horizontal gap between the two compound eyes when viewed in the XY dimension, to provide an allometric control in our statistical analyses. As was the case for the volumetric reconstructions, all eye anatomical data were $\log_{10}$ transformed before any analyses.

4.iv. Statistical analysis

To test for allometric scaling differences, linear models were constructed where each measurement of interest was scaled against IOD with species, sex and their interaction included as additional independent variables. If significant species or sex effects were

detected, post-hoc pairwise scaling comparisons were made by building standardised major

axis regressions using the sma function in the *smatr* package (Warton *et al.* 2012). This

function tests for group level deviation from a common scaling relationship between two

traits, modelled as log y = βlog x + α. We first tested for conservation in the allometric slope

(β), deviation from which would suggest an effect of species or sex in how each dependent

variable interacts with *IOD*. If equal slopes were found, the presence of a 'grade shift'

(change in α) along the y-axis was examined, a common indicator of adaptive sensory

divergence (Farnworth & Montgomery, 2022; Kruska, 2005; Montgomery *et al.* 2016;

Sylvester *et al.* 2011). Facet diameter (*D*) did not scale with *IOD* (or central brain or body

size) suggesting that any differences between species occur independently of allometric

scaling. We therefore tested for differences using the absolute, untransformed *D* values.

Intraindividual differences in eye structure between dorsal, frontal, and ventral eye regions

were analysed by applying the same statistical methods used for the eyeshine data, described

above.

5. *Neuroanatomical comparisons*

5.i. Immunohistochemistry

Brain dissections were conducted under HEPES-buffered saline (HBS; 150mM NaCl; 5mM

$CaCl_2$, 25mM sucrose; 10mM HEPES; pH 7.4) with brains subsequently fixed in zinc

formaldehyde solution (ZnFA; 0.25% [18.4mM] $ZnCl_2$; 0.788% [135mM] NaCl; 1.2%

[35mM] sucrose, 1% formaldehyde) for 16-20 hours under agitation, as in Ott (2008). Brains

were washed three times in HBS and placed in Dent's solution (80% methanol/20% DMSO)

for ~two hours under agitation. They were then placed in 100% methanol for one hour at

room temperature before being transferred to fresh methanol and stored at -20 °C until later

use.

Rehydrated brains were embedded in agarose gel (5% UltraPure agarose power in deionised

water) and sectioned using a vibrating blade microtome with an 80 μm interval (Leica

VT1200, Wetzlar, Germany). Sections were then washed six times in PBS-TX (0.5% Triton

X-100 detergent in phosphate-buffered saline), five minutes each, before being incubated in

5% normal goat serum dissolved in PBS-TX (NGS-PBS-TX; NGS; New England BioLabs,

Hitchin, Hertfordshire, UK) for a minimum of one hour at room temperature. Samples were

then stained with anti-allatostatin (Antibody 5F10; Developmental Studies Hybridoma Bank,

University of Iowa, Iowa City, IA, RRID: AB_528076) at a 1:100 dilution in fresh NGS-

PBS-TX and incubated for 24 hours at room temperature under agitation. Allatostatin is a

neuropeptide expressed in distinct cell types within the optic lobes of insects and crustaceans

(Kreissl *et al.* 2010; Lin *et al.* 2021; Sivasubramanian & Sood, 2003) and provides additional

anatomical detail of these brain regions. Slices then underwent 6 x 30 minute PBS-TX

washes before introducing the secondary Cy2-conjugated anti-mouse antibody (Jackson

ImmunoResearch Cat No. 115-225-146, RRID: AB_2307343, West Grove, PA) at a 1:100

dilution in fresh NGS-PBS-TX. Samples were left in the dark at room temperature for a

further 24 hours, before undergoing 6 x 30 minute PBS-TX washes and added to 60%

glycerol in PBS and left overnight. Slices were then transferred to 80% glycerol in PBS

before mounting on glass slides with excess 80% glycerol solution under a cover slip, sealed

with nail varnish.

5.ii. Dextran tracing and wholemount immunohistochemistry

In a separate sample of individuals, we performed *in vivo* dextran injections to explore neural

connections between the optic lobe and structures within the central brain. Live butterflies

were kept in custom-made slotted plastic holders and immobilised using dental wax. A small

window was made in or just behind the compound eye, so part of the optic lobe was exposed.

For some individuals, a window was made above the dorsal central brain instead to allow

access to the mushroom bodies. A few crystals of dextran tetramethylrhodamine

(Invitrogen^{TM} D1868,Fisher Scientific, Leicestershire, UK) were dissolved in bovine serum

albumin (BSA, Merck Life Science UK Ltd, Gillingham, Dorset, UK) and placed on the tip

of a horizontally pulled glass electrode (tip diameter ~200 µm). Dextran was manually

inserted into either both optic lobes or both mushroom bodies calyces under red-filtered light

using a dissection microscope. Any remaining dextran on the brain surface was rinsed off

with Ringer's solution (Ringer; 150 mM NaCl, 3mM CaCl$_2$, 3mM KCl, 2mM MgCl$_2$, 10mM

HEPES, 5mM glucose, 20mM NaOH, pH 6.9). The intact butterfly head was then left

submerged in Ringer's solution for a minimum of four hours at room temperature to allow

the dextran to be transported to the terminal projection sites of affected neurons. The brain

was then dissected and stored using the protocols described above.

To reveal neuropil structure, injected brains were stained as wholemounts using indirect

immunolabelling against synapsin (Antibody 3C11; Developmental Studies Hybridoma

Bank, University of Iowa, Iowa City, IA, RRID: AB_2315424), a conserved insect protein

expressed at presynaptic regions (Brandt *et al.* 2005; Heinze & Reppert, 2011; Klagges *et al.*

1996; Montgomery & Ott, 2015). Brain samples were rehydrated in a decreasing methanol-

Tris buffer dilution series (90%, 70%, 50%, 30%, 0%, pH 7.4) for ten minutes each. Samples

were incubated in NGS (NGS; New England BioLabs, Hitchin, Hertfordshire, UK) diluted in

0.1 M PBS (pH 7.4) and 1% DMSO for two hours at room temperature, before the primary

antibody was added at a 1:30 dilution with fresh NGS-PBSd and left at 4°C for 3.5 days

under agitation. Non-bound primary antibody was removed after 3 x two hour PBSd washes

before introducing the secondary Cy2-conjugated anti-mouse antibody at a 1:100 dilution in

NGS-PBSd and left in the dark at 4°C for a further 2.5 days under agitation. Samples were

then washed in glycerol diluted in 0.1 M Tris buffer (1% DMSO) in an increasing dilution

series (1%, 2%, 4%, for two hours each, 8%, 15%, 30%, 50%, 60%, 70%, 80%, for one hour

each) under agitation. Samples were dehydrated and washed in 100% ethanol three times, for

30 minutes each, before clarification by underlaying the ethanol with methyl salicylate.

5.iii. Confocal microscopy

All immunostained brains were imaged on a confocal laser-scanning microscope (Leica SP5-

AOBS/SP5-II, Leica Microsystem, Mannheim, Germany) at the University of Bristol's

Wolfson Bioimaging Facility, using either a 10x 0.4 NA or 20x 0.7 NA objective lens (Leica

Material Nos. 506285, 506513, Leica Microsystem). Single-capture images of wholemounts

and sections were achieved with a 488 nm argon laser (20% intensity), and an x-y resolution

of 512x512 pixels. For dextran-traced individuals, an additional 580 nm argon laser channel

(20% intensity) was included and scanned sequentially with the 488 nm laser to visualise the

passage of dextran dye against the synapsin-stained background.

5.iv Volumetric comparisons

Volumetric reconstructions of the four primary optic neuropils (lamina, medulla, lobula plate

and lobula) were obtained from the micro-CT scans using the same methods described above

(Fig. 7a). Paired neuropils were doubled and log_{10} transformed and analysed using the same

statistical methods as the eye volumetric data. To test for associations between physically and

functionally linked visual structures, a covariance matrix for the volumetric data was built by

running multiple linear regressions with each structure of interest as a dependent variable and

species as a random effect. We then tested for species and sex differences after accounting for

these covariances by including species and sex as additional independent variables in these

regressions.

Results

1. Evidence of duplication and habitat-related relaxation of selection in UV opsins

We recovered single copies of UV, B and LW sensitive opsins in the genomes of all four

ithomiine species, except for *Tithorea* where two putative full-length UV opsin sequences

containing no stop codons were identified. Our selection analyses revealed the UV opsin of

the shade dwelling *Greta* to show a significantly lower rate of nonsynonymous to

synonymous nucleotide base substitutions relative to all assigned 'open habitat' reference

branches (Fig. 2b; UV opsin, relaxation parameter, k = 0.17, χ2 test p = 0.016) when

contrasted against the null model where all test and reference branches are assumed to be

under the same selection pressures, indicating relaxed selection (see Methods). This null

model was found to hold true when the same analyses were repeated on the B and LW opsin

sequences (B opsin, k = 22.21, p = 0.314; LW opsin, k = 0.290, p = 0.174). The duplicated

UV opsin of *Tithorea* also showed signatures of significant relaxed selection (Fig. 2c; k =

0.000, p < 0.001), potentially indicative of ongoing pseudogenization. Our results therefore

may suggest shifts in selective regime operating at the molecular level, mediated by both

habitat preference and gene duplication, which together might drive further variation in how

visual systems respond to light.

2. The response to light varies across the eye and between species

We found significant variation in the retinal mosaic (Fig. 3) between species in the frontal

region (yellow:Red reflecting frontal ommatidia, χ^2 = 16.678, df =3, p = 0.001). *Greta,*

Mechanitis and *Tithorea* all had distinguishable red and yellow reflecting facets, where

Tithorea contained a greater proportion of red-reflecting ommatidia. *Methona* displayed more

homogeneous orange eyeshine patterns (which were classified as red in the above analysis)

(Fig. 3a, Table S4A). Therefore, all four species show evidence of a red-sensitive spectral

channel. The number of shining ommatidia within each pseudopupil was greatest in the

frontal region (dorso-frontal, t = 12.830, df = 37, p < 0.001; fronto-ventral, t = 3.540, df = 37,

p = 0.003; Table S4D), suggesting that this is where spatial resolution is at its highest. The

dorsal region contained the fewest shining facets.

When testing for differences in frontal pupillary response time (RT), we found evidence of

significant interspecific variation in response time (F = 15.870, df = 3, p < 0.001) with no sex

or interaction effects between species or sex (Fig. 3b, see Table S4B). Post-hoc analysis

revealed that overall, *Greta* and *Mechanitis* had longer response times regardless of the

length of dark adaptation (Fig. 3c, Table S4B). Response time decreased after five minutes of

exposure to a flashing light stimulus (t = 2.354, df = 46, p = 0.011), but no differences in this

habituation time were found between species, sex, or their interaction (see Table S4C).

Finally, the pupillary response of ventral ommatidia was significantly longer than those in the

dorsal (t = 5.460, df = 46, p < 0.001) and frontal (t = 5.121, df = 46, p < 0.001) eye regions

with no significant species, sex, or interaction effects (Fig. 3d). Significant species

differences were only found when testing for dorso-frontal variation (F = 11.627, df = 3, p <

0.001) (Table S4E).

3. Eye structure varies between species, sexes, and eye regions

Micro-CT scans revealed quantifiable differences in overall eye size and structure between

species (Fig. 4a,b). Each ommatidium within the apposition compound eye consists of a

corneal lens and a deep crystalline cone which directs light through to the light-sensitive

rhabdom where the photoreceptor cells are located (Fig. 1a, 4c,d). Relative eye volume varied

significantly between species (F = 6.232, df = 3, p = 0.007) when scaled against *IOD*.

Allometric analysis also revealed that these differences were a result of grade-shifts along the

y-axis (Wald χ^2 = 8.483, df = 3, p = 0.037, (Fig. 5a) with *Greta* investing more in eye volume

than the similarly sized *Mechanitis* (Table S6), for example. Significant grade-shifts were

also observed between *Methona* and *Tithorea*, with the latter sharing the same scaling

relationship as its co-mimic, *Mechanitis* (Fig. 5a).

Frontal and ventral ommatidial length (*L)* also varied between species (frontal region, F =

4.292, df = 3, p = 0.024; ventral region, F = 4.048, df = 3, p =0.029) with significant, non-

allometric interspecific grade-shifts found for both eye regions (frontal, Wald χ^2 = 23.08, df =

3, p < 0.001, (Fig. 5b; ventral, Wald χ^2 = 15.23, df = 3, p = 0.002). For frontal ommatidia,

pairwise comparisons revealed *Greta* to have relatively longer ommatidia than tiger-stripe co-

mimics *Mechanitis* and *Tithorea* which showed a conserved scaling relationship (Table S6C).

Interommatidial angle ($\Delta\Phi$) did not show any significant species differences when scaled

against *IOD*, suggesting that variation in this trait is largely determined by variation in head

size (Table S6A). Testing for differences in the absolute, untransformed values for facet

diameter, *D*, revealed significant interspecific variation for the frontal and ventral eye regions

(frontal, F = 8.429, df = 3, p= 0.006, (Fig. 5c; ventral, F = 6.373, df = 3, p = 0.015). Post-hoc

comparisons of these models showed that the majority of significant pairwise contrasts were

with *Mechanitis,* which had consistently the narrowest facets (Fig. 5c, Table S6B).

Across the four species, evidence of sexual dimorphism was observed for relative eye volume

(F = 17.358, df = 1, p = 0.001) with males displaying an increased investment in eye size

(Wald χ^2 = 12.37, df = 1, p < 0.001, (Fig. 5d). However, our sample sizes were insufficient to

detect whether these differences in eye volume were explained by any other measured eye

anatomical traits (Fig. 5e,f). We found no interactions between sex and species for any

anatomical variables.

Finally, comparing intraindividual variation in eye anatomy, we found significant variation

between the dorsal, frontal, and ventral eye regions from the micro-CT scans (Fig. 5g-i).

After multiple test correction, no species, sex, or interaction effects were found, with the

exception of *D* (where dorso-frontal differences between species were observed for females

but not males, see Table S6E for details)*,* suggesting these differences are mostly consistent

across all individuals and species. Dorsal ommatidia were shorter and narrower than frontal

and ventral ommatidia (Table S6E) and the $\Delta\Phi$ of dorsal ommatidia was significantly greater

than the other two eye regions (dorso-frontal, t = 8.949, df = 22, p < 0.001; dorso-ventral, t =

5.342, df = 22, p < 0.001).

4. Gross composition of optic lobe neuropils is generally conserved across species but there

is evidence of sexual dimorphism in the presence of the ventral lobula

Staining against allatostatin revealed the retinotopic organisation of the optic lobe,

particularly within the lamina and medulla where the respective cartridge and column

modules are clearly visible (Fig. 6a). Distinct layers of neuronal organisation are present in

the medulla, consisting of two major outer and inner divisions, separated by the serpentine

layer, as seen in other ithomiines (Montgomery & Ott, 2015). Within each division are

several further striations where a maximum of seven medulla layers were clearly observed in

the scans of all four species (Fig. 7a). The lobula plate and the lobula have two and three

layers respectively, as observed in other Lepidoptera (e.g. Hamanaka *et al.* 2012). Two

smaller neuropils, the accessory medulla, found against the anteromedial edge of the medulla,

and the ventral lobula, an ovoid ventromedial structure which forms a close physical

association with the lobula, were more clearly visibile in whole mount anti-synapsin stains.

Our samples suggest that the ventral lobula, which is smaller in ithomiines than in other

butterflies where it has been identified (Wainwright & Montgomery, 2022), is sexually

dimorphic in *Greta* and *Mechanitis,* being totally absent in all males of these species (Fig.

6b). In *Papilio xuthus* and Heliconiini butterflies, the ventral lobula appears to act as a relay

centre between the optic lobe and the mushroom body calyx (Couto et al 2023; Kinoshita *et*

al. 2015). However, our dextran injections provided no evidence that the ventral lobula serves

this purpose in ithomiines. The ventral lobula appears to receive inputs via the lobula, without

innervation in the lobula plate (Fig. 6c), reminiscent of *Drosophila* medulla projection (Tm)

neurons which receive direct synaptic inputs from photoreceptors and relay this directly to

the lobula (Borst, 2009; Lin *et al.* 2015). Our dextran injections suggested the ventral lobula

then projects to the bulb in the central brain, a structure which forms part of the anterior

visual pathway in *Drosophila* (Hardcastle *et al.* 2021; Lovick *et al.* 2017; Omoto *et al.* 2017).

Dextran staining in the anterior optic tubercle, the main optic neuropil in the central brain,

was also found for some individuals, but the precise neuronal inputs to this region are not

clear from our confocal scans. Injections into the mushroom body calyces also provided no

evidence that these structures receive projections from the ventral lobula.

5. *Despite a conserved composition, interspecific and intersexual variation in the visual*

pathway extends to investment in visual neuropils

Volumetric data from segmented micro-CT scans allowed further quantitative volumetric

comparisons of the main visual neuropils (Fig. 7). Two-way ANOVAs revealed significant

interspecific differences for all four structures (lamina, $F = 4.617$, $df = 3$, $p = 0.019$; medulla,

$F_3 = 12.543$, $p < 0.001$; lobula plate, $F_3 = 11.806$, $p < 0.001$; lobula, $F_3 = 14.180$, $p < 0.001$;

optic lobe, $F_3 = 11.192$, $p < 0.001$) with no sex or interaction effects found for any neuropils.

Similar to our comparisons of eye volume, allometric analysis revealed all species effects to

be the result of grade-shifts in visual investment (lamina, Wald $\chi^2_3 = 17.86$, p < 0.001;

medulla, Wald $\chi^2_3 = 43.42$, p < 0.001; lobula plate, Wald $\chi^2_3 = 39.27$, p < 0.001; lobula, Wald

$\chi^2_3 = 40.80$, p < 0.001; optic lobe, Wald $\chi^2_3 = 30.68$, p < 0.001; (Fig. 7b). For the medulla,

lobula plate, lobula, and optic lobe as a whole, significant pairwise comparisons commonly

involved *Greta*, which consistently invested more in visual neuropil, for a given brain size,

than the other three species, when accounting for *IOD* (Table S6).

A covariance matrix revealed significant correlations between the size of some of the optic

neuropils and the eye (see Fig. S2). After controlling for this covariance and including

species as an additional independent variable, the significant species effects remained for the

eye (F = 14.232, df = 3, p < 0.001) and the lobula plate (F = 3.530, df = 3, p = 0.046).

Interspecies differences in eye size therefore appear, on the most part, to evolve in concert

with investment in optic neuropils. Interestingly, interactions between functionally related

neuropils appear to be obscuring sex effects as significant sex differences in the medulla were

revealed after controlling for this covariance (F = 13.333, df = 1, p = 0.003) with males

investing in larger medullas. This is consistent with our observed patterns of sexually

dimorphic eye investment (Fig. 5d), which also remained after controlling for this covariance

(F = 49.124, df = 1, p < 0.001).

Discussion

Specialization in visual systems, likely reflects selection pressures imposed by ecological

needs, which act at the functional, physiological, anatomical and molecular level (e.g.

Bartholomée *et al.* 2023; Gonzalez-Bellido *et al.* 2011; Hofmann *et al.* 2009; Huber *et al.*

1997; Scales & Butler, 2016; Sugawara *et al.* 2005; Zhao *et al.* 2009). By measuring

physiological, anatomical, and molecular variation in the eyes and visual neuropils of four

closely butterfly species, we demonstrate multiple dimensions of variation in the visual

system of Ithomiini. The 26-Myr-old adaptive radiation of these butterflies has been linked to

microhabitat partitioning (Chazot *et al.* 2019; Elias *et al.* 2008; Hill, 2010), suggesting

species are exposed to contrasting patterns of sensory information. By identifying aspects of

the visual system that differ between species, we provide a foundation for future work linking

variation in sensory ecology and sensory systems. Below we discuss our results with this eco-

evolutionary framework in mind.

Duplication and relaxed selection in UV opsins

Selection analyses on recovered visual opsin sequences revealed evidence of relaxation of

selection. We found evidence of relaxed selection in the coding sequence of the UV-sensitive

visual opsin in *Greta*, relative to nymphalid butterfly species living in open habitats (Fig. 2b;

see Methods). Photosynthesising vegetation tends to absorb short wavelengths, particularly

UV, making the light environment of the forest understory less rich in UV radiation than

more exposed forest microhabitats (Endler, 1993; Théry, 2001). UV light is known to play a

major role in butterfly inter/intraspecific communication and navigation (Briscoe *et al.* 2010;

Brunton & Majerus, 1995; Froy *et al.* 2003; Sauman *et al.* 2005) but these cues may be less

reliable for species occupying such densely shaded forest. Similar patterns were not observed

for the blue and long wavelength sensitive (B and LW respectively) visual opsins. This

relaxation of selective constraint might indicate a diminished role of UV-orientated

behaviours in *Greta*, in comparison to its close relatives.

In contrast, we identify a duplication of the UV opsin in *Tithorea*, making this the second

independent duplication of this gene identified in butterflies (McCulloch *et al.* 2017; two

additional duplications of this gene have been identified in non-Papilionoid Lepidoptera but

with no known functional role, see Sondhi *et al.* 2021). In *Heliconius* butterflies, which

exhibit an independent UV opsin duplication, the duplicated gene experienced positive

selection in some lineages but relaxed selection in others, with the latter leading to

pseudogenization or down-regulated expression of this opsin in photoreceptor cells. While it

is tempting to link UV duplication to increased reliance on UV cues in *Tithorea*, the

duplicated copy also appears to have experienced relaxed selection, which raises questions

over its functionality (Fig. 2c). Nevertheless, our data illustrate the potential for divergent

repertoires of opsin genes, and selection regimes governing their evolution, across Ithomiini.

Larger scale interspecific comparisons could provide opportunities to link this variation to

preferences for different light environments.

Response to temporal and spatial light variation

Despite evidence of visual opsin conservation across all four species, photoreceptor

sensitivity can also be shifted by changes in eye physiology. Our eyeshine images revealed a

diversity of retinal mosaic patterns, caused by fixed combinations of photosensitive

rhodopsins with additional screening pigments (Briscoe, 2008; Briscoe & Bernard, 2005;

Stavenga & Arikawa, 2006). These screening pigments create the red-orange reflecting

ommatidia seen in our four species (Fig. 3a), providing these species with an additional

spectral channel for discriminating longer wavelengths (Belušič *et al.* 2021; Zaccardi *et al.*

2006).

Intraindividual variation between dorsal, frontal, and ventral compound eye regions likely

mirror spatial differences in sensory cue abundance and variability (Fig. 3A). Dorsal facets

are primarily yellow-green in colour whereas frontal and ventral facets are orange-red. This

indicates a specialisation for short wavelength sensitivity in dorsal ommatidia, which are

exposed to a greater abundance of these wavelengths in downwelling light (Briscoe, 2008;

Stalleicken *et al.* 2006). In addition, shorter ommatidia found within the dorsal region is a

way of minimising self-screening for improved polarization vision, which other butterflies

have been shown to utilise for orientation and navigation, particularly within the dorsal rim

area (Kinoshita & Arikawa, 2014; Labhart & Meyer, 1999; Meyer & Labhart, 1992; Nilsson

et al. 1987; Reppert *et al.* 2004) (Fig. 5h). Although the dorsal rim area was not delineated in

our image data, larger interommatidial angles in the dorsal area might hint that this region is

optimised for enhanced contrast detection rather than improved visual acuity. The latter

would be more beneficial in the frontal and ventral regions to optimise the localisation of

hostplants, foodplants and mates (Bergman *et al.* 2021; Labhart & Meyer, 1999; Labhart *et*

al. 2009; Land, 1989) (Fig. 5g). A greater abundance of downwelling light hitting the dorsal

ommatidia (Stavenga, 2002) could also explain the narrower facets found within this region

(Fig. 5i). In our ophthalmoscopy experiment, we found that the ventral region showed longer

pupillary response times which might reflect lower temporal variability in upwelling light

intensity (Fig. 3d). Dorsal facing ommatidia must respond to rapid fluctuations in

downwelling solar irradiance caused by gaps in the forest canopy whereas frontal ommatidia

must respond quickly to changes in other types of visual contrast, such as contrast variation

created by optic flow when the animal is in flight to reduce motion blur. This role in

stabilising visual processing during flight potentially explains the faster response times in

these eye regions (Endler, 1993; Grittner *et al.* 2022; Palermo & Theobald, 2019; Théry,

2001). However, differences in pupillary kinetics between red and yellow reflecting

ommatidia, which may further interact with variation in rhabdom and photoreceptor cross-

sectional area, might also explain this result.

Comparing differences in frontal pupillary response time revealed that *Greta* and *Mechanitis*

show slower responses times when exposed to a bright light stimulus following dark

adaptation (Fig. 3b,c). Responding more slowly to fluctuations in light conditions may reflect

a history of evolution within more homogeneous environments, and a strong preference for

particular light conditions. Indeed, *Greta* tends to occupy constantly shaded forest understory

(Elias & Joron, 2015; Willmott & Mallet, 2004)*, while *Mechanitis* might consistently occupy

sunlit environments and therefore respond less rapidly to changes in light intensity than those

from patchier forest where *Methona* and *Tithorea* are found*. In addition, the longer, thinner

wings of *Methona* are indicative of greater flight speeds (Hill, 2021), which might select for

faster physiological responses due to enhanced optic flow and contrast change during flight

(Hill, 2021). However, we cannot rule out a degree of phylogenetic inertia in predicting these

differences, given the closer relationships between these latter two species (Chazot *et al.*

2019). Larger species comparisons coupled with fine-scale ecological data (e.g. flight speed,

light temporal variability, canopy cover) are required to disentangle the factors driving eye

physiological variation but the well documented natural history of ithomiines (e.g. Beccaloni,

1997; Brown & Freitas, 1994; Willmott & Mallet, 2004) make them a prime model system

for tackling these questions in future work.

Interspecific differences in eye and optic lobe structure

Across our four species, we identified extensive interspecific differences in optical and neural

architecture which could not be explained solely by variation in overall head size or by

evolutionary relatedness. Differences were observed for relative eye size, ommatidial length,

facet diameter and the relative size of all four measured optic neuropils. In contrast, the lack

of interspecific differences in interommatidial angle ($\Delta\Phi$) suggest that differences in visual

acuity can be fully explained by differences in overall head size (as in Jander & Jander, 2002;

Rutowski *et al.* 2009).

Larger relative eye size in *Greta* could, at least partially, be explained by enhanced

ommatidial length in the frontal and ventral eye regions of this species (Fig. 5a,b). Longer

rhabdoms provide a greater photoreceptive surface, and longer crystalline cones would

enhance focal length. Together, this would optimise the eye for increased light sensitivity at

greater distances (Greiner *et al.* 2004a; Warrant & McIntyre, 1993), which may be required

in low-light conditions. The large relative volume of the lamina in *Greta* is consistent with

the hypothesis that this species is enhancing either the signal-to-noise ratio for improved

achromatic visual sensitivity and/or the amplitude of spectral opponency for improved colour

vision (Fig. 7b; Matsushita *et al.* 2022; Sterling & Laughlin, 2015; Stöckl, 2022). The lamina

is retinotopically organised, with lamina monopolar cells (LMCs) within neural cartridges

receiving visual input from individual ommatidia (Fig. 6a). These LMCs spatially integrate

signals for enhancing light sensitivity (Greiner *et al.* 2005; Stöckl *et al.* 2016, 2020). Larger

lamina volumes observed in *Greta* therefore suggests that this species devotes greater

neuronal resources to processing each visual pixel. Indeed, these patterns of investment are

even more extreme in the remaining three optic neuropil where *Greta* shows consistently

larger visual neuropil than the other three species, controlling for head size (Fig. 7b).

Therefore, increases in relative eye size appear to positively correlate with increases in neural

processing (see also Garamszegi *et al.* 2002; Corral-López *et al.* 2017).

The relatively larger eyes, optic lobes and longer ommatidia of *Greta* might have evolved to

enhance visual sensitivity, reflecting the shaded low-light forest understory where this species

and its co-mimics are usually found (Elias *et al.* 2008; Willmott & Mallet, 2004; Willmott *et*

al. 2017). In dense forest environments, overall illuminance is likely to be lower in

comparison to the more sunlit forest frequented by *Mechanitis, Tithorea,* and *Methona*, which

all fly at higher elevations (Beccaloni, 1997; Elias *et al.* 2008; Endler, 1993, Hill, 2010). We

note that these changes in relative investment may also arise to maintain functional visual

sensitivity rather than increase it, given the constraints posed by a smaller head size

(Rutowski *et al.* 2009). For example, the mean facet diameter (*D*) of *Greta* is similar to those

of *Methona* and *Tithorea* despite these two species having a body size 7.12 and 4.13 times

larger than that of *Greta* respectively (Fig. 5c). Hence, to overcome allometric size

constraints and maintain visual performance, facet diameter must vary independently of head

size.

Sexual dimorphism in the visual system

Patterns of eye sexual dimorphism within Ithomiini match previous findings from other

Lepidoptera (e.g. Everett *et al.* 2012; Meyer-Rochow & Lau, 2008; Ziemba & Rutowski,

2000) and arthropods (e.g. Zeil, 1983) where relative eye size is larger in males. Other studies

have proposed this to reflect greater visual sensitivity for male-limited visual behaviours such

as the searching and localisation of potential mates and territorial defence (Bergman *et al.*

2021; Everett *et al.* 2012; McClure *et al.* 2019; Pliske, 1975; Rutowski, 2000) (Fig. 5d,f).

This enhanced activity may also select for males to become more visually sensitive to the

presence of predators (Everett *et al.* 2012). Unlike relative eye size, sexual dimorphism was

not found for structures within the optic lobe, except for the medulla but this is only revealed

after controlling for covariance between optic neuropils (Fig. S2). Therefore, intersexual

differences in visual ecology might promote changes in visual perception without

downstream shifts in visual processing (e.g. by changing the number of lamina cartridges).

However, we also uncovered a rare case of dimorphism in the presence or absence of an

apparently butterfly-specific optic neuropil – the ventral lobula (Fig. 6b). This small neuropil

was absent in the ten *Greta* and seven *Mechanitis* males we examined, but present in both

sexes of *Methona* and *Tithorea*. In other butterflies, the ventral lobula appears to be a relay

centre directing visual projection neurons to the mushroom body calyx (Kinoshita *et al.* 2015;

Couto et al. 2023). However, our own dextran neural tracing revealed no evidence of this

pathway in ithomiines (Fig. 6c). Dextran fluorescence is instead detected in the bulb of the

central brain which might indicate that the ithomiine ventral lobula has different functional

roles which, in the case of *Greta* and *Mechanitis*, presumably coordinates female-specific

behaviours. In other insects, the bulb receives projections from the anterior optic tubercle and

relays these signals to the ellipsoid body of the central complex, a collection of neuropils

known to coordinate navigation and orientation behaviours (Hardcastle *et al.* 2021; Heinze *et*

al. 2013; Lovick *et al.* 2017; Omoto *et al.* 2017). However, the small size of the ithomiine

ventral lobula in comparison to other studied butterflies suggests overall lower functional

performance in this neuropil (Montgomery & Ott, 2015; Wainwright & Montgomery, 2022).

Nevertheless, this work provides striking evidence of neuroanatomical sexual dimorphism,

further highlighting the lability of the nervous system to vary across short ecological and

temporal scales.

Summary

Our qualitative and quantitative analysis reveals variation between and within species which

potentially reflect the different ecological conditions they occupy. Previous comparative

work in a wild community of ithomiine butterflies mirrors these findings, having shown that

small-scale ecological shifts can predict adaptive changes in other aspects of visual

neuroanatomy (Wainwright & Montgomery, 2022). Our current work further integrates

information on eye structure, physiology, and molecular evolution. We present multiple lines of evidence that extensive visual system variation can exist within diverse insect radiations, hinting at the evolutionary lability of these systems to rapidly develop specialisations to distinct visual ecological niches, with selection potentially acting at both the perceptual. processing, and molecular level. We found physiological and anatomical differences between species, within species and within individuals, as well as evidence of gene duplications and relaxed selection in the visual opsins and suggest various explanatory hypotheses for these results by linking our findings with knowledge of ithomiine natural history, behaviour, and ecology. Larger comparative analyses across a greater number of species will enable the discrimination of adaptive processes from phylogenetic effects and repeated examples of microhabitat divergence and convergence within single communities make Ithomiini an ideal model system for identifying the ecological drivers of eye, brain, and visual pigment diversification in the wild.

Acknowledgements

We are grateful to Alanna Kelly and Tom Pitman for support at the University of Bristol's Old Park Hill Greenhouse facility, and to Amaia Alcalde for assisting in butterfly rearing. We are also thankful to the University of Bristol's XTM Facility, Wolfson Bioimaging Facility and High Performance Computing Facility for micro-CT, confocal microscope and bioinformatics support respectively. We also thank Nick Chazot for providing the phylogeny used in Fig. 1b, Antoine Couto for his assistance with the dextran tracing and to Jessica Foley for her support with aspects of the immunohistochemistry and microscopy protocols.

Competing interests

The authors declare no competing interests.

Author contributions

Conceptualisation: JBW, SHM; Methodology: JBW, EMS, EAB, FC, MJH, SHM; Validation: NWR, SHM; Formal analysis: JBW; Investigation: JBW, CS, MC, DP, EMS; Resources: EMS, EAB, FC; MJH; Data curation: JBW; Writing – original draft: JBW; Writing – review and editing: JBW, EMS, EAB, FC, MJH, NWR, SHM; Visualisation: JBW,

EAB; Supervision: NWR, SHM; Project administration: JBW, SHM; Funding acquisition:

JBW, SHM.

Funding

This work was supported by the Natural Environment Research Council (GW4+ NERC DTP

to JBW and a NERC IRF NE/N014936/1 to SHM), and the Air Force Research

Laboratory/Air Force Office of Scientific Research (AFOSR) through the European Office of

Aerospace Research and Development (EOARD) grant FA9550-18-1-7005.

Data availability

All raw data can be found in Table S1-3 and all opsin sequence alignment and gene tree files

will be uploaded to Dryad Digital Repository upon acceptance. All opsin sequences have

been uploaded GenBank (accession IDs can be found in Table S1). All pseudopupil video

recordings and micro-CT scans are available from Zenodo

https://doi.org/10.5281/zenodo.8154379 (Wainwright *et al.* 2023).

References

Arikawa, K., Pirih, P. & Stavenga, D. G. (2009). Rhabdom constriction enhances filtering by

the red screening pigment in th eye of the Eastern Paled Clouded yellow butterfly, *Colias*

erate (Pieridae), *Journal of Experimental Biology*, **212**, 2057-2064.

Bartholomée, O., Dwyer, C., Tichit, P., Caplat, P., Baird, E. & Smith, H. G. (2023). Shining a

light on species coexistence: visual traits drive bumblebee communities, *Proceedings of the*

Royal Society B Biological Sciences, **290**, Article no. 20222548.

Bates, H. W. (1862). XXXII. Contributions to an insect fauna of the Amazon valley

(Lepidoptera: Heliconidae), *Transactions of the Linnean Society London*, **23**, 495-566.

Beccaloni, G. W. (1997). Vertical stratification of ithomiine butterfly (Nymphalidae:

Ithomiinae) mimicry complexes: the relationship between adult flight height and larva host-

plant height, *Biological Journal of the Linnean Society*, **62**, 313-341.

Belušič, G., Ilić, M., Meglič, A. & Pirih, P. (2021). Red-green opponency in the long visual

fibre photoreceptors of brushfoot butterflies (Nymphalidae), *Proceedings of the Royal Society*

B-Biological Sciences, **288**, Article no. 20211560.

Bergman, M., Smolka, J., Nilsson, D-E. & Kelber, A. (2021). Seeing the world through the

eyes of a butterfly: visual ecology of the territorial males of *Pararge aegeria* (Lepidoptera:

Nymphalidae), *Journal of Comparative Physiology A*, **207**, 701-713.

Borst, A. (2009). *Drosophila's* view on insect vision, *Current Biology*, **19**, 36-47.

Brandt, R., Rohlfing, T., Rybak, J., Krofczik, S., Maye, A., Westerhoff, M., Hege, H-C. &

Menzel, R. (2005). Three-dimensional average-shape atlas of the honeybee brain and its

applications, *Journal of Comparative Neurology*, **492**, 1-19.

Briscoe, A. D. (2008). Reconstructing the ancestral butterfly eye: focus on the opsins,

Journal of Experimental Biology, **211**, 1805-1813.

Briscoe, A. D. & Bernard, G. D. (2005). Eyeshine and spectral tuning of long wavelength-

sensitive rhodopsins: no evidence for red-sensitive photoreceptors among five Nymphalini

butterfly species, *Journal of Experimental Biology*, **208**, 687-696.

Briscoe, A. D. & Chittka, L. (2001). The evolution of color vision in insects, *Annual Review*

of Entomology, **46**, 471-510.

Briscoe, A. D., Bybee, S. M., Bernard, G. D., Yuan, F., Sison-Mangus, M. P., Reed, R. D.,

Warren, A. D., Llorente-Bousquets, J. & Chiao, C-C. (2010). Positive selection of a

duplicated UV-sensitive visual pigment coincides with wing pigment evolution in *Heliconius*

butterflies, *Proceedings of the National Academy of Sciences,* **107**, 3628-3633.

Brodrick, E. A., Roberts, N. W., Sumner-Rooney, L., Schlepütz, C. M. & How, M. J. (2020).

Light adaptation mechanisms in the eye of the fiddler crab *Afruca tangeri*, *Journal of*

Comparative Neurology, **529**, 616-634.

Brown, Jr. K. S. & Freitas, A. V. L. (1994). Juvenile stages of Ithomiinae: overview and

systematics (Lepidoptera: Nymphalidae), *Tropical Lepidoptera*, **5**, 9-20.

Brunton, C. F. A. & Majerus, M. E. N. (1995). Ultraviolet colours in butterflies: intra- or

inter-specific communication?, *Proceedings of the Royal Society B Biological Sciences*, **260**,

199-204.

Chazot, N., Willmott, K. R., Lamas, G., Freitas, A. V. L., Piron-Prunier, F., Arias, C. F.,

Mallet, J., De-Silva, D. L. & Elias, M. (2019). Renewed diversification following Miocene

landscape turnover in a Neotropical butterfly radiation, *Global Ecology and Biogeography*,

28, 1118-1132.

Cicconardi, F. C., Morris, B. J. & Montgomery, S. H. (2023). Comparative antennal

transcriptomics in Ithomiini butterflies reveals novel sex-specific genes and hyper-diverse

inter-species expression pattern, *in prep.*

Corral-López, A., Garate-Olaizola, M., Buechel, S. D., Kolm, N. & Kotrschal, A. (2017). On

the role of body size, brain size, and eye size in visual acuity, *Behavioral Ecology and*

Sociobiology, **71**, Article no. 179.

Couto, A., Wainwright, J. B., Morris, B. J. & Montgomery, S. H. (2020). Linking ecological

specialisations to adaptations in butterfly brains and sensory systems, *Current Opinion in*

Insect Science, **42**, 55-60.

Couto, A., Young, F. J., Atzeni, D., Marty, S., Melo-Flórez, L., Hebberecht, L., Monllor, M.,

Neal, C., Cicconardi, F., McMillan, W. O., & Montgomery, S. H. (2023). Rapid expansion

and visual specialization of learning and memory centers in Heliconiini butterflies,

Nature Communications. **14**, Article no. 4024.

Cronin, T. W., Johnsen, S., Marshall, N. J. & Warrant, E. J. (2014). *Visual Ecology*,

Princeton University Press, Princeton, USA.

Dacke, M., Nordström, P., Scholtz, C. & Warrant, E. (2002). A specialised dorsal rim area for

polarized light detection in the compound eye of the scarab beetle *Pachysoma striatum,*

Journal of Comparative Physiology A, **188**, 211-216.

Elias, M., Gompert, Z., Jiggins, C. & Willmott, K. (2008). Mutualistic interactions drive

ecological niche convergence in a diverse butterfly community, *Plos Biology*, **6**, 2642-2649.

Elias, M. & Joron, M. (2015). Mimicry in *Heliconius* and Ithomiini butterflies: The profound

consequences of an adaptation, *BIO Web of Conferences,* **4**, Article no. 0008.

Endler, J. A. (1993). The color of light in forests and its implications, *Ecological*

Monographs, **63**, 1-27.

Endler, J. A., Westcott, D. A., Madden, J. R. & Robson, T. (2005). Animal visual systems

and the evolution of color patterns: sensory processing illuminates signal evolution,

Evolution, **59**, 1795-1818.

Everett, A., Tong, X., Briscoe, A. D. & Monteiro, A. (2012). Phenotypic plasticity in opsin

expression in a butterfly compound eye complements sex role reversal, *BMC Evolutionary*

Biology, **12**, Article no. 232.

Fain, G. L., Hardie, R. & Laughlin, S. B. (2010). Phototransduction and the evolution of

photoreceptors, *Current Biology*, **20**, 114-124.

Farnworth, M. S. & Montgomery, S. H. (2022). Complexity of biological scaling suggests an

absence of systematic trade-offs between sensory modalities in *Drosophila*, *Nature*

Communications, **13**, Article no. 2944.

Feuda, R., Marlétaz, F., Bentley, M. A. & Holland, P. W. H. (2016). Conservation,

duplication, and divergence of five opsin genes in insect evolution, *Genome Biology And*

Evolution, **8**, 579-587.

Franceschini, N & Kirschfeld, K. (1971). Étude optique in vivo des éléments photorécepteurs

dans l'oeil compose de *Drosophila*, *Kybernetik*, **8**, 1-13.

Frederiksen, R. & Warrant, E. J. (2008). Visual sensitivity in the crepuscular owl butterfly

Caligo memnon and the diurnal blue morpho *Morpho peleides* a clue to explain the evolution

of nocturnal apposition eyes?, *Journal of Experimental Biology*, **211**, 844-851.

Frentiu, F. D., Bernard, G. D., Cuevas, C. I., Sison-Mangus, M. P., Prudic, K. L. & Briscoe,

A. D. (2007). Adaptive evolution of color vision as seen through the eyes of butterflies,

Proceedings of the National Academy of Sciences, **104**, 8634-8640.

Froy, O., Gotter, A. L., Casselman, A. L. & Reppert, S. M. (2003). Illuminating the circadian

clock in monarch butterfly migration, *Science*, **300**, 1303-1305.

Garamszegi, L. Z., Møller, A. P. & Erritzoe, J. (2002). Coevolving avian eye size and brain

size in relation to prey capture and nocturnality, *Proceedings of the Royal Society B*

Biological Sciences, **269**, 961-967.

Gompert, Z., Willmott, K. & Elias, M. (2011). Heterogeneity in predator micro-habitat use

and the maintenance of Müllerian mimicry diversity, *Journal of Theoretical Biology*, **281**, 39-

46.

Gonzalez-Bellido, P. T., Wardill, T. J. & Juusola, M. (2011). Compound eyes and retinal

information processing in miniature diptean species match their specific ecological demands,

Proceedings of the National Academy of Sciences, **108**, 4224-4229.

Gouy, M., Tannier, E., Comte, N. & Parsons, D. P. (2021). Seaview Version 5: A

multiplatform software for multiple sequence alignment, nolecular phylogenetic analyses,

and tree reconciliation, *Methods in Molecular Biology*, **1198**, 241-260.

Greiner, B. (2006). Adaptations for nocturnal vision in insect apposition eyes, *International*

Review of Cytology, **250**, 1-46.

Greiner, B., Ribi, W. A. & Warrant, E. J. (2004a). Retinal and optical adaptations for

nocturnal vision in the halictid bee *Megalopta genalis*, *Cell and Tissue Research,* **316**, 377-

390.

Greiner, B., Ribi, W. A. & Warrant, E. J. (2005). A neural network to improve dim-light

vision? Dendritic fields of first-order interneurons in the nocturnal bee *Megalopta genalis*,

Cell and Tissue Research, **322**, 313-320.

Greiner, B., Ribi, W. A., Wcislo, W. T. & Warrant, E. J. (2004b). Neural organisation in the

first optic ganglion of the nocturnal bee *Megalopta genalis*, *Cell and Tissue Research*, **318**,

429-437.

Grittner, R., Baird, E. & Stöckl, A. (2022). Spatial tuning of translational optic flow

responses in hawkmoths of varying body size, *Journal of Comparative Physiology A*, **208**,

279-296.

Hardcastle, B. J., Omoto, J. J., Kandimalla, P., Nguyen, B-C, M., Keles, M. F., Boyd, N. K.,

Hartenstein, V. & Frye, M. A. (2021). A visual pathway for skylight polarization processing

in *Drosophila*, *eLife*, **10**, Article no. e63225.

Hardie, R. C. (1986). The photoreceptor array of the dipteran retina, *Trends in*

Neurosciences, **9**, 235-244.

Hamanaka, Y., Kinoshita, M., Homberg, U. & Arikawa, K. (2012). Immunocytochemical

localization of amines and GABA in the optic lobe of the butterfly, *Papilio xuthus, PLoS*

One, **7**, Article no. e41109.

Hebberecht, L., Wainwright, J. B., Thompson, C., Kershenbaum, S., McMillan, W. O. &

Montgomery, S. H. (2023). Plasticity and genetic effects contribute to different axes of neural

divergence in a community of mimetic *Heliconius* butterflies, *Journal of Evolutionary*

Biology,**32**, 1116-1132.

Heinze, S., Florman, J., Asokaraj, S., El Jundi, B. & Reppert, S. M. (2013). Anatomical basis

of sun compass navigation II: the neuronal composition of the central complex of the

monarch butterfly, *Journal of Comparative Neurology*, **521**, 267-298.

Heinze, S. & Reppert, S. M. (2012). Anatomical basis of sun compass navigation I: The

general layout of the monarch butterfly brain, *Journal of Comparative Neurology*, **520**, 1599-

1628.

Hill, R. I. (2010). Habitat segregation among mimetic ithomiine butterflies (Nymphalidae),

Evolutionary Ecology, **24**, 273-285.

Hill, R. I. (2021). Convergent flight morphology among Müllerian mimic mutualists,

Evolution, **75**, 2460-2479.

Hoang, D. T., Chernomor, O., von Haeseler, A., Minh, B. Q. & Vinh, L. S. (2018). UFBoot2:

improving the ultrafast bootstrap approximation, *Molecular Biology and Evolution*, **35**, 518-

522.

Hofmann, C. M., O'Quin, K. E., Marshall, N. J., Cronin, T. W., Seehausen, O. & Carleton, K. L. (2009). The eyes have it: Regulatory and structural changes both underlie cichlid visual pigment diversity, *PLOS Biology,* **7**, Article no. e1000266.

Jander, U. & Jander, R. (2002). Allometry and resolution of bee eyes (Apoidea), *Arthropod Structure and Development*, **30**, 179-193.

Kall, L., Krogh, A. & Sonnhammer, E. L. L. (2007). Advantages of combined transmembrane topology and signal peptide prediction – the Phobius web server, *Nucleic Acids Research*, **35**, 429-432.

Kalyaanamoorthy, S., Minh, B. Q., Wong, T. K. F., von Haeseler, A. & Jermiin, L. S. (2017). ModelFinder: fast model selection for accurate phylogenetic estimates, *Nature Methods*, **14**, 587-589.

Kinoshita, M. & Arikawa, K. (2014). Color and polarization vision in foraging *Papilio*, *Journal of Comparative Physiology A*, **200**. 513-526.

Kinoshita, M., Shimohigasshi, M., Tominaga, Y., Arikawa, K. & Homberg, U. (2015). Topographically distinct visual and olfactory inputs to the mushroom body in the swallowtail butterfly, *Papilio xuths*, *Journal of Comparative Neurology*, **523**, 162-182.

Klagges, B. R. E., Heimbeck, G., Godenschwege, T. A., Hofbauer, A., Pflugfelder, G. O., Reifegerste, R., Reisch, D., Schaupp, M., Buchner, S. & Buchner, E. (1996). Invertebrate synapsins: A single gene codes for several isoforms in *Drosophila*, *Journal of Neuroscience*, **16**, 3154-3165.

Kosakovsky Pond, S. L., Murrell, B., Fourment, M., Frost, S. D., Delport, W. & Scheffler, K. (2011). A random effects branch-site model for detecting episodic diversifying selection, *Molecular Biology and Evolution*, **28**, 3033-3043.

Kosakovsky Pond, S. L. K., Poon, A. F. Y., Velazquez, R., Weaver, S., Hepler, N. L., Murrell, B., Shank, S. D., Magalis, B. R., Bouvier, D., Nekrutenko, A., Wisotsky, S., Spielman, S. J., Frost, S. D. W. & Muse, S. V. (2020). HyPhy 2.5 – a customizable platform for evolutionary hypothesis testing using phylogenies, *Molecular Biology and Evolution*, **37**, 295-299.

Kreissl, S., Strasser, C. & Galizia, C. G. (2010). Allatostatin immunoreactivity in the honeybee brain, *Journal of Comparative Neurology*, **518**, 1391-1417.

Kruska, D. C. T. (2005). On the evolutionary significance of encephalization in some eutherian mammals: effects of adaptive radiation, domestication, and fertilization, *Brain, Behaviour and Evolution*, **65**, 73-108.

Labhart, T., Baumann, F. & Bernard, G. D. (2009). Specialized ommatidia of the

polarization-sensitive dorsal rim area in the eye of monarch butterflies have non-functional

reflecting tapeta, *Cell and Tissue Research*, **338**, 391-400.

Labhart, T. & Meyer, E. P. (1999). Detectors for polarized skylight in insects: a survey of

ommatidial specializations in the dorsal rim area of the compound eye, *Microscopy Research*

and Technique, **47**, 368-379.

Land, M. F. (1989). Variations in the structure and design of compound eyes, In: Stavenga,

D. G. & Hardie, R. C. *Facets of Vision* (pp 90-111), Springer, Berlin, Heidelberg.

Land, M. F. (1997). Visual acuity in insects, *Annual Review of Entomology*, **42**, 147-177.

Land, M. F. & Nilsson, D. E. (2012). *Animal Eyes*, Oxford University Press, Oxford.

Laughlin, S. B., de Rutyer van Steveninck, R. R. & Anderson, J. C. (1998). The metabolic

cost of neural information, *Nature Neuroscience,* **1**, 36-41.

Lehrer, M. (1998). Looking all around: honeybees use different cues in different eye regions,

Journal of Experimental Biology, **201**, 3275-3292.

Lin, C., Hoving, H-J, T., Cronin, T. W. & Osborn, K. J. (2021). Strange eyes, stranger brains:

exceptional diversity of optic lobe organization in midwater crustaceans, *Proceedings of the*

Royal Society B-Biological Sciences, **288**, Article no. 20210216.

Lin, T-Y., Luo, J., Shinomiya, K., Ting, C-Y., Lu, Z., Meinertzhagen, I. A., Lee, C-H.

(2015). Mapping chromatic pathways in the *Drosophila* visual system, *Journal of*

Comparative Physiology, **524**, 213-227.

Lovick, J. K., Omoto, J. J., Ngo, K. T. & Hartenstein, V. (2017). Development of the anterior

visual input pathway to the *Drosophila* central complex*, Journal of Comparative Neurology*,

525, 3458-3475.

Lythgoe, J. N. (1979). *The Ecology of Vision*, Oxford: Clarendon. 244 pp.

Matsushita, A., Stewart, F., Ilić, M., Chen, P-J., Wakita, D., Miyazaki, N., Murata, K.,

Kinoshita, M., Belušič, G. & Arikawa, K. (2022). Connectome of the lamina reveals the

circuit for early color processing in the visual pathway of a butterfly, *Current Biology*, **32**,

2291-2299.

McClure, M., Mahrouche, L., Houssin, C., Monllor, M., Le Poul, Y., Frèrot, B., Furtos, A.,

Elias, M. (2019). Does divergent selection predict the evolution of mate preference and

reproductive isolation in the tropical butterfly genus *Melinaea* (Nymphalidae: Ithomiini)?,

Journal of Animal Ecology, **88**, 940-952.

Mappes, M. & Homberg, U. (2004). Behavioral analysis of polarization vision in tethered

flying locusts, *Journal of Comparative Physiology A*, **190**, 61-68.

McCulloch, K. J., Yuan, F., Zhen, Y., Aardema, M. L., Smith, G., Llorente-Bousquets, J., Andolfatto, P. & Briscoe, A. D. (2017). Sexual dimorphism and retinal mosaic diversification following the evolution of a violet receptor in butterflies, *Molecular Biology and Evolution*, **34**, 2271-2284.

Meyer, E. P. & Labhart, T. (1992). Morphological specializations of dorsal rim ommatidia in the compound eye of dragonflies and damselflies (Odonata), *Cell and Tissue Research*, **272**, 17-22.

Meyer-Rochow, V. B. & Lau, T. F. (2008). Sexual dimorphism in the compound eye of the moth *Operophtera brumata* (Lepidoptera, Geometridae), *Invertebrate Biology*, **127**, 201-216.

Montgomery, S. H., Merrill, R. M. & Ott, S. R. (2016). Brain composition in *Heliconius* butterflies, posteclosion growth and experience-dependent neuropil plasticity, *Journal of Comparative Neurology*, **524**, 1747-1769.

Montgomery, S. H., Mundy, N. I. & Barton, R. A. (2016). Brain evolution and development: adaptation, allometry and constraint, *Proceedings of the Royal Society B Biological Sciences*, **283**, Article no. 20160433.

Montgomery, S. H. & Ott, S. R. (2015). Brain composition in *Godyris zavaleta*, a dirurnal butterfly, reflects an increased reliance on olfactory information, *Journal of Comparative Neurology*, **523**, 869-891.

Montgomery, S. H., Rossi, M., McMillan, W. O. & Merrill, R. M. (2021). Neural divergence and hybrid disruption between ecologically isolated *Heliconius* butterflies, *Proceedings of the National Academy of Sciences of the United States of America,* **118**, Article no. e2015102118.

Moran, D., Softley, R. & Warrant, E. J. (2015). The energetic cost of vision and the evolution of eyeless Mexican cavefish, *Science Advances*, **1**, Article no. e1500363.

Morris, B. J., Couto, A., Aydin, A. & Montgomery, S. H. (2021). Re-emergence and diversification of a specialized antennal lobe morphology in ithomiine butterflies, *Evolution*, **75**, 3191-3202.

Müller, F. (1879). *Ituna* and *Thyridia;* A remarkable case of mimicry in butterflies, *Transactions of the Entomological Society London*, **1879**, xx-xxix.

Narendra, A., Alkaladi, A., Raderschall, C. A., Robson, S. K. A. & Ribi, W. A. (2013). Compound eye adaptations for diurnal and nocturnal lifestyle in the intertidal ant, *Polyrhachis sokolova*, *PLoS ONE*, **8**, Article no. e76015.

Nilsson, D. E., Labhart, T. & Meyer, E. P. (1987). Photoreceptor design and optical

properties affecting polarization sensitivity in ants and crickets, *Journal of Comparative*

Physiology A, **161**, 491-501.

Nilsson, D-E., Land, M. F. & Howard, J. (1988). Optics of the butterfly eye, *Journal of*

Comaprative Physiology A, **162**, 341-366.

Nilsson, D. E. & Smolka, J. (2021). Quantifying biologically essential aspects of

environmental light, *Journal of the Royal Society Interface*, **18**, Article no. 20210184.

Nilsson, D-E., Smolka, J. & Bok, M. (2022). The vertical light-gradient and its potential

impact on animal distribution and behavior, *Frontiers in Ecology and Evolution*, **10**, Article

no. 951328.

Niven, J. E., Anderson, J. C. & Laughlin, S. B. (2007). Fly photoreceptors demonstrate

energy-information trade-offs in neural coding, *PLoS Biology*, **5**, Article no. e116.

Omasits, U., Ahrens, C. H., Müller, S. & Wollscheid, B. (2014). Protter: interactive protein

feature visualization and integration with experimental proteomic data, *Bioinformatics*, **30**,

884-886.

Omoto, J. J., Keles, M. F., Nguyen, B. M., Bolanos, C., Lovick, J. K., Frye, M. A. &

Hartenstein, V. (2017). Visual input to the *Drosophila* central complex by developmentally

and functionally distinct neuronal populations, *Current Biology*, **404**, 2-20.

Osorio, D., Averof, M. & Bacon, J. P. (1995). Arthropod evolution: great brains, beautiful

bodies, *Trends in Ecology and Evolution*, **10**, 449-454.

Ott, S. R. (2008). Confocal microscopy in large insect brains: Zinc-formaldehyde fixation

improves synapsin immunostaining and preservation of morphology in whole-mounts,

Journal of Neuroscience Methods, **172**, 220-230.

Palermo, N. & Theobald, J. (2019). Fruit flies increase attention to their frontal visual field

during fast forward optic flow, *Biology Letters*, **15**, Article no. 20180767.

Pliske, T E. (1975). Courtship behaviour and use of chemical communication by males of

certain species of ithomiine butterflies (Nymphalidae – Lepidoptera), *Annals of the*

Entomological Society of America, **68**, 935-942.

Qiu, X., Vanhoutte, K. J. A., Stavenga, D. G. & Arikawa, K. (2002). Ommatidial

heterogeneity in the compound eye of the male small white, *Pieris rapae crucivora*, *Cell and*

Tissue Research, **307**, 371-379.

R Core Team (2016). R: A language and environment for statistical computing: R Foundation

for Statistical Computing, Vienna, Austria. URL: https://www.R-project.org/

Ranwez, V., Douzery, E. J. P., Cambon, C., Chantret, N. & Delsuc, F. (2018). MACSE v2: toolkit for the alignment of coding sequences accounting for frameshifts and stop codons, *Molecular Biology and Evolution*, **35**, 2582-2584.

Reppert, S. M., Zhu, H. & White, R. H. (2004). Polarized light helps monarch butterflies navigate, *Current Biology*, **14**, 155-158.

Rutowski, R. L. (2000). Variation of eye size in butterflies: inter- and intraspecific patterns, *Journal of Zoology*, **252**, 187-195.

Rutowski, R. L., Gislén, L. & Warrant, E. J. (2009). Visual acuity and sensitivity increase allometrically with body size in butterflies, *Arthropod Structure and Development,* **38**, 91-100.

Sauman, I., Briscoe, A. D., Zhu, H., Shi, D., Froy, O., Stalleicken, J., Yuan, Q., Casselman, A. & Reppert, S. M. (2005). Connecting the navigational clock to sun compass input in monarch butterfly brain, *Neuron*, **46**, 457-467.

Scales, J. A. & Butler, M. A. (2016). The relationship between microhabitat use, allometry and functional variation in the eyes of Hawaiian *Megalagrion* damselfies, *Functional Ecology*, **30**, 356-368.

Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T., Preibisch, S., Rueden, C., Saalfeld, S., Schmid, B., Tinevez, J-Y., White, D. J., Hartenstein, V., Eliceiri, K., Tomancak, P. & Cardona, A. (2012). Fiji: an opec-source platform for biological-image analysis, *Nature Methods*, **9**, 676-682.

Seymoure, B. M., McMillan, W. O. & Rutowski, R. (2015). Peripheral eye dimensions in Longwing (*Heliconius*) butterflies vary with body size and sex but not light environment nor mimicry ring, *The Journal of Research on the Lepidoptera*, **48**, 83-92.

Sivasubramanian, P & Sood, P. P. (2003). Allatostatin-like immunoreactivity in the optic lobe of the fly *Sarophaga bullata*, *Cellular and Molecular Biology*, **49**, 641-644.

Sinakevitch, I., Douglass, J. K., Scholtz, G., Loesel, R. & Strausfeld, N. J. (2003). Consevred and convergent organization in the optic lobes of insects and isopods, with reference to other crustacean taxa, *Journal of Comparative Neurology*, **467**, 150-172.

Slater, G. S. C. & Birney, E. (2005). Automated generation of heuristics for biological sequence comparison, *BMC Bioinformatics*, **6**, 1-11.

Smith, D. B., Bernhardt, G., Raine, N. E., Abel, R. L., Sykes, D., Ahmned, F., Pedroso, I. & Gill, R. J. (2016). Exploring miniature insect brains using micro-CT scanning techniques, *Scientific Reports*, **6**, Article no. 21768.

Sondhi, Y., Ellis, E. A., Bybee, S. M., Theobald, J. C. & Kawahara, A. Y. (2021). Light

environment drives evolution of color vision genes in butterflies and moths, *Communications*

Biology, **4**, Article no. 177.

Somanathan, H., Kelber, A., Borges, R. M., Wallén, R. & Warrant, E. J. (2009). Visual

ecology of Indian carpenter bees II: adaptations of eyes and ocelli to nocturnal and diurnal

lifestyles, *Journal of Comparative Physiology A*, **195**, 571-583.

Stalleicken, J., Labhart, T., Mouritsen, H. (2006). Physiological characterization of the

compound eye in monarch butterflies with focus on the dorsal rim area, *Journal of*

Comparative Physiology A, **192**, 321-331.

Stansbury, M. S. & Moczek, A. P. (2013). The evolvability of arthropods. In *Arthropod*

biology and evolution: Molecules, development, morphology (pp. 479-493). Berlin

Heidelberg: Springer-Verlag.

Stavenga, D. G. (1979). Pseudopupils of compound eyes. In: Autrum H (ed) *Handbook of*

sensory physiology, vol VII/6A. Springer, Berlin Heidelberg, New York, pp 357-439.

Stavenga, D. G. (1992). Eye regionalization and spectral tuning of retinal pigments in insects,

Trends in Neurosciences, **15**, 213-218.

Stavenga, D. G. (2002). Reflections on colourful ommatidia of butterfly eyes, *Journal of*

Experimental Biology, **205**, 1077-1085.

Stavenga, D. G. & Arikawa, K. (2006). Evolution of color and vision of butterflies,

Arthropod Structure and Development, **35**, 307-318.

Stavenga, D. G., Kinoshita, M., Yang, E. C. & Arikawa, K. (2001). Retinal regionalization

and heterogeneity of butterfly eyes, *Naturwissenschaften*, **88**, 477-481.

Stavenga, D. G., Numan, J. A J., Tinbergen, J. & Kuiper, J. W. (1977). Insect pupil

mechanisms. II. Pigment migration in retinula cells of butterflies, *Journal of Comparative*

Physiology A, **113**, 73-93.

Sterling, P. & Laughlin, S. (2015). *Principles of Neural Design*, MIT Press, Cambridge, MA.

Stöckl, A. (2022). Connectomics: From form to colour vision, *Current Biology*, **32**, 457-481.

St□ckl, A. L., Heinze, S., Charalabidis, A., El Jundi, B., Warrant, E. & Kelber, A. (2016a).

Differential investment in visual and olfactory brain areas reflects behavioural choices in

hawk moths, *Scientific Reports*, **6**, 1-10.

Stöckl, A. L., O'Carroll, D. C. & Warrant, E. J. (2020). Hawkmoth lamina monopolar cells

act as dynamic spatial filters to optimize vision at different light levels, *Science Advances*, **6**,

Article no. eaaz8645.

Stöckl, A. L., Ribi, W. A. & Warrant, E. J. (2016b). Adaptations for nocturnal and diurnal vision in the hawkmoth lamina, *Journal of Comparative Neurology,* **524**, 160-175.

Strausfeld, N. J. (2009). Brain organization and the origin of insects: an assessment, *Proceedings of the Royal Society B-Biological Sciences*, **276**, 1929-1937.

Strausfeld, N. J. & Nässel, D. R. (1980). Neuroarchitecture of brain regions that subserve the compound eyes of crustacea and insects, 102-112, In *Autrum H (eds) Handbook of sensory physiology*, Berlin, Springer-Verlag, 200p.

Sugawara, T., Terai, Y., Imai, H., Turner, G. F., Koblmuller, S., Sturmbauer, C., Shicada, Y. & Okada, N. (2005). Parallelism of amino acid changes at the RH1 affecting spectral sensitivity among deep-water cichlids from lakes Tanganyika and Malawi, *Proceedings of the Nation Academy of Sciences of the United States of America,* **102**, 5448-5453.

Swart, P., Wicklein, M., Sykes, D., Ahmed, F. & Krapp, H. G. (2016). A quantitative comparison of micro-CT preparations in Dipteran flies, *Scientific Reports*, **6**, Article no. 39380.

Sylvester, J. B., Pottin, K., Streelman, J. T. (2011). Integrated brain diversification along the early neuraxes, *Brain, Behaviour and Evolution*, **78**, 237-247.

Terakita, A. (2005). The opsins, *Genome Biology*, **6**, 213.

Théry, M. (2001). Forest light and its influence on habitat selection¸ *Plant Ecology*, **153**, 251-261.

Trifinopoulos, J., Nguyen, L-T., von Haeseler, A. & Minh, B. Q. (2016). Q-IQ-TREE: a fast online phylogenetic tool for maximum likelihood analysis, *Nucleic Acids Research*, **44**, 232-235.

Turlure, C., Schtickzelle, N., Van Dyck, H., Seymoure, B. & Rutowski, R. (2016). Flight morphology, compound eye structure and dispersal in the bog and the cranberry fritillary butterflies: an inter- and intraspecific comparison, *PLoS One*, **11**, Article no. e0158073.

Wainwright, J. B. & Montgomery, S. H. (2022). Neuroanatomical shifts mirror patterns of ecological divergence in three diverse clades of mimetic butterflies, *Evolution*, **76**, 1806-1820.

Wainwright, J. B., Schofield, C., Conway, M., Phillips, D., Martin-Silverstone, E., Brodrick, E. A., Cicconardi, F., How, M. J., Roberts, N. W. & Montgomery, S. H. (2023). Multiple axes of visual system diversity in Ithomiini, an ecologically diverse tribe of mimetic butterflies – Ophthalmoscope video recordings and microCT scansm, Zenodo Digital Repository, (https://doi.org/10.5281/zenodo.8154379).

Warrant, E. J. (2001). The design of compound eyes and the illumination of natural habitats,

In: Barth, F. G. & Schmid, A. *Ecology of Sensing* (pp 187-213), Springer, Berlin, Heidelberg.

Warrant, E. J. (2017). The remarkable visual capacities of nocturnal insects: vision at the

limits with small eyes and tiny brains, *Philosophical Transactions of the Royal Society B*

Biological Sciences, **372**, Article no. 20160063.

Warrant, E. J., Kelber, A., Gislén, A., Greiner, B., Ribi, W. & Wcislo, W. T. (2004).

Nocturnal vision and landmark orientation in a tropical halictid bee, *Current Biology,* **14**,

1309-1318.

Warrant, E. J. & McIntyre, P. D. (1993). Arthropod eye design and the physical limits to

spatial resolving power, *Progress in Neurobiology*, **40**, 413-461.

Warton, D. I., Duursma, R. A., Falster, D. S. & Taskinen, S. (2012). smatr 3 – an R package

for estimation and inference about allometric lines, *Methods in Ecology and Evolution*, **3**,

257-259.

Waterhouse, A., Bertoni, M., Bienert, S., Studer, G., Tauriello, G., Gumienny, R., Heer, F. T.,

de Beer, T. A. P., Rempfer, C., Bordoli, L., Lepore, R. & Schwede, T. (2018). SWISS-

MODEL: homology modelling of protein structures and complexes, *Nucleic Acids Research*,

46, 296-303.

Wertheim, J. O., Murrell, B., Smith, M. D., Kosakovsky Pond, S. L. & Scheffler, K. (2015).

RELAX: detecting relaxed selection in a phylogenetic framework, *Molecular Biology and*

Evolution, **32**, 820-832.

White, R. H., Xu, H., Münch, T. A., Bennett, R. R. & Grable, E. A. (2003). The retina of

Manduca sexta: rhodopsin expression, the mosaic of green-, blue- and UV- sensitive

photoreceptors, and regional specialization, *Journal of Experimental Biology*, **206**, 3337-

3348.

Willmott, K. R. & Mallet, J. (2004). Correlations between adult mimicry and larval host

plants in ithomiine butterflies, *Proceedings of the Royal Society B Biological Sciences*, **271**,

266-269.

Willmott, K. R., Willmott, J. C. R., Elias, M. & Jiggins, C. D. (2017). Maintaining mimicry

diversity: optimal warning colour patterns differ among microhabitats in Amazonian

clearwing butterflies, *Proceedings of the Royal Society B-Biological Sciences*, **284**, Article

no. 20170744.

Yack, J. E., Johnson, S. E., Brown, S. G. & Warrant, E. J. (2007). The eyes of *Macrosoma sp.*

(Lepidoptera: Hedyloidea): a nocturnal butterfly with superposition optics, *Arthropod*

Structure & Development, **36**, 11-22.

Yilmaz, A., Hempel de Ibarra, N. & Kelber, A. (2022). High diversity of arthropod colour

vision: from genes to ecology, *Philosophical Transactions of the Royal Society B Biological*

Sciences, **377**, Article no. 20210273.

Zeil, J. (1983). Sexual dimorphism in the visual system of flies: The compound eyes and

neural superposition in Bibionidae (Diptera), *Journal of Comparative Physiology A*, **150**,

379-393.

Zhao, H., Ru, B., Teeling, E. C., Faulkes, C. G., Zhang, S. & Rossiter, S. J. (2009).

Rhodopsin molecular evolution in mammals inhabiting low light environments, *PLoS One,* **4**,

Article no. e8326.

Zhan, S. & Reppert, S. M. (2012). MonarchBase: the monarch butterfly genome database,

Nucleic Acids Research, **41**, 758-763.

Ziemba, K. S. & Rutowski, R. L. (2000). Sexual dimorphism in eye morphology in a

butterfly (*Asterocampa leilia*, Lepidoptera, Nymphalidae), *Psyche*, **103**, 25-36.

Zufall, F., Schmitt, M. & Menzel, R. (1989). Spectral and polarized light sensitivity of

photoreceptors in the compound eye of the cricket (*Gryllus bimaculatus*), *Journal of*

Comparative Physiology A, **164**, 597-608.

1292

1293   **Figure legends**

1294

1295   **Fig. 1**: **An overview of butterfly visual systems and ecology of the four ithomiine study**

1296   **species.** (a) Schematic diagram of a butterfly head representing the different levels of sensory

1297   organisation within the visual system. (b) Microhabitat segregation in ithomiines coupled

1298   with a pruned molecular phylogeny calibrated by Chazot *et al.* (2019) to show the

1299   evolutionary relationships between the four study species.

1300

Fig. 2: Signatures of relaxed selection in ithomiine UV visual opsins. (a) Maximum

likelihood nucleotide gene tree created using IQ-TREE which includes all UV opsin

sequences used in our analyses. Ithomiine branches are highlighted in blue for which the

ultrafast bootstrap values calculated from 10000 replicates are displayed at the nodes. (b)

Comparison of ω distributions between the *Greta morgane* UV test branch (turquoise) and all

other reference branches under the alternative model of relaxed selection. The branch tips

leading to the duplicated UV2 opsins of *Heliconius melpomene*and *Tithorea harmonia* were

left as unspecified in the RELAX analysis. (c) Comparison of ω distributions from the newly

discovered UV2 opsin of *Tithorea harmonia* under the alternative relaxed selection model

when compared to all other reference branches. Branches leading to the *Greta morgana* UV

and *Heliconius melpomene* UV2 opsins were left unspecified. In both (b) and (c), sites under

purifying ($\omega > 1$) or positive ($\omega < 1$) selection in the reference move towards neutrality in the

test branch ($\omega = 1$). Arrows indicate that when comparing sites in the prespecified test and

reference branches, all three ω distributions used in the BS-REL model (corresponding to

sites under strong purifying, weakly purifying, and positive selection) shift closer towards

neutrality in the test branches. Superimposed on both (b) and (c) is a 3D protein model of the

Greta morgane UV and *Tithorea harmonia* UV opsin respectively, created using Swiss-

Model (Waterhouse *et al.* 2018) with the jumping spider rhodopsin-1 as the template.

Prediction of alpha helices are shown in different colours and areas between the top and

bottom dotted layers indicate transmembrane predictions.

1321

Fig. 3: Ithomiine eyeshine variation (n = 48, 54.17% female)**.** (a) Ophthalmoscope images

of the dorsal, frontal, and ventral luminous pseudopupil within a single individual of all four

study species after ten minutes of dark adaptation. (b) Interspecific differences in frontal

pupillary response time. (c) Frontal pupillary response time after different lengths of dark

adaptation. Points show means for each species (n = 12 per species, 50-58% female) with

error bars representing standard deviation. (d) Intraindividual differences in pupillary

response time for the dorsal, frontal, and ventral eye regions. For all boxplots, medians (thick

horizontal bars), interquartile ranges (boxes), values within 1.5 interquartile ranges of the box

edges (whiskers) and possible outliers (data points outside the whiskers) are plotted.

1331

Fig. 4: An eye and brain anatomical overview for the ithomiine species using micro-CT

data (n = 23, 52.17% female). (a) Volume rendered whole-head images from micro-CT scans

of *Methona confusa* from the anterior (i) and posterior (ii) with surface reconstructions of the

eye and the four primary optic neuropils of interest superimposed on the right. (b) Anterior (i)

and posterior (ii) surface reconstructions of the eye (EYE), lamina (LAM), medulla (ME),

lobula (LOB) and lobula plate (LOP) of *Methona confusa* (left) and *Greta morgane* (right),

shown to scale. (c) Iodine-stained micro-CT X-ray tomography frontal sections of *Methona*

confusa taken at progressively posterior positions through the head, moving from left to right.

Additional labels are the cornea, the areas containing crystalline cones (CC) and

photoreceptor cells (PRC) and an exemplification of our measure of interocular distance

(*IOD*). (d) Further exemplifications of our measures of facet diameter (*D*), ommatidial length

(*L*) and interommatidial angle (*ΔΦ*) from sections of the same *Methona confusa* individual.

Scale bars = 200 μm.

1345

Fig. 5: Eye anatomical variation in four ithomiine species from micro-CT measurements

(n = 23, 52.17% female)**.** (a-c) Interspecific differences in total eye volume (EYE, a),

ommatidial length (*L*, b) and facet diameter (*D,* c). (d-f) Intersexual variation in EYE (d), *L*

(e) and *D* (f). For EYE and *L*, log-transformed values are scaled against interocular distance

(*IOD*) where an "α shift" denotes a grade shift in the relationship between these two variables

from our SMATR analysis. (g-i) Intraindividual differences in interommatidial angle (*ΔΦ,* g),

L (h) and *D* (i) between the dorsal, frontal, and ventral eye regions. For all boxplots, medians

(thick horizontal bars), interquartile ranges (boxes), values within 1.5 interquartile ranges of

the box edges (whiskers) and possible outliers (data points outside the whiskers) are plotted.

NS P > 0.05, *P < 0.05, **P < 0.01, ***P < 0.001.

1356

Fig. 6: Detailed neuroanatomy from immunostained brain samples (anti-allatostatin

labelling: n = 24, 54.17% female; dextran injections: n = 24, 62.5% male). (a) Confocal

images of anti-allatostatin labelled frontal vibratome sections (thickness, 80 μm) taken at

approximately equivalent regions of the brain of *Greta morgane* (i), *Mechanitis polymnia* (ii),

Tithorea harmonia (iii), and *Methona confusa* (iv). Divisions in the medulla (ME), lobula

plate (LOP) and lobula (LOB) are shown using thick black lines and finer-scale layering

within the medulla is shown in orange. (b-c) Sexual dimorphism in the presence/absence of

the ventral lobula, a butterfly-specific optic neuropil. (b) Confocal sections of anti-synapsin

labelled wholemounts of a male (i) and female (ii) brain of *Mechanitis polymnia*. (c)

Confocal section of a single female *Greta morgane* brain, co-labelled against synapsin

(green) and dextran injected neurons (magenta) originating from the medial optic lobe (i) and

terminating in the bulb of the central brain (ii). All scale bars = 100 μm. (LAM: lamina;

LAMc: lamina cartridges; ME: medulla; MEc: medulla columns; iME: inner medulla; oME:

outer medulla; LOP: lobula plate; iLOP: inner lobula plate; oLOP: outer lobula plate; LOB:

lobula; iLOB: inner lobula; mLOB: middle lobula; oLOB: outer lobula; vLOB: ventral

lobula; AOTU: anterior optic tubercle; BU: bulb; CB: central body; MB-pe: mushroom body

peduncle).

1374

Fig. 7: Neuroanatomical variation in four ithomiine species from micro-CT

measurements (n = 23, 52.17% female)**.** (a) Posterior surface reconstructions of the four

main optic lobe neuropils (LAM: lamina, ME: medulla, LOP: lobula plate, LOB: lobula) for

all four species shown to scale. Scale bar = 200 μm. (b) Interspecific scaling relationships

between interocular distance (IOD) and the lamina (i), medulla (ii), lobula plate (iii), and

lobula (iv). An "α shift" denotes a grade shift in the relationship between the two variables

from our SMATR analysis. NS P > 0.05, *P < 0.05, **P < 0.01, ***P < 0.001.

a)

Visual information →

1. Facet
2. Ommatidia
3. Lamina
4. Medulla
5. Lobula plate
6. Lobula
7. Central brain

Perception — Primary visual processing — Sensory integration

b)

Species

Mechanitis polymnia

Greta morgane

Methona confusa

Tithorea harmonia

2 my

Microhabitat

Patchy/open forest

Shaded understory

(a)

Agraulis vanillae EU449018 UV
Dione juno HM366553 UV
Heliconius erato MF035588 UV2
Heliconius melpomene GQ451902 UV2
Heliconius erato MF035587 UV1
Heliconius melpomene GQ451901 UV1
Eueides isabella MF035500 UV
Dryas iulia GU324674 UV
Speyeria mormonia EU449019 UV
Limenitis weidemeyerii EU449015 UV
Limenitis archippus EU449016 UV
Greta morgane UV
64
Methona confusa UV
56
Tithorea harmonia UV1
98 100
Tithorea harmonia UV2
Mechanitis polymnia UV
Danaus plexippus AAU07979 UV
Danaus gilippus EU449017 UV
Coenonympha tullia EU449020 UV
Euphydryas chalcedona EU449014 UV
Vanessa cardui AF414074 UV

0.1

(b)

Greta morgane UV opsin
Reference branches

Greta morgane UV opsin

ω

(c)

Tithorea harmonia UV2 opsin
Reference branches

Tithorea harmonia UV2 opsin

ω

(a)

LAM
ME
LOP
LOB

Greta morgane

Mechanitis polymnia

Tithorea harmonia

Methona confusa

(b)

(i)

Species
● *Greta morgane*
● *Mechanitis polymnia*
● *Tithorea harmonia*
● *Methona confusa*

$\log_{10}$(lamina volume)

$\log_{10}$(IOD)

*** alpha shift

(ii)

$\log_{10}$(medulla volume)

$\log_{10}$(IOD)

*** alpha shift

(iii)

$\log_{10}$(lobula plate volume)

$\log_{10}$(IOD)

*** alpha shift

(iv)

$\log_{10}$(lobula volume)

$\log_{10}$(IOD)

*** alpha shift