Physiological and metabolic insights into the first cultured anaerobic

representative of deep-sea *Planctomycetes* bacteria

Rikuan Zheng[1,2,4], Chong Wang[1,2,4], Rui Liu[1,2,4], Ruining Cai[1,2,3,4], Chaomin Sun[1,2,3,4*]

[1]CAS and Shandong Province Key Laboratory of Experimental Marine Biology &

Center of Deep Sea Research, Institute of Oceanology, Chinese Academy of

Sciences, Qingdao, China

[2]Laboratory for Marine Biology and Biotechnology, Qingdao National Laboratory

for Marine Science and Technology, Qingdao, China

[3]College of Earth Science, University of Chinese Academy of Sciences, Beijing,

China

[4]Center of Ocean Mega-Science, Chinese Academy of Sciences, Qingdao, China

[*]Corresponding author

Chaomin Sun   Tel.: +86 532 82898857; fax: +86 532 82898857.

14   E-mail address: sunchaomin@qdio.ac.cn

15   **Key words:** *Planctomycetes*, deep sea, cultivation, nitrogen metabolism,

16   bacteriophages

17   **Running title:** A novel deep-sea *Planctomycetes* bacterium

Abstract

Planctomycetes bacteria are ubiquitously distributed across various biospheres and play key roles in global element cycles. However, few deep-sea *Planctomycetes* members have been cultivated, limiting our understanding of *Planctomycetes* in the deep biosphere. Here, we have successfully cultured a novel strain of *Planctomycetes* (strain ZRK32) from a cold seep sediment, by using an enriched medium supplemented with rifampicin and different sources of nitrogen. Our genomic, physiological and phylogenetic analyses indicate that strain ZRK32 is a novel species, which we propose be named: *Poriferisphaera heterotrophicis*. We show that strain ZRK32 replicates using a budding mode of division. Based on the combined results from growth assays and transcriptomic analyses, we found that rich nutrients, or supplementation with NO_3^- or NH_4^+ promoted the growth of strain ZRK32 by facilitating energy production through the tricarboxylic acid (TCA) cycle and the Embden–Meyerhof–Parnas (EMP) glycolysis pathway. Moreover, supplementation with NO_3^- or NH_4^+ induced strain ZRK32 to release a bacteriophage in a chronic manner, without host cell lysis. This bacteriophage then enabled strain ZRK32, and another marine bacterium that we studied, to metabolize nitrogen through the function of auxiliary metabolic genes (AMGs). Overall, these findings expand our understanding of deep-sea *Planctomycetes* bacteria, while highlighting their ability to metabolize nitrogen when reprogrammed by chronic viruses.

Introduction

Planctomycetes bacteria are ubiquitous in many environments, including lakes (Pollet et al., 2011), wetlands (Dedysh and Ivanova, 2019), soil (Buckley et al., 2006), freshwater (Brümmer et al., 2004), oceanic waters, and abyssal sediments (Woebken et al., 2007; Goffredi and Orphan, 2010), where they are critical for carbon and nitrogen cycling; although, the specific pathways used are unknown (Wiegand et al., 2018). Although *Planctomycetes* bacteria are highly abundant in nature, relatively few have been cultivated; therefore, the unexplored groups lack cultured and characterized representatives (Fuerst and Sagulenko, 2011). Currently, all taxonomically described *Planctomycetes* with validly published names can be divided into two recognized classes: *Planctomycetia* and *Phycisphaerae* (Fukunaga et al., 2009; Wiegand et al., 2018). *Candidatus* Brocadiae is considered a third class within the phylum *Planctomycetes*, although pure cultures are not yet available for its members and its class-level status remains to be defined (Kartal et al., 2012; Kartal et al., 2013). The majority of the species isolated from the phylum *Planctomycetes* are from the class, *Planctomycetia* — a class that consists of four orders: *Planctomycetales*, *Pirellulales*, *Gemmatales*, and *Isosphaerales* (Dedysh et al., 2020). In contrast, only few members from the *Phycisphaerae* class have been cultured; the class *Phycisphaerae* contains three orders: *Phycisphaerales* (Fukunaga et al., 2009), *Tepidisphaerales* (Kovaleva et al., 2015), and *Sedimentisphaerales* (Spring et al., 2018).

Planctomycetes bacteria have fascinating physiological characteristics (Wiegand et al., 2018). For decades, *Planctomycetes* bacteria have blurred the lines between prokaryotes and eukaryotes. *Planctomycetes* bacteria possess several uncommon traits when compared with typical bacteria: they have a compartmentalized cell plan, an enlarged periplasm, a tightly folded nucleus-like structure, an endocytosis-like method of uptake, and a FtsZ-free method of cell division (Fuerst and Webb, 1991; Lindsay et al., 1997; Lonhienne et al., 2010; Wiegand et al., 2018). The unique cellular structures in *Planctomycetes* bacteria have stretched and challenged our understanding of the concept of a 'prokaryote'. *Planctomycetes* bacteria also exhibit a

diverse range of respiration and cell division methods. Members of the class *Planctomycetia* are mostly aerobic and divide by budding, while *Phycisphaerae* members are mostly anaerobic and divide by binary fission (Rivas-Marń et al., 2016; Wiegand et al., 2018; Pradel et al., 2020; Wiegand et al., 2020b). However, a recent report showed that *Poriferisphaera corsica* KS4 — a novel strain from class *Phycisphaerae* — was aerobic and that it might divide by budding (Kallscheuer et al., 2020b); this is different from previous reports and suggests that there are many physiological and cellular *Planctomycetes* bacteria characteristics yet to be discovered.

The deep sea is where life may have originated from and where stepwise evolution occurred (Orgel, 1998); it is also where a large number of uncultured microorganisms live (Zheng et al., 2021b). Among these microbes, *Planctomycetes* might dominate deep-sea sediments (Wiegand et al., 2018). For example, in the Gulf of Mexico, the phylum *Planctomycetes* accounts for 28% of all bacteria, where they seem to be involved in the nitrogen cycle, and the breakdown of organic detrital matter, which is delivered to the sediment as marine snow (Vigneron et al., 2017). Unfortunately, only few *Planctomycetes* bacteria from deep-sea environments have been cultured (Wang et al., 2020; Wiegand et al., 2020b), limiting our understanding of their characteristics (e.g., material metabolism, element cycling, and ecological role) (Kulichevskaya et al., 2020).

Here, we successfully cultured a novel member of *Planctomycetes* (strain ZRK32) from a deep-sea subsurface sediment. We found that strain ZRK32 used a budding mode of division. We also found that supplementation with rich nutrients, and either NO_3^- or NH_4^+, promoted strain ZRK32 growth. Moreover, the presence of NO_3^- or NH_4^+ induced strain ZRK32 to release a bacteriophage in a chronic manner — a process that does not kill the host cell (i.e., the host bacteria continues to grow despite phage reproduction) (Liu et al., 2022). This bacteriophage reprogrammed strain ZRK32, and another marine bacterium we studied, to metabolize nitrogen through the action of auxiliary metabolic genes (AMGs).

Results

Isolation, morphology, and phylogenetic analysis of a novel strain of _Planctomycetes_ isolated from a deep-sea cold seep

We enriched deep-sea _Planctomycetes_ bacteria using a basal medium supplemented with rifampicin and inorganic nitrogen sources ($NaNO_3$ and NH_4Cl). These enriched samples were plated on to agar slants in Hungate tubes, and then colonies with distinct morphologies were selected and cultivated (Figure. 1A). Some colonies were noted to be from the phylum _Planctomycetes_, based on their 16S rRNA sequences. Among these, strain ZRK32 was selected for further study as it grew faster than other strains. Following negative staining and TEM observation, we observed that strain ZRK32 cells were spherical (with an average diameter of 0.4-1.0 μm), and had a single polar flagellum (Figure. 1B). Moreover, we found that the mother and daughter cells of strain ZRK32 had distinct sizes at the stage of cell division (Figure. 1C and Figure. S1), indicating that strain ZRK32 was dividing asymmetrically (i.e., through budding). Ultrathin whole-cell sections showed that strain ZRK32 possessed a condensed and intact nucleoid-like structure, and a complex extended membrane structure (Figure. 1D).

To confirm the taxonomy of strain ZRK32, we performed phylogenetic analyses. The maximum likelihood tree of 16S rRNA indicated that strain ZRK32 was from the genus _Poriferisphaera_ and that it formed an independent phyletic line with the strain _Poriferisphaera corsica_ KS4^T (Figure. 1E). The genome tree also suggested that this novel clade was a sister-strain of strain KS4^T, which belongs to the genus _Poriferisphaera_ (Figure. S2). Based on its genomic (Figure. S3 and Table S1), physiological (Figure. S4) and phylogenetic characteristics, strain ZRK32 was distinguishable from _Poriferisphaera corsica_ KS4^T, which is currently the only species of the genus _Poriferisphaera_ with a validly published name. We therefore propose that strain ZRK32 represents a novel species in the genus _Poriferisphaera_, for which the name _Poriferisphaera heterotrophicis_ sp. nov. is proposed.

Figure. 1. Isolation, morphology, and phylogenetic analysis of *P. heterotrophicis* ZRK32. (A) Diagram showing the strategy used to isolate the *Planctomycetes* bacteria. (B, C) TEM observation of strain ZRK32. (D) TEM observation of ultrathin sections of cells from strain ZRK32. Abbreviations: CM, outer membrane; Pi, cytoplasm; R, ribosome; N, nucleoid; ICM, cytoplasmic membrane; Py, periplasm; V, vesicle-like organelles. (E) Phylogenetic analysis of strain ZRK32. Phylogenetic placement of strain ZRK32 within the phylum *Planctomycetes*, based on almost complete 16S rRNA gene sequences. The NCBI accession number for each 16S rRNA gene is indicated after each corresponding strain's name. The tree was inferred and reconstructed using the maximum likelihood criterion, with bootstrap values (%) > 80; these are indicated at the base of each node with a gray dot (expressed as a percentage from 1,000 replications). The 16S rRNA gene sequence of *Actinoplanes derwentensis* LA107[T] was used as the outgroup. Bar, 0.1 substitutions per nucleotide position.

Rich nutrients promote *P. heterotrophicis* ZRK32 growth

The growth rate of strain ZRK32 increased when it was cultured in a rich medium (containing ten times more yeast extract than basal medium) (Figure. 2A). To gain

further insight into its metabolic characteristics, we performed transcriptomic analyses

of strain ZRK32 grown in the rich medium and strain ZRK32 grown in the basal

medium. The results showed that the expression of many genes involved in the TCA

cycle (Figure. 2B, C) and the EMP glycolysis pathway (Figure. S5A, B) (which both

contribute to energy production) were up-regulated. In addition, the expression of

genes encoding NADH-ubiquinone oxidoreductase and flagellum assembly-related

proteins were also up-regulated (Figure. 2D, E). To verify the transcriptomic data, we

performed qRT-PCR assays, which showed the same gene expression variation,

consistent with the transcriptomic results (Figure. S6). Based on the combined results

of the growth assay and transcriptomic analyses, we concluded that strain ZRK32

growth is better when cultured using nutrient-rich media.

Figure. 2. Growth assay and transcriptomic analysis of *P. heterotrophicis* ZRK32 strains

cultivated in basal medium and rich medium. (A) Growth curves of ZRK32 strains cultivated

in basal medium and rich medium. (B) Diagram of the TCA cycle. The gene numbers shown in this schematic are the same as those shown in panel C. Transcriptomics-based heat map showing the relative expression levels of genes associated with the TCA cycle (C), NADH-quinone oxidoreductase (D), and flagellar assembly (E) of strain ZRK32 cultivated in rich medium (Rich) compared with strain cutivated in basal medium (Basal). The numbers in panels C, D, and E represent the fold change of gene expression (by using the $\log_2$ value).

P. heterotrophicis ZRK32 replicates using a budding mode of division

After reviewing more than 600 TEM photos, we confirmed that strain ZRK32 divided by budding — a method also reported in other *Planctomycetes* bacteria (Figure. 1C and Figure. S1) (Wiegand et al., 2020b). Remarkably, during the early stages of budding in strain ZRK32, the extracellular membrane extended and formed a bulge, which grew until it was a similar size to the mother cell (Figure. 3A, B, panels 1 to 4). The genetic materials within the nucleoid then duplicated and divided equally between the mother and daughter cells, along with other cytoplasmic contents (Figure. 3A, B, panels 5 to 8). The daughter cell then separated from the mother cell, completing cell division.

Next, we investigated whether genes associated with budding method of cell division were present in the genome of strain ZRK32, and whether they were functional during bacterial growth. We performed genomic and transcriptomic analyses of strain ZRK32. We did not detect the cell division protein, FtsZ, in strain ZRK32, but we did identify other Fts-related proteins (e.g., FtsJ, FtsK, FtsL, FtsW, FtsX, and FtsY) and rod shape-determining proteins (MreB, MreC, and MreD); the expression of genes encoding these Fts-related proteins were upregulated in the ZRK32 strains cultured in the rich medium compared with strains cultured in basal medium (Figure. 3C).

Figure. 3. The mode of cell division utilized by *P. heterotrophicis* ZRK32. (A) Ultrathin TEM sections showing the process of polar budding division (panels 1-8) in strain ZRK32. Images representing the different phases of cell division are shown. Scale bars are 200 nm in panels A and B. (B) The proposed model of cell division of strain ZRK32 based on the TEM observation shown in panel B. The numbers in panels A and B correspond to the same phase of division. (C) Transcriptomics-based heat map showing the differentially expressed genes that encode different key proteins associated with cell division in strain ZRK32. The numbers in panel A represent the fold change of gene expression (by using the $\log_2$ value).

Effects of NO_3^-, NH_4^+, and NO_2^- on *P. heterotrophicis* ZRK32 growth

As *Planctomycetes* bacteria are involved in nitrogen cycling, we tested the effects of different nitrogen-containing substances (including NO_3^-, NH_4^+, and NO_2^-) on strain

ZRK32 growth. These assays showed that adding NO_3^- or NH_4^+ to the culture medium

increased strain ZRK32 growth, while adding NO_2^- inhibited growth (Figure. 4A).

The concentration of NO_3^- decreased from ~21 mM to ~6 mM, and then to ~4 mM

after strain ZRK32 had been incubating for four days and six days, respectively. The

concentration of NH_4^+ increased from ~0 mM to ~11 mM, and then to ~7 mM after

strain ZRK32 had been incubating for four days and six days, respectively. These

results strongly suggest that strain ZRK32 can effectively convert NO_3^- to NH_4^+

(Figure. 4B). In addition, when strain ZRK32 was incubated in the rich medium

supplemented with NH_4^+ for four days and six days, the concentration of NH_4^+

decreased from ~19 mM to ~11 mM, and then to ~4 mM, with no change in the

concentrations of NO_3^- and NO_2^- (Figure. 4C). To investigate nitrogen metabolism in

strain ZRK32, we analyzed the strain ZRK32 genome and found that it contained a

complete nitrate reduction pathway and key genes responsible for the conversion of

ammonia to glutamate (Figure. 4D), which explains the results in figures 4B and C.

Subsequently, we performed transcriptome sequencing analysis and found that the

genes encoding nitrate reductase (NapA and NapB), nitrite reductase (NirB),

glutamine synthetase, and glutamate synthase were all simultaneously up-regulated in

the presence of NO_3^-. We also found that the genes encoding glutamine synthetase and

glutamate synthase were up-regulated in the presence of NH_4^+ (Figure. 4E). However,

no differential expression was observed for the gene, *nirD* in the presence of NO_3^- or

NH_4^+, even though this gene encodes a nitrite reductase. Moreover, we observed that

some genes involved in the TCA cycle (Figure. 4F), the EMP glycolysis pathway

(Figure. S5C), and genes encoding NADH-ubiquinone oxidoreductase-related

proteins (Figure. 4G) were upregulated in the presence of NO_3^- or NH_4^+, but

down-regulated in the presence of NO_2^-. We also observed that the expression of

many genes associated with flagellum assembly were up-regulated when NO_3^- or

NH_4^+ were supplemented into the culture medium (Figure. 4H). We observed similar

trends in our qRT-PCR results (Figure. S7), which validated our RNA-seq results.

Figure. 4. Nitrogen metabolism assays of *P. heterotrophicis* ZRK32. (A) Growth curves of ZRK32 strains cultivated in the rich medium alone and cultivated in rich medium supplemented with either 20 mM NO_3^-, 20 mM NH_4^+, or 20 mM NO_2^-. (B) The dynamics of concentrations of NO_3^-, NH_4^+, and NO_2^- in strains of ZRK32 cultivated in the rich medium supplemented with 20 mM NO_3^-. (C) The dynamics of concentrations of NO_3^-, NH_4^+, and NO_2^- in strains of ZRK32 cultivated in the rich medium supplemented with 20 mM NH_4^+. (D) The predicted nitrogen metabolism pathway of strain ZRK32. Abbreviations: NapA, periplasmic nitrate reductase; NapB, periplasmic nitrate reductase, electron transfer subunit; NirB, nitrite reductase (NADH) large subunit; NirD, nitrite reductase (NADH) small subunit; GlnA, glutamine synthetase; GLT1,

glutamate synthase. Transcriptomics-based heat map showing the relative expression levels of
genes associated with nitrogen metabolism (E), the TCA cycle (F), NADH-quinone
oxidoreductase (G), and flagellar assembly (H) in strains of ZRK32 cultivated in the rich medium
supplemented with different inorganic nitrogen sources (20 mM NO_3^-, 20 mM NH_4^+ or 20 mM
NO_2^-) compared with strains cultivated in the rich medium alone. "Rich" indicates rich medium.
"NO_3^-, NH_4^+, and NO_2^-" indicate rich medium supplemented with 20 mM NO_3^-, 20 mM NH_4^+, and
20 mM NO_2^-, respectively. The numbers in panels E, F, G, and H represent the fold change of gene
expression (by using the $\log_2$ value).

NO_3^- and NH_4^+ induce the release of a chronic bacteriophage in *P. heterotrophicis* strain ZRK32

Bacteriophages are widely distributed across oceans and can regulate nitrogen

metabolism in their host (Cassman et al., 2012; Monier et al., 2017; Gazitúa et al.,

2021; Wang et al., 2022). We therefore investigated whether bacteriophages affected

nitrogen metabolism in strain ZRK32. TEM observations showed that phage-like

structures (hexagonal phages, ~30 nm) were present in cell suspensions of the ZRK32

strain that had been cultured using nutrient-rich medium supplemented with either

NO_3^- or NH_4^+ (Figure. 5A, panels II and III). In contrast, no phage-like structures

were observed in the cell suspensions of ZRK32 strain that was cultured in the rich

medium alone (without NO_3^- or NH_4^+) (Figure. 5A, panel I). This suggests that the

presence of NO_3^- or NH_4^+ stimulated the release of the bacteriophages from strain

ZRK32. Most chronic bacteriophages do not negatively affect their host's growth

when cultivated in the laboratory (Alarcón-Schumacher et al., 2022). Consistently, the

replication and release of the bacteriophages from strain ZRK32 did not kill the host

cell, consistent well with the key feature of chronic bacteriophages (Howard-Varona

et al., 2017). By comparing genomic sequences, we confirmed that the genome of the

phage induced by NO_3^- was the same as the phage induced by NH_4^+. When we

compared this phage genome (Phage-ZRK32, 21.9 kb) (Figure. 5B) with the host

genome (strain ZRK32) (using Galaxy Version 2.6.0 (https://galaxy.pasteur.fr/)

(Afgan et al., 2018) with the NCBI BLASTN method), we found that the

Phage-ZRK32 genome was outside of the host chromosome; this indicates that this

chronic bacteriophage is extrachromosomal, which is consistent with previous reports

(Chevallereau et al., 2022). In addition to the genes encoding phage-associated

proteins, the genome of Phage-ZRK32 also has many auxiliary metabolic genes
(AMGs), which encode glutamine amidotransferase, amidoligase, glutathione
synthase, and gamma-glutamylcyclotransferase (Figure. 5B). Although, some genes
(including genes encoding amidoligase, glutathione synthase and
gamma-glutamylcyclotransferase) were absent from the strain ZRK32 genome.

Figure. 5. Observation and functional assay of the chronic bacteriophage induced by NO_3^- or
NH_4^+ from *P. heterotrophicis* ZRK32. (A) TEM observation of phages extracted from the cell
suspensions of ZRK32 strains that cultured in either the rich medium alone, or rich medium
supplemented with 20 mM of either NO_3^- or NH_4^+. (A, Panel I) No phage-like particles were
observed in the cell suspensions from the ZRK32 strain cultured in the rich medium. (A, Panels II
and III) Hexagonal phages (indicated with yellow arrows) observed in the cell suspensions from
the ZRK32 strains cultured in the rich medium supplemented with 20 mM of either NO_3^- or NH_4^+.
The scale bar is 200 nm. (B) A diagram showing the genomic composition of Phage-ZRK32. The

arrows represent different ORFs and the direction of transcription. The main putative gene products of this phage are shown, and the numbers in brackets indicate the numbers of amino acids. Hypothetical proteins are indicated by gray arrows, structural modules are indicated by green arrows, nucleotide metabolism is indicated by blue-gray arrows, the replication module is indicated by gold arrows, and AMGs are indicated by red arrows. The size of the phage genome is shown beside the gene cluster. (C) Bacterial growth curve showing the growth rate of strains of *Pseudomonas stutzeri* 273 cultivated in either oligotrophic medium, oligotrophic medium supplemented with Phage-ZRK32, oligotrophic medium supplemented with 20 mM NO_3^-, or oligotrophic medium supplemented with 20 mM NO_3^- and Phage-ZRK32. (D) Bacterial growth curve showing the growth rate of strains of *Pseudomonas stutzeri* 273 cultivated in either oligotrophic medium, oligotrophic medium supplemented with Phage-ZRK32, oligotrophic medium supplemented with 20 mM NH_4^+, or oligotrophic medium supplemented with 20 mM NH_4^+ and Phage-ZRK32. "Oligo" indicates oligotrophic medium.

To verify whether these AMGs were functional, we investigated whether Phage-ZRK32 was capable of reprogramming nitrogen metabolism and promoting growth in other marine bacteria. We selected the aerobic marine bacterium, *Pseudomonas stutzeri* 273 (Wu et al., 2016), and examined the effects of Phage-ZRK32 on its growth. This showed that Phage-ZRK32 promoted *Pseudomonas stutzeri* 273 growth by facilitating the metabolism and utilization of NO_3^- and NH_4^+ (Figure. 5C, D). In particular, adding Phage-ZRK32 and NH_4^+ into the oligotrophic medium resulted in an approximate 3–8-fold increase in growth compared with strains cultivated without NH_4^+ or Phage-ZRK32 supplementation (Figure. 5D). The AMGs that encode amidoligase and gamma-glutamylcyclotransferase were absent from the *Pseudomonas stutzeri* 273 genome, even though it contains a complete nitrate reduction pathway and some genes responsible for converting ammonia to glutamate. Therefore, the Phage-ZRK32 AMGs might facilitate nitrogen metabolism and amino acid generation in *Pseudomonas stutzeri* 273 in a similar way to *P. heterotrophicis* ZRK32.

Discussion

Until recently, most research on *Planctomycetes* has focused on strains in freshwater and shallow ocean environments (Bondoso et al., 2011; Kallscheuer et al., 2020b), with few studies on deep-sea strains; this is likely due to the logistical difficulties associated with sampling and cultivating these strains. The availability of cultured and

characterized representatives for many phylogenetic clades within the phylum *Planctomycetes* are lacking (Dedysh et al., 2021). Therefore, more approaches and media types, like those used in our study (Figure. 1A), should be developed to obtain *Planctomycetes* bacteria from different deep-sea environments. The vast majority of *Planctomycetes* members have been isolated using an oligotrophic medium supplemented with N-acetyl glucosamine (Kaboré et al., 2020; Kallscheuer et al., 2020a; Peeters et al., 2020; Salbreiter et al., 2020; Wiegand et al., 2020a); use of this medium previously resulted in the breakthrough isolation of 79 *Planctomycetes* strains (Wiegand et al., 2020b). However, we observed that strains of *P. heterotrophicis* ZRK32 grew much better when cultivated in a rich medium compared with strains cultivated in the basal medium (Figure. 2A). We also found that *N*-acetyl glucosamine did not stimulate the growth of strain ZRK32. These findings have revealed that strain ZRK32 prefers a nutrient-rich medium, which is different from most other *Planctomycetes* bacteria (Wiegand et al., 2018; Wiegand et al., 2020b). When cultured in a nutrient-rich medium, the expression of genes associated with the TCA cycle and the EMP glycolysis pathway were up-regulated (Figure. 2B-D, Figure. S5B and Figure. S6), suggesting that strain ZRK32 uses and metabolizes nutrients to produce energy, which it then uses to grow (Figure. S8) (Zheng et al., 2021b).

Strain ZRK32 possesses a condensed and intact nucleoid-like structure and a complex membrane structure (Figure. 1D), which is similar to other reported *Planctomycetes* bacteria (Kaushik et al., 2020; Dedysh et al., 2021). In addition, TEM observation of ultrathin sections of cells from strain ZRK32 showed some eukaryote-like structures (Figure. S9, panels 1 to 6). Although it is impossible to judge accurately what these structures are based on current methods, the internal structures of *Planctomycetes* bacteria are nevertheless fascinating. We also observed some vesicle-like (Figure. S9, panels 7 to 8) and vacuole-like structures (Figure. S9, panel 9) in strain ZRK32, similar to those observed in other *Planctomycetes* members (Wiegand et al., 2018). Vacuoles store cellular components (such as proteins and sugars etc.,) and play essential roles in plant responses to different biotic/abiotic signaling pathways (Zhang et al., 2014). The presence of vacuoles in strain ZRK32

suggests that *Planctomycetes* bacteria might have adopted a eukaryotic mechanism for

nutrient metabolism and signal transduction.

Most species within the class *Planctomycetia* divide by budding, and species

within the class *Phycisphaerae* divide by binary fission (Fukunaga et al., 2009; Yoon

et al., 2014; Kovaleva et al., 2015; Kallscheuer et al., 2020a; Pradel et al., 2020;

Salbreiter et al., 2020; Wiegand et al., 2020b). However, the *Poriferisphaera* strain

ZRK32 (Figure. 1C and Figure. S1) was demonstrated to divide by budding, similar to

the *Poriferisphaera* strain KS4 proposed by Kallscheuer et *al* (2020) (Kallscheuer et

al., 2020b), suggesting that members within the class *Phycisphaerae* might divide

through both binary fission and budding. It is noteworthy that *P. heterotrophicis*

ZRK32 forms an empty cell framework first, followed by the entry of the cellular

contents (Figure. 3A, B), which is different from the typical budding mode observed

in yeast, where the cellular framework and contents both extend simultaneously

(Herskowitz, 1988). During microbial cell division, members of the Fts family of

proteins (tubulin homologs) usually assemble at the future site of cell division,

forming a contractile ring known as the Z ring (Wiegand et al., 2018). We observed

that the expression of numerous Fts-related proteins (FtsJ, FtsK, FtsL, FtsW, FtsX,

and FtsY) in strain ZRK32 were upregulated when the strain had been cultivated in

the rich medium (Figure. 3C). However, the FtsZ protein was absent from strain

ZRK32, which is consistent with the proposal that the *ftsZ* gene is absent from

Planctomycetes genomes (Jogler et al., 2012). In addition, we also found that some

genes that encoded rod shape-determining proteins (MreB, MreC, and MreD) were

present in strain ZRK32, and that their expression was upregulated in the strains

cultivated in the rich medium (Figure. 3C). MreB, usually participates in the

formation and degradation of peptidoglycan, which ultimately determines bacterial

cell shape (Rohs and Bernhardt, 2021).

Planctomycetes bacteria are major players in the global nitrogen cycling and

perform important reactions, such as the anaerobic ammonium oxidation process that

oxidizes NH_4^+ to N_2 gas, using NO_2^- as an electron acceptor (Strous et al., 1998;

Oshiki et al., 2016; Wiegand et al., 2018). To date, most studies on anaerobic

ammonium oxidation have been related to the monophyletic group (*Candidatus*

Brocadiae) in the phylum *Planctomycetes* (Strous et al., 1999), with only few reports

on the process of nitrogen metabolism in other *Planctomycetes* bacteria. Here, we

found that NO_3^- and NH_4^+ promoted the growth of strain ZRK32, while NO_2^- inhibited

its growth (Figure. 4A). Based on our data and results from others, we speculate that

NO_3^- might act as a terminal electron acceptor in the respiratory electron transport

chain of strain ZRK32. In addition, NO_2^- might react with the iron-sulfur proteins in

strain ZRK32 to form iron-nitric oxide complexes, which then inactivate iron-sulfur

enzymes and inhibit the growth of strain ZRK32 (Reddy et al., 1983). We speculate

that strain ZRK32 converts NO_3^- to NH_4^+ (Figure. 4B, C), enabling the NH_4^+ created

to enter the glutamate metabolic pathway (Figure. 4D, E) — a pathway that is closely

associated with several processes, including nitrogen metabolism, the TCA cycle, the

EMP glycolysis pathway, and amino acid metabolism (Figure. S8). Consistently, in

the presence of NO_3^- or NH_4^+, genes associated with the TCA cycle and the EMP

glycolysis pathway were upregulated (Figure. 4F, Figure. S5C and Figure. S7). The

presence of rich nutrients, and either NO_3^- or NH_4^+ also stimulated the expression of

genes encoding the NADH-ubiquinone oxidoreductase complex (Figures. 2D and 4G).

This complex couples the oxidation of NADH and the reduction of ubiquinone to

generate a proton gradient, which is then used for ATP synthesis (Reda et al., 2008).

Notably, large number of genes associated with flagellum assembly were also

up-regulated in strain ZRK32 (Figures. 2E and 4H). Flagellum-mediated motility is

beneficial for bacteria, not only because it allows them to respond quickly to an

ever-changing environment but also because it enables them to seek and acquire

nutrients for survival (Wadhams and Armitage, 2004; Zheng et al., 2021a). Thus,

strain ZRK32 might regulate the formation and motility of its flagellum to accelerate

the absorption and utilization of nutrients available in its environment.

One of our most exciting results was that NO_3^- or NH_4^+ could induce the release

of a chronic bacteriophage (Phage-ZRK32) from strain ZRK32 (Figure. 5A). The

Phage-ZRK32 was capable of facilitating nitrogen metabolism and amino acid

metabolism in strain ZRK32 through the function of AMGs, which incorporate nitrogen into certain amino acids (including glutamate, cysteine and glycine) (Figure. 5B) (Orlowski and Meister, 1970; Mouilleron and Golinelli-Pimpaneau, 2007; Iyer et al., 2009; Lu, 2013). Phylogenetic analyses were performed (Figures. S10-13) using AMG amino acid sequences (including amidoligase, glutamine amidotransferase, gamma-glutamylcyclotransferase, and glutathione synthase) from Phage-ZRK32, in addition to the same amino acid sequences from related phages and their bacterial hosts. The results showed that these AMGs may have been acquired from *Pseudomonas* via horizontal gene transfer (Matilla et al., 2014; Zhou et al., 2019). Consistently, Phage-ZRK32 promoted the growth of a marine *Pseudomonas* bacterium (*Pseudomonas stutzeri* 273) in the presence of either NO_3^- or NH_4^+ (Figure. 5C, D). Given that the AMGs encoding amidoligase and gamma-glutamylcyclotransferase were absent in the genome of *Pseudomonas stutzeri* 273, we speculate that Phage-ZRK32 might promote the growth of *Pseudomonas stutzeri* 273 by facilitating nitrogen metabolism and amino acid generation, as in the *Planctomycetes* strain ZRK32. Most bacteriophage life-cycles are described as a lytic or lysogenic cycle (Du Toit, 2017). Currently, more attention has been given to the chronic life cycle, where bacterial growth continues despite phage reproduction (Hoffmann Berling and Maze, 1964), and the progeny of these phage particles are released from host cells via extrusion or budding without killing the host (Putzrath and Maniloff, 1977; Russel, 1991; Marvin et al., 2014). Undoubtedly, Phage-ZRK32, which was induced by either NO_3^- or NH_4^+ in *Planctomycetes* strain ZRK32, is a chronic bacteriophage. Our findings provide a novel insight into nitrogen metabolism in *Planctomycetes* bacteria and provide a suitable model to study the interactions between *Planctomycetes* and viruses.

Materials and Methods

Enrichment and cultivation of deep-sea *Planctomycetes* bacteria

To isolate and cultivate *Planctomycetes* bacteria, 2 g deep-sea sediment samples were added to a 500 mL anaerobic bottle containing 400 mL basal medium (1.0 g/L yeast

extract, 20.0 g/L NaCl, 1.0 g/L CH$_3$COONa, 1.0 g/L NaHCO$_3$, 0.5 g/L KH$_2$PO$_4$, 0.2

g/L MgSO$_4$·7H$_2$O, 0.7 g/L cysteine hydrochloride, 500 μL/L 0.1 % (w/v) resazurin,

1.0 L sterilized distilled water, pH 7.0) supplemented with 1.0 g/L NH$_4$Cl and 1.0 g/L

NaNO$_3$, ensuring headspace volume was retained above the liquid surface. The

medium was prepared under a 100% N$_2$ gas phase and sterilized by autoclaving at

115 ℃ for 30 minutes; following this, rifampicin (100 μg/mL) was added. The

inoculated media were anaerobically incubated at either 4 ℃ or 28 ℃ for one month.

The basal medium supplemented with 1.0 g/L NH$_4$Cl, 1.0 g/L NaNO$_3$, and 15 g/L agar

was evenly spread on to the inside wall of a Hungate tube, which formed a thin layer

of medium for the bacteria to grow. After this, 50 μL of the enriched culture was

anaerobically transferred into an anaerobic roll tube and then spread on to the medium

layer. These tubes were also anaerobically cultured at either 4 ℃ or 28 ℃ for ten days.

Single colonies growing at 28 ℃ were selected using sterilized bamboo sticks; they

were then cultured in the 15 mL Hungate tube containing 10 mL basal medium

(supplemented with 1.0 g/L NH$_4$Cl and 1.0 g/L NaNO$_3$) at 28 ℃ for seven days under

a 100% N$_2$ atmosphere. One strain was identified as a member of the phylum

Planctomycetes, but was noted to have less than 98% 16S rRNA gene sequence

similarity to other cultured strains; this strain was therefore selected to be cultivated.

Strain ZRK32 was selected and purified by repeating the Hungate roll-tube method.

The purity of strain ZRK32 was confirmed regularly by observation using a

transmission electron microscope (TEM) and by repeating partial sequencing of the

16S rRNA gene. As strain ZRK32 grew slowly in basal medium, we used a rich

culture medium (10.0 g/L yeast extract, 20.0 g/L NaCl, 1.0 g/L CH$_3$COONa, 1.0 g/L

NaHCO$_3$, 0.5 g/L KH$_2$PO$_4$, 0.2 g/L MgSO$_4$·7H$_2$O, 0.7 g/L cysteine hydrochloride, 500

μL/L 0.1 % (w/v) resazurin, 1.0 L sterilized distilled water, pH 7.0).

TEM observation

To observe the morphological characteristics of strain ZRK32, 10 mL culture was

collected by centrifuging at 5000 × g for ten minutes. Cells were then washed three

times with PBS buffer (137 mM NaCl, 2.7 mM KCl, 10 mM Na$_2$HPO$_4$, 1.8 mM

KH$_2$PO$_4$, 1 L sterile water, pH 7.4). Finally, the cells were suspended in 20 μL PBS

buffer, and then transferred onto copper grids coated with a carbon film by immersing

the grids in the cell suspension for 30 minutes (Zheng et al., 2021b). To observe the

ultrastructure of strain ZRK32, ultrathin-sections were prepared using methods

previously described (Graham and Orenstein, 2007). Briefly, 500 mL of cells

(cultured for six days at 28 ℃) were collected by centrifuging at 5000 × g for 20

minutes, and then washing three times with PBS buffer. The cells were then preserved

in 2.5% (v/v) glutaraldehyde for 12 hours at 4 °C and then dehydrated using different

ethanol concentrations (30%, 50%, 70%, 90%, and 100%) for ten minutes each time.

The cells were then embedded in a plastic resin. Finally, 50-70 nm ultrathin sections

were produced using an ultramicrotome (Leica EM UC7, Germany) and then stained

using uranyl acetate and lead citrate. All samples were examined under TEM

(HT7700, Hitachi, Japan).

Phylogenetic analysis

To construct a maximum likelihood 16S rRNA phylogenetic tree, the full-length 16S

rRNA gene sequences of strain ZRK32 and other related taxa were extracted from

their corresponding genomes (www.ncbi.nlm.nih.gov/). The maximum likelihood

genome phylogenetic tree was constructed from a concatenated alignment of 37

protein-coding genes (Wu et al., 2013) (extracted from the genomes using Phylosift

v1.0.1) (Darling et al., 2014); all genes were present in a single copy and were

universally distributed in both archaea and bacteria (Table S2). All phylogenetic trees

were constructed using the W-IQ-TREE web server (http://iqtree.cibiv.univie.ac.at)

(Trifinopoulos et al., 2016) using the "GTR+F+I+G4" model, and the Interactive Tree

of Life (iTOL v5) online tool (Letunic and Bork, 2021) was used to edit the

phylogenetic trees.

Growth assays of strain ZRK32

To assess the effects of nutrient-rich media on strain ZRK32 growth, we set up

different cultures using either rich medium or basal medium. To assess the effects of

different inorganic nitrogen sources (20 mM NO$_3^-$, 20 mM NH$_4^+$, and 20 mM NO$_2^-$)

on strain ZRK32 growth, we used a rich culture medium supplemented with the nitrogen sources mentioned above. For each growth assay, 15 mL of strain ZRK32 culture was inoculated in a 2 L Hungate bottle containing 1.5 L of the respective media. All Hungate bottles were anaerobically incubated at 28 °C. Bacterial growth was monitored by measuring daily OD_{600} values via a microplate reader until cell growth reached a stationary phase. Three replicates were performed for each condition. The concentrations of NO_3^-, NH_4^+, and NO_2^- were determined using a continuous flow analyzer (SKALAR-SAN++, Netherlands), which has an analytical precision of 6%.

Transcriptomics

For transcriptomic sequencing, strains of ZRK32 were cultured in 1.5 L of either basal medium, rich medium, or rich medium supplemented with different nitrogen sources (20 mM NO_3^-, 20 mM NH_4^+, or 20 mM NO_2^-) at 28 °C for six days. Three biological replicates were cultured for each condition. Following this, cells from the three replicates were mixed and then collected for transcriptomic sequencing by Novogene (Tianjin, China), as previously described (Zheng et al., 2021b; Zheng et al., 2022). Detailed protocols describing library preparation, clustering, sequencing, and data analyses are described in the supplementary information. All heat maps were made by HemI 1.0.3.3.

Isolation of bacteriophages

Isolation of the bacteriophages was performed using similar methods to those described previously, but with some modifications (Yamamoto et al., 1970; Tseng et al., 1990; Kim and Blaschek, 1991). Strain ZRK32 was inoculated in either rich medium, or rich medium supplemented with 20 mM NO_3^- or 20 mM NH_4^+, and then incubated at 28 °C for six days. Different cultures were collected by centrifuging at 8000 $\times$ g at 4 °C for 20 minutes; this was repeated three times. The supernatant was filtered through a 0.22 μm millipore filter (Pall Supor, USA), and then 1 M NaCl was added to lyse the residual bacteria. The supernatant was collected by centrifuging at 8000 $\times$ g at 4 °C for 20 minutes. The phage particles were immediately precipitated

with 100 g/L polyethylene glycol (PEG8000) at 4 ℃ for two hours, and collected by centrifuging at 10000 × g at 4 ℃ for 20 minutes. The phage particles were then suspended in SM buffer (0.01% gelatin, 50 mM Tris-HCl, 100 mM NaCl, and 10 mM MgSO$_4$). The suspension was then extracted three times using an equal volume of chloroform (Lin et al., 2012) and collected by centrifuging at 4000 × g at 4 ℃ for 20 minutes. Finally, the clean phage particles were obtained. Detailed protocols describing the genome sequencing of the bacteriophages are described in the supplementary information.

Growth assay of *Pseudomonas stutzeri* 273 cultured in oligotrophic medium supplemented with different nitrogen sources and Phage-ZRK32

The assistance role of the bacteriophage induced from strain ZRK32 was tested in another deep-sea bacterium *Pseudomonas stutzeri* 273 (Wu et al., 2016). Specifically, 50 μL freshly incubated *Pseudomonas stutzeri* 273 cells were inoculated in 5 mL of either oligotrophic medium (10 g/L NaCl, 0.1 g/L yeast extract, 1 L sterilized distilled water), oligotrophic medium supplemented with 20 μl/L Phage-ZRK32 (without the extraction by chloroform), oligotrophic medium supplemented with 20 mM NO$_3^-$ or NH$_4^+$, oligotrophic medium supplemented with 20 mM of either NO$_3^-$ or NH$_4^+$, or 20 μl/L Phage-ZRK32. The cultures were then incubated for 24 hours at 28 ℃. Three biological replicates for each culture condition were performed. The progress of the bacterial growth was monitored by measuring OD$_{600}$ values using a microplate reader (Infinite M1000 Pro, Switzerland) every four hours until cell growth reached a stationary phase.

Data availability

The full-length 16S rRNA gene sequence of strain ZRK32 has been deposited at GenBank under the accession number MW376756. The complete genome sequence of strain ZRK32 has been deposited at GenBank under the accession number CP066225. The raw sequencing reads from the transcriptomics analyses of ZRK32 strains cultured with different concentrations of yeast extract and nitrogen sources

have been deposited into the NCBI Short Read Archive (accession numbers: PRJNA768630 and PRJNA694614, respectively). The genome sequence of Phage-ZRK32 has been deposited into the GenBank database with the accession number OP650935.

Acknowledgements

This work was funded by the Major Research Plan of the National Natural Science Foundation (Grant No. 92051107), the NSFC Innovative Group Grant (No. 42221005), Science and Technology Innovation Project of Laoshan Laboratory (Grant No. LSKJ202203103), Shandong Provincial Natural Science Foundation (ZR2021ZD28), Strategic Priority Research Program of the Chinese Academy of Sciences (Grant No. XDA22050301), Key deployment projects of Center of Ocean Mega-Science of the Chinese Academy of Sciences (Grant No. COMS2020Q04), and Key Collaborative Research Program of the Alliance of International Science Organizations (Grant No. ANSO-CR-KP-2022-08) for Chaomin Sun. We thank Dr. Diana Walsh from Life Science Editors for her great effort to improve the writing quality of our manuscript.

Author contributions

RZ, CW, and CS conceived and designed the study; RZ and CW conducted most of the experiments; RL collected the samples from the deep-sea cold seep; RC helped to construct evolutionary trees; RZ, CW, and CS lead the writing of the manuscript; all authors contributed to and reviewed the manuscript.

Conflict of interest

The authors declare that they have no competing interests.

References

Afgan, E., Baker, D., Batut, B., van den Beek, M., Bouvier, D., Cech, M., Chilton, J., Clements, D., Coraor, N., Grüning, B.A., Guerler, A., Hillman-Jackson, J., Hiltemann, S., Jalili, V., Rasche, H., Soranzo, N., Goecks, J., Taylor, J., Nekrutenko, A., and Blankenberg, D. The Galaxy platform for accessible, reproducible and collaborative biomedical analyses: 2018 update. *Nucleic Acids Res.* 2018;46:W537-W544.

Alarcón-Schumacher, T., Naor, A., Gophna, U., and Erdmann, S. Isolation of a virus
causing a chronic infection in the archaeal model organism *Haloferax volcanii* reveals
antiviral activities of a provirus. *Proc Natl Acad Sci USA.* 2022;119:e2205037119.

Bondoso, J., Albuquerque, L., Nobre, M.F., Lobo-da-Cunha, A., da Costa, M.S., and
Lage, O.M. *Aquisphaera giovannonii* gen. nov., sp. nov., a planctomycete isolated
from a freshwater aquarium. *Int J Syst Evol Microbiol.* 2011;61:2844-2850.

Brümmer, I.H., Felske, A.D., and Wagner-Döbler, I. Diversity and seasonal changes
of uncultured *Planctomycetales* in river biofilms. *Appl Environ Microbiol.*
2004;70:5094-5101.

Buckley, D.H., Huangyutitham, V., Nelson, T.A., Rumberger, A., and Thies, J.E.
Diversity of *Planctomycetes* in soil in relation to soil history and environmental
heterogeneity. *Appl Environ Microbiol.* 2006;72:4522-4531.

Cassman, N., Prieto-Davó, A., Walsh, K., Silva, G.G., Angly, F., Akhter, S., Barott,
K., Busch, J., McDole, T., Haggerty, J.M., Willner, D., Alarcón, G., Ulloa, O.,
DeLong, E.F., Dutilh, B.E., Rohwer, F., and Dinsdale, E.A. Oxygen minimum zones
harbour novel viral communities with low diversity. *Environ Microbiol.*
2012;14:3043-3065.

Chevallereau, A., Pons, B.J., van Houte, S., and Westra, E.R. Interactions between
bacterial and phage communities in natural environments. *Nat Rev Microbiol.*
2022;20:49-62.

Darling, A.E., Jospin, G., Lowe, E., Matsen, F.I.V., Bik, H.M., and Eisen, J.A.
PhyloSift: phylogenetic analysis of genomes and metagenomes. *Peerj.* 2014;2:e243.

Dedysh, S.N., and Ivanova, A.A. *Planctomycetes* in boreal and subarctic wetlands:
diversity patterns and potential ecological functions. *FEMS Microbiol Ecol.* 2019;95.

Dedysh, S.N., Kulichevskaya, I.S., Beletsky, A.V., Ivanova, A.A., Rijpstra, W.I.C.,
Damsté, J.S.S., Mardanov, A.V., and Ravin, N.V. *Lacipirellula parvula* gen. nov., sp.
nov., representing a lineage of planctomycetes widespread in low-oxygen habitats,
description of the family *Lacipirellulaceae* fam. nov. and proposal of the orders
Pirellulales ord. nov., *Gemmatales* ord. nov. and *Isosphaerales* ord. nov. *Syst Appl*
Microbiol. 2020;43:126050.

Dedysh, S.N., Beletsky, A.V., Ivanova, A.A., Kulichevskaya, I.S., Suzina, N.E.,
Philippov, D.A., Rakitin, A.L., Mardanov, A.V., and Ravin, N.V. Wide distribution of
Phycisphaera-like planctomycetes from WD2101 soil group in peatlands and genome
analysis of the first cultivated representative. *Environ Microbiol.* 2021;23:1510-1526.

Du Toit, A. Viral infection: The language of phages. *Nat Rev Microbiol.*
2017;15:134-135.

Fuerst, J.A., and Webb, R.I. Membrane-bounded nucleoid in the eubacterium
Gemmata obscuriglobus. Proc Natl Acad Sci U S A. 1991;88:8184-8188.

Fuerst, J.A., and Sagulenko, E. Beyond the bacterium: planctomycetes challenge our
concepts of microbial structure and function. *Nat Rev Microbiol.* 2011;9:403-413.

Fukunaga, Y., Kurahashi, M., Sakiyama, Y., Ohuchi, M., Yokota, A., and Harayama,
S. *Phycisphaera mikurensis* gen. nov., sp. nov., isolated from a marine alga, and
proposal of *Phycisphaeraceae* fam. nov., *Phycisphaerales* ord. nov. and

Phycisphaerae classis nov. in the phylum *Planctomycetes*. *J Gen Appl Microbiol.* 2009;55:267-275.

Gazitúa, M.C., Vik, D.R., Roux, S., Gregory, A.C., Bolduc, B., Widner, B., Mulholland, M.R., Hallam, S.J., Ulloa, O., and Sullivan, M.B. Potential virus-mediated nitrogen cycling in oxygen-depleted oceanic waters. *ISME J.* 2021;15:981-998.

Goffredi, S.K., and Orphan, V.J. Bacterial community shifts in taxa and diversity in response to localized organic loading in the deep sea. *Environ Microbiol.* 2010;12:344-363.

Graham, L., and Orenstein, J.M. Processing tissue and cells for transmission electron microscopy in diagnostic pathology and research. *Nat Protoc.* 2007;2:2439-2450.

Herskowitz, I. Life cycle of the budding yeast *Saccharomyces cerevisiae*. *Microbiol Rev.* 1988;52:536-553.

Hoffmann Berling, H., and Maze, R. Release of male-specific bacteriophages from surviving host bacteria. *Virology.* 1964;22:305-313.

Howard-Varona, C., Hargreaves, K.R., Abedon, S.T., and Sullivan, M.B. Lysogeny in nature: mechanisms, impact and ecology of temperate phages. *ISME J.* 2017;11:1511-1520.

Iyer, L.M., Abhiman, S., Maxwell Burroughs, A., and Aravind, L. Amidoligases with ATP-grasp, glutamine synthetase-like and acetyltransferase-like domains: synthesis of novel metabolites and peptide modifications of proteins. *Mol Biosyst.* 2009;5:1636-1660.

Jogler, C., Waldmann, J., Huang, X., Jogler, M., Glöckner, F.O., Mascher, T., and Kolter, R. Identification of proteins likely to be involved in morphogenesis, cell division, and signal transduction in *Planctomycetes* by comparative genomics. *J Bacteriol.* 2012;194:6419-6430.

Kaboré, O.D., Godreuil, S., and Drancourt, M. *Planctomycetes* as host-associated bacteria: a perspective that holds promise for their future isolations, by mimicking their native environmental niches in clinical microbiology laboratories. *Front Cell Infect Microbiol.* 2020;10:519301.

Kallscheuer, N., Jogler, M., Wiegand, S., Peeters, S.H., Heuer, A., Boedeker, C., Jetten, M.S.M., Rohde, M., and Jogler, C. *Rubinisphaera italica* sp. nov. isolated from a hydrothermal area in the Tyrrhenian Sea close to the volcanic island Panarea. *Antonie Van Leeuwenhoek.* 2020a;113:1727-1736.

Kallscheuer, N., Wiegand, S., Kohn, T., Boedeker, C., Jeske, O., Rast, P., Müller, R.W., Brümmer, F., Heuer, A., Jetten, M.S.M., Rohde, M., Jogler, M., and Jogler, C. Cultivation-independent analysis of the bacterial community associated with the calcareous sponge clathrina clathrus and isolation of *Poriferisphaera corsica* gen. nov., sp. nov., belonging to the barely studied class *Phycisphaerae* in the phylum *Planctomycetes*. *Front Microbiol.* 2020b;11:602250.

Kartal, B., van Niftrik, L., Keltjens, J.T., Op den Camp, H.J., and Jetten, M.S. Anammox--growth physiology, cell biology, and metabolism. *Adv Microb Physiol.* 2012;60:211-262.

Kartal, B., de Almeida, N.M., Maalcke, W.J., Op den Camp, H.J., Jetten, M.S., and
Keltjens, J.T. How to make a living from anaerobic ammonium oxidation. *FEMS*
microbiology reviews. 2013;37:428-461.

Kaushik, R., Sharma, M., Gaurav, K., Jagadeeshwari, U., Shabbir, A., Sasikala, C.,
Ramana, C.V., and Pandit, M.K. *Paludisphaera soli* sp. nov., a new member of the
family *Isosphaeraceae* isolated from high altitude soil in the Western Himalaya.
Antonie Van Leeuwenhoek. 2020;113:1663-1674.

Kim, A.Y., and Blaschek, H.P. Isolation and characterization of a filamentous
viruslike particle from *Clostridium acetobutylicum* NCIB 6444. *J Bacteriol.*
1991;173:530-535.

Kovaleva, O.L., Merkel, A.Y., Novikov, A.A., Baslerov, R.V., Toshchakov, S.V., and
Bonch-Osmolovskaya, E.A. *Tepidisphaera mucosa* gen. nov., sp. nov., a moderately
thermophilic member of the class *Phycisphaerae* in the phylum *Planctomycetes*, and
proposal of a new family, *Tepidisphaeraceae* fam. nov., and a new order,
Tepidisphaerales ord. nov. *Int J Syst Evol Microbiol.* 2015;65:549-555.

Kulichevskaya, I.S., Ivanova, A.A., Naumoff, D.G., Beletsky, A.V., Rijpstra, W.I.C.,
Sinninghe Damsté, J.S., Mardanov, A.V., Ravin, N.V., and Dedysh, S.N.
Frigoriglobus tundricola gen. nov., sp. nov., a psychrotolerant cellulolytic
planctomycete of the family *Gemmataceae* from a littoral tundra wetland. *Syst Appl*
Microbiol. 2020;43:126129.

Letunic, I., and Bork, P. Interactive Tree Of Life (iTOL) v5: an online tool for
phylogenetic tree display and annotation. *Nucleic Acids Res.* 2021;49:W293-W296.

Lin, Y.R., Chiu, C.W., Chang, F.Y., and Lin, C.S. Characterization of a new phage,
termed phi A318, which is specific for *Vibrio alginolyticus*. *Arch Virol.*
2012;157:917-926.

Lindsay, M.R., Webb, R.I., and Fuerst, J.A. Pirellulosomes: a new type of
membrane-bounded cell compartment in planctomycete bacteria of the genus
Pirellula. *Microbiology (Reading).* 1997;143:739-748.

Liu, Y., Alexeeva, S., Bachmann, H., Guerra Martínez, J.A., Yeremenko, N., Abee, T.,
and Smid, E.J. Chronic release of tailless phage particles from *Lactococcus lactis*.
Appl Environ Microbiol. 2022;88:e0148321.

Lonhienne, T.G., Sagulenko, E., Webb, R.I., Lee, K.C., Franke, J., Devos, D.P.,
Nouwens, A., Carroll, B.J., and Fuerst, J.A. Endocytosis-like protein uptake in the
bacterium *Gemmata obscuriglobus*. *Proc Natl Acad Sci U S A.*
2010;107:12883-12888.

Lu, S.C. Glutathione synthesis. *Biochim Biophys Acta.* 2013;1830:3143-3153.

Marvin, D.A., Symmons, M.F., and Straus, S.K. Structure and assembly of
filamentous bacteriophages. *Prog Biophys Mol Bio.* 2014;114:80-122.

Matilla, M.A., Fang, X., and Salmond, G.P. Viunalikeviruses are environmentally
common agents of horizontal gene transfer in pathogens and biocontrol bacteria.
ISME J. 2014;8:2143-2147.

Monier, A., Chambouvet, A., Milner, D.S., Attah, V., Terrado, R., Lovejoy, C.,
Moreau, H., Santoro, A.E., Derelle, É., and Richards, T.A. Host-derived viral

transporter protein for nitrogen uptake in infected marine phytoplankton. *Proc Natl Acad Sci U S A.* 2017;114:E7489-e7498.

Mouilleron, S., and Golinelli-Pimpaneau, B. Conformational changes in ammonia-channeling glutamine amidotransferases. *Curr Opin Struct Biol.* 2007;17:653-664.

Orgel, L.E. The origin of life - a review of facts and speculations. *Trends Biochem Sci.* 1998;23:491-495.

Orlowski, M., and Meister, A. The gamma-glutamyl cycle: a possible transport system for amino acids. *Proc Natl Acad Sci U S A.* 1970;67:1248-1255.

Oshiki, M., Satoh, H., and Okabe, S. Ecology and physiology of anaerobic ammonium oxidizing bacteria. *Environ Microbiol.* 2016;18:2784-2796.

Peeters, S.H., Wiegand, S., Kallscheuer, N., Jogler, M., Heuer, A., Jetten, M.S.M., Rast, P., Boedeker, C., Rohde, M., and Jogler, C. Three marine strains constitute the novel genus and species *Crateriforma conspicua* in the phylum *Planctomycetes*. *Antonie Van Leeuwenhoek.* 2020;113:1797-1809.

Pollet, T., Tadonléké, R.D., and Humbert, J.F. Spatiotemporal changes in the structure and composition of a less-abundant bacterial phylum (*Planctomycetes*) in two perialpine lakes. *Appl Environ Microbiol.* 2011;77:4811-4821.

Pradel, N., Fardeau, M.L., Tindall, B.J., and Spring, S. *Anaerohalosphaera lusitana* gen. nov., sp. nov., and *Limihaloglobus sulfuriphilus* gen. nov., sp. nov., isolated from solar saltern sediments, and proposal of *Anaerohalosphaeraceae* fam. nov. within the order *Sedimentisphaerales*. *Int J Syst Evol Microbiol.* 2020;70:1321-1330.

Putzrath, R.M., and Maniloff, J. Growth of an enveloped mycoplasmavirus and establishment of a carrier state. *J Virol.* 1977;22:308-314.

Reda, T., Barker, C.D., and Hirst, J. Reduction of the iron-sulfur clusters in mitochondrial NADH:ubiquinone oxidoreductase (complex I) by EuII-DTPA, a very low potential reductant. *Biochemistry.* 2008;47:8885-8893.

Reddy, D., Lancaster, J.R., Jr., and Cornforth, D.P. Nitrite inhibition of *Clostridium botulinum*: electron spin resonance detection of iron-nitric oxide complexes. *Science.* 1983;221:769-770.

Rivas-Marín, E., Canosa, I., and Devos, D.P. Evolutionary cell biology of division mode in the bacterial *Planctomycetes-Verrucomicrobia- Chlamydiae* superphylum. *Front Microbiol.* 2016;7:1964.

Rohs, P.D.A., and Bernhardt, T.G. Growth and division of the peptidoglycan matrix. *Annu Rev Microbiol.* 2021;75:315-336.

Russel, M. Filamentous phage assembly. *Mol Microbiol.* 1991;5:1607-1613.

Salbreiter, M., Waqqas, M., Jogler, M., Kallscheuer, N., Wiegand, S., Peeters, S.H., Heuer, A., Jetten, M.S.M., Boedeker, C., Rast, P., Rohde, M., and Jogler, C. Three Planctomycetes isolated from biotic surfaces in the Mediterranean Sea and the Pacific Ocean constitute the novel species *Symmachiella dynata* gen. nov., sp. nov. and *Symmachiella macrocystis* sp. nov. *Antonie Van Leeuwenhoek.* 2020;113:1965-1977.

Spring, S., Bunk, B., Spröer, C., Rohde, M., and Klenk, H.P. Genome biology of a novel lineage of planctomycetes widespread in anoxic aquatic environments. *Environ Microbiol.* 2018;20:2438-2455.

Strous, M., Heijnen, J.J., Kuenen, J.G., and Jetten, M.S.M. The sequencing batch reactor as a powerful tool for the study of slowly growing anaerobic ammonium-oxidizing microorganisms. *Appl Microbiol Biotechnol.* 1998;50:589-596.

Strous, M., Fuerst, J.A., Kramer, E.H., Logemann, S., Muyzer, G., van de Pas-Schoonen, K.T., Webb, R., Kuenen, J.G., and Jetten, M.S. Missing lithotroph identified as new planctomycete. *Nature.* 1999;400:446-449.

Trifinopoulos, J., Nguyen, L.T., von Haeseler, A., and Minh, B.Q. W-IQ-TREE: a fast online phylogenetic tool for maximum likelihood analysis. *Nucleic Acids Res.* 2016;44:W232-W235.

Tseng, Y.H., Lo, M.C., Lin, K.C., Pan, C.C., and Chang, R.Y. Characterization of filamentous bacteriophage phi Lf from *Xanthomonas campestris* pv. campestris. *J Gen Virol.* 1990;71:1881-1884.

Vigneron, A., Alsop, E.B., Cruaud, P., Philibert, G., King, B., Baksmaty, L., Lavallee, D., Lomans, B.P., Kyrpides, N.C., Head, I.M., and Tsesmetzis, N. Comparative metagenomics of hydrocarbon and methane seeps of the Gulf of Mexico. *Sci Rep.* 2017;7:16015.

Wadhams, G.H., and Armitage, J.P. Making sense of it all: bacterial chemotaxis. *Nat Rev Mol Cell Biol.* 2004;5:1024-1037.

Wang, J., Ruan, C.J., Song, L., Li, A., Zhu, Y.X., Zheng, X.W., Wang, L., Lu, Z.J., Huang, Y., Du, W.B., Zhou, Y.G., Huang, L., and Dai, X. *Gimesia benthica* sp. nov., a planctomycete isolated from a deep-sea water sample of the Northwest Indian Ocean. *Int J Syst Evol Micr.* 2020;70:4384-4389.

Wang, S., Yang, Y., and Jing, J. A synthesis of viral contribution to marine nitrogen cycling. *Front Microbiol.* 2022;13:834581.

Wiegand, S., Jogler, M., and Jogler, C. On the maverick *Planctomycetes*. *FEMS Microbiol Rev.* 2018;42:739-760.

Wiegand, S., Jogler, M., Boedeker, C., Heuer, A., Rast, P., Peeters, S.H., Jetten, M.S.M., Kaster, A.K., Rohde, M., Kallscheuer, N., and Jogler, C. Additions to the genus *Gimesia*: description of *Gimesia alba* sp. nov., *Gimesia algae* sp. nov., *Gimesia aquarii* sp. nov., *Gimesia aquatilis* sp. nov., *Gimesia fumaroli* sp. nov. and *Gimesia panareensis* sp. nov., isolated from aquatic habitats of the Northern Hemisphere. *Antonie Van Leeuwenhoek.* 2020a;113:1999-2018.

Wiegand, S., Jogler, M., Boedeker, C., Pinto, D., Vollmers, J., Rivas-Marń, E., Kohn, T., Peeters, S.H., Heuer, A., Rast, P., Oberbeckmann, S., Bunk, B., Jeske, O., Meyerdierks, A., Storesund, J.E., Kallscheuer, N., Lücker, S., Lage, O.M., Pohl, T., Merkel, B.J., Hornburger, P., Müller, R.W., Brümmer, F., Labrenz, M., Spormann, A.M., Op den Camp, H.J.M., Overmann, J., Amann, R., Jetten, M.S.M., Mascher, T., Medema, M.H., Devos, D.P., Kaster, A.K., Øvreås, L., Rohde, M., Galperin, M.Y., and Jogler, C. Cultivation and functional characterization of 79 planctomycetes uncovers their unique biology. *Nat Microbiol.* 2020b;5:126-140.

Woebken, D., Teeling, H., Wecker, P., Dumitriu, A., Kostadinov, I., Delong, E.F., Amann, R., and Glöckner, F.O. Fosmids of novel marine *Planctomycetes* from the Namibian and Oregon coast upwelling systems and their cross-comparison with planctomycete genomes. *ISME J.* 2007;1:419-435.

Wu, D.Y., Jospin, G., and Eisen, J.A. Systematic identification of gene families for
use as "markers" for phylogenetic and phylogeny-driven ecological studies of
Bacteria and Archaea and their major subgroups. *Plos One.* 2013;8:e77033.
Wu, S., Liu, G., Jin, W., Xiu, P., and Sun, C. Antibiofilm and anti-Infection of a
marine bacterial exopolysaccharide against *Pseudomonas aeruginosa*. *Front*
Microbiol. 2016;7:102.
Yamamoto, K.R., Alberts, B.M., Benzinger, R., Lawhorne, L., and Treiber, G. Rapid
bacteriophage sedimentation in the presence of polyethylene glycol and its application
to large-scale virus purification. 1970;40:734-744.
Yoon, J., Jang, J.H., and Kasai, H. *Algisphaera agarilytica* gen. nov., sp. nov., a novel
representative of the class *Phycisphaerae* within the phylum *Planctomycetes* isolated
from a marine alga. *Antonie Van Leeuwenhoek.* 2014;105:317-324.
Zhang, C., Hicks, G.R., and Raikhel, N.V. Plant vacuole morphology and vacuolar
trafficking. *Front Plant Sci.* 2014;5:476.
Zheng, R., Wu, S., and Sun, C. MerF is a novel regulator of deep-sea *Pseudomonas*
stutzeri flagellum biogenesis and motility. *Environ Microbiol.* 2021a;23:110-125.
Zheng, R., Cai, R., Wang, C., Liu, R., and Sun, C. Characterization of the first
cultured representative of "*Candidatus* Thermofonsia" Clade 2 within *Chloroflexi*
reveals its phototrophic lifestyle. *mBio.* 2022;13:e0028722.
Zheng, R., Liu, R., Shan, Y., Cai, R., Liu, G., and Sun, C. Characterization of the first
cultured free-living representative of *Candidatus* Izemoplasma uncovers its unique
biology. *ISME J.* 2021b;15:2676-2691.
Zhou, K., Zhang, R., Sun, J., Zhang, W., Tian, R.M., Chen, C., Kawagucci, S., and
Xu, Y. Potential interactions between clade SUP05 sulfur-oxidizing bacteria and
phages in hydrothermal vent sponges. *Appl Environ Microbiol.* 2019;85:1-20.
821