Title page

Multi-level processing of emotions in life motion signals

revealed through pupil responses

Tian Yuan[1,2,3], Li Wang[1,2,3]*, Yi Jiang[1,2,3]

[1]State Key Laboratory of Brain and Cognitive Science, CAS Center for Excellence in

Brain Science and Intelligence Technology, Institute of Psychology, Chinese Academy

of Sciences, 16 Lincui Road, Beijing 100101, China; [2]Department of Psychology,

University of Chinese Academy of Sciences, 19A Yuquan Road, Beijing 100049, China;

[3]Chinese Institute for Brain Research, 26 Science Park Road, Beijing 102206, China

Running title: pupil response to emotion

Key words: pupil size, biological motion, emotion processing, local

*Corresponding authors: Li Wang (wangli@psych.ac.cn)

Institute of Psychology, Chinese Academy of Sciences, 16 Lincui Road, Beijing

100101, China

Abstract

Perceiving emotions from the movements of other biological entities is critical for
human survival and interpersonal interactions. Here, we report that emotional
information conveyed by point-light biological motion (BM) triggered automatic
physiological responses as reflected in pupil size. Specifically, happy BM evoked larger
pupil size than neutral and sad BM, while sad BM induced a smaller pupil response
than neutral BM. Moreover, this happy over sad pupil dilation effect is negatively
correlated with individual autistic traits. Notably, emotional BM with only local motion
features retained could also exert modulations on pupils. Compared with intact BM,
both happy and sad local BM evoked stronger pupil responses than neutral local BM
starting from an earlier timepoint, with no difference between the happy and sad
conditions. These results revealed a fine-grained pupil-related emotional modulation
induced by intact BM and a coarse but rapid modulation by local BM, demonstrating
multi-level processing of emotions in life motion signals. Taken together, our findings
shed new light on the mechanisms underlying BM emotion processing, and highlight
the potential of utilizing the emotion-modulated pupil response to facilitate the
diagnosis of social cognitive disorders.
35

Introduction

Perceiving and interpreting emotions from various social signals is critical for human social functioning, which enables us to infer the intentions of our conspecifics and further facilitates interpersonal interactions. Facial expressions present the most common non-verbal social communicative signals regarding others' affective states and intentions (Frith, 2009). In addition to faces, the movements of biological organisms serve as another essential type of social signals carrying significant emotional information, and this information remained salient even from a far distance (de Gelder, 2006). The human visual system is highly sensitive to such signals that we can readily decipher emotions from biological motion (BM), even when it was portrayed by several point lights attached to the major joints (Johansson, 1973; Troje, 2008). Moreover, it has been found that happy point-light walkers were recognized faster and more accurately than sad, angry or neutral walkers, demonstrating a happiness superiority (Lee & Kim, 2016; Spencer et al., 2016). While these studies provided some insights into the emotion processing mechanism of BM, they relied mainly on the explicit identification and active evaluation of BM emotions. Importantly, the encoding of emotional information also involved a rather automatic and implicit process that is independent of the participant's explicit identifications (Critchley et al., 2000; Lange et al., 2003; Okon-Singer et al., 2013; Shafer et al., 2012). Notably, this implicit aspect of emotion processing could be even more effective in probing individual differences and social deficits (Kana et al., 2015; Keifer et al., 2020; Kovarski et al., 2018; Wong et al., 2008), as it requires the intuitive processing of emotions that could not be learned (U. Frith, 2004). For example, research has shown that individuals with autistic disorders showed altered neural activities during implicit but not explicit emotion processing of natural scenes (Kana et al., 2015). These observations highlighted the importance of using objective measurements to investigate the implicit and automatic aspect of BM emotion processing.

The pupil response hence serves as a promising approach for reliably unfolding the implicit emotion processing of BM as it adds no extra task requirements to the cognitive process (Burley et al., 2017). In particular, pupil size is related to the activity of the automatic nervous system mediated by the locus coeruleus norepinephrine (LC-NE), which not only responds to physical light, but also reflects the underlying cognitive state (Joshi & Gold, 2019). Moreover, it could spontaneously capture the current subjective state and is thus useful for revealing the time course of the related cognitive processing (de Gee et al., 2014; Graves et al., 2021; Kloosterman et al., 2015;

Oliva & Anikin, 2018). Recently, this measurement has been introduced to the field of
emotion perception, and emerging evidence has suggested that emotions conveyed by
social signals (e.g., faces, voices) could modulate pupil responses. For instance, happy
and angry dynamic faces elicited a pupil dilation effect as compared with sad and
neutral faces (Burley & Daughters, 2020; Prunty et al., 2021). Noticeably, the point-
light BM, as another critical type of social signals, also conveyed salient affective
information. Besides, its emotion processing mechanism is closely connected with that
of faces (Alaerts et al., 2011; Becker et al., 2011; Lee & Kim, 2016). However, it
remains unequivocal whether the pupil also responds to the emotional information
carried by the minimalistic point-light walker display. Such physiological observation
can faithfully uncover the implicit emotion processing mechanism of BM, and it also
expands the existing line of inquiry on the emotion processing of social cues.

To fill this gap, the current study implemented the pupil recording technique
together with the passive viewing paradigm to explore the automatic and implicit
emotion processing mechanism of BM. We first investigated the pupil responses to
point-light walkers with different emotions (i.e., happy, sad and neutral). In addition to
intact emotional BM sequences, we also tested scrambled emotional BM sequences,
which lack the gestalt of a global figure but preserve the same local motion components
as the intact walkers. Recent studies have shown that such local motion signals play an
essential role in conveying biologically salient information such as animacy, walking
direction (Chang & Troje, 2008, 2009; Troje & Westhoff, 2006). This study went
further to investigate whether this local BM signal could convey emotional information
and elicit corresponding pupil responses. Moreover, given that emotion perception from
social signals is generally impaired in individuals with autism (Harms et al., 2010;
Hubert et al., 2006), we also took the individual autistic traits into consideration by
measuring the autistic tendencies in normal populations with the Autistic Quotient (AQ)
questionnaire (Baron-Cohen et al., 2001).

Results

Experiment 1: Intact emotional BM

In Experiment 1, we investigated whether emotional BM could exert influences on
pupil responses. A one-way repeated measures ANOVA was conducted on the mean
pupil size for each emotional condition (happy, sad, neutral) obtained by collapsing the
pupillometry across all time points. The results showed a significant main effect of
emotional condition ($F(1.4, 32.2) = 8.93$, $p = .002$, $\eta_p^2 = 0.28$; Greenhouse-Geisser

corrected, Fig.1A). On average, the happy BM induced a significantly larger pupil response than the neutral BM ($t(23) = 2.73$ $p = .024$, Cohen's $d = 0.56$, 95% CI for the mean difference = [0.037, 0.266]; Holm-corrected, Fig.1A). In contrast, the sad BM evoked a significantly smaller pupil size than the neutral BM (sad vs. neutral: $t(23) = -2.43$, $p = .024$, Cohen's $d = 0.50$, 95% CI for the mean difference = [-0.35, -0.03]; Holm-corrected, Fig.1A). Moreover, the happy BM evoked a significantly larger pupil size than the sad BM ($t(23) = 3.34$, $p = .009$, Cohen's $d = 0.68$, 95% confidence interval (CI) for the mean difference = [0.13, 0.55]; Holm-corrected, Fig.1A), which echoes with former studies showing a happiness advantage in BM processing (Actis-Grosso et al., 2015; Lee & Kim, 2016; Spencer et al., 2016; Yuan et al., 2023). Importantly, the observed happy over sad dilation effect is negatively correlated with the individual autistic traits ($r(23) = -0.47$, $p = .022$, 95% CI for the mean difference = [-0.73, -0.08], Fig.1B), indicating a compromised ability to perceive emotions from BM among individuals with higher autism tendencies. The results of the cluster-based permutation analysis showed that the happy BM induced a significant pupil dilation effect than the neutral BM from 1750 ms to 3200 ms (see Fig. 2A). Conversely, the sad BM evoked a significantly smaller pupil response than the neutral BM, which starts from 1900 ms until the end of the stimulus presentation (see Fig. 2A). Notably, the happy BM evoked a significantly larger pupil response as compared to the sad BM in a longstanding time window, ranging from 1200 ms until the disappearance of the display (see Fig. 2A). These results together demonstrated a pupil dilation effect caused by happy information embedded in the minimized point-light walker, reflecting that emotional BM can modulate pupil responses. Moreover, the happy and sad BM exerted differential modulation on the pupil responses: the happy BM evoked a larger pupil response than the neutral BM, while the sad BM evoked smaller pupil size as compared to the neutral one.

Experiment 2: Inverted emotional BM

To rule out the possibility that the difference in low-level visual features rather than the emotional information per se might account for the obtained emotional modulation effect, we presented observers with the inverted BM stimuli that shared the exact perceptual features with their upright counterparts in Experiment 2. An identical one-way repeated measures ANOVA was conducted, while no significant main effect of emotional condition on pupil responses was observed in inverted BM ($F(2, 46) = 0.95$, $p = .396$, $\eta_p^2 = 0.04$, Fig. 1C, Fig. 2B). Besides, the cluster-based permutation analysis also observed no significantly different time points among different emotional

conditions. Critically, the mixed 2 (orientation: upright, inverted) × 3 (emotional
condition: happy, sad, neutral) ANOVA on the average pupil size obtained in
Experiment 1 and Experiment 2 showed a significant interaction between orientation
and emotional condition ($F(2, 92) = 4.53$, $p = .013$, $\eta_p^2 = 0.09$). This further indicated
that the observed emotional modulation on pupil responses did not arise from low-level
visual differences.

Experiment 3: Nonbiological Motion

Given the ample evidence showing that local motion feature is essential for the
perception of biologically significant information from BM (Chang & Troje, 2008,
2009; Troje & Westhoff, 2006; Wang & Jiang, 2012), we moved forward to explore
the role of local motion feature in the emotion processing of BM. In Experiment 3, we
presented observers with non-BM stimuli, which were derived from the fragments
identical to emotional BM but with critical local characteristic removed (Chang & Troje,
2009). Results of the one-way repeated ANOVA on average pupil size showed no
significant main effect of emotional condition ($F(1.6, 35.7) = 0.02$, $p = .964$, $\eta_p^2 = 0.00$;
Greenhouse-Geisser corrected). Besides, the permutation analysis also found no
significantly different time points in pupil responses to the happy, sad, and neutral
conditions (see Fig. 1D, Fig. 2C). Moreover, combining the results obtained from
Experiment 1 and Experiment 3, we found a significant interaction between stimulus
type (upright, non-biological) and emotional condition (happy, sad, neutral) ($F(2, 92)$
$= 7.38$, $p = .001$, $\eta_p^2 = 0.14$). These findings together showed that the removal of local
characteristic greatly disrupted the emotional modulations on observers' pupil
responses, providing conceivable evidence for the critical contribution of local motion
feature in emotion perception from the BM signal.

Experiment 4: Local emotional BM

In Experiment 4, we went further to examine whether local BM alone could carry
emotional information and exert a similar modulation effect on pupil size. In particular,
we adopted the well-established local BM stimuli, the scrambled BM, whose local
motion feature was retained while the global configuration information was completely
disrupted. Both the explicit and implicit emotion processing of the scrambled BM were
investigated to provide a thorough view of emotion perception from local BM.
Specifically, a group of observers viewed the scrambled BM and made explicit
behavioral judgments on the emotional information contained in scrambled BM. The
results showed that observers could successfully recognize the emotions contained in

scrambled BMs with the average accuracy reaching significantly above 50% (M ± SD
= 83% ± 2.8%, $p < .001$). This indicated that scrambled BM conveyed recognizable
emotional information.

Furthermore, we investigated the pupil responses to scrambled happy, sad, and
neutral BMs to explore the automatic and implicit processing of local emotional BM.
The one-way repeated measures ANOVA on the mean pupil size indicated a significant
main effect of emotional condition ($F(2, 46) = 6.52$, $p = .003$, $\eta_p^2 = 0.22$, Fig. 1E).
Follow-up analysis showed that the scrambled happy BM induced a significantly larger
pupil response than the scrambled neutral BM ($t(23) = 3.26$, $p = .008$, Cohen's $d = 0.68$,
95% CI for the mean difference = [0.14, 0.59]; Holm-corrected, Fig. 1E), which is
similar to that observed in Experiment 1. Notably, the scrambled sad BM also induced
a larger pupil size than the neutral one ($t(23) = 3.35$, $p = .008$, Cohen's $d = 0.67$, 95%
CI for the mean difference = [0.14, 0.59]; Holm-corrected, Fig. 1E). Moreover, no
difference in pupil size is observed between the happy and sad conditions ($t(23) = -0.20$,
$p = .842$, Cohen's $d = 0.04$, 95% CI for the mean difference = [-0.28, 0.23]; Holm-
corrected, Fig. 1E). The consecutive cluster-based permutation analysis further showed
that both the scrambled happy and sad BM evoked larger pupil size than the scrambled
neutral BM. Importantly, such effect appeared in a rather early time window and could
last for a quite long time (happy vs. neutral: 450 ms -3550 ms; sad vs. neutral: 350 ms-
4000 ms, see Fig. 2D). The observed effect could not be accounted for by perceptual
differences (e.g., speed), as the happy and sad scrambled BM differed the most in low-
level features, whereas they induced similar dilation effects on pupil sizes. An
additional mixed 2 (BM type: intact, scrambled) × 3 (emotional condition: happy, sad,
neutral) ANOVA on average pupil size was conducted to examine whether or not this
emotional modulation effect of local BM varied from that of intact BM reported in
Experiment 1. Results revealed a significant interaction between BM type and
emotional condition ($F(2, 92) = 3.22$, $p = .045$, $\eta_p^2 = 0.07$), indicating that emotions in
intact and local BM exerted differential modulations on pupil size. Overall, these
findings showed that the local BM could convey significant emotional information,
which could induce a pupil dilation effect regardless of its exact type. As compared to
the intact BM, the emotional modulation observed in local BM is different because it
occurred faster and is rather coarse.

Discussion

Life motion signals convey salient emotional information that is crucial for human survival and social interactions (Johansson, 1973; Troje, 2008). Here, we reported that such emotional clues carried by the point-light BM walker could exert a modulation effect on pupil responses. Specifically, the happy BM significantly dilated pupils as compared to the neutral BM, and the sad BM evoked smaller pupil responses than the neutral BM, showing distinct emotional modulations on pupil responses which depend on the emotional contents. Moreover, this emotional modulation of pupil responses could not be explained by the low-level differences, as viewing inverted BMs failed to induce such effects. Importantly, when the emotional BM was deprived of the local motion feature through the removal of accelerations, it failed to induce any modulation on pupil responses. Furthermore, the scrambled emotional BM with only local motion feature retained could still produce a modulation effect on pupil size. Noticeably, this modulation is rapid but rather coarse: viewing the scrambled happy and sad BM would evoke greater pupil size than the scrambled neutral BM in a relatively early time window, while no significant difference was observed between the scrambled happy and sad BM. Taken together, these findings revealed multi-level processing of emotions in life motion signals: the global emotional BM modulated pupil size in a fine-grained manner that discriminates each type of emotion, and the local emotional BM exerted a coarse but rapid modulation on pupil size.

The emotion processing mechanisms of BM have been investigated in former studies using explicit behavioral detection and recognition paradigm. It has been reported that happy BM is more rapidly identified and detected as compared to sad and neutral BM (Actis-Grosso et al., 2015; Lee & Kim, 2016; Spencer et al., 2016), while other research observed no such superiority (Chouchourelou et al., 2006). Here, the present study adopted the novel and objective pupillometry index to investigate the emotion processing mechanisms of BM from a physiological aspect. The pupil index, as a direct reflection of individual arousal level and cognitive state, served as a powerful tool for faithfully and automatically reflecting the implicit emotion processing mechanism of BM. Our results showed a significant pupil dilation effect for happy over sad and neutral BM, showing a happiness superiority that could occur during the implicit emotion processing of BM. This finding resonates well with the happiness superiority reported in the explicit emotion processing of BM (Lee & Kim, 2016; Spencer et al., 2016; Yuan et al., 2023). More importantly, it provided a physiological account for the happiness superiority by demonstrating that happy BM evoked a more heightened state in the automatic nervous system than sad and neutral BM. The activity of this system has long been associated with the level of attention engagement into related cognitive

processes (Eckstein et al., 2017; Eldar et al., 2013). Hence, our findings suggest that
happy BM may be more effective in capturing attention as compared to neutral BM,
while sad BM could be less efficient than neutral BM. This observation parallels with
a previous study, which demonstrated that infants looked more at the happy BM walker
when displayed in pair with the neutral walker, whereas they attended less to the sad
walker as compared to the neutral one (Ogren et al., 2019). Notably, the happy over sad
dilation effect was negatively correlated with autistic traits: individuals with greater
autistic tendencies showed decreased sensitivities to emotions in BM. Given that pupil
measurement does not require any explicit verbal reports, it is easily attainable in
children and even infants. Thus, the pupil dilation effect obtained here may serve as a
sensitive and reliable physiological marker for detecting early social cognitive disorders.

Remarkably, our findings highlighted the central role of local motion feature in
modulating pupil responses towards emotional information contained in BM. Previous
studies have shown that human visual system is highly sensitive to the local BM
stimulus, whose global configural information is completely deprived through spatially
scrambling (Troje & Westhoff, 2006). For example, it has been demonstrated that
scrambled BM could perform as intact BM in lengthening subjective temporal
perception (Wang & Jiang, 2012). Besides, such local motion feature also served as a
basic pre-attentive feature in visual search (Wang et al., 2010) and could further induce
a significant reflexive attentional effect (Wang et al., 2014; Yu et al., 2020). Moreover,
the deprivation of the local BM feature would greatly disrupt the perception of the
contained biologically salient information, such as animacy and walking direction
(Chang & Troje, 2008, 2009; Yu et al., 2020). Here, we extended this line of research
to the emotional domain by demonstrating that the local BM component is also critical
for the processing of emotional information: when the local BM feature is disrupted,
the modulation of emotions on pupil size completely disappeared. Furthermore, the
emotional BM that retained only local motion features could still exert a salient
modulation effect on pupil size. In particular, the happy and sad local BM induced a
significant pupil dilation effect as compared to the neutral local BM stimuli.
Intriguingly, this dilation effect is independent of the specific emotion category but
reflects a general activation caused by the affective salient information. In addition, this
non-selective pupil dilation effect in local BM appeared in a relatively early time
window, indicating that the extraction of emotions from local BM is rapid. Taken
together, these findings identified a coarse but rapid emotion processing mechanism in
local BM that could promptly detect the emotional information therein without the aid
of the global shape.

Notably, our findings revealed the existence of distinct emotional modulations in
intact and local BM, respectively, as evidenced by variations in pupil responses. In
particular, the intact emotional point-light walker exerted a relatively late but fine-
grained modulation on pupil responses, while the local BM induced a quick but non-
selective emotional modulation on pupil sizes. The dissociated emotion processing
mechanisms could be well understood regarding its analogous function to another
important type of emotional signal (i.e., facial expression). Specifically, it has been
argued that there exist two parallel pathways for facial expression processing: a slow
route that conveys fine-grained facial features along cortical areas to the amygdala, and
a rapid subcortical pathway that directly transfers emotional information to amygdala
without detailed analysis, which is known as the "quick and dirty" route (Garrido et al.,
2012; Johnson, 2005; Méndez-Bértolo et al., 2016; Vuilleumier et al., 2003). It is
probable that the emotion processing of the local and intact point-light BM potentially
functions in a manner similar to that of the faces, with the former serving as a rapid
detection mechanism that automatically captures emotionally significant information
conveyed by animate agents without detailed analysis and the latter aiding in more
specific emotion identification based on fine-grained analyses of their motion pattern
and global shape. Compatible with this view, recent studies have reported that the
emotion processing of face and BM was very similar. For example, they both showed
a happiness superiority in visual detection (Becker et al., 2011; Lee & Kim, 2016;
Nackaerts et al., 2012) and guiding social attention (Yuan et al., 2023). Moreover, their
emotion processing involved potentially overlapping brain regions (e.g., amygdala,
STS) (Alaerts et al., 2014; Engell & Haxby, 2007; Peelen et al., 2007; Pessoa &
Adolphs, 2010). Overall, our findings, together with the former evidence, suggest
dissociated emotion processing mechanisms for BM in the human brain akin to the
dual-route model reported in faces, and further imply a general "emotional brain" that
can be shared by different types of social signals. Still, future studies are needed to
implement the neuroimaging techniques to directly identify the brain regions involved
in the emotion processing of local and global BM signals.

To conclude, the current study clearly demonstrated that the emotional information
conveyed by point-light BM modulated pupil responses. The intact BM exerted a fine-
grained emotional modulation on pupil sizes, while disrupting the contained local
motion characteristic would deteriorate the observed modulations. Moreover, BM with
only local motion features retained could exert a fast but rather coarse modulation on
pupillometry. These findings together highlight the critical role of local motion feature
in BM emotion processing, and further reveal the multi-level mechanisms subserving

emotion processing of BM. Notably, the observed emotional modulation in intact BM
is associated with individual autistic traits, suggesting the potential of utilizing the
emotion-modulated pupil responses to facilitate the diagnosis of social cognitive
disorders.

Methods

Participants

A total of 120 participants (44 males, 76 females) ranging from 18 to 29 years old (M
± SD = 23.1 ± 2.5) were recruited, with 24 in each experiment. All of the participants
had normal or corrected-to-normal vision and gave written informed consent in
accordance with the procedure and protocols approved by the institutional review board
of the Institute of Psychology, Chinese Academy of Sciences. They were all naïve to
the purpose of the experiments. Prior power analyses (*F* tests, repeated measures, within
factors) using G*Power (Version 3.1.9.4; (Faul et al., 2007)) indicated that a sample
size of 24 participants would afford 80% power with alpha at .05 to detect a moderate
main effect ($f = 0.27$). This sample size was comparable to previous studies with similar
designs (Nakakoga et al., 2020).

Stimuli

Stimuli were displayed using MATLAB (Mathworks, Inc.) together with the
Psychtoolbox extensions (Brainard, 1997; Pelli, 1997) on a 23-inch LED monitor (1920
× 1080 at 60 Hz) with gray background (RGB: 100, 100, 100). The biological motion
(BM) stimuli were adopted from Troje (Troje, 2008). Each comprised 15 point-light
dots depicting the motions of the head, pelvis, thorax and major joints (i.e., shoulders,
elbows, wrists, hips, knees, and ankles). Its emotional state was indexed by a
normalized Z score on an axis that reflects the differences between happy and sad
walkers in terms of a linear classifier. The scores were computed within a 10-
dimensional sub-space spanned by the first 10 principal components based on a Fourier-
based representation of observers' emotional ratings of 80 actual walkers (half male)
(Troje, 2008). We adopted the neutral walker that scored 0 on the linear axis, together
with the happy walker 6 SDs into the happy part of the axis and the sad walker 6 SDs
into the sad part of the axis (see https://www.biomotionlab.ca/html5-bml-walker/ for
an interactive animation). Besides, we turned the BM walkers 45 degrees leftwards or
rightwards to maximize the visibility of expressive bodily cues (Roether et al., 2009).
In Experiment 1, the upright emotional (happy, sad, or neutral) BM walkers with two

walking directions (45° leftwards or rightwards) were used. In Experiment 2, the stimuli were replaced with their inverted counterparts created by mirror flipping the upright BM vertically. In Experiment 3, we presented observers with the non-biological motion stimuli whose local BM characteristic was disrupted through acceleration removal. More specifically, each individual dot moved along the original path with a constant speed equal to the average speed of the dot (Chang & Troje, 2009). Such manipulation disrupted the local motion feature of the BM stimuli but kept the trajectories of individual point lights unchanged. In Experiment 4, observers viewed the scrambled BM stimuli, which were created by randomly relocating the point light dots within the region occupied by the original BM walker. In this manner, the local motion feature was preserved while the global configuration of the BM stimulus was entirely disrupted (Troje & Westhoff, 2006). Please refer to Supplementary Materials for the demostrations displaying the motion stimuli used in Experiments 1-4.

Procedure

Participants were seated at a viewing distance of 60 cm to the computer screen with their heads on a chinrest to minimize their head movements. Their pupil size and eye position of the left eye were recorded using a video-based iView X Hi-Speed system (SMI, Berlin, Germany) at 500 Hz. Each trial began with a central fixation cross (0.2° × 0.2°) with variable duration (800-1200 ms), followed by a happy/sad/neutral point-light walker (half leftwards and half rightwards) presented centrally for 4000 ms. Participants were required to passively view the BM stimulus and continued the procedure by pressing the space bar. After that, a blank screen was displayed for 3000 ms (Fig. 3). There were 4 blocks of 30 trials, and participants were given a short break after every block. Besides, we also administered a 9-point calibration of the eye-tracker followed by a validation stage before each experimental block to ensure the data quality. Participants were told to maintain fixation on the center of the screen and not to blink during the stimuli presentation. The experimental procedure of Experiment 2 is identical to Experiment 1, except that the BM stimuli were changed to their inverted counterparts. In Experiment 3, we instead presented participants with the non-biological motion displays whose local motion feature was deprived through acceleration removal. In Experiment 4, we adopted the scrambled emotional BM stimuli, and investigated both the explicit and the implicit emotion processing of scrambled BM. Specifically, a group of participants was recruited to make emotional judgements on the scrambled BM in order to investigate whether the emotional information carried by local BM signals could be explicitly identified. Another group

of participants was enrolled for the pupil recording experiment, which followed the identical experimental procedure of Experiments 1-3. At the end of all experiments, participants were required to complete the 50-point Autism-spectrum Questionnaire (AQ), which measured the degree of autistic traits in the normal population (Baron-Cohen et al., 2001).

Data Analysis

The raw pupil data for each trial was first screened to remove eye blinks (either replaced by linear interpolation or with this trial discarded). Then, trials with pupil size deviating ± 3 SDs from the mean were excluded from further analysis. Finally, the pupil size data were down-sampled to 20 Hz and baseline-corrected for each trial by subtracting the mean pupil size during the 200 ms pre-stimulus period. We computed the average pupil size for each emotional condition (happy, sad, or neutral) obtained by collapsing the pupillometry across all time points. Besides, to depict the time course of pupil responses towards emotional BM, we further conducted a consecutive paired-sample *t*-test across all time points comparing different emotional conditions. The cluster-based permutation analysis was applied to avoid potential problems brought about by multiple comparisons (Einhauser et al., 2008). In this analysis, the computed *t*-values neighboring in time that exceeded a threshold ($p < 0.05$) were defined as clusters, and then summed to produce a cluster mass. The cluster mass was compared with a null distribution, which was generated by 2000 random permutations of the pupil data from different conditions. If the cluster mass fell beyond 95% of the null distribution ($\alpha = 0.05$), it was deemed to be statistically significantly different (Einhauser et al., 2008). The pupil size was analyzed and reported in arbitrary unit (a.u.) without transforming into the actual unit (mm), as the relative change of the pupil size was of main interest.

Acknowledgements

This research was supported by grants from the Ministry of Science and Technology of China (STI2030-Major Projects 2022ZD0205100 and 2021ZD0203800), the National Natural Science Foundation of China (No. 31830037), the Strategic Priority Research Program (No. XDB32010300), the Interdisciplinary Innovation Team (JCTD-2021-06), the Science Foundation of Institute of Psychology, Chinese Academy of Sciences, and the Fundamental Research Funds for the Central Universities. We thank Professor Nikolaus F. Troje for kindly providing us with the visual stimuli. We report how we determined our sample size, all data exclusions (if any), all manipulations, and all measures in the study. This study's design and its analysis were not pre-registered. All

data, materials, and analysis code used in the current study could be accessed at Science Data Bank (https://doi.org/10.57760/sciencedb.psych.00125). The authors declared no conflicts of interest with respect to the authorship or the publication of this article.

References:

Actis-Grosso, R., Bossi, F., & Ricciardelli, P. (2015). Emotion recognition through static faces and moving bodies: a comparison between typically developed adults and individuals with high level of autistic traits. *Frontiers in Psychology*, *6*, 1570. https://doi.org/10.3389/fpsyg.2015.01570

Alaerts, K., Nackaerts, E., Meyns, P., Swinnen, S. P., & Wenderoth, N. (2011). Action and emotion recognition from point light displays: an investigation of gender differences. *PLoS ONE*, *6*(6), e20989.

Alaerts, K., Woolley, D. G., Steyaert, J., Di Martino, A., Swinnen, S. P., & Wenderoth, N. (2014). Underconnectivity of the superior temporal sulcus predicts emotion recognition deficits in autism. *Social Cognitive and Affective Neuroscience*, *9*(10), 1589–1600. https://doi.org/10.1093/scan/nst156

Baron-Cohen, S., Wheelwright, S., Skinner, R., Martin, J., & Clubley, E. (2001). The autism-spectrum quotient (aq): evidence from asperger syndrome/high-functioning autism, malesand females, scientists and mathematicians. *Journal of Autism and Developmental Disorders*. https://doi.org/10.1023/a:1005653411471

Becker, D. V., Anderson, U. S., Mortensen, C. R., Neufeld, S. L., & Neel, R. (2011). The face in the crowd effect unconfounded: happy faces, not angry faces, are more efficiently detected in single- and multiple-target visual search tasks. *Journal of Experimental Psychology: General*, *140*(4), 637–659. https://doi.org/10.1037/a0024060

Brainard, D. (1997). The psychophysics toolbox. *Spatial Vision*, *10*(4), 433–436.

Burley, D. T., & Daughters, K. (2020). The effect of oxytocin on pupil response to naturalistic dynamic facial expressions. *Hormones and Behavior*, *125*, 104837. https://doi.org/10.1016/j.yhbeh.2020.104837

Burley, D. T., Gray, N. S., & Snowden, R. J. (2017). As far as the eye can see: relationship between psychopathic traits and pupil response to affective stimuli. *PLOS ONE*, *12*(1), e0167436. https://doi.org/10.1371/journal.pone.0167436

Chang, D. H. F., & Troje, N. F. (2008). Perception of animacy and direction from local biological motion signals. *Journal of Vision*, *8*(5), 3. https://doi.org/10.1167/8.5.3

Chang, D. H. F., & Troje, N. F. (2009). Acceleration carries the local inversion effect in biological motion perception. *Journal of Vision*, *9*(1), 19–19. https://doi.org/10.1167/9.1.19

Chouchourelou, A., Matsuka, T., Harber, K., & Shiffrar, M. (2006). The visual analysis of emotional actions. *Social Neuroscience*, *1*(1), 63–74. https://doi.org/10.1080/17470910600630599

Critchley, H., Daly, E., Phillips, M., Brammer, M., Bullmore, E., Williams, S., Van Amelsvoort, T., Robertson, D., David, A., & Murphy, D. (2000). Explicit and implicit neural mechanisms for processing of social information from facial expressions: a functional magnetic resonance imaging study. *Human Brain Mapping*, *9*(2), 93–105. https://doi.org/10.1002/(sici)1097-0193(200002)9:2<93::aid-hbm4>3.0.co;2-z

de Gee, J. W., Knapen, T., & Donner, T. H. (2014). Decision-related pupil dilation reflects upcoming choice and individual bias. *Proceedings of the National Academy of Sciences*, *111*(5), E618–E625. https://doi.org/10.1073/pnas.1317557111

de Gelder, B. (2006). Towards the neurobiology of emotional body language. *Nature Reviews Neuroscience*, *7*(3), 242–249. https://doi.org/10.1038/nrn1872

Eckstein, M. K., Guerra-Carrillo, B., Miller Singley, A. T., & Bunge, S. A. (2017). Beyond eye gaze: what else can eyetracking reveal about cognition and cognitive development? *Developmental Cognitive Neuroscience*, *25*, 69–91. https://doi.org/10.1016/j.dcn.2016.11.001

Einhauser, W., Stout, J., Koch, C., & Carter, O. (2008). Pupil dilation reflects perceptual selection and predicts subsequent stability in perceptual rivalry. *Proceedings of the National Academy of Sciences*, *105*(5), 1704–1709. https://doi.org/10.1073/pnas.0707727105

Eldar, E., Cohen, J. D., & Niv, Y. (2013). The effects of neural gain on attention and learning. *Nature Neuroscience*, *16*(8), 1146–1153. https://doi.org/10.1038/nn.3428

Engell, A. D., & Haxby, J. V. (2007). Facial expression and gaze-direction in human superior temporal sulcus. *Neuropsychologia*, *45*(14), 3234–3241. https://doi.org/10.1016/j.neuropsychologia.2007.06.022

Faul, F., Erdfelder, E., Lang, A.-G., & Buchner, A. (2007). G*Power 3: a flexible statistical power analysis program for the social, behavioral, and biomedical sciences. *Behavior Research Methods*, *39*(2), 175–191. https://doi.org/10.3758/bf03193146

Frith, C. (2009). Role of facial expressions in social interactions. *Philosophical Transactions of the Royal Society B: Biological Sciences*, *364*(1535), 3453–3458. https://doi.org/10.1098/rstb.2009.0142

Frith, U. (2004). Emanuel miller lecture: confusions and controversies about asperger syndrome. *Journal of Child Psychology and Psychiatry*, *45*(4), 672–686. https://doi.org/10.1111/j.1469-7610.2004.00262.x

Garrido, M. I., Barnes, G. R., Sahani, M., & Dolan, R. J. (2012). Functional evidence for a dual route to amygdala. *Current Biology*, *22*(2), 129–134. https://doi.org/10.1016/j.cub.2011.11.056

Graves, J. E., Egré, P., Pressnitzer, D., & de Gardelle, V. (2021). An implicit representation of stimulus ambiguity in pupil size. *Proceedings of the National Academy of Sciences*, *118*(48), e2107997118.https://doi.org/10.1073/pnas.2107997118

Harms, M. B., Martin, A., & Wallace, G. L. (2010). Facial emotion recognition in autism spectrum disorders: a review of behavioral and neuroimaging studies. *Neuropsychology Review*, *20*(3), 290–322. https://doi.org/10.1007/s11065-010-9138-6

Hubert, B., Wicker, B., Moore, D. G., Monfardini, E., Duverger, H., Fonséca, D. D., & Deruelle, C. (2006). Brief report: recognition of emotional and non-emotional biological motion in individuals with autistic spectrum disorders. *Journal of Autism and Developmental Disorders*, *37*(7), 1386–1392. https://doi.org/10.1007/s10803-006-0275-y

Johansson, G. (1973). Visual perception of biological motion and a model for its analysis. *Perception & Psychophysics*, *14*(2), 201–211. https://doi.org/10.3758/bf03212378

Johnson, M. H. (2005). Subcortical face processing. *Nature Reviews Neuroscience*, *6*(10), 766–774. https://doi.org/10.1038/nrn1766

Joshi, S., & Gold, J. I. (2019). Pupil size as a window on neural substrates of cognition. *Trends in Cognitive Sciences*, *24*(6), 466–480. https://doi.org/10.31234/osf.io/dvsme

Kana, R. K., Patriquin, M. A., Black, B. S., Channell, M. M., & Wicker, B. (2015). Altered medial frontal and superior temporal response to implicit processing of emotions in autism. *Autism Research*, *9*(1), 55–66. https://doi.org/10.1002/aur.1496

Keifer, C. M., Mikami, A. Y., Morris, J. P., Libsack, E. J., & Lerner, M. D. (2020). Prediction of social behavior in autism spectrum disorders: explicit versus implicit social cognition. *Autism*, *24*(7), 1758–1772. https://doi.org/10.1177/1362361320922058

Kloosterman, N. A., Meindertsma, T., van Loon, A. M., Lamme, V. A. F., Bonneh, Y. S., & Donner, T. H. (2015). Pupil size tracks perceptual content and surprise. *European Journal of Neuroscience*, *41*(8), 1068–1078. https://doi.org/10.1111/ejn.12859

Kovarski, K., Mennella, R., Wong, S. M., Dunkley, B. T., Taylor, M. J., & Batty, M. (2018). Enhanced early visual responses during implicit emotional faces processing in autism spectrum disorder. *Journal of Autism and Developmental Disorders*, *49*(3), 871–886. https://doi.org/10.1007/s10803-018-3787-3

Lange, K., Williams, L. M., Young, A. W., Bullmore, E. T., Brammer, M. J., Williams, S. C. R., Gray, J. A., & Phillips, M. L. (2003). Task instructions modulate neural responses to fearful facial expressions. *Biological Psychiatry*, *53*(3), 226–232. https://doi.org/10.1016/s0006-3223(02)01455-5

Lee, H., & Kim, J. (2016). Facilitating effects of emotion on the perception of biological motion: evidence for a happiness superiority effect. *Perception*, *46*(6), 679–697. https://doi.org/10.1177/0301006616681809

Méndez-Bértolo, C., Moratti, S., Toledano, R., Lopez-Sosa, F., Martínez-Alvarez, R., Mah, Y. H., Vuilleumier, P., Gil-Nagel, A., & Strange, B. A. (2016). A fast pathway for fear in human amygdala. *Nature Neuroscience*, *19*(8), 1041–1049. https://doi.org/10.1038/nn.4324

Nackaerts, E., Wagemans, J., Helsen, W., Swinnen, S. P., Wenderoth, N., & Alaerts, K. (2012). Recognizing biological motion and emotions from point-light displays in autism spectrum disorders. *PLoS ONE*, *7*(9), e44473. https://doi.org/10.1371/journal.pone.0044473

Nakakoga, S., Higashi, H., Muramatsu, J., Nakauchi, S., & Minami, T. (2020). Asymmetrical characteristics of emotional responses to pictures and sounds: evidence from pupillometry. *PLOS ONE*, *15*(4), e0230775. https://doi.org/10.1371/journal.pone.0230775

Ogren, M., Kaplan, B., Peng, Y., Johnson, K. L., & Johnson, S. P. (2019). Motion or emotion: infants discriminate emotional biological motion based on low-level visual information. *Infant Behavior and Development*, *57*, 101324. https://doi.org/10.1016/j.infbeh.2019.04.006

Okon-Singer, H., Lichtenstein-Vidne, L., & Cohen, N. (2013). Dynamic modulation of emotional processing. *Biological Psychology*, *92*(3), 480–491. https://doi.org/10.1016/j.biopsycho.2012.05.010

Oliva, M., & Anikin, A. (2018). Pupil dilation reflects the time course of emotion recognition in human vocalizations. *Scientific Reports*, *8*(1), 4871. https://doi.org/10.1038/s41598-018-23265-x

Peelen, M. V., Atkinson, A. P., Andersson, F., & Vuilleumier, P. (2007). Emotional modulation of body-selective visual areas. *Social Cognitive and Affective Neuroscience*, *2*(4), 274–283. https://doi.org/10.1093/scan/nsm023

Pelli, D. (1997). The videotoolbox software for visual psychophysics: transforming numbers into movies. *Spatial Vision*.

Pessoa, L., & Adolphs, R. (2010). Emotion processing and the amygdala: from a "low road" to "many roads" of evaluating biological significance. *Nature Reviews Neuroscience*, *11*(11), 773–782. https://doi.org/10.1038/nrn2920

Prunty, J. E., Keemink, J. R., & Kelly, D. J. (2021). Infants show pupil dilatory responses to happy and angry facial expressions. *Developmental Science*, *25*(2). https://doi.org/10.1111/desc.13182

Roether, C. L., Omlor, L., Christensen, A., & Giese, M. A. (2009). Critical features for the perception of emotion from gait. *Journal of Vision*, *9*(6), 15–15. https://doi.org/10.1167/9.6.15

Shafer, A. T., Matveychuk, D., Penney, T., O'Hare, A. J., Stokes, J., & Dolcos, F. (2012). Processing of emotional distraction is both automatic and modulated by attention: evidence from an event-related fmri investigation. *Journal of Cognitive Neuroscience*, *24*(5), 1233–1252. https://doi.org/10.1162/jocn_a_00206

Spencer, J. M. Y., Sekuler, A. B., Bennett, P. J., Giese, M. A., & Pilz, K. S. (2016). Effects of aging on identifying emotions conveyed by point-light walkers. *Psychology and Aging*, *31*(1), 126–138. https://doi.org/10.1037/a0040009

Troje, N. F. (2008). *Retrieving Information from Human Movement Patterns* (pp. 308–334). Oxford University PressNew York. https://doi.org/10.1093/acprof:oso/9780195188370.003.0014

Troje, N. F., & Westhoff, C. (2006). The inversion effect in biological motion perception: evidence for a "life detector"? *Current Biology*, *16*(8), 821–824. https://doi.org/10.1016/j.cub.2006.03.022

Vuilleumier, P., Armony, J. L., Driver, J., & Dolan, R. J. (2003). Distinct spatial frequency sensitivities for processing faces and emotional expressions. *Nature Neuroscience*, *6*(6), 624–631. https://doi.org/10.1038/nn1057

Wang, L., & Jiang, Y. (2012). Life motion signals lengthen perceived temporal duration. *Proceedings of the National Academy of Sciences*, *109*(11), E673–E677. https://doi.org/10.1073/pnas.1115515109

Wang, L., Yang, X., Shi, J., & Jiang, Y. (2014). The feet have it: local biological motion cues trigger reflexive attentional orienting in the brain. *NeuroImage*, *84*, 217–224. https://doi.org/10.1016/j.neuroimage.2013.08.041

Wang, L., Zhang, K., He, S., & Jiang, Y. (2010). Searching for life motion signals. *Psychological Science*, *21*(8), 1083–1089. https://doi.org/10.1177/0956797610376072

Wong, T. K. W., Fung, P. C. W., Chua, S. E., & McAlonan, G. M. (2008). Abnormal spatiotemporal processing of emotional facial expressions in childhood autism: dipole source analysis of event-related potentials. *European Journal of Neuroscience*, *28*(2), 407–416. https://doi.org/10.1111/j.1460-9568.2008.06328.x

Yu, Y., Ji, H., Wang, L., & Jiang, Y. (2020). Cross-modal social attention triggered by biological motion cues. *Journal of Vision*, *20*(10), 21. https://doi.org/10.1167/jov.20.10.21

Yuan, T., Ji, H., Wang, L., & Jiang, Y. (2023). Happy is stronger than sad: Emotional information modulates social attention. *Emotion*, *23*(4), 1061–1074. https://doi.org/10.1037/emo0001145

Figures:

Fig. 1. Normalized mean pupil responses using the neutral condition as baseline, plotted against happy and sad

conditions, and the relevant correlation. (A) In Experiment 1, the group average pupil response to happy intact BM

is significantly larger than that to sad and neutral BM, while the pupil size induced by sad BM is significantly smaller

than that evoked by neutral BM. (B) Moreover, a significant negative correlation was found between the happy over

sad pupil dilation effect and individual autistic quotient (AQ). (C) In Experiment 2, no significant differences in

pupil responses were observed for inverted BM. (E) In Experiment 3, when the biological characteristic was deprived

from the emotional BM, it failed to induce any modulations on pupil sizes. (D) In Experiment 4, both the happy and

sad local BM induced a significantly larger pupil size than neutral local BM, with no significant difference between

the happy and sad condition. Green dots in the scatter plot indicate the individual data and the shaded region indicate

the 95% confidence interval. All the pupil data are in arbitrary units (a.u.). Error bars showed standard errors of the

mean. * $p < .05$, ** $p < .01$.

Fig. 2. Time course of pupil responses to happy, sad, and neutral BM in Experiments 1-4. Solid lines represent pupil diameter under each emotional condition as a function of time (happy: red; sad: blue; neutral: gray); shaded areas represent the SEM between participants; colored horizontal lines indicate periods during which there are statistically significant differences among conditions at $p < 0.05$; and black horizontal lines indicate significant differences after cluster-based permutation correction. All the pupil data are in arbitrary units (a.u.) (A) In Experiment 1, the happy BM evoked larger pupil response as compared to the sad and neutral BM, and the sad BM evoked smaller pupil size than the neutral BM. (B) In Experiment 2, the inverted BM failed to produce such emotional modulation effects. (C) In Experiment 3, the emotional BM that is deprived of the local motion feature exerted no emotional modulation on pupil responses. (D) In Experiment 4, both the happy and sad local BM induced a larger pupil response than the neutral local BM, and such dilation effect started from a relatively early time point.

Fig. 3. Schematic representation of the experimental procedure and the stimuli in Experiment 1. A happy/sad/neutral BM walker turning 45 degrees leftwards or rightwards was presented at the center of the screen for 4000 ms. Participants were instructed to maintain their fixation on the BM stimuli during stimulus presentation and to continue the procedure through key pressing.