1 Title page

Multi-level processing of emotions in life motion signals revealed through pupil responses

Tian Yuan[1,2,3], Li Wang[1,2,3]*, Yi Jiang[1,2,3]

[1]State Key Laboratory of Brain and Cognitive Science, CAS Center for Excellence in

Brain Science and Intelligence Technology, Institute of Psychology, Chinese Academy

of Sciences, 16 Lincui Road, Beijing 100101, China; [2]Department of Psychology,

University of Chinese Academy of Sciences, 19A Yuquan Road, Beijing 100049, China;

[3]Chinese Institute for Brain Research, 26 Science Park Road, Beijing 102206, China

Running title: Pupil Response to Emotion

Key words: pupil size, biological motion, emotion processing, local

*Corresponding authors: Li Wang (wangli@psych.ac.cn)

Institute of Psychology, Chinese Academy of Sciences, 16 Lincui Road, Beijing

100101, China

Abstract

Perceiving emotions from the movements of other biological entities is critical for
human survival and interpersonal interactions. Here, we report that emotional
information conveyed by point-light biological motion (BM) triggered automatic
physiological responses as reflected in pupil size. Specifically, happy BM evoked larger
pupil size than neutral and sad BM, while sad BM induced a smaller pupil response
than neutral BM. Moreover, this happy over sad pupil dilation effect is negatively
correlated with individual autistic traits. Notably, emotional BM with only local motion
features retained could also exert modulations on pupils. Compared with intact BM,
both happy and sad local BM evoked stronger pupil responses than neutral local BM
starting from an earlier timepoint, with no difference between the happy and sad
conditions. These results revealed a fine-grained pupil-related emotional modulation
induced by intact BM and a coarse but rapid modulation by local BM, demonstrating
multi-level processing of emotions in life motion signals. Taken together, our findings
shed new light on BM emotion processing, and highlight the potential of utilizing the
emotion-modulated pupil response to facilitate the diagnosis of social cognitive
disorders.

Introduction

Perceiving and interpreting emotions from various social signals is critical for human social functioning, which enables us to infer the intentions of our conspecifics and further facilitates interpersonal interactions. Facial expressions present the most common non-verbal social communicative signals regarding others' affective states and intentions (C. Frith, 2009). In addition to faces, the movement of biological organisms serves as another essential type of social signal carrying significant emotional information, and this information remained salient even from a far distance (de Gelder, 2006). The human visual system is highly sensitive to such signals that we can readily decipher emotions from biological motion (BM), even when it was portrayed by several point lights attached to the major joints (Johansson, 1973; Troje, 2008). Moreover, it has been found that happy point-light walkers were recognized faster and more accurately than sad, angry or neutral walkers, demonstrating a happiness superiority (Lee & Kim, 2016; Spencer et al., 2016). While these studies provided some insights into the emotion processing mechanism of BM, they relied mainly on the explicit identification and active evaluation of BM emotions. Importantly, the encoding of emotional information also involves a rather automatic and implicit process that is independent of the participant's explicit identifications (Critchley et al., 2000; Lange et al., 2003; Okon-Singer et al., 2013; Shafer et al., 2012). Notably, this implicit aspect of emotion processing could be even more effective in probing individual differences and social deficits (Kana et al., 2015; Keifer et al., 2020; Kovarski et al., 2018; Wong et al., 2008), as it requires the intuitive processing of emotions that could not be learned (U. Frith, 2004). For example, research has shown that individuals with autistic disorders showed altered neural activities during implicit but not explicit emotion processing of natural scenes (Kana et al., 2015). These observations highlighted the importance of using objective measurements to investigate the implicit and automatic aspect of BM emotion processing.

The pupil response hence serves as a promising approach for reliably unfolding the implicit emotion processing of BM as it adds no extra task requirements to the cognitive process (Burley et al., 2017). In particular, pupil size is related to the activity of the automatic nervous system mediated by the locus coeruleus norepinephrine (LC-NE), which not only responds to physical light, but also reflects the underlying cognitive state (Joshi & Gold, 2019). Moreover, it could spontaneously capture the current subjective state and is thus useful for revealing the time course of the related cognitive processing (de Gee et al., 2014; Graves et al., 2021; Kloosterman et al., 2015;

Oliva & Anikin, 2018). Recently, this measurement has been introduced to the field of
emotion perception, and emerging evidence has suggested that emotions conveyed by
social signals (e.g., faces, voices) could modulate pupil responses. For instance, happy
and angry dynamic faces elicited a pupil dilation effect as compared with sad and
neutral faces (Burley & Daughters, 2020; Prunty et al., 2021). Noticeably, the point-
light BM, as another critical type of social signal, also conveyed salient affective
information. Besides, its emotion processing mechanism is closely connected with that
of faces (Alaerts et al., 2011; Becker et al., 2011; Lee & Kim, 2016; Yuan et al., 2024).
However, it remains unequivocal whether the pupil also responds to the emotional
information carried by the minimalistic point-light walker display. Such physiological
observation can faithfully uncover the implicit emotion processing of BM, and it also
expands the existing line of inquiry on the emotion processing of social cues.

To fill this gap, the current study implemented the pupil recording technique
together with the passive viewing paradigm to explore the automatic and implicit
emotion processing of BM. We first investigated the pupil responses to point-light
walkers with different emotions (i.e., happy, sad, and neutral). In addition to intact
emotional BM sequences, we also tested scrambled emotional BM sequences, which
lack the gestalt of a global figure but preserve the same local motion components as the
intact walkers. Recent studies have shown that such local motion signals play an
essential role in conveying biologically salient information such as animacy, and
walking direction (Chang & Troje, 2008, 2009; Troje & Westhoff, 2006). This study
went further to investigate whether this local BM signal could convey emotional
information and elicit corresponding pupil responses. Moreover, given that emotion
perception from social signals is generally impaired in individuals with autism (Harms
et al., 2010; Hubert et al., 2006), we also took the individual autistic traits into
consideration by measuring the autistic tendencies in normal populations with the
Autistic Quotient (AQ) questionnaire (Baron-Cohen et al., 2001).

Results

Experiment 1a-b: Intact emotional BM

In Experiment 1a, we investigated whether emotional BM could automatically exert
influences on pupil responses. Specifically, participants were instructed to passively
view the intact happy/sad/neutral BM with their pupil responses being recorded
simultaneously (see Fig.1). A cluster-based permutation analysis was applied to
illustrate the time course of emotional modulations on pupil responses. Results showed

that the happy BM induced a significant pupil dilation effect than the neutral BM from 1750 ms to 3200 ms (see Fig.2A). Conversely, the sad BM evoked a significantly smaller pupil response than the neutral BM, which starts from 1850 ms until the end of the stimulus presentation (see Fig. 2A). Moreover, the happy BM evoked a significantly larger pupil response as compared to the sad BM in a longstanding time window, ranging from 1200 ms until the disappearance of the display (see Fig.2A).

We then computed the mean pupil size by collapsing the pupillometry across all time points, ranging from the onset of the stimuli to the end of the stimuli presentation. A one-way repeated measures ANOVA was conducted on the mean pupil size for each emotional condition (happy, sad, neutral), and the results showed a significant main effect of emotional condition ($F(1.4, 32.2) = 8.92$, $p = .002$, $\eta_p^2 = 0.28$; Greenhouse-Geisser corrected, Fig.3A). On average, the happy BM induced a significantly larger pupil response than the neutral BM ($t(23) = 2.73$ $p = .024$, Cohen's $d = 0.56$, 95% confidence interval (CI) for the mean difference = [0.04, 0.27]; Holm-corrected, $p = .036$ after Bonferroni correction, Fig.3A). In contrast, the sad BM evoked a significantly smaller pupil size than the neutral BM (sad vs. neutral: $t(23) = -2.43$, $p = .024$, Cohen's $d = 0.50$, 95% CI for the mean difference = [-0.35, -0.03]; Holm-corrected, $p = .071$ after Bonferroni correction, Fig.3A). Moreover, the happy BM evoked a significantly larger pupil size than the sad BM ($t(23) = 3.34$, $p = .009$, Cohen's $d = 0.68$, 95% CI for the mean difference = [0.13, 0.55]; Holm-corrected, $p = .009$ after Bonferroni correction, Fig.3A), which echoes with former studies showing a happiness advantage in BM processing (Actis-Grosso et al., 2015; Lee & Kim, 2016; Spencer et al., 2016; Yuan et al., 2023). Importantly, the observed happy over sad dilation effect is negatively correlated with the individual autistic traits ($r(23) = -0.47$, $p = .022$, 95% CI for the mean difference = [-0.73, -0.08], Fig. 4A), indicating a compromised ability to perceive emotions from BM among individuals with higher autism tendencies. No significant correlations were found between AQ and other pupil modulation effects (Fig. 4B-C). Additionally, no significant correlations were observed between AQ and the original pupil responses induced by happy/neutral/sad BM (see Supplementary Material for more details).

To strengthen the correlation results, we conducted a replication experiment (Experiment 1b) and added a test-retest examination to further assess the reliability of our measurements. Specifically, a new group of participants were recruited to perform the identical task, and they were asked to return to the lab for a retest. The results again revealed the main effect of emotional condition in both the first test ($F(2, 46) = 12.0$, p

< .001, η_p^2 = 0.34, Fig.3B) and the second test ($F(2, 46)$ = 14.8, p < .001, η_p^2 = 0.39,

Fig.3C). The happy BM induced a significantly larger pupil response than the neutral

BM (First Test: $t(23)$ = 2.60, p = .022, Cohen's d = 0.53, 95% CI for the mean difference

= [0.02, 0.14], Holm-corrected, p = .048 after Bonferroni correction, Fig.3B; Second

Test: $t(23)$ = 3.36, p = .005, Cohen's d = 0.68, 95% CI for the mean difference = [0.06,

0.24], Holm-corrected, p = .008 after Bonferroni correction, Fig.3C). On the contrary,

the sad BM induced a significantly smaller pupil response than the neutral BM (First

Test: $t(23)$ = -2.77, p = .022, Cohen's d = 0.57, 95% CI for the mean difference = [-

0.19, -0.03], Holm-corrected, p = .033 after Bonferroni correction, Fig.3B; Second Test:

$t(23)$ = -3.19, p = .005, Cohen's d = 0.65, 95% CI for the mean difference = [-0.24, -

0.05], Holm-corrected, p = .012 after Bonferroni correction, Fig.3C). Besides, the

happy BM induced significantly larger pupil response than the sad BM (First Test: $t(23)$

= 4.23, p < .001, Cohen's d = 0.86, 95% CI for the mean difference = [0.10, 0.28],

Holm-corrected, p < .001 after Bonferroni correction, Fig.3B; Second Test: $t(23)$ = 4.26,

p < .001, Cohen's d = 0.87, 95% CI for the mean difference = [0.15, 0.44], Holm-

corrected, p < .001 after Bonferroni correction, Fig.3C). The results of the cluster-based

permutation analysis were also similar (see Supplementary Material for more details).

Notably, we successfully replicated the negative correlation between the happy

over sad dilation effect and individual autistic traits ($r(23)$ = -0.46, p = .023, 95% CI

for the mean difference = [-0.73, -0.07], Fig.4D) in the first test of Experiment 1b. No

other significant correlations were found (see Fig.4E-F). Moreover, in the second test,

such a correlation was similarly found and was even stronger ($r(23)$ = -0.61, p = .002,

95% CI for the mean difference = [-0.81, -0.27], Fig.4G). A test-retest reliability

analysis was performed on the happy over sad pupil dilation effect and the AQ score,

and the results showed robust correlations (r(happy-sad pupil size)= 0.56; r(AQ)= 0.90)

and strong test-retest reliabilities (α(happy-sad pupil size)= 0.60; α(AQ)= 0.82).

Furthermore, in the second test, we've additionally observed a significant negative

correlation between AQ and the happy minus neutral pupil dilation effect ($r(23)$ = -0.44,

p = .032, 95% CI for the mean difference = [-0.72, -0.04], Fig.4H), and a significant

positive correlation between the sad minus neutral pupil size and AQ ($r(23)$ = 0.50, p

= .014, 95% CI for the mean difference = [0.12, 0.75], Fig.4I). This indicated that the

overall correlation between happy over sad dilation effect and AQ was the aggregate

outcome of the diminished pupil modulations by happy and sad BM for high AQ

individuals.

Overall, these results demonstrated a robust and replicable modulation of emotions
embedded in the minimized point-light walker on pupil sizes: the happy BM evoked a
larger pupil response than the neutral BM, while the sad BM evoked a smaller pupil
size as compared to the neutral one. Importantly, a significant negative correlation
between AQ and happy over sad pupil dilation effect was consistently found, and such
effect was caused by a general attenuation in BM emotion perception sensitivity among
individuals with high autistic tendencies.

Experiment 2: Inverted emotional BM

To rule out the possibility that the difference in low-level visual features rather than the
emotional information per se might account for the obtained emotional modulation
effect, we presented observers with the inverted BM stimuli that shared the exact
perceptual features with their upright counterparts in Experiment 2. The cluster-based
permutation analysis observed no significantly different time points among three
emotional conditions (see Fig. 2B). An identical one-way repeated measures ANOVA
was conducted, while no significant main effect of emotional condition on pupil
responses was observed in inverted BM ($F(2, 46) = 0.95$, $p = .396$, $\eta_p^2 = 0.04$, Fig. 3D).
Besides, no significant correlations between AQ and pupil modulations were found
(AQ with happy-sad pupil size: $r(23) = 0.14$, $p = .512$, 95% CI for the mean difference
= [-0.28, 0.52]; AQ with happy-neutral pupil size: $r(23) = 0.26$, $p = .227$, 95% CI for
the mean difference = [-0.16, 0.60]; AQ with sad-neutral pupil size: $r(23) = 0.19$, p
= .376, 95% CI for the mean difference = [-0.23, 0.55]). Critically, the mixed 2
(orientation: upright, inverted) × 3 (emotional condition: happy, sad, neutral) ANOVA
on the average pupil size obtained in Experiment 1a and Experiment 2 showed a
significant interaction between orientation and emotional condition ($F(2, 92) = 4.38$, p
= .013, $\eta_p^2 = 0.09$), which could be attributed to the diminishment of emotional
modulation in inverted BM. This indicated that the observed emotional modulation of
pupil responses did not arise from low-level visual differences.

Experiment 3: Nonbiological Motion

Given the ample evidence showing that local motion feature is essential for the
perception of biologically significant information from BM (Chang & Troje, 2008,
2009; Troje & Westhoff, 2006; L. Wang & Jiang, 2012), we moved forward to explore
the role of local motion feature in the emotion processing of BM. In Experiment 3, we
presented observers with non-BM stimuli, which were derived from the fragments
identical to emotional BM but with critical local characteristics removed (Chang &

Troje, 2009). The permutation analysis found no significantly different time points in
pupil responses to the happy, sad, and neutral conditions (see Fig. 2D). Results of the
one-way repeated ANOVA on the average pupil sizes also showed no significant main
effect of emotional condition ($F(1.6, 35.7) = 0.02$, $p = .964$, $\eta_p^2 = 0.00$; Greenhouse-
Geisser corrected, see Fig. 3C). Similarly, no pupil modulation effects significantly
correlated with AQ (AQ with happy-sad pupil size: $r(23) = -0.03$, $p = .896$, 95% CI for
the mean difference = [-0.43, 0.38]; AQ with happy-neutral pupil size: $r(23) = 0.01$, p
$= .956$, 95% CI for the mean difference = [-0.39, 0.41]; AQ with sad-neutral pupil size:
$r(23) = 0.06$, $p = .777$, 95% CI for the mean difference = [-0.35, 0.45]). Moreover,
combining the results obtained from Experiment 1a and Experiment 3, we found a
significant interaction between stimulus type (intact, nonbiological) and emotional
condition (happy, sad, neutral) ($F(2, 92) = 3.22$, $p = .045$, $\eta_p^2 = 0.07$), which was caused
by the lack of emotional modulation in nonbiological stimuli. These findings together
showed that the removal of local characteristics greatly disrupted the emotional
modulations on observers' pupil responses, providing conceivable evidence for the
critical contribution of local motion features in emotion perception from the BM signal.

Experiment 4: Local emotional BM

In Experiment 4, we went further to examine whether local BM alone could carry
emotional information and exert a similar modulation effect on pupil size. In particular,
we adopted the well-established local BM stimuli, the scrambled BM, whose local
motion feature was retained while the global configuration information was completely
disrupted. Both the explicit and implicit emotion processing of the scrambled BM were
investigated to provide a thorough view of emotion perception from local BM.
Specifically, a group of observers viewed the scrambled BM and made explicit
behavioral judgments on the emotional information contained in the scrambled BM.
The results showed that observers could successfully recognize the emotions contained
in scrambled BMs with the average accuracy reaching significantly above 50% (M ±
SD = 83% ± 2.8%, $p < .001$). This indicated that scrambled BM conveyed recognizable
emotional information.

Furthermore, we investigated the pupil responses to scrambled happy, sad, and
neutral BMs to explore the automatic and implicit processing of local emotional BM.
The consecutive cluster-based permutation analysis further showed that both the
scrambled happy and sad BM evoked larger pupil sizes than the scrambled neutral BM.
Importantly, such effect appeared in a rather early time window and could last for a

quite long time (happy vs. neutral: 450 ms -3550 ms; sad vs. neutral: 350 ms-4000 ms,

see Fig. 2D). The one-way repeated measures ANOVA on the mean pupil size indicated

a significant main effect of emotional condition (F(2, 46) = 6.47, p = .003, η_p^2 = 0.22,

Fig. 3F). Follow-up analysis showed that the scrambled happy BM induced a

significantly larger pupil response than the scrambled neutral BM (t(23) = 3.22, p = .008,

Cohen's d = 0.66, 95% CI for the mean difference = [0.12, 0.53]; Holm-corrected, p

= .011 after Bonferroni correction, Fig. 3F), which is similar to that observed in

Experiment 1a-b. Notably, the scrambled sad BM also induced a larger pupil size than

the neutral one (t(23) = 3.41, p = .007, Cohen's d = 0.70, 95% CI for the mean difference

= [0.12, 0.53]; Holm-corrected, p = .007 after Bonferroni correction, Fig. 3F).

Moreover, no difference in pupil size is observed between the happy and sad conditions

(t(23) = -0.07, p = .948, Cohen's d = 0.01, 95% CI for the mean difference = [-0.23,

0.24]; Holm-corrected, p = 1.000 after Bonferroni correction, Fig. 3F). The observed

effect could not be accounted for by perceptual differences (e.g., speed), as the happy

and sad scrambled BM differed the most in low-level features, whereas they induced

similar dilation effects on pupil sizes. Again, no significant correlations were observed

between AQ and pupil modulation effects (AQ with happy-sad pupil size: r(23) = -0.21,

p = .32, 95% CI for the mean difference = [-0.57, 0.21]; AQ with happy-neutral pupil

size: r(23) = -0.13, p = .56, 95% CI for the mean difference = [-0.50, 0.29]; AQ with

sad-neutral pupil size: r(23) = 0.13, p = .56, 95% CI for the mean difference = [-0.29,

0.50]). Importantly, a mixed 2 (BM type: intact, local) × 3 (emotional condition: happy,

sad, neutral) ANOVA on average pupil size was conducted to examine whether or not

this emotional modulation effect of local BM varied from that of intact BM reported in

Experiment 1a-b. Results revealed a significant interaction between BM type and

emotional condition (F(2, 92) = 7.76, p < .001, η_p^2 = 0.14), indicating that emotions in

intact and local BM exerted differential modulations on pupil size. This interaction is

primarily driven by the vanishment of pupil modulation between happy and sad local

BM. Overall, these findings showed that the local BM could convey significant

emotional information, which could induce a pupil dilation effect regardless of its exact

type. As compared to the intact BM, the emotional modulation observed in local BM is

different because it occurs faster and is rather coarse.

Discussion

Life motion signals convey salient emotional information that is crucial for human

survival and social interactions (Johansson, 1973; Troje, 2008). Here, we reported that

such emotional clues carried by the point-light BM walker could exert a modulation

effect on pupil responses. Specifically, the happy BM significantly dilated pupils as compared to the neutral BM, and the sad BM evoked smaller pupil responses than the neutral BM, showing distinct emotional modulations on pupil responses that depend on the emotional contents. Moreover, this emotional modulation of pupil responses could not be explained by the low-level differences, as viewing inverted BMs failed to induce such effects. Importantly, when the emotional BM was deprived of the local motion feature through the removal of accelerations, it failed to induce any modulation on pupil responses. Furthermore, the scrambled emotional BM with only the local motion feature retained could still produce a modulation effect on pupil size. Noticeably, this modulation is rapid but rather coarse: viewing the scrambled happy and sad BM would evoke greater pupil size than the scrambled neutral BM in a relatively early time window, while no significant difference was observed between the scrambled happy and sad BM. Taken together, these findings revealed multi-level processing of emotions in life motion signals: the global emotional BM modulated pupil size in a fine-grained manner that discriminates each type of emotion, and the local emotional BM exerted a coarse but rapid modulation on pupil size.

The emotion processing of BM has been investigated in former studies using explicit behavioral detection and recognition paradigm. It has been reported that happy BM is more rapidly identified and detected as compared to sad and neutral BM (Actis-Grosso et al., 2015; Lee & Kim, 2016; Spencer et al., 2016), while other research observed no such superiority (Chouchourelou et al., 2006). Here, the present study adopted the novel and objective pupillometry index to investigate the emotion processing of BM from a physiological aspect. The pupil index, as a direct reflection of individual arousal level and cognitive state, served as a powerful tool for faithfully and automatically reflecting the implicit processing of emotions in BM. Previous studies have reported larger pupil responses towards emotional images as compared to neutral images, with no significant differences observed between the positive and negative conditions (Bradley & Lang, 2015; Bradley et al., 2008; Snowden et al., 2016). However, these studies mostly adopted complex emotional scene images that conveyed rather general emotional information. When it comes to the specific emotion cues (e.g., happy, sad) delivered by our conspecifics through biologically salient signals (e.g., faces, gestures, voices), the results became intermixed. Some studies reported pupil dilatory effects towards fearful, disgusted, angry, sad faces but not happy faces (Burley et al., 2017). On the contrary, other studies observed larger pupil responses for happy as compared to sad, fearful, and surprised faces (Aktar et al., 2018; Burley & Daughters, 2020; Jessen et al., 2016; Prunty et al., 2021). These conflicting results could be due to the low-level confounds

of emotional faces (e.g., eye size) (Carsten et al., 2019; Harrison et al., 2006). Similar to faces, BM also conveyed salient clues concerning the emotional states of our interactive partners. However, they were highly simplified, and deprived of various irrelevant visual confounders (e.g., body shape). Here, we reported that the happy BM induced a stronger pupil response than the neutral and sad BM, lending support to the happy dilation effect observed with faces (Burley & Daughters, 2020; Prunty et al., 2021). Moreover, it helps ameliorate the concern regarding the low-level confounding factors by identifying similar pupil modulations in another type of social signal with distinctive perceptual features.

It has long been documented that pupil size reflects the degree of cognitive effort and attention input (Joshi & Gold, 2019; van der Wel & van Steenbergen, 2018), and indexes the noradrenalin activity in emotion processing structures like amygdala (Dal Monte et al., 2015; Harrison et al., 2006; Liddell et al., 2005). Thus, the happy dilation effect suggests that the happy BM may be more effective in capturing attention and evoking emotional arousal than the neutral and sad BM, which echoes with the happiness superiority reported in the explicit processing of BM (Lee & Kim, 2016; Spencer et al., 2016; Yuan et al., 2023). Instead, the sad constriction effect potentially indicates a disadvantage for sad BM in attracting visual attention and evoking emotional arousal than neutral BM. In line with this, it has been found that infants looked more at the happy BM walker when displayed in pair with the neutral walker, whereas they attended less to the sad walker as compared to the neutral one (Ogren et al., 2019). Besides, it has been revealed by neural studies that, the happy emotion evoked stronger activities in emotionally relevant brain regions including the amygdala, the extrastriate body area (EBA), and the fusiform body area (FBA), while the sad emotion failed to induce such effects (Peelen et al., 2007; Ross et al., 2019). Still, future studies are needed to provide further evidence regarding the happy advantage and the sad disadvantage in attracting visual attention. For example, in addition to pupil size, the microsaccades also provided valuable insights into attention processes (Baumeler et al., 2020; Engbert & Kliegl, 2003; Meyberg et al., 2017), and potentially involved shared neural circuits with pupil responses (Hafed et al., 2009; Hafed & Krauzlis, 2012; C.-A. Wang et al., 2012). Future studies could combine the microsaccade index with pupil size to provide a more thorough understanding of BM emotion processing.

Notably, the happy over sad dilation effect was negatively correlated with autistic traits: individuals with greater autistic tendencies showed decreased sensitivities to emotions in BM. In fact, the perception of biological motion (BM) has long been

considered an important hallmark of social cognition, and abundant studies reported
that individuals with social cognitive deficits (e.g., ASD) were impaired in BM
perception (Blake et al., 2003; Freitag et al., 2008; Klin et al., 2009; Nackaerts et al.,
2012). More recently, it has been pointed out that the extraction of more complex social
information (e.g., emotions, intentions) from BM, as compared to basic BM
recognitions, could be more effective in detecting ASDs (Federici et al., 2020;
Koldewyn et al., 2009; Parron et al., 2008; Todorova et al., 2019). Specifically, a meta-
analysis found that the effect size expanded nearly twice when the task required
emotion recognition as compared to simple perception/detection (Todorova et al., 2019).
However, for the high-functioning ASD individuals, it has been reported that they
showed comparable performance with the control group in explicitly labelling BM
emotions, while their responses were rather delayed (Mazzoni et al., 2021). This
suggests that ASD individuals could adopt compensatory strategies to complete the
explicit BM labelling task, while their automatic responses remained impaired. Such an
observation highlights the importance of using more objective measures that do not rely
on subjective reports to investigate the intrinsic perception of emotions from BM and
its relationship with ASD-related social deficits. The current study thus introduced
pupil size measurement to this field, and we combined it with the passive viewing task
to investigate the more automatic aspect of BM emotion processing. In addition to
diagnostic ASDs, the non-clinical general population also manifested autistic
tendencies that followed normal distribution and demonstrated substantial heritability
(Hoekstra et al., 2007). Here, we focused on the autistic tendencies in the general
population, and our results showed that the automatic emotion processing of BM stimuli
was impaired in individuals with high autistic tendencies, lending support to previous
studies (Hubert et al., 2006; Nackaerts et al., 2012; Parron et al., 2008). The more
detailed test-retest examination further confirmed such a correlation and illustrated a
general diminishment in BM emotion perception ability (happy and sad) for high AQ
individuals. Given that pupil measurement does not require any explicit verbal reports,
it is easily attainable in children and even infants. Thus, the pupil modulation effect
obtained here may serve as a sensitive and reliable physiological marker for detecting
early social cognitive disorders in both clinical and non-clinical populations.

Remarkably, our findings highlighted the central role of local motion feature in
modulating pupil responses towards emotional information contained in BM. Previous
studies have shown that human visual system is highly sensitive to the local BM
stimulus, whose global configural information is completely deprived through spatially
scrambling (Troje & Westhoff, 2006). For example, it has been demonstrated that

scrambled BM could perform as intact BM in lengthening subjective temporal perception (L. Wang & Jiang, 2012). Besides, such local motion feature also served as a basic pre-attentive feature in visual search (L. Wang et al., 2010) and could further induce a significant reflexive attentional effect (L. Wang et al., 2014; Yu et al., 2020). Moreover, the deprivation of the local BM feature would greatly disrupt the perception of the contained biologically salient information, such as animacy and walking direction (Chang & Troje, 2008, 2009; Yu et al., 2020). Here, we extended this line of research to the emotional domain by demonstrating that the local BM component is also critical for the processing of emotional information: when the local BM feature is disrupted, the modulation of emotions on pupil size completely disappears. Furthermore, the emotional BM that retained only local motion features could still exert a salient modulation effect on pupil size. In particular, the happy and sad local BM induced a significant pupil dilation effect as compared to the neutral local BM stimuli. Intriguingly, this dilation effect is independent of the specific emotion category but reflects a general activation caused by the affective salient information. In addition, this non-selective pupil dilation effect in local BM appeared in a relatively early time window, indicating that the extraction of emotions from local BM is rapid. Taken together, these findings identified a coarse but rapid emotion processing mechanism in local BM that could promptly detect the emotional information therein without the aid of the global shape.

Moreover, our findings revealed the existence of distinct emotional modulations in intact and local BM, respectively, as evidenced by variations in pupil responses. In particular, the happy intact BM induced a significant pupil dilation effect as compared to the neutral one, whereas the sad intact BM led to pupil constriction. Interestingly, both the happy and sad local BM resulted in pupil dilation effects relative to the neutral local BM, and such effect occurred at a relatively early time point. This distinctive pupil modulation effects observed in intact and local BM could be elucidated by a multi-level emotion processing mechanism. Specifically, even though both the intact and local BM conveyed important life information, the local BM is deprived of the global configural information (Chang & Troje, 2008, 2009; Simion et al., 2008). Thus, its emotion processing potentially occurred at a more basic and preliminary level, responding to the general affective salient information without engaging in further detailed analysis. Importantly, similar dissociated emotion processing phenomenon has been observed in another important type of emotional signal with analogous function (i.e., facial expression). For example, happy and fearful faces induced differential amygdala activations when perceived consciously, yet they induced comparable amygdala

activations when suppressed (Williams et al., 2004). Moreover, it has been argued that there exist two parallel pathways for facial expression processing: a slow route that conveys fine-grained facial features along cortical areas to the amygdala, and a rapid subcortical pathway that directly transfers emotional information to amygdala without detailed analysis, which is known as the "quick and dirty" route (Garrido et al., 2012; Johnson, 2005; Méndez-Bértolo et al., 2016; Vuilleumier et al., 2003). It is probable that the emotion processing of the local and intact point-light BM potentially functions in a manner similar to that of the faces, with the former serving as a primary detection mechanism that automatically captures emotionally significant information conveyed by animate agents without detailed analysis, and the latter aiding in more specific emotion identification based on fine-grained analyses of their motion pattern and global shape. Compatible with this view, recent studies have reported that the emotion processing of face and BM was very similar. For example, they both showed a happiness superiority in visual detection (Becker et al., 2011; Lee & Kim, 2016; Nackaerts et al., 2012) and guiding social attention (Yuan et al., 2023). Moreover, their emotion processing involved potentially overlapping brain regions (e.g., amygdala, STS) (Alaerts et al., 2014; Engell & Haxby, 2007; Peelen et al., 2007; Pessoa & Adolphs, 2010). Overall, our findings, together with the former evidence, suggest dissociated emotion processing for BM in the human brain akin to the dual-route model reported in faces, and further imply a general "emotional brain" that can be shared by different types of social signals. Still, future studies are needed to implement neuroimaging techniques to directly identify the brain regions involved in the emotion processing of local and global BM signals.

To conclude, the current study clearly demonstrated that the emotional information conveyed by point-light BM modulated pupil responses. The intact BM exerted a fine-grained emotional modulation on pupil sizes, while disrupting the contained local motion characteristic would deteriorate the observed modulations. Moreover, BM with only local motion features retained could exert a fast but rather coarse modulation on pupillometry. These findings together highlight the critical role of local motion feature in BM emotion processing, and further reveal the multi-level emotion processing of BM. Notably, the observed emotional modulation in intact BM is associated with individual autistic traits, suggesting the potential of utilizing the emotion-modulated pupil responses to facilitate the diagnosis of social cognitive disorders.

Methods

Participants

A total of 144 participants (52 males, 92 females) ranging from 18 to 30 years old (M

$\pm$ SD = 23.0 $\pm$ 2.6) were recruited, with 24 in Experiment 1a, 24 in Experiment 1b, 24

in Experiment 2, 24 in Experiment 3, 24 in the behavioral part of Experiment 4, and 24

in the eye recoding part of Experiment 4. All of the participants had normal or

corrected-to-normal vision and gave written informed consent in accordance with the

procedure and protocols approved by the institutional review board of the Institute of

Psychology, Chinese Academy of Sciences. They were all naïve to the purpose of the

experiments. Prior power analyses (*F* tests, repeated measures, within factors) using

G*Power (Version 3.1.9.4; (Faul et al., 2007)) indicated that a sample size of 24

participants would afford 80% power with alpha at .05 to detect a moderate main effect

(*f* = 0.27). This sample size was comparable to previous studies with similar designs

(Nakakoga et al., 2020).

Stimuli

Stimuli were displayed using MATLAB (Mathworks, Inc.) together with the

Psychtoolbox extensions (Brainard, 1997; Pelli, 1997) on a 23-inch LED monitor (1920

$\times$ 1080 at 60 Hz) with gray background (RGB: 100, 100, 100). The biological motion

(BM) stimuli were adopted from Troje (Troje, 2008). Each comprised 15 point-light

dots depicting the motions of the head, pelvis, thorax and major joints (i.e., shoulders,

elbows, wrists, hips, knees, and ankles). Its emotional state was indexed by a

normalized Z score on an axis that reflects the differences between happy and sad

walkers in terms of a linear classifier. The scores were computed within a 10-

dimensional sub-space spanned by the first 10 principal components based on a Fourier-

based representation of observers' emotional ratings of 80 actual walkers (half male)

(Troje, 2008). We adopted the neutral walker that scored 0 on the linear axis, together

with the happy walker 6 SDs into the happy part of the axis and the sad walker 6 SDs

into the sad part of the axis (see https://www.biomotionlab.ca/html5-bml-walker/ for

an interactive animation). Besides, we turned the BM walkers 45 degrees leftwards or

rightwards to maximize the visibility of expressive bodily cues (Roether et al., 2009).

In Experiment 1a-1b, the upright emotional (happy, sad, or neutral) BM walkers with

two walking directions (45° leftwards or rightwards) were used. The velocity was 5.76

pixels/frame for the happy BM, 4.14 pixels/frame for the neutral BM, and 3.21

pixels/frame for the sad BM. This difference in velocity profile was considered an

important signature for conveying emotional information, as the happy walker was

characterized by a larger step pace and longer arm swing, and the sad walker would

instead exhibit a slouching gait with short slow strides and smaller arm movement

(Barliya et al., 2012; Chouchourelou et al., 2006; Halovic & Kroos, 2018; Roether et

al., 2009). In Experiment 2, the stimuli were replaced with their inverted counterparts

created by mirror flipping the upright BM vertically. In Experiment 3, we presented

observers with the non-biological motion stimuli whose local BM characteristic was

disrupted through acceleration removal. More specifically, each individual dot moved

along the original path with a constant speed equal to the average speed of the dot

(Chang & Troje, 2009). Such manipulation disrupted the local motion feature of the

BM stimuli but kept the trajectories of individual point lights unchanged. In Experiment

4, observers viewed the scrambled BM stimuli, which were created by randomly

relocating the point light dots within the region occupied by the original BM walker. In

this manner, the local motion feature was preserved while the global configuration of

the BM stimulus was entirely disrupted (Troje & Westhoff, 2006). Please refer to

Supplementary Materials for the demonstrations displaying the motion stimuli used in

Experiments 1-4.

Procedure

Participants were seated at a viewing distance of 60 cm to the computer screen with

their heads on a chinrest to minimize their head movements. Their pupil size and eye

position of the left eye were recorded using a video-based iView X Hi-Speed system

(SMI, Berlin, Germany) at 500 Hz. In Experiment 1a, each trial began with a central

fixation cross ($0.2° \times 0.2°$) with variable duration (800-1200 ms), followed by an

upright happy/sad/neutral point-light walker (half leftwards and half rightwards)

presented centrally for 4000 ms. Participants were required to passively view the BM

stimulus and continue the procedure by pressing the space bar. After that, a blank screen

was displayed for 3000 ms (Fig. 1). There were 4 blocks of 30 trials, and participants

were given a short break after every block. Besides, we also administered a 9-point

calibration of the eye-tracker followed by a validation stage before each experimental

block to ensure the data quality. Participants were told to maintain fixation on the center

of the screen and not to blink during the stimuli presentation. Importantly, a replication

experiment of Experiment 1a (Experiment 1b) was conducted, which followed the same

protocol, with the change being that participants were asked to return to the lab for a

retest after at least seven days. Experiment 2 is identical to Experiment 1a, except that

the BM stimuli were changed to their inverted counterparts. In Experiment 3, we instead

presented participants with the non-biological motion displays whose local motion

feature was deprived through acceleration removal. In Experiment 4, we adopted the

scrambled emotional BM stimuli, and investigated both the explicit and the implicit emotion processing of scrambled BM. Note that a group of participants was recruited to perform emotional judgments on the scrambled BM in order to investigate whether the emotional information carried by local BM signals could be explicitly identified. Another group of participants was enrolled for the pupil recording experiment, which followed the identical experimental procedure of Experiments 1-3. At the end of all experiments, participants were required to complete the 50-point Autism-spectrum Questionnaire (AQ), which measured the degree of autistic traits in the normal population (Baron-Cohen et al., 2001).

Data Analysis

The raw pupil data for each trial was first screened to remove eye blinks (either replaced by linear interpolation or with this trial discarded). Then, trials with pupil size deviating ± 3 SDs from the mean were excluded from further analysis. Finally, the pupil size data were down-sampled to 20 Hz and baseline-corrected for each trial by subtracting the mean pupil size during the 200 ms pre-stimulus period. We computed the average pupil size for each emotional condition (happy, sad, or neutral) obtained by collapsing the pupillometry across all time points. Besides, to depict the time course of pupil responses towards emotional BM, we further conducted a consecutive paired-sample *t*-test across all time points comparing different emotional conditions. The cluster-based permutation analysis was applied to avoid potential problems brought about by multiple comparisons (Einhauser et al., 2008). In this analysis, the computed *t*-values neighboring in time that exceeded a threshold ($p < 0.05$) were defined as clusters, and then summed to produce a cluster mass. The cluster mass was compared with a null distribution, which was generated by 2000 random permutations of the pupil data from different conditions. If the cluster mass fell beyond 95% of the null distribution (α = 0.05), it was deemed to be statistically significantly different (Einhauser et al., 2008). The pupil size was analyzed and reported in arbitrary units (a.u.) without transforming into the actual unit (mm), as the relative change of the pupil size was of main interest.

Acknowledgements

This research was supported by grants from the Ministry of Science and Technology of China (STI2030-Major Projects 2021ZD0203800 and 2022ZD0205100), the National Natural Science Foundation of China (No. 31830037, 32371106), the Strategic Priority Research Program (No. XDB32010300), the Interdisciplinary Innovation Team (JCTD-2021-06), the Science Foundation of Institute of Psychology, Chinese Academy of

Sciences, and the Fundamental Research Funds for the Central Universities. We thank Professor Nikolaus F. Troje for kindly providing us with the visual stimuli. We report how we determined our sample size, all data exclusions (if any), all manipulations, and all measures in the study. This study's design and its analysis were not pre-registered. All data, materials, and analysis code used in the current study could be accessed at Science Data Bank (https://doi.org/10.57760/sciencedb.psych.00125). The authors declared no conflicts of interest with respect to the authorship or the publication of this article.

References:

Actis-Grosso, R., Bossi, F., & Ricciardelli, P. (2015). Emotion recognition through static faces and moving bodies: a comparison between typically developed adults and individuals with high level of autistic traits. *Frontiers in Psychology*, *6*, 1570. https://doi.org/10.3389/fpsyg.2015.01570

Aktar, E., Mandell, D. J., de Vente, W., Majdandžić, M., Oort, F. J., van Renswoude, D. R., Raijmakers, M. E. J., & Bögels, S. M. (2018). Parental negative emotions are related to behavioral and pupillary correlates of infants' attention to facial expressions of emotion. *Infant Behavior and Development, 53*, 101–111. https://doi.org/10.1016/j.infbeh.2018.07.004

Alaerts, K., Nackaerts, E., Meyns, P., Swinnen, S. P., & Wenderoth, N. (2011). Action and emotion recognition from point light displays: an investigation of gender differences. *PLoS ONE*, *6*(6), e20989.

Alaerts, K., Woolley, D. G., Steyaert, J., Di Martino, A., Swinnen, S. P., & Wenderoth, N. (2014). Underconnectivity of the superior temporal sulcus predicts emotion recognition deficits in autism. *Social Cognitive and Affective Neuroscience*, *9*(10), 1589–1600. https://doi.org/10.1093/scan/nst156

Barliya, A., Omlor, L., Giese, M. A., Berthoz, A., & Flash, T. (2012). Expression of emotion in the kinematics of locomotion. *Experimental Brain Research, 225*(2), 159 – 176. https://doi.org/10.1007/s00221-012-3357-4

Baron-Cohen, S., Wheelwright, S., Skinner, R., Martin, J., & Clubley, E. (2001). The autism-spectrum quotient (aq): evidence from asperger syndrome/high-functioning autism, malesand females, scientists and mathematicians. *Journal of Autism and Developmental Disorders*. https://doi.org/10.1023/a:1005653411471

Baumeler, D., Schönhammer, J. G., & Born, S. (2020). Microsaccade dynamics in the attentional repulsion effect. *Vision Research, 170*, 46–52. https://doi.org/10.1016/j.visres.2020.03.009

Becker, D. V., Anderson, U. S., Mortensen, C. R., Neufeld, S. L., & Neel, R. (2011). The face in the crowd effect unconfounded: happy faces, not angry faces, are more efficiently detected in

single- and multiple-target visual search tasks. *Journal of Experimental Psychology: General*, *140*(4), 637–659. https://doi.org/10.1037/a0024060

Blake, R., Turner, L. M., Smoski, M. J., Pozdol, S. L., & Stone, W. L. (2003). Visual recognition of biological motion is impaired in children with autism. *Psychological Science, 14*(2), 151–157. https://doi.org/10.1111/1467-9280.01434

Bradley, M. M., & Lang, P. J. (2015). Memory, emotion, and pupil diameter: repetition of natural scenes. *Psychophysiology, 52*(9), 1186–1193. https://doi.org/10.1111/psyp.12442

Bradley, M. M., Miccoli, L., Escrig, M. A., & Lang, P. J. (2008). The pupil as a measure of emotional arousal and autonomic activation. *Psychophysiology, 45*(4), 602–607. https://doi.org/10.1111/j.1469-8986.2008.00654.x

Brainard, D. (1997). The psychophysics toolbox. *Spatial Vision*, *10*(4), 433–436.

Burley, D. T., & Daughters, K. (2020). The effect of oxytocin on pupil response to naturalistic dynamic facial expressions. *Hormones and Behavior*, *125*, 104837. https://doi.org/10.10 16/j.yhbeh.2020.104837

Burley, D. T., Gray, N. S., & Snowden, R. J. (2017). As far as the eye can see: relationship between psychopathic traits and pupil response to affective stimuli. *PLOS ONE*, *12*(1), e0167436. https://doi.org/10.1371/journal.pone.0167436

Carsten, T., Desmet, C., Krebs, R. M., & Brass, M. (2019). Pupillary contagion is independent of the emotional expression of the face. *Emotion, 19*(8), 1343 – 1352. https://doi.org/10.1037/emo0000503

Chang, D. H. F., & Troje, N. F. (2008). Perception of animacy and direction from local biological motion signals. *Journal of Vision*, *8*(5), 3. https://doi.org/10.1167/8.5.3

Chang, D. H. F., & Troje, N. F. (2009). Acceleration carries the local inversion effect in biological motion perception. *Journal of Vision*, *9*(1), 19–19. https://doi.org/10.1167/9.1.19

Chouchourelou, A., Matsuka, T., Harber, K., & Shiffrar, M. (2006). The visual analysis of emotional actions. *Social Neuroscience*, *1*(1), 63–74. https://doi.org/10.1080/17470910600630599

Critchley, H., Daly, E., Phillips, M., Brammer, M., Bullmore, E., Williams, S., Van Amelsvoort, T., Robertson, D., David, A., & Murphy, D. (2000). Explicit and implicit neural mechanisms for processing of social information from facial expressions: a functional magnetic resonance imaging study. *Human Brain Mapping*, *9*(2), 93–105. https://doi.org/10.1002/(sici)1097-0193(200002)9:2<93::aid-hbm4>3.0.co;2-z

Dal Monte, O., Costa, V. D., Noble, P. L., Murray, E. A., & Averbeck, B. B. (2015). Amygdala lesions in rhesus macaques decrease attention to threat. *Nature Communications, 6*(1). https://doi.org/10.1038/ncomms10161

de Gee, J. W., Knapen, T., & Donner, T. H. (2014). Decision-related pupil dilation reflects upcoming choice and individual bias. *Proceedings of the National Academy of Sciences*, *111*(5), E618–E625. https://doi.org/10.1073/pnas.1317557111

de Gelder, B. (2006). Towards the neurobiology of emotional body language. *Nature Reviews Neuroscience*, *7*(3), 242–249. https://doi.org/10.1038/nrn1872

Einhauser, W., Stout, J., Koch, C., & Carter, O. (2008). Pupil dilation reflects perceptual selection and predicts subsequent stability in perceptual rivalry. *Proceedings of the National Academy of Sciences*, *105*(5), 1704–1709. https://doi.org/10.1073/pnas.0707727105

Engbert, R., & Kliegl, R. (2003). Microsaccades uncover the orientation of covert attention. *Vision Research, 43*(9), 1035–1045. https://doi.org/10.1016/s0042-6989(03)00084-1

Engell, A. D., & Haxby, J. V. (2007). Facial expression and gaze-direction in human superior temporal sulcus. *Neuropsychologia*, *45*(14), 3234–3241. https://doi.org/10.1016/j.neuropsychologia.2007.06.022

Faul, F., Erdfelder, E., Lang, A.-G., & Buchner, A. (2007). G*Power 3: a flexible statistical power analysis program for the social, behavioral, and biomedical sciences. *Behavior Research Methods*, *39*(2), 175–191. https://doi.org/10.3758/bf03193146

Federici, A., Parma, V., Vicovaro, M., Radassao, L., Casartelli, L., & Ronconi, L. (2020). Anomalous perception of biological motion in autism: a conceptual review and meta-analysis. *Scientific Reports, 10*(1). https://doi.org/10.1038/s41598-020-61252-3

Freitag, C. M., Konrad, C., Häberlen, M., Kleser, C., von Gontard, A., Reith, W., Troje, N. F., & Krick, C. (2008). Perception of biological motion in autism spectrum disorders. *Neuropsychologia, 46*(5), 1480 – 1494. https://doi.org/10.1016/j.neuropsychologia.2007.12.025

Frith, C. (2009). Role of facial expressions in social interactions. *Philosophical Transactions of the Royal Society B: Biological Sciences*, *364*(1535), 3453–3458. https://doi.org/10.1098/rstb.2009.0142

Frith, U. (2004). Emanuel miller lecture: confusions and controversies about asperger syndrome. *Journal of Child Psychology and Psychiatry*, *45*(4), 672–686. https://doi.org/10.1111/j.1469-7610.2004.00262.x

Garrido, M. I., Barnes, G. R., Sahani, M., & Dolan, R. J. (2012). Functional evidence for a dual route to amygdala. *Current Biology*, *22*(2), 129–134. https://doi.org/10.1016/j.cub.2011.11.056

Graves, J. E., Egré, P., Pressnitzer, D., & de Gardelle, V. (2021). An implicit representation of stimulus ambiguity in pupil size. *Proceedings of the National Academy of Sciences*, *118*(48), e2107997118.https://doi.org/10.1073/pnas.2107997118

Hafed, Z. M., Goffart, L., & Krauzlis, R. J. (2009). A neural mechanism for microsaccade generation in the primate superior colliculus. *Science, 323*(5916), 940 – 943. https://doi.org/10.1126/science.1166112

Hafed, Z. M., & Krauzlis, R. J. (2012). Similarity of superior colliculus involvement in microsaccade and saccade generation. *Journal of neurophysiology, 107*(7), 1904-1916.

Halovic, S., & Kroos, C. (2018). Not all is noticed: kinematic cues of emotion-specific gait. *Human Movement Science, 57*, 478–488. https://doi.org/10.1016/j.humov.2017.11.008

Harms, M. B., Martin, A., & Wallace, G. L. (2010). Facial emotion recognition in autism spectrum disorders: a review of behavioral and neuroimaging studies. *Neuropsychology Review*, *20*(3), 290–322. https://doi.org/10.1007/s11065-010-9138-6

Harrison, N. A., Singer, T., Rotshtein, P., Dolan, R. J., & Critchley, H. D. (2006). Pupillary contagion: central mechanisms engaged in sadness processing. *Social Cognitive and Affective Neuroscience, 1*(1), 5–17. https://doi.org/10.1093/scan/nsl006

Hoekstra, R. A., Bartels, M., Verweij, C. J. H., & Boomsma, D. I. (2007). Heritability of autistic traits in the general population. *Archives of Pediatrics & Adolescent Medicine, 161*(4), 372. https://doi.org/10.1001/archpedi.161.4.372

Hubert, B., Wicker, B., Moore, D. G., Monfardini, E., Duverger, H., Fonséca, D. D., & Deruelle, C. (2006). Brief report: recognition of emotional and non-emotional biological motion in individuals with autistic spectrum disorders. *Journal of Autism and Developmental Disorders*, *37*(7), 1386–1392. https://doi.org/10.1007/s10803-006-0275-y

Jessen, S., Altvater-Mackensen, N., & Grossmann, T. (2016). Pupillary responses reveal infants' discrimination of facial emotions independent of conscious perception. *Cognition, 150*, 163–169. https://doi.org/10.1016/j.cognition.2016.02.010

Johansson, G. (1973). Visual perception of biological motion and a model for its analysis. *Perception & Psychophysics*, *14*(2), 201–211. https://doi.org/10.3758/bf03212378

Johnson, M. H. (2005). Subcortical face processing. *Nature Reviews Neuroscience*, *6*(10), 766–774. https://doi.org/10.1038/nrn1766

Joshi, S., & Gold, J. I. (2019). Pupil size as a window on neural substrates of cognition. *Trends in Cognitive Sciences*, *24*(6), 466–480. https://doi.org/10.31234/osf.io/dvsme

Kana, R. K., Patriquin, M. A., Black, B. S., Channell, M. M., & Wicker, B. (2015). Altered medial frontal and superior temporal response to implicit processing of emotions in autism. *Autism Research*, *9*(1), 55–66. https://doi.org/10.1002/aur.1496

Keifer, C. M., Mikami, A. Y., Morris, J. P., Libsack, E. J., & Lerner, M. D. (2020). Prediction of social behavior in autism spectrum disorders: explicit versus implicit social cognition. *Autism*, *24*(7), 1758–1772. https://doi.org/10.1177/1362361320922058

Klin, A., Lin, D. J., Gorrindo, P., Ramsay, G., & Jones, W. (2009). Two-year-olds with autism orient to non-social contingencies rather than biological motion. *Nature, 459*(7244), 257–261. https://doi.org/10.1038/nature07868

Kloosterman, N. A., Meindertsma, T., van Loon, A. M., Lamme, V. A. F., Bonneh, Y. S., & Donner, T. H. (2015). Pupil size tracks perceptual content and surprise. *European Journal of Neuroscience*, *41*(8), 1068–1078. https://doi.org/10.1111/ejn.12859

Koldewyn, K., Whitney, D., & Rivera, S. M. (2009). The psychophysics of visual motion and global
form processing in autism. *Brain, 133*(2), 599–610. https://doi.org/10.1093/brain/awp272

Kovarski, K., Mennella, R., Wong, S. M., Dunkley, B. T., Taylor, M. J., & Batty, M. (2018).
Enhanced early visual responses during implicit emotional faces processing in autism spectrum
disorder. *Journal of Autism and Developmental Disorders*, *49*(3), 871–886.
https://doi.org/10.1007/s10803-018-3787-3

Lange, K., Williams, L. M., Young, A. W., Bullmore, E. T., Brammer, M. J., Williams, S. C. R.,
Gray, J. A., & Phillips, M. L. (2003). Task instructions modulate neural responses to fearful
facial expressions. *Biological Psychiatry*, *53*(3), 226–232. https://doi.org/10.1016/s0006-
3223(02)01455-5

Lee, H., & Kim, J. (2016). Facilitating effects of emotion on the perception of biological motion:
evidence for a happiness superiority effect. *Perception*, *46*(6), 679–697.
https://doi.org/10.1177/0301006616681809

Liddell, B. J., Brown, K. J., Kemp, A. H., Barton, M. J., Das, P., Peduto, A., Gordon, E., & Williams,
L. M. (2005). A direct brainstem–amygdala–cortical 'alarm' system for subliminal signals of
fear. *NeuroImage, 24*(1), 235–243.

Mazzoni, N., Ricciardelli, P., Actis-Grosso, R., & Venuti, P. (2021). Difficulties in recognising
dynamic but not static emotional body movements in autism spectrum disorder. *Journal of
Autism and Developmental Disorders, 52*(3), 1092–1105. https://doi.org/10.1007/s10803-021-
05015-7

Méndez-Bértolo, C., Moratti, S., Toledano, R., Lopez-Sosa, F., Martínez-Alvarez, R., Mah, Y. H.,
Vuilleumier, P., Gil-Nagel, A., & Strange, B. A. (2016). A fast pathway for fear in human
amygdala. *Nature Neuroscience*, *19*(8), 1041–1049. https://doi.org/10.1038/nn.4324

Meyberg, S., Sinn, P., Engbert, R., & Sommer, W. (2017). Revising the link between microsaccades
and the spatial cueing of voluntary attention. *Vision Research, 133*, 47 – 60.
https://doi.org/10.1016/j.visres.2017.01.001

Nackaerts, E., Wagemans, J., Helsen, W., Swinnen, S. P., Wenderoth, N., & Alaerts, K. (2012).
Recognizing biological motion and emotions from point-light displays in autism spectrum
disorders. *PLoS ONE*, *7*(9), e44473. https://doi.org/10.1371/journal.pone.0044473

Nakakoga, S., Higashi, H., Muramatsu, J., Nakauchi, S., & Minami, T. (2020). Asymmetrical
characteristics of emotional responses to pictures and sounds: evidence from pupillometry.
PLOS ONE, *15*(4), e0230775. https://doi.org/10.1371/journal.pone.0230775

Ogren, M., Kaplan, B., Peng, Y., Johnson, K. L., & Johnson, S. P. (2019). Motion or emotion:
infants discriminate emotional biological motion based on low-level visual information. *Infant
Behavior and Development*, *57*, 101324. https://doi.org/10.1016/j.infbeh.2019.04.006

Okon-Singer, H., Lichtenstein-Vidne, L., & Cohen, N. (2013). Dynamic modulation of emotional
processing. *Biological Psychology*, *92*(3), 480–491. https://doi.org/10.1016/j.biopsy
cho.2012.05.010

Oliva, M., & Anikin, A. (2018). Pupil dilation reflects the time course of emotion recognition in
human vocalizations. *Scientific Reports*, *8*(1), 4871. https://doi.org/10.1038/s41598-018-
23265-x

Parron, C., Da Fonseca, D., Santos, A., Moore, D. G., Monfardini, E., & Deruelle, C. (2008).
Recognition of biological motion in children with autistic spectrum disorders. *Autism, 12*(3),
261–274. https://doi.org/10.1177/1362361307089520

Peelen, M. V., Atkinson, A. P., Andersson, F., & Vuilleumier, P. (2007). Emotional modulation of
body-selective visual areas. *Social Cognitive and Affective Neuroscience*, *2*(4), 274–283.
https://doi.org/10.1093/scan/nsm023

Pelli, D. (1997). The videotoolbox software for visual psychophysics: transforming numbers into
movies. *Spatial Vision*.

Pessoa, L., & Adolphs, R. (2010). Emotion processing and the amygdala: from a "low road" to
"many roads" of evaluating biological significance. *Nature Reviews Neuroscience*, *11*(11),
773–782. https://doi.org/10.1038/nrn2920

Prunty, J. E., Keemink, J. R., & Kelly, D. J. (2021). Infants show pupil dilatory responses to happy
and angry facial expressions. *Developmental Science*, *25*(2). https://doi.org/10.11
11/desc.13182

Roether, C. L., Omlor, L., Christensen, A., & Giese, M. A. (2009). Critical features for the
perception of emotion from gait. *Journal of Vision*, *9*(6), 15–15. https://doi.org/10.1167/9.6.15

Ross, P., de Gelder, B., Crabbe, F., & Grosbras, M. H. (2019). Emotion modulation of the body-
selective areas in the developing brain. *Developmental Cognitive Neuroscience, 38*, 100660.

Shafer, A. T., Matveychuk, D., Penney, T., O'Hare, A. J., Stokes, J., & Dolcos, F. (2012). Processing
of emotional distraction is both automatic and modulated by attention: evidence from an event-
related fmri investigation. *Journal of Cognitive Neuroscience*, *24*(5), 1233–1252.
https://doi.org/10.1162/jocn_a_00206

Simion, F., Regolin, L., & Bulf, H. (2008). A predisposition for biological motion in the newborn
baby. *Proceedings of the National Academy of Sciences, 105*(2), 809 – 813.
https://doi.org/10.1073/pnas.0707021105

Snowden, R. J., O'Farrell, K. R., Burley, D., Erichsen, J. T., Newton, N. V., & Gray, N. S. (2016).
The pupil's response to affective pictures: role of image duration, habituation, and viewing
mode. *Psychophysiology, 53*(8), 1217–1223. https://doi.org/10.1111/psyp.12668

Spencer, J. M. Y., Sekuler, A. B., Bennett, P. J., Giese, M. A., & Pilz, K. S. (2016). Effects of aging
on identifying emotions conveyed by point-light walkers. *Psychology and Aging*, *31*(1), 126–
138. https://doi.org/10.1037/a0040009

Todorova, G. K., Hatton, R. E. M., & Pollick, F. E. (2019). Biological motion perception in autism
spectrum disorder: a meta-analysis. *Molecular Autism, 10*(1). https://doi.org/10.1186/s13229-
019-0299-8

Troje, N. F. (2008). *Retrieving Information from Human Movement Patterns* (pp. 308–334). Oxford
University PressNew York. https://doi.org/10.1093/acprof:oso/9780195188370.003.0014

Troje, N. F., & Westhoff, C. (2006). The inversion effect in biological motion perception: evidence
for a "life detector"? *Current Biology*, *16*(8), 821–824. https://doi.org/10.10
16/j.cub.2006.03.022

van der Wel, P., & van Steenbergen, H. (2018). Pupil dilation as an index of effort in cognitive
control tasks: a review. *Psychonomic Bulletin & Review, 25*(6), 2005 – 2015.
https://doi.org/10.3758/s13423-018-1432-y

Vuilleumier, P., Armony, J. L., Driver, J., & Dolan, R. J. (2003). Distinct spatial frequency
sensitivities for processing faces and emotional expressions. *Nature Neuroscience*, *6*(6), 624–
631. https://doi.org/10.1038/nn1057

Wang, C.-A., Boehnke, S. E., White, B. J., & Munoz, D. P. (2012). Microstimulation of the monkey
superior colliculus induces pupil dilation without evoking saccades. *Journal of Neuroscience,*
32(11), 3629–3636. https://doi.org/10.1523/jneurosci.5512-11.2012

Wang, L., & Jiang, Y. (2012). Life motion signals lengthen perceived temporal duration.
Proceedings of the National Academy of Sciences, *109*(11), E673–E677. https://doi.org/10.10
73/pnas.1115515109

Wang, L., Yang, X., Shi, J., & Jiang, Y. (2014). The feet have it: local biological motion cues trigger
reflexive attentional orienting in the brain. *NeuroImage*, *84*, 217–224.
https://doi.org/10.1016/j.neuroimage.2013.08.041

Wang, L., Zhang, K., He, S., & Jiang, Y. (2010). Searching for life motion signals. *Psychological*
Science, *21*(8), 1083–1089. https://doi.org/10.1177/0956797610376072

Williams, M. A., Morris, A. P., McGlone, F., Abbott, D. F., & Mattingley, J. B. (2004). Amygdala
responses to fearful and happy facial expressions under conditions of binocular suppression.
Journal of Neuroscience, 24(12), 2898-2904.

Wong, T. K. W., Fung, P. C. W., Chua, S. E., & McAlonan, G. M. (2008). Abnormal spatiotemporal
processing of emotional facial expressions in childhood autism: dipole source analysis of
event-related potentials. *European Journal of Neuroscience*, *28*(2), 407–416.
https://doi.org/10.1111/j.1460-9568.2008.06328.x

Yu, Y., Ji, H., Wang, L., & Jiang, Y. (2020). Cross-modal social attention triggered by biological
motion cues. *Journal of Vision*, *20*(10), 21. https://doi.org/10.1167/jov.20.10.21

Yuan, T., Ji, H., Wang, L., & Jiang, Y. (2023). Happy is stronger than sad: Emotional information
modulates social attention. *Emotion*, *23*(4), 1061–1074. https://doi.org/10.1037/emo0001145

Yuan, T., Wang, L., & Jiang, Y. (2024). Cross-channel adaptation reveals shared emotion

representation from face and biological motion. *Emotion*, in press.

Figures:

Fig. 1. Schematic representation of the experimental procedure and the stimuli in Experiment 1a. A
happy/sad/neutral BM walker turning 45 degrees leftwards or rightwards was presented at the center of the screen
for 4000 ms. Participants were instructed to maintain their fixation on the BM stimuli during stimulus presentation
and to continue the procedure through key pressing.
820

Fig. 2. Time course of pupil responses to happy, sad, and neutral BM in Experiments 1-4. Solid lines represent pupil diameter under each emotional condition as a function of time (happy: red; sad: blue; neutral: gray); shaded areas represent the SEM between participants; colored horizontal lines indicate periods during which there are statistically significant differences among conditions at $p < 0.05$; and black horizontal lines indicate significant differences after cluster-based permutation correction. All the pupil data are in arbitrary units (a.u.) (A) In Experiment 1a, the happy BM evoked larger pupil response as compared to the sad and neutral BM, and the sad BM evoked smaller pupil size than the neutral BM. (B) In Experiment 2, the inverted BM failed to produce such emotional modulation effects. (C) In Experiment 3, the emotional BM that is deprived of the local motion feature exerted no emotional modulation on pupil responses. (D) In Experiment 4, both the happy and sad local BM induced a larger pupil response than the neutral local BM, and such dilation effect started from a relatively early time point.

Fig. 3. Normalized mean pupil responses using the neutral condition as baseline, plotted against happy and sad conditions, and the relevant correlation. (A) In Experiment 1a, the group average pupil response to happy intact BM is significantly larger than that to sad and neutral BM, while the pupil size induced by sad BM is significantly smaller than that evoked by neutral BM. (B-C) Moreover, such an emotional modulation on pupil sizes was again identified in the test and retest of the replication experiment (Experiment 1b). (D) In Experiment 2, no significant differences in pupil responses were observed for inverted BM. (E) In Experiment 3, when the biological characteristic was deprived from the emotional BM, it failed to induce any modulations on pupil sizes. (F) In Experiment 4, both the happy and sad local BM induced a significantly larger pupil size than neutral local BM, with no significant difference between the happy and sad condition. All the pupil data are in arbitrary units (a.u.). Each point represents one individual data. Error bars showed standard errors of the mean. $* p < .05$, $** p < .01$.

Fig. 4. Correlation results for pupil modulations and AQ scores in Experiment 1a and its replication experiment (Experiment 1b). (A) In Experiment 1a, a significant negative correlation was found between the happy over sad pupil dilation effect and individual autistic quotient (AQ). (B-C) No other significant correlations were found. (D-F) The first test of Experiment 1b replicated the negative correlation between happy over sad pupil dilation effect and AQ. Similarly, no other significant correlations were found. (G) In the second test, the negative correlation between happy over sad pupil dilation effect and AQ was similarly observed and even stronger. (H-I) Moreover, we've additionally observed a significant positive correlation between AQ and the happy minus neutral pupil dilation effect, and a significant negative correlation between AQ and the sad minus neutral pupil constriction effect.