Adaptation of the *Mycobacterium tuberculosis* transcriptome to biofilm growth

Madison A. Youngblom [1,2], Tracy M. Smith [2*], Caitlin S. Pepperell [2,3#]

[1] Microbiology Doctoral Training Program, University of Wisconsin-Madison, Wisconsin, USA

[2] Department of Medical Microbiology and Immunology, School of Medicine and Public Health, University of

Madison-Wisconsin, Wisconsin, USA

[3] Department of Medicine (Infectious Diseases), School of Medicine and Public Health, University of Wisconsin-

Madison, Wisconsin, USA

Running Title: Evolving the *M. tb* biofilm transcriptome

*Present address: Tracy M. Smith, New York Genome Center, New York, USA

#Address correspondence to cspepper@medicine.wisc.edu

Abstract

Mycobacterium tuberculosis (*M. tb*), the causative agent of tuberculosis (TB), is a leading global cause

of death from infectious disease. Biofilms are increasingly recognized as a relevant growth form during

M. tb infection and may impede treatment by enabling bacterial drug and immune tolerance. *M. tb* has a

complicated regulatory network that has been well-characterized for many relevant disease states,

including dormancy and hypoxia. However, despite its importance, our knowledge of the genes and

pathways involved in biofilm formation is limited. Here we characterize the biofilm transcriptomes of

fully virulent clinical isolates and find that the regulatory systems underlying biofilm growth vary widely

between strains and are also distinct from regulatory programs associated with other environmental

cues. We used experimental evolution to investigate changes to the transcriptome during adaptation to

biofilm growth and found that the application of a uniform selection pressure resulted in loss of strain-to-

strain variation in gene expression, resulting in a more uniform biofilm transcriptome. The adaptive

trajectories of transcriptomes were shaped by the genetic background of the *M. tb* population leading to

convergence on a sub-lineage specific transcriptome. We identified widespread upregulation of non-

coding RNA (ncRNA) as a common feature of the biofilm transcriptome and hypothesize that ncRNA

function in genome-wide modulation of gene expression, thereby facilitating rapid regulatory responses

to new environments. These results reveal a new facet of the *M. tb* regulatory system and provide

valuable insight into how *M. tb* adapts to new environments.

Importance

Understanding mechanisms of resistance and tolerance in *Mycobacterium tuberculosis* (*M. tb*) can help us develop new treatments that capitalize on *M. tb*'s vulnerabilities. Here we used transcriptomics to study both the regulation of biofilm formation in clinical isolates as well as how those regulatory systems adapt to new environments. We find that closely related clinical populations have diverse strategies for growth under biofilm conditions, and that genetic background plays a large role in determining the trajectory of evolution. These results have implications for future treatment strategies that may be informed by our knowledge of the evolutionary constraints on strain(s) from an individual infection. This work provides new information about the mechanisms of biofilm formation in *M. tb* and outlines a framework for population level approaches for studying bacterial adaptation.

Introduction

Mycobacterium tuberculosis (*M. tb*), the causative agent of tuberculosis (TB), is a globally distributed and highly prevalent pathogen. *M. tb* is uniquely adapted to its environment within human tissues, and is adept at evading both antibiotic treatment and the immune system, resulting in recalcitrant, difficult to treat infections. Increasing drug resistance among populations of *M. tb* is a global health concern: in 2021 there were over 10 million new infections, ~500,000 of which were multi-drug resistant (*1*). Our ability to treat infections in the face of ever-worsening antibiotic resistance hinges on our understanding of how *M. tb* responds and adapts to the various selective pressures encountered during infection and transmission.

M. tb's regulatory system is highly interconnected (*2, 3*) and is structured such that genome-wide patterns of gene expression can be changed rapidly via master regulatory elements (*2, 4*). Features of this regulatory system have been well-studied under a number of conditions relevant to *M. tb* pathogenesis including dormancy (*5*) and hypoxia (*6–8*). These studies have outlined regulatory programs used by *M. tb* to adapt to these conditions including the DosR regulon, which is responsible for coordinating gene expression during non-replicative dormancy. Non-replicative dormancy is thought to allow *M. tb* to persist during latent infection.

Studies of human autopsies (*9–11*) and animal models (*12*) have identified aggregated *M. tb* cells that resemble biofilms, suggesting that biofilms are an important growth form for *M. tb* during natural infection. Biofilms contribute to infection persistence via immune evasion and drug tolerance (*12–15*).

Yet, we know very little about biofilm formation, particularly the regulatory processes governing this

growth form. A single transcriptomics study of *M. tb* pellicle biofilms has been published, in which the

authors identified a role for isonitrile lipopeptide (INLP) in biofilm growth (*16*). In a recent paper we

describe adaptation of *M. tb* clinical isolates to biofilm growth, using an evolve and re-sequence

approach. We discovered that the mutational pathway to enhanced biofilm growth varied between

clinical isolates (*17*). These results led us to hypothesize that the regulatory mechanisms underlying

pellicle biofilm formation exhibit similar variation.

Here we use experimental evolution and high-coverage transcriptomics to delineate transcriptional

responses to pellicle biofilm growth across six clinical strains belonging to two *M. tb* sub-lineages of

lineage 4 (L4.9 and L4.2). We further quantify changes to the transcriptome after application of a

uniform selective pressure for pellicle growth. We found these relatively closely related clinical strains of

M. tb to have diverse transcriptional responses to biofilm growth. The application of a uniform selective

pressure for pellicle growth caused the strains to converge on a shared transcriptome characterized by

wide scale downregulation of gene expression. Within this global pattern we could differentiate two

subtypes that were associated with bacterial sub-lineage. Strains from the same sub-lineage evolved a

characteristic transcriptome, although the mutational path to this shared transcriptome varied from

strain to strain. This implies that pre-existing genetic differences, even among closely related strains,

have a strong impact on transcriptome adaptation. We further found that the *M. tb* biofilm transcriptome

is characterized by widespread upregulation of non-coding RNA (ncRNA) and hypothesize that

expression of ncRNA contributes to genome-wide gene expression modulation.

100 **Results**

Figure 1: Protocol for the serial passage of *M. tb* pellicle biofilms and subsequent transcriptomic comparisons. Ancestral (clinical) populations were grown in planktonic cultures, then grown as pellicle biofilms (passage 1) before passaging 8-12 times. This passaged (evolved) population was taken from a biofilm culture and grown planktonically again. We performed RNA sequencing on all ancestral and evolved populations, grown under both biofilm and planktonic conditions.

RNA sequencing of clinical and experimental M. tb populations

We previously passaged six clinical populations of *M. tb* (isolated from sputum and not subjected to

single-colony passage) under selective pressure to grow as pellicle biofilms as described in T. M. Smith

et al., 2022. We identified phenotypic changes during adaptation to biofilm growth including changes in

cell morphology, increased extracellular matrix (ECM) production and alterations of growth rate. We

also identified genetic changes accompanying enhanced biofilm growth that varied according to the

genetic background of the population (*17*).

Here, we sought to assess how serial passage affected the transcriptome. In order to investigate the

impacts of biofilm adaptation on global gene expression, we performed RNA sequencing on the

ancestral and evolved populations grown under biofilm and planktonic conditions (Figure 1). Two

biological replicates were sequenced for each combination of population, genotype, and condition –

excepting ancestral planktonic MT49, for which only one replicate was sequenced (Table S1). We

extracted between 0.4 and 7 μg of total RNA for each sample and sequenced to high coverage:

between 50-120 million reads per sample (Table S1). Sequencing reads were quality checked,

trimmed, and aligned to the *M. tb* H37Rv reference genome (NCBI accession NC_000962.3). For

calculating expression counts we used HTSeq (*18*) and a custom annotation file (see Methods).

Differential expression analysis between conditions (biofilm and planktonic) and across passaging

(ancestral and evolved) was performed in R using DESeq2 (*19*).

Figure 2: Differentiation of transcriptomes between closely related populations. Left: Whole-genome
sequence phylogeny of ancestral populations used for passaging. Populations fall into two major L4 sub-
lineages: L4.9 (purple) and L4.4 (orange) with SNP distances shown along each branch. Right: Number of
significantly differentially expressed genes (DEGs) between biofilm and planktonic conditions for each ancestral
population.

Figure 3: Genetic background affects transcriptional mechanisms of biofilm growth. A) Experimental diagram highlighting comparator populations: ancestral populations grown as pellicle biofilms are compared to the same populations grown in planktonic cultures. B) Principal component analysis (PCA) of normalized, log-transformed gene expression for ancestral populations. Each point is a biological replicate. Results demonstrate diversity among ancestral transcriptomes, with intermingling of planktonic and biofilm samples. C) Volcano plot summarizing differential expression between ancestral populations grown as biofilms and as planktonic cultures. Log transformed adjusted p-values plotted against the log2 fold change for each gene.

Genes that did not have significant differential expression are shown in grey. Very few genes are significantly differentially expressed when all populations are analyzed together. D) Matrix of individual DEGs shared across ancestral populations. A total of 3874 DEGs are plotted according to whether the gene is upregulated (red), downregulated (blue) or not significantly differentially expressed (white) in each population. *Only 4% of downregulated genes are shared by at least 5 populations while no upregulated genes are shared across more than 4 populations.

Genetic background shapes the biofilm transcriptome

We compared the transcriptomes of ancestral bacterial populations grown under biofilm and planktonic

conditions. Genetic background appears to have a substantial impact on these transcriptomes, which

were consistent among biological replicates but differentiated between populations (Figure 2), with

intermingling of planktonic and biofilm samples (Figure 3B, S1A). Differences among genetic

backgrounds were also evident in the transcriptomic response to biofilm growth. When analyzed

individually, our ancestral populations had between ~ 500-3000 DEGs in comparisons of biofilm with

planktonic conditions (Figure 2, Supplementary Data 1). However, we detected only 143 DEGs with

log2 fold changes > +/-2 when all populations were analyzed together (Figure 3C), compared to an

average of 625 genes per population with log2 fold changes > +/-2 when populations were analyzed

separately (Supplementary Data 1). There is very little overlap in DEGs between populations: only 4%

of downregulated genes are shared by at least 5 populations and no upregulated genes are shared

across more than 4 populations (Figure 3D). To assess whether the effect of genetic background was

reflective of phenotypic differences between ancestral populations, we overlayed wet weight

measurements (the simplest marker of biofilm phenotype which measures the total biomass of the

biofilm) onto normalized, log-transformed gene expression (Figure S1B). We did not identify a simple

relationship between transcriptomic signature and biofilm mass, as samples with similar biomass were

spread across transcriptomic space summarized in the PCA. Overall, our results indicate the biofilm

transcriptome is affected by subtle differentiation between genetic backgrounds separated by just 10s-

100s of SNPs (Figure 2).

Figure 4: Transcriptional responses to biofilm growth following biofilm passaging. A) Experimental diagram highlighting comparator populations: evolved populations grown as pellicle biofilms are compared to the same populations grown in planktonic cultures. Analyses are parallel to those of ancestral populations in Figure 3. B) Principal component analysis (PCA) of normalized, log-transformed gene expression for evolved populations grown as biofilms and planktonic cultures. Biofilm passaging resulted in clear separation of biofilm and planktonic transcriptomes. These transcriptomes were not clearly separated at baseline (Figure 3B) C) Volcano plot summarizing differential expression between evolved populations grown as biofilms and as

planktonic cultures. Log transformed adjusted p-values plotted against the log2 fold change for each gene. Genes that did not have significant differential expression are shown in grey. Biofilm transcriptomes were characterized by broad-scale downregulation of gene expression. D) Matrix of individual DEGs shared across evolved populations. A total of 3568 DEGs are plotted according to whether that gene is upregulated (red), downregulated (blue) or not significantly differentially expressed (white) in each population. 17% of downregulated genes are shared by at least 5 populations (*), while 6% of upregulated genes are shared by at least 5 populations (**). This contrasts with ancestral populations, which did not share any upregulated genes and only shared 4% of downregulated genes across 5 or more populations (Figure 3D).

Biofilm passaging drives uniformity of the biofilm transcriptome and widespread repression of gene
expression

Prior to biofilm passaging, we observed substantial diversity among biofilm transcriptomes in our
sample (Figure 2, 3). The application of a uniform selection pressure resulted in a more uniform biofilm
transcriptome across genetic backgrounds, with clear separation of samples from biofilm and planktonic
conditions: approximately 76% of variance in our samples can be attributed to growth condition
(represented by PC1, Figure 4B). When all populations were analyzed together, approximately 75% of
the genome was involved in the transcriptomic response to biofilm conditions, with ~ 3600 genes
differentially expressed in comparison with planktonic growth (Figure 4C, Supplementary Data 2). In
comparison with ancestral populations, the evolved biofilm transcriptome was characterized by even
more extensive repression of gene expression (Figure 3C, 4C). Looking at the expression of individual
genes, it is obvious that gene expression in evolved populations is more homogenous than in ancestral
populations (Figure S2). This greater uniformity is also reflected in a larger proportion of shared,
downregulated DEGs across populations: 17% in the evolved versus 4% in the ancestral populations
(Figure 3D, 4D). We also observed a small proportion (6%) of DEGs that were upregulated across 5
populations, suggesting uniform recruitment of a cadre of genes under laboratory conditions (Figure
4D).

Notably, there is one sample that is an outlier with respect to DEGs as well as its overall gene
expression profile. This sample, obtained from the biofilm-passaged population of MT31 grown under
planktonic conditions, has a gene expression profile more similar to a biofilm than to other planktonic
samples (Figure 4B, S2B). If we classify this sample as planktonic, we identify few DEGs identified
within this population (Figure 4D). If we instead group planktonic MT31 with the biofilm samples, we
see a stronger contrast between conditions, and a much higher number of DEGs for MT31, many of
which are shared with the other biofilm populations (Figure S3). After extended passaging as a biofilm,

we attempted to grow our populations planktonically again and while most populations reverted to their

ancestral planktonic transcriptomes, this replicate did not. Interestingly, this population was not

defective for planktonic growth (as measured by OD600) and in fact evolved a higher rate of planktonic

growth in response to biofilm passaging (*17*). One feature of these clinical strains is that unlike H37Rv

they are not well adapted to laboratory growth and often develop biofilm-like "clumps" in planktonic

culture, even in the presence of detergents. This phenomenon may explain why this particular replicate

had a biofilm-like transcriptome, despite being grown in planktonic culture. We hypothesize that MT31's

transcriptome failed to shift to a planktonic pattern because it lost transcriptomic flexibility as a result of

biofilm passaging, and/or that such flexibility was unnecessary because the biofilm transcriptome did

not impose any fitness costs under planktonic growth conditions.

Figure 5: Genetic background shapes adaptive trajectories. A) Principal component analysis (PCA) of normalized, log-transformed expression counts from ancestral and evolved biofilm populations. Arrows are drawn between corresponding ancestral and evolved populations, indicating the trajectory of evolution across passaging. Points are colored by sub-lineage of the ancestral population. B) PCA of normalized, log-transformed expression counts from ancestral and evolved populations grown under planktonic conditions. Like panel A, arrows are drawn between corresponding ancestral and evolved populations and are colored by sub-lineage. Biofilm passaging reduced transcriptome diversity, with populations converging on a sub-lineage-specific signature (A); this pattern was not observed under planktonic growth conditions (B). C) Bar plot of DEGs comparing evolved and ancestral biofilm populations broken down by sub-lineage. There are 680 DEGs shared between sub-lineages, and 405 and 200 DEGs unique to L4.4 and L4.9, respectively. Opacity of the bar indicates the subset of DEGs that are either up (dark) or downregulated (light) after passaging. The pellicle biofilm transcriptome is characterized by broad scale downregulation of gene expression with a smaller

complement of genes that are upregulated in a sub-lineage specific manner. D) Differential expression of DEGs unique to each sub-lineage, from the comparison of evolved to ancestral biofilm populations. Log transformed adjusted p-values plotted against the log2 fold change for each gene. Genes that did not have significant differential expression are shown in grey. Lineage 4.9 evolved a larger complement of upregulated genes in response to biofilm passaging.

Genetic background shapes adaptive trajectories

To further investigate how patterns of biofilm gene expression changed after passaging, we compared

transcriptomes of *M. tb* passaged under selection for biofilm growth with corresponding ancestral

populations (Figure 5). These analyses revealed three striking patterns. First, passaging under

selection for biofilm growth reduced overall transcriptome diversity as summarized in a PCA (Figure

5A). Second, convergence of transcriptomes occurred in a lineage-specific pattern, with the evolution of

clearly differentiated groups that corresponded with sub-lineage designations (Figure 5A). Finally, we

found this pattern to be specific to biofilm growth as the transcriptomes of evolved populations grown

planktonically remained intermingled and diverse (Figure 5B). These results indicate that transcriptomic

responses were specific to the selection pressure applied, and that adaptation to this new environment

was constrained by bacterial genetic background.

Gene-by-gene analyses of the transcriptome also revealed lineage-specific patterns. As noted above,

biofilm growth was associated with widespread downregulation of gene expression (compared to

exponential growth in planktonic culture). Biofilm passaging reinforced this pattern, with comparisons of

evolved and ancestral populations revealing further genome-wide downregulation (Figures 5C, 5D). All

differentially expressed genes (DEGs) that were shared among sub-lineages were downregulated

(Figure 5C). Outside of these shared DEGs, the sub-lineages exhibited distinct patterns with sub-

lineage 4.9 showing a larger proportion (39%) of upregulated genes in its DEG repertoire than sub-

lineage 4.4 (6%) (Figure 5C, 5D, Supplementary Data 3). We found previously that *M. tb* populations

within sub-lineage 4.9 all evolved large duplications (*17*), raising the possibility that the relatively large

proportion of upregulated genes observed here arose from the duplication. However, the majority (55%)

of L4.9-specific, upregulated DEGs lie outside the duplicated region (Supplementary Data 3). Thus, the

increased proportion of upregulated genes in L4.9 does not appear to arise directly from increased

gene dosage from the genomic duplication. It is also noteworthy that while all populations in L4.4

evolved similar transcriptomes (Figure 5A), we did not identify any genomic mutations that were shared

across these populations (*17*). Together, these observations implicate a complex relationship between

genetic backgrounds, novel mutations, transcriptional responses, and phenotypes. It appears that the

application of a uniform selection pressure can result in similar phenotypic adaptation (in this case

enhanced biofilm growth) via distinct mutational paths, that result in similar transcriptional responses.

Thus, when exposed to a new environment, sub-populations of *M. tb* may adapt similarly using different

paths to the same goal.

Figure 6: Genome duplication results in complex patterns of gene expression. A) Experimental diagram highlighting comparator populations: comparison of transcriptomes between evolved and ancestral biofilm populations. B) Top: large tandem duplication that evolved in populations MT31 and MT55 under selection to grow as a biofilm. Bottom: coordinates of the duplication vary slightly between populations, with an average duplicated length of 175 kb. C) Left: log2 fold changes in gene expression between evolved and ancestral biofilm populations plotted against the position in the genome. Each point is a single gene, colored according to whether that gene lies inside of the duplicated region for that population. Right: Boxplots of log2 fold changes in gene expression, for genes outside (light blue) and inside (dark blue) the duplication. Genes within the duplication were significantly more upregulated in both populations (Mann Whitney U test with Benjamini-Hochberg correction, *p* < 0.0001).

Expression of large tandem duplication

In our passaging experiment, a near identical genomic duplication emerged repeatedly and specifically

in association with biofilm passage, providing powerful evidence for its association with biofilm growth

(*17*). We have termed this the MMMC duplication, an umbrella term for duplications starting around

3.1Mb (*Rv3100)* with lengths from 150-350 kb, which have been described before both *in vitro* (*20–23*),

and *in vivo* (*24*). In our experiments, the independently arising MMMC duplications became fixed in

biofilm passaged populations and were stable for up to 20 passages, whereas they were absent from

populations passaged as planktonic cultures (*17*). Here we sought to investigate the impact of a large-

scale duplication on the biofilm transcriptome, thereby illuminating potential mechanisms for its impact

on biofilm growth. A more in-depth analysis of the coordinates of the MMMC duplications using sliding-

window DNA sequencing coverage revealed them to be larger than we had originally reported (~175

versus 120 kb) using another method of structural variant identification (*17*). The duplication has slightly

different coordinates in each population, but spans from position ~3.54 to 3.76 Mb, with an average

length of 175 kb, and includes genes *hpx through PPE56* (Figure 6B).

A priori, we might expect that a second copy of a given gene would increase expression by 2x,

corresponding to a log2 fold change (L2FC) value of 1. Such a modest fold change is not likely to be

identified as statistically significant and in fact only 39 duplicated genes were significantly upregulated

in either MT31 or MT55 (Supplementary Data 3). To further investigate impacts of the MMMC

duplication on gene expression, we plotted L2FC values between evolved and ancestral biofilm

populations (Figure 6A) for all genes across the genome and compared changes in duplicated genes to

non-duplicated genes (Figure 6C). Overall patterns of L2FC values seem to differ between the two

populations: MT55 has greater differentiation between ancestral and evolved populations, resulting in

L2FC values of higher magnitude than MT31 (Figure 6C). The distribution of L2FC values within the

duplication is shifted upward relative to genes outside the duplication, a difference that was statistically

significant under both biofilm and planktonic growth conditions (Mann Whitney U test with Benjamini-

Hochberg correction, $p < 0.0001$) for both MT31 and MT55 (Figure 6C, Figure S4B). However, we did

not observe a simple two-fold increase in gene expression within the duplication: genes exhibited a

range of values with some strongly down- and up-regulated. Thus, the transcriptomic impacts of the

gene duplication appear to be complex, rather than being simply characterized by a doubling of gene

dosage.

We expected expression changes of genes within the duplication to be specific to evolved populations

of MT31 and MT55. However, we found two other populations with expression changes within this

region: expression was overall downregulated in evolved populations of MT49 and MT72 under biofilm

conditions (Figure S4A) and in MT49 under planktonic conditions (Figure S4B). The difference was

relatively subtle for MT72 but striking for MT49 (Figure S5A). Coverage plots from DNA sequencing

reads did not reveal a genomic deletion in this region for MT49 (Figure S5B). Using coverage of RNA

sequencing in this region we observed that the ancestral population had relatively high expression of

genes in this region compared to the surrounding area (Figure S5B). Thus, it appears that the

difference in gene expression at this locus arises from increased expression in the ancestral population,

rather than decreased expression in the evolved population. Our hypothesis is that a transient MMMC

duplication arose in the ancestral population during the process of growing the population as a biofilm

for RNA extraction and sequencing. This is further supported by RNA coverage of ~1.25x within this

region indicating that the duplication was present at intermediate frequency when the sample was

sequenced (Figure S5B).

Our observation that duplications in this region of the *M. tb* genome appear repeatedly in association

with a selective pressure to grow as a biofilm (*17*) suggest that genes within the duplication may be

important for robust biofilm formation. Our hypothesis that this adaptation affects fitness via gene

dosage appears to be supported by the transcriptomic data, although the relationship between

duplicated genes and increased expression is complex. To our knowledge, these data represent the

first in-depth description of the effects of large-scale genome duplications on gene expression in

bacteria, expanding our understanding of these variants in the broader context of strategies for

bacterial adaptation to new environments.

Figure 7: Expression of ncRNAs and sORFs mirror coding regions after passaging. A) Volcano plot summarizing differential expression of ncRNAs and sORFs in evolved populations. Log transformed adjusted p-values plotted against the log2 fold change for each gene. Genes that did not have significant differential expression are shown in grey. B) Heatmap of normalized, log-transformed expression counts for ncRNAs and sORFs with significant differential expression in evolved populations. Each column is a single sample from an

evolved population, grown either as a biofilm or in a planktonic culture. Each row is a gene labeled either ncRNA or sORF. Expression values for each gene are normalized to the mean across samples. Samples are clustered by Euclidean distance and plotted as a tree at the top of the heatmap. C) Principal component analysis (PCA) of normalized, log-transformed expression counts of ncRNAs and sORFs from ancestral and evolved biofilm populations. Arrows are drawn between corresponding ancestral and evolved populations, indicating the trajectory of evolution across passaging. Points are colored by sub-lineage of the ancestral population. The effects of biofilm passage on ncRNAs and sORFs mirror that of coding regions: reduced transcriptome diversity and widespread downregulation with populations converging on a sub-lineage-specific signature.

Non-coding RNA and small proteins are a part of the adaptive transcriptome

Advances in transcriptomic and proteomic methods have revealed an abundance of non-coding RNA

(ncRNA) (*25–31*) and small open reading frames (sORFs) (Shell et al., 2015; C. Smith et al., 2022) in

the *M. tb* genome. In *M. tb*, ncRNA are important for gene regulation and adaptation (*34*) and virulence

(*35*). In other bacteria, diversity in the non-coding transcriptome is hypothesized to be a driver of inter-

strain divergence and adaptation (*36*). We hypothesize that these features may similarly play a key role

in *M. tb* adaptation to biofilm growth. We included a custom list of ncRNA and sORFs (Supplementary

Data 4) in our annotation file used for all differential expression analyses in Figures 2-6. To observe

subtle patterns in expression of these non-canonical features we separated them from the rest of the

open reading frames (ORFs) and looked at their differential expression across biofilm passaging, and in

the evolved populations.

We observed that in general, sORFs and ncRNA follow similar patterns to coding regions: there is more

downregulation (both in the number of the features and the magnitude of differential expression) than

upregulation (Figure 7A,B), and across passaging the expression of these features converge in a

genetic background dependent manner (Figure 7C), a phenomenon that does not hold true under

planktonic growth conditions (Figure S6). This reflects what we see with transcriptome-wide adaptation

to biofilm growth, which is that specific, directional changes in gene expression as a result of the

applied selective pressure (biofilm growth in this case), are only observed when that selective pressure

is maintained (Figure 5A,B). One pattern that was particularly striking was the upregulation of ncRNAs

in evolved biofilms: when all populations were analyzed together, ncRNA represent 33% of all strongly

upregulated features (L2FC > 2, Supplementary Data 2) despite comprising only 12% of all features.

Integrating the analysis of all populations with those of individual populations, we find 14 ncRNA that

are strongly upregulated in at least 5 of our individual populations (Table 1). Comparing the distribution

of log2 fold change values of ncRNA with other feature types (ORFs and sORFs), we found that

ncRNAs were significantly more likely to be upregulated in ancestral biofilms (Figure 8A), as well as

being significantly more likely to be upregulated as a result of passaging (Figure 8B). Our results

demonstrate that upregulation of ncRNA seems to be a common feature of the *M. tb* biofilm

transcriptome, and that non-canonical features such as small proteins and non-coding RNAs play an

important role in bacterial adaptation to new environments.

Table 1: Most significantly downregulated (< -5 L2FC) and upregulated (> 2 L2FC) ncRNAs in evolved
populations. Features included in this list are significantly differentially expressed when all populations were
analyzed together, as well as in at least five individual populations. Log2 fold change (L2FC) values given for
analysis of all populations together.
323

Feature	L2FC	Upstream gene	Upstream gene L2FC	Downstream gene	Downstream gene L2FC
ncRv10896	3.15	*gltA2*	-3.31	*Rv0897c*	-0.72
ncRv2240	3.11	*Rv2240c*	1.77	*aceE*	-0.97
ncRv3911Ac	3.05	*sigM*	2.56	*ncRv3911Bc*	0.82
ncRv1969c	2.93	*mce3D*	1.96	*lprM*	1.59
ncRv2379	2.9	*mbtF*	1.24	*mbtE*	0.76
ncRv0386c	2.76	*Rv0386*	0.58	*purT*	0.86
ncRv3480	2.74	*Rv3480c*	-0.16	*Rv3481c*	1.67
ncRv3507c	2.73	*PE_PGRS53*	0.52	*PE_PGRS54*	-1.09
ncRv3793c	2.71	*embC*	0.88	*embA*	0.81
ncRv2611	2.69	*Rv2611c*	0.99	*pgsA1*	1.58
ncRv1118	2.54	*Rv1118c*	1.5	*Rv1119c*	0.36
ncRv0664c	2.48	*vapB8*	1.09	*vapC8*	0.94
ncRv10186c	2.32	*ncRv0186*	-0.63	*bglS*	-0.05
ncRv1835A	2.24	*Rv1835c*	1.08	*ncRv1835B*	0.1
ncRv3917c	-6.22	*parB*	-2.03	*Rv3916c*	-2.38
ncRv12929	-6.84	*Rv2929*	-0.73	*fadD26*	-3.93
ncRv11221	-7	*sigE*	-5.56	*rseA*	-2.55
ncRv13460c	-7.74	*rpsM*	-5.78	*rpmJ*	-5.74
ssr	-8.90	*Rv3099c*	-1.92	*Rv3098A*	-2.75
ncRv1954	-9.49	*Rv1954c*	0.57	*higB*	-3.67

324
325

326
327

Figure 8: Upregulation of ncRNA is a common feature of *M. tb* biofilm growth. A) Log2 fold change (L2FC) values from ancestral populations by feature type. ncRNA have significantly higher L2FC values than ORFs and sORFs (Mann Whitney U Test with Benjamini-Hochberg correction, *p* < 0.0001) indicating they are more likely to be upregulated in biofilms. B) L2FC values between evolved and ancestral biofilms by feature type. ncRNA expression increased significantly more than either ORFs or sORFs after biofilm passage (Mann Whitney U Test with Benjamini-Hochberg correction, *p* < 0.0001).

328
329
330

Discussion

Here we have characterized the diversity of transcriptomes among closely related clinical strains of *M. tb*. We observed genome-wide changes in gene expression in response to the application of a specific selective pressure, resulting in convergence on a biofilm transcriptome that was structured by lineage. Characterization of the pellicle transcriptome revealed a complex relationship between gene dosage, gene expression and the upregulation of non-coding RNA. Overall, this work highlights the complexity of the regulatory system in *M. tb* and provides a framework for considering the transcriptome as a target of selection.

M. tb biofilm transcriptome: one size does not fit all

Little is known about the structure and function of *M. tb* biofilms (*37*), and even less about the transcriptional signatures that define this growth form. To our knowledge there is just one prior study of pellicle biofilm transcriptomics (*16*), which pinpointed a role for isonitrile lipopeptide (INLP) in biofilm growth. In their study of an attenuated strain of *M. tuberculosis* (mc^{2}7000), Richards et al. discovered that a 6-gene cluster termed the INLP synthase complex (*PPE1, Rv0097, fcoT, fadD10, Rv0100, nrp*) was highly upregulated during pellicle growth and they performed further experimental work to identify INLP in mycobacterial biofilms.

In the present study of closely related, fully virulent clinical strains of *M. tb*, we found that the pellicle biofilm transcriptome exhibited substantial variation among strains (Figures 2, 3). This variation is evident in the number of genes differentially expressed under biofilm growth conditions (Figure 2), as well as the identity of genes subject to differential expression. As a result, there are few differentially expressed genes that are shared among strains (Figure 3C, 3D) and strain-to-strain variation dwarfs transcriptomic signals differentiating planktonic and biofilm growth (Figure 3B). Among all this variation, we did identify one strain that exhibited a similar pattern to the strain studied by Richards et al. Four out of the six genes in the INLP synthase complex were upregulated and in the 98th percentile of differentially expressed genes for the strain MT55 (Supplementary Data 1). Genes in the complex were not upregulated to any significant degree in any of the other strains in our study indicating that INLP may be a significant component of the extracellular matrix in some, but not all strains. Integrating our results with those of Richards et al., we conclude that natural populations of *M. tb* exhibit substantial variation in the regulatory mechanisms underlying biofilm growth, such that there appear to be multiple paths to this complex phenotype.

Flattening and global repression in evolved populations

The application of a uniform selection pressure resulted in evolution of a more uniform transcriptome, loss of strain-to-strain variation, and disambiguation of gene expression under conditions of biofilm growth (Figure 4). The evolved transcriptome is characterized by widespread repression of gene expression, a pattern that was evident in ancestral populations but markedly intensified after passaging (Figure 3C, 4C). Of the ~3600 genes differentially expressed across all pellicle passaged populations, there are similar numbers of down and upregulated genes (Figure 4C), however, the magnitude of the most downregulated genes exceeded 3x the magnitude of the most highly upregulated genes (Supplementary Data 2). Similar patterns of widespread repression have been identified previously, in studies of bacteria isolated from sputum (*38*, *39*) and in vitro models of dormancy (*5*). Where the pellicle model differs from sputum and dormancy, however, is in the global regulatory schemes that dictate the transcriptional program. The DosR regulon controls a group of genes that are activated under anaerobic conditions to trigger entry into a non-replicating dormant state (*40*). Expression of the DosR regulon is a feature of the dormancy transcriptome (*5*) and has also been shown to be characterize the sputum transcriptome (*38*, *41*, *42*). Data from Richards et al. show that the DosR regulon was largely downregulated in their pellicle model, relative to exponential growth (*16*). Here, we found that except for one strain (ancestral MT72), all ancestral and evolved biofilms showed significant downregulation of DosR regulon genes (Supplementary Data 1,2). Thus, it appears that while gene expression during pellicle growth shares some similarities to sputum and dormancy, the mechanisms of adaptation to pellicle growth are strain variable, and appear to occur most commonly by a distinct, DosR-independent mechanism.

Signatures of pellicle adaptation in the non-coding transcriptome

Growth as a biofilm was evident in the non-coding transcriptome, in addition to the coding transcriptome. Expression of non-coding RNAs (ncRNAs) was increased relative to that of genes during biofilm growth of ancestral populations (Figure 8A). Evolution under selective pressure for biofilm growth reinforced this pattern with ncRNA being significantly more upregulated than other loci after passaging (Figure 8B). Considering both ncRNAs and small open reading frames (sORFs), adaptation to pellicle growth resulted in emergence of a uniform biofilm transcriptome that was clearly distinguishable from planktonic conditions (Figure 7). We identified 14 ncRNAs that were significantly upregulated (L2FC > 2) and 6 that were significantly downregulated (L2FC < -5) in our pellicle

passaged populations (Table 1). These are candidate loci mediating shifts in gene expression that
enable *M. tb* growth as a biofilm. It has been shown that upregulation of even a single ncRNA can
induce widespread gene repression (*25, 28*). There are many targets of ncRNA regulation including
genes involved in lipid metabolism (*26*), response to iron-limitation (*28*) and two-component regulatory
systems (2CRS) (*30*). Therefore, it is hypothesized that in *M. tb,* ncRNA mediate a rapid response to
new environmental stimuli, whereby changes in expression of a small number of ncRNA can have large
impacts by regulation of many other genes. A similar phenomenon was recently uncovered in *C. difficile*
where ncRNAs were shown to control initiation of sporulation (*43*). We show here that ncRNAs are
upregulated in our pellicle biofilm model and hypothesize that upregulation of ncRNAs contributes to
the widespread changes in gene expression (particularly downregulation) we see in pellicle passaged
populations, a phenomenon we have termed Non-Coding Regulatory Reprogramming (NCRR). Future
work isolating the downstream effects of individual ncRNA on expression may further illuminate the role
of NCRR in the complex regulatory network of *M. tb*.
412
413   *How do transcriptomes evolve?*
414
Experimental evolution studies have shown that many adaptive mutations in bacteria are predicted to
have a regulatory effect. Studies of *Escherichia coli* (*44–46*) and *Pseudomonas aeruginosa* (*47*)
indicate that rapid adaptation to new environments can be accomplished with a few well-placed
mutations in regulatory genes, a pattern we observed in our experiment evolving *M. tb* (*17*). What is
less well-known however, is the impact of these mutations on the transcriptional landscape. We sought
to characterize the effects of these adaptive mutations on the transcriptome and found the most
obvious pattern to emerge after passaging was the intensification of pre-existing patterns observed in
our ancestral populations. Selection for pellicle growth resulted in stronger downregulation of genes
(Figure 3C, 4C), reduction of sample-to-sample diversity (Figure 3B, 4B) and further upregulation of
ncRNA (Figure 8B). Of particular interest is the reduction of diversity between biofilm transcriptomes
after passaging, an indication that these populations converged onto a shared transcriptome under the
same selective pressure. These findings extend observations from previous studies illustrating parallel
transcriptome adaptation after imposition of a uniform selection pressure. Under laboratory conditions,
convergence towards a shared genic transcriptome has been observed several times in bacteria (*47–*
50) and even in fungus (*51*). One study of *P. aeruginosa* transcriptomes from an explanted lung
showed that genetically distinct strains exhibited minimal variability in their transcriptomes (*52*),
indicating that this phenomenon likely holds true for bacterial adaptation during infection. Here, we have
shown that the phenomenon applies to *M. tb*, and that it also extends to the non-coding transcriptome.

We identified a striking phenomenon beyond this convergence on a shared transcriptome. Our findings

suggest that the adaptive trajectories of both the coding and non-coding transcriptomes are shaped by

the genetic background of the bacterial population (Figures 5A, 7B). Populations with differentiated

transcriptomes at baseline evolved shared, lineage-specific transcriptomes under selection for biofilm

growth. This pattern was specific to biofilm growth, as neither convergence nor an effect of genetic

background were evident during planktonic growth of evolved populations (Figures 5B, S6). There are

several, not mutually exclusive, explanations of these phenomena. In our prior study describing

genomic adaptations during this experiment, we found evidence that strain genetic background affects

the types of mutations observed during selection for pellicle growth (*17*). Convergence on a strain-

specific transcriptome could thus reflect the mutational trajectory associated with each genetic

background, since the majority of observed mutations appeared to act by a regulatory mechanism (*17*).

We discuss this possibility in more detail below.

Transcriptomic consequences of mutations

One of the mutations that exhibited an association with lineage is a 175 kb duplication we termed the

MMMC duplication. The MMMC duplication has been observed previously, and structural variants

appear to be common in *M. tb* populations, likely facilitated by recombination among IS elements (*22*).

We previously found the MMMC duplication to be specifically associated with pellicle adaptation and

the L4.9 background. Here, we found that *M. tb* populations from the L4.9 background, known to have

developed the MMMC duplication, converged on a similar transcriptome (Figure 5A).

The evolutionary history of *M. tb* is littered with evidence of structural variants, particularly duplications,

which have formed some of the most important gene families like the Type VII secretion systems (*53*)

and PE/PPE genes (*54*). In addition to the creation of new gene functions, duplications are also known

to increase fitness through gene dosage effects. For example, an attenuated strain of *M. tb* missing

ESX-3 effectors *esxG* and *esxH* developed duplications of a set of paralogous effectors which, when

overexpressed as a result of duplication, compensate for the loss of *esxG* and *esxH* (*55*). Despite their

prevalence in *M. tb*, transcriptional impacts of large structural variants are not well understood. A prior

study of a genomic deletion found that it was associated with loss of expression of genes within the

deletion (*56*), but to our knowledge there are no prior studies investigating transcriptional impacts of

large duplications. Here we sought to identify the transcriptional impacts of the MMMC duplication. We

identified a two-fold increase in gene dosage from the MMMC (*17*), thus we might naively expect the

duplication to result in a two-fold increase in expression of genes within the duplication. However, this

was not the case. Effects of the duplication on gene expression appeared to be highly variable, with

some genes exhibiting expected two-fold increases (log2 fold change of 1) and others with as much as

9-fold increases (log2 fold change of ~3) and 50-fold decreases (log2 fold change of ~ -5) (Figure 6C,

Supplementary Data 3). These findings illuminate a complex relationship between mutation, gene

expression, and phenotype that belies a simple one to one map between gene dose and gene

expression. Our initial hypothesis for the mechanism underlying this complex pattern is that the

duplication of such a large region in the genome interrupts the function of trans-regulatory elements

that are either encoded within, or act on loci within the duplication. Duplication of binding sites for a

trans-regulatory element outside the duplicated region would effectively 'dilute' the effect of that

regulator. Conversely, increased gene dosage of regulators within the duplication may affect

expression of genes within and outside the duplication. In these scenarios the compounded effects of

increased gene dosage and complex regulatory architecture could lead to highly unpredictable

expression of genes in a large-scale duplication. Further analysis of the specific regulators and

regulatory binding sites within the MMMC duplication may help to elucidate patterns in the expression

of duplicated genes.

Another possible explanation for the apparently complex relationship observed between gene

expression and gene dosage is modification of the duplication's impact by concomitant mutations.

However, the data do not support this hypothesis. The two strains from L4.9 did not share any

mutations other than MMMC (*17*), and yet they evolved similar transcriptomes (Figure 5A). In fact,

populations within L4.4 also converged on similar coding and non-coding transcriptomes (Figure 5A,

7B), and they do not share any mutations (*17*). Thus, convergence of the transcriptome does not

appear to be driven by mutational convergence: different mutations on similar genetic backgrounds can

result in similar transcriptomes. Transcriptome diversity has been described previously among clinical

isolates of *M. tb* (*57–59*). Our observations here suggest that this diversity reflects both canalization

(the tendency of a population to produce the same phenotype regardless of genotype) arising from

genetic structure, including neutral haplotype and lineage structure, as well as variation in selection

pressures encountered during natural infection.

Implications & future directions

We know that biofilms represent heterogeneous populations, with sub-populations that differ in

genotype and functional specialization (*60*). Recent work using spatial transcriptomics has also begun

to reveal the ways in which bacterial communities coordinate gene expression to form biofilms (*61, 62*).

Our analyses in this work have compressed the diversity of the *M. tb* biofilm transcriptome within this

population to a single dimension. Future work studying the diversity of gene expression within biofilm

sub-populations will give us more insight into the form and function of *M. tb* biofilms.

In this study we have characterized both the patterns of gene expression within *M. tb* biofilms, as well

as how those patterns change upon the application of a complex selective pressure. We discovered a

cadre of upregulated ncRNA which we believe to be involved in genome-wide gene expression

modulation, a phenomenon we call Non-Coding Regulatory Reprogramming (NCRR). We have further

illustrated the complex effects on expression of large structural variants. These results give us valuable

insight into how *M. tb* adapts to new environments. The complexity of the selection pressure applied in

this experiment, as well as our focus on characterizing different strains of *M. tb* make these results

particularly relevant to the study of natural infections and adaptation of *M. tb* under within-host selection

pressures.

Materials and Methods

Bacterial strains and growth conditions

Six clinical populations (MT31, MT49, MT55, MT72, MT345, and MT540) were initially isolated from

sputum samples and passaged repeatedly as pellicle biofilms as previously described (*17*). Ancestral

(prior to biofilm passaging) and evolved (after biofilm passaging) populations were grown both as

planktonic cultures and biofilms for RNA sequencing (Figure 1). Planktonic cultures of *M. tuberculosis*

were inoculated from freezer stocks into 7H9OTG (Middlebrook 7H9 broth [HiMedia, #M198] containing

0.2% w/v glycerol, 10% v/v OADC supplement [oleic acid, albumin, D-glucose, catalase; Becton

Dickinson, # B12351] and 0.05% w/v Tween-80) and incubated at 37^0C with 5% CO_2 to an OD600 ~1

before RNA extraction. For biofilm growth, 250 μL of planktonic culture was inoculated into 25mL

Sauton's medium (for 1L: 0.5g KH_2PO_4, 0.5g $MgSO_4$, 4g L-Asparagine, 2g Citric acid, 0.05g Ammonium

Iron (III) citrate, 60mL glycerol, adjust pH to 7.0 with NaOH) containing 0.1% w/v $ZnSO_4$, in a 250 mL

flask (Corning, #430281) and incubated at 37^0C, with 5% CO_2, without shaking, with a tight-fitting cap.

After 3 weeks the lid was loosened to allow gas exchange and the cultures grown for an additional 2

weeks (for a total of 5 weeks of growth) before RNA extraction.

RNA extraction

Biofilm samples were harvested by first removing and discarding the media below the biofilm. The

biofilm was resuspended in fresh Sauton's medium, pelleted at 5,000 xg for 10 minutes, then

resuspended in 3 mL of RNAprotect for bacterial cells (Qiagen, #76506). Planktonic cultures were

similarly pelleted and resuspended in RNAprotect. Aliquots (biological replicates) of the RNAprotect

suspensions were frozen at -80°C before extraction. Before RNA extraction, RNAprotect suspensions

were thawed and pelleted at 5,000 xg for 10 minutes before discarding the supernatant. The pellet was

then resuspended in 100 μL of 100 mg/mL lysozyme (Sigma, #L7651) and 150 μL of Proteinase K

(Qiagen, #19131), vortexed, and incubated at 37°C for 30 minutes (vortexed every 10 minutes). The

remainder of the extraction was performed with a modified version of the Illustra RNAspin mini RNA

isolation kit (GE Healthcare, #25-0500-71) protocol: 350 μL of buffer RA1, 3.5 μL of β-mercaptoethanol

(Sigma, #M6250) and 25-50 mg of acid-washed glass beads (Sigma, #G8772) were added to the cell

suspension. Tubes were vortexed at top speed for a total of 3 minutes (6 x 30 sec vortexes with 1

minute ice rest between vortexes) and then centrifuged at 11,000 xg for 1 minute. The supernatant was

removed before proceeding with step 3 of section 7.3 of the kit protocol. RNA quantification was

performed by Qubit using the RNA HS Assay Kit (Invigrogen, #Q32852).

Library preparation & sequencing

Total RNA submitted to the University of Wisconsin-Madison Biotechnology Center was assayed for

purity and integrity via the NanoDrop One Spectrophotometer and Agilent 2100 Bioanalyzer,

respectively. For library preparation, ribosomal RNA was removed from total RNA using the RiboMinus

Eukaryote System v2 (ThermoFisher, #A15026) kit following the directions provided by the

manufacturer. Using the TruSeq Stranded RNA kit (Illumina, Inc., Carlsbad, CA), the ribo-depleted RNA

was fragmented using divalent cations under elevated temperature. Fragmented RNA was copied into

first stranded cDNA using SuperScript II Reverse Transcriptase (Invitrogen, Carlsbad, California, USA)

and random primers. Second strand cDNA was synthesized using a modified dNTP mix (dTTP

replaced with dUTP), DNA Polymerase I, and RNase H. Double-stranded cDNA was cleaned up with

AMPure XP Beads (1.8X) (Agencourt, Beckman Coulter). The cDNA products were incubated with

Klenow DNA Polymerase to add a single 'A' nucleotide to the 3' end of the blunt DNA fragments.

Unique dual indexes (UDI) were ligated to the DNA fragments and cleaned up with two rounds of

AMPure XP beads (0.8X). Adapter ligated DNA was amplified by PCR and cleaned up with AMPure XP

beads (0.8X). Final libraries were assessed for size and quantity using an Agilent DNA1000 chip and

Qubit dsDNA HS Assay Kit (Invitrogen, Carlsbad, California, USA), respectively. Libraries were

standardized to 2nM. Paired-end 150bp sequencing was performed, using standard SBS chemistry on

an Illumina NovaSeq6000 sequencer. Images were analyzed using the standard Illumina Pipeline,

version 1.8.2.

Custom annotation file

A custom annotation file (available at github.com/myoungblom/mtb_ExpEvo_RNA) was made using the

standard annotations for reference strain H37Rv (NCBI accession GCA_000195955.2). To these

annotations we added non-coding RNA (ncRNA) compiled from recent publications (*25–31*) as well as

unpublished data from collaborators (*63*). Finally we included subset of the small open reading frames

(sORFs) discovered by C. Smith et al., 2022: only sORFs identified in multiple technical replicates

using Ribo-RET, that were antisense to or not overlapping any other genes were included (*64*). The

resulting annotation file contains a total of 579 ncRNA and 300 sORFs (Supplementary Data 4).

RNA sequencing data analysis

Raw RNA-sequencing data was processed using an in-house pipeline (code available at

github.com/myoungblom/RNAseq). Briefly, raw data was checked for quality with FastQC v0.11.8

(http://www.bioinformatics.babraham.ac.uk/projects/fastqc) and trimmed using Trimmomatic v0.39 (*65*).

Trimmed reads were mapped to the H37Rv reference genome (NCBI accession NC_000962.3) using

BWA-MEM v0.7.17 (*66*), and Samtools v1.17 (*67*) view and sort were used to process SAM and BAM

files. Assembly quality was determined using Qualimap v2.2.1 BamQC and RNAseq tools (*68*). Finally,

gene expression was counted using HTSeq counts v1.99.2 (*18*) using the custom annotation file

described above with flags '—nonunique none' and '-s reverse' to exclude reads mapping to more than

one feature and to indicate a reverse-stranded library prep, respectively.

Differential expression analysis

Differential expression analysis was performed in R v4.2.2 (*69*) using DESeq2 v1.30.0 (*19*). First, gene

expression counts from HTSeq were loaded using the function 'DESeqDataSetFromHTSeqCount', and

genes with '0' counts in all samples were removed. A regularized-log transformation was applied to the

raw expression counts using the function 'rlog' which normalizes counts with respect to library size –

these counts were used for visualizations including PCA plots and heatmaps. Significantly differentially

expressed genes (DEGs) were identified for each comparison using the function 'DESeq', filtering

results by *p*-value (with Benjamini-Hochberg correction) of < 0.05. For analyses of all populations, all

samples were analyzed with DESeq together using either the growth condition (biofilm or planktonic) or

the genotype (evolved or ancestral) as the design factor. For analyses of individual populations, only

samples from a given population were included in the DESeq analysis, using the same design criteria

as listed above. Scripts for differential expression analyses and figures are available here:

github.com/myoungblom/mtb_ExpEvo_RNA.

Phylogenetic tree

Phylogenetic tree of ancestral populations was inferred using RAxML v8.2.3 as described previously

(*17*) and plotted using ggtree (*70*).

Sliding coverage plots

Sliding window DNA and RNA sequencing coverage was calculated from sorted BAM files created from

the mapping pipeline described above, using Samtools bedcov (*67*) with a window size of 20,000 bp

and a step size of 5,000 bp (code available at github.com/myoungblom/mtb_ExpEvo). Relative

coverage was calculated by dividing each window coverage by the average coverage across the

assembly – as calculated by Qualimap BamQC (*68*).

Data Availability

All RNA sequencing data used for these analyses has been deposited to the NCBI SRA under

accession PRJNA927269. Code for processing of raw RNA sequencing data available at

github.com/myoungblom/RNAseq. Expression counts and differential expression analysis scripts are

available at github.com/myoungblom/mtb_ExpEvo_RNA.

Acknowledgements

MAY was funded by National Science Foundation Graduate Research Fellowship Program under grant

No. DGE-1747503. This work was supported by NIH grant R01-AI113287. The funders had no role in

study design, data collection and interpretation, or the decision to submit the work for publication.

The authors thank the University of Wisconsin-Madison Biotechnology Center for sequencing services,

and Dr. Sarah Fortune's lab for their assistance in compiling a list of non-coding RNA.

Author Contributions

631   MAY: Data curation, formal analysis, methodology, software, visualization, writing – original draft,

632   writing – review & editing

633   TMS: Conceptualization, data curation, investigation, writing – review & editing

634   CSP: Conceptualization, funding acquisition, project administration, supervision, writing – original draft,

635   writing – review & editing

References

1. Global tuberculosis report 2021, (available at https://www.who.int/publications-detail-redirect/9789240037021).

2. R. Chauhan, J. Ravi, P. Datta, T. Chen, D. Schnappinger, K. E. Bassler, G. Balázsi, M. L. Gennaro, Reconstruction and topological characterization of the sigma factor regulatory network of Mycobacterium tuberculosis. *Nat Commun.* **7**, 11062 (2016).

3. J. E. Galagan, K. Minch, M. Peterson, A. Lyubetskaya, E. Azizi, L. Sweet, A. Gomes, T. Rustad, G. Dolganov, I. Glotova, T. Abeel, C. Mahwinney, A. D. Kennedy, R. Allard, W. Brabant, A. Krueger, S. Jaini, B. Honda, W.-H. Yu, M. J. Hickey, J. Zucker, C. Garay, B. Weiner, P. Sisk, C. Stolte, J. K. Winkler, Y. Van de Peer, P. Iazzetti, D. Camacho, J. Dreyfuss, Y. Liu, A. Dorhoi, H.-J. Mollenkopf, P. Drogaris, J. Lamontagne, Y. Zhou, J. Piquenot, S. T. Park, S. Raman, S. H. E. Kaufmann, R. P. Mohney, D. Chelsky, D. B. Moody, D. R. Sherman, G. K. Schoolnik, The Mycobacterium tuberculosis regulatory network and hypoxia. *Nature.* **499**, 178–183 (2013).

4. P. V. Parvati Sai Arun, S. K. Miryala, A. Rana, S. Kurukuti, Y. Akhter, S. Yellaboina, System-wide coordinates of higher order functions in host-pathogen environment upon Mycobacterium tuberculosis infection. *Sci Rep.* **8**, 5079 (2018).

5. M. Kundu, J. Basu, Applications of Transcriptomics and Proteomics for Understanding Dormancy and Resuscitation in Mycobacterium tuberculosis. *Frontiers in Microbiology.* **12** (2021) (available at https://www.frontiersin.org/articles/10.3389/fmicb.2021.642487).

6. E. Iona, M. Pardini, A. Mustazzolu, G. Piccaro, R. Nisini, L. Fattorini, F. Giannoni, Mycobacterium tuberculosis gene expression at different stages of hypoxia-induced dormancy and upon resuscitation. *J Microbiol.* **54**, 565–572 (2016).

7. J. Jiang, C. Lin, J. Zhang, Y. Wang, L. Shen, K. Yang, W. Xiao, Y. Li, L. Zhang, J. Liu, Transcriptome Changes of Mycobacterium marinum in the Process of Resuscitation From Hypoxia-Induced Dormancy. *Front. Genet.* **10** (2020), doi:10.3389/fgene.2019.01359.

8. T. R. Rustad, M. I. Harrell, R. Liao, D. R. Sherman, The Enduring Hypoxic Response of Mycobacterium tuberculosis. *PLOS ONE.* **3**, e1502 (2008).

9. W. Nyka, The chromophobic tubercle bacilli and the problem of endogenous reactivation of tuberculosis. *Mater Med Pol.* **9**, 175–185 (1977).

10. W. Nyka, STUDIES ON MYCOBACTERIUM TUBERCULOSIS IN LESIONS OF THE HUMAN LUNG. A NEW METHOD OF STAINING TUBERCLE BACILLI IN TISSUE SECTIONS. *Am Rev Respir Dis.* **88**, 670–679 (1963).

11. W. Nyka, E. F. O'Neill, A new approach to the study of non-acid-fast mycobacteria. *Ann N Y Acad Sci.* **174**, 862–871 (1970).

12. P. Chakraborty, S. Bajeli, D. Kaushal, B. D. Radotra, A. Kumar, Biofilm formation in the lung contributes to virulence and drug tolerance of Mycobacterium tuberculosis. *Nat Commun.* **12**, 1606 (2021).

13. D. F. Ackart, L. Hascall-Dove, S. M. Caceres, N. M. Kirk, B. K. Podell, C. Melander, I. M. Orme, J. G. Leid, J. A. Nick, R. J. Basaraba, Expression of antimicrobial drug tolerance by attached communities of Mycobacterium tuberculosis. *Pathog Dis.* **70**, 359–369 (2014).

14. A. K. Ojha, A. D. Baughn, D. Sambandan, T. Hsu, X. Trivelli, Y. Guerardel, A. Alahari, L. Kremer, W. R.
Jacobs, G. F. Hatfull, Growth of Mycobacterium tuberculosis biofilms containing free mycolic acids and
harbouring drug-tolerant bacteria. *Molecular Microbiology*. **69**, 164–174 (2008).

15. A. Trivedi, P. S. Mavi, D. Bhatt, A. Kumar, Thiol reductive stress induces cellulose-anchored biofilm formation
in Mycobacterium tuberculosis. *Nature Communications*. **7**, 1–15 (2016).

16. J. P. Richards, W. Cai, N. A. Zill, W. Zhang, A. K. Ojha, Adaptation of Mycobacterium tuberculosis to Biofilm
Growth Is Genetically Linked to Drug Tolerance. *Antimicrobial Agents and Chemotherapy*. **63** (2019),
doi:10.1128/AAC.01213-19.

17. T. M. Smith, M. A. Youngblom, J. F. Kernien, M. A. Mohamed, S. S. Fry, L. L. Bohr, T. D. Mortimer, M. B.
O'Neill, C. S. Pepperell, Rapid adaptation of a complex trait during experimental evolution of Mycobacterium
tuberculosis. *eLife*. **11**, e78454 (2022).

18. S. Anders, P. T. Pyl, W. Huber, HTSeq—a Python framework to work with high-throughput sequencing data.
Bioinformatics. **31**, 166–169 (2015).

19. M. I. Love, W. Huber, S. Anders, Moderated estimation of fold change and dispersion for RNA-seq data with
DESeq2. *Genome Biology*. **15**, 550 (2014).

20. J. S. Abrahams, M. R. Weigand, N. Ring, I. MacArthur, J. Etty, S. Peng, M. M. Williams, B. Bready, A. P.
Catalano, J. R. Davis, M. D. Kaiser, J. S. Oliver, J. M. Sage, S. Bagby, M. L. Tondella, A. R. Gorringe, A.
2022 Preston, Towards comprehensive understanding of bacterial genetic diversity: large-scale
amplifications in Bordetella pertussis and Mycobacterium tuberculosis. *Microbial Genomics*. **8**, 000761
(2022).

21. P. Domenech, A. Rog, J. Moolji, N. Radomski, A. Fallow, L. Leon-Solis, J. Bowes, M. A. Behr, M. B. Reed,
Origins of a 350-Kilobase Genomic Duplication in Mycobacterium tuberculosis and Its Impact on Virulence.
Infection and Immunity. **82**, 2902–2912 (2014).

22. P. Domenech, G. S. Kolly, L. Leon-Solis, A. Fallow, M. B. Reed, Massive Gene Duplication Event among
Clinical Isolates of the Mycobacterium tuberculosis W/Beijing Family. *Journal of Bacteriology*. **192**, 4562–
4570 (2010).

23. B. Weiner, J. Gomez, T. C. Victor, R. M. Warren, A. Sloutsky, B. B. Plikaytis, J. E. Posey, P. D. van Helden,
N. C. G. van Pittius, M. Koehrsen, P. Sisk, C. Stolte, J. White, S. Gagneux, B. Birren, D. Hung, M. Murray, J.
Galagan, Independent Large Scale Duplications in Multiple M. tuberculosis Lineages Overlapping the Same
Genomic Region. *PLOS ONE*. **7**, e26038 (2012).

24. K. Hjort, P. Jurén, J. C. Toro, S. Hoffner, D. I. Andersson, L. Sandegren, Dynamics of Extensive Drug
Resistance Evolution of Mycobacterium tuberculosis in a Single Patient During 9 Years of Disease and
Treatment. *The Journal of Infectious Diseases*. **225**, 1011–1020 (2022).

25. K. B. Arnvig, I. Comas, N. R. Thomson, J. Houghton, H. I. Boshoff, N. J. Croucher, G. Rose, T. T. Perkins, J.
Parkhill, G. Dougan, D. B. Young, Sequence-Based Analysis Uncovers an Abundance of Non-Coding RNA
in the Total Transcriptome of Mycobacterium tuberculosis. *PLoS Pathog*. **7** (2011),
doi:10.1371/journal.ppat.1002342.

26. K. B. Arnvig, D. B. Young, Identification of small RNAs in Mycobacterium tuberculosis. *Mol Microbiol*. **73**,
397–408 (2009).

27. J. M. DiChiara, L. M. Contreras-Martinez, J. Livny, D. Smith, K. A. McDonough, M. Belfort, Multiple small
RNAs identified in Mycobacterium bovis BCG are also expressed in Mycobacterium tuberculosis and
Mycobacterium smegmatis. *Nucleic Acids Research*. **38**, 4067–4078 (2010).

28. E. R. Gerrick, T. Barbier, M. R. Chase, R. Xu, J. François, V. H. Lin, M. J. Szucs, J. M. Rock, R. Ahmad, B.
Tjaden, J. Livny, S. M. Fortune, Small RNA profiling in Mycobacterium tuberculosis identifies MrsI as
necessary for an anticipatory iron sparing response. *PNAS*. **115**, 6464–6469 (2018).

29. A. M. McGuire, B. Weiner, S. T. Park, I. Wapinski, S. Raman, G. Dolganov, M. Peterson, R. Riley, J. Zucker,
T. Abeel, J. White, P. Sisk, C. Stolte, M. Koehrsen, R. T. Yamamoto, M. Iacobelli-Martinez, M. J. Kidd, A. M.
Maer, G. K. Schoolnik, A. Regev, J. Galagan, Comparative analysis of mycobacterium and related
actinomycetes yields insight into the evolution of mycobacterium tuberculosis pathogenesis. *BMC*
Genomics. **13**, 120 (2012).

30. P. Miotto, F. Forti, A. Ambrosi, D. Pellin, D. F. Veiga, G. Balazsi, M. L. Gennaro, C. D. Serio, D. Ghisotti, D.
M. Cirillo, Genome-Wide Discovery of Small RNAs in Mycobacterium tuberculosis. *PLOS ONE*. **7**, e51950
(2012).

31. C.-H. Tsai, C. Baranowski, J. Livny, K. A. McDonough, J. T. Wade, L. M. Contreras, Identification of Novel
sRNAs in Mycobacterial Species. *PLOS ONE*. **8**, e79411 (2013).

32. S. S. Shell, J. Wang, P. Lapierre, M. Mir, M. R. Chase, M. M. Pyle, R. Gawande, R. Ahmad, D. A. Sarracino,
T. R. Ioerger, S. M. Fortune, K. M. Derbyshire, J. T. Wade, T. A. Gray, Leaderless Transcripts and Small
Proteins Are Common Features of the Mycobacterial Translational Landscape. *PLOS Genetics*. **11**,
e1005641 (2015).

33. C. Smith, J. G. Canestrari, A. J. Wang, M. M. Champion, K. M. Derbyshire, T. A. Gray, J. T. Wade, Pervasive
translation in Mycobacterium tuberculosis. *eLife*. **11**, e73980 (2022).

34. A. A. Ostrik, T. L. Azhikina, E. G. Salina, Small Noncoding RNAs and Their Role in the Pathogenesis of
Mycobacterium tuberculosis Infection. *Biochemistry Moscow*. **86**, S109–S119 (2021).

35. G. Shepelkova, V. Evstifeev, M. Averbakch Jr., I. Sivokozov, A. Ergeshov, T. Azhikina, V. Yeremeev, Small
Noncoding RNAs MTS0997 and MTS1338 Affect the Adaptation and Virulence of Mycobacterium
tuberculosis. *Microbiology Research*. **12**, 186–195 (2021).

36. M. Kopf, S. Klähn, I. Scholz, W. R. Hess, B. Voß, Variations in the non-coding transcriptome as a driver of
inter-strain divergence and physiological adaptation in bacteria. *Sci Rep*. **5**, 9560 (2015).

37. J. P. Richards, A. K. Ojha, *Microbiology Spectrum*, in press, doi:10.1128/microbiolspec.MGM2-0004-2013.

38. N. J. Garton, S. J. Waddell, A. L. Sherratt, S.-M. Lee, R. J. Smith, C. Senner, J. Hinds, K. Rajakumar, R. A.
Adegbola, G. S. Besra, P. D. Butcher, M. R. Barer, Cytological and Transcript Analyses Reveal Fat and
Lazy Persister-Like Bacilli in Tuberculous Sputum. *PLOS Medicine*. **5**, e75 (2008).

39. S. Sharma, M. B. Ryndak, A. N. Aggarwal, R. Yadav, S. Sethi, S. Masih, S. Laal, I. Verma, Transcriptome
analysis of mycobacteria in sputum samples of pulmonary tuberculosis patients. *PLoS One*. **12** (2017),
doi:10.1371/journal.pone.0173508.

40. R. L. Leistikow, R. A. Morton, I. L. Bartek, I. Frimpong, K. Wagner, M. I. Voskuil, The Mycobacterium
tuberculosis DosR regulon assists in metabolic homeostasis and enables rapid recovery from nonrespiring
dormancy. *J Bacteriol*. **192**, 1662–1670 (2010).

41. I. Honeyborne, T. D. McHugh, I. Kuittinen, A. Cichonska, D. Evangelopoulos, K. Ronacher, P. D. van Helden,
S. H. Gillespie, D. Fernandez-Reyes, G. Walzl, J. Rousu, P. D. Butcher, S. J. Waddell, Profiling persistent
tubercle bacilli from patient sputa during therapy predicts early drug efficacy. *BMC Medicine*. **14**, 68 (2016).

42. R. P. J. Lai, T. Cortes, S. Marais, N. Rockwood, M. L. Burke, A. Garza-Garcia, S. Horswell, A. K. Sesay, A. O'Garra, D. B. Young, R. J. Wilkinson, Transcriptomic Characterization of Tuberculous Sputum Reveals a Host Warburg Effect and Microbial Cholesterol Catabolism. *mBio.* **12**, e01766-21 (2021).

43. M. Fuchs, V. Lamm-Schmidt, T. Lenče, J. Sulzer, A. Bublitz, J. Wackenreuter, M. Gerovac, T. Strowig, F. Faber, A network of small RNAs regulates sporulation initiation in Clostridioides difficile. *The EMBO Journal.* **n/a**, e112858 (2023).

44. N. Philippe, E. Crozat, R. E. Lenski, D. Schneider, Evolution of global regulatory networks during a long-term experiment with Escherichia coli. *BioEssays.* **29**, 846–860 (2007).

45. P. E. Puentes-Téllez, Á. T. Kovács, O. P. Kuipers, J. D. van Elsas, Comparative genomics and transcriptomics analysis of experimentally evolved Escherichia coli MC1000 in complex environments. *Environ Microbiol.* **16**, 856–870 (2014).

46. L. Wang, B. Spira, Z. Zhou, L. Feng, R. P. Maharjan, X. Li, F. Li, C. McKenzie, P. R. Reeves, T. Ferenci, Divergence involving global regulatory gene mutations in an Escherichia coli population evolving under phosphate limitation. *Genome Biol Evol.* **2**, 478–487 (2010).

47. A. Dötsch, M. Schniederjans, A. Khaledi, K. Hornischer, S. Schulz, A. Bielecka, D. Eckweiler, S. Pohl, S. Häussler, The Pseudomonas aeruginosa Transcriptional Landscape Is Shaped by Environmental Heterogeneity and Genetic Variation. *mBio.* **6**, e00749 (2015).

48. T. F. Cooper, D. E. Rozen, R. E. Lenski, Parallel changes in gene expression after 20,000 generations of evolution in Escherichiacoli. *Proc Natl Acad Sci U S A.* **100**, 1072–1077 (2003).

49. S. S. Fong, A. R. Joyce, B. Ø. Palsson, Parallel adaptive evolution cultures of Escherichia coli lead to convergent growth phenotypes with different gene expression states. *Genome Res.* **15**, 1365–1372 (2005).

50. J. G. Thöming, J. Tomasch, M. Preusse, M. Koska, N. Grahl, S. Pohl, S. D. Willger, V. Kaever, M. Müsken, S. Häussler, Parallel evolutionary paths to produce more than one Pseudomonas aeruginosa biofilm phenotype. *npj Biofilms Microbiomes.* **6**, 1–13 (2020).

51. A. J. Jallet, A. Le Rouzic, A. Genissel, Evolution and Plasticity of the Transcriptome Under Temperature Fluctuations in the Fungal Plant Pathogen Zymoseptoria tritici. *Frontiers in Microbiology.* **11** (2020) (available at https://www.frontiersin.org/article/10.3389/fmicb.2020.573829).

52. A. Kordes, M. Preusse, S. D. Willger, P. Braubach, D. Jonigk, A. Haverich, G. Warnecke, S. Häussler, Genetically diverse Pseudomonas aeruginosa populations display similar transcriptomic profiles in a cystic fibrosis explanted lung. *Nat Commun.* **10**, 3397 (2019).

53. T. D. Mortimer, A. M. Weber, C. S. Pepperell, Evolutionary Thrift: Mycobacteria Repurpose Plasmid Diversity during Adaptation of Type VII Secretion Systems. *Genome Biology and Evolution.* **9**, 398–413 (2017).

54. N. C. Gey van Pittius, S. L. Sampson, H. Lee, Y. Kim, P. D. van Helden, R. M. Warren, Evolution and expansion of the Mycobacterium tuberculosis PE and PPE multigene families and their association with the duplication of the ESAT-6 (esx) gene cluster regions. *BMC Evolutionary Biology.* **6**, 95 (2006).

55. L. Wang, E. Asare, A. C. Shetty, F. Sanchez-Tumbaco, M. R. Edwards, R. Saranathan, B. Weinrick, J. Xu, B. Chen, A. Bénard, G. Dougan, D. W. Leung, G. K. Amarasinghe, J. Chan, C. F. Basler, W. R. Jacobs, J. M. Tufariello, Multiple genetic paths including massive gene amplification allow Mycobacterium tuberculosis to overcome loss of ESX-3 secretion system substrates. *Proceedings of the National Academy of Sciences.* **119**, e2112608119 (2022).

56. C. Duncan, F. B. Jamieson, J. Troudt, L. Izzo, H. Bielefeldt-Ohmann, A. Izzo, C. Mehaffy, Whole
transcriptomic and proteomic analyses of an isogenic M. tuberculosis clinical strain with a naturally occurring
15 Kb genomic deletion. *PLOS ONE*. **12**, e0179996 (2017).

57. Á. Chiner-Oms, M. Berney, C. Boinett, F. González-Candelas, D. B. Young, S. Gagneux, W. R. Jacobs, J.
Parkhill, T. Cortes, I. Comas, Genome-wide mutational biases fuel transcriptional diversity in the
Mycobacterium tuberculosis complex. *Nat Commun*. **10**, 3994 (2019).

58. S. Homolka, S. Niemann, D. G. Russell, K. H. Rohde, Functional Genetic Diversity among Mycobacterium
tuberculosis Complex Clinical Isolates: Delineation of Conserved Core and Lineage-Specific Transcriptomes
during Intracellular Survival. *PLoS Pathog*. **6**, e1000988 (2010).

59. G. Rose, T. Cortes, I. Comas, M. Coscolla, S. Gagneux, D. B. Young, Mapping of Genotype–Phenotype
Diversity among Clinical Isolates of Mycobacterium tuberculosis by Sequence-Based Transcriptional
Profiling. *Genome Biology and Evolution*. **5**, 1849–1862 (2013).

60. J. Jo, A. Price-Whelan, L. E. P. Dietrich, Gradients and consequences of heterogeneity in biofilms. *Nat Rev
Microbiol*. **20**, 593–607 (2022).

61. D. Dar, N. Dar, L. Cai, D. K. Newman, Spatial transcriptomics of planktonic and sessile bacterial populations
at single-cell resolution. *Science*. **373**, eabi4882 (2021).

62. T. Wang, P. Shen, Y. He, Y. Zhang, J. Liu, Spatial transcriptome uncovers rich coordination of metabolism in
E. coli K12 biofilm. *Nat Chem Biol*, 1–11 (2023).

63. T. Barbier, E. R. Gerrick, Unpublished Data (2023).

64. C. Smith, J. G. Canestrari, A. J. Wang, M. M. Champion, K. M. Derbyshire, T. A. Gray, J. T. Wade,
"Pervasive Translation in Mycobacterium tuberculosis" (2021), p. 665208.

65. A. M. Bolger, M. Lohse, B. Usadel, Trimmomatic: a flexible trimmer for Illumina sequence data.
Bioinformatics. **30**, 2114–2120 (2014).

66. H. Li, Aligning sequence reads, clone sequences and assembly contigs with BWA-MEM. *arXiv:1303.3997 [q-
bio]* (2013) (available at http://arxiv.org/abs/1303.3997).

67. H. Li, B. Handsaker, A. Wysoker, T. Fennell, J. Ruan, N. Homer, G. Marth, G. Abecasis, R. Durbin, 1000
Genome Project Data Processing Subgroup, The Sequence Alignment/Map format and SAMtools.
Bioinformatics. **25**, 2078–2079 (2009).

68. K. Okonechnikov, A. Conesa, F. García-Alcalde, Qualimap 2: advanced multi-sample quality control for high-
throughput sequencing data. *Bioinformatics*. **32**, 292–294 (2016).

69. R Core Team, *R: A language and environment for statistical computing* (Vienna, Austria, 2022;
https://www.R-project.org/).

70. G. Yu, D. K. Smith, H. Zhu, Y. Guan, T. T.-Y. Lam, ggtree: an r package for visualization and annotation of
phylogenetic trees with their covariates and other associated data. *Methods Ecol Evol*. **8**, 28–36 (2017).

 Supplement

 Table S1: RNA sequencing sample information. Passage number 1 refers to the ancestral population and
 evolved populations were sequenced after either 8 or 12 passages. Number of mapped reads and percentage of
 reads assigned to a feature were calculated using Qualimap v2.2.1(*68*).

Sample Number	Population	Passage Number	Condition	RNA sequenced (ug)	Mapped reads (millions)	Reads assigned to feature (millions)
1	MT31	1	Planktonic	1.29	73	66
2	MT31	1	Planktonic	1.62	55	50
3	MT31	12	Planktonic	7.04	62	36
4	MT31	12	Planktonic	1.58	53	50
5	MT31	1	Biofilm	0.95	51	32
6	MT31	1	Biofilm	0.98	63	55
7	MT31	12	Biofilm	2.64	79	54
8	MT31	12	Biofilm	3.18	99	39
10	MT49	1	Planktonic	1.52	69	64
11	MT49	12	Planktonic	3.42	76	73
12	MT49	12	Planktonic	2.49	69	56
13	MT49	1	Biofilm	1.54	52	30
14	MT49	1	Biofilm	3.21	117	59
15	MT49	12	Biofilm	2.19	70	34
16	MT49	12	Biofilm	0.95	83	29
17	MT55	1	Planktonic	1.5	69	64
18	MT55	1	Planktonic	0.82	54	50
19	MT55	8	Planktonic	0.44	64	57
20	MT55	8	Planktonic	3.4	75	72
21	MT55	1	Biofilm	0.39	73	70
22	MT55	1	Biofilm	7.33	82	81
23	MT55	8	Biofilm	1.15	75	31
24	MT55	8	Biofilm	1.40	77	32
25	MT72	1	Planktonic	3.15	43	38
26	MT72	1	Planktonic	1.89	62	54
27	MT72	8	Planktonic	1.07	70	58
28	MT72	8	Planktonic	0.72	55	52
29	MT72	1	Biofilm	1.73	52	49
30	MT72	1	Biofilm	4.22	63	54
31	MT72	8	Biofilm	2.24	64	39
32	MT72	8	Biofilm	1.16	54	33
33	MT345	1	Planktonic	0.60	71	58
34	MT345	1	Planktonic	1.14	72	69
35	MT345	8	Planktonic	1.8	73	71

36	MT345	8	Planktonic	0.87	51	49
37	MT345	1	Biofilm	1.05	87	35
38	MT345	1	Biofilm	1.32	87	28
39	MT345	8	Biofilm	0.85	68	28
40	MT345	8	Biofilm	1.2	90	34
41	MT540	1	Planktonic	0.54	69	58
42	MT540	1	Planktonic	1.56	71	68
43	MT540	8	Planktonic	2.73	51	48
44	MT540	8	Planktonic	2.88	72	65
45	MT540	1	Biofilm	2.67	76	74
46	MT540	1	Biofilm	1.91	61	59
47	MT540	8	Biofilm	1.07	63	20
48	MT540	8	Biofilm	1.46	80	33

843
844
845
846

Figure S1: A) Principal component analysis (PCA) of normalized, log-transformed gene expression counts from ancestral populations. Each point represents the total gene expression of a single sample, where the shape indicates the growth condition of the sample, and the color indicates the population. B) PCA (same as in A) colored this time by biofilm wet weight. There is no clear correlation – either among biofilm samples or planktonic samples – between gene expression patterns and ancestral biofilm phenotype (as measured by wet weight).

Figure S2: Heatmaps of normalized, log-transformed expression counts all genes in the genome. Each column is a single sample from an ancestral (A) or evolved (B) population, grown either as a biofilm or in a planktonic culture. Each row is a gene. Heatmap colored by expression values for each gene which are normalized to the mean across samples. Samples are clustered by Euclidean distance and plotted as a tree at the top of the heatmap. Evolved populations have much more uniform biofilm transcriptomes.

848
849
850
851

Figure S3: Treating evolved MT31 outlier as a biofilm. A) Top: Experimental diagram highlighting comparator populations: evolved populations grown as pellicle biofilms are compared to the same populations grown in planktonic cultures. Bottom: Principal component analysis (PCA) of normalized, log-transformed gene expression for evolved populations grown as biofilms and planktonic cultures. B) Differential expression of DEGs between evolved populations grown as biofilms and as planktonic cultures. Log transformed adjusted p-values plotted against the log2 fold change for each gene. Genes that did not have significant differential expression are shown in grey. C) Heatmap of normalized, log-transformed expression counts for DEGs shown in panel B. Each column is a single sample from an evolved population, grown either as a biofilm or in a planktonic culture – the sample identified as an outlier in Figure 4 is highlighted by an arrow. Each row is a DEG. Expression values for each gene are normalized to the mean across samples. Samples are clustered by Euclidean distance and plotted as a tree at the top of the heatmap. D) Matrix of individual DEGs shared across evolved populations. A total of 4373 DEGs are plotted according to if that gene is upregulated (red), downregulated (blue) or not significantly differentially expressed (white) in each population. 24% of downregulated genes are shared by at least 5 populations (*), while 16% of upregulated genes are shared by at least 5 populations (**).

Figure S4: A) Log-2 fold changes (L2FC) for all genes in each biofilm population, separated by their presence in one or both duplications acquired by MT31 and MT55. Mann Whitney U test with Benjamini-Hochberg correction shows significantly higher L2FC values for duplicated genes in MT31 and MT55 (same data shown in Figure 6C), and significantly lower L2FC values for duplicated genes in MT49 and MT72. Comparison is between evolved populations grown as a biofilm and ancestral populations grown as biofilms. B) Same as A, but comparing pellicle evolved populations grown as planktonic cultures, to ancestral populations grown as planktonic cultures. Mann Whitney U test with Benjamini-Hochberg correction shows significantly higher L2FC values for duplicated genes in MT31 and MT55 and significantly lower L2FC values for duplicated genes in MT49.

Figure S5: A) log2 fold change (L2FC) values between evolved and ancestral biofilm populations of MT49. Shown is each gene within the region surrounding the duplication which arose in MT31 and MT55. Points are colored according to whether they lie inside of the duplicated region. B) Top: Relative coverage of DNA sequencing for the ancestral (passage 0) and evolved (passage 12) populations of MT49. Bottom: Relative coverage of RNA sequencing for ancestral and evolved populations of MT49. Two lines per genotype indicate two biological replicates.

Figure S6: Principal component analysis (PCA) of normalized, log-transformed expression counts of ncRNAs and sORFs from ancestral and evolved populations grown in planktonic culture. Arrows are drawn between corresponding ancestral and evolved populations, indicating the trajectory of evolution across passaging. Points are colored by sub-lineage of the ancestral population.

856
857
Supplementary Data
859
860 **SupplementaryData_1.xlsx**: DEGs comparing ancestral populations grown as biofilms, to the same
861 populations grown under planktonic conditions. DEGs provided for multiple analyses: all populations
862 and each population individually.
863
864 **SupplementaryData_2.xlsx**: DEGs comparing pellicle passaged (evolved) populations grown as
865 biofilms, to the same populations grown under planktonic conditions. DEGs provided for multiple
866 analyses: all populations and each population individually.
867
868 **SupplementaryData_3.xlsx**: DEGs comparing pellicle passaged (evolved) populations grown as
869 biofilms, to ancestral populations grown as biofilms. DEGs provided for multiple analyses: all
870 populations, each population individually and for each of the two sub-lineages in our sample (L4.4 and
871 L4.9).
872
873 **SupplementaryData_4.xlsx:** Non-coding RNAs (ncRNAs) and small open reading frames (sORFs)
874 included to make the custom annotation file.