Reference genome of the ant *Lasius platythorax*

Author: Barbara Feldmeyer[1], Nadège Guiglielmoni[2], Joseph Kirangwa[2], Florian Menzel[3], Judit Salces-Ortiz[4], Rosa Fernández[4], Elena Buena Atienza[5,6], Claudio Ciofi[7], Maria Angela Diroma[7], Alessio Iannucci[7], Chiara Natali[7], Ann M. Mc Cartney[8], Olaf Riess[5,6], Nicolas Casadei[5,6], Ann-Marie Waldvogel[2*]

Senckenberg Biodiversity and Climate Research Centre (SBiK-F), Molecular Ecology, Frankfurt, Frankfurt, Germany
Institute of Zoology, Universität zu Köln, Köln, Germany
Institute of Organismic and Molecular Evolution, Johannes Gutenberg University Mainz, Mainz, Germany
Metazoa Phylogenomics Lab, Institute of Evolutionary Biology (CSIC-University Pompeu Fabra), Barcelona, Spain
Institute of Medical Genetics and Applied Genomics, University of Tübingen, Tübingen, Germany
NGS Competence Center Tübingen, Tübingen, Germany
Department of Biology, University of Florence, Sesto Fiorentino (FI), Italy
Genomics Institute, University of California Santa Cruz, USA

ORCID IDs
Barbara Feldmeyer: 0000-0002-0413-7245
Nadège Guiglielmoni:0000-0002-6185-1592
Joseph Kirangwa: 0000-0002-9393-0500
Florian Menzel: 0000-0002-9673-3668
Judit Salces-Ortiz: 0000-0001-5202-7676
Rosa Fernández: 0000-0002-4719-6640
Elena Buena Atienza: 0000-0002-9890-1960
Claudio Ciofi: 0000-0001-8537-8659
Maria Angela Diroma: 0000-0003-1427-8946
Alessio Iannucci: 0000-0001-7729-4412
Chiara Natali: 0000-0002-0293-171X
Ann M. Mc Cartney: 0000-0003-3191-3200
Olaf Riess: 0000-0002-7011-1369
Nicolas Casadei: 0000-0003-2209-0580
Ann-Marie Waldvogel: 0000-0003-2637-0766

corresponding author: Barbara Feldmeyer

Abstract

Ants are a highly diversified insect family of the order Hymenoptera, with many fascinating characteristics such as eusociality, chemical communication, farming, or social parasitism. Moreover, ants frequent a wide variety of habitats from dry deserts, grasslands, and savannas to cold temperate forests. The ability of ants to inhabit such diverse habitat ranges demonstrates their adaptability and ecological resilience. However, little is known about the genetic underpinnings of this vast array of traits and their adaptive potential. Here, we generated a high-quality genome assembly for the ant species *Lasius platythorax* using long-read PacBio HiFi in combination with chromatin conformation capture (Hi-C) sequencing. We successfully assembled the genome into 15 chromosome-level scaffolds ranging from 7.9 to 19.2 Mb and encompassing 204.6 Mb out of 235.3 Mb (total assembly), and a BUSCO score of 86% (Hymenoptera_odb10). Comparative genomics between the two sister species will provide insights into the genomic basis of trait differentiation.

Introduction

Ants inhabit nearly all terrestrial habitats and fulfill multiple ecosystem functions (Elizalde et al. 2020, Wills & Landis 2018), and have many characteristics of interest including social structure, division of labour, morphological and behavioural diversity, and intricate communication systems (Hölldobler and Wilson 1990). The genus *Lasius* (Formicidae: Formicinae) is common and species-rich in many Palearctic habitats and is scientifically significant due to extraordinary behaviours, species interactions, ecological services, but also invasive species dynamics (Hölldobler and Wilson 1990; Cremer et al. 2008; Seifert 2018; Stukalyuk et al. 2020). Some *Lasius* species have evolved a social parasitic lifestyle, where one eusocial species depends on the labour force of another (Buschinger 2009; Savolainen and Vepsalainen 2003). Moreover, fungiculture generates a powerful system for investigating the origin and maintenance of mutualism (Mueller et al. 2005). In ants, fungiculture has evolved at least twice independently: in attines which are members of the Myrmicinae that culture the fungi for food, and in *Lasius* ants that utilise fungi to construct composite nest walls (Maschwitz and Hölldobler 1970; Mueller and Gerardo 2002; Schlick-Steiner et al. 2008). Within the genus *Lasius,* both social parasitism and fungiculture evolved twice (Boudinot, Borowiec, and Prebus 2022; Maruyama et al. 2008). Moreover, *Lasius* ants are highly diverse in their signal and defense chemistry (Akino and Yamaoka 2005; Kern et al. 1997; Wu and Mori 1991; Yamaoka and Akino 2000). Ants of the genus *Lasius* are widely distributed throughout most of the northern hemisphere, from the Indomalaya, Nearctic, Neotropical, Palearctic (Quque and Bles 2021).

Lasius platythorax is a small brownish ant, which is closely related to *Lasius niger* and was recognised as a separate species in only 1991 (Seifert 1991). While *L. niger* is among the most common species throughout Central Europe, inhabiting grasslands and park areas, *L. platythorax* inhabits deciduous forests, bogs, and fens. *L. platythorax* prefers habitats that are more humid and less sunny in comparison to its sister species. These sibling ant species can be separated using morphometric characters and ecological parameters (Seifert 2007). In addition to inhabiting different niches, *L. platythorax* also differs behaviourally, displaying a higher level of aggression than *L. niger* (Seifert 2007). Interestingly, the cuticular hydrocarbons

of *L. platythorax* and *L. niger* also differ significantly. Cuticular hydrocarbons play an important
role in ant communication, particularly in recognition of nestmates, and in waterproofing and
defense against pathogens (Lenoir et al. 2009). *L. platythorax* cuticular hydrocarbons are
characterised by a high abundance of dimethyl and trimethyl alkanes with relatively high chain
lengths (mostly between C33 and C37), and so are longer than those of *L. niger* (Wittke,
Baumgart, and Menzel 2022; Baumgart et al. 2022).
95
To date, genomic research on *Lasius* ants has been limited, with only a single *Lasius* genome
available for the black garden ant *L. niger* (Konorov et al. 2017). A broader understanding of
genomic diversity, intricate communication systems, and adaptive potential within the *Lasius*
genus requires the availability of multiple genome assemblies representing different species.
Here, we present a reference genome of *L. platythorax*, generated via a combination of long-
read Pacific Biosciences (PacBio) HiFi sequencing and chromatin conformation capture (Hi-
C). This genome will provide invaluable opportunities to investigate the genomic basis of
ecological, behavioural, and chemical differentiation among closely related *Lasius* species as
well as gain insights into the mechanisms driving their adaptive potential and species
diversification.

Materials and Methods

Sample collection and DNA extraction

Lasius platythorax is a small, brownish ant. It is distributed across Central Europe and is
typically found in different kinds of woodland, bogs, and fens and avoids urban habitats. The
species is monogynous and colonies can contain several thousands of workers (Seifert 1991).
One *L. platythorax* colony was sampled in the Lennebergwald, close to Mainz in May 2021.
Brood and workers were collected from this single colony for the long-read and Hi-C
sequencing effort. DNA was isolated using the Qiagen MagAttract HMW DNA Kit from a single
larva.

RNA extraction

116 RNA was extracted with the Zymo Direct-zol RNA miniprep kit, from a pool of ten pooled larvae
117 and five pooled workers, according to the manufacturer's instructions. RNA quality and
118 quantity were determined using the Nanodrop 2000 and stored at -80°C.

Long-read sequencing

120 HiFi libraries were prepared with the Express 2.0 Template kit (Pacific Biosciences, Menlo
121 Park, CA, USA) from a single larva and sequenced on a Sequel II/Sequel IIe instrument with
122 30h movie time.
123 HiFi reads were generated using SMRT Link (v10; (Pacific Biosciences, Menlo Park, CA,
124 USA) with default parameters.

Hi-C library and sequencing

Hi-C libraries were prepared from a single larva using Arima Genomics HiC kit, adhering to
the manufacturer's guidelines with specific adjustments, including extended lysis because of
the thick cuticle and overnight chromatin digestion. The sample was flash-frozen and stored

at -70ºC before processing. DNA fragmentation was performed using a Bioruptor Pico
(Diagenode), and libraries were generated with the KAPA Hyper Prep kit (P/N: KK8504).
Library concentration and fragment size distribution were assessed using a Qubit Fluorometer
(Thermo Fisher Scientific) and a Bioanalyzer with the DNA 1000 Kit (Agilent Technologies),
respectively. Arima Hi-C libraries were pooled, denatured, and loaded for paired-end
sequencing on Illumina NovaSeq 6000 system (NovaSeq 6000 S1 Reagent Kit v1.5, 300
cycles).

RNA sequencing

137   Library preparation for full-length mRNASeq was performed using the NEB Ultra II Directional
138   RNA Library Prep with NEBNext Poly(A) mRNA Magenetic Isolation Module and 100ng total
139   RNA as starting material. Sequencing was performed on an Illumina NovaSeq 6000 device
140   with 2x150bp paired-end sequencing protocol and >50M reads per sample.
141

Genome assembly

PacBio HiFi reads were assembled using wtdbg2 v2.5 (Ruan and Li 2020) with parameters -
x ccs -g 200m. Hi-C reads were trimmed using TrimGalore v0.6.10 (Krueger et al. 2023),
mapped to the draft assembly using bowtie2 v2.2.5 (Langmead and Salzberg 2012) and
preprocessed using hicstuff v3.1.5 (Matthey-Doret et al. 2020) with parameters -e DpnII,HinfI
-m iterative. The assembly was then scaffolded based on Hi-C contacts using instaGRAAL
v0.1.6 (Baudry et al. 2020) no-opengl branch with parameters --level 4 --cycles 50 and curated
using the module instagraal-polish with parameters -m polishing -j NNNNNNNNNN. Gaps
were filled using TGS-GapCloser v1.1.1 (Xu et al. 2020) with parameters --ne --minmap_arg
'-x map-hifi'. PacBio HiFi reads were mapped using minimap2 v2.24 (Li 2018) with parameters
--secondary=no --MD -ax map-hifi and the output was provided to HyPo v1.0.3 (Kundu, Casey,
and Sung 2019) for polishing with parameters -k ccs -s 200m -c 70. k-mer completeness was
assessed with KAT comp v2.4.2 (Mapleson et al. 2017) and orthologue completeness was
evaluated using BUSCO v5.0.0 (Manni et al. 2021) against the Hymenoptera odb10 and the
Insecta odb10 lineages in genome mode. Meryl v1.3 and Merqury v1.3 (Rhie et al. 2020)
were run with default parameters using the PacBio HiFi reads. The scaffolds were aligned
against the nucleotide database using the Basic Local Alignment Search Tool (BLAST)
v2.13.0 (Altschul et al. 1990) with parameters -outfmt "6 qseqid staxids bitscore std sscinames
scomnames" -max_hsps 1 -evalue 1e-25. The PacBio HiFi reads were mapped against the
scaffolds using minimap2 v2.24 with parameter -ax map-hifi and the alignment was sorted
using SAMtools v1.17. The outputs of minimap2, BLAST, and BUSCO were provided as input
to BlobTools2 v4.1.5. The Hi-C contact map of the main scaffolds was built using hicstuff as
previously described and visualized using the module hicstuff view with parameter -b 500.
165

Repeat masking and gene annotation

We first used RepeatModeler v2.0.4 (Flynn et al. 2020) to generate a custom repeat library.
We then ran two rounds of RepeatMasker v4.1.4 (Tarailo-Graovac and Chen 2009), the first
round with the custom repeat library, search engine "ncbi". The second round of
RepeatMasker was run with -species "arthropoda". The RepeatMasker output tables were
merged to summarize the abundance or RE categories with the ProcessRepeats function of

RepeatMasker. We used HiSat2 (Kim et al. 2019) to map transcriptome reads to the genome
and used the .bam file as RNA-seq evidence. BRAKER3 (Gabriel et al. 2023) was used to
annotate the softmasked genome using RNA-seq evidence, OrhtoDB11 protein evidence
(Kuznetsov et al. 2023) as well as setting the -AUGUSTUS_ab_initio option. BUSCO v5.2.2
(Manni et al. 2021) against the Hymenoptera odb10 and the Insecta odb10 lineages in protein
mode was used to assess the completeness of the annotation.

Results

The genome of *Lasius platythorax* was assembled from 25.0 Gb of PacBio HiFi reads with an
N50 of 11.3 kb and 273 million pairs of Hi-C reads. Initial assembly with wtdbg2 yielded 3,986
contigs with a total size of 235.3 Mb and an N50 of 106 kb (Table 1). 79% of the Hi-C reads
were mapped to this draft assembly. Hi-C scaffolding using instaGRAAL resulted in 1,374
scaffolds with 15 chromosome-level scaffolds ranging from 7.9 to 19.2 Mb and encompassing
204.6 Mb out of 235.3 Mb (total assembly, Figure 1). Gap filling and polishing closed 204 gaps
to reach 3,332 gaps and increased the assembly size to 235.8 Mb. The final 15 chromosome-
level scaffolds have BUSCO scores of 85.6% (0.4% duplicated features) against Hymenoptera
odb10 and 90.1% (0.8% duplicated features) against Insecta odb10. *k*-mer analysis does not
show uncollapsed haplotypes (Figure 2) and the QV score of 43.7 indicates high overall
accuracy. No contaminants were identified (Figure 3).
Approximately 23% of the genome assembly consists of repetitive elements, which is
comparable to other ant genomes (Hartke et al. 2019). These 23% were mainly annotated as
simple repeats (3.23%) and interspersed repeats (18.72%) (Table 2). Within the latter, the
amount of retroelements (6.02%), DNA transposons (5.88%), and unclassified repeats
(6.83%) is comparable. The annotation with BRAKER3 (Gabriel et al. 2023) using a
combination of RNA-seq and protein evidence as well as ab-initio gene prediction resulted in
the annotation of 17,976 genes. A total of 1208 (88%) complete gene models among 1367
conserved orthologs were identified by BUSCO (Table 2).

Conclusion

199 This newly generated *L. platythorax* genome will provide valuable opportunities to investigate
200 the genomic basis of niche, behavioural and chemical differentiation among closely related
201 *Lasius* species and gain insights into the mechanisms driving their adaptive potential and
202 species diversification.

Data availability

204 The data are part of the ERGA pilot project. Long-read PacBio HiFi data are accessible via
205 BioProject: PRJNA1054215, and BioSample: SRX23153489. Illumina RNAseq and Illumina
206 HiC raw sequences, as well as the assembly and annotation were uploaded to ENA with the
207 accession: PRJEB72067. TolIDs=iyLasPlat10-17:001-008.

Acknowledgments

This genome project is part of the DeRGA pilot study and we acknowledge coordination and
support by Dr. Astrid Böhne and Dr. Philipp Schiffer. We thank Stephan Ossowski for

bioinformatic support with the HiFi data. We acknowledge collaboration with the ERGA Pilot
Project Coordinators, particularly Dr. Alice Mouton and Dr. Giulio Formenti. We furthermore
acknowledge and thank all supplier partners that have kindly donated kits, reagents to the
ERGA pilot Library Preparation Hubs to support species to produce the generation of high-
quality genomes and annotations. This support has been key to embedding a culture of
diversity, equity, inclusion, and justice in the Pilot Project. Specifically we want to thank Arima
Genomics, Integrated DNA Technologies (IDT), Agilent Technologies, Fisher Scientific Spain
and Illumina Inc.

Conflict of interest

220 The authors declare no conflict of interest

Funding information

HiFi sequencing was performed with the support of the DFG-funded NGS Competence Center
Tübingen (INST 37/1049-1). Hi-C library preparation and sequencing was supported by a DFG
ENP grant to PS (grant number: 434028868) and Ramón y Cajal fellowship (grant agreement
no. RYC-2017-22492 funded by MCIN/AEI /10.13039/501100011033 and ESF 'Investing in
your future'), project PID2019-108824GA-I00 funded by MCIN/AEI/10.13039/501100011033,
and by the European Research Council (ERC) under the European's Union's Horizon 2020
research and innovation program (grant agreement no. 948281). JK's position was funded by
a DFG ENP grant (grant number: 434028868). NG's position was first funded through a DFG
grant (458953049) and subsequently through the European Union's Horizon Europe research
and innovation programme under the Marie Skłodowska-Curie grant agreement No
101110569.

References

Akino, Toshiharu, and Ryohei Yamaoka. 2005. "Trail Discrimination Signal of Lasius
Japonicus (Hymenoptera: Formicidae)." *CHEMOECOLOGY* 15(1):21–30. doi:
10.1007/s00049-005-0288-6.
Altschul, SF, Gish, W, Miller, W, Myers, EW, and DJ Lipman. 1990. Basic local alignment
search tool. Journal Molecular Bioliology 215(3):403-10. doi: 10.1016/S0022-
2836(05)80360-2.
Baudry, Lyam, Nadège Guiglielmoni, Hervé Marie-Nelly, Alexandre Cormier, Martial
Marbouty, Komlan Avia, Yann Loe Mie, Olivier Godfroy, Lieven Sterck, J. Mark Cock,
Christophe Zimmer, Susana M. Coelho, and Romain Koszul. 2020. "InstaGRAAL:
Chromosome-Level Quality Scaffolding of Genomes Using a Proximity Ligation-
Based Scaffolder." *Genome Biology* 21(1):148. doi: 10.1186/s13059-020-02041-z.
Boudinot, Brendon, Marek Borowiec, and Matthew Prebus. 2022. "Phylogeny, Evolution, and
Classification of the Ant Genus Lasius, the Tribe Lasiini and the Subfamily
Formicinae (Hymenoptera: Formicidae)." *Systematic Entomology* 47:113–51. doi:
10.1111/syen.12522.
Buschinger, Alfred. 2009. "Social Parasitism among Ants: A Review (Hymenoptera:
Formicidae)." *Myrmecological News* 12:219–35.
Cremer, Sylvia, Line V. Ugelvig, Falko P. Drijfhout, Birgit C. Schlick-Steiner, Florian M.
Steiner, Bernhard Seifert, David P. Hughes, Andreas Schulz, Klaus S. Petersen,
Heino Konrad, Christian Stauffer, Kadri Kiran, Xavier Espadaler, Patrizia d'Ettorre,

Nihat Aktaç, Jørgen Eilenberg, Graeme R. Jones, David R. Nash, Jes S. Pedersen,
and Jacobus J. Boomsma. 2008. "The Evolution of Invasiveness in Garden Ants."
PLOS ONE 3(12):e3838. doi: 10.1371/journal.pone.0003838.
Flynn, Jullien M., Robert Hubley, Clément Goubert, Jeb Rosen, Andrew G. Clark, Cédric
Feschotte, and Arian F. Smit. 2020. "RepeatModeler2 for Automated Genomic
Discovery of Transposable Element Families." *Proceedings of the National Academy*
of Sciences of the United States of America 117(17):9451–57. doi:
10.1073/pnas.1921046117.
Gabriel, Lars, Tomas Bruna, Katharina Hoff, Matthis Ebel, Alexandre Lomsadze, Mark
Borodovsky, and Mario Stanke. 2023. *BRAKER3: Fully Automated Genome*
Annotation Using RNA-Seq and Protein Evidence with GeneMark-ETP, AUGUSTUS
and TSEBRA.
Hölldobler, Bert, and Edward O. Wilson. 1990. *The Ants:* Cambridge, MA: Belknap Press.
Kern, Friedrich, Rüdiger W. Klein, Edelgard Janssen, Hans-Jürgen Bestmann, Athula B.
Attygalle, Doris Schäfer, and Ulrich Maschwitz. 1997. "Mellein, a Trail Pheromone
Component of the Ant Lasius Fuliginosus." *Journal of Chemical Ecology* 23(3):779–
92. doi: 10.1023/B:JOEC.0000006410.35938.49.
Kim, Daehwan, Joseph M. Paggi, Chanhee Park, Christopher Bennett, and Steven L.
Salzberg. 2019. "Graph-Based Genome Alignment and Genotyping with HISAT2 and
HISAT-Genotype." *Nature Biotechnology* 37(8):907–15. doi: 10.1038/s41587-019-
0201-4.
Konorov, Evgenii A., Mikhail A. Nikitin, Kirill V. Mikhailov, Sergey N. Lysenkov, Mikhail
Belenky, Peter L. Chang, Sergey V. Nuzhdin, and Victoria A. Scobeyeva. 2017.
"Genomic Exaptation Enables Lasius Niger Adaptation to Urban Environments." *BMC*
Evolutionary Biology 17(1):39. doi: 10.1186/s12862-016-0867-x.
Krueger, Felix, Frankie James, Phil Ewels, Ebrahim Afyounian, Michael Weinstein, Benjamin
Schuster-Boeckler, Gert Hulselmans, and sclamons. 2023.
"FelixKrueger/TrimGalore: V0.6.10 - Add Default Decompression Path."
Kundu, Ritu, Joshua Casey, and Wing-Kin Sung. 2019. *HyPo: Super Fast & Accurate*
Polisher for Long Read Genome Assemblies.
Kuznetsov, Dmitry, Fredrik Tegenfeldt, Mosè Manni, Mathieu Seppey, Matthew Berkeley,
Evgenia V. Kriventseva, and Evgeny M. Zdobnov. 2023. "OrthoDB V11: Annotation
of Orthologs in the Widest Sampling of Organismal Diversity." *Nucleic Acids*
Research 51(D1):D445–51. doi: 10.1093/nar/gkac998.
Baumgart, Lars, Wittke, Matti, Morsbach S, Abou B, and Florian Menzel. 2022. "Why Do
Ants Differ in Acclimatory Ability? Biophysical Mechanisms behind Cuticular
Hydrocarbon Acclimation across Species." *The Journal of Experimental Biology*
225(16). doi: 10.1242/jeb.243847.
Langmead, Ben, and Steven L. Salzberg. 2012. "Fast Gapped-Read Alignment with Bowtie
2." *Nature Methods* 9(4):357–59. doi: 10.1038/nmeth.1923.
Lenoir, Alain, Stéphanie Depickère, Séverine Devers, Jean-Philippe Christidès, and Claire
Detrain. 2009. "Hydrocarbons in the Ant Lasius Niger: From the Cuticle to the Nest
and Home Range Marking." *Journal of Chemical Ecology* 35(8):913–21. doi:
10.1007/s10886-009-9669-6.
Li, Heng. 2018. "Minimap2: Pairwise Alignment for Nucleotide Sequences." *Bioinformatics*
(Oxford, England) 34(18):3094–3100. doi: 10.1093/bioinformatics/bty191.
Manni, Mosè, Matthew R. Berkeley, Mathieu Seppey, Felipe A. Simão, and Evgeny M.
Zdobnov. 2021. "BUSCO Update: Novel and Streamlined Workflows along with
Broader and Deeper Phylogenetic Coverage for Scoring of Eukaryotic, Prokaryotic,
and Viral Genomes." *Molecular Biology and Evolution* 38(10):4647–54. doi:
10.1093/molbev/msab199.
Mapleson, Daniel, Gonzalo Garcia Accinelli, George Kettleborough, Jonathan Wright, and
Bernardo J. Clavijo. 2017. "KAT: A K-Mer Analysis Toolkit to Quality Control NGS
Datasets and Genome Assemblies." *Bioinformatics (Oxford, England)* 33(4):574–76.
doi: 10.1093/bioinformatics/btw663.

Maruyama, Munetoshi, Florian M. Steiner, Christian Stauffer, Toshiharu Akino, Ross H. Crozier, and Birgit C. Schlick-Steiner. 2008. "A DNA and Morphology Based Phylogenetic Framework of the Ant Genus Lasius with Hypotheses for the Evolution of Social Parasitism and Fungiculture." *BMC Evolutionary Biology* 8(1):237. doi: 10.1186/1471-2148-8-237.

Maschwitz, Ulrich, and Bert Hölldobler. 1970. "Der Kartonnestbau bei Lasius fuliginosus Latr. (Hym. Formicidae)." *Zeitschrift für vergleichende Physiologie* 66(2):176–89. doi: 10.1007/BF00297777.

Matthey-Doret, Cyril, Lyam Baudry, Amaury Bignaud, Axel Cournac, Rémi Montagne, Nadège Guiglielmoni, Théo Foutel-Rodier, and Vittore F. Scolari. 2020. "hicstuff: Simple library/pipeline to generate and handle Hi-C data." Zenodo. *http://doi. org/10.5281/zenodo, 4066363*.

Mueller, Ulrich G., and Nicole Gerardo. 2002. "Fungus-Farming Insects: Multiple Origins and Diverse Evolutionary Histories." *Proceedings of the National Academy of Sciences* 99(24):15247–49. doi: 10.1073/pnas.242594799.

Mueller, Ulrich G., Nicole M. Gerardo, Duur K. Aanen, Diana L. Six, and Ted R. Schultz. 2005. "The Evolution of Agriculture in Insects." *Annual Review of Ecology, Evolution, and Systematics* 36(1):563–95. doi: 10.1146/annurev.ecolsys.36.102003.152626.

Quque, Martin, and Olivier Bles. 2021. "Lasius." Pp. 555–58 in *Encyclopedia of Social Insects*, edited by C. K. Starr. Cham: Springer International Publishing.

Rhie, Arang, Walenz, Brian P., Koren, Sergey and Adam M. Phillippy. 2020. Merqury: reference-free quality, completeness, and phasing assessment for genome assemblies. Genome Biology 21(1):245. doi: 10.1186/s13059-020-02134-9.

Ruan, Jue, and Heng Li. 2020. "Fast and Accurate Long-Read Assembly with Wtdbg2." *Nature Methods* 17(2):155–58. doi: 10.1038/s41592-019-0669-3.

Savolainen, Riitta, and Kari Vepsalainen. 2003. "Sympatric Speciation through Intraspecific Social Parasitism." *Proceedings of the National Academy of Sciences of the United States of America* 100(12):7169–74. doi: 10.1073/pnas.1036825100.

Schlick-Steiner, Birgit C., Florian M. Steiner, Heino Konrad, Bernhard Seifert, Erhard Christian, Karl Moder, Christian Stauffer, and Ross H. Crozier. 2008. "Specificity and Transmission Mosaic of Ant Nest-Wall Fungi." *Proceedings of the National Academy of Sciences of the United States of America* 105(3):940–43.

Seifert, Bernhard. 1991. "Lasius Platythorax n. Sp. a Widespread Sibling Species of Lasius Niger (Hymenoptera: Formicidae)." *Entomologia Generalis* 69–81. doi: 10.1127/entom.gen/16/1991/69.

Seifert, Bernhard. 2007. *Die Ameisen Mittel- und Nordeuropas.* Görlitz / Tauer: Lutra Verl.- u. Vertr.-ges.

Seifert, Bernhard. 2018. *The ants of Central and North Europe.* 1. Auflage. Boxberg: lutra.

Stukalyuk, Stanislav V., Mykola S. Kozyr, Maksym V. Netsvetov, and Vitaliy V. Zhuravlev. 2020. "Effect of the Invasive Phanerophytes and Associated Aphids on the Ant (Hymenoptera, Formicidae) Assemblages." doi: 10.5281/ZENODO.4192900.

Tarailo-Graovac, Maja, and Nansheng Chen. 2009. "Using RepeatMasker to Identify Repetitive Elements in Genomic Sequences." *Current Protocols in Bioinformatics* Chapter 4:4.10.1-4.10.14. doi: 10.1002/0471250953.bi0410s25.

Wittke, Marti, Lucas Baumgart, and Florian Menzel. 2022. "Acclimation in Ants: Interference of Communication and Waterproofing through Cuticular Hydrocarbons in a Multifunctional Trait." *Functional Ecology* 36(8):1973–85. doi: 10.1111/1365-2435.14104.

Wu, Jiang, and Kenji Mori. 1991. "Synthesis of the Enantiomers of 2,6-Dimethyl-6-Hepten-1-Ol, a Mandibular Gland Secretion of the Male Ant Lasius Niger." *Agricultural and Biological Chemistry* 55(10):2667–68. doi: 10.1080/00021369.1991.10870994.

Xu, Mengyang, Lidong Guo, Shengqiang Gu, Ou Wang, Rui Zhang, Brock A. Peters, Guangyi Fan, Xin Liu, Xun Xu, Li Deng, and Yongwei Zhang. 2020. "TGS-GapCloser: A Fast and Accurate Gap Closer for Large Genomes with Low Coverage of Error-Prone Long Reads." *GigaScience* 9(9):giaa094. doi: 10.1093/gigascience/giaa094.

365 Yamaoka, Ryohei, and Toshiharu Akino. 2000. "Evidence for Volatile and Contact Signals of
366   Nestmate Recognition in the Black Shining Ant, Lasius Fulginosus." *Entomological*
367   *Science* 3:1–8.
368

369
370 **Figure 1:** Hi-C contact map of 15 chromosome-level scaffolds.
371

372
373   **Figure 2.** *k*-mer comparison of the HiFi reads v. the final scaffolds.

Figure 3. BlobTools analysis of the final scaffolds. The sequences were either flagged as "Arthropoda" or "no hit", and do not exhibit drastic divergence in coverage or GC content.

379 **Table 1:** Genome features of *Lasius platythorax* including the mitochondrial and plastid
380 genomes.

Assembly ID	*L. platythorax* genome v1
Total length (bp)	235,769,877
No. of contigs	3,986
No. of scaffolds	1,374
Longest scaffold (bp)	19,197,288
N50 (kb)	15,532,059
L50	7
GC content (%)	36.12

382 **Table 2:** Annotation features of the *L. platythorax* genome.

Genome annotation	
Repeat content (%)	23.29
Retrotransposons	6.02
DNA transposons	5.88
Simple repeats	3.23
Unclassified	6.83
No. gene models	17,976
BUSCO (insecta_odb10)	C:88.4%[S:71.8%,D:16.6%],F:1.0%,M:10.6% n:1367