Trophic diversity and evolution in Enantiornithes: a synthesis including new insights from Bohaiornithidae

Case Vincent Miller[1,*], Michael Pittman[2,*], Xiaoli Wang[3,4], Xiaoting Zheng[3,4] and Jen A. Bright[5]

[1]Department of Earth Sciences, The University of Hong Kong, Pokfulam, Hong Kong SAR, China.

[2]School of Life Sciences, The Chinese University of Hong Kong, Shatin, Hong Kong SAR, China.

[3]Institute of Geology and Paleontology, Linyi University, Linyi City, Shandong, 276005, China.

[4]Shandong Tianyu Museum of Nature, Pingyi, Shandong, 273300, China.

[5]Department of Biological and Marine Sciences, University of Hull, Hull HU6 7RX, UK

*Corresponding authors

Abstract

The "opposite birds" Enantiornithines were the dominant birds of the Mesozoic, but our understanding of their ecology is still tenuous. In particular, diets of enantiornithine species have remained speculative until recently. While this new work has been effective at determining diet within groups of enantiornithines, diet data thus far has been too sparse to comment on larger trends in the diversity and evolution of enantiornithine ecology. We introduce new data on the enantiornithine family Bohaiornithidae, famous for their large size and strong teeth and claws. In tandem with previously-published data on the earlier-diverging pengornithids and later-diverging longipterygids, we comment on the breadth of enantiornithine ecology and potential patterns in which it evolved.

Body mass, jaw mechanical advantage, finite element analysis of the jaw, and traditional morphometrics of the claws and skull are compared between bohaiornithids and living birds. The sample size for living bird body mass is over ten times larger than previous studies on longipterygid and pengornithid diet, with implications in interpreting their results. We find bohaiornithids to be ecologically diverse: *Bohaiornis* and *Parabohaiornis* are similar to living plant-eating birds; *Longusunguis* resembles raptorial carnivores; *Zhouornis* is similar to both fruit-eating birds and generalist feeders; and *Shenqiornis* and *Sulcavis* plausibly ate fish, plants, or a mix of both. This ecological diversity is wider than any other enantiornithine family studied previously, which may be driven by strengthening of the jaw relative to other early birds. This strong jaw would allow bohaiornithids to eat harder foods than other birds at the time, but their jaws were weaker than most "strong-jawed" living birds.

With these reconstructions of diet in Bohaiornithidae, there is quantitative support for enantiornithines inhabiting nearly every trophic level. By combining these reconstructions with

past dietary predictions for Longipterygidae and Pengornithidae, we predict the ancestral enantiornithine bird to have been a generalist which ate a wide variety of foods. This would suggest that the ecological diversity of enantiornithine birds represents specialisation in taking foods their ancestors were already eating, rather than many dramatic changes in diet. However, more quantitative data from across the enantiornithine tree is needed to refine this prediction. By the Early Cretaceous, enantiornithine birds had diversified into a variety of ecological niches in a similar way to crown birds after the K-Pg extinction, adding to the body of evidence that traits unique to crown birds (e.g. a toothless beak or cranial kinesis) cannot completely explain their ecological success.

Introduction

The diet of crown birds is extremely broad, and dietary evolution within the crown is a complex mosaic [1]. However, dietary evolution in birds outside the crown is poorly understood, largely because the diet of most Mesozoic birds remains speculative [2]. Some recent studies have begun to elucidate this matter [3-7]. The most progress has been made among enantiornithines [8-10], the most abundant and speciose birds in the Cretaceous [11]. This progress only amounts to an examination of 12 of the over 100 described enantiornithine species [11], though. This has limited any large-scale examinations of the overall trophic diversity of enantiornithines and the patterns in which they diversified. Ideally, the next step to answering these questions is to examine a large group of enantiornithines which are phylogenetically intermediate to the previously-studied families. The enantiornithine family Bohaiornithidae fits both of these requirements, and thus serves as an ideal stepping stone to a large-scale understanding of enantiornithine ecology.

Bohaiornithids are iconic among enantiornithines for their robust teeth, large claws, large

size [12], and iridescent colouration [13]. Their robust teeth have led the clade to be interpreted

as durophagous [14, 15, 16 pg. 83, 17], while their large claws and large body size lead to

raptorial interpretations [12, 16 pg. 270, 18]. Bohaiornithidae has a troubled taxonomic history,

with authors proposing the clade actually represents an evolutionary grade [19, 20] or at least a

group in need of phylogenetic redefinition [21]. But across the literature the original six taxa

referred to Bohaiornithidae (*Bohaiornis*, *Longusunguis*, *Parabohaiornis*, *Shenqiornis*, *Sulcavis*,

and *Zhouornis* [12]; Fig. 2) have consistently resolved as closely-related taxa (Table S1), so for

the purposes of this work we will refer to this group as a clade (see Methods for further

justification). With the recently-described *Beiguornis* [22], Bohaiornithidae is the most speciose

family of enantiornithine birds. Given their unusual morphology and apparent ecological

success, Bohaiornithidae presents an ideal study topic for Mesozoic bird diet.

To investigate bohaiornithid diet, we utilise four quantitative diet proxies: body mass,

mechanical advantage (MA) and related functional indices of the jaws, finite element analysis

(FEA) modelling the jaws during a bite, and traditional morphometric (TM) analysis of claw

shape and size. Size has a strong effect on birds' diets [23-25], so estimating the mass of extinct

birds [26] helps narrow dietary possibilities. Functional indices are ratios of measures of an

animal that inform the mechanical efficiency of body parts to exert or withstand certain forces.

Most commonly, the functional index used in ecology is MA of the jaw, looking at trade-offs

between the speed and force [27-29]. Herein three versions of MA are collected alongside three

other functional indices which have previously discriminated animal diet [2, 30]. Finite element

analysis is an engineering tool used to model forces acting on irregular structures [31]. Here it is

used to model bird jaws during a bite. If a model experiences less strain under the same relative

load as another model, its shape can withstand a greater force before failure. By maintaining a

constant relative load for FEA models, models can be compared in terms of relative strength [32,

33]. TM describes the shape of animal parts with measurements relevant to their ecological role

[34]. Here, measurements are taken of the curvature and relative size of pedal unguals which can

differentiate between non-raptorial and raptorial birds [35-38] and, potentially, between specific

types of raptorial behaviour [8, 36, 39].

These proxies are of little use without reference values, so comparative data is taken from

nearly 200 extant birds [8, 9] including tinamous, flamingos, turacos, strisores, and songbirds

among others. These birds are also ecologically diverse, with diet categories based on the

EltonTraits 1.0 database of bird diet [40]. Table S2 provides cut-offs for diet assignment. Claw

shape is not expected to correlate with diet but the use of talons, following the classification of

[9] separating raptorial birds into raptors taking small prey (which can be completely encircled in

the pes) and those hunting large prey (which cannot be encircled by the pes). With the recent

publication of the bird database AVONET [41], we are able to expand our dataset comparing

mass and diet to 8,758 birds of the 9,994 recognized species of birds.

We expand the framework used in our past works [8, 9] by including traditional

morphometrics of the skull. This line of evidence was recently used successfully in

reconstructing the diet of longipterygid enantiornithines [10], so we expect it to be useful here as

well. We incorporate the extant data from [10], collect new data for bohaiornithids, and analyse

the data with a modified version of their methodology. Notably, we use alternate sources of body

mass and diet data and adjust the data in an attempt to remove the effects of body size.

After each proxy is analysed, they are synthesised into a set of likely diets agreed upon

by different evidence. Combining multiple lines of evidence allows for more precise and

confident diet assignments than any single line can provide. Using this framework [2] we

quantitatively test hypotheses of durophagy and raptorial behaviour in bohaiornithids.

Once dietary predictions are made for Bohaiornithidae, we can begin to examine large-

scale trends in enantiornithine ecology via ancestral state reconstruction. Each of the diet proxies

above can be predicted for the common ancestor of Enantiornithes and placed into the same

framework as any individual species, to create a diet hypothesis for the common ancestor.

However, one would reasonably question if this sample size is large enough to intuit an ancestral

diet reconstruction. To better visualize the uncertainty of our relatively small sample size, we

also include a larger tree with qualitative diet assignments. This tree assumes published diet

hypotheses based on purely qualitative data are correct as a test of what sampling density is

necessary to produce confident ancestral state reconstructions of diet.

Results

Body Mass

Bird masses separated into 4 significantly different combinations of diet categories using

phylogenetic HSD [8, 42]: a. nectarivores; b. granivores and invertivores; c. frugivores and

generalists; and d. folivores and tetrapod hunters. Piscivores and scavengers were not

significantly different from groups b or c. Optimizing the Youden index [43], cut-off points

between these diet groups are as follows: between a and b, 9 g (95% confidence interval [CI] 8–

10 g); between b and c, 56 g (95% CI 41–63 g); and between c and d, 265 g (95% CI 162–303

g). Mass of birds by diet and cut-off points between them are visualised in Fig. 1.

Estimated bohaiornithid body masses are provided in Table 1. Masses range from 91 g to

905 g, x̄ = 287 g, though the largest estimate (*Zhouornis* CNUVB-903) is an outlier. With the

larger *Zhouornis* excluded, masses range from 91 to 406 g, x̄ = 248 g.

Mechanical Advantage and Functional Indices

Graphs of MA results are available in Fig. S1 (univariate) and Fig. 3 (multivariate), with 3D

graphs in supplemental files. Posterior predictions of bohaiornithid diet from FDA are provided

in Table 2. Extant MA and functional index results are unchanged from [9].

Bohaiornithids generally have low anterior jaw-closing mechanical advantage (AMA),

posterior jaw-closing mechanical advantage (PMA), relative articular offset (AO), and relative

maximum mandibular height (MMH) (Fig. S1A–D,G,H,J). Most bohaiornithids also have a low

jaw-opening mechanical advantage (OMA), but *Bohaiornis* has a high OMA. Other functional

indices are intermediate across bohaiornithids.

In PCA, *Longusunguis*, *Shenqiornis*, and *Sulcavis*, all plot near one another in a region

inhabited by invertivores and generalists. This is driven by their low jaw-closing MA (Fig. S2A).

Parabohaiornis, with a slightly higher jaw-closing MA, plots nearby but closer to frugivores.

Bohaiornis plots in an unoccupied region nearest frugivores and tetrapod hunters, separating

from other bohaiornithids by its high upper jaw OMA (Fig. S2A). *Zhouornis* plots between

Bohaiornis and other bohaiornithids near invertivores and granivores due to its high ACH.

In FDA, bohaiornithids other than *Bohaiornis* plot together in an indeterminate region

inhabited by all diets but folivores and husking granivores. This mirrors their high affinity with

generalist feeders (Table 2). *Bohaiornis* plots far from other bohaiornithids and the extant-

inhabited area of the functional phylomorphospace along DA2 (Fig. S2B). The reason for this is

unclear. The only functional index in which *Bohaiornis* differs from other bohaiornithids is

OMA (Fig. S1), which is loaded primarily on DA1 (Fig. S2B). pFDA does not provide

meaningful results when applied to functional index data, likely due to its poor explanatory

power of the extant data [9].

Finite Element Analysis

Graphs of FEA results are available in Fig. 4 (mesh-weighted arithmetic mean [MWAM] strain

[44]) and Fig. 5 (multivariate strain intervals [45]), with 3D graphs in supplemental files.

Posterior predictions of bohaiornithid diet from FDA are provided in Table 3. Extant FEA results

are unchanged from [9].

MWAM strain of bohaiornithids is low (89–156 µε, x̄ = 128 µε), with *Bohaiornis* and

Zhouornis experiencing less strain under loading that any extant carnivorous bird in this study

(min = 105 µε).

In PCA, *Bohaiornis* inhabits a region of the strain-space that is only inhabited by extant

herbivores and *Zhouornis* inhabits a space that is exclusively non-carnivorous. *Longusunguis*,

Parabohaiornis, and *Sulcavis* inhabit a region where all diets overlap. *Shenqiornis* inhabits a

nearby region with slightly less occupation by invertivores, piscivores, and frugivores.

Bohaiornithids tend to display heterogeneous strain, i.e. more model area under high or low

strain rather than under intermediate strain.

In FDA, bohaiornithids tend to plot in uninhabited areas of the strain-space (Fig. 5B).

Thus, the high confidence of posterior predictions from FEA (Table 3) reflect the nearest extant

diet group. As previously noted for this dataset [8], groups with similar distance from the origin

are not meaningfully different in jaw strength, so all bohaiornithids, not just specific taxa, should

be interpreted as plotting similarly to folivores, frugivores, nectarivores, piscivores, and

scavengers. The exception is *Longusunguis*, the bohaiornithid with the second weakest jaw

whose affinities are strongest with generalist feeders. pFDA does not provide meaningful results

when applied to FEA intervals data, likely due to the low phylogenetic signal in these data [9].

Traditional Morphometrics

Pes

Graphs of pedal TM results are available in Fig. 6, with character weights in Fig. l and 3D graphs

in supplemental files. Posterior predictions of bohaiornithid pedal ecology from FDA and pFDA

are provided in Table 4. Extant pedal TM results are unchanged from [9]. Bohaiornithids

generally spread across the phylomorphospace. In PCA, *Sulcavis* plots among large raptors and

both *Parabohaiornis* individuals plot among non-raptorial perching birds (though the latter

separate into an adjacent uninhabited space with PC3 included, see supplemental files).

Longusunguis IVPP V18693, both specimens of *Bohaiornis*, and both specimens of *Zhouornis*

plot in an indeterminate region of primarily small raptors and non-raptorial perching birds.

Longusunguis in particular also plots near tinamous, which are ground birds.

In FDA, *Parabohaiornis* shows a consistent affinity with non-raptorial perching birds

(Table 4). Both specimens of *Bohaiornis* and the younger specimen of *Zhouornis* show affinity

with small raptors as well as non-raptorial perching birds, with the older *Zhouornis* showing

more affinity with ground birds than perching birds (Table 4). *Longusunguis* and *Sulcavis* both

specimens show affinity with large raptors, with Longusunguis also similar to non-raptorial

perchers and Sulcavis to small raptors (Table 4).

While pFDA results differ from [9] due to the use of a new phylogeny and additional

extant data (see Methods), the differences are minimal. The new optimal λ of the extant data is

0.33 (0.31 in [9]), and axis loadings are nearly identical (Fig. lC). In pFDA, *Bohaiornis* LPM

B00167 shows affinity with small raptors, *Longusunguis* with large raptors, and *Zhouornis*

CNUVB-903 with non-raptorial perching birds (Table 4). The remaining bohaiornithid fossils

plot outside the phylomorphospace occupied by extant taxa (Fig. 6C).

Skull

A graph of skull TM as described in its original application to enantiornithines [10] is available

in Fig. S4A. However, as seen in Fig. S4B and discussed at length in the Methods, skull length

appears to be a size proxy that is redundant given the mass data examined in this work. Instead,

we compare relative rostrum length as used in [10] to relative skull length (i.e. skull length

normalised to body size; Fig. 7). Bohaiornithids display distinctly short skulls (relative skull

length < 1.6), with *Zhouornis* displaying the shortest skull relative to body mass of any taxon

sampled. Their relative rostrum lengths fall in the middle of the extant range.

Ancestral State Reconstruction

Qualitative ancestral state reconstruction of enantiornithine diet is presented in Fig. 8. Numerical

likelihoods of each node are available in the supplemental information. Quantitative ancestral

states are provided for body mass (Fig. S5), MA and functional indices (Fig. S6), FEA MWAM

strain (Fig. S7), and pedal TM (Fig. S8). Due to software limitations, these presentations have

polytomies non-randomly resolved and thus are only one possible reconstruction. Table 5

provides more valid values of each quantitative trait for the common ancestor of Enantiornithes

by averaging results from 10,000 random polytomy resolutions. To facilitate interpreting

multivariate traits of the common ancestor of Enantiornithes, LDA posterior probabilities of the

MA data in Table 5 are provided in Table 6 and a PCA graph of the pedal TM data is provided in

Fig. S9.

Discussion

Body Mass

Bohaiornithid masses are most consistent with folivory, frugivory, generalist feeding, piscivory,

scavenging, or tetrapod hunting. Mean estimates of mass for most bohaiornithid specimens

(Table 1) are near the 265 g split between frugivores + generalists + piscivores + scavengers

(group c) and folivores + tetrapod hunters (group d) (Fig. 1), and the uncertainties of both the

split and the masses cause all but two bohaiornithid specimens to fall within both groups c and d.

The exceptions are the indefinite bohaiornithid CUGB P1202, firmly within group c but also

highly immature, and *Zhouornis* CNUVB-903 whose estimated mass is firmly within group d.

The largest specimens of *Bohaiornis*, *Shenqiornis*, *Sulcavis*, and *Zhouornis* have mean mass

estimates which are greater than 265 g, which we interpret as stronger affinity with folivory and

tetrapod hunting than other bohaiornithids.

We note that every bohaiornithid is less massive than any avian obligate scavenger, and

facultative scavenging in birds as small as even the largest bohaiornithid, *Zhouornis* CNUVB-

903, is facilitated by anthropogenic waste [46, 47] and thus may not represent a natural state.

Coupled with past work proposing very large body size is a prerequisite for obligate scavenging

[48], we consider scavenging less likely in bohaiornithids than the other diets mentioned above.

Mechanical Advantage and Functional Indices

As previously observed [8, 9], skull functional indices only effectively separate folivores,

husking granivores, and scavengers in the functional morphospace (Fig. 3). This means dietary

resolution is poor from this line of evidence. However, bohaiornithids also separate into two

distinct functional guilds: *Bohaiornis* and *Parabohaiornis* in one with distinctly high OMA; and

Longusunguis, *Shenqiornis*, *Sulcavis*, and *Zhouornis* with OMA values more in line with the

average extant bird.

Bohaiornis has unusually high OMA (0.16 lower and 0.33 upper). Folivores have a

diagnostically high OMA ($\bar{x}$ = 0.10 lower, 0.27 upper; Fig. S1E–F), making this the strongest

affinity for this taxon (Table 2). Notably though, it has a PMA below any extant folivore (0.16

lower jaw PMA and 0.09 upper jaw PMA, vs a folivore minimum of 0.24 lower jaw PMA and

0.18 upper jaw PMA; Fig. S1C–D). Note that OMA is strongly affected by the length of the

cranium, which may be under different developmental constraints in bohaiornithids (see Skull

Traditional Morphometrics Discussion). However, if anything one would expect bohaiornithids

to have shorter crania and thus lower OMA than an equivalent crown bird, making a distinctly

high OMA even more impactful. *Parabohaiornis* has considerable affinity with swallowing

granivores and folivores (Table 2), respectfully due to its relatively high AMA and PMA and

lower jaw OMA (Fig. S1A–F).

The remaining bohaiornithids plot within the region of undifferentiated diets (Fig. 3) and

have the strongest affinity with dietary generalists (Table 2), previously interpreted as a "default"

prediction in longipterygids [8]. These taxa, however, do show greater affinity with frugivores

and less with invertivores and piscivores compared to pengornithids (though *Shenqiornis* has

nearly the same affinity with piscivores), bringing all three diets into roughly equal likelihood

after generalist feeding. The exception is *Sulcavis*, whose slightly higher OMA increases affinity

with folivores rather than frugivores.

Durophagy, as hypothesized in bohaiornithids [14, 15, 16 pg. 83, 17], is poorly-defined.

Thus we cannot clearly define a "durophage" diet category to test this hypothesis. Rather, we can

test it indirectly by comparing bohaiornithid jaw performance to that of husking granivores and

parrots. Husking granivores are labelled as such in this work based on regular recorded

behaviour of cracking hard seeds (Table S2), and parrots are the archetypical avian durophages

with adaptations across the clade for withstanding high bite forces [49]. The jaw-closing

mechanical advantage of *Parabohaiornis* is higher than the nectarivorous brown lory

(*Chalcopsitta duivenbodei*; upper jaw AMA 0.12 in lory vs 0.15 in *Parabohaiornis*), but smaller

than every other parrot and husking granivore. The average for upper jaw AMA for parrots

(0.23) and husking granivores (0.24) is much higher. The maximum relative jaw height in

bohaiornithids is also below the minimum in parrots, and the same is true for husking granivores

in the lower jaw. The average MCH of husking granivores and bohaiornithids is nearly the same

(respectively 0.35 and 0.34). In short, the upper jaw in bohaiornithids is somewhat robust, but the

upper and lower jaws lack force production adaptations seen in extant avian durophages.

Finite Element Analysis

Bohaiornithid jaws are stronger than both longipterygids' and pengornithids'. Under the same

construction and loading conditions, longipterygid jaws ranged from 259 to 354 $\mu\varepsilon$ [8] and

pengornithid jaws from 190 to 275 $\mu\varepsilon$ [9]. Bohaiornithids, in contrast, have jaw strain ranging

from 89 to 156 $\mu\varepsilon$ (Fig. 4). While they do separate from these other enantiornithine families, the

intermediate strength of their jaws compared to extant birds means they overlap with most diets

in multivariate space (Fig. 5). Thus, dietary diagnosis from FEA is less precise in bohaiornithids

than previously-studied enantiornithines.

In lieu of a clear separation in multivariate space, we can still consider diets unlikely for a

taxon if the MWAM strain of the jaw under loading falls outside the range of any birds with that

diet. However, unlike in Longipterygidae in which the fossil taxa's jaws are weaker than many

extant diet groups [8], bohaiornithid jaws are stronger than most extant birds'. Rather than being

unable to process a given food as in a weaker jaw, we interpret stronger jaws as "overbuilt" for a

given diet. This implies the jaw evolved under pressures for more durable food, but does not

necessarily prohibit consumption of more compliant foods. *Bohaiornis*, *Shenqiornis*, *Sulcavis*,

and *Zhouornis* experience less strain than any swallowing granivore in the extant dataset;

Bohaiornis, *Shenqiornis*, and *Zhouornis* also experience less than any tetrapod hunter; and

Bohaiornis and *Zhouornis* experience less than any carnivore (Fig. 4).

In multivariate space (Fig. 5A) it is apparent that bohaiornithid jaws tend to have a

heterogeneous strain distribution. This contrasts with both scavengers and tetrapod hunters,

which have a diagnostically homogenous strain distribution. To a lesser extent frugivores also

tend to have homogenous strain, though the African green pigeon *Treron calva* is notably the

nearest neighbour of *Zhouornis*.

As mentioned in the previous section, comparison to husking granivores and parrots

warrants additional attention to evaluate hypotheses of durophagy. Bohaiornithids have overall

low jaw strain under loading (89–156 µε, x□ = 128 µε). This is on par with husking granivores

(78–161 µε, x□ = 123 µε), but weaker than parrots (57–95 µε, x□ = 78 µε). Husking granivores

plot across PC2 in Fig. 5, but parrots and bohaiornithids both experience mainly heterogeneous

strain patterns. The locations of high and low strains differ between the two groups, though.

Parrots tend to experience high strain at the rostral tip of the jaw, while the posterior end of the

jaw is under minimal strain. Conversely bohaiornithids tend to experience high strains along the

dorsal edge of the jaw, particularly near the dentary/surangular suture, and low strains along the

ventral edge of the jaw. This may be partially explained by the lack of a keratinous beak in

bohaiornithids, which acts as a strain sink in the rostral jaw of beaked animals [50].

Bohaiornithids have jaws of similar overall strength to husking granivores but weaker than

parrots, and the patterns of strain they experience during a bite are not analogous to either group.

Traditional Morphometrics

Pes

As previously proposed [12], most bohaiornithid talons plot near some extant raptorial birds

(Fig. 6) and have some affinity with them in FDA and pFDA (Table 4). However, most taxa also

have some level of non-raptorial affinity.

Longusunguis is the most consistently raptorial bohaiornithid. It plots within or near large

raptors in every analysis (Fig. 6) and is consistently recovered as likely to be a large raptor

(Table 4). The joints of digits I and II are somewhat hinged (= ginglymoid *sensu* [51]), though

digits III and IV are unhinged [52] indicating grasping ability focussed in the first two digits as

in many extant accipiters [36]. As noted by [12] the tarsometatarsus of *Longusunguis* and other

bohaiornithids is relatively short and robust, typically interpreted as increasing grip strength at

the cost of speed [36, 39, 53] and more common among raptorial avians in ambush predators

[39].

Sulcavis has the strongest raptorial affinity in PCA (Fig. 6A) and has some raptorial

affinity in FDA and pFDA (Fig. 6B; Table 4). Unexpected for a raptorial bird, the specimen

displays a relatively small and straight digit II ungual. Conversely, digit I is the most enlarged

and recurved of any bohaiornithid and among the most enlarged and recurved in the dataset

overall. Enlargement of the hallux without enlargement of any opposing digit is uncommon in

extant birds, mostly occurring in larks (Alaudidae) in which the digit I is also nearly straight.

Digit II is damaged in the holotype of *Sulcavis* [15], and it may be that our estimate of its extent

does not reflect its actual enlargement and curvature. If this is the case then it likely used its feet

raptorially as we hypothesise for *Longusunguis*, but if not then this may indicate some unique

use for digit I other than raptorial grasping.

Zhouornis and *Bohaiornis* can reasonably be interpreted either as small raptors or non-

raptorial perchers. Both have strong affinity with small raptors in FDA (Fig. 6B, Table 4) but

also plot near ground birds and non-raptorial birds in PCA (Fig. 6A). Their claw curvature

(average 91°) is dissimilar from ground birds, rendering that diagnosis less likely. Their

phalanges are moderately hinged [18, 54-56] which indicates some level of grasping adaptation,

useful both for raptors and non-raptorial perchers. The more mature specimen of each bird (B2

and Z1 in Fig. 6A; Fig. s) has claws less similar to raptors and more similar to non-raptorial

perchers. Digits I, II, and IV maintain similar proportions across ontogeny within each species,

but in both species digit III increases in relative size in the more mature specimens (less

confident in *Bohaiornis* as pes preservation is poor in IVPP V17963 [18]), explaining the shift

towards non-raptorial birds.

Parabohaiornis consistently resolves as a non-raptorial perching bird across all analyses

(Fig. 6, Table 4). The taxon has a relatively small digit I compared to other bohaiornithids,

driving this finding. Published image resolution is too poor to tell how hinged their phalanges are

[12].

Skull

Skull traditional morphometrics, while useful in determining longipterygid diet [10], appear less

useful when examining bohaiornithids. Bohaiornithids have rostra to skull length ratios clustered

around 0.8, a region inhabited by birds of all diets except nectarivores and granivores.

Bohaiornithids also do not conform to the general correlation of increasing rostrum length and

skull length. If anything, those with longer skulls have shorter rostra.

We offer two possible interpretations. Firstly, some bohaiornithids may have skull

functions not discussed here. *Bohaiornis*, *Longusunguis*, and *Parabohaiornis*, do plot at the edge

of the extant space near folivores (*Opisthocomus hoazin* and two phasianids), frugivores (two

psittacoids), and a generalist (*Numida meleagris*). *Shenqiornis*, *Sulcavis*, and *Zhouornis*,

however, plot outside the extant space and may represent an ecology dissimilar from any

sampled extant bird.

Alternatively, this may indicate different developmental constraints on the skull in

bohaiornithids (and possibly enantiornithines as a whole). One hypothesis proposes that crown

birds ancestrally have relatively larger brains than their earlier-diverging counterparts [57],

which may necessitate relatively longer skulls to house the larger brain. However, a

subsequently-discovered enantiornithine basicranium displays many crown-bird-like

characteristics and has been used to argue a crown-bird-like endocranium was present in the

common ancestor of Ornithothoraces [58]. Given this uncertainty, we use skull TM only

tentatively to interpret ecology in bohaiornithid taxa and generally recommend caution in

interpreting non-crown bird ecology from measures including the cranium.

Bohaiornithid Ecology and Evolution

By combining the individual lines of evidence discussed above, more precise diet diagnoses can

be obtained. Table 7 summarizes these diagnoses.

Bohaiornis has jaws highly reminiscent of avian folivores (note we use this term to refer

to animals primarily consuming any non-reproductive plant tissues, not strictly leaves), its

relative skull and rostrum size is highly similar to them, and its body mass that falls within the

folivore range as well. As noted in the MA discussion, the most uniquely folivorous trait of this

taxon is a high OMA. While the PMA *Bohaiornis* is below that of any studied folivore, its jaw

strength in FEA is also similar to that of folivores. The other groups it plots near in FEA function

space (Fig. 5A), husking granivores and nectarivores, tend to be much less massive than

Bohaiornis. An herbivorous diet would imply the claws were not used to kill prey, which is

plausible for *Bohaiornis* (Fig. 6). Perching/arboreal lifestyles for living folivorous birds are

uncommon, and these birds are typically weak fliers [59]. The hoatzin (*Opisthocomus hoazin*)

and southern screamer (*Chauna torquata*) have both similar OMA (respectively 0.37 upper 0.07

lower and 0.28 upper 0.13 lower) and MWAM strain (both 121 µε) to *Bohaiornis* (upper OMA

0.33, lower OMA 0.16, MWAM strain 104 µε). Both of these extant folivores typically fly only

short distances with continuous flapping [60, 61] which is also the flight style predicted for

bohaiornithids [19]. These two taxa serve as the most likely extant analogues for *Bohaiornis*;

either birds which spent most of their lives feeding and climbing in trees like the hoatzin [60] or

more ground-based foragers who use trees as a refuge when resting or threatened [61].

Bohaiornis IVPP 17963 plots near the hoatzin in pedal TM (Fig. 6A), so the former lifestyle

seems more likely for at least a mature *Bohaiornis*.

Parabohaiornis has strong herbivorous affinities, though they are less clear than those in

Bohaiornis. Its skull is most similar to swallowing granivores in both MA (Table 2) and FEA

(Table 3), though it is more massive than most extant granivorous birds (178–406 g, vs upper

cutoff of 53 g). Sandgrouse (Pteroclidae) may serve as an extant analogy as swallowing

granivores in the 200–300 g mass range [62]. The two sandgrouse in this study, *Pterocles*

exustus and *Pterocles orientalis*, have overall similar MA (e.g. upper jaw AMA 0.18 and 0.19,

0.15 in *Parabohaiornis*) and skull robusticity (ACH 0.24 and 0.23, 0.25 in *Parabohaiornis*), but

their jaws are weaker under loading in FEA (MWAM strain 311 and 273 µɛ, 156 µɛ in

Parabohaiornis). Also of note, sandgrouse are generally terrestrial while the unguals of

Parabohaiornis indicate an arboreal lifestyle. Alternatively, *Parabohaiornis* shows the next most

affinity with folivores, and this diagnosis is more consistent with its large body mass and

relatively short skull and intermediate-length rostrum. Like *Bohaiornis*, *Parabohaiornis* is

adapted for non-raptorial perching and would likely be analogous to a hoatzin if folivorous.

Generalist feeding is recovered as possible for Parabohaiornis in MA and FEA, but consistently

less so than granivory and folivory so it is treated here as a "default prediction" [8] and not

discussed at length.

The crop and gizzard (i.e. ventriculus or gastric mill) play major roles in folivory and

granivory for extant avians, serving as a site to process the tough plant matter during digestion

[59, 63]. Crops and gizzards have previously been inferred to be absent in Enantiornithes [4],

which could potentially be strong evidence against folivory or granivory in *Bohaiornis* and

Parabohaiornis. Two (non-exclusive) possibilities may still allow for effective herbivory in

these taxa: oral processing and hindgut specialisation. Most groups of herbivorous lizards have

minimal postcranial digestive specialisation, and instead are able to orally process plants to a

level where nutrients can be efficiently extracted [64]. The lack of any accessory cusps as seen in

many herbivorous lizard clades [65] is unsurprising in bohaiornithids, given the simplification of

enamel at the Avialae node [66]. Better analogues are skinks (Scincidae) and tegus (Teiidae),

which retain conical dentition which is simply more robust than their carnivorous and

omnivorous relatives [65]. Bohaiornithids, notably, have some of the most robust teeth within

Enantiornithes [14]. This may have been an adaptation or exaptation for processing plant matter.

Additionally, several groups of extant folivorous birds have specialised hindguts to process plant

matter. Both emus [67] and geese [68] have regions of their digestive tract specialized in

fermenting fibrous foods, and anseriform [69, 70] and galliform [71-73] birds are known to

increase their intestinal lengths when consuming high fibre diets, which allows for a higher

percentage to be digested. While we lack any direct evidence of these adaptations in

Bohaiornithidae, we contest there is evidence for specialised hindgut digestion of plant material

in enantiornithines more broadly. Structures now recognised as plant propagules [74] have been

identified in seven enantiornithine specimens [75]. Of these, four specimens are figured (STM

10-45, STM 29-8 [76], STM 10-12 [75], and STM 11-80 [77]), and in these figured specimens

the propagules are preserved either in the lowest thoracic (STM 10-45 and STM 11-80) or pelvic

(STM 29-8 and STM 10-12) region. As previously noted, consumulite preservation is most likely

with long gut retention times [2], so preservation of these plant propagules specifically in the

region inhabited by the hindgut in life implies a prolonged retention in this area where plant

matter could ferment and break down. So, despite likely lacking a crop and gizzard, *Bohaiornis*

and *Parabohaiornis* could have sustained folivory with oral processing of plant matter and/or

specialised fermentation in the hindgut.

434    Alternatively, as noted in the FEA discussion, our diagnosis of *Bohaiornis* and

435 *Parabohaiornis* as herbivorous is based on the relatively high strength and efficiency of their

436 jaws. These adaptations indicate evolutionary pressure for the ability to consume plant material,

437 but not necessarily a regular habit of doing so. "Fallback foods", foods avoided during normal

438 feeding but essential to survival when alternatives are unobtainable, are well-documented

439 evolutionary drivers of extant animals' functional morphology [78]. The Jehol climate is

440 believed to have been highly variable and affected by local volcanic activity [79], so specialised

441 herbivory may have been restricted to times when this instability limited other food resources. If

442 this is the case, then specialized oral or hindgut processing may not have been necessary to allow

443 *Bohaiornis* and *Paraboahiornis* to survive short-term resource scarcity.

444    *Zhouornis* has the strongest jaw among bohaiornithids (Fig. 4), with total evidence

445 pointing to either a frugivorous or generalist lifestyle. Only frugivores, generalists, husking

446 granivores, and nectarivores inhabit a region of the FEA strain space where jaw strain is as low

447 and heterogeneous as in *Zhouornis*. This space is generally sparsely occupied and thus seems

448 ecologically meaningful. However, *Zhouornis* is also the largest bohaiornithid (mean estimate

449 758 g for the largest specimen), and very few extant granivorous or nectarivorous birds reach this

450 size (Fig. 1). Frugivory and generalist feeding are also recovered as likely from MA posterior

451 predictions (Table 2), so these two diets are considered most likely for *Zhouornis*. Pedal TM

452 results could support either diet, with talons similar to both raptors taking small prey (potentially

453 useful for a generalist) and non-raptorial perching birds (as expected of a frugivore). If

454 *Zhouornis* was indeed a generalist it would potentially have an interesting niche overlap with that

455 proposed for *Pengornis* [9]. *Pengornis* shows signs of generalist feeding with ability to

456 raptorially take large prey. While *Zhouornis*' pes more resembles that of raptors taking prey that

is small relative to body size, *Zhouornis*' absolute body size is much larger than *Pengornis*

(mean mass estimate 437 g [26]). This may represent a different approach to taking advantage of

the same prey resources. We predict *Zhouornis* would be capable of macrocarnivory but less

suited for it than *Pengornis*; the barred owl (*Strix varia*) has a similar body mass and claw shape

to *Zhouornis*, and while it is capable of taking large prey like grouse and rabbits it preferentially

takes mouse-sized prey [80].

Longusunguis, *Shenqiornis*, and *Sulcavis* have less clear diagnoses. Piscivory and

generalist feeding are recovered as likely for all three, with folivory also likely for *Shenqiornis*

and *Sulcavis*. Unlike *Parabohaiornis*, however, there is no clear hierarchy of these possibilities.

Much of the uncertainty stems from FEA giving little useful data, with these taxa having

intermediate MWAM strains when loaded (Fig. 4) and inhabiting a region of the strain-space

shared by most diets (Fig. 5A) only ruling out scavenging and tetrapod hunting. As mentioned

above, their intermediate mass can also only rule out granivory, invertivory, and nectarivory. We

are therefore left with MA to discriminate between the remaining diets, which has poor

predictive power in extant birds. Generalist feeding and piscivory are recovered as somewhat

likely for all three from MA (Table 2), though as previously proposed this could represent a

default prediction [8]. Folivory can also not be ruled out in any of these taxa due to uncertainty

in quadrate placement (Table S3). Pedal TM, however, points to folivory being unlikely in

Longusunguis as its talons appear adapted for taking large animal prey. *Longusunguis* notably

has more recurved teeth than other bohaiornithids [figures 4 and 6 in 17] which is associated

with carnivory [81]. Skull TM of *Longusunguis* are most similar to folivorous birds, but given

the uncertainty in what developmental constraints are at play we consider folivory overall less

likely in *Longusunguis*. Claw shape may also point to carnivory in *Sulcavis*, though as discussed

above its raptorial affinities are questionable. *Shenqiornis* only preserves two disarticulated

talons [82], and while these superficially resemble the claws of *Zhouornis* we cannot

quantitatively reject hypotheses of the pes not being adapted for prey capture, and in turn

hypotheses of folivory, in *Shenqiornis* or *Sulcavis* (*contra* our previous carnivorous assignment

of *Shenqiornis* [2]). In any case, even with four lines of evidence the dietary diagnosis of these

three taxa remains tentative, and investigating additional lines of evidence (e.g. dental

microwear, muscle reconstruction, or sTable S1sotope analysis [2]) is necessary.

Beiguornis preserves neither a skull nor talons [22], so only mass data is available. As

with most bohaiornithids, the only diets which can be considered unlikely based on its mass are

invertivory, granivory, and nectarivory. Further ecological information will require new

techniques or more complete specimens.

Our results support hypotheses of raptorial behaviour [2, 12, 16 pg. 270, 18] in

bohaiornithids, though not for every taxon. We interpret *Longusunguis* as a small predator

adapted to take prey too large to be completely encircled by the pes, and *Zhouornis* as potentially

a large predator whose talons were adapted to take prey which could be fully encircled by the

pes. *Sulcavis* has a greatly enlarged and recurved digit I as expected in a raptor, but lacks any

enlarged opposed digit that could be used to grip. As noted above, this could be taphonomy

obscuring the full extent of digit II or a different, unaccounted for use of the pes. The talons of

Parabohaiornis are most akin to non-raptorial perching birds, and while the talons of *Bohaiornis*

are between those of raptors and non-raptors we interpret them as non-raptorial given the taxon's

herbivorous skull adaptations. Note that the alleged rangle (raptorial gastroliths) used as previous

evidence of raptorial behaviour in *Bohaiornis* [18] has been reinterpreted as post-mortem mineral

growth [83]. Specimens of *Shenqiornis* preserving talons are needed to opine on whether it was

adapted for raptorial behaviour. We previously interpreted *Shenqiornis* as carnivorous from a

combination of its MA and prior interpretations of all other bohaiornithids as raptorial [2], but

with the variety of pedal ecologies identified in this study we now consider its diet indeterminate.

We also support hypotheses of durophagy [14, 15, 16 pg. 83, 17] in bohaiornithids,

though with the caveat that their adaptations are less extreme than what one usually considers an

avian durophage. The first proposal of durophagy in a bohaiornithid is made in contrast to

proposing most other enantiornithines as "general insectivore[s], limited to small arthropods with

softer cuticles" [14 pg. 153]. We support Bohaiornithids as durophagous in the sense that they

have higher jaw strengths than other modelled enantiornithines (see above). They appear adapted

for taking foods harder and tougher than soft-bodied arthropods, but the same could be said for

pengornithids. In living birds, "durophagy" typically refers to cracking hard foods like seeds or

nuts within the beak [84]. As examined in the MA and FEA discussion sections, bohaiornithids

resemble extant husking granivores in both the relative height of their upper jaw and the overall

strain under loading of their lower jaw; however, they lack the adaptations for bite force

production of husking granivores, and fall short of parrots in all potential durophagy metrics.

Bohaiornithids, therefore, do not display adaptations for taking foods as hard as extant birds

commonly considered durophages.

Based on our findings herein, we encourage caution in future works using hypothesized

diet as a variable. Notably when [17] investigates avialan tooth shape in relation to diet,

bohaiornithids are placed into the single "durophage" diet category and are its only members.

Bohaiornithid tooth shape clustered tightly in the study's phylomorphospaces [figures 4–6 in 17],

but the ecological diversity within the group we present here suggests this clustering is driven by

phylogeny rather than, as proposed by the study, diet.

Evolutionary History of Enantiornithine Ecology

Our data suggest that bohaiornithids cover a greater ecological breadth than previously-studied

enantiornithine families (Fig. 8). Longipterygids are reconstructed as mainly invertivorous with

variability in foraging environment [8] (though our mass results here suggest the hypothesis of

Longipteryx as a piscivore cannot be rejected, *contra* [8]), and pengornithids are reconstructed as

generally piscivorous with *Pengornis* diversifying into a more generalist and macrocarnivorous

diet [9]. In contrast, *Bohaiornis* and *Parabohaiornis* show adaptations reminiscent of extant

herbivores while *Longusunguis* and possibly *Zhouornis* were adapted for taking vertebrate prey.

The differences in the mass and talon shape of the latter pair further imply partitioning of the

carnivorous niche.

The increased dietary breadth in Bohaiornithidae appears to stem from increased strength

in the bohaiornithid skull. In particular, Longipterygidae and Bohaiornithidae may represent

opposite evolutionary trajectories from the skull architecture of Pengornithidae, which is

considered plesiomorphic for enantiornithines [85], and from the ancestral enantiornithine skull

(Table 5). Past works [12, 19] have qualitatively described bohaiornithid skulls and teeth as

robust, though when using ACH to quantify robusticity bohaiornithids are not particularly

remarkable. While there is a large difference between longipterygids (ACH 0.09–0.11, $\bar{x}$ = 0.10

[86]) and pengornithids (ACH 0.22–0.23, $\bar{x}$ = 0.23 [87]), bohaiornithid skulls (ACH 0.21–0.26,

$\bar{x}$ = 0.24) are only slightly more robust than those of pengornithids. Ancestral state

reconstruction (Fig. S7K) predicts Bohaiornithidae and Pengornithidae converged upon this level

of robusticity from an ancestor whose ACH was between them and Longipterygidae. What we

find more noteworthy than robusticity is a distinct stepwise increase in jaw strength between

Jehol enantiornithine families: longipterygids have the weakest jaws (259–345 μɛ, $\bar{x}$ = 309 μɛ

[86]), with pengornithid jaws stronger (190–275 µε, x̄ = 222 µε [87]), and bohaiornithids'

stronger still (89–156 µε, x̄ = 128 µε). Ancestral state reconstruction supports bohaiornithids

and longipterygids diversifying from a common ancestor with pengornithid-like jaw strength

(Fig. S7), though a lack of jaw strength data from families outside these three limits the

reliability of this reconstruction. An increase in in jaw strength may have allowed

Bohaiornithidae to process a wider variety of difficult-to-process foods than contemporary

groups and ecologically diversify, but the poor resolution of relationships within and near the

clade [85, 88] (Table S1) makes this possibility difficult to test. For instance the skull of

Fortunguavis, while overall poorly-preserved, does appear to have a robust surangular and

pterygoid [89] and is commonly recovered as sister to Bohaiornithidae. Strengthening of the

skull may therefore be present in more "bohaiornithid-like" enantiornithines (*sensu* [20]) and

even convergent between them. *Shenqiornis*, the geologically oldest bohaiornithid, has a jaw

strength intermediate within bohaiornithids (Fig. 4) which does imply the skull strengthening

adaptations in Bohaiornithidae at least predate 124 Ma. Longipterygids, conversely, were limited

by their weak and gracile jaws to relatively soft and compliant food sources [8]. However,

whether the gracile jaws of longipterygids were adaptations to specialise in soft and compliant

prey, or had unrelated benefits and coincidentally constrained their diet, remains unclear.

Bohaiornithids and Pengornithids are notably larger than other Jehol enantiornithines [2].

While we interpret only *Longusunguis* and possibly *Zhouornis* as inhabiting similar niches to

pengornithids, this size increase does make invertivory unlikely in every pengornithid and

bohaiornithid. The food web of the Jehol Biota is generally understood to be dominated by small

animals [90, 91], particularly in the Jiufotang formation in which bohaiornithids and

pengornithids are most diverse. Accordingly, one would expect interspecific competition for

invertebrates to be high in this ecosystem. Bohaiornithidae and Pengornithidae (alongside the

ornithurine family Songlingornithidae [2]), then, may be lineages which attempted to alleviate

invertivorous competition by diversifying into consuming vertebrate or plant prey. Alternatively,

size increase is also commonly interpreted as defence against predation. *Microraptor* was

definitely capable of preying on smaller enantiornithines [92] and the Jehol biota had many

carnivores larger than *Microraptor* [90, 91]. These possibilities need not be mutually exclusive:

the more abundant large carnivores in the Yixian formation [90] may have provided pressure to

grow larger when these clades first emerged, and the subsequent reduction in large carnivores by

the Jiufotang formation [90] may have allowed them to further diversify. Indeed, ancestral state

reconstruction (Fig. S5) predicts a concurrent increase in body mass for Bohaiornithidae and

Pengornithidae near 125 Ma. Interestingly, though, it recovers the large mass of pengornithids as

convergent, not present ancestrally in the clade. This is driven by the earliest pengornithid,

Eopengornis, being smaller than its more recent relatives. No bohaiornithid is known from this

timeframe, so we cannot yet comment on if their increased body mass is similarly convergent or

ancestral to Bohaiornithidae as a whole.

Given the above trends, we can begin to speculate on the ancestral enantiornithine diet.

The common ancestor of Enantiornithes likely had a body mass between 120 and 160 g (Fig. S5,

Table 5), pointing to frugivory or generalist feeding. Its MA and jaw robusticity was likely

intermediate to low (Fig. S6), overall resembling extant generalists, piscivores, and invertivores

(Table 6). Its jaw strain during a bite is estimated at between 220 $\mu\varepsilon$ and 275 $\mu\varepsilon$ (Table 5), within

the range of Pengornithidae (Fig. S7) and thus consistent with folivory, swallowing granivory,

generalist feeding, and invertivory [9]. The claws of the common ancestor of Enantiornithes are

shaped in a way that is dissimilar from any extant bird included in this study, plotting between

non-raptorial perchers and shrikes (Fig. S9). This is reminiscent of *Eopengornis*, and this claw

shape has been interpreted as arboreal and allowing for very limited prey manipulation [9].

Overall, these lines of evidence point to the common ancestor of Enantiornithes being an

arboreal generalist feeder. If this is the case, then the numerous ecological specialisations seen

within Enantiornithes likely represent lineages already exploiting a resource further specialising

in it, rather than dramatic trophic shifts. One enantiornithine lineage, that leading to both

Longipterygidae and *Eoalulavis*, does appear to be ancestrally invertivorous (Fig. 8). Thus the

purportedly vertivorous avisaurids [93, 94], late-diverging members of this lineage, would

indeed represent a distinct increase in trophic level. However, as illustrated in Fig. 8, the diet of

many of the earliest-diverging enantiornithines remains obscure, and these taxa strongly affect

the interpretation of the ancestral state. Our quantitative reconstruction of the common ancestor

of Enantiornithes in many ways resembles a smaller pengornithid, but that is to be expected

when this family is much closer to the root than other taxa included in MA, FEA, and pedal TM

reconstructions. This highlights the importance of further ecological study of the earliest-

diverging enantiornithine taxa.

While the ancestral enantiornithine diet remains somewhat speculative (Fig. 8), it is clear

that much ecological change and diversification occurred within Enantiornithes by 120 Ma. If

current estimates of the Ornithothoraces node at 145 Ma [95, 96] are correct, then this level of

ecological diversity took only 25 million years to achieve. For comparison, Neoaves (containing

most trophic diversity in crown birds [1]) had only just diverged 25 million years after the origin

of crown birds [97]. This suggests, then, that adaptations within crown Aves like edentulism [17,

98] or cranial kinesis [99] cannot fully explain the rate or extent of ecological diversification of

crown birds after the K-Pg extinction (in agreement with several recent studies [57, 100]). In

fact, crown birds may have been slower to occupy the same ecological breadth as their Mesozoic

counterparts. This reframes some questions in evolutionary biology: while the question of what

made crown birds uniquely fit to survive the K-Pg extinction [57, 99, 101] is still relevant, to

find the traits which allowed for their ecological success and diversity we should look at

Ornithothoraces as a whole. For instance, Ornithothoraces is defined by a variety of adaptations

for improving flight capabilities [102]. Flight allows an organism easy access to multiple vertical

strata within an environment, and flighted vertebrates tend to have larger home ranges than

terrestrial vertebrates [103, 104]. Both of these factors provide volant animals with a greater

variety and amount of resources, and may be a driver of repeated rapid diversification in

ornithothoracine lineages.

Methods

Taxonomic Reference

We refer to extant taxa based on their genus and species in the Birds of the World database for

consistency [105]. Within data files, taxa are referred to by their name in the data source

(Skullsite Bird Skull Collection [106] or museum specimen designation). Comments in data files

note where these identifications differ from Birds of the World or the bird diet database

EltonTraits 1.0 [40]. Designations and relationships of fossil clades are based on [107], though

use of "Bohaiornithidae" in this work is necessarily imprecise due to the instability of the clade

(Table S1, [21]). For reasons explained below, we used Bohaiornithidae to refer to the six taxa

referred when the clade was originally defined [12] plus *Beiguornis* [22].

Taxa Included in Bohaiornithidae

The taxa included within Bohaiornithidae vary between studies. A summary of taxa included in the group in multiple phylogenies is provided in Table S1. The original definition of Bohaiornithidae is "The most recent common ancestor of *Shenqiornis mengi* and *Bohaiornis guoi*, and all its descendants" [12], but in practice this definition is used less strictly. Several studies [22, fig. 3B in 107, 108, 109] place the Bohaiornithidae node at an earlier diverging position than the strict clade definition in order to include *Zhouornis*, and others [19, 21, fig. 5a in 88, 110] recover *Shenqiornis* as so early-diverging that most or all enantiornithines would be considered bohaiornithids by the strict definition. Liu et al. [21] call for a redefinition of the clade after these findings, but do not provide a formal one. They place the Bohaiornithidae node on its phylogeny with a set of synapomorphies, but include *Gretcheniao* and *Junornis* in the clade which is not supported by subsequent studies [20, 22, 85]. Overall, the six original taxa proposed as members of Bohaiornithidae by [12] tend to be recovered as closely-related to *Bohaiornis*. *Longusunguis* is recovered in a clade with *Bohaiornis* in 16 of 22 (73%) studies, *Sulcavis* and *Zhouornis* in 18 of 23 (78%), *Shenqiornis* in 18 of 22 (82%), and *Parabohaiornis* in 19 of 21 (90%). Conversely, other taxa recovered in Bohaiornithidae tend to only rarely resolve near *Bohaiornis*. *Linyiornis* is recovered in a clade with *Bohaiornis* in 1 of 9 (11%) studies, *Eoenantiornis* in 3 of 21 (14%), and *Fortunguavis* in 4 of 19 (21%). *Gretcheniao* and *Musivavis*, both described as bohaiornithid-like but not bohaiornithid, are never recovered close to *Bohaiornis* (closest in [21] where *Gretcheniao* is in a clade sister to the six original bohaiornithids plus BMNHC-Ph1204). *Beiguornis* has only been included in one study [22] where it is recovered as a bohaiornithid, so it is tentatively included in the group pending future studies.

Phylogenetic Tree Topology

Extant bird phylogenetic trees in this study are trimmed versions of the maximum clade

credibility supertree used in [111]. This supertree follows [112], a tree time-scaled using

Bayesian uncorrelated relaxed molecular clock data from 15 genes in 6,663 avian species

constrained by seven fossil taxa. These data were mapped onto the backbone of [113] which used

Bayesian uncorrelated relaxed molecular clock data from 259 genes in 200 species constrained

by 19 fossil taxa. The tree files and R code to merge them were taken from the supplement of

[111].

All grafted bohaiornithid branch lengths were scaled linearly so that the total length of

the avian portion of the tree was equal to 94 Ma following the estimate of [97]. The

Ornithothoraces node was placed at 145 Ma after Bayesian morphological clock analysis of two

independent character sets [95, 96]. For the bohaiornithid portion of the tree, a topology

recovered by two recent analyses [85, 88] was used where *Bohaiornis* and *Parabohaiornis* are

sister taxa and the remaining bohaiornithids are in a basal polytomy. The position of *Beiguornis*

is considered uncertain as the phylogeny in its description [22] differs from this topology, but it

was not included in any analyses where phylogeny was incorporated.

Bohaiornithid Fossil Dating

Shenqiornis is referred to the Qiaotou Formation [82]. This has been correlated to the

Dawangzhangzi bed of the Yixian Formation [114], which is within the "Upper Undivided

Yixian Formation" of [115] dated to 124 Ma. All other bohaiornithids are referred to the

Jiufotang Formation. Most of these [12, 21] are from the Xiaotaizi/Lamadong locality dated to

119 Ma [116]. *Longusunguis* IVPP V18693 from the Lingyuan locality [52], *Zhouornis* BMNHC

Ph756 from the Xiaoyugou locality [56], and *Zhouornis* CNUVB 903 of uncertain provenance

[55] are referred to 121 Ma as the median age of the Jiufotang Formation [116].

Taxonomic Status of BMNHC-Ph1204

BMNHC-Ph1204 was previously identified to Bohaiornithidae indet. [21], though we herein

refine the diagnosis to cf. *Sulcavis*. As noted in [21], the large number of dentary teeth in

BMNHC-Ph1204 is only seen in *Sulcavis* among known bohaiornithids, and the maxillary

process of the nasal and interclavicular angle of the specimen are also consistent with those seen

in *Sulcavis*. The final defining feature of BMNHC-Ph1204 noted by [21] is the lack of a fenestra

in the maxilla, and the maxilla is poorly preserved in the *Sulcavis* holotype with no fenestra

visible [15]. Most of the synapomorphies of *Sulcavis* are present in BMNHC-Ph1204: teeth with

a flat lingual margin creating a 'D-shaped' cross-section [21 p. 5]; a broad nasal with a short,

rostrally-directed maxillary process [21 p. 3]; the caudal-most process of the synsacrum

extending far caudally to the vertebra's articular surface [fig. 8 in 21]; a long and delicate

acromion process on the scapula [fig. 7 in 21]; a convex lateral margin of the coracoid [fig. 8 in

21]; and an alular claw larger than that on the major digit [fig. 9 in 21]. The only missing

synapomorphy is enamel with longitudinal ridges, autapomorphic for *Sulcavis* [15]. BMNHC-

Ph1204 is less mature than the *Sulcavis* holotype (Fig. S10; [15, 21]), and ontogenetic change in

enamel ornamentation has been noted previously in reptiles [117, 118], so it is possible that this

is an ontogenetic difference. The same can be said of the additional carpal element in BMNHC-

Ph1204, as carpometacarpal fusion is believed to occur relatively late in bohaiornithid ontogeny

[12, 21]. Therefore, with this uncertainty from ontogeny, we refer BMNHC-Ph1204 to cf.

Sulcavis. Should this specimen later be referred to its own taxon, the similarities above imply the

new taxon would be very closely-related to *Sulcavis*, and this specimen would still be the most

appropriate for filling in missing pieces of the skull of *Sulcavis*, as in Fig. 2.

Bohaiornithid Skull Reconstruction for MA and FEA

Reconstructions of bohaiornithid skulls are provided in Fig. 2. No complete bohaiornithid skulls

are currently known, so digital reconstruction was necessary to create workable biological

models [119]. Reconstructions of a given taxon used material from that taxon where possible, but

each reconstruction required at least one bone from another taxon. Ideally these replacement

bones would come from a taxon's closest relative, but the relationships within Bohaiornithidae

are unsTable 1side from the sister relationship of *Bohaiornis* and *Parabohaiornis*. For

reconstruction purposes, we treated *Shenqiornis* and *Sulcavis* as sister taxa due to them having

the next most frequent sister pairing among bohaiornithids [20, 56, 89] and the superficially

similar shallow angle of their rostra. *Longusunguis* and *Zhouornis* have the most unstable

placement within Bohaiornithidae, so supplemental parts were taken from taxa arbitrarily.

Fortunately these taxa each had two moderately complete skulls, so little material from other

taxa was needed.

Only *Bohaiornis* IVPP V17963 preserves the cranioventral region of the skull, so this

region in all reconstructions is based on this. This affects jaw-opening mechanical advantage

(OMA) of the upper jaw, and thus this functional index should be interpreted cautiously. We

interpret the bone in *Bohaiornis* LPM B00167 previously labelled as the postorbital [54] to be

the quadratojugal, as its shape is more similar to the short and weakly-forked quadratojugal

typical of enantiornithines [120] than the elongate postorbital of *Longusunguis* IVPP V18693

[52], *Parabohaiornis* IVPP V28398 [121], *Sulcavis* BMNHC Ph-805 [15], and cf. *Sulcavis*

BMNHC Ph-1204 [21]. The alternative is that this bone is in fact the postorbital, meaning

Bohaiornis (and possibly *Shenqiornis* and *Zhouornis*) lacked a complete postorbital bar.

The jugal and postorbital bones of *Longusunguis* are elongate and believed to be in

contact in life [52], which would increase the overall rigidity and stability of the skull. This is

also very likely in *Parabohaiornis* and *Sulcavis*, which preserve both a dorsally-curving jugal

and elongate postorbital (Fig. 2C,D). We reconstruct these bones as also in contact in other

bohaiornithids (Fig. 2), though this is less certain. Notably if we are incorrect in our

reinterpretation of a bone in LPM B00167 identified as the postorbital [54] as the quadratojugal,

then this much smaller element could not contact the jugal in life.

Bohaiornithid Ontogeny

Ontogeny is an important dimension to account for when discussing the morphology and ecology

of extinct animals, but the ontogeny of Bohaiornithidae (and enantiornithines in general) is

poorly understood. Hu and O'Connor [122] devised a useful "character stage" system for

tracking enantiornithine maturity based on the fusion of compound bones. Bohaiornithids are

aged using this framework in Fig. S10. Hu and O'Connor [122] combine the fusion of the

proximal metatarsus, the tibiotarsus, and carpometacarpus into a single Stage 3 (presumably due

to a lack of resolution). *Longusunguis* IVPP V18693 has a fused tibiotarsus and unfused

tarsometatarsus and carpometacarpus and *Sulcavis* BMNHC-Ph805 has a fused tibiotarsus and

tarsometatarsus but unfused carpometacarpus, indicating the tibiotarsus fused first and the

carpometacarpus last within this family. We thus subdivide the Stage 3 of [122] into 3a

(formation of the tibiotarsus), 3b (proximal fusion of the tarsometatarsus), and 3c (fusion of the

carpometacarpus) in bohaiornithids.

The above stages only approximate biological maturity, however. Histological data from

two bones has been used to suggest that *Zhouornis* CNUVB-0903 had achieved sexual maturity

but not skeletal maturity [55]; a recent study [123] has shown that histological maturity can vary

greatly within individual enantiornithines but upholds the diagnosis of CNUVB-0903 as an adult.

The indeterminate bohaiornithid CUGB P1202 is considered the least mature published member

of the family [13]. It was proposed as sexually mature due to its elongate tail feathers (with the

assumption they served a sexual display purpose) [13], but similar feathers in decidedly juvenile

enantiornithines (STM 34-7 [124] and 34-9, IVPP V15564 [125]) call this into doubt. By

correlating the histological maturity of [123] with the character stage system (Table S4), we see

that subadulthood begins at or before character stage 1 and full maturity occurs after completion

of character stage 3. We thus consider all bohaiornithids aside from the two mentioned above to

be mature subadults or young adults, as proposed previously [12, 21, 52].

Extant Sampling

Five additional taxa were added to the ungual TM dataset of [9] to increase the sample of

anisodactyl non-raptorial perching birds. Target taxa were identified from the literature as birds

with well-recorded perching behaviour and no record of grasping or manipulating prey with the

pes. This list of target taxa was sent in a formal loan request to Serina Brady of Carnegie

Museum of Natural History, who provided photographs of all available specimens fitting the

criteria.

Analytical Techniques

Body Mass

Extant bird masses were taken from the bird ecology database AVONET [41], and combined

with bird diet data from EltonTraits 1.0 [40] following Table S2. Granivores were not split into

husking and swallowing granivores; past work [8, 9] did not find the two subcategories to

meaningfully differ in body mass. Average masses of groups were compared via phylogenetic

HSD (pairwise function of R package RRPP [42] used to compare group means). Diagnostic cut-

off values between groups were found with the R package OptimalCutpoints [126] version 1.1-5

by optimizing the Youden index, with bootstrap estimation of 95% confidence intervals using the

R package boot [127] version 1.3-28.

Pedal Traditional Morphometrics, Mechanical Advantage, and Finite Element Analysis

The extant MA and FEA data in this study is unchanged from [9], and data collection from

fossils follows the same procedures as well. Pedal TM data are measured from scale photographs

where available, though the five new included taxa were photographed without scale. All digits

from a single individual were measured from a single photograph, so their relative scale is

preserved. Our TM data is based solely on angles and length ratios, so this lack of scale should

not be an issue. MA data were measured from the reconstructions in Fig. 2 in CorelDraw X8.

FEA models based on the reconstructions in Fig. 2 were created and solved within HyperWorks

2022 Student Edition (*HyperMesh* and *Optistruct*, Altair Engineering, Inc., USA). All analyses

of the data were performed in R version 4.1.2 [128], with scripts available from Mendeley Data

(doi 10.17632/7xtpbv27zh.1).

Skull Traditional Morphometrics

Clark et al. [10] recently used skull traditional morphometrics to great effect in both

differentiating extant bird diets and diagnosing the diet of longipterygid enantiornithines. Thus,

we incorporate their data and methodology here as well to expand the evidence supporting our

diet diagnoses. Measurements of skull and rostrum length for extant birds are unchanged from

[10]. Clark et al. [10] do not diagram their measurement landmarks, and the caudal landmark of

the rostral length as "the caudal margin of the lacrimal (i.e., the rostral margin of the orbit)" does

not specify a dorsoventral height. So, for all new rostrum length measurements of bohaiornithids,

we measured to the dorsoventral midpoint of the caudal margin of the ventral ramus of the

lacrimal.

Diet assignments of extant taxa were changed from those used by Clark et al. [10] to

those in Table S2 for consistency following information in EltonTraits 1.0 [40]. Of the 408

extant bird samples included in the analysis of [10], 61 were removed due to not meeting the

definition of any diet category in Table S2 (they were "in between" diets as defined here, often

with 50% of the diet from two sources). Thus the final skull morphometric dataset consisted of

347 samples across 298 species. We also use mass estimates for taxa from AVONET [41] rather

than those listed by Clark et al. [10], for consistency within this work.

Clark et al. [10] visualise (their figure 4) and generally interpret their skull

morphometrics in terms of proportional rostral length (i.e. rostrum length divided by skull

length) vs. $\log_{10}$ skull length. We have modified both of these factors in Fig. 7. Firstly, we $\log_{10}$-

transform both rostrum and skull length before taking their ratio, as this better normalises the

distribution of proportional rostral length. Actual distribution of the data along this axis is

minimally affected. Secondly, we use a relative skull length instead of absolute skull length.

Absolute skull length is essentially a size proxy; Fig. S4 shows skull length and body mass

creating similar distributions, with the position of diet groups maintained relative to one another

while spread more evenly along the x-axis when using body mass than absolute skull length.

This work already discusses body size as a diet proxy at length, so including it in morphometric

discussion would be redundant. Through $\log_{10}$-$\log_{10}$ regression, we found that skull length

generally increases with body mass with a scaling exponent of 2.75 (slightly negative allometry,

$R^2 = 0.63$) for the extant birds sampled. Thus we calculated relative skull length as:

$$relative\ skull\ length = \frac{(log_{10}(skull\ length))^{2.7482}}{log_{10}(body\ mass)}$$

Relative skull length approximates how much longer or shorter a given bird's skull is than the

expected length for a bird of the same body mass.

Ancestral State Reconstruction

The backbone for the enantiornithine tree topology used in ancestral state reconstruction follows

[85]. This tree contains 34 of the 56 species used in ancestral state reconstruction. Additional

taxa were grafted onto this backbone following [20-22, 88, 110, 129-132]. Some taxa have only

been included in one phylogenetic analysis, which placed said taxa within Bohaiornithidae,

Longipterygidae, or Pengornithidae (e.g. *Noguerornis* placed within Longipterygidae [129]).

These taxa all lack the synapomorphies used to formally define the relevant clade, and their

placements within them are likely artefacts of sharing a few characters while most data is

missing. We therefore maintained monophyly of the Bohaiornithidae, Longipterygidae, and

Pengornithidae by placing these ambiguous taxa as sister to the family they were placed within.

Time-scaling was performed using the R package paleotree [133]. All enantiornithine taxa were

placed at their earliest occurrence, with species divergence arbitrarily chosen to take 1,000 years.

The enantiornithine phylogeny was rooted at 144 Ma after [95, 96]. Ages of localities follow

[115, 116, 134-143], with precise notes in the data repository. If the horizon of a specimen was

in doubt, it was placed at the median age of the formation.

The diet of the common ancestor of Enantiornithes was reconstructed in two ways:

qualitatively, directly labelling diets onto taxa to reconstruct the diet of ancestral nodes; and

quantitatively, reconstructing ancestral body mass, jaw MA and functional indices, jaw FEA

MWAM strain and interval values, and pedal TM values, and predicting ancestral diet from these

reconstructed values. The qualitative set includes quantitative diet reconstructions here and in [8,

9], all qualitative hypotheses of diet in enantiornithines (Table S5), and all enantiornithines with

mass data (approximating enantiornithines known from substantial fossil material, as they need

to be somewhat complete for mass estimation; *Cuspirostrisornis*, *Dalingheornis*, *Gracilornis*,

Jibeinia, *Longchengornis*, *Microenantiornis*, and *Paraprotopteryx* were excluded due to never

being included in a phylogenetic analysis). The latter were coded with "Unknown" diet to

illustrate uncertainty in the reconstruction. As hypothesised diets are less precise than the diet

categories used elsewhere in this study, they were lumped together as herbivores (folivores,

frugivores, granivores, and nectarivores), invertivores, omnivores (generalists), or vertivores

(piscivores, tetrapod hunters, and scavengers). 14 of 30 (47%) of diets in this reconstruction are

supported by preserved meal evidence or quantitative diet proxies. A total of 56 taxa are included

in the qualitative ancestral state reconstruction.

Quantitative ancestral state reconstruction includes data from the present study and our

two other quantitative analyses of enantiornithines [8, 9], as well as pedal TM data for

Fortunguavis from [144] and mass data from [2, 26]. Only MWAM strain of FEA models was

reconstructed, not full intervals data, as we found the validity of reconstructing an individual

strain interval to be dubious. If pedal TM data was available for multiple specimens of a genus,

we favoured the most mature specimen to represent the genus. If mass data was available for

multiple specimens of a genus, we favoured the largest mass estimate as an assumption of it

representing the most mature sample. The large-toothed *Longipteryx* morphotype [8] was

assumed more mature than the small-toothed morphotype and used for all reconstructions. A

total of 44 taxa are used in ancestral state reconstruction of body mass, 13 in upper jaw MA, 9 in

lower jaw MA, 13 in FEA, and 16 in pedal TM.

Ancestral states were estimated using the fastAnc() function in Phytools 0.7-90 [145]. As

polytomies are present in the enantiornithine tree used, ancestral states calculated 10,000 times

with random resolutions of the polytomies and averaged. As some diets were uncertain (e.g.

Zhouornis may be either herbivorous or a generalist), and fastAnc() cannot accept probabilistic

qualitative states for terminal taxa, we coded diets quantitatively as percent likelihood for each

diet.

Acknowledgements

The authors would like to thank Serina Brady, Chase Mendenhall, Stephen Rogers (Carnegie

Museum of Natural History), Andrew Kratter, and David Steadman (Florida Museum of Natural

History) for their assistance and expertise in selecting physical specimens for this study. We

would also like to thank Kathryn C. Gamble DVM, MS, Dipl ACZM, Dip ECZM (ZHM),

Veterinary Advisor Coraciiformes/Bucerotiformes TAG for providing coraciiform radiographs

used in this study. We also thank Gavin Thomas and Ryan Felice for their insight in constructing

the consensus phylogeny of extant birds. CVM is supported by a Postgraduate Scholarship from

The University of Hong Kong (HKU PGS). MP is supported by the Research Grant Council of

Hong Kong's General Research Fund (17120920; 17103315; 17105221) and the School of Life

Sciences at The Chinese University of Hong Kong. X.W. is supported by the Taishan Scholars

Program of Shandong Province (Ts20190954).

Declaration of Interests

882 The authors declare no competing interests.

References

1. Felice, R.N., et al., *Dietary niche and the evolution of cranial morphology in birds.* Proceedings of the Royal Society B: Biological Sciences, 2019. **286**(1897): p. 20182677.
2. Miller, C.V. and M. Pittman, *The diet of early birds based on modern and fossil evidence and a new framework for its reconstruction.* Biological Reviews, 2021. **96**(5): p. 2058-2112.
3. O'Connor, J.K., *The trophic habits of early birds.* Palaeogeography, Palaeoclimatology, Palaeoecology, 2019. **513**: p. 178-195.
4. O'Connor, J.K. and Z. Zhou, *The evolution of the modern avian digestive system: insights from paravian fossils from the Yanliao and Jehol biotas.* Palaeontology, 2019. **63**: p. 13-27.
5. Miller, C.V., et al., *Disassociated rhamphotheca of fossil bird Confuciusornis informs early beak reconstruction, stress regime, and developmental patterns.* Communications Biology, 2020. **3**: p. 519.
6. Hu, H., et al., *Earliest evidence for fruit consumption and potential seed dispersal by birds.* eLife, 2022. **11**: p. e74751.
7. Marugán-Lobón, J. and L.M. Chiappe, *Ontogenetic niche shifts in the Mesozoic bird Confuciusornis sanctus.* Current Biology, 2022. **32**(7): p. 1629-1634.e2.
8. Miller, C.V., et al., *Diet of Mesozoic toothed birds (Longipterygidae) inferred from quantitative analysis of extant avian diet proxies.* BMC Biology, 2022. **20**: p. 101.
9. Miller, C.V., et al., *Quantitative investigation of Mesozoic toothed birds (Pengornithidae) diet reveals earliest evidence of macrocarnivory in birds.* iScience, 2023. **26**(3): p. 106211.
10. Clark, A.D., et al., *Reconstructing the dietary habits and trophic positions of the Longipterygidae (Aves: Enantiornithes) using neontological and comparative morphological methods.* PeerJ, 2023: p. 15139.
11. Pittman, M., et al., *The fossil record of Mesozoic and Paleocene pennaraptorans*, in *Pennaraptoran Theropod Dinosaurs: Past Progress and New Frontiers*, M. Pittman and X. Xu, Editors. 2020, Bulletin of the American Museum of Natural History. p. 37-96.
12. Wang, M., et al., *A new diverse enantiornithine family (Bohaiornithidae fam. nov.) from the Lower Cretaceous of China with information from two new species.* Vertebrata PalAsiatica, 2014. **52**: p. 31-76.
13. Peteya, J.A., et al., *The plumage and colouration of an enantiornithine bird from the Early Cretaceous of China.* Palaeontology, 2017. **60**(1): p. 55-71.
14. O'Connor, J.K. and L.M. Chiappe, *A revision of enantiornithine (Aves: Ornithothoraces) skull morphology.* Journal of Systematic Palaeontology, 2011. **9**(1): p. 135-157.
15. O'Connor, J.K., et al., *A new enantiornithine from the Yixian Formation with the first recognized avian enamel specialization.* Journal of Vertebrate Paleontology, 2013. **33**(1): p. 1-12.
16. Chiappe, L.M. and Q. Meng, *Birds of Stone: Chinese Avian Fossils from the Age of Dinosaurs.* Vol. 9. 2016, Baltimore, MD, USA: John Hopkins University Press.
17. Zhou, Y.C., et al., *Evolution of tooth crown shape in Mesozoic birds, and its adaptive significance with respect to diet.* Palaeoworld, 2021. **30**(4): p. 724-736.
18. Li, Z.H., et al., *A new specimen of large-bodied basal enantiornithine Bohaiornis from the Early Cretaceous of China and the inference of feeding ecology in Mesozoic birds.* Journal of Paleontology, 2014. **88**(1): p. 99-108.

929   19.   Chiappe, L.M., et al., *New Bohaiornis-like bird from the Early Cretaceous of China:*
930   *enantiornithine interrelationships and flight performance.* PeerJ, 2019. **7**: p. e7846.
931   20.   Wang, X., et al., *A new bohaiornithid-like bird from the Lower Cretaceous of China fills*
932   *a gap in enantiornithine disparity.* Journal of Paleontology, 2022. **96**(4): p. 961 - 976.
933   21.   Liu, D., et al., *Cranial and dental morphology in a bohaiornithid enantiornithine with*
934   *information on its tooth replacement pattern.* Cretaceous Research, 2022: p. 105021.
935   22.   Wang, X., et al., *The first enantiornithine bird from the Lower Cretaceous Longjiang*
936   *Formation in the Great Khingan Range of Inner Mongolia.* Acta Geologica Sinica, 2022.
937   **96**(2): p. 337-348.
938   23.   Navalón, G., et al., *The evolutionary relationship among beak shape, mechanical*
939   *advantage, and feeding ecology in modern birds.* Evolution, 2018. **73**(3): p. 422-435.
940   24.   Pigot, A.L., et al., *Macroevolutionary convergence connects morphological form to*
941   *ecological function in birds.* Nature Ecology & Evolution, 2020. **4**: p. 230-239.
942   25.   Natale, R. and G.J. Slater, *The effects of foraging ecology and allometry on avian skull*
943   *shape vary across levels of phylogeny.* The American Naturalist, 2022.
944   26.   Serrano, F.J., P. Palmqvist, and J.L. Sanz, *Multivariate analysis of neognath skeletal*
945   *measurements: implications for body mass estimation in Mesozoic birds.* Zoological
946   Journal of the Linnean Society, 2015. **173**(4): p. 929-955.
947   27.   Stayton, C.T., *Testing hypotheses of convergence with multivariate data: morphological*
948   *and functional convergence among herbivorous lizards.* Evolution, 2006. **60**(4): p. 824-
949   841.
950   28.   Corbin, C.E., L.K. Lowenberger, and B.L. Gray, *Linkage and trade☐off in trophic*
951   *morphology and behavioural performance of birds.* Functional ecology, 2015. **29**(6): p.
952   808-815.
953   29.   Adams, N.F., et al., *Functional tests of the competitive exclusion hypothesis for*
954   *multituberculate extinction.* Royal Society Open Science, 2019. **6**(3): p. 181536.
955   30.   Ma, W., et al., *Functional morphology of the oviraptorosaur and scansoriopterygid skull*,
956   in *Pennaraptoran Theropod Dinosaurs: Past Progress and New Frontiers*, M. Pittman
957   and X. Xu, Editors. 2020, Bulletin of the American Museum of Natural History. p. 229-
958   249.
959   31.   Bathe, K.-J., *Finite Element Procedures*. 2 ed. 2014, Watertown, MA, USA: Bathe,
960   Klaus-Jürgen.
961   32.   Dumont, E.R., I.R. Grosse, and G.J. Slater, *Requirements for comparing the performance*
962   *of finite element models of biological structures.* Journal of Theoretical Biology, 2009.
963   **256**(1): p. 96-103.
964   33.   Bright, J.A., *A review of paleontological finite element models and their validity.* Journal
965   of Paleontology, 2014. **88**(4): p. 760-769.
966   34.   Marcus, L.F., *Traditional Morphometrics*, in *Proceedings of the Michigan*
967   *morphometrics workshop*, F.J. Rohlf and F.L. Bookstein, Editors. 1990, University of
968   Michigan Museum of Zoology: Ann Arbor, MI, USA. p. 77-122.
969   35.   Csermely, D. and O. Rossi, *Bird claws and bird of prey talons: where is the difference?*
970   Italian Journal of Zoology, 2006. **73**(01): p. 43-53.
971   36.   Fowler, D.W., E.A. Freedman, and J.B. Scannella, *Predatory functional morphology in*
972   *raptors: interdigital variation in talon size is related to prey restraint and immobilisation*
973   *technique.* PLOS ONE, 2009. **4**(11): p. e7999.

37.   Csermely, D., O. Rossi, and F. Nasi, *Comparison of claw geometrical characteristics*
among birds of prey and non-raptorial birds. Italian Journal of Zoology, 2012. **79**(3): p.
410-433.
38.   Cobb, S.E. and W.I. Sellers, *Inferring lifestyle for Aves and Theropoda: a model based*
on curvatures of extant avian ungual bones. PLOS ONE, 2020. **15**(2): p. e0211173.
39.   Einoder, L.D. and A.M.M. Richardson, *Aspects of the hindlimb morphology of some*
Australian birds of prey: a comparative and quantitative study. The Auk: Ornithological
Advances, 2007. **124**(3): p. 773-788.
40.   Wilman, H., et al., *EltonTraits 1.0: Species☐level foraging attributes of the world's birds*
and mammals. Ecology, 2014. **95**(7): p. 2027.
41.   Tobias, J.A., et al., *AVONET: morphological, ecological and geographical data for all*
birds. Ecology Letters, 2022. **25**(3): p. 581-597.
42.   Collyer, M.L. and D.C. Adams, *RRPP: An r package for fitting linear models to*
high☐dimensional data using residual randomization. Methods in Ecology and
Evolution, 2018. **9**: p. 1772-1779.
43.   Fluss, R., D. Faraggi, and B. Reiser, *Estimation of the Youden Index and its associated*
cutoff point. Biometrical Journal, 2005. **47**(4): p. 458-472.
44.   Marcé-Nogué, J., et al., *Accounting for differences in element size and homogeneity when*
comparing finite element models: armadillos as a case study. Palaeontologia Electronica,
2016. **19**(2): p. 2T.
45.   Marcé-Nogué, J., et al., *The intervals method: a new approach to analyse finite element*
outputs using multivariate statistics. PeerJ, 2017. **5**: p. e3793.
46.   Thomson, V.K., et al., *Carrion preference in Australian coastal raptors: effects of*
urbanisation on scavenging. The Sunbird, 2016. **46**(1): p. 16-28.
47.   Mazumdar, S., D. Ghose, and G.K. Saha, *Offal dumping sites influence the relative*
abundance and roosting site selection of Black Kites (Milvus migrans govinda) in urban
landscape: a study from Kolkata metropolis, India. Environmental Monitoring and
Assessment, 2018. **190**(1): p. 1-10.
48.   Ruxton, G.D. and D.C. Houston, *Obligate vertebrate scavengers must be large soaring*
fliers. Journal of Theoretical Biology, 2004. **228**(3): p. 431-436.
49.   Bright, J.A., et al., *The multifactorial nature of beak and skull shape evolution in parrots*
and cockatoos (Psittaciformes). BMC Evolutionary Biology, 2019. **19**(1): p. 104.
50.   Lautenschlager, S., et al., *Edentulism, beaks, and biomechanical innovations in the*
evolution of theropod dinosaurs. Proceedings of the National Academy of Sciences,
2013. **110**(51): p. 20657-20662.
51.   Fowler, D.W., et al., *The predatory ecology of Deinonychus and the origin of flapping in*
birds. PLOS ONE, 2011. **6**(12): p. e28964.
52.   Hu, H., et al., *New anatomical information on the bohaiornithid Longusunguis and the*
presence of a plesiomorphic diapsid skull in Enantiornithes. Journal of Systematic
Palaeontology, 2020. **18**(18): p. 1481-1495.
53.   Ward, A.B., P.D. Weigl, and R.M. Conroy, *Functional morphology of raptor hindlimbs:*
implications for resource partitioning. The Auk: Ornithological Advances, 2002. **119**(4):
p. 1052-1063.
54.   Hu, D., et al., *A new enantiornithine bird from the Lower Cretaceous of western*
Liaoning, China. Journal of Vertebrate Paleontology, 2011. **31**(1): p. 154-161.

55.   Zhang, Z.H., et al., *A large bird from the Early Cretaceous of China: new information on*
the skull of enantiornithines. Journal of Vertebrate Paleontology, 2013. **33**(5): p. 1176-
1189.
56.   Zhang, Y.G., et al., *New information on the anatomy of the Chinese Early Cretaceous*
Bohaiornithidae (Aves: Enantiornithes) from a subadult specimen of Zhouornis hani.
PeerJ, 2014. **2**: p. e407.
57.   Torres, C.R., M.A. Norell, and J.A. Clarke, *Bird neurocranial and body mass evolution*
across the end-Cretaceous mass extinction: The avian brain shape left other dinosaurs
behind. Science Advances, 2021. **7**(31): p. eabg7099.
58.   Chiappe, L.M., et al., *Fossil basicranium clarifies the origin of the avian central nervous*
system and inner ear. Proceedings of the Royal Society B: Biological Sciences, 2022.
289: p. 20221398.
59.   Dudley, R. and G.J. Vermeij, *Do the power requirements of flapping flight constrain*
folivory in flying animals? Functional Ecology, 1992. **6**: p. 101-104.
60.   Billerman, S.M., *Hoatzin (Opisthocomus hoazin), version 1.0*, in *Birds of the World*, S.M.
Billerman, et al., Editors. 2020, Cornell Lab of Ornithology: Ithaca, NY, USA.
61.   Brady, S., *Southern Screamer (Chauna torquata), version 1.0*, in *Birds of the World*, T.S.
Schulenberg, Editor. 2020, Cornell Lab of Ornithology: Ithaca, NY, USA.
62.   Winkler, D.W., S.M. Billerman, and I.J. Lovette, *Sandgrouse (Pteroclidae), version 1.0*,
in *Birds of the World*, S.M. Billerman, et al., Editors. 2020, Cornell Lab of Ornithology:
Ithaca, NY, USA.
63.   Gionfriddo, J.P. and L.B. Best, *Grit Use by Birds*, in *Current Ornithology*, V. Nolan,
E.D. Ketterson, and C.F. Thompson, Editors. 1999, Springer US: Boston, MA. p. 89-148.
64.   Cooper Jr, W.E. and L.J. Vitt, *Distribution, extent, and evolution of plant consumption by*
lizards. Journal of Zoology, 2002. **257**(4): p. 487-517.
65.   Melstrom, K.M., *The relationship between diet and tooth complexity in living*
dentigerous saurians. Journal of Morphology, 2017. **278**(4): p. 500-522.
66.   Li, Z., et al., *Ultramicrostructural reductions in teeth: implications for dietary transition*
from non-avian dinosaurs to birds. BMC Evolutionary Biology, 2020. **20**: p. 46.
67.   Herd, R.M. and T.J. Dawson, *Fiber digestion in the emu, Dromaius novaehollandiae, a*
large bird with a simple gut and high rates of passage. Physiological Zoology, 1984.
57(1): p. 70-84.
68.   Buchsbaum, R., J. Wilson, and I. Valiela, *Digestibility of plant constitutents by Canada*
Geese and Atlantic Brant. Ecology, 1986. **67**(2): p. 386-393.
69.   Ankney, C.D. and A.D. Afton, *Bioenergetics of breeding Northern Shovelers: diet,*
nutrient reserves, clutch size, and incubation. The Condor, 1988. **90**(2): p. 459-472.
70.   Kehoe, F.P., C.D. Ankney, and R.T. Alisauskas, *Effects of dietary fiber and diet diversity*
on digestive organs of captive Mallards (Anas platyrhynchos). Canadian Journal of
Zoology, 1988. **66**(7): p. 1597-1602.
71.   Gasaway, W.C., *Seasonal variation in diet, volatile fatty acid production and size of the*
cecum of rock ptarmigan. Comparative Biochemistry and Physiology Part A: Physiology,
1976. **53**(1): p. 109-114.
72.   Moss, R., *Gut size and the digestion of fibrous diets by tetraonid birds.* Journal of
Experimental Zoology, 1989. **252**(S3): p. 61-65.
73.   Remington, T.E., *Why do grouse have ceca? A test of the fiber digestion theory.* Journal
of Experimental Zoology, 1989. **252**(S3): p. 87-94.

74. Mayr, G., et al., *Reanalysis of putative ovarian follicles suggests that Early Cretaceous*
birds were feeding not breeding. Scientific Reports, 2020. **10**: p. 19035.
75. O'Connor, J.K., et al., *Ovarian follicles shed new light on dinosaur reproduction during*
the transition towards birds. National Science Review, 2014. **1**(1): p. 15-17.
76. Zheng, X., et al., *Preservation of ovarian follicles reveals early evolution of avian*
reproductive behaviour. Nature, 2013. **495**: p. 507-511.
77. Wang, Y., et al., *A new Jehol enantiornithine bird with three-dimensional preservation*
and ovarian follicles. Journal of Vertebrate Paleontology, 2016. **36**(2): p. e1054496.
78. Marshall, A.J. and R.W. Wrangham, *Evolutionary consequences of fallback foods.*
International Journal of Primatology, 2007. **28**(6): p. 1219-1235.
79. Xu, X., et al., *Study on the Jehol Biota: recent advances and future prospects.* Science
China Earth Sciences, 2020. **63**: p. 757-773.
80. Mazur, K.M. and P.C. James, *Barred Owl (Strix varia), version 1.1*, in *Birds of the*
World, A.F. Poole and F.B. Gill, Editors. 2021, Cornell Lab of Ornithology: Ithaca, NY,
USA.
81. Button, D.J. and L.E. Zanno, *Repeated evolution of divergent modes of herbivory in non-*
avian dinosaurs. Current Biology, 2020. **30**(1): p. 158-168.
82. Wang, X.R., et al., *New Species of Enantiornithes (Aves: Ornithothoraces) from the*
Qiaotou Formation in Northern Hebei, China. Acta Geologica Sinica (English Edition),
2010. **84**(2): p. 247-256.
83. Liu, S., et al., *Investigating possible gastroliths in a referred specimen of Bohaiornis guoi*
(Aves: Enantiornithes). Frontiers in Earth Science, 2021. **9**: p. 62.
84. Soons, J., et al., *Mechanical stress, fracture risk and beak evolution in Darwin's ground*
finches (Geospiza). Philosophical Transactions of the Royal Society B: Biological
Sciences, 2010. **365**(1543): p. 1093-1098.
85. Wang, M., et al., *An Early Cretaceous enantiornithine bird with a pintail.* Current
Biology, 2021. **31**(21): p. 4845-4852.e2.
86. Miller, C.V., et al., *Diet of Mesozoic toothed birds (Longipterygidae) inferred from*
quantitative analysis of extant avian diet proxies. Mendeley Data, 2022.
87. Miller, C.V., et al., *Quantitative investigation of Mesozoic toothed birds (Pengornithidae)*
diet reveals earliest evidence of macrocarnivory in birds. Mendeley Data, 2023.
88. Li, Z., et al., *Novel evolution of a hyper□elongated tongue in a Cretaceous*
enantiornithine from China and the evolution of the hyolingual apparatus and feeding in
birds. Journal of Anatomy, 2021. **240**(4): p. 627-638.
89. Wang, M., J.K. O'Connor, and Z.H. Zhou, *A new robust enantiornithine bird from the*
Lower Cretaceous of China with scansorial adaptations. Journal of Vertebrate
Paleontology, 2014. **34**(3): p. 657-671.
90. Matsukawa, M., et al., *The Early Cretaceous terrestrial ecosystems of the Jehol Biota*
based on food-web and energy-flow models. Biological Journal of the Linnean Society,
2014. **113**(3): p. 836-853.
91. O'Connor, J., et al., *Microraptor with ingested lizard suggests non-specialized digestive*
function. Current Biology, 2019. **29**(14): p. 2423-2429. e2.
92. O'Connor, J., Z. Zhou, and X. Xu, *Additional specimen of Microraptor provides unique*
evidence of dinosaurs preying on birds. Proceedings of the National Academy of
Sciences, 2011. **108**(49): p. 19662-19665.

93. Chiappe, L.M., *Enantiornithine (Aves) tarsometatarsi from the Cretaceous Lecho*
Formation of northwestern Argentina. American Museum novitates, 1993. **3083**.
94. Chiappe, L.M. and J.O. Calvo, *Neuquenornis volans, a new Late Cretaceous bird*
(Enantiornithes: Avisauridae) from Patagonia, Argentina. Journal of Vertebrate
Paleontology, 1994. **14**(2): p. 230-246.
95. Cau, A., *The assembly of the avian body plan: a 160-million-year long process.*
Bollettino della Società Paleontologica Italiana, 2018. **57**(1): p. 1-25.
96. Wang, M., et al., *The patterns and modes of the evolution of disparity in Mesozoic birds.*
Proceedings of the Royal Society B, 2021. **288**(1944): p. 20203105.
97. Kuhl, H., et al., *An unbiased molecular approach using 3'-UTRs resolves the avian*
family-level tree of life. Molecular Biology and Evolution, 2020. **38**(1): p. 108-127.
98. Wang, S., et al., *Tooth vs beak: the evolutionary developmental control of the avian*
feeding apparatus, in *Pennaraptoran Theropod Dinosaurs: Past Progress and New*
Frontiers, M. Pittman and X. Xu, Editors. 2020, Bulletin of the American Museum of
Natural History. p. 205-228.
99. Bhullar, B.A.S., et al., *How to make a bird skull: major transitions in the evolution of the*
avian cranium, paedomorphosis, and the beak as a surrogate hand. Integrative and
Comparative Biology, 2016. **56**(3): p. 389-403.
100. Brocklehurst, N. and D.J. Field, *Macroevolutionary dynamics of dentition in Mesozoic*
birds reveal no long-term selection towards tooth loss. iScience, 2021. **24**(3): p. 102243.
101. Field, D.J., et al., *Early evolution of modern birds structured by global forest collapse at*
the end-Cretaceous mass extinction. Current Biology, 2018. **28**(11): p. 1825-1831.
102. Mayr, G., *Ornithothoraces and the Origin of Sustained Flapping Flight Capabilities*, in
Avian Evolution: The Fossil Record of Birds and its Paleobiological Significance, M.J.
Benton, Editor. 2016, John Wiley & Sons: Oxford, UK. p. 50-55.
103. Flaherty, E.A., M. Ben-David, and W.P. Smith, *Quadrupedal locomotor performance in*
two species of arboreal squirrels: predicting energy savings of gliding. Journal of
Comparative Physiology B, 2010. **180**: p. 1067-1078.
104. Dodge, S., et al., *Environmental drivers of variability in the movement ecology of turkey*
vultures (Cathartes aura) in North and South America. Philosophical Transactions of the
Royal Society B: Biological Sciences, 2014. **369**: p. 20130195.
105. Billerman, S.M., et al. *Birds of the World*. 2023 https://birdsoftheworld.org/bow/home.
106. Van Gestel, W. and J. Jansen. *Skullsite Bird Skull Repository*. 2020 http://skullsite.com/.
107. Pittman, M., et al., *Pennaraptoran systematics*, in *Pennaraptoran Theropod Dinosaurs:*
Past Progress and New Frontiers, M. Pittman and X. Xu, Editors. 2020, Bulletin of the
American Museum of Natural History. p. 7-36.
108. Wang, M., H. Hu, and Z. Li, *A new small enantiornithine bird from the Jehol Biota, with*
implications for early evolution of avian skull morphology. Journal of Systematic
Palaeontology, 2015. **14**(6): p. 481-497.
109. Wang, M. and D. Liu, *Taxonomical reappraisal of Cathayornithidae (Aves:*
Enantiornithes). Journal of Systematic Palaeontology, 2016. **14**(1): p. 29-47.
110. O'Connor, P.M., et al., *Late Cretaceous bird from Madagascar reveals unique*
development of beaks. Nature, 2020. **588**: p. 272–276.
111. Cooney, C.R., et al., *Mega-evolutionary dynamics of the adaptive radiation of birds.*
Nature, 2017. **542**: p. 344-347.

112. Jetz, W., et al., *The global diversity of birds in space and time.* Nature, 2012. **491**: p. 444-448.

113. Prum, R.O., et al., *A comprehensive phylogeny of birds (Aves) using targeted next-generation DNA sequencing.* Nature, 2015. **526**: p. 569-573.

114. Jin, F., et al., *On the horizon of Protopteryx and the early vertebrate fossil assemblages of the Jehol Biota.* Chinese Science Bulletin, 2008. **53**(18): p. 2820-2827.

115. Zhong, Y., et al., *High-precision geochronological constraints on the duration of 'Dinosaur Pompeii' and the Yixian Formation.* National Science Review, 2021. **8**(6): p. nwab063.

116. Yu, Z., et al., *New geochronological constraints for the Lower Cretaceous Jiufotang Formation in Jianchang Basin, NE China, and their implications for the late Jehol Biota.* Palaeogeography, Palaeoclimatology, Palaeoecology, 2021. **583**: p. 110657.

117. Thies, D. and A. Broschinski, *Teeth of a small durophagous crocodile from the Late Jurassic (Kimmeridgian) of North Germany.* Geologische Beiträge Hannover, 2001. **2**: p. 65-70.

118. Street, H., A. LeBlanc, and M. Caldwell, *A histological investigation of dental crown characters used in mosasaur phylogenetic analyses.* Vertebrate Anatomy Morphology Palaeontology, 2021. **9**(1): p. 83-94.

119. Lautenschlager, S., *Reconstructing the past: methods and techniques for the digital restoration of fossils.* Royal Society Open Science, 2016. **3**(10): p. 160342.

120. Wang, M. and H. Hu, *A comparative morphological study of the jugal and quadratojugal in early birds and their dinosaurian relatives.* The Anatomical Record: Advances in Integrative Anatomy and Evolutionary Biology, 2017. **300**(1): p. 62-75.

121. Wang, M., *A new specimen of Parabohaiornis martini (Avialae: Enantiornithes) sheds light on early avian skull evolution.* Vertebrata Palasiatica, 2023.

122. Hu, H. and J.K. O'Connor, *First species of Enantiornithes from Sihedang elucidates skeletal development in Early Cretaceous enantiornithines.* Journal of Systematic Palaeontology, 2017. **15**(11): p. 909-926.

123. Atterholt, J., et al., *Intraskeletal osteohistovariability reveals complex growth strategies in a Late Cretaceous enantiornithine.* Frontiers in Earth Science, 2021. **9**: p. 640220.

124. Zheng, X.T., et al., *Insight into the early evolution of the avian sternum from juvenile enantiornithines.* Nature Communications, 2012. **3**: p. 1116.

125. O'Connor, J.K., et al., *First report of immature feathers in juvenile enantiornithines from the Early Cretaceous Jehol avifauna.* Vertebrata PalAsiatica, 2020. **59**: p. 24-44.

126. López-Ratón, M., et al., *OptimalCutpoints: an R package for selecting optimal cutpoints in diagnostic tests.* Journal of Statistical Software, 2014. **61**(8).

127. Davison, A.C. and D.V. Hinkley, *Bootstrap Methods and their Application.* 1997, Cambridge, UK: Cambridge University Press.

128. R Core Team, *R: A language and environment for statistical computing.* 2020, R Foundation for Statistical Computing: Vienna, Austria.

129. O'Connor, J.K., *A systematic review of Enantiornithes (Aves: Ornithothoraces).* 2009, University of Southern California.

130. Li, L., et al., *A new enantiornithine bird from the Lower Cretaceous Jiufotang Formation in Jinzhou Area, Western Liaoning Province, China.* 地质学报（英文版）, 2012. **86**(5): p. 1039-1044.

131. Kurochkin, E.N., S. Chatterjee, and K.E. Mikhailov, *An embryonic enantiornithine bird*
and associated eggs from the Cretaceous of Mongolia. Paleontological Journal, 2013.
47(11): p. 1252-1269.
132. Atterholt, J., J.H. Hutchison, and J.K. O'Connor, *The most complete enantiornithine from*
North America and a phylogenetic analysis of the Avisauridae. PeerJ, 2018. **6**: p. e5910.
133. Bapst, D.W., *paleotree: an R package for paleontological and phylogenetic analyses of*
evolution. Methods in Ecology and Evolution, 2012. **3**(5): p. 803-807.
134. Chen, P., et al., *Geological ages of dinosaur-track-bearing formations in China.*
Cretaceous Research, 2006. **27**(1): p. 22-32.
135. Fregenal-Martínez, M., et al., *The stratigraphic record of the Late Jurassic-Early*
Cretaceous rifting in the Alto Tajo-Serranía de Cuenca region (Iberian Ranges, Spain).
Revista de la Sociedad Geológica de España, 2017. **30**(1): p. 113-142.
136. Lockley, M.G., et al., *Large theropod and small sauropod trackmakers from the Lower*
Cretaceous Jingchuan Formation, Inner Mongolia, China. Cretaceous Research, 2018.
92: p. 150-167.
137. Porfiri, J.D., et al., *A new megaraptoran theropod dinosaur from the Upper Cretaceous*
Bajo de la Carpa Formation of northwestern Patagonia. Cretaceous Research, 2018. **89**:
p. 302-319.
138. Chen, G., et al., *Astronomical cycles and variations in sediment accumulation rate of the*
terrestrial lower Cretaceous Xiagou Formation from the Jiuquan Basin, NW China.
Cretaceous Research, 2020. **109**: p. 104156.
139. Yang, S., et al., *The appearance and duration of the Jehol Biota: Constraint from SIMS*
U-Pb zircon dating for the Huajiying Formation in northern China. Proceedings of the
National Academy of Sciences, 2020. **117**(25): p. 14299-14305.
140. Napoli, J.G., et al., *A new dromaeosaurid (Dinosauria: Coelurosauria) from Khulsan,*
central Mongolia. American Museum Novitates, 2021. **2021**(3982): p. 1-47.
141. Galobart, À., J. Párraga, and A. Gil-Delgado, *The spread of fossil heritage: How to*
valorise the lithographic limestones of the La Pedrera de Meià and La Cabroa Quarries
in the Orígens Unesco Global Geopark (Southern Pyrenees, Catalonia). Geoheritage,
2022. **14**(1): p. 34.
142. Montano, D., et al., *Depositional age models in lacustrine systems from zircon and*
carbonate U-Pb geochronology. Sedimentology, 2022. **69**(6): p. 2507-2534.
143. Ramezani, J., et al., *Calibrating the zenith of dinosaur diversity in the Campanian of the*
Western Interior Basin by CA-ID-TIMS U–Pb geochronology. Scientific Reports, 2022.
12: p. 16026.
144. Pittman, M., et al., *Exceptional preservation and foot structure reveal ecological*
transitions and lifestyles of early theropod flyers. Nature Communications, 2022. **13**(1):
p. 7684.
145. Revell, L.J., *phytools: an R package for phylogenetic comparative biology (and other*
things). Methods in Ecology and Evolution, 2012. **3**(2): p. 217-223.
146. Sanz, J.L., et al., *An Early Cretaceous bird from Spain and its implications for the*
evolution of avian flight. Nature, 1996. **382**: p. 442-445.

Figures

 Figure 1

 Bohaiornithid skull reconstructions used for MA and FEA calculations in this study.

 Reconstructions are of *Bohaiornis* (A), *Longusunguis* (B), *Parabohaiornis* (C), *Shenqiornis* (D),

 Sulcavis (E), and *Zhouornis* (F). Different colours indicate elements taken from different

 individual specimens. All sclerotic rings are based on *Longipteryx* specimen BMNHC Ph-930B.

 See the Methods section for more details on reconstruction. Scale for each reconstruction is

 based on the individual which makes up the largest portion of the reconstruction.

Figure 2
Violin plots of bird mass by diet, arranged in order of ascending mean mass. Masses were tested
for significant differences via phylogenetic HSD. Diets marked with the same letter are not
significantly different from one another. Cut-off points between significantly different mass
groups (black lines, with 95% CIs as grey lines) were calculated by optimizing the Youden index
and plotted. Note that, unlike in other diet treatments herein, granivores are not separated into
husking and swallowing granivores. Diet abbreviations: Tetra Hunt, Tetrapod Hunter.

1258

1259

1260

Figure 3

Functional phylomorphospace of MA and functional index data, grouped by diet. Grey lines

indicate phylogenetic relationships. Data are presented with PCA (A) and FDA (B). See Fig. S2

for character weights and Table 2 for FDA posterior predictions. Diet abbreviations: GranivoreH,

Husking Granivore; GranivoreS, Swallowing Granivore; Tetra Hunt, Tetrapod Hunter. Fossil

taxon abbreviations: B, *Bohaiornis*; L, *Longusunguis*; P, *Parabohaiornis*; Sh, *Shenqiornis*; Su,

Sulcavis; Z, *Zhouornis*.

1268

Figure 4

Violin plots of MWAM strain of FEA models, organized by diet. Extant diets ascend in average

MWAM strain from left to right. MWAM strains were tested for significant differences via

phylogenetic HSD. Diets marked with the same letter are not significantly different from one

another. Diet abbreviations: GranivoreH, Husking Granivore; GranivoreS, Swallowing

Granivore; Tetra Hunt, Tetrapod Hunter. Fossil taxon abbreviations: B, *Bohaiornis*; L,

Longusunguis; P, *Parabohaiornis*; Sh, *Shenqiornis*; Su, *Sulcavis*; Z, *Zhouornis*.

1276

Figure 5
Phylogenetic strain-space of total maximum in-plane principal strain intervals for extant and
fossil bird lower jaw finite element models, grouped by diet. MWAM strain is mapped overtop
the data. Grey lines indicate phylogenetic relationships. Data are presented with PCA (A) and
FDA (B). In PCA (A) overall strain increases along PC1, and strain heterogeneity (i.e. lower
areas of intermediate strain) increases along PC2. In FDA (B), DA1 and DA2 have loadings of
various similar low-strain intervals, with high-strain intervals clustering near the origin. See
Table 3 for FDA posterior predictions. Diet abbreviations: GranivoreH, Husking Granivore;
GranivoreS, Swallowing Granivore; Tetra Hunt, Tetrapod Hunter. Fossil taxon abbreviations: B,
Bohaiornis; L, *Longusunguis*; P, *Parabohaiornis*; Sh, *Shenqiornis*; Su, *Sulcavis*; Z, *Zhouornis*.

1287

Figure 6
Phylomorphospace of extant and fossil bird claw shape from pedal TM, grouped by pedal
ecological category. Grey lines indicate phylogenetic relationships. Data are presented with PCA
(A), FDA (B), and pFDA (C). See Fig. l for character weights and Table 4 for FDA and pFDA
posterior predictions. Category abbreviations: Large Raptor, raptor taking prey which does not fit
in the foot; Small Raptor, raptor taking prey which can fit in the foot. Fossil taxon abbreviations:
B1, *Bohaiornis* LPM B00167; B2, *Bohaiornis* IVPP V17963, L, *Longusunguis* IVPP V18693;
P1, *Parabohaiornis* IVPP V18690; P2, *Parabohaiornis* IVPP V18691; Su, *Sulcavis* BMNH Ph-
805; Z1, *Zhouornis* CNUVB-0903, Z2, *Zhouornis* BMNHC Ph 756.

1297

Figure 7
Phylomorphospace of extant and fossil bird skull proportions, grouped by diet. Grey lines
indicate phylogenetic relationships. The data presented are modified from [10], see Fig. S4 for
data more directly comparable to that study. Diet abbreviation: Tetra Hunt, Tetrapod Hunter.
Fossil taxon abbreviations: B, *Bohaiornis*; L, *Longusunguis*; P, *Parabohaiornis*; Sh, *Shenqiornis*;
Su, *Sulcavis*; Z, *Zhouornis*.

1304

Figure 8
Ancestral state reconstruction of enantiornithine diet. Phylogeny is presented as not time-scaled
for node visibility. All enantiornithine taxa with dietary hypotheses are included, as well as
enantiornithines complete enough to create robust mass estimates [8, 9]. Taxa with bold names
have diet assigned based on preserved meals (*Eoalulavis* [146]) or quantitative diet proxies [8,
9]. Remaining diets assignments are based on qualitative morphology and depositional setting.
The diet of the common ancestor of Enantiornithes remains obscure, though many late-diverging
enantiornithines are recovered as ancestrally invertivorous.

Protopteryx
Parapengornis
Chiappeavis
Yuanchuavis
Pengornis
Eopengornis
Yuanjiawaornis
Monoenantiornis
Parabohaiornis
Bohaiornis
Zhouornis
Sulcavis
Shenqiornis
Longusunguis
Huoshanornis
Otogornis
Orienantius
Fortunguavis
Falcatakely
Eoenantiornis
Brevirostruavis
Linyiornis
Grabauornis
Parvavis
Junornis
Gretcheniao
Shengjingornis
Rapaxavis
Longirostravis
Shanweiniao
Longipteryx
Boluochia
Noguerornis
Dapingfangornis
Vescornis
Gobipteryx
Pterygornis
Musivavis
Dunhuangia
Piscivorenantiornis
Shangyang
Houornis
Eocathayornis
Halimornis
Elsornis
Gobipipus
Alethoalaornis
Mirarce
Gettyia
Avisaurus
Soroavisaurus
Neuquenornis
Concornis
Eoalulavis
Liaoningornis
Cathayornis

Pengornithidae

Bohaiornithidae

Longipterygidae

Gobipterygidae

Avisauridae

Diet

- Herbivory
- Invertivory
- Omnivory
- Vertivory
- Unknown

1313

Tables

Table 1

Masses for bohaiornithid taxa based on the regression equations of [26]. Most masses were previously reported in [2], though masses for the juvenile bohaiornithid CUGB P1202 [13], *Beiguornis khinganensis* MHGU-F307/8 [22], and cf. *Sulcavis* BMNHC-Ph1204 [21] are newly-calculated in this study from literature images.

Taxon	Specimen	Ontogenetic Stage	Mean Mass Estimate (g)	Min Mass Estimate (g)	Max Mass Estimate (g)
Bohaiornithidae indet.	CUGB P1202	1	113	91	135
Beiguornis khinganensis	MHGU-F307/8	2a	260	210	310
Bohaiornis guoi	IVPP V17963	3c	300	242	358
B. guoi	LPM B00167	2b	249	201	298
Longusunguis kurochkini	IVPP V17964	2b	171	137	204
L. kurochkini	IVPP V18693	3a	237	191	283
Parabohaiornis martini	IVPP V18691	2b	221	178	263
Parabohaiornis martini	IVPP V28398	3b	323	260	386
Shenqiornis mengi	DNHM D2950/1	2a	340	274	406
Sulcavis geeorum	BMNHC-Ph805	3b	333	268	397
cf. *Sulcavis*	BMNHC-Ph1204	2a	171	137	204
Zhouornis hani	BMNHC-Ph756	2a	253	204	303
Z. hani	CNUVB-903	3c	758	611	905

Table 2

Posterior probabilities predicting bohaiornithid diet by FDA from MA and functional indices of extant bird jaws. Values with green backgrounds are more likely, values with red backgrounds are less likely. All bohaiornithids have high affinity with generalists and low affinity with husking granivores, with other affinities varying by taxon. Diet abbreviations: GranivoreH, Husking Granivore; GranivoreS, Swallowing Granivore; Tetra Hunt, Tetrapod Hunter.

Taxon	Folivore	Frugivore	Generalist	GranivoreH	GranivoreS	Invertivore	Nectarivore	Piscivore	Scavenger	Tetra Hunt
Bohaiornis	9.99E-01	6.43E-06	2.92E-04	5.97E-13	2.15E-04	5.43E-06	1.62E-05	2.00E-07	1.74E-10	7.78E-08
Longusunguis	1.53E-02	1.81E-01	5.39E-01	5.48E-07	4.38E-03	1.57E-01	3.05E-02	6.91E-02	9.85E-05	3.95E-03
Parabohaiornis	2.44E-01	6.72E-04	2.34E-01	2.49E-07	3.64E-01	1.05E-02	2.42E-04	1.49E-02	1.32E-01	7.69E-05
Shenqiornis	7.32E-03	1.12E-02	6.45E-01	1.81E-06	2.00E-03	5.15E-02	8.12E-03	2.65E-01	9.50E-03	5.24E-04
Sulcavis	7.28E-02	1.58E-02	8.22E-01	5.46E-07	4.12E-03	2.43E-02	3.39E-03	5.68E-02	5.74E-04	6.25E-04
Zhouornis	3.20E-02	1.61E-01	3.11E-01	9.82E-07	2.30E-02	2.69E-01	5.42E-02	7.86E-02	9.64E-04	6.99E-02

Table 3

Posterior probabilities predicting bohaiornithid diet by FDA from FEA following the intervals method [45]. Values with green backgrounds are more likely, values with red backgrounds are less likely. Bohaiornithid affinities varying considerably between taxa, only universally not resembling tetrapod hunters. Diet abbreviations: GranivoreH, Husking Granivore; GranivoreS, Swallowing Granivore; Tetra Hunt, Tetrapod Hunter.

Taxon	Folivore	Frugivore	Generalist	GranivoreH	GranivoreS	Invertivore	Nectarivore	Piscivore	Scavenger	Tetra Hunt
Bohaiornis	3.44E-10	9.33E-12	1.73E-07	2.25E-23	1.00E+00	7.09E-07	4.67E-19	1.43E-10	8.82E-06	3.37E-19
Longusunguis	3.05E-01	1.15E-13	6.95E-01	1.56E-26	2.26E-15	3.58E-09	4.35E-26	1.51E-11	3.47E-29	1.07E-20
Parabohaiornis	8.49E-02	3.30E-15	2.08E-13	3.50E-24	5.52E-01	1.18E-04	4.67E-04	3.63E-01	3.12E-13	1.13E-06
Shenqiornis	2.28E-08	1.45E-23	9.39E-09	4.31E-11	2.28E-13	2.47E-05	3.61E-02	9.58E-01	6.11E-03	2.81E-20
Sulcavis	6.82E-06	1.91E-16	1.19E-03	9.94E-01	5.99E-18	5.65E-09	4.71E-03	1.22E-07	4.68E-35	1.55E-22
Zhouornis	5.69E-50	1.00E+00	5.55E-20	4.89E-13	4.06E-12	2.08E-18	8.99E-13	4.74E-20	1.13E-53	2.52E-17

Table 4

Posterior probabilities predicting bohaiornithid pedal ecology by FDA and pFDA from TM of extant bird claws. Values with green backgrounds are more likely, values with red backgrounds are less likely. Bohaiornithid affinities vary considerably by taxon and between FDA and pFDA. Diet abbreviations: GranivoreH, Husking Granivore; GranivoreS, Swallowing Granivore; Tetra Hunt, Tetrapod Hunter. Category abbreviations: Large Raptor, raptor taking prey which does not fit in the foot; Small Raptor, raptor taking prey which can fit in the foot.

Test	Taxon	Ground	Perch	Large Raptor	Small Raptor	Shrike
FDA	*Bohaiornis* LPM B00167	2.45E-02	4.03E-01	2.69E-01	3.03E-01	4.60E-04
	Bohaiornis IVPP V17963	1.92E-02	2.52E-01	2.31E-04	7.23E-01	5.78E-03
	Longusunguis	3.65E-02	4.93E-01	3.76E-01	9.45E-02	4.42E-07
	Parabohaiornis IVPP V18690	2.94E-05	8.57E-01	1.49E-03	5.57E-02	8.63E-02
	Parabohaiornis IVPP V18691	2.41E-05	9.66E-01	2.66E-02	4.41E-03	2.85E-03
	Sulcavis	2.03E-01	1.80E-02	2.17E-01	5.62E-01	2.93E-09
	Zhouornis CNUVB-0903	1.22E-01	3.62E-01	6.10E-02	4.07E-01	4.83E-02
	Zhouornis BMNHC Ph 756	5.46E-01	1.41E-02	1.72E-02	4.22E-01	1.33E-04
pFDA	*Bohaiornis* LPM B00167	0.00E+00	6.09E-02	1.29E-01	8.10E-01	4.57E-05
	Bohaiornis IVPP V17963	0.00E+00	6.21E-07	4.58E-06	4.87E-12	1.00E+00
	Longusunguis	0.00E+00	2.66E-03	9.94E-01	2.82E-03	6.46E-04
	Parabohaiornis IVPP V18690	0.00E+00	1.98E-04	2.10E-01	7.90E-01	3.27E-12
	Parabohaiornis IVPP V18691	0.00E+00	2.68E-02	3.05E-01	6.68E-01	3.28E-07
	Sulcavis	0.00E+00	2.87E-04	3.92E-05	3.27E-05	1.00E+00
	Zhouornis CNUVB-0903	0.00E+00	4.94E-01	2.84E-01	2.19E-01	3.00E-03
	Zhouornis BMNHC Ph 756	0.00E+00	3.10E-01	5.23E-03	3.96E-03	6.80E-01

Table 5

Quantitative ancestral states of the common ancestor of Enantiornithines. Mean values are the average of 10,000 random tree permutations where polytomies are randomly resolved. 95% confidence intervals are given as the 2.5% and 97.5% quantiles of the permutations. As these values are based on multiple possible topologies of the enantiornithine tree, we consider them more valid than those visualised in Figs. 12–16.

Line of Evidence	Trait	Mean	95% CI Low	95% CI High
Mass	Body Mass (g)	134	119	159
MA	AMAup	0.13	0.12	0.13
	PMAup	0.18	0.17	0.19
	OMAup	0.20	0.19	0.21
	AOup	0.10	0.09	0.11
	MCHup	0.32	0.32	0.33
	ACHup	0.19	0.19	0.20
	AMAlow	0.20	0.19	0.21
	PMAlow	0.31	0.29	0.32
	OMAlow	0.09	0.08	0.10
	AOlow	0.09	0.06	0.11
	MMHlow	0.11	0.11	0.12
	AMHlow	0.07	0.06	0.07
FEA	MWAM Strain (με)	231	221	236
pedal TM	DI/DIII Ratio	0.76	0.76	0.76
	DII/DIII Ratio	0.85	0.85	0.85
	DIV/DIII Ratio	0.60	0.60	0.60
	DI Angle (°)	120	120	120
	DII Angle (°)	98	98	98
	DIII Angle (°)	90	90	90
	DIV Angle (°)	71	71	71

Table 6

Posterior probability of diet in the common ancestor of Enantiornithes based on MA and functional indices. Generalist feeding is the most likely, followed by piscivory and invertivory. Diet abbreviations: GranivoreH, Husking Granivore; GranivoreS, Swallowing Granivore; Tetra Hunt, Tetrapod Hunter.

Folivore	Frugivore	Generalist	GranivoreH	GranivoreS	Invertivore	Nectarivore	Piscivore	Scavenger	Tetra Hunt
4.82E-03	9.09E-03	6.98E-01	1.32E-07	1.42E-02	9.69E-02	3.96E-03	1.50E-01	2.16E-02	6.96E-04

Table 7

Summary table of interpretations of each line of evidence used herein. Body mass, MA, and FEA inform diet. Pedal TM informs the use of the pes or lack thereof in feeding. See relevant Discussion sections for additional details Bolded diets are agreed upon by all available diet proxies. Bolded pedal ecologies are either help discriminate diet (carnivory in *Longusunguis*) or are supported over other possibilities by diet information (non-raptorial perching in *Bohaiornis*). Given the uncertain application of skull TM to bohaiornithids, diets from this proxy are not bolded.

Line of Evidence	Taxon	Likely Diets/Ecologies	Unlikely Diets/Ecologies
Body Mass	*Beiguornis*	Folivore, Frugivore, Generalist, Piscivore, Tetrapod Hunter	Granivore, Invertivore, Nectarivore
	Bohaiornis	**Folivore**, Frugivore, Generalist, Piscivore, Tetrapod Hunter	Granivore, Invertivore, Nectarivore
	Longusunguis	Folivore, Frugivore, **Generalist**, **Piscivore**, Tetrapod Hunter	Granivore, Invertivore, Nectarivore
	Parabohaiornis	**Folivore**, Frugivore, **Generalist**, Piscivore, Tetrapod Hunter	Granivore, Invertivore, Nectarivore
	Shenqiornis	**Folivore**, Frugivore, **Generalist**, **Piscivore**, Tetrapod Hunter	Granivore, Invertivore, Nectarivore
	Sulcavis	**Folivore**, Frugivore, **Generalist**, **Piscivore**, Tetrapod Hunter	Granivore, Invertivore, Nectarivore
	Zhouornis	Folivore, **Frugivore**, **Generalist**, Piscivore, Scavenger, Tetrapod Hunter	Granivore, Invertivore, Nectarivore
Mechanical Advantage	*Bohaiornis*	**Folivore**, Generalist, Swallowing Granivore	Husking Granivore, Scavenger
	Longusunguis	Frugivore, **Generalist**, Invertivore, **Piscivore**	Husking Granivore, Scavenger

	Parabohaiornis	**Folivore**, **Generalist**, Scavenger, Swallowing Granivore	Husking Granivore, Tetrapod Hunter
	Shenqiornis	**Folivore**, Frugivore, **Generalist**, Invertivore, **Piscivore**	Husking Granivore, Tetrapod Hunter
	Sulcavis	**Folivore**, **Generalist**, Invertivore, **Piscivore**	Husking Granivore, Scavenger
	Zhouornis	**Frugivore**, **Generalist**, Invertivore, Piscivore	Husking Granivore, Scavenger
Finite Element Analysis	*Bohaiornis*	**Folivore**, Husking Granivore, Nectarivore	Swallowing Granivore, Invertivore, Piscivore, Scavenger, Tetrapod Hunter
	Longusunguis	Folivore, Frugivore, **Generalist**, Husking Granivore, Swallowing Granivore, Invertivore, Nectarivore, **Piscivore**	Scavenger, Tetrapod Hunter
	Parabohaiornis	**Folivore**, Frugivore, **Generalist**, Husking Granivore, Swallowing Granivore, Invertivore, Nectarivore, Piscivore	Scavenger, Tetrapod Hunter
	Shenqiornis	**Folivore**, Frugivore, **Generalist**, Husking Granivore, Invertivore, Nectarivore, **Piscivore**	Swallowing Granivore, Scavenger, Tetrapod Hunter
	Sulcavis	**Folivore**, Frugivore, **Generalist**, Husking Granivore, Invertivore, Nectarivore, **Piscivore**	Swallowing Granivore, Scavenger, Tetrapod Hunter
	Zhouornis	**Frugivore**, **Generalist**, Husking Granivore, Nectarivore	Swallowing Granivore, Invertivore, Piscivore, Scavenger, Tetrapod Hunter
Pedal Traditional Morphometrics	*Bohaiornis*	**Non-Raptorial Perching**, Small Raptor	Ground Bird, Shrike-like
	Longusunguis	**Large Raptor**	Ground Bird, Shrike-like
	Parabohaiornis	Non-Raptorial Perching	Ground Bird
	Sulcavis	Large Raptor, unique ecology not in dataset	Shrike-like
	Zhouornis	Small Raptor, Non-Raptorial Perching	Ground Bird, Shrike-like
Skull Traditional Morphometrics	*Bohaiornis*	Folivore, Frugivore, Generalist	Granivore, Nectarivore
	Longusunguis	Folivore, Frugivore, Generalist	Granivore, Nectarivore
	Parabohaiornis	Folivore, Frugivore, Generalist	Granivore, Nectarivore
	Shenqiornis	Unique ecology not in dataset	All sampled diets
	Sulcavis	Unique ecology not in dataset	All sampled diets
	Zhouornis	Unique ecology not in dataset	All sampled diets