RNA polymerase III is involved in regulating *Plasmodium falciparum* virulence

Gretchen Diffendall,[1,2] Aurélie Claës,[1] Anna Barcons-Simon,[1,2,3] Prince Nyarko,[4] Florent Dingli,[5] Miguel Santos,[6] Damarys Loew,[5] Antoine Claessens,[4,7] and Artur Scherf [1,*]

Institut Pasteur, [1]Universite Paris Cité, INSERM U1201, CNRS EMR9195, Paris, France.
[2]Sorbonne Université Ecole doctorale Complexité du Vivant ED515, Paris, France.
[3]Biomedical Center, Division of Physiological Chemistry, Faculty of Medicine, Ludwig-Maximilians-Universität München, Munich 82152, Germany
[4] Laboratory of Pathogen–Host Interaction (LPHI), CNRS, University of Montpellier, France
[5]Institut Curie, PSL Research University, Centre de Recherche, CurieCoreTech Mass Spectrometry Proteomics, Paris, France
[6] Instituto de Medicina Molecular João Lobo Antunes, Faculdade de Medicina da Universidade de Lisboa, Lisboa, Portugal
[7] LPHI, MIVEGEC, CNRS, INSERM, University of Montpellier, France

*Corresponding author: artur.scherf@pasteur.fr

Abstract:

While often undetected and untreated, persistent seasonal asymptomatic malaria infections remain a global public health problem. Despite the presence of parasites in the peripheral blood, no symptoms develop. Disease severity is correlated with the levels of infected red blood cells (iRBCs) adhering within blood vessels. Changes in iRBC adhesion capacity have been linked to seasonal asymptomatic malaria infections, however how this is occurring is still unknown. Here we present evidence that RNA polymerase III (RNA Pol III) transcription in *Plasmodium falciparum* is downregulated in field isolates obtained from asymptomatic individuals during the dry season. Through experiments with in vitro cultured parasites, we have uncovered an RNA Pol III-dependent mechanism that controls pathogen proliferation and expression of a major virulence factor in response to external stimuli. Our findings establish a connection between *P. falciparum* cytoadhesion and a non-coding RNA family transcribed by Pol III. Additionally, we have identified *P. falciparum* Maf1 as a pivotal regulator of Pol III transcription, both for maintaining cellular homeostasis and responding adaptively to external signals. These results introduce a novel perspective that contributes to our understanding of *P. falciparum* virulence. Furthermore, they establish a connection between this regulatory process and the occurrence of seasonal asymptomatic malaria infections.

Introduction:

The parasite *Plasmodium falciparum* is responsible for the deadliest form of human malaria that annually affects over 200 million people, with 619,000 fatal cases, majorly African children under the age of 5. The disease is transmitted to its human host during a blood meal by the parasites' vector, *Anopheles* mosquitoes. Disease symptoms are seen while the parasite multiplies asexually within the host RBCs. During this time, variant surface antigens (VSAs) are exported and displayed on the surface of the iRBCs (Wahlgren *et al*, 2017). During the ~48-hour asexual intraerythrocytic developmental cycle (IDC), the parasite develops through different morphological stages: ring (circulating), trophozoite, and schizont (sequestered in capillaries). Expression of one type of VSA that is linked to immune evasion and pathogenesis, termed *P. falciparum* erythrocyte membrane protein 1 (PfEMP1) (Leech *et al*, 1984), can bind to a wide range of receptors on endothelial cells such as CD36. PfEMP1 binding mediates adhesion to the vascular endothelium within the host, thereby preventing mature iRBCs from traveling to, and being cleared by, the spleen (Smith *et al*, 2013). *Plasmodium* parasites replicate via schizogony inside the host RBCs generating the infectious form, termed merozoites. Upon bursting of the iRBC, merozoites invade new RBCs, thus enabling the cycle to restart. The number of merozoites per schizont has been shown to reduce in unfavorable conditions as a way to diminish the total parasite load and therefore overall disease severity (Mancio-Silva *et al*, 2017b).

P. falciparum chronic infection relies largely on mutually exclusive expression of PfEMP1 surface adhesins, that are encoded by the *var* virulence gene family (Scherf *et al*, 1998; Smith *et al*, 1995; Guizetti & Scherf, 2013). The expression of *var* genes, while under tight epigenetic regulation by the parasite, is additionally controlled by a family of ncRNA, termed RUF6. *P. falciparum* 3D7 encodes 15 RUF6 genes dispersed over several chromosomes located adjacent to central *var* genes (Gardner *et al*, 2002). RUF6 ncRNA have been observed to associate with the active *var* gene in *trans* and transcriptional repression disrupts the monoallelic expression of *var* genes by down-regulating the entire *var* gene family (Barcons-Simon *et al*, 2020; Guizetti *et al*, 2016). Genetically modified parasites that express the entire RUF6 gene family show a general upregulation of *var* genes (Zhang *et al*, 2014; Fan *et al*, 2020).

P. falciparum virulence is linked to cytoadhesion and sequestration as a result of reduced

clearance of iRBCs in the spleen and increased microcapillary obstruction and local

inflammation (Miller *et al*, 2002). Substantial sequestration of iRBCs in critical target organs

has been associated with severe malaria (Miller *et al*, 2013; Guillochon *et al*, 2022). In long-

term asymptomatic infections, found during prolonged low transmission dry seasons in

malaria-endemic regions, decreased cytoadhesion and longer circulation time of iRBCs in the

blood has been observed (Andrade *et al*, 2020). It is not fully understood how asymptomatic

infections develop and are maintained nor is a molecular understanding of host and parasite

factors, that underlies disease severity during the dry and wet seasons (reviewed in (Zhang &

Deitsch, 2022)). Disease severity appears to be influenced by a variety of factors derived from

the human host, parasite, and environment. *Plasmodium* parasites are auxotrophic for various

nutrients including glucose, certain types of amino acids, and lipids, which must be obtained

from the host. Physiological conditions in the host, like diabetes, pregnancy, diet, and

immunity can change the metabolic environment the parasites are exposed to. Numerous

studies conclude that host immunity is a contributing factor to the maintenance of

asymptomatic infections (Kimenyi *et al*, 2019). While it is proposed that *P. falciparum*

virulence regulation contributes to decreased endothelial binding, the mechanism and factors

involved remains to be determined.

Almost all eukaryotes use conserved nutrient sensing pathways, including mTOR, GCN2,

GCN5, Rgt2/Snf3, and Snf1/AMPK signaling cascades (reviewed in (Kumar *et al*, 2021)).

Most notably, the target of rapamycin complex (TORC) pathway involves the signaling of

external factors to control many cellular processes (Wullschleger *et al*, 2006). Specifically,

TORC participates in the regulation of RNA Pol III activity via the phosphorylation of its

inhibitor, Maf1. When nutrients are available and mTOR kinase is active, Maf1 is

hyperphosphorylated resulting in an inactive cytoplasmic state. Stress-induced Maf1

dephosphorylation results in nuclear localization and inhibition of RNA Pol III (Pluta *et al*,

2001; Rollins *et al*, 2007; Vannini *et al*, 2010). This pathway responds to a range of positive

and negative external stimuli, most notably the presence of amino acids, that drive or inhibit

cellular growth. The majority of the TORC pathway components are lost in the apicomplexan

lineage (McLean & Jacobs-Lorena, 2017; Serfontein *et al*, 2010; van Dam *et al*, 2011). While

P. falciparum encodes none of the core TORC1 components, it does contain a homologue to

Maf1 (McLean & Jacobs-Lorena, 2017). The biology of *P. falciparum* Pf-Maf1 is still poorly

characterized.

Here, we present data showing inhibition of RNA Pol III transcription in field isolates from
asymptomatic individuals during the dry season. We present evidence for *P. falciparum* Maf1
as a negative regulator of RNA Pol III-transcribed genes including tRNA and RUF6 ncRNA,
and demonstrate that this mechanism is responsive to external stimuli using *in vitro* culture.
Inhibition of *P. falciparum* cytoadherence, via the regulation of RNA Pol III activity,
represents a new paradigm contributing to *P. falciparum* virulence and links this process to
seasonal asymptomatic malaria infections.

Results:

RUF6 ncRNA is downregulated in asymptomatic individuals during the dry season

We and others have previously established the role of the 15-member RUF6 ncRNA gene
family in *P. falciparum var* gene expression (Barcons-Simon *et al*, 2020; Diffendall *et al*,
2022; Fan *et al*, 2020; Guizetti *et al*, 2016). RUF6 ncRNA protein complex associates with
the active *var* gene expression site (Diffendall *et al*, 2022; Guizetti *et al*, 2016), and
knockdown of the ncRNA RUF6 gene family dramatically reduced *var* gene transcription
(Barcons-Simon *et al*, 2020). Bioinformatic analysis identified highly conserved RNA Pol III
binding elements (conserved A- and B-box sequence) within RUF6 genes (Guizetti *et al*,
2016) (Figure S1A), hinting at a role for RNA Pol III in virulence gene regulation. Thus, we
determined the effects of RNA Pol III inhibition on gene transcription in synchronized wild
type parasites. We observed downregulation of tRNA and RUF6 transcription after 21 hours
of treatment with a commercially available RNA Pol III inhibitor (Figure S1B). As expected,
no changes were observed in the transcription of an RNA Pol II-regulated gene, *uce*
(ubiquitin-conjugating enzyme, PF3D7_0812600) (Figure S1B). As predicted from the RUF6
sequence, our data provide direct experimental evidence for RNA Pol III-dependent
transcription of RUF6 genes.

We next wanted to determine if RNA Pol III-transcribed genes, specifically RUF6 ncRNA,
varied between parasites causing different clinical states of malaria infection, since previous
studies did not include RNA Pol III-transcribed genes for their data analysis (Andrade *et al*,
2020). We first re-analyzed raw RNA-seq data, from a previous study that compared parasites
infecting individuals during the dry versus wet season in a malaria endemic region of Mali
(Andrade *et al*, 2020), to include these genes. We were able to retrieve transcriptional data for
two RNA Pol III-transcribed tRNAs (tRNA Leucine Pf3D7_API05400 and tRNA Asparagine
Pf3D7_API05500, Figure S2A), both of which show significantly lower levels in parasites

infecting asymptomatic individuals during the dry season (n = 12) when compared to
parasites infecting symptomatic individuals during the wet season (n = 12).

We further investigated whether RNA Pol III modulates gene transcription in field isolates of
P. falciparum from different individuals infected during the wet and dry season in malaria
endemic regions (The Gambia). In this pilot study, we analyzed *P. falciparum* steady state
levels of RNA in venous blood samples taken from individuals with asymptomatic
parasitemia during the dry season (n = 17), as well as mildly symptomatic malaria infections
during the wet season (n = 14) in The Gambia (Collins *et al*, 2022; Fogang *et al*, 2023). RT-
qPCR of total RNA showed that levels of tRNAs (Asparagine and Valine) were significantly
lower (p < 0.05) in parasites from asymptomatic infections during the dry season compared to
parasites causing symptomatic infections during the wet season (Figure 1A, B). Likewise,
RUF6 ncRNA was significantly downregulated (p < 0.005) in parasites from asymptomatic
infections during the dry season when compared to symptomatic infections during the wet
season (Figure 1C). Expression was normalized to a housekeeping gene encoding fructose
bisphosphate aldolase (Pf3D7_1444800).

The same previous study, mentioned above, also showed that *var* gene transcription varied in
parasites infecting individuals during the dry versus wet seasons (Andrade *et al*, 2020). We
observed a significantly lower level of *var* gene transcription (p < 0.005) in parasites taken
from asymptomatic individuals during the dry season compared to those taken from
symptomatic individuals during the wet season (Figure 1D). The observed lower transcript
levels of *var* genes may be a direct result of reduced RUF6 ncRNA levels. Taken together,
our results reveal that levels of RNA Pol III-transcribed genes, particularly the regulatory
RUF6 ncRNA, varies significantly between two different clinical states and seasons.

The environmental factor, magnesium, plays a regulatory role in genes transcribed by
RNA Polymerase III

In studies of malaria infections, variations in metabolite levels were observed between
symptomatic and asymptomatic cases across wet and dry seasons. However, the data did not
pinpoint any particular factor that could be reliably linked to a specific condition of the
disease (Andrade *et al*, 2020). Other clinical studies investigated the correlation between
macro- and micro-mineral concentrations and disease severity. In one study, it was discovered

that levels of serum $MgCl_2$ were lower in cases of more severe malaria infections (Innocent *et*

al, 2013). We chose to analyze plasma samples from individuals participating in the same

study as referenced in Figure 1, focusing on magnesium concentrations. Our findings show a

significant increase in $MgCl_2$ levels in asymptomatic individuals during the dry season

compared to symptomatic individuals in the wet season, as depicted in Figure 2A.

Our subsequent aim was to delve into the potential molecular mechanisms connecting

external factors, like magnesium, with the changes in RUF6 ncRNA levels as shown in Figure

2B. For this, we employed in vitro cultured *P. falciparum* asexual blood stage parasites. We

supplemented culture medium with $MgCl_2$, noted for its varying levels in individuals with

malaria (see Figure 2A). In addition, we mimicked nutrient deprivation with isoleucine-

deprived culture medium (as was described in (Babbitt *et al*, 2012; McLean & Jacobs-Lorena,

2017)), fever conditions with incubation at 40ºC, and food starvation with low glucose levels

(Jensen *et al*, 1983; Fang *et al*, 2004). We evaluated the steady state levels of RNA Pol I-

(rRNA A1), RNA Pol II- (UCE and the active *var*), and RNA Pol III- (tRNA Valine and

RUF6 ncRNA) transcribed genes in clonal parasites at late ring stage. We observed that high

temperature and low glucose conditions affected gene transcription of RNA Pol I, II and Pol

III (Figure S2B). Deprivation of isoleucine and $MgCl_2$ supplementation seemingly affected

only RNA Pol III-transcribed genes with the exception of the RNA Pol II-transcribed active

var gene, Pf3D7_1240900, likely as a result of decreased RUF6 ncRNA as was reported

earlier (Barcons-Simon *et al*, 2020).

To explore the underlying molecular pathway that leads to the downregulation of RNA Pol III

activity in malaria parasites, we continued using $MgCl_2$ supplementation. We selected a

concentration of 3mM total $MgCl_2$, based on reports indicating it does not inhibit the growth

of *P. falciparum* asexual blood stage and is within a physiological range (Jahnen-Dechent &

Ketteler, 2012; Hess *et al*, 1995). Additionally, we verified lower magnesium concentrations

and show as low as 1mM total $MgCl_2$, observed in healthy individuals, impacts both Pol III-

transcribed RUF6 and tRNA, while not affecting Pol II-transcribed UCE (Figure S3B). RNA-

seq was done on control and $MgCl_2$ supplemented cultured parasites during two time points

of the asexual blood stage parasites (ring and trophozoite) to further confirm that RNA Pol III

is the main target (Figure 2C, D). RNA-seq data was compared to the microarray time course

data in (Bozdech *et al*, 2003) as in (Lemieux *et al*, 2009), which provided a statistical

estimation of cell cycle progression at 12 and 24 hpi in both control and MgCl$_2$ supplemented

parasites (Figure S3B).

These data indicate that any differences in transcription were due to MgCl$_2$ supplementation

and not differences in cell cycle progression. During ring stage (Figure 2C), 77% of the

down-regulated genes contain typical RNA Pol III promoter sequences, A- and B-box

consensus (from Figure S1A). A total of 27 ncRNA genes were downregulated (logFC < 1.95,

FDR < 0.05) including all 15 RUF6 ncRNA, 6 tRNAs, 1 spliceosomal RNA, and 4 snoRNA.

Additionally, 12 protein coding genes were up-regulated with no conserved pathway nor

molecular or biological process. During trophozoite stage (Figure 2D), 74% of the down-

regulated genes have typical RNA Pol III promoter sequences. This demonstrates that MgCl$_2$

supplementation predominantly inhibits RNA Pol III-transcribed genes, including the entire

RUF6 gene family.

Nuclear PfMaf1 protein is essential to regulate RNA Pol III

In almost all eukaryotes, the target of rapamycin complex (TORC) pathway has been shown

to participate in the regulation of RNA Pol III activity via the phosphorylation of its only

known inhibitor, Maf1 (Figure 3A, reviewed in (Wullschleger *et al*, 2006)). Upon

phosphatase activation, modulated in response to external stimuli such as low nutrient

availability, Maf1 shuttles to the nucleus to repress RNA Pol III activity (Michels *et al*, 2010).

Because Maf1 is conserved in the apicomplexan lineage, a predicted essential blood stage

protein in *P. falciparum*, we set out to explore if *P. falciparum* Maf1 (PfMaf1) functions

similarly to model eukaryotes. We generated a PfMaf1 parasite line tagged with a 3HA tag

and a ligand-controlled destabilization domain (ddFKBP), PfMaf1-FKBP (Figure 3B top).

PfMaf1-FKBP transfected parasite clonal populations were validated for proper integration

and confirmation that the system was working, shown by western blot for addition and

removal of the ligand, Shield-1, over the course of 3 cycles (Figure 3B bottom). We observed

that upon removal of Shield-1, PfMaf1 levels were shown to decrease by approximately 57%

in total extracts after one cycle and almost complete degradation was achieved after 2 cycles.

Maf1 in Plasmodium species seemingly affects asexual blood stage proliferation making

classical gene KO experiments unattainable (McLean & Jacobs-Lorena, 2017; Zhang *et al*,

2018), pointing to a regulatory role of the putative RNA Pol III inhibitor under normal *in*

vitro culture growth conditions. Once PfMaf1 is depleted (Figure 3B bottom, 3 cycles -

Shield), transcription of tRNA Valine and RUF6 ncRNA increased dramatically by 3-fold (Figure 3C), possibly perturbing cellular homeostasis. RNA Pol II-transcribed genes were unaffected. Parasite growth was assessed over the course of 5 days for two PfMaf1-FKBP clones cultured in the presence or absence of PfMaf1 (+/- Shield-1, removed one cycle before day 0, Figure 3D). A significant difference in growth was achieved, most evidently in parasites cultured without Shield for 3 cycles, consistent with transposon mutagenesis scoring PfMaf1 essential (Zhang *et al*, 2018). Dead parasites were not detected by Giemsa staining even up to 8 cycles without Shield-1. We confirmed that differences in merozoite numbers per schizont contributed to the change in growth rate between the control and KD PfMaf1 parasites after the 3rd cycle KD (Figure 3E).

Additionally, we demonstrated that nuclear PfMaf1 interacts with the RNA Pol III protein complex by performing co-immunoprecipitation followed by quantitative mass spectrometry (Co-IP MS) of cytoplasmic and nuclear fractions of PfMaf1-HA-ddFKBP. Each fraction was split to include a control with no antibody added. Analysis of the quantitative mass spectrometry data revealed unique interactome of cytoplasmic PfMaf1 and nuclear PfMaf1 in their respective samples compared to their controls (Figure S3A, B). Gene Ontology (GO) analysis showed that PfMaf1-associated proteins uniquely found in the nuclear fraction, and neither of the two controls nor cytoplasmic fraction, are significantly represented by the biological function category of "RNA Polymerase III transcription" (p = 0.00175) with two subunits of RNA Pol III (Pf3D7_1206600 and Pf3D7_1329000). (GO) analysis of significant and unique proteins found only in the cytoplasmic fractions, and not nuclear, are significantly represented by the molecular function category of "protein kinase activator" (p = 0.0143, Pf3D7_1103100). Additionally, two protein phosphatases (Pf3D7_1127000 and Pf3D7_1464600) were significantly enriched in the cytoplasmic fraction (Figure S4C). Figure 3F illustrates the cytoplasmic and nuclear enzymes associated with PfMaf1, which could play a role in sensing environmental shifts that lead to the inhibition of Pol III.

Magnesium modulates cytoadhesion through PfMaf1-regulated RNA Pol III transcription

We further investigated if nuclear PfMaf1 activity could be regulated, in response to external factors, to lead to changes in RNA Pol III activity using *P. falciparum* wildtype parasites cultured *in vitro*. The *P. falciparum* cellular localization of PfMaf1 was investigated in response to external stimuli (+MgCl$_2$). Parasites were tightly synchronized, split into control,

addition of $MgCl_2$ and harvested during late ring stage. In contrast to several eukaryotic
model systems, under normal culture conditions we observed PfMaf1 in cytoplasmic and
nuclear fractions using immunoprecipitation followed by western blot analysis (Figure 4A,
S4A). Nuclear PfMaf1 levels increased upon addition of $MgCl_2$ compared to control nuclear
PfMaf1 (Figure 4A). Immunofluorescence (IF) assays showed that PfMaf1 forms foci-like
aggregates in the cytoplasm and near the nuclear periphery in both culture conditions (Figure
S4B).
275
Next, we used the PfMaf1-FKBP transfected parasite lines, to show that the downregulation
of RNA Pol III-transcribed genes, triggered by external stimuli, is dependent on PfMaf1. We
used the PfMaf1 KD parasites (2nd cycle, see Figure 3B) in control conditions and $MgCl_2$
supplementation to the growth medium in ring stage parasites for RT-qPCR (Figure 4B). We
confirmed the previously observed down-regulation of RNA Pol III-transcribed tRNA Valine
and RUF6 upon $MgCl_2$ supplementation in two independent PfMaf1-FKBP parasite clones.
Notably, when PfMaf1 degradation was induced, the addition of $MgCl_2$ did not affect Pol III
transcription. These results demonstrate that PfMaf1 mediates the observed decrease in RNA
Pol III-transcribed genes with $MgCl_2$ supplementation.
285
286 Previous attempts to develop a knockout (KO) line did not yield success, as highlighted by
287 McLean & Jacobs-Lorena in 2017. However, our strain with an inducible system for
288 degrading PfMaf1 protein has unveiled a twofold function of PfMaf1: i) to help maintain Pol
289 III transcription at levels essential for optimal parasite proliferation, and ii) to serve as a key
290 element in an environmental sensing pathway that directly controls Pol III activity.
291
Finally, we set out to link *P. falciparum* virulence gene expression with decreased RUF6
transcription, by investigating the effect of $MgCl_2$ supplementation on cytoadherence of 3D7
iRBCs. We found that in static binding assays to the endothelial receptor CD36, bound to
plastic dishes, adhesion of iRBCs was reduced by 50% with $MgCl_2$ supplementation (Figure
4C). This was also confirmed at the protein level using antibodies directed against the
conserved intracellular ATS domain of PfEMP1 (Figure 4D). Our findings, which
demonstrate the inducible inhibition of cytoadhesive capacity, are particularly significant for
asymptomatic *P. falciparum* infections, as illustrated in Figure 4E.
300
Discussion

Our study reveals a regulatory mechanism in *P. falciparum* involving RNA Polymerase III,
which plays a pivotal role in the parasite's virulence. This discovery illuminates a previously
unidentified adaptive molecular process in the parasite's cytoadhesion and proliferation. We
propose a connection with asymptomatic infections prevalent during the dry season in African
regions, a period characterized by reduced mosquito transmission. In our analysis, we
compared RNA from parasites in symptomatic individuals during the wet season to those in
asymptomatic individuals in the dry season. In this comparison, we noted a reduction in the
levels of tRNAs and RUF6 ncRNA, with the latter being recognized for its involvement in
regulating *var* gene expression, as illustrated in Figures 1A-C. To date, RNA Polymerase III
(Pol III) has not been acknowledged as a factor influencing the virulence of *P. falciparum* in
mild or asymptomatic malaria infections. Additionally, our research revealed a link between
elevated serum $MgCl_2$ levels and asymptomatic malaria infections, as shown in Figure 2A.
The normal concentration range for $MgCl_2$ in human serum is between [0.7–1.0mM], while
mild hypermagnesemia is characterized by levels ranging from [2.2-3.5mM]. In our pilot
study, the average concentration was observed to be around [0.7mM] during the wet season
and increased to [1.1mM] in the dry season.

To investigate the potential mechanistic link between the changes in $MgCl_2$ levels and Pol III
activity, we established an in vitro culture assay for *P. falciparum* that enabled us to delve
into the molecular mechanism underpinning the environment-dependent inhibition of Pol III.
We experimented with various concentrations of $MgCl_2$ to assess its effect on Pol III activity.
The standard culture medium we used contains [0.5mM] of $MgCl_2$. Our observations
indicated a progressive inhibition of Pol III activity in the range of [1-3mM]. To further
explore the molecular mechanisms involved, we opted to use a concentration of 3 mmol/L in
our continued studies. Quantitative RT-PCR analysis showed inhibition of Pol III
transcription of tRNA and the RUF6 ncRNA gene family. Subsequent RNA seq data
validated that transcriptional inhibition, induced by $MgCl_2$ supplementation, is primarily
limited to canonical RNA Pol III-transcribed genes, which include A and B box containing
sequences, and notably the RUF6 gene, crucial for the activation of *var* genes. Our findings
established that the inhibition of *var* gene expression, mediated by Pol III, led to a decrease in
the cytoadherence of infected red blood cells (iRBCs) when supplemented with $MgCl_2$, as
shown in Figure 4C. We noted a reduction of over 50% in binding to the endothelial receptor
CD36 in a static binding assay. It is expected that the inhibition of iRBC cytoadherence

would be even more pronounced under the physiological shear stress typically found in
human capillaries (Crabb *et al*, 1997).

Having established the role of Pol III in the expression of virulence genes, we subsequently
showed a link between environmental factors and Maf1, which is the only known eukaryotic
repressor of Pol III. We created an inducible system designed to specifically reduce PfMaf1
protein levels, enabling us to explore its influence on Pol III activity. Proteomic analysis
through mass spectrometry of PfMaf1's interactome showed its association with various
nuclear RNA Pol III subunits, reinforcing its function as a repressor of Pol III in *P.*
falciparum. PfMaf1 showed nuclear presence under standard growth conditions. Nuclear
shuttling usually increases under specific conditions (stress, nutrient starvation) by changing
the phosphorylation state of the Maf1 protein in eukaryotic organisms (Michels *et al*, 2010).
Our study supports a model in which a critical baseline amount of PfMaf1 in the nucleus is
essential to modulate the expression of tRNAs by RNA Pol III during the multinucleated
blood stage of the parasite. This regulation is essential for the parasite to balance its energy
resources efficiently while producing large numbers of new infective forms, known as
merozoites. Under optimal growth conditions, a single parasite undergoes several rounds of
genome replication, forming approximately 30 nuclei within a shared cytoplasm. When
PfMaf1 is completely removed, the transcription levels of tRNA Valine and RUF6 ncRNA
increase more than threefold, while the transcription level of a Pol II-mediated housekeeping
gene UCE remains unchanged. This significant elevation of tRNA transcription beyond
physiological levels could potentially lead to an energy deficit within the cell, resulting in the
observed reduction in the parasite's multiplication rate, as demonstrated in Figure 3C and E.
In clinical isolates, parasite densities have been associated with malaria pathogenesis (Nyarko
& Claessens, 2021; Thomson-Luque *et al*, 2021) and low parasite densities are also a feature
of asymptomatic malaria (Murray *et al*, 2017).

Nutrient availability in the host has been demonstrated to have a profound effect on the
replication of parasites using *P. berghei* as a rodent malaria model (Mancio-Silva *et al*,
2017a). Reports from cultured blood stage *P. falciparum* parasites showed that amino acid
deficiency can alter parasite growth and survival to stress (McLean & Jacobs-Lorena, 2017;
Marreiros *et al*, 2023). Furthermore, a recent report suggested that nutrients involved in S-
adenosylmethionine (SAM) metabolism can also affect *var* gene switching in cultured
parasites (Schneider *et al*, 2023). Individuals infected with malaria, compared to healthy

individuals, were found to have decreased levels (up to 54%) of plasma free amino acids,
which could be the result of a nutrient-reduced diet (Leopold *et al*, 2019). In fact, dietary
intake, including energy, protein, iron, zinc, calcium and folate, was found to decrease
significantly in individuals during the dry season in areas of Africa (M'Kaibi *et al*, 2015).
Blood serum magnesium levels can fluctuate in humans and were found to decrease with
increasing disease severity (Innocent *et al*, 2013). In our study, plasma magnesium levels
were significantly increased in asymptomatic individuals during the dry season compared to
symptomatic individuals during the wet season (Figure 2A). Wet season samples were within
the range of normal serum magnesium levels, whereas dry season samples were higher. It is
noteworthy that intracellular magnesium was found to increase in yeast as a result of calorie
restriction, defined as a decrease in dietary glucose intake (Abraham *et al*, 2016), indicating
that our $MgCl_2$ supplementation could be mimicking low glucose conditions and thus, low
nutrient availability. Although interactions between human malaria and malnutrition have
been studied for many years, the evidence linking an effect between the two remains
inconclusive at the mechanistic level. Our study sets the stage for exploring whether there are
additional external stimuli, beyond magnesium chloride, that could activate the regulatory
pathway of RNA Polymerase III. Given the lack of a conventional TOR pathway in malaria
parasites (Serfontein *et al*, 2010; van Dam *et al*, 2011), the exact signaling pathway that
activates PfMaf1 remains unknown. Our mass spectrometry analysis of PfMaf1's cytoplasmic
binding partners has identified interactions with enzymes, such as phosphatases and kinases,
which could potentially influence Maf1 activity. This discovery opens up new avenues for
future research into the environmental sensing mechanisms that function upstream of PfMaf1.
391

392 A traditional view posits that the decreased cytoadherence seen in dry-season parasites may
393 be affected by host immunity, which impairs the parasites' adhesion capabilities. Although
394 experimental evidence supporting this theory is missing, our research introduces a novel
395 perspective to this critical topic by highlighting the role of metabolic changes in modulating
396 virulence gene expression.
397

In summary, our research demonstrates that RNA Polymerase III activity, regulated by
environmental factors, plays a crucial role in the proliferation of parasites and in reducing the
cytoadhesive capacity of *P. falciparum*. This discovery unveils a previously unknown
molecular process, significantly enhancing our understanding of subclinical parasite

persistence. The insights gained from this study could pave the way for novel strategies aimed

at preventing severe malaria by promoting reduced pathogen virulence.

Materials and Methods:

Parasite and serum samples from malaria infected patients

Venous blood draw of different infected individuals in The Gambia during the dry and wet

seasons was collected as previously described in (Collins *et al*, 2022; Fogang *et al*, 2023).

The study protocol was reviewed and approved by the Gambia Government/MRC Joint

Ethics Committee (SCC 1476, SCC 1318, L2015.50) and by the London School of Hygiene

& Tropical Medicine ethics committee (Ref 10982). The field studies were also approved by

local administrative representatives, the village chiefs. Written informed consent was obtained

from participants over 18 years old and from parents/guardians for participants under 18

years. Written assent was obtained from all individuals aged 12-17 years. The dry season

months include January, February, March, April, and May. The wet season months include

November and October. For additional information about each sample, see Supplementary

Table 2.

Polymerase III inhibition assay

423   Parasites were treated with 50 μM of RNA Pol III inhibitor CAS 577784-91-9 (Calbiochem,

424   Merck) after sorbitol treatment at 3 ±3hpi and RNA was harvested at 24hpi in parallel with

425   untreated and control samples. The control was treated with the same volume of DMSO

426   added to the inhibitor treated flasks of stock solution.

Clinical isolate RT-qPCR

Total RNA was extracted from TRizol using an miRNeasy minikit and performing on-column

DNase treatment (Qiagen) and continued as previously described in ((Collins *et al*, 2022)).

Transcript levels were shown by using the following primers: RUF6, Valine tRNA,

Asparagine tRNA, and *var* DBLalpha were normalized to the reference gene, fructose-

bisphosphate aldolase (PF3D7_1444800, Figure 2A, Table S1). The starting quantity means

from three replicates were extrapolated from a standard curve of serial dilutions of 3D7

genomic DNA.

Plasma magnesium concentration assay

Plasma magnesium levels were determined using a commercial magnesium assay kit (Sigma-

Aldrich MAK026) from individuals from the same study as in (Collins *et al*, 2022; Fogang *et*

al, 2023). Complete information on tested individual samples can be found in Supplementary

Table 2. The protocol was followed as described in the kit and read on a Synergy 2 microplate

reader for spectrophotometric reading.

Parasite culture and synchronization

Asexual blood stage 3D7 *P. falciparum* parasites were cultured as previously described in

(Lopez-Rubio *et al*, 2009). Parasites were cultured in human RBCs (obtained from the

Etablissement Français du Sang with approval number HS 2019-24803) in RPMI-1640

culture medium (Thermo Fisher 11875) supplemented with 10% v/v Albumax I (Thermo

Fisher 11020), hypoxanthine (0.1 mM final concentration, C.C.Pro Z-41-M) and 10 mg

gentamicin (Sigma G1397) at 4% hematocrit and under 5% O_2, 3% CO_2 at 37 °C. Static

parasite development was monitored by Giemsa staining. Parasites were synchronized by

sorbitol (5%, Sigma S6021) lysis at ring stage, plasmagel (Plasmion, Fresenius Kabi)

enrichment of late stages 24 h later, and an additional sorbitol lysis 3 h after plasmagel

enrichment. The 0 h time point was considered to be 1.5 h after plasmagel enrichment.

Parasites were harvested at 1–5% parasitemia.

External stimuli induction

Synchronized parasites were divided into control, addition of magnesium chloride ($MgCl_2$),

isoleucine-deficient, low glucose, and 40 degree Celsius and harvested at 18hpi. Parasites

exposed to an addition of $MgCl_2$ were supplemented with 0.5mM, 1.5mM, and 2.5 mM

$MgCl_2$, for a final concentration of [1mM], [1.5mM], [2mM], and [3mM] including [0.5mM]

found in RPMI. Isoleucine-deficient medium consisted of 10.3 g/liter RPMI 1640 isoleucine

(Ile) Drop-out medium (United States BioLogicals; catalog no. R9014), supplemented with

2.0 g/liter $NaHCO_3$, 6.0 g/liter HEPES, 10% v/v Albumax I (Thermo Fisher 11020),

hypoxanthine (0.1 mM final concentration, C.C.Pro Z-41-M) and 10 mg gentamicin (Sigma

G1397). Low glucose (0.25mg/mL) RPMI was made by adding glucose (Dextrose) to

glucose-free media: 2.979 g HEPES + 50 mL albumax + 10 mL hypoxanthine + 200 μL

468	gentamycin to final volume of 500 mL with glucose free RPMI (11879). The pH was adjusted
469	to 7.3 with NaOH and filter sterilized. 40-degree Celsius samples were incubated in an
470	adjacent incubator set to 40 degrees Celsius under 5% O_2, 3% CO_2. Parasites were then
471	harvested with 0.075% Saponin lysis at ~2-5% parasitemia for RNA, genomic DNA, and
472	protein extraction at 18hpi.
473	

RNA isolation and reverse transcription-quantitative PCR (RT-qPCR)

RNA was isolated from synchronized parasite cultures harvested at 18hpi after saponin lysis in 0.075% saponin in PBS, followed by one wash in Dulbecco's phosphate-buffered saline (DPBS, Thermo Fisher 14190) and resuspension in the QIAzol reagent. Total RNA was extracted using an miRNeasy minikit and performing on-column DNase treatment (Qiagen). Reverse transcription from total RNA was achieved using SuperScript VILO (Thermo Fisher Scientific) and random hexamer primers. cDNA levels were quantified by quantitative PCR in the CFX384 real time PCR detection system (BioRad) using Power SYBR Green PCR Master Mix (Applied Biosystems) and primers from a previous study (Guizetti *et al*, 2016). Starting quantity means of three replicates were extrapolated from a standard curve of serial dilutions of genomic DNA. Transcript levels were shown by using the following primers: RUF6, Valine tRNA, Alanine tRNA, Asparagine tRNA, *var* 58 (PF3D7_1240900), and *var* DBLalpha were normalized to the reference gene, fructose-bisphosphate aldolase (PF3D7_1444800).

Stranded RNA sequencing and analysis

Infected RBCs containing synchronized (12 and 24hpi) parasites were lysed in 0.075% saponin (Sigma S7900) in DPBS at 37°C. The parasite cell pellet was washed once with DPBS and then resuspended in 700 µL QIAzol reagent (Qiagen 79306). Total RNA was subjected to rRNA depletion to ensure ncRNA and mRNA capture using the RiboCop rRNA Depletion Kit (Lexogen) prior to strand-specific RNA-seq library preparation using the TruSeq Stranded RNA LT Kit (Illumina) with the KAPA HiFi polymerase (Kapa Biosystems) for the PCR amplification. Multiplexed libraries were subjected to 150 bp paired-end sequencing on a NextSeq 500 platform (Illumina). Sequenced reads (150 bp paired end) were mapped to the *P. falciparum* genome (Gardner *et al*, 2002) (plasmoDB.org, version 3, release 57) using "bwa mem" (Li & Durbin, 2009) allowing a read to align only once to the reference genome (option "–c 1"). Alignments were subsequently filtered for duplicates and a mapping

quality ≥ 20 using samtools (Li & Durbin, 2009). Three biological replicates for -MgCl$_2$ and

+MgCl$_2$ samples were analyzed for both timepoints.

Generation of PfMaf1 strains

All cloning was performed using KAPA HiFi DNA Polymerase (Roche 07958846001), In-

Fusion HD Cloning Kit (Clontech 639649), and XL10-Gold Ultracompetent *E. coli* (Agilent

Technologies 200315). Transgenic pSLI parasites were generated as previously described in

(Birnbaum *et al*, 2017) with the following modifications: GFP was replaced with a 3HA tag

and a ddFKBP domain was added after the protein of interest, PfMaf1. For localization and

knock down studies, the last 500-1000 bp of target gene, PfMaf1 Pf3D7_0416500, was

cloned into pSLI-3HA-ddFKBP. Each sequence started with an in-frame stop codon but the

stop codon at the end of the gene was removed. 50 μg of plasmid DNA was transfected into

ring stage 3D7 *P. falciparum* parasites using the protocol described elsewhere (Hasenkamp *et*

al, 2013). Transfected parasites were selected with constant drug selection pressure of 2.56

nM WR99210 (Jacobus Pharmaceuticals) to obtain a cell line containing the episomal

plasmid. A second drug selection using 400 μg/ml of G418 was done to select for integrants.

Once parasites emerged, gDNA of each integration cell line was collected using a commercial

kit (DNeasy Blood & Tissue Kit) and checked by PCR to show that integration occurred at

the correct locus. Both genome and vector specific primers for the 5' and 3' region were used

so that the PCR product would cover the plasmid/genome junction. Vector primers used were

the same as described (Birnbaum *et al*, 2017). Once proper size gel bands from PCR were

seen, parasites were cloned by limiting dilution, and the targeted genomic locus was

sequenced to confirm tag and FKBP integration.

Flow cytometry

Two different pSLI-Maf1-FKBP parasite clones were tightly synchronized and diluted to

0.2% parasitemia (5% hematocrit) at ring stage. One cycle before, Shield-1 was removed

from half of the culture. The growth curve was performed in a 96-well plate (200 μl culture

per well) with three technical replicates of two biological replicates per condition per clone.

At 0 h, 24 h, 48 h, 72 h, and 96 h, 5 μl of the culture was stained in 95 μl of DPBS

supplemented with 2× Sybr Green I (Ozyme; stock = 10,000×) for 30 min at room

temperature, diluted 20-fold in D-PBS (final volume = 200 μl), and the Sybr Green

fluorescence measured in a Guava easyCyte Flow Cytometer (EMD Millipore). 30000 events

were counted in duplicate to establish an accurate parasitemia value for each culture. Data

was analyzed using the InCyte software (EMD Millipore).

Western blot analysis

Shield-1 was removed for 3 consecutive cycles to monitor the degradation of PfMaf1. Total

protein extracts were prepared from trophozoite stages for control and -Shield-1 for 3 cycles.

Additionally, membrane extracts were prepared for control and $MgCl_2$ supplementation from

parasites isolated from plasmion enrichment. iRBCs were washed once with DPBS at 37°C

and lysed with 0.075% saponin (Sigma S7900) in DPBS at 37°C. Parasites were washed once

with DPBS, resuspended in 1 ml cytoplasmic lysis buffer (25 mM Tris–HCl pH 7.5, 10 mM

NaCl, 1.5 mM $MgCl_2$, 1% IGEPAL CA-630, and 1× protease inhibitor cocktail ["PI", Roche

11836170001]) at 4°C, and incubated on ice for 30 min. Cells were further homogenized with

a chilled glass douncer, and the cytoplasmic lysate was cleared with centrifugation (13,500 g,

10 min, 4°C). The pellet (containing the nuclei) was resuspended in 100 μl nuclear extraction

buffer (25 mM Tris–HCl pH 7.5, 600 mM NaCl, 1.5 mM $MgCl_2$, 1% IGEPAL CA-630, PI) at

4°C and sonicated for 10 cycles with 30 s (on/off) intervals (5 min total sonication time) in a

Diagenode Pico Bioruptor at 4°C. The nuclear lysate was cleared with centrifugation (13,500

g, 10 min, 4°C). Membrane extracts were prepared by resuspending parasite pellets in NETT

buffer (50mM Tris pH8, 150mM NaCl, 5mM EDTA, 1% IGEPAL CA-630, PI) and

incubated at 4°C for 10 min. Supernatant was removed after centrifugation (13,500 g,

10 min, 4°C) and the pellet was resuspended in Tris-saline buffer (50mM Tris pH8, 150mM

NaCl, 2% SDS, PI) and sonicated for 6 cycles with 30 s (on/off) intervals (3 min total

sonication time) in a Diagenode Pico Bioruptor at 4°C. The membrane lysates were cleared

with centrifugation (13,500 g, 10 min, 4°C). All protein samples were supplemented with

NuPage Sample Buffer (Thermo Fisher NP0008) and NuPage Reducing Agent (Thermo

Fisher NP0004) and denatured for 5 min at 95°C. Proteins were separated on a 4-15% TGX

(Tris-Glycine eXtended) (Bio-Rad) and transferred to a PVDF membrane. The membrane was

blocked for 1 h with 5% milk in PBST (PBS, 0.1% Tween 20) at 25°C. HA-tagged proteins,

and histone H3 were detected with anti-HA (Abcam 9110, 1:1,000 in 5% milk-PBST) and

anti-H3 (Abcam ab1791, 1:1,000 in 5% milk-PBST) primary antibodies, respectively,

followed by donkey anti-rabbit secondary antibody conjugated to horseradish peroxidase

("HRP", Sigma GENA934, 1:5,000 in 5% milk-PBST). Anti-ATS and anti-NapL were

detected as previously described (Nacer *et al*, 2015; Dawn *et al*, 2014). Aldolase was detected

with anti-aldolase-HRP (Abcam ab38905, 1:5,000 in 5% milk-PBST). HRP signal was

developed with SuperSignal West Pico chemiluminescent substrate (Thermo Fisher 34080)

and imaged with a ChemiDoc XRS+ (Bio-Rad).

Merozoite number per schizont analysis:

Mature schizonts were assessed by microscopic analysis of Giemsa-stained smears and

manually quantified using ImageJ software, as reported previously (Marreiros *et al*, 2023).

100 segmented schizonts with clearly individualized merozoites containing a single hemozoin

crystal were quantified per condition.

Co-Immunoprecipitation followed by Mass spectrometry (Co-IP-MS)

PfMaf1-HA-ddFKBP tagged parasites (*n* = 5 biological replicates) were synchronized. At

24hpi, each culture (1.5×10^9 parasites) was centrifuged and RBCs were lysed with six

volumes of 0.15% saponin in DPBS for 5 min at 4°C. Parasites were centrifuged at 4,000 *g*

for 5 min at 4°C, and the pellet was washed twice with DPBS at 4°C. Parasites were then

cross-linked with 1% formaldehyde for 15 min at room temperature and quenched with 125

mM glycine for 5 min on ice. Cross-linked parasites were washed twice with DPBS and then

resuspended in 900uL of cytoplasmic lysis buffer (10 mM Tris–HCl pH 7.5, 1 mM EDTA,

0.65% IGEPAL CA-630, 10mM NaCl) supplemented with protease inhibitors (Thermo Fisher

78440) at 4°C and incubated with rotation for 30 min at 4°C. Extracts were centrifuged for 10

min at 2000g at 4°C and the cleared cytoplasmic supernatant was removed and kept on ice.

The nuclear pellet was resuspended in 900 μL nuclear lysis buffer (10 mM Tris–HCl pH 7.5,

500 mM NaCl, 1 mM EDTA, 1% sodium deoxycholate, 0.1% SDS, 1% IGEPAL CA-630, PI)

at 4°C and transferred to 1.5 ml sonication tubes (300 μL per tube, DiagenodeC30010016).

Samples were sonicated for 5 min (30 s on/off) in a Diagenode Pico Bioruptor at 4°C. Lysates

were then centrifuged for 10 min at 13,500g at 4°C and supernatant was transferred to a fresh

tube. Cytoplasmic fractions were mixed with 2:3 ratio of cytoplasmic dilution buffer (10 mM

Tris-HCl pH 7.5, 150 mM NaCl, 0.5 mmM EDTA) and nuclear supernatants were mixed with

1:3 ratio of nuclear dilution buffer (10 mM Tris-HCl pH 7.5, 0.5 mmM EDTA). Cytoplasmic

and nuclear fraction supernatants were incubated with 1ug of α-HA antibody (Abcam 9110)

and 25uL Protein G Magnetic Dynabeads (Invitrogen), pre-incubated for a minimum of 2

hours and washed twice with dilution buffer, overnight with rotation at 4°C. The next day, the

beads were collected on a magnet and the supernatant was removed. While on the magnetic

stand, beads were washed twice with 500 μL wash buffer (10 mM Tris-HCl pH 7.5, 150 mM

NaCl, 0.5 mM EDTA, 0.05% NP40), once with 25 mM NH_4HCO_3 (Sigma 09830) buffer,

and then transferred to new tube. Finally, the beads were resuspended in 100 µl of 25 mM

NH$_4$HCO$_3$ (Sigma 09830) and digested by adding 0.2 µg of trypsin-LysC (Promega) for

1 h at 37 °C. Samples were then loaded into custom-made C18 StageTips packed by

stacking one AttractSPE® disk (#SPE-Disks-Bio-C18-100.47.20 Affinisep) in a 200 µL

micropipette tip for desalting. Peptides were eluted using a ratio of 40:60

CH3CN:H2O + 0.1% formic acid and vacuum concentrated to dryness with a SpeedVac

apparatus. Peptides were reconstituted in 10 of injection buffer in 0.3% trifluoroacetic acid

(TFA) before liquid chromatography-tandem mass spectrometry (LC-MS/MS) analysis.

Online LC was performed with an RSLCnano system (Ultimate 3000, Thermo Scientific)

coupled to an Orbitrap Eclipse mass spectrometer (Thermo Scientific). Peptides were trapped

on a 2 cm nanoviper Precolumn (i.d. 75 µm, C18 Acclaim PepMap^{TM} 100, Thermo Scientific)

at a flow rate of 3.0 µL/min in buffer A (2/98 MeCN/H2O in 0.1% formic acid) for 4 min to

desalt and concentrate the samples. Separation was performed on a 50 cm nanoviper column

(i.d. 75 µm, C18, Acclaim PepMap^{TM} RSLC, 2 µm, 100Å, Thermo Scientific) regulated to a

temperature of 50°C with a linear gradient from 2% to 25% buffer B (100% MeCN in 0.1%

formic acid) at a flow rate of 300 nL/min over 91 min. MS1 data were collected in the

Orbitrap (120,000 resolution; maximum injection time 60 ms; AGC 4 x 10^5). Charges states

between 2 and 7 were required for MS2 analysis, and a 60 s dynamic exclusion window was

used. MS2 scan were performed in the ion trap in rapid mode with HCD fragmentation

(isolation window 1.2 Da; NCE 30%; maximum injection time 60 ms; AGC 104)

For identification, the data were searched against the *Homo Sapiens* (UP000005640_9606)

and the *Plasmodium falciparum 3D7* (UP000001450_36329) UniProt databases using Sequest

HT through proteome discoverer (version 2.4). Enzyme specificity was set to trypsin and a

maximum of two miss cleavages sites were allowed. Oxidized methionine, Met-loss, Met-

loss-Acetyl and N-terminal acetylation were set as variable modifications.

Carbamidomethylation of cysteins were set as fixed modification. Maximum allowed mass

deviation was set to 10 ppm for monoisotopic precursor ions and 0.6 Da for MS/MS peaks.

The resulting files were further processed using myProMS v3.9.3 (https://github.com/bioinfo-

pf-curie/myproms) (Poullet *et al*, 2007) FDR calculation used Percolator (The *et al*, 2016) and

was set to 1% at the peptide level for the whole study. The label free quantification was

performed by peptide Extracted Ion Chromatograms (XICs), reextracted across all conditions

and computed with MassChroQ version 2.2.21 (Valot *et al*, 2011) For protein quantification,

XICs from proteotypic peptides shared between compared conditions (TopN matching) and

missed cleavages were allowed. Median and scale normalization was applied on the total

signal to correct the XICs for each biological replicate (n = 5 in each condition). To estimate

the significance of the change in protein abundance, a linear model (adjusted on peptides and

biological replicates) was performed, and p-values were adjusted using the Benjamini–

Hochberg FDR procedure. Proteins with at least 2 distinct peptides in 3 replicates of a same

state, a 2-fold enrichment and an adjusted p-value $\leq$ 0.05 were considered significantly

enriched in sample comparisons. Proteins unique to a condition were also considered if they

matched the peptides criteria.

Immunofluorescence assay

pSLI-Maf1-FKBP control, +MgCl$_2$ parasites were used with rat anti-HA (Roche 3F10)

antibodies. 10 µl of iRBCs were washed with PBS and fixed with for 30 min in 0.0075%

Glutaraldehyde/4% PFA/DPBS. After DPBS washing, parasites were permeabilized with

0.1% TritonX100/PBS for 10-15 min before quenching free aldehyde groups with NH$_4$Cl

solution for 10 min. Next, parasites were blocked with 3% BSA–DPBS for 30 min. Primary

antibody incubation (1:500 dilution) lasted for 1 h before three washes with PBS, and

secondary antibody incubation for 30-60 min, Alexa Fluor 488-conjugated anti-mouse IgG

(Invitrogen) diluted 1:2,000 in 4% BSA–DPBS. After three final washes in DPBS, cells were

mounted in Vectashield containing DAPI for nuclear staining. Images were captured using a

Delta Vision Elite microscope (GE Healthcare). Image overlays were generated using Fiji

(Schindelin *et al*, 2012).

Co-Immunoprecipitation followed by western blot (Co-IP-WB)

PfMaf1-HA-ddFKBP tagged parasites were synchronized and split into control and MgCl$_2$

supplementation. At 18hpi, each culture (1.5×10^9 parasites) was centrifuged and RBCs were

lysed with six volumes of 0.15% saponin in DPBS for 5 min at 4°C. Parasites were

centrifuged at 4,000 *g* for 5 min at 4°C, and the pellet was washed twice with DPBS at 4°C.

The parasite pellets were then resuspended in 900uL of cytoplasmic lysis buffer (10 mM

Tris–HCl pH 7.5, 1 mM EDTA, 0.65% IGEPAL CA-630, 10mM NaCl) supplemented with

protease inhibitors (Thermo Fisher 78440) at 4°C and incubated with rotation for 30 min at

4°C. Extracts were centrifuged for 10 min at 2000g at 4°C and the cleared cytoplasmic

supernatant was removed and kept on ice. The nuclear pellet was resuspended in 200 µL

nuclear lysis buffer (10 mM Tris–HCl pH 7.5, 500 mM NaCl, 1 mM EDTA, 1% sodium

deoxycholate, 0.1% SDS, 1% IGEPAL CA-630, PI) at 4°C and transferred to 1.5 ml

sonication tubes (300 µL per tube, DiagenodeC30010016). Samples were sonicated for 5 min

(30 s on/off) in a Diagenode Pico Bioruptor at 4°C. Lysates were then centrifuged for 10 min

at 13,500g at 4°C and supernatant was transferred to a fresh tube. Cytoplasmic fractions were

mixed with 2:3 ratio of cytoplasmic dilution buffer (10 mM Tris-HCl pH 7.5, 150 mM NaCl,

0.5 mmM EDTA) and nuclear supernatants were mixed with 1:3 ratio of nuclear dilution

buffer (10 mM Tris-HCl pH 7.5, 0.5 mmM EDTA). Cytoplasmic and nuclear fraction

supernatants were incubated with 1ug of α-HA antibody (Abcam 9110) and 25uL Protein G

Magnetic Dynabeads (Invitrogen), pre-incubated for a minimum of 2 hours and washed twice

with dilution buffer, overnight with rotation at 4°C. The next day, the beads were collected on

a magnet and the supernatant was removed. While on the magnetic stand, beads were washed

twice with 500 μL wash buffer (10 mM Tris-HCl pH 7.5, 150 mM NaCl, 0.5 mM EDTA,

0.05% NP40), once with 25 mM NH_4HCO_3 (Sigma 09830) buffer, and then transferred to

new tube. Finally, the beads were resuspended in 20μl 2x SDS sample buffer and reducing

agent   before   5   min   at   95°C.

The beads were separated with magnet and the supernatant was transferred to a new tube.

Western blot protocol was followed as mentioned earlier.

Static cytoadhesion binding assay

Mature stage iRBCs for 3D7 control and supplementation of $MgCl_2$ ([3mM] total

concentration) for 1 cycle were used for cytoadhesion binding assays. Receptors diluted in

PBS (10 μg/mL target CD36 or negative control 1% BSA) were incubated overnight at 4°C in

labeled petri dishes. After the dishes were blocked for 30 minutes at 37°C with 1% BSA/PBS,

parasitemia was measured with a Guava easyCyte Flow Cytometer (EMD Millipore) after

trophozoites and schizonts iRBCs were isolated by plasmagel (Plasmion, Fresenius Kabi)

enrichment. Adjusted amounts of iRBCs were resuspended in binding medium (RPMI 1640

powder (Gibco 51800: W/L-Glutamine W/O $NaHCO_3$) with HEPES pH 7) for a

concentration of 2.2×10^8 iRBCs/mL. iRBCs/binding medium was added to each petri dish

and incubated for 2 hours at 37°C. After, unbound cells were removed and the dishes were

washed 5 times with binding medium by carefully tilting from side to side. Adherent iRBCs

were counted using 40x lens with a Nikon ECLIPSE TE200 in 10 randomly selected fields

(each with 0.2 μm^2) before 2 our fixation with 2% glutaraldehyde (G5882; Sigma)/PBS. After

fixation, giemsa staining was done to confirm percentage of bound iRBCs. A total of 3

biological replicates were performed and the results were expressed as the number of bound

iRBCs per 0.2 mm^2 of target receptor monolayer.

Statistical analysis

All statistical analyses were performed using GraphPad Prism version 9.1.0 (216) for Mac. To test for a normal distribution of the data, the Shapiro-Wilk normality test was used. To test for significance between two groups, a two-sided independent-samples t test was used. Gene ontology enrichments were calculated using the build-in tool at https://plasmoDB.org.

Estimation of cell cycle progression

RNA-seq-based cell cycle progression for control and $MgCl_2$ supplementation was estimated in R by comparing the normalized expression values (i.e., RPKM, reads per kilobase per exon per one million mapped reads) of each sample to the microarray data from Bozdech et al (2003) (Data ref: Bozdech et al, 2003) using the statistical model as in Lemieux et al (2009).

Data availability

The data generation in this study is available in the following database:

The mass spectrometry proteomics data have been deposited to the ProteomeXchange Consortium via the PRIDE (Perez-Riverol *et al*, 2022) partner repository with the dataset identifier PXD040576" (Username: reviewer_pxd040576@ebi.ac.uk & Password: wlw5Mlc6).

Acknowledgments

We thank L. Mancio Silva, E. Real and J. Bryant for critical reading this manuscript. This work was supported by the Laboratoire d'Excellence (LabEx) ParaFrap [ANR-11-LABX-0024], ERC AdG PlasmoSilencing (670301), ANR-JC (18-CE15-0009-01). We thank Camille Cohen and Marta Miera Maluenda for technical assistance with this project. With financial support from "la Région Île-de-France" (N°EX061034) and ITMO Cancer of Aviesan and INCa on funds administered by Inserm (N°21CQ016-00) for MS analysis.

Author contributions

Conceptualization A.S, GD; Methodology: GD, AClaës, AB, PBN; Investigation GD, AClaës, AB; Analysis: GD, AB, FD, DL; Writing: GD and A.S; Funding acquisition: A.S, AC

Conflict of interest

737 The funders had no role in the study design, data collection, and interpretation or decision to

738 submit the work for publication. The authors declare that they have no conflict of interest.

739
740
741

743
Figure 1. RNA Pol III-transcribed genes are downregulated in asymptomatic individuals during the dry
season. Steady state RNA levels as quantified by RT-qPCR using primers to tRNA Asparagine
(Pf3D7_0714700) (A) and tRNA Valine (Pf3D7_0312600) (B) as well as RNA Pol III-transcribed RUF6
ncRNA (C). DBLalpha primers were used to detect RNA Pol II-transcribed *var* genes (D). Normalized
expression is shown using fructose-bisphosphate aldolase (FBA Pf3D7_1444800) as the reference gene in
symptomatic individuals during the wet season (n = 14[+]) and asymptomatic individuals during the dry season (n
= 17[+]). [+] with the exception of Asparagine tRNA wet symptomatic (n=11) and dry asymptomatic (n=12), and
DBLalpha wet symptomatic (n = 12) and dry asymptomatic (n = 16). Box plots indicate the mean with standard
deviation. Wilcoxon matched-pairs signed rank test was done to determine significance (* indicates $P < 0.05$ and
** indicates $P < 0.005$).
754
755
756

757
758
Figure 2. External factors modulate RNA Pol III-transcribed genes. (A) Plasma magnesium levels are
significantly increased in asymptomatic individuals during the dry season compared to symptomatic individuals
during the wet season. Concentration is shown in nmol/μL (mM) (B) Schematic showing underlying molecular
mechanism summary using *in vitro* cultured *P. falciparum*. (C, D) Volcano plot showing log2 (fold change, FC)
against −log10 (p-value) of transcripts identified by RNA-seq analysis of 3D7 control and addition of $MgCl_2$
harvested during ring (C) and trophozoite stages (D). Expressed transcripts from three replicates between control
and addition of $MgCl_2$ that are significantly up-regulated are highlighted in red while significantly down-
regulated RNA Pol III genes are highlighted in blue (FDR corrected p-value of <0.05) and a FC≥±1.95) with
examples listed as text. Black dots indicate non-significant transcripts with a FC≤2.0.
768

769
770
Figure 3. Nuclear PfMaf1 is essential to regulate RNA Pol III. (A) Illustration of the TORC1-dependent
cellular localization of Maf1 protein in unfavorable conditions in typical eukaryotic organisms compared *P.*
falciparum that has no TORC1 pathway. (B, top) Illustration of recombinant PfMaf1 with a 3HA tag followed
by a ddFKBP domain to allow for knockdown studies. (B, bottom) Western blot analysis for PfMaf1 in total
extracts in pSLI-Maf1-FKBP transfected parasites at 24hpi after 1, 2, and 3 cycles without addition of Shield-1
(-) and control, with addition of Shield-1 (+). Aldolase levels are also shown. Representative of 3 replicates. (C)
Transcript levels as quantified by RT-qPCR using the same primers in (Figure 2B) in parasites harvested at
18hpi in control group and without Shield for 2 cycles. Error bars are displayed from 3 biological replicates.
Statistical significance was determined by two-tailed Student's t-test (** $p<0.005$). (D) Growth curve over 5
days of clonal pSLI-Maf1-FKBP parasites for 2 conditions: in the presence or absence of Shield-1. Uninfected
red blood cells ("blood" in red) serve as reference of background. Error bars indicate standard deviation of three
technical replicates in different blood from two different clones (n = 6). (E) Data is represented as box-whisker
plot of mean merozoite number per schizont ±SD (Mann-Whitney), with the with the median represented at the
center line. Boxplots show the data of 100 segmented schizonts counted per condition (n = 100). Statistical
significance was determined by two-tailed Student's t-test (*** $p<0.001$). Representative giemsa images are

shown to the right for + and – Shield. (F) Visual representation of Co-IP MS analysis of cytoplasmic and nuclear
PfMaf1. Labeled proteins represent important significant and unique proteins in cytoplasmic and nuclear
fractions not found in either of the controls.

809
810
Figure 4. External factors modulate virulence through PfMaf1-regulated RNA Pol III inhibition
(A) Immunoprecipitation western blot analysis for cytoplasmic and nuclear extracts for PfMaf1 expression in
PfMaf1-FKBP transfected parasites with addition of $MgCl_2$ ([3mM] total) and control parasites harvested at
18hpi. Controls are shown in Figure S4B. (B) Transcript levels as quantified by RT-qPCR using primers to
tRNA Valine (Pf3D7_0312600), RUF6 ncRNA, *var* DBLalpha, and normalized to FBA (Pf3D7_1444800) for 4
conditions: control (black), addition of $MgCl_2$ (green), KD of PfMaf1 (blue), and KD of PfMaf1 and addition of
$MgCl_2$ (orange). Error bars are displayed from 3 biological replicates. Statistical significance was determined by
two-tailed Student's t-test (** $p<0.005$). (C) Cytoadhesion binding assay data is represented as box-whisker plot
of mean number of bound iRBC ±SD (Mann-Whitney) mm^2, with the median represented at the center line.
Boxplots show the data of 3 biological replicates (n = 3). Statistical significance was determined by two-tailed
Student's t-test (**** $p<0.001$). Representative images are shown below for 3D7 and 3D7 +$MgCl_2$. (D) Western
blot analysis for extracts for ATS-PfEMP1 expression in 3D7 control parasites and with addition of $MgCl_2$
([3mM] total) harvested after plasmion. NapL control levels are also shown. Representative of 3 replicates. (E)
Schematic showing summary of study linking decreased cytoadherence, associated with disease severity, with
increased RNA Pol III-inhibition, triggered in response to external factors.
826
827
828
829

A

B

Figure S1. (A) Multiple sequence alignment of highly conserved genes encoding GC-rich ncRNA elements. 15 members of RUF6 gene family aligned by Clustal Omega (http://www.ebi.ac.uk) and presented in Jalview (http://www.jalview.org). Degree of conservation per base and consensus sequence are displayed below. Black lines show position of potential A- and B-box consensus motifs (as assessed in (Guizetti *et al*, 2016)). (B) RT-qPCR shows transcript levels for tRNA Asparagine, tRNA Valine, RUF6 ncRNA and a housekeeping gene encoding ubiquitin-conjugating enzyme in synchronized wildtype parasites at 24 hpi that were untreated or treated with RNA Pol III inhibitor or DMSO. Transcript levels are normalized to fructose-bisphosphate aldolase (PF3D7_1444800) transcript levels. Mean□SEM of two independent experiments are shown. Statistical significance was determined by two-tailed Student's t-test (* $p < 0.05$).

A

B

856
Figure S2. (A) tRNA Leucine (Pf3D7_API05400) and tRNA Asparagine (Pf3D7_API05500) transcripts
identified by RNA-Seq analysis from ((Andrade *et al*, 2020)) for wet symptomatic (n = 12) and dry
asymptomatic (n = 12) individuals. Statistical significance was determined by two-tailed Student's t-test (*
p<0.05). (B) Transcript levels as quantified by RT-qPCR on 3D7 parasites harvested during late ring stage for
control parasites, parasites in the absence of isoleucine (w/o Ile), at 40 degrees Celsius, at low-glucose levels
(0.5mg/mL), and presence of additional $MgCl_2$ ([3mM] total). Primers were used for Pol II-transcribed ubiquitin-
conjugating enzyme (UCE Pf3D7_0812600), Pol I-transcribed rRNA A1, Pol III-transcribed RUF6 ncRNA, Pol
III-transcribed tRNA Valine (Pf3D7_0312600), and Pol II-transcribed active *var* gene (Pf3D7_1240900).
Results are normalized to an RNA Pol II-transcribed reference gene FBA (Pf3D7_1444800) and presented as
fold change/control. Error bars are displayed from 3 biological replicates. Statistical significance was determined
by two-tailed Student's t-test (* p<0.05).

A

B

878
879
Figure S3. (A) Transcript levels as quantified by RT-qPCR on 3D7 parasites harvested during late ring stage for
control parasites and parasites in the presence of additional $MgCl_2$ levels ([1mM], [1.5mM], [2mM], and [3mM]
total). Primers were used for Pol II-transcribed ubiquitin-conjugating enzyme (UCE Pf3D7_0812600), Pol III-
transcribed RUF6 ncRNA, Pol III-transcribed tRNA Valine (Pf3D7_0312600) and Alanine (PF3D7_0411500),
and Pol II-transcribed *var* DBLalpha. Results are normalized to an RNA Pol II-transcribed reference gene FBA
(Pf3D7_1444800) and presented as fold change/control. Error bars are displayed from 3 biological replicates.
(B) Cell cycle progression estimation of a wildtype 3D7 clone in the absence ('C') or presence ('Mg') of $MgCl_2$
supplementation. RNA-seq data from synchronized parasites harvested at 12 and 24 hpi were compared to
microarray data (from (Bozdech *et al*, 2003) as in (Lemieux *et al*, 2009). Replicates are represented with circles.

A   Cytoplasmic PfMaf1 vs Cytoplasmic Control

B   Nuclear PfMaf1 vs Nuclear Control

C

Gene ID	Ab Cytoplasm/Control Cytoplasm						Total peptides in set	MW (kDa)	Description
	Ratio	Log2(Ratio)	Adj. p-value	Dist. pept. used	Pept. used	Distrib. dist. pep. in replic.			
PF3D7_1464600	2,263481181	1,178543312	1,577E-06	6	31	4, 5, 5, 4, 3/0, 4, 3, 2, 1	70	170,5	Serine/threonine protein phosphatase UIS2, putative
PF3D7_1127000	2,838513427	1,505135566	0,00160101	2	16	1, 2, 2, 2, 2/1, 0, 2, 2, 2	17	34,5	Protein phosphatase, putative

Gene ID	Ab Nuclear/Control Nuclear						Total peptides in set	MW (kDa)	Description
	Ratio	Log2(Ratio)	Adj. p-value	Dist. pept. used	Pept. used	Distrib. dist. pep. in replic.			
PF3D7_0507700	4,544185096	2,184021601	0,00072787	3	18	3, 2, 3, 2, 2/2, 0, 2, 0, 2	32	62,6	Nuclear protein localization protein 4, putative
PF3D7_1206600	1000	1000		6	13	2, 3, 3, 3, 2/0, 0, 0, 0, 0	13	167,3	DNA-directed RNA polymerase subunit beta
PF3D7_1329000	1000	1000		5	13	2, 3, 1, 3, 2/1, 1, 0, 0, 0	13	274,8	DNA-directed RNA polymerase subunit

D

898
899
900 **Figure S4. PfMaf1 cytoplasmic and nuclear interactome.** Co-IP MS volcano plot of enrichment for all 5
901 replicates for cytoplasmic (A) and nuclear (B) PfMaf1 vs control proteins are indicated and labeled. Each dot

represents a protein, and its size corresponds to the sum of peptides from both conditions used to quantify the ratio of enrichment. x-axis = $\log_2$(fold-change), y-axis = $-\log_{10}$(p-value), horizontal red line indicates adjusted p-value = 0.05, and vertical green lines indicate absolute fold-change = 2.0. Side panels indicate proteins uniquely identified in either sample (y-axis = number of peptides per 100 amino acids) with a minimum of 2 distinct peptides in 3 replicates of a same state. (C) Table showing values for significantly and uniquely enriched proteins from both extracts as labeled in (Figure 3F). (D) Volcano plot showing the distribution of significant and unique proteins in cytoplasmic and nuclear fractions not found in either of the controls. Each dot represents a protein, and its size corresponds to the sum of peptides from both conditions used to quantify the ratio of enrichment. x-axis = $\log_2$(fold-change), y-axis = $-\log_{10}$(p-value), horizontal red line indicates adjusted p-value = 0.05, and vertical green lines indicate absolute fold-change = 2.0. Side panels indicate proteins uniquely identified in either sample (y-axis = number of peptides per 100 amino acids) with a minimum of 2 distinct peptides in 3 replicates of a same state.

A

B

924
925
Figure S5. (A) Immunoprecipitation western blot analysis for cytoplasmic and nuclear extracts for PfMaf1
expression in PfMaf1-FKBP transfected parasites with addition of MgCl$_2$ ([3mM] total) and control parasites
harvested at 18hpi from Figure 4A. Anti-HA PfMaf1, aldolase and histone H3 are shown from input and flow-
through (FT). (B) Representative immunofluorescence images show brightfield, Dapi, anti-HA PfMaf1, and
Dapi-HA merge for PfMaf1 in control and addition of MgCl$_2$ ([3mM] total) in parasites harvested and fixed at
late ring stage.

932
933

934 **Supplementary Table 1. qPCR analysis primer pairs**
935

Primer pair	Forward	Reverse
Pool RUF6 A RUF6 B	5'-AAGCTGCCTCAGTAGCCCA-3'	5'-AAAAATTGCGCCACCCCC-3'
	5'-AAGCTGCCCCAGTAGCCCA-3'	5'-AAAAATTGCGCCGCCCCC-3'
rRNA A1 (from (Mancio-Silva *et al*, 2010))	5'-TGTTTTCTTTTTTCTAAGTTT-3'	5'-TCACCTCATTTGAAGCAA-3'
tRNA Alanine (PF3D7_0411500)	5'-GGGCAGGTGGTGTAGTGG-3'	5'-TGCTGGACAGACGGGGAATT-3'
tRNA Asparagine (PF3D7_0714700)	5'-GCAAGTATTTCCGCCTGTCA-3'	5'-GAATTGAACCCGGGTCTTCC-3'
tRNA Valine (Pf3D7_0312600)	5'-GCGGGCATGGTCTAGTGG-3'	5'-ACTACGGGCACCGAGGATC-3'
FBA fructose-bisphosphate aldolase (PF3D7_1444800)	5'-TGTACCACCAGCCTTACCAG-3'	5'-TTCCTTGCCATGTGTTCAAT-3'
UCE ubiquitin-conjugating enzyme (Pf3D_70812600)	5'-TAACAGCCCAGCGAATCAAG-3'	5'-CGGCATCTTCTTCAGCTTTCTG-3'
var gene (PF3D7_1240900)	5'-CAAAATGGTAGTGATGGTGGTCG-3'	5'-CCCCCTGCTTTATTATCTTTCGTC-3'
Pool *var* DBLalpha	5'-GCACGAACTTTTGCA-3'	5'-GCCCCATTCGTCGAACC-3'
	5'-GCACGCAGTTTTGCA-3'	5'-GCCCCATTCCTCGAACC-3'

936
937
938

939
940 References:
941

Abraham KJ, Chan JN, Salvi JS, Ho B, Hall A, Vidya E, Guo R, Killackey SA, Liu N, Lee
JE, *et al* (2016) Intersection of calorie restriction and magnesium in the suppression of
genome-destabilizing RNA-DNA hybrids. *Nucleic Acids Res Oct* 14;44(18):8870-
8884

Andrade CM, Fleckenstein H & Thomson-Luque R (2020) Increased circulation time of
Plasmodium falciparum underlies persistent asymptomatic infection in the dry season.
Nat Med 26: 1929–1940

Babbitt SE, Altenhofen L, Cobbold SA, Istvan ES, Fennell C, Doerig C, Llinás M &
Goldberg DE (2012) *Plasmodium falciparum* responds to amino acid starvation by
entering into a hibernatory state. *Proc Natl Acad Sci USA* 109

Barcons-Simon A, Cordon-Obras C, Guizetti J, Bryant JM & Scherf A (2020) CRISPR
Interference of a Clonally Variant GC-Rich Noncoding RNA Family Leads to General
Repression of var Genes in Plasmodium falciparum. *mBio* 11

Birnbaum J, Flemming S & Reichard N (2017) A genetic system to study Plasmodium
falciparum protein function. *Nat Methods* 14: 450–456

Bozdech Z, Llinás M, Pulliam BL, Wong ED, Zhu J & DeRisi JL (2003) The Transcriptome
of the Intraerythrocytic Developmental Cycle of Plasmodium falciparum. *PLoS Biol* 1:
e5

Collins KA, Ceesay S, Drammeh S, Jaiteh FK, Guery MA, Lanke K, Grignard L, Stone W,
Conway DJ, D'Alessandro U, *et al* (2022) A Cohort Study on the Duration of
Plasmodium falciparum Infections During the Dry Season in The Gambia. *The*
Journal of Infectious Diseases 226: 128–137

Crabb BS, Cooke BM, Reeder JC, Waller RF, Caruana SR, Davern KM, Wickham ME,
Brown GV, Coppel RL & Cowman AF (1997) Targeted Gene Disruption Shows That
Knobs Enable Malaria-Infected Red Cells to Cytoadhere under Physiological Shear
Stress. *Cell* 89: 287–296

van Dam TJP, Zwartkruis FJT, Bos JL & Snel B (2011) Evolution of the TOR Pathway. *J*
Mol Evol 73: 209–220

Dawn A, Singh S, More KR, Siddiqui FA, Pachikara N, Ramdani G, Langsley G & Chitnis
CE (2014) The Central Role of cAMP in Regulating Plasmodium falciparum
Merozoite Invasion of Human Erythrocytes. *PLoS Pathog* 10: e1004520

Diffendall GM, Barcons-Simon A, Baumgarten S, Dingli F, Loew D & Scherf A (2022)
Discovery of RUF6 ncRNA–interacting proteins involved in *P. falciparum* immune
evasion. *Life Sci Alliance* 6: e202201577

Fan Y, Shen S, Wei G, Tang J, Zhao Y, Wang F, He X, Guo G, Shang X, Yu X, *et al* (2020)
Rrp6 Regulates Heterochromatic Gene Silencing via ncRNA RUF6 Decay in Malaria
Parasites. *mBio* 11: e01110-20

Fang J, Sullivan M & McCutchan TF (2004) The Effects of Glucose Concentration on the
Reciprocal Regulation of rRNA Promoters in Plasmodium falciparum. *Journal of*
Biological Chemistry 279: 720–725

Fogang B, Lellouche L, Ceesay S, Drammeh S, Jaiteh FK, Guery M-A, Landier J, Haanappel
C, Froberg J, Conway D, *et al* (2023) Asymptomatic *Plasmodium falciparum* Carriage
at the End of the Dry Season is Associated with Subsequent Infection and Clinical
Malaria in Eastern Gambia Epidemiology

Gardner MJ, Hall N, Fung E, White O, Berriman M, Hyman RW, Carlton JM, Pain A, Nelson
KE, Bowman S, *et al* (2002) Genome sequence of the human malaria parasite
Plasmodium falciparum. *Nature* 419: 498–511

Guillochon E, Fraering J, Joste V, Kamaliddin C, Vianou B, Houzé L, Baudrin LG, Faucher
JF, Aubouy A, Houzé S, *et al* (2022) Transcriptome Analysis of *Plasmodium*
falciparum Isolates From Benin Reveals Specific Gene Expression Associated With
Cerebral Malaria. *The Journal of Infectious Diseases* 225: 2187–2196

Guizetti J, Barcons-Simon A & Scherf A (2016) Trans-acting GC-rich non-coding RNA at
var expression site modulates gene counting in malaria parasite. *Nucleic Acids Res* 44:
9710–9718

Guizetti J & Scherf A (2013) Silence, activate, poise and switch! Mechanisms of antigenic
variation in Plasmodium falciparum. *Cell Microbiology* 15: 718–726

Hasenkamp S, Merrick CJ & Horrocks P (2013) A quantitative analysis of Plasmodium
falciparum transfection using DNA-loaded erythrocytes. *Mol Biochem Parasitol* 187:
117–120

Hess FI, Kilian A, Söllner W, Nothdurft HD, Pröll S & Löscher T (1995) Plasmodium
falciparum and Plasmodium berghei: Effect of Magnesium on the Development of
Parasitemia. *Experimental Parasitology* 80: 196–193

Innocent O, Ejovi O & Charles E (2013) Levels of iron and magnesium in serum of
Plasmodium falciparum malarial infected children in Abraka, Delta State, Nigeria. *J*
Invest Biochem 2: 62

Jahnen-Dechent W & Ketteler M (2012) Magnesium basics. *Clinical Kidney Journal* 5: i3–
i14

Jensen MD, Conley M & Helstowski LD (1983) Culture of Plasmodium falciparum: The Role
of pH, Glucose, and Lactate. *The Journal of Parasitology* 69: 1060

Kimenyi KM, Wamae K & Ochola-Oyier LI (2019) Understanding P. falciparum
Asymptomatic Infections: A Proposition for a Transcriptomic Approach. *Front*
Immunol 10: 2398

Kumar M, Skillman K & Duraisingh MT (2021) Linking nutrient sensing and gene
expression in *Plasmodium falciparum* blood-stage parasites. *Mol Microbiol* 115: 891–
900

Leech JH, Barnwell JW, Miller LH & Howard RJ (1984) Identification of a strain-specific
malarial antigen exposed on the surface of plasmodium falciparum-infected
erythrocytes. *J Exp Med* 159: 1567–1575

Lemieux JE, Gomez-Escobar N, Feller A, Carret C, Amambua-Ngwa A, Pinches R, Day F,
Kyes SA, Conway DJ, Holmes CC, *et al* (2009) Statistical estimation of cell-cycle
progression and lineage commitment in *Plasmodium falciparum* reveals a
homogeneous pattern of transcription in ex vivo culture. *Proc Natl Acad Sci USA* 106:
7559–7564

Leopold SJ, Apinan S, Ghose A, Kingston HW, Plewes KA, Hossain A, Dutta AK, Paul S,
Barua A, Sattar A, *et al* (2019) Amino acid derangements in adults with severe
falciparum malaria. *Sci Rep* 9: 6602

Li H & Durbin R (2009) Fast and accurate short read alignment with BurrowsWheeler
transform. *Bioinformatics* 25: 1754–1760

Lopez-Rubio JJ, Mancio-Silva L & Scherf A (2009) Genome-wide analysis of
heterochromatin associates clonally variant gene regulation with perinuclear
repressive centers in malaria parasites. *Cell Host Microbe* 5: 179–190

Mancio-Silva L, Slavic K, Grilo Ruivo MT, Grosso AR, Modrzynska KK, Vera IM, Sales-
Dias J, Gomes AR, MacPherson CR, Crozet P, *et al* (2017a) Nutrient sensing
modulates malaria parasite virulence. *Nature* 547: 213–216

Mancio-Silva L, Slavic K, Grilo Ruivo MT, Grosso AR, Modrzynska KK, Vera IM, Sales-
Dias J, Gomes AR, MacPherson CR, Crozet P, *et al* (2017b) Nutrient sensing
modulates malaria parasite virulence. *Nature* 547: 213–216

Mancio-Silva L, Zhang Q, Scheidig-Benatar C & Scherf A (2010) Clustering of dispersed
ribosomal DNA and its role in gene regulation and chromosome-end associations in
malaria parasites. *Proc Natl Acad Sci USA* 107: 15117–15122

Marreiros IM, Marques S, Parreira A, Mastrodomenico V, Mounce BC, Harris CT, Kafsack
BF, Billker O, Zuzarte-Luís V & Mota MM (2023) A non-canonical sensing pathway
mediates Plasmodium adaptation to amino acid deficiency. *Commun Biol* 6: 205

McLean KJ & Jacobs-Lorena M (2017) Plasmodium falciparum Maf1 Confers Survival upon
Amino Acid Starvation. *mBio* 8: 02317–16

Michels AA, Robitaille AM, Buczynski-Ruchonnet D, Hodroj W, Reina JH, Hall MN &
Hernandez N (2010) mTORC1 Directly Phosphorylates and Regulates Human MAF1.
Mol Cell Biol 30: 3749–3757

Miller LH, Ackerman HC, Su X & Wellems TE (2013) Malaria biology and disease
pathogenesis: insights for new treatments. *Nat Med* 19: 156–167

Miller LH, Baruch DI, Marsh K & Doumbo OK (2002) The pathogenic basis of malaria.
Nature 415: 673–679

M'Kaibi FK, Steyn NP, Ochola S & Du Plessis L (2015) Effects of agricultural biodiversity
and seasonal rain on dietary adequacy and household food security in rural areas of
Kenya. *BMC Public Health* 15: 422

Murray L, Stewart LB, Tarr SJ, Ahouidi AD, Diakite M, Amambua-Ngwa A & Conway DJ
(2017) Multiplication rate variation in the human malaria parasite Plasmodium
falciparum. *Sci Rep* 7: 6436

Nacer A, Claes A, Roberts A, Scheidig-Benatar C, Sakamoto H, Ghorbal M, Lopez-Rubio J
& Mattei D (2015) Discovery of a novel and conserved Plasmodium falciparum
exported protein that is important for adhesion of PfEMP1 at the surface of infected
erythrocytes. *Cell Microbiol* 17: 1205–1216

Nyarko PB & Claessens A (2021) Understanding Host–Pathogen–Vector Interactions with
Chronic Asymptomatic Malaria Infections. *Trends in Parasitology* 37: 195–204

Perez-Riverol Y, Bai J, Bandla C, García-Seisdedos D, Hewapathirana S, Kamatchinathan S,
Kundu DJ, Prakash A, Frericks-Zipper A, Eisenacher M, *et al* (2022) The PRIDE
database resources in 2022: a hub for mass spectrometry-based proteomics evidences.
Nucleic Acids Research 50: D543–D552

Pluta K, Lefebvre O, Martin NC, Smagowicz WJ, Stanford DR, Ellis H SR, AK S, A B, & M.
(2001) Maf1p, a negative effector of RNA polymerase III in Saccharomyces
cerevisiae. *Mol Cell Biol* 21: 5031–5040

Poullet P, Carpentier S & Barillot E (2007) myProMS, a web server for management and
validation of mass spectrometry-based proteomic data. *Proteomics* 7: 2553–2556

Rollins J, Veras I, Cabarcas S, Willis I & Schramm L (2007) Human Maf1 negatively
regulates RNA Polymerase III transcription via the TFIIB family members Brf1 and
Brf2. *Int J Biol Sci*: 292–302

Scherf A, Hernandez-Rivas R, Buffet P, Bottius E, Benatar C, Pouvelle B, Gysin J & Lanzer
M (1998) Antigenic variation in malaria: in situ switching, relaxed and mutually
exclusive transcription of var genes during intra-erythrocytic development in
Plasmodium falciparum. *EMBO J* 17: 5418–5426

Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T, Preibisch S,
Rueden C, Saalfeld S, Schmid B, *et al* (2012) Fiji: an open-source platform for
biological-image analysis. *Nat Methods* 9: 676–682

Schneider VM, Visone JE, Harris CT, Florini F, Hadjimichael E, Zhang X, Gross MR, Rhee
KY, Ben Mamoun C, Kafsack BFC, *et al* (2023) The human malaria parasite
Plasmodium falciparum can sense environmental changes and respond by antigenic
switching. *Proc Natl Acad Sci USA* 120: e2302152120

Serfontein J, Nisbet RER, Howe CJ & de Vries PJ (2010) Evolution of the TSC1/TSC2-TOR
Signaling Pathway. *Sci Signal* 3

Smith JD, Chitnis CE, Craig AG, Roberts DJ, Hudson-Taylor DE, Peterson DSP, R. N, I. C &
Miller LH (1995) Switches in expression of plasmodium falciparum var genes

correlate with changes in antigenic and cytoadherent phenotypes of infected
erythrocytes. *Cell* 82: 101–110

Smith JD, Rowe JA, Higgins MK & Lavstsen T (2013) Malaria's deadly grip: cytoadhesion
of Plasmodium falciparum-infected erythrocytes: Plasmodium falciparum
cytoadhesion and var genes. *Cell Microbiol* 15: 1976–1983

The M, MacCoss MJ, Noble WS & Käll L (2016) Fast and Accurate Protein False Discovery
Rates on Large-Scale Proteomics Data Sets with Percolator 3.0. *J Am Soc Mass*
Spectrom 27: 1719–1727

Thomson-Luque R, Votborg-Novél L, Ndovie W, Andrade CM, Niangaly M, Attipa C, Lima
NF, Coulibaly D, Doumtabe D, Guindo B, *et al* (2021) Plasmodium falciparum
transcription in different clinical presentations of malaria associates with circulation
time of infected erythrocytes. *Nat Commun* 12: 4711

Valot B, Langella O, Nano E & Zivy M (2011) MassChroQ: a versatile tool for mass
spectrometry quantification. *Proteomics* 11: 3572–3577

Vannini A, Ringel R, Kusser AG, Berninghausen O, Kassavetis GA & Cramer P (2010)
Molecular basis of RNA polymerase III transcription repression by Maf1. *Cell* 143: 70

Wahlgren M, Goel S & Akhouri RR (2017) Variant surface antigens of Plasmodium
falciparum and their roles in severe malaria. *Nat Rev Microbiol*

Wullschleger S, Loewith R & Hall MN (2006) TOR Signaling in Growth and Metabolism.
Cell 124: 471–484

Zhang M, Wang C, Otto TD, Oberstaller J, Liao X, Adapa U SR, K B, IF C, D M, M B, *et al*
(2018) Uncovering the essential genes of the human malaria parasite Plasmodium
falciparum by saturation mutagenesis. *Science*

Zhang Q, Siegel TN, Martins RM, Wang F, Cao J, Gao Q, Cheng X, Jiang L, Hon CC,
Scheidig-Benatar C, *et al* (2014) Exonuclease-mediated degradation of nascent RNA
silences genes linked to severe malaria. *Nature* 513: 431–435

Zhang X & Deitsch KW (2022) The mystery of persistent, asymptomatic Plasmodium
falciparum infections. *Current Opinion in Microbiology* 70: 102231

1121

1122
1123
1124