RNA polymerase III is involved in regulating *Plasmodium falciparum* virulence

Gretchen Diffendall,[1,2] Aurélie Claës,[1] Anna Barcons-Simon,[1,2, 3] Prince Nyarko,[4] Florent Dingli,[5] Damarys Loew,[5] Antoine Claessens,[4,6] and Artur Scherf [1, *]

Institut Pasteur, [1]Universite Paris Cité, INSERM U1201, CNRS EMR9195, Paris, France.
[2]Sorbonne Université Ecole doctorale Complexité du Vivant ED515, Paris, France.
[3]Biomedical Center, Division of Physiological Chemistry, Faculty of Medicine, Ludwig-Maximilians-Universität München, Munich 82152, Germany
[4] Laboratory of Pathogen–Host Interaction (LPHI), CNRS, University of Montpellier, France
[5]Institut Curie, PSL Research University, Centre de Recherche, CurieCoreTech Mass Spectrometry Proteomics, Paris, France
[6] LPHI, MIVEGEC, CNRS, INSERM, University of Montpellier, France

*Corresponding author: artur.scherf@pasteur.fr

Summary:

Malaria pathogenesis is linked to parasite sequestration in critical target organs and is regulated seasonally in malaria-endemic regions, suggesting environmental sensing might regulate disease severity. How this occurs at the molecular level and the contributing host and parasite factors remain unclear. Here, we report that *P. falciparum* RNA polymerase III (RNA Pol III) transcription is downregulated in field isolates from asymptomatic patients. By using *in vitro* cultured parasites, we identified an RNA Pol III-dependent mechanism that governs expression of a major virulence factor in response to external stimuli. We establish a link between *P. falciparum* cytoadhesion and a Pol III-transcribed non-coding RNA family. Furthermore, we identify *P. falciparum* Maf1 as a critical regulator of Pol III transcription for cellular homeostasis and in adaptive responsive to external stimuli. Our results identify an RNA Pol III-dependent reduction in parasite cytoadherence capacity and link this process to asymptomatic malaria infection.

Introduction:

The parasite *Plasmodium falciparum* is responsible for the deadliest form of human malaria that annually affects over 200 million people, with 619,000 fatal cases, majorly African children under the age of 5 (W.H.O., 2022). The disease is transmitted to its human host during a blood meal by the parasites' vector, *Anopheles* mosquitoes. Disease symptoms are seen while the parasite multiplies asexually within the host RBCs. During this time, variant surface antigens (VSAs) are exported and displayed on the surface of the iRBCs (Wahlgren *et al*, 2017). During

the ~48-hour asexual intraerythrocytic developmental cycle (IDC), the parasite develops through different morphological stages: ring (circulating), trophozoite, and schizont (sequestered in capillaries). Expression of one type of VSA that is linked to immune evasion and pathogenesis, termed *P. falciparum* erythrocyte membrane protein 1 (PfEMP1) (Leech *et al*, 1984), can bind to a wide range of receptors on endothelial cells such as CD36. PfEMP1 binding mediates adhesion to the vascular endothelium within the host, thereby preventing mature iRBCs from traveling to, and being cleared by, the spleen (Smith *et al*, 2013). *Plasmodium* parasites replicate via schizogony inside the host RBCs generating the infectious form, termed merozoites. Upon bursting of the iRBC, merozoites invade new RBCs, thus enabling the cycle to restart. The number of merozoites per schizont has been shown to reduce in unfavorable conditions as a way to diminish the total parasite load and therefore overall disease severity (Mancio-Silva *et al*, 2017b).

P. falciparum chronic infection relies largely on mutually exclusive expression of PfEMP1 surface adhesins, that are encoded by the *var* virulence gene family (Scherf *et al*, 1998; Smith *et al*, 1995; Guizetti & Scherf, 2013). The expression of *var* genes, while under tight epigenetic regulation by the parasite, is additionally controlled by a family of ncRNA, termed RUF6. *P. falciparum* 3D7 encodes 15 RUF6 genes dispersed over several chromosomes located adjacent to central *var* genes (Gardner *et al*, 2002). RUF6 ncRNA have been observed to associate with the active *var* gene in *trans* and transcriptional repression disrupts the monoallelic expression of *var* genes by down-regulating the entire *var* gene family (Barcons-Simon *et al*, 2020; Guizetti *et al*, 2016). Genetically modified parasites that express the entire RUF6 gene family show a general upregulation of *var* genes (Zhang *et al*, 2014; Fan *et al*, 2020).

P. falciparum virulence is linked to cytoadhesion and sequestration as a result of reduced clearance of iRBCs in the spleen and increased microcapillary obstruction and local inflammation (Miller *et al*, 2002). Substantial sequestration of iRBCs in critical target organs has been associated with severe malaria (Miller *et al*, 2013; Guillochon *et al*, 2022). In long-term asymptomatic infections, found during prolonged low transmission dry seasons in malaria-endemic regions, decreased cytoadhesion and longer circulation time of iRBCs in the blood has been observed (Andrade *et al*, 2020). It is not fully understood how asymptomatic infections develop and are maintained nor is a molecular understanding of host and parasite factors, that underlies disease severity during the dry and wet seasons (reviewed in (Zhang & Deitsch, 2022)). Disease severity appears to be influenced by a variety of factors derived from the human

host, parasite, and environment. *Plasmodium* parasites are auxotrophic for various nutrients including glucose, certain types of amino acids, and lipids, which must be obtained from the host. Physiological conditions in the host, like diabetes, pregnancy, diet, and immunity can change the metabolic environment the parasites are exposed to. Numerous studies conclude that host immunity is a contributing factor to the maintenance of asymptomatic infections (Kimenyi *et al*, 2019). While it is proposed that *P. falciparum* virulence regulation contributes to decreased endothelial binding, the mechanism and factors involved remains to be determined.

Almost all eukaryotes use conserved nutrient sensing pathways, including mTOR, GCN2, GCN5, Rgt2/Snf3, and Snf1/AMPK signaling cascades (reviewed in (Kumar *et al*, 2021)). Most notably, the target of rapamycin complex (TORC) pathway involves the signaling of external factors to control many cellular processes (Wullschleger *et al*, 2006). Specifically, TORC participates in the regulation of RNA Pol III activity via the phosphorylation of its inhibitor, Maf1. When nutrients are available and mTOR kinase is active, Maf1 is hyperphosphorylated resulting in an inactive cytoplasmic state. Stress-induced Maf1 dephosphorylation results in nuclear localization and inhibition of RNA Pol III (Pluta *et al*, 2001; Rollins *et al*, 2007; Vannini *et al*, 2010). This pathway responds to a range of positive and negative external stimuli, most notably the presence of amino acids, that drive or inhibit cellular growth. The majority of the TORC pathway components are lost in the apicomplexan lineage (McLean & Jacobs-Lorena, 2017; Serfontein *et al*, 2010; van Dam *et al*, 2011). While *P. falciparum* encodes none of the core TORC1 components, it does contain a homologue to Maf1 (McLean & Jacobs-Lorena, 2017). The biology of *P. falciparum* Pf-Maf1 is still poorly characterized.

Here, we present data showing inhibition of RNA Pol III transcription in field isolates from asymptomatic individuals during the dry season. We present evidence for *P. falciparum* Maf1 as a negative regulator of RNA Pol III-transcribed genes including tRNA and RUF6 ncRNA, and demonstrate that this mechanism is responsive to external stimuli using *in vitro* culture. Inhibition of *P. falciparum* cytoadherence, via the regulation of RNA Pol III activity, represents a new paradigm contributing to *P. falciparum* virulence and links this process to asymptomatic malaria parasite persistence.

Results:

RNA Pol III-transcribed genes are downregulated in asymptomatic individuals

We and others have previously established the role of the 15-member RUF6 ncRNA gene
family in *P. falciparum var* gene expression (Barcons-Simon *et al*, 2020; Diffendall *et al*, 2022;
Fan *et al*, 2020; Guizetti *et al*, 2016). RUF6 ncRNA protein complex associates with the active
var gene expression site (Diffendall *et al*, 2022; Guizetti *et al*, 2016), and knockdown of the
ncRNA RUF6 gene family dramatically reduced *var* gene transcription (Barcons-Simon *et al*,
2020). Bioinformatic analysis identified highly conserved RNA Pol III binding elements
(conserved A- and B-box sequence) within RUF6 genes (Guizetti *et al*, 2016) (Figure S1A),
hinting at a role for RNA Pol III in virulence gene regulation. Thus, we determined the effects
of RNA Pol III inhibition on gene transcription in synchronized wild type parasites. We
observed downregulation of tRNA and RUF6 transcription after 20 hours of treatment with a
commercially available RNA Pol III inhibitor (Figure S1B). As expected, no changes were
observed in the transcription of an RNA Pol II-regulated gene, *uce* (ubiquitin-conjugating
enzyme, PF3D7_0812600) (Figure S1B). As predicted from the RUF6 sequence, our data
provide direct experimental evidence for RNA Pol III-dependent transcription of RUF6 genes.
116
We next wanted to determine if RNA Pol III-transcribed genes, specifically RUF6 ncRNA,
varied between parasites causing different clinical states of malaria infection, since previous
studies did not include RNA Pol III-transcribed genes for their data analysis. We first re-
analyzed raw RNA-seq data, from a previous study that compared parasites infecting
individuals during the dry versus wet season in a malaria endemic region of Mali (Andrade *et*
al, 2020), to include these genes. We were able to retrieve transcriptional data for two RNA Pol
III-transcribed tRNAs (tRNA Leucine Pf3D7_API05400 and tRNA Asparagine
Pf3D7_API05500), both of which show significantly lower levels in parasites infecting
asymptomatic individuals during the dry season (n = 12) when compared to parasites infecting
symptomatic individuals during the wet season (n = 12).
127
We further investigated whether RNA Pol III modulates gene transcription in field isolates of
P. falciparum from different individuals infected during the wet and dry season in malaria
endemic regions. In this pilot study, we analyzed *P. falciparum* steady state levels of RNA in
venous blood samples taken from individuals with asymptomatic parasitemia during the dry
season (n = 17) and wet season (n = 12), as well as mildly symptomatic malaria infections
during the wet season (n = 14) in The Gambia (Collins *et al*, 2022). RT-qPCR of total RNA
showed that levels of tRNAs (Asparagine and Valine) were significantly lower (p < 0.05) in
parasites from asymptomatic infections compared to parasites causing symptomatic infections

during the wet season (Figure 1A, B). Likewise, RUF6 ncRNA was significantly downregulated ($p < 0.005$) in parasites from asymptomatic infections (dry season) when compared to symptomatic infections (wet season) (Figure 1C). Expression was normalized to a housekeeping gene encoding fructose bisphosphate aldolase (Pf3D7_1444800).

The same previous study, mentioned above, also showed that *var* gene transcription varied in parasites infecting individuals during the dry versus wet seasons (Andrade *et al*, 2020). We observed a significantly lower level of *var* gene transcription ($p < 0.005$) in parasites taken from asymptomatic individuals during the dry season compared to those taken from symptomatic individuals during the wet season (Figure 1D). The observed lower transcript levels of *var* genes may be a direct cause of reduced RUF6 ncRNA levels. Taken together, our results reveal that levels of RNA Pol III-transcribed genes, particularly the regulatory RUF6 ncRNA, is significantly different between parasites causing two different clinical states.

Environmental factors regulate RNA Pol III-transcribed genes

Having established that RUF6 ncRNA decreases in parasite samples from asymptomatic individuals (Figure 1C), we sought to explore the underlying molecular mechanism, using *in vitro* cultured *P. falciparum* asexual blood stage parasites (Figure 2A). We mimicked nutrient deprivation with isoleucine-deprived culture medium (as was described in (Babbitt *et al*, 2012; McLean & Jacobs-Lorena, 2017)), fever conditions with incubation at 40ºC, and food starvation with low glucose levels (Jensen *et al*, 1983; Fang *et al*, 2004). Additionally, we used magnesium chloride ($MgCl_2$) supplementation, an abundant cation involved in signaling pathways and essential biological processes including protein synthesis and regulating blood glucose levels with varying levels in malaria-infected individuals (Oluboyo *et al*, 2018; Kanellopoulou *et al*, 2019). We evaluated the steady state levels of RNA Pol I- (rRNA A1), RNA Pol II- (UCE and the active *var*), and RNA Pol III- (tRNA Valine and RUF6 ncRNA) transcribed genes in clonal parasites at late ring stage. We observed that high temperature and low glucose conditions affected gene transcription of RNA Pol I, II and Pol III (Figure 2B). Deprivation of isoleucine and $MgCl_2$ supplementation seemingly affected only RNA Pol III-transcribed genes with the exception of the RNA Pol II-transcribed active *var* gene, Pf3D7_1240900, likely as a result of decreased RUF6 ncRNA as was reported earlier (Barcons-Simon *et al*, 2020).

To explore the underlying molecular pathway that leads to the downregulation of RNA Pol III activity in malaria parasites, we continued using $MgCl_2$ supplementation. RNA-seq was done

on control and +MgCl$_2$ cultured parasites during two time points of the asexual blood stage parasites (ring and trophozoite) to further confirm that RNA Pol III is the main target (Figure 2C, D). RNA-seq data was compared to the microarray time course data in (Bozdech *et al*, 2003) as in (Lemieux *et al*, 2009), which provided a statistical estimation of cell cycle progression at 12 and 24 hpi in both control and MgCl$_2$ supplemented parasites (Figure S2B). These data indicate that any differences in transcription were due to MgCl$_2$ supplementation and not differences in cell cycle progression. During ring stage (Figure 2C), 77% of the down-regulated genes contain typical RNA Pol III promoter sequences, A- and B-box consensus (from Figure S1A). A total of 27 ncRNA genes were downregulated (logFC < 1.95, FDR < 0.05) including all 15 RUF6 ncRNA, 6 tRNAs, 1 spliceosomal RNA, and 4 snoRNA. Additionally, 12 protein coding genes were up-regulated with no conserved pathway nor molecular or biological process. During trophozoite stage (Figure 2D), 74% of the down-regulated genes have typical RNA Pol III promoter sequences. This demonstrates that MgCl$_2$ supplementation primarily inhibits RNA Pol III-transcribed genes, including the entire RUF6 gene family.

Nuclear PfMaf1 protein is essential to regulate RNA Pol III

In almost all eukaryotes, the target of rapamycin complex (TORC) pathway has been shown to participate in the regulation of RNA Pol III activity via the phosphorylation of its only known inhibitor, Maf1 (Figure 3A, reviewed in (Wullschleger *et al*, 2006)). Upon phosphatase activation, modulated in response to external stimuli such as low nutrient availability, Maf1 shuttles to the nucleus to repress RNA Pol III activity (Michels *et al*, 2010). Because Maf1 is conserved in the apicomplexan lineage, a predicted essential blood stage protein in *P. falciparum*, we set out to explore if *P. falciparum* Maf1 (PfMaf1) functions similarly to model eukaryotes. We generated a PfMaf1 parasite line tagged with a 3HA tag and a ligand-controlled destabilization domain (ddFKBP), PfMaf1-FKBP (Figure 3B top). PfMaf1-FKBP transfected parasite clonal populations were validated for proper integration and confirmation that the system was working, shown by western blot for addition and removal of the ligand, Shield-1, over the course of 3 cycles (Figure 3B bottom). We observed that upon removal of Shield-1, PfMaf1 levels were shown to decrease by approximately 57% in total extracts after one cycle and almost complete degradation was achieved after 2 cycles.

Plasmodial Maf1 seemingly affects asexual blood stage proliferation making classical gene KO experiments unattainable (McLean & Jacobs-Lorena, 2017; Zhang *et al*, 2018), pointing to a

regulatory role of the putative RNA Pol III inhibitor under normal *in vitro* culture growth
conditions. Once PfMaf1 is depleted (Figure 3B bottom, 3 cycles -Shield), transcription of
tRNA Valine and RUF6 ncRNA increased dramatically by 3-fold (Figure 3C), possibly
perturbing cellular homeostasis. RNA Pol II-transcribed genes were unaffected. Parasite growth
was assessed over the course of 5 days for two PfMaf1-FKBP clones cultured in the presence
or absence of PfMaf1 (+/- Shield-1, removed one cycle before day 0, Figure 3D). A significant
difference in growth was achieved, most evidently in parasites cultured without Shield for 3
cycles, consistent with transposon mutagenesis scoring PfMaf1 essential (Zhang *et al*, 2018).
Dead parasites were not detected by Giemsa staining even up to 8 cycles without Shield-1. We
confirmed that differences in merozoite numbers per schizont contributed to the change in
growth rate between the control and KD PfMaf1 parasites after the 3^{rd} cycle KD (Figure 3E).
215
Additionally, we confirmed that nuclear PfMaf1 interreacts with the RNA Pol III protein
complex by performing co-immunoprecipitation followed by quantitative mass spectrometry
(Co-IP MS) of cytoplasmic and nuclear fractions of PfMaf1-HA-ddFKBP (Figure 3F, Figure
S3). Each fraction was split to include a control with no antibody added. Analysis of the
quantitative mass spectrometry data revealed unique interactome of cytoplasmic PfMaf1 and
nuclear PfMaf1 in their respective samples compared to their controls (Figure S3A, B). Gene
Ontology (GO) analysis showed that PfMaf1-associated proteins uniquely found in the nuclear
fraction, and neither of the two controls nor cytoplasmic fraction, are significantly represented
by the biological function category of "RNA Polymerase III transcription" ($p = 0.00175$) with
two subunits of RNA Pol III (Pf3D7_1206600 and Pf3D7_1329000). (GO) analysis of
significant and unique proteins found only in the cytoplasmic fractions, and not nuclear, are
significantly represented by the molecular function category of "protein kinase activator" ($p =
0.0143$, Pf3D7_1103100). Additionally, two protein phosphatases (Pf3D7_1127000 and
Pf3D7_1464600) were significantly enriched in the cytoplasmic fraction (Figure S3C).
230
231 While previous attempts to achieve a KO line were unsuccessful (McLean & Jacobs-Lorena,
232 2017), here we have generated an inducible PfMaf1 protein degradation system to highlight its
233 housekeeping role in maintaining RNA Pol III transcription at levels to maintain optimal
234 parasite proliferation.
235
MgCl₂ modulates cytoadhesion through PfMaf1-regulated RNA Pol III transcription

We further investigated if nuclear PfMaf1 activity could be regulated, in response to external

factors, to lead to changes in RNA Pol III activity using *P. falciparum* wildtype parasites

cultured *in vitro*. The *P. falciparum* cellular localization of PfMaf1 was investigated in response

to external stimuli ($+MgCl_2$). Parasites were tightly synchronized, split into control, addition of

$MgCl_2$ and harvested during late ring stage. In contrast to several eukaryotic model systems,

under normal culture conditions we observed PfMaf1 in cytoplasmic and nuclear fractions

using immunoprecipitation followed by western blot analysis (Figure 4A, S4A). Nuclear

PfMaf1 levels increased upon addition of $MgCl_2$ compared to control nuclear PfMaf1 (Figure

4A). Immunofluorescence (IF) assays showed that PfMaf1 forms foci-like aggregates in the

cytoplasm and near the nuclear periphery in both culture conditions (Figure S4B).

Next, we used the PfMaf1-FKBP transfected parasite lines, to show that the downregulation of

RNA Pol III-transcribed genes, triggered by external stimuli, is dependent on PfMaf1. We used

the PfMaf1 KD parasites (2^{nd} cycle, see Figure 3B) in control conditions and $MgCl_2$

supplementation to the growth medium in ring stage parasites for RT-qPCR (Figure 4B). We

confirmed the previously observed down-regulation of RNA Pol III-transcribed tRNA Valine

and RUF6 upon $MgCl_2$ supplementation in two independent PfMaf1-FKBP parasite clones.

Notably, when PfMaf1 degradation was induced, the addition of $MgCl_2$ did not affect Pol III

transcription. These results demonstrate that PfMaf1 mediates the observed decrease in RNA

Pol III-transcribed genes with $MgCl_2$ supplementation.

Finally, we set out to link *P. falciparum* virulence gene expression with decreased RUF6

transcription, by investigating the effect of $MgCl_2$ supplementation on cytoadherence of 3D7

iRBCs. We found that in static binding assays to the endothelial receptor CD36, bound to plastic

dishes, adhesion of iRBCs was reduced by 50% with $MgCl_2$ supplementation (Figure 4C). This

was also confirmed at the protein level using antibodies directed against the conserved

intracellular ATS domain of PfEMP1 (Figure 4D). Altogether, our results of inducible reduced

cytoadhesive capacity is highly relevant for asymptomatic *P. falciparum* infections (Figure 4E).

We show that parasites are able to respond to external stimuli, by controlling PfMaf1 activity,

in order to regulate RNA Pol III transcription and disease severity.

Discussion

In this study, we identified a *P. falciparum* RNA Pol III-dependent regulatory mechanism of parasite virulence that sheds light on the adaptive process of parasite cytoadhesion and proliferation. It suggests a link to asymptomatic infections during the dry season in African regions, when mosquito transmission is halted. By comparing parasite RNA isolated from symptomatic individuals during the wet season with asymptomatic individuals during the dry season, we observed reduced levels of tRNAs and RUF6 ncRNA, a known regulator of *var* gene expression (Figure 1A-C). RNA Pol III has so far not been implicated as a modulator of *P. falciparum* virulence in mild or asymptomatic malaria infections. We present evidence that changes to the parasite environment can signal the observed changes in RNA Pol III activity. To explore how this occurs at the molecular level and to identify contributing host and parasite factors, we developed an *in vitro* culture assay for *P. falciparum* that allowed us to uncover molecular events driving environment-dependent RNA Pol III inhibition (Figure 2A). Several environmental factors, including nutrient starvation and $MgCl_2$ supplementation, were identified that inhibit RNA Pol III transcription of tRNA and the ncRNA RUF6 gene family using qRT-PCR. RNA-seq data confirmed that transcriptional inhibition, with $MgCl_2$ supplementation, is specific almost solely to canonical RNA Pol III-transcribed genes (A and B box containing sequences) including RUF6 that is essential for *var* gene activation. We confirmed that Pol III-mediated *var* gene transcription inhibition reduced cytoadherence of iRBCs upon supplementation of $MgCl_2$ (Figure 4C). We observed >50% reduction in binding to the endothelial receptor CD36 using a static binding assay. We anticipate that inhibition of iRBC cytoadherence would most likely be even more striking under physiological shear stress that occurs in human capillaries (Crabb *et al*, 1997).

Having established a role of Pol III in virulence gene expression, we demonstrate a link between environmental factors and the only known eukaryotic repressor of Pol III, Maf1. Mass spectrometry analysis of the protein interactome of PfMaf1 revealed the binding to several nuclear RNA Pol III subunits, supporting its role as repressor of Pol III. Furthermore, we generated an inducible protein knock down of PfMaf1 to establish its role as regulator of RNA Pol III activity induced by $MgCl_2$ (Figure 4B). Notably, *P. falciparum* PfMaf1 protein showed nuclear presence under standard growth conditions. Nuclear shuttling usually increases under specific conditions (stress, nutrient starvation) by changing the phosphorylation state of the Maf1 protein (Michels *et al*, 2010). Our experimental data support a model where a critical basal level of nuclear PfMaf1 is needed to dampen RNA Pol III expression of tRNAs at the multinucleated blood stage, to adjust its energy resources to produce high numbers of new

infective forms (merozoites). A single parasite replicates its genome several times over, generating approximately 30 nuclei within a shared cytoplasm under optimal growth conditions. Upon complete removal of PfMaf1, tRNA Valine and RUF6 ncRNA, but not Pol II-transcribed *var* and UCE, increased by over 3-fold their normal transcript levels. Drastically raising tRNA transcription over physiological levels could lead to a shortage in cellular energy levels causing the observed decrease in parasite multiplication rate (Figure 3C, E). In clinical isolates, parasite densities have been associated with malaria pathogenesis (Nyarko & Claessens, 2021; Thomson-Luque *et al*, 2021) and low parasite densities are also a feature of asymptomatic malaria (Murray *et al*, 2017).

Nutrient availability in the host has been demonstrated to have a profound effect on the replication of parasites using *P. berghei* as a rodent malaria model (Mancio-Silva *et al*, 2017a). Reports from cultured blood stage *P. falciparum* parasites showed that amino acid deficiency can alter parasite growth and survival to stress (McLean & Jacobs-Lorena, 2017; Marreiros *et al*, 2023). Furthermore, a recent report suggested that nutrients involved in S-adenosylmethionine (SAM) metabolism can also affect *var* gene switching (Schneider *et al*, 2023). Individuals infected with malaria, compared to healthy individuals, were found to have decreased levels (up to 54%) of plasma free amino acids, which could be the result of a nutrient-reduced diet (Leopold *et al*, 2019).

In fact, dietary intake, including energy, protein, iron, zinc, calcium and folate, was found to decrease significantly in individuals during the dry season in areas of Africa (M'Kaibi *et al*, 2015). Blood serum magnesium levels can fluctuate in humans and were found to be over three-fold higher in individuals infected with malaria (Garba & Ubom, 2006). It is noteworthy that intracellular magnesium was found to increase as a result of calorie restriction, defined as a decrease in dietary glucose intake (Abraham *et al*, 2016), indicating that our $MgCl_2$ supplementation could be mimicking low glucose conditions and thus, low nutrient availability. Still, while interactions between human malaria and malnutrition have been studied for many years, the evidence linking an effect between the two remains inconclusive at the mechanistic level. In the absence of a canonical TOR pathway in malaria parasites (Serfontein *et al*, 2010; van Dam *et al*, 2011), the signaling pathway triggering the activation of PfMaf1 remains unknown. Our mass spectrometry analysis of the cytoplasmic PfMaf1 binding partners identifies interactions with potential regulatory enzymes of Maf1 activity (phosphatases and

kinases). This result opens avenues to explore, in the future, the environmental sensing machinery upstream of PfMaf1.

Our work on cultured parasites has established that the parasite RNA Pol III machinery can sense various environmental factors. We assume that host metabolic changes, in particular those that arise during the dry season may impact RNA Pol III activity and contribute to persistent asymptomatic infections. It is tempting to speculate that mediators of the host immune response may modulate the parasite RNA Pol III activity as well. *P. falciparum* induces a strong inflammatory response in humans with the release of soluble mediators like cytokines and chemokines. Several human and mouse studies have shown that macrophage RNA Pol III activity can be modulated and inhibited by an inflammatory response or by intracellular *Leishmania* infection (Rana *et al*, 2011; Graczyk *et al*, 2015). Based on those reports, we hypothesize that soluble host immune factors could impact *P. falciparum* RNA Pol III regulation, thereby linking host immunity to parasite RNA Pol III downregulation in establishing asymptomatic infections.

In conclusion, this work shows the involvement of RNA Pol III in the reduced cytoadhesive capacity of *P. falciparum* in that way unraveling an unprecedented molecular process that is highly relevant for the understanding of subclinical parasite persistence. This knowledge may lead to new strategies to prevent severe malaria by inducing asymptomatic malaria.

Materials and Methods:

Polymerase III inhibition assay

Parasites were treated with 50 μM of RNA Pol III inhibitor CAS 577784-91-9 (Calbiochem, Merck) after sorbitol treatment at 3 ±3hpi and RNA was harvested at 24hpi in parallel with untreated and control samples. The control was treated with the same volume of DMSO added to the inhibitor treated flasks of stock solution.

Clinical isolate RT-qPCR

Venous blood draw of different infected individuals in The Gambia during the dry and wet seasons was collected as previously described in ((Collins *et al*, 2022)). The dry season months include January, February, March, April, and May. The wet season months include November and October. Total RNA was extracted from TRizol using an miRNeasy minikit and performing

on-column DNase treatment (Qiagen) and continued as previously described in ((Collins *et al*,

2022)). Transcript levels were shown by using the following primers: RUF6, Valine tRNA,

Asparagine tRNA, and *var* DBLalpha were normalized to the reference gene, fructose-

bisphosphate aldolase (PF3D7_1444800, Figure 2A). The starting quantity means from three

replicates were extrapolated from a standard curve of serial dilutions of 3D7 genomic DNA.

Parasite culture and synchronization

Asexual blood stage 3D7 *P. falciparum* parasites were cultured as previously described in

(Lopez-Rubio *et al*, 2009). Parasites were cultured in human RBCs (obtained from the

Etablissement Français du Sang with approval number HS 2019-24803) in RPMI-1640 culture

medium (Thermo Fisher 11875) supplemented with 10% v/v Albumax I (Thermo Fisher

11020), hypoxanthine (0.1 mM final concentration, C.C.Pro Z-41-M) and 10 mg gentamicin

(Sigma G1397) at 4% hematocrit and under 5% O_2, 3% CO_2 at 37 °C. Static parasite

development was monitored by Giemsa staining. Parasites were synchronized by sorbitol (5%,

Sigma S6021) lysis at ring stage, plasmagel (Plasmion, Fresenius Kabi) enrichment of late

stages 24 h later, and an additional sorbitol lysis 3 h after plasmagel enrichment. The 0 h time

point was considered to be 1.5 h after plasmagel enrichment. Parasites were harvested at 1–5%

parasitemia.

External stimuli induction

Synchronized parasites were divided into control, addition of magnesium chloride ($MgCl_2$),

isoleucine-deficient, low glucose, and 40 degree Celsius and harvested at 18hpi. Parasites

exposed to an addition of $MgCl_2$ were supplemented with 2.5 mM $MgCl_2$, for a final

concentration of [3mM] including [0.5mM] found in RPMI. We chose [3mM] total $MgCl_2$

concentration as it was reported not to affect growth of *P. falciparum* asexual blood stage after

48 hours and within a physiological range (Jahnen-Dechent & Ketteler, 2012; Hess *et al*, 1995).

Isoleucine-deficient medium consisted of 10.3 g/liter RPMI 1640 isoleucine (Ile) Drop-out

medium (United States BioLogicals; catalog no. R9014), supplemented with 2.0 g/liter

$NaHCO_3$, 6.0 g/liter HEPES, 10% v/v Albumax I (Thermo Fisher 11020), hypoxanthine (0.1

mM final concentration, C.C.Pro Z-41-M) and 10 mg gentamicin (Sigma G1397). Low glucose

(0.25mg/mL) RPMI by adding glucose (Dextrose) to glucose-free media: 2.979 g HEPES + 50

mL albumax + 10 mL hypoxanthine + 200 μL gentamycin to final volume of 500 mL with

glucose free RPMI (11879). Adjust pH to 7.3 with NaOH and filter sterilize. 40-degree Celsius

samples were incubated in an adjacent incubator set to 40 degrees Celsius under 5% O_2, 3%

CO_2. Parasites were then harvested with 0.075% Saponin lysis at ~2-5% parasitemia for RNA,
genomic DNA, and protein extraction at 18hpi.
405

RNA isolation and reverse transcription-quantitative PCR (RT-qPCR)

RNA was isolated from synchronized parasite cultures harvested at 18hpi after saponin lysis in
0.075% saponin in PBS, followed by one wash in Dulbecco's phosphate-buffered saline (DPBS,
Thermo Fisher 14190) and resuspension in the QIAzol reagent. Total RNA was extracted using
an miRNeasy minikit and performing on-column DNase treatment (Qiagen). Reverse
transcription from total RNA was achieved using SuperScript VILO (Thermo Fisher Scientific)
and random hexamer primers. cDNA levels were quantified by quantitative PCR in the CFX384
real time PCR detection system (BioRad) using Power SYBR Green PCR Master Mix (Applied
Biosystems) and primers from a previous study (Guizetti *et al*, 2016). Starting quantity means
of three replicates were extrapolated from a standard curve of serial dilutions of genomic DNA.
Transcript levels were shown by using the following primers: RUF6, Valine tRNA, Alanine
tRNA, Asparagine tRNA, *var* 58 (PF3D7_1240900), and *var* DBLalpha were normalized to
the reference gene, fructose-bisphosphate aldolase (PF3D7_1444800). The starting quantity
means from three replicates were extrapolated from a standard curve of serial dilutions of
genomic DNA.
421

Stranded RNA sequencing and analysis

Infected RBCs containing synchronized (12 and 24hpi) parasites were lysed in 0.075% saponin
(Sigma S7900) in DPBS at 37°C. The parasite cell pellet was washed once with DPBS and then
resuspended in 700 μL QIAzol reagent (Qiagen 79306). Total RNA was subjected to rRNA
depletion to ensure ncRNA and mRNA capture using the RiboCop rRNA Depletion Kit
(Lexogen) prior to strand-specific RNA-seq library preparation using the TruSeq Stranded
RNA LT Kit (Illumina) with the KAPA HiFi polymerase (Kapa Biosystems) for the PCR
amplification. Multiplexed libraries were subjected to 150 bp paired-end sequencing on a
NextSeq 500 platform (Illumina). Sequenced reads (150 bp paired end) were mapped to the *P.*
falciparum genome (Gardner *et al*, 2002) (plasmoDB.org, version 3, release 57) using "bwa
mem" (Li & Durbin, 2009) allowing a read to align only once to the reference genome (option
"–c 1"). Alignments were subsequently filtered for duplicates and a mapping quality ≥ 20 using
samtools (Li & Durbin, 2009). Three biological replicates for -$MgCl_2$ and +$MgCl_2$ samples
were analyzed for both timepoints.
436

Generation of PfMaf1 strains

All cloning was performed using KAPA HiFi DNA Polymerase (Roche 07958846001), In-Fusion HD Cloning Kit (Clontech 639649), and XL10-Gold Ultracompetent *E. coli* (Agilent Technologies 200315). Transgenic pSLI parasites were generated as previously described in (Birnbaum *et al*, 2017) with the following modifications: GFP was replaced with a 3HA tag and a ddFKBP domain was added after the protein of interest, PfMaf1. For localization and knock down studies, the last 500-1000 bp of target gene, PfMaf1 Pf3D7_0416500, was cloned into pSLI-3HA-ddFKBP. Each sequence started with an in-frame stop codon but the stop codon at the end of the gene was removed. 50 μg of plasmid DNA was transfected into ring stage 3D7 *P. falciparum* parasites using the protocol described elsewhere (Hasenkamp *et al*, 2013). Transfected parasites were selected with constant drug selection pressure of 2.56 nM WR99210 (Jacobus Pharmaceuticals) to obtain a cell line containing the episomal plasmid. A second drug selection using 400 μg/ml of G418 was done to select for integrants. Once parasites emerged, gDNA of each integration cell line was collected using a commercial kit (DNeasy Blood & Tissue Kit) and checked by PCR to show that integration occurred at the correct locus. Both genome and vector specific primers for the 5' and 3' region were used so that the PCR product would cover the plasmid/genome junction. Vector primers used were the same as described (Birnbaum *et al*, 2017). Once proper size gel bands from PCR were seen, parasites were cloned by limiting dilution, and the targeted genomic locus was sequenced to confirm tag and FKBP integration.

Flow cytometry

Two different pSLI-Maf1-FKBP parasite clones were tightly synchronized and diluted to 0.2% parasitemia (5% hematocrit) at ring stage. One cycle before, Shield-1 was removed from half of the culture. The growth curve was performed in a 96-well plate (200 μl culture per well) with three technical replicates of two biological replicates per condition per clone. At 0 h, 24 h, 48 h, 72 h, and 96 h, 5 μl of the culture was stained in 95 μl of DPBS supplemented with 2× Sybr Green I (Ozyme; stock = 10,000×) for 30 min at room temperature, diluted 20-fold in D-PBS (final volume = 200 μl), and the Sybr Green fluorescence measured in a Guava easyCyte Flow Cytometer (EMD Millipore). 30000 events were counted in duplicate to establish an accurate parasitemia value for each culture. Data was analyzed using the InCyte software (EMD Millipore).

Western blot analysis

Shield-1 was removed for 3 consecutive cycles to monitor the degradation of PfMaf1. Total protein extracts were prepared from trophozoite stages for control and -Shield-1 for 3 cycles. Additionally, membrane extracts were prepared for control and $MgCl_2$ supplementation from parasites isolated from plasmion enrichment. iRBCs were washed once with DPBS at 37°C and lysed with 0.075% saponin (Sigma S7900) in DPBS at 37°C. Parasites were washed once with DPBS, resuspended in 1 ml cytoplasmic lysis buffer (25 mM Tris–HCl pH 7.5, 10 mM NaCl, 1.5 mM $MgCl_2$, 1% IGEPAL CA-630, and 1× protease inhibitor cocktail ["PI", Roche 11836170001]) at 4°C, and incubated on ice for 30 min. Cells were further homogenized with a chilled glass douncer, and the cytoplasmic lysate was cleared with centrifugation (13,500 g, 10 min, 4°C). The pellet (containing the nuclei) was resuspended in 100 µl nuclear extraction buffer (25 mM Tris–HCl pH 7.5, 600 mM NaCl, 1.5 mM $MgCl_2$, 1% IGEPAL CA-630, PI) at 4°C and sonicated for 10 cycles with 30 s (on/off) intervals (5 min total sonication time) in a Diagenode Pico Bioruptor at 4°C. The nuclear lysate was cleared with centrifugation (13,500 g, 10 min, 4°C). Membrane extracts were prepared by resuspending parasite pellets in NETT buffer (50mM Tris pH8, 150mM NaCl, 5mM EDTA, 1% IGEPAL CA-630, PI) and incubated at 4°C for 10 min. Supernatant was removed after centrifugation (13,500 g, 10 min, 4°C) and the pellet was resuspended in Tris-saline buffer (50mM Tris pH8, 150mM NaCl, 2% SDS, PI) and sonicated for 6 cycles with 30 s (on/off) intervals (3 min total sonication time) in a Diagenode Pico Bioruptor at 4°C. The membrane lysates were cleared with centrifugation (13,500 g, 10 min, 4°C). All protein samples were supplemented with NuPage Sample Buffer (Thermo Fisher NP0008) and NuPage Reducing Agent (Thermo Fisher NP0004) and denatured for 5 min at 95°C. Proteins were separated on a 4-15% TGX (Tris-Glycine eXtended) (Bio-Rad) and transferred to a PVDF membrane. The membrane was blocked for 1 h with 5% milk in PBST (PBS, 0.1% Tween 20) at 25°C. HA-tagged proteins, and histone H3 were detected with anti-HA (Abcam 9110, 1:1,000 in 5% milk-PBST) and anti-H3 (Abcam ab1791, 1:1,000 in 5% milk-PBST) primary antibodies, respectively, followed by donkey anti-rabbit secondary antibody conjugated to horseradish peroxidase ("HRP", Sigma GENA934, 1:5,000 in 5% milk-PBST). Anti-ATS and anti-NapL were detected as previously described (Nacer *et al*, 2015; Dawn *et al*, 2014). Aldolase was detected with anti-aldolase-HRP (Abcam ab38905, 1:5,000 in 5% milk-PBST). HRP signal was developed with SuperSignal West Pico chemiluminescent substrate (Thermo Fisher 34080) and imaged with a ChemiDoc XRS+ (Bio-Rad).

Merozoite number per schizont analysis:

504 Mature schizonts were assessed by microscopic analysis of Giemsa-stained smears and
505 manually quantified using ImageJ software, as reported previously (Marreiros *et al*, 2023). 100
506 segmented schizonts with clearly individualized merozoites containing a single hemozoin
507 crystal were quantified per condition.

Co-Immunoprecipitation followed by Mass spectrometry (Co-IP-MS)

PfMaf1-HA-ddFKBP tagged parasites ($n = 5$ biological replicates) were synchronized. At
24hpi, each culture (1.5×10^9 parasites) was centrifuged and RBCs were lysed with six volumes
of 0.15% saponin in DPBS for 5 min at 4°C. Parasites were centrifuged at 4,000 g for 5 min at
4°C, and the pellet was washed twice with DPBS at 4°C. Parasites were then cross-linked with
1% formaldehyde for 15 min at room temperature and quenched with 125 mM glycine for 5
min on ice. Cross-linked parasites were washed twice with DPBS and then resuspended in
900uL of cytoplasmic lysis buffer (10 mM Tris–HCl pH 7.5, 1 mM EDTA, 0.65% IGEPAL
CA-630, 10mM NaCl) supplemented with protease inhibitors (Thermo Fisher 78440) at 4°C
and incubated with rotation for 30 min at 4°C. Extracts were centrifuged for 10 min at 2000g
at 4°C and the cleared cytoplasmic supernatant was removed and kept on ice. The nuclear pellet
was resuspended in 900 µL nuclear lysis buffer (10 mM Tris–HCl pH 7.5, 500 mM NaCl, 1
mM EDTA, 1% sodium deoxycholate, 0.1% SDS, 1% IGEPAL CA-630, PI) at 4°C and
transferred to 1.5 ml sonication tubes (300 µL per tube, DiagenodeC30010016). Samples were
sonicated for 5 min (30 s on/off) in a Diagenode Pico Bioruptor at 4°C. Lysates were then
centrifuged for 10 min at 13,500g at 4°C and supernatant was transferred to a fresh tube.
Cytoplasmic fractions were mixed with 2:3 ratio of cytoplasmic dilution buffer (10 mM Tris-
HCl pH 7.5, 150 mM NaCl, 0.5 mmM EDTA) and nuclear supernatants were mixed with 1:3
ratio of nuclear dilution buffer (10 mM Tris-HCl pH 7.5, 0.5 mmM EDTA). Cytoplasmic and
nuclear fraction supernatants were incubated with 1ug of α-HA antibody (Abcam 9110) and
25uL Protein G Magnetic Dynabeads (Invitrogen), pre-incubated for a minimum of 2 hours and
washed twice with dilution buffer, overnight with rotation at 4°C. The next day, the beads were
collected on a magnet and the supernatant was removed. While on the magnetic stand, beads
were washed twice with 500 µL wash buffer (10 mM Tris-HCl pH 7.5, 150 mM NaCl, 0.5 mM
EDTA, 0.05% NP40), once with 25 mM NH4HCO3 (Sigma 09830) buffer, and then transferred
to new tube. Finally, the beads were resuspended in 100 µl of 25 mM NH4HCO3 (Sigma 09830)
and digested by adding 0.2 µg of trypsin-LysC (Promega) for 1 h at 37 °C. Samples were then
loaded into custom-made C18 StageTips packed by stacking one AttractSPE® disk (#SPE-
Disks-Bio-C18-100.47.20 Affinisep) in a 200 µL micropipette tip for desalting. Peptides were

eluted using a ratio of 40:60 CH3CN:H2O + 0.1% formic acid and vacuum concentrated to dryness with a SpeedVac apparatus. Peptides were reconstituted in 10 of injection buffer in 0.3% trifluoroacetic acid (TFA) before liquid chromatography-tandem mass spectrometry (LC-MS/MS) analysis. Online LC was performed with an RSLCnano system (Ultimate 3000, Thermo Scientific) coupled to an Orbitrap Eclipse mass spectrometer (Thermo Scientific). Peptides were trapped on a 2 cm nanoviper Precolumn (i.d. 75 μm, C18 Acclaim PepMap^{TM} 100, Thermo Scientific) at a flow rate of 3.0 μL/min in buffer A (2/98 MeCN/H2O in 0.1% formic acid) for 4 min to desalt and concentrate the samples. Separation was performed on a 50 cm nanoviper column (i.d. 75 μm, C18, Acclaim PepMap^{TM} RSLC, 2 μm, 100Å, Thermo Scientific) regulated to a temperature of 50°C with a linear gradient from 2% to 25% buffer B (100% MeCN in 0.1% formic acid) at a flow rate of 300 nL/min over 91 min. MS1 data were collected in the Orbitrap (120,000 resolution; maximum injection time 60 ms; AGC 4 x 10^5). Charges states between 2 and 7 were required for MS2 analysis, and a 60 s dynamic exclusion window was used. MS2 scan were performed in the ion trap in rapid mode with HCD fragmentation (isolation window 1.2 Da; NCE 30%; maximum injection time 60 ms; AGC 104) For identification, the data were searched against the *Homo Sapiens* (UP000005640_9606) and the *Plasmodium Falciparum 3D7* (UP000001450_36329) UniProt databases using Sequest HT through proteome discoverer (version 2.4). Enzyme specificity was set to trypsin and a maximum of two miss cleavages sites were allowed. Oxidized methionine, Met-loss, Met-loss-Acetyl and N-terminal acetylation were set as variable modifications. Carbamidomethylation of cysteins were set as fixed modification. Maximum allowed mass deviation was set to 10 ppm for monoisotopic precursor ions and 0.6 Da for MS/MS peaks. The resulting files were further processed using myProMS v3.9.3 (https://github.com/bioinfo-pf-curie/myproms) (Poullet *et al*, 2007) FDR calculation used Percolator (The *et al*, 2016) and was set to 1% at the peptide level for the whole study. The label free quantification was performed by peptide Extracted Ion Chromatograms (XICs), reextracted across all conditions and computed with MassChroQ version 2.2.21 (Valot *et al*, 2011) For protein quantification, XICs from proteotypic peptides shared between compared conditions (TopN matching) and missed cleavages were allowed. Median and scale normalization was applied on the total signal to correct the XICs for each biological replicate (n = 5 in each condition). To estimate the significance of the change in protein abundance, a linear model (adjusted on peptides and biological replicates) was performed, and p-values were adjusted using the Benjamini–Hochberg FDR procedure. Proteins with at least 2 distinct peptides in 3 replicates of a same state, a 2-fold enrichment and

571 an adjusted p-value ≤ 0.05 were considered significantly enriched in sample comparisons.

572 Proteins unique to a condition were also considered if they matched the peptides criteria.

Immunofluorescence assay

pSLI-Maf1-FKBP control, +MgCl$_2$, and low glucose parasites were used with rat anti-HA

(Roche 3F10) antibodies. 10 µl of iRBCs were washed with PBS and fixed with for 30 min in

0.0075% Glutaraldehyde/4% PFA/DPBS. After DPBS washing, parasites were permeabilized

with 0.1% TritonX100/PBS for 10-15 min before quenching free aldehyde groups with NH$_4$Cl

solution for 10 min. Next, parasites were blocked with 3% BSA–DPBS for 30 min. Primary

antibody incubation (1:500 dilution) lasted for 1 h before three washes with PBS, and secondary

antibody incubation for 30-60 min, Alexa Fluor 488-conjugated anti-mouse IgG (Invitrogen)

diluted 1:2,000 in 4% BSA–DPBS. After three final washes in DPBS, cells were mounted in

Vectashield containing DAPI for nuclear staining. Images were captured using a Delta Vision

Elite microscope (GE Healthcare). Image overlays were generated using Fiji (Schindelin *et al*,

2012).

Co-Immunoprecipitation followed by western blot (Co-IP-WB)

PfMaf1-HA-ddFKBP tagged parasites were synchronized and split into control and MgCl$_2$

supplementation. At 18hpi, each culture (1.5×10^9 parasites) was centrifuged and RBCs were

lysed with six volumes of 0.15% saponin in DPBS for 5 min at 4°C. Parasites were centrifuged

at 4,000 *g* for 5 min at 4°C, and the pellet was washed twice with DPBS at 4°C. The parasite

pellets were then resuspended in 900uL of cytoplasmic lysis buffer (10 mM Tris–HCl pH 7.5,

1 mM EDTA, 0.65% IGEPAL CA-630, 10mM NaCl) supplemented with protease inhibitors

(Thermo Fisher 78440) at 4°C and incubated with rotation for 30 min at 4°C. Extracts were

centrifuged for 10 min at 2000g at 4°C and the cleared cytoplasmic supernatant was removed

and kept on ice. The nuclear pellet was resuspended in 200 µL nuclear lysis buffer (10 mM

Tris–HCl pH 7.5, 500 mM NaCl, 1 mM EDTA, 1% sodium deoxycholate, 0.1% SDS, 1%

IGEPAL CA-630, PI) at 4°C and transferred to 1.5 ml sonication tubes (300 µL per tube,

DiagenodeC30010016). Samples were sonicated for 5 min (30 s on/off) in a Diagenode Pico

Bioruptor at 4°C. Lysates were then centrifuged for 10 min at 13,500g at 4°C and supernatant

was transferred to a fresh tube. Cytoplasmic fractions were mixed with 2:3 ratio of cytoplasmic

dilution buffer (10 mM Tris-HCl pH 7.5, 150 mM NaCl, 0.5 mmM EDTA) and nuclear

supernatants were mixed with 1:3 ratio of nuclear dilution buffer (10 mM Tris-HCl pH 7.5, 0.5

mmM EDTA). Cytoplasmic and nuclear fraction supernatants were incubated with 1ug of α-

HA antibody (Abcam 9110) and 25uL Protein G Magnetic Dynabeads (Invitrogen), pre-

incubated for a minimum of 2 hours and washed twice with dilution buffer, overnight with

rotation at 4°C. The next day, the beads were collected on a magnet and the supernatant was

removed. While on the magnetic stand, beads were washed twice with 500 μL wash buffer (10

mM Tris-HCl pH 7.5, 150 mM NaCl, 0.5 mM EDTA, 0.05% NP40), once with 25 mM

NH_4HCO_3 (Sigma 09830) buffer, and then transferred to new tube. Finally, the beads were

resuspended in 20μl 2x SDS sample buffer and reducing agent before 5 min at 95°C.

The beads were separated with magnet and the supernatant was transferred to a new tube.

Western blot protocol was followed as mentioned earlier.

Static cytoadhesion binding assay

Mature stage iRBCs for 3D7 control and supplementation of $MgCl_2$ ([3mM] total

concentration) for 1 cycle were used for cytoadhesion binding assays. Receptors diluted in PBS

(10 µg/mL target CD36 or negative control 1% BSA) were incubated overnight at 4°C in

labeled petri dishes. After the dishes were blocked for 30 minutes at 37°C with 1% BSA/PBS,

parasitemia was measured with a Guava easyCyte Flow Cytometer (EMD Millipore) after

trophozoites and schizonts iRBCs were isolated by plasmagel (Plasmion, Fresenius Kabi)

enrichment. Adjusted amounts of iRBCs were resuspended in binding medium (RPMI 1640

powder (Gibco 51800: W/L-Glutamine W/O $NaHCO_3$) with HEPES pH 7) for a concentration

of 2.2×10^8 iRBCs/mL. iRBCs/binding medium was added to each petri dish and incubated for

2 hours at 37°C. After, unbound cells were removed and the dishes were washed 5 times with

binding medium by carefully tilting from side to side. Adherent iRBCs were counted using 40x

lens with a Nikon ECLIPSE TE200 in 10 randomly selected fields (each with 0.2 μm^2) before

2 our fixation with 2% glutaraldehyde (G5882; Sigma)/PBS. After fixation, giemsa staining

was done to confirm percentage of bound iRBCs. A total of 3 biological replicates were

performed and the results were expressed as the number of bound iRBCs per 0.2 mm^2 of target

receptor monolayer.

Statistical analysis

All statistical analyses were performed using GraphPad Prism version 9.1.0 (216) for Mac. To

test for a normal distribution of the data, the Shapiro-Wilk normality test was used. To test for

significance between two groups, a two-sided independent-samples t test was used. Gene

ontology enrichments were calculated using the build-in tool at https://plasmoDB.org.

Estimation of cell cycle progression

RNA-seq-based cell cycle progression for control and MgCl$_2$ supplementation was estimated in R by comparing the normalized expression values (i.e., RPKM, reads per kilobase per exon per one million mapped reads) of each sample to the microarray data from Bozdech et al (2003) (Data ref: Bozdech et al, 2003) using the statistical model as in Lemieux et al (2009).

Data availability

The data generation in this study is available in the following database:

The mass spectrometry proteomics data have been deposited to the ProteomeXchange Consortium via the PRIDE (Perez-Riverol *et al*, 2022) partner repository with the dataset identifier PXD040576" (Username: reviewer_pxd040576@ebi.ac.uk & Password: wlw5Mlc6).

Acknowledgments

We thank L. Mancio Silva, E. Real and J. Bryant for critical reading this manuscript. This work was supported by the Laboratoire d'Excellence (LabEx) ParaFrap [ANR-11-LABX-0024], ERC AdG PlasmoSilencing (670301), ANR-JC (18-CE15-0009-01). We thank Miguel Santos, Camille Cohen and Marta Miera Maluenda for technical assistance with this project. With financial support from "la Région Île-de-France" (N°EX061034) and ITMO Cancer of Aviesan and INCa on funds administered by Inserm (N°21CQ016-00) for MS analysis.

Author contributions

Conceptualization A.S, GD; Methodology: GD, AClaës, AB, PBN; Investigation GD, AClaës, AB; Analysis: GD, AB, FD, DL; Writing: GD and A.S; Funding acquisition: A.S, AC

Conflict of interest

The funders had no role in the study design, data collection, and interpretation or decision to submit the work for publication. The authors declare that they have no conflict of interest.

671
Figure 1. RNA Pol III-transcribed genes are downregulated in asymptomatic individuals. Steady state RNA
levels as quantified by RT-qPCR using primers to tRNA Asparagine (Pf3D7_0714700) (A) and tRNA Valine
(Pf3D7_0312600) (B) as well as RNA Pol III-transcribed RUF6 ncRNA (C). DBLalpha primers were used to
detect RNA Pol II-transcribed *var* genes (D). Normalized expression is shown using fructose-bisphosphate
aldolase (FBA Pf3D7_1444800) as the reference gene in symptomatic individuals during the wet season (n = 14[+])
and asymptomatic individuals during the wet season (n = 12[+]) and dry season (n = 17[+]). [+] with the exception of
Asparagine tRNA wet symptomatic (n=11) and dry asymptomatic (n=12), Valine tRNA wet asymptomatic (n =
11), RUF6 wet asymptomatic (n = 4) and DBLalpha wet symptomatic (n = 12), wet asymptomatic (n = 9), and dry
asymptomatic (n = 16). Box plots indicate the mean with standard deviation. Wilcoxon matched-pairs signed rank
test was done to determine significance (* indicates $P < 0.05$ and ** indicates $P < 0.005$).

682
683
684

A

B

C   D

685
686
Figure 2. External factors modulate RNA Pol III-transcribed genes. (A) Schematic showing underlying
molecular mechanism summary using *in vitro* cultured *P. falciparum.* (B) Transcript levels as quantified by RT-
qPCR on 3D7 parasites harvested during late ring stage for control parasites, parasites in the absence of isoleucine
(w/o Ile), at 40 degrees Celsius, at low-glucose levels (0.5mg/mL), and presence of additional $MgCl_2$ ([3mM]
total). Primers were used for Pol II-transcribed ubiquitin-conjugating enzyme (UCE Pf3D7_0812600), Pol I-
transcribed rRNA A1, Pol III-transcribed RUF6 ncRNA, Pol III-transcribed tRNA Valine (Pf3D7_0312600), and
Pol II-transcribed active *var* gene (Pf3D7_1240900). Results are normalized to an RNA Pol II-transcribed
reference gene FBA (Pf3D7_1444800) and presented as fold change/control. Error bars are displayed from 3
biological replicates. Statistical significance was determined by two-tailed Student's t-test (* p<0.05). (C, D)
Volcano plot showing log2 (fold change, FC) against −log10 (p-value) of transcripts identified by RNA-seq

analysis of 3D7 control and addition of $MgCl_2$ harvested during ring (C) and trophozoite stages (D). Expressed
transcripts from three replicates between control and addition of $MgCl_2$ that are significantly up-regulated are
highlighted in red while significantly down-regulated RNA Pol III genes are highlighted in blue (FDR corrected
p-value of <0.05) and a FC $\geq \pm$ 1.95) with examples listed as text. Black dots indicate non-significant transcripts
with a FC $\leq$ 2.0.
702

703
704
Figure 3. Nuclear PfMaf1 is essential to regulate RNA Pol III. (A) Illustration of the TORC1-dependent cellular
localization of Maf1 protein in unfavorable conditions in typical eukaryotic organisms compared *P. falciparum*
that has no TORC1 pathway. (B, top) Illustration of plasmid design for PfMaf1 with a 3HA tag followed by a
ddFKBP domain to allow for knockdown studies. (B, bottom) Western blot analysis for PfMaf1 in total extracts
in pSLI-Maf1-FKBP transfected parasites at 24hpi after 1, 2, and 3 cycles without addition of Shield-1 (-) and
control, with addition of Shield-1 (+). Aldolase levels are also shown. (C) Transcript levels as quantified by RT-
qPCR using the same primers in (Figure 2B) in parasites harvested at 18hpi in control group and without Shield
for 2 cycles. Error bars are displayed from 3 biological replicates. Statistical significance was determined by two-
tailed Student's t-test (** p<0.005). (D) Growth curve over 5 days of clonal pSLI-Maf1-FKBP parasites for 2
conditions: in the presence or absence of Shield-1. Uninfected red blood cells ("blood" in red) serve as reference
of background. Error bars indicate standard deviation of three technical replicates in different blood from two
different clones (n = 6). (E) Data is represented as box-whisker plot of mean merozoite number per schizont ±SD
(Mann-Whitney), with the with the median represented at the center line. Boxplots show the data of 100 segmented
schizonts counted per condition (n = 100). Statistical significance was determined by two-tailed Student's t-test
(*** p<0.001). Representative giemsa images are shown to the right for + and – Shield. (F) Volcano plot showing

the distribution of significant and unique proteins in cytoplasmic and nuclear fractions not found in either of the
controls. Each dot represents a protein, and its size corresponds to the sum of peptides from both conditions used
to quantify the ratio of enrichment. x-axis = $\log_2$(fold-change), y-axis = $-\log_{10}$(p-value), horizontal red line
indicates adjusted p-value = 0.05, and vertical green lines indicate absolute fold-change = 2.0. Side panels indicate
proteins uniquely identified in either sample (y-axis = number of peptides per 100 amino acids) with a minimum
of 2 distinct peptides in 3 replicates of a same state.

752
753
Figure 4. External factors modulate virulence through PfMaf1-regulated RNA Pol III inhibition
(A) Immunoprecipitation western blot analysis for cytoplasmic and nuclear extracts for PfMaf1 expression in
PfMaf1-FKBP transfected parasites with addition of $MgCl_2$ ([3mM] total) and control parasites harvested at 18hpi.
Controls are shown in Figure S4B. (B) Transcript levels as quantified by RT-qPCR using primers to tRNA Valine
(Pf3D7_0312600), RUF6 ncRNA, *var* DBLalpha, and normalized to FBA (Pf3D7_1444800) for 4 conditions:
control (black), addition of $MgCl_2$ (green), KD of PfMaf1 (blue), and KD of PfMaf1 and addition of $MgCl_2$
(orange). Error bars are displayed from 3 biological replicates. Statistical significance was determined by two-
tailed Student's t-test (** $p<0.005$). (C) Cytoadhesion binding assay data is represented as box-whisker plot of
mean number of bound iRBC ±SD (Mann-Whitney) mm^2, with the median represented at the center line. Boxplots
show the data of 3 biological replicates (n = 3). Statistical significance was determined by two-tailed Student's t-
test (**** $p<0.001$). Representative images are shown below for 3D7 and 3D7 +$MgCl_2$. (D) Western blot analysis
for extracts for ATS-PfEMP1 expression in 3D7 control parasites and with addition of $MgCl_2$ ([3mM] total)
harvested after plasmin. NapL control levels are also shown. (E) Schematic showing summary of study linking
decreased cytoadherence, associated with disease severity, with increased RNA Pol III-inhibition, triggered in
response to external factors.
769
770
771
772

A

B

773
Figure S1. (A) Multiple sequence alignment of highly conserved genes encoding GC-rich ncRNA elements. 15
members of RUF6 gene family aligned by Clustal Omega (http://www.ebi.ac.uk) and presented in Jalview
(http://www.jalview.org). Degree of conservation per base and consensus sequence are displayed below. Black
lines show position of potential A- and B-box consensus motifs (as assessed in (Guizetti *et al*, 2016)). (B) RT-
qPCR shows transcript levels for tRNA Asparagine, tRNA Valine, RUF6 ncRNA and a housekeeping gene
encoding ubiquitin-conjugating enzyme in synchronized wildtype parasites at 24 hpi that were untreated or treated
with RNA Pol III inhibitor or DMSO. Transcript levels are normalized to fructose-bisphosphate aldolase
(PF3D7_1444800) transcript levels. Mean SEM of two independent experiments are shown. Statistical
significance was determined by two-tailed Student's t-test (* $p < 0.05$).

A

B

783
784
Figure S2. (A) tRNA Leucine (Pf3D7_API05400) and tRNA Asparagine (Pf3D7_API05500) transcripts
identified by RNA-Seq analysis from ((Andrade *et al*, 2020)) for wet symptomatic (n = 12) and dry asymptomatic
(n = 12) individuals. Statistical significance was determined by two-tailed Student's t-test (* p<0.05). (B) Cell
cycle progression estimation of a wildtype 3D7 clone in the absence ('C') or presence ('Mg') of MgCl$_2$
supplementation. RNA-seq data from synchronized parasites harvested at 12 and 24 hpi were compared to
microarray data (from (Bozdech *et al*, 2003)) as in (Lemieux *et al*, 2009). Replicates are represented with circles.
791
792
793
794

A Cytoplasmic PfMaf1 vs Cytoplasmic Control

B Nuclear PfMaf1 vs Nuclear Control

C

Gene ID	Ab Cytoplasm/Control Cytoplasm						Total peptides in set	MW (kDa)	Description
	Ratio	Log2(Ratio)	Adj. p-value	Dist. pept. used	Pept. used	Distrib. dist. pep. in replic.			
PF3D7_1464600	2,263481181	1,178543312	1,577E-06	6	31	4, 5, 5, 4, 3/0, 4, 3, 2, 1	70	170,5	Serine/threonine protein phosphatase UIS2, putative
PF3D7_1127000	2,838513427	1,505135566	0,00160101	2	16	1, 2, 2, 2, 2/1, 0, 2, 2, 2	17	34,5	Protein phosphatase, putative

Gene ID	Ab Nuclear/Control Nuclear						Total peptides in set	MW (kDa)	Description
	Ratio	Log2(Ratio)	Adj. p-value	Dist. pept. used	Pept. used	Distrib. dist. pep. in replic.			
PF3D7_0507700	4,544185096	2,184021601	0,00072787	3	18	3, 2, 3, 2, 2/2, 0, 2, 0, 2	32	62,6	Nuclear protein localization protein 4, putative
PF3D7_1206600	1000	1000		6	13	2, 3, 3, 3, 2/0, 0, 0, 0, 0	13	167,3	DNA-directed RNA polymerase subunit beta
PF3D7_1329000	1000	1000		5	13	2, 3, 1, 3, 2/1, 1, 0, 0, 0	13	274,8	DNA-directed RNA polymerase subunit

Figure S3. PfMaf1 cytoplasmic and nuclear interactome. Co-IP MS volcano plot of enrichment for all 5 replicates for cytoplasmic (A) and nuclear (B) PfMaf1 vs control proteins are indicated and labeled. Each dot represents a protein, and its size corresponds to the sum of peptides from both conditions used to quantify the ratio of enrichment. x-axis = log$_2$(fold-change), y-axis = $-$log$_{10}$(p-value), horizontal red line indicates adjusted p-value = 0.05, and vertical green lines indicate absolute fold-change = 2.0. Side panels indicate proteins uniquely identified in either sample (y-axis = number of peptides per 100 amino acids) with a minimum of 2 distinct peptides in 3 replicates of a same state. (C) Table showing values for significantly and uniquely enriched proteins from both extracts as labeled in (Figure 3F).

A

B

807
808
Figure S4. (A) Immunoprecipitation western blot analysis for cytoplasmic and nuclear extracts for PfMaf1
expression in PfMaf1-FKBP transfected parasites with addition of $MgCl_2$ ([3mM] total) and control parasites
harvested at 18hpi. Anti-HA PfMaf1, aldolase and histone H3 are shown from input and flow-through (FT). (B)
Representative immunofluorescence images show brightfield, Dapi, anti-HA PfMaf1, and Dapi-HA merge for
PfMaf1 in control, addition of $MgCl_2$ ([3mM] total), and low glucose (0.5mg/mL) in parasites harvested and fixed
at late ring stage.

815
816

817 **Supplementary Table 1. qPCR analysis primer pairs**
818

Primer pair	Forward	Reverse
Pool RUF6 A RUF6 B	5'-AAGCTGCCTCAGTAGCCCA-3'	5'-AAAAATTGCGCCACCCCC-3'
	5'-AAGCTGCCCCAGTAGCCCA-3'	5'-AAAAATTGCGCCGCCCCC-3'
rRNA A1 (from (Mancio-Silva *et al*, 2010))	5'-TGTTTTCTTTTTTCTAAGTTT-3'	5'-TCACCTCATTTGAAGCAA-3'
tRNA Alanine (PF3D7_0411500)	5'-GGGCAGGTGGTGTAGTGG-3'	5'-TGCTGGACAGACGGGGAATT-3'
tRNA Asparagine (PF3D7_0714700)	5'-GCAAGTATTTCCGCCTGTCA-3'	5'-GAATTGAACCCGGGTCTTCC-3'
tRNA Valine (Pf3D7_0312600)	5'-GCGGGCATGGTCTAGTGG-3'	5'-ACTACGGGCACCGAGGATC-3'
FBA fructose- bisphosphate aldolase (PF3D7_1444800)	5'-TGTACCACCAGCCTTACCAG-3'	5'-TTCCTTGCCATGTGTTCAAT-3'
UCE ubiquitin- conjugating enzyme (Pf3D_70812600)	5'-TAACAGCCCAGCGAATCAAG-3'	5'-CGGCATCTTCTTCAGCTTTCTG-3'
var gene (PF3D7_1240900)	5'-CAAAATGGTAGTGATGGTGGTCG-3'	5'-CCCCCTGCTTTATTATCTTTCGTC-3'
Pool *var* DBLalpha	5'-GCACGAACTTTTGCA-3'	5'-GCCCCATTCGTCGAACC-3'
	5'-GCACGCAGTTTTGCA-3'	5'-GCCCCATTCCTCGAACC-3'

820   References:
821

Abraham KJ, Chan JN, Salvi JS, Ho B, Hall A, Vidya E, Guo R, Killackey SA, Liu N, Lee
JE, *et al* (2016) Intersection of calorie restriction and magnesium in the suppression of
genome-destabilizing RNA-DNA hybrids. *Nucleic Acids Res Oct* 14;44(18):8870-
8884

Andrade CM, Fleckenstein H & Thomson-Luque R (2020) Increased circulation time of
Plasmodium falciparum underlies persistent asymptomatic infection in the dry season.
Nat Med 26: 1929–1940

Babbitt SE, Altenhofen L, Cobbold SA, Istvan ES, Fennell C, Doerig C, Llinás M &
Goldberg DE (2012) *Plasmodium falciparum* responds to amino acid starvation by
entering into a hibernatory state. *Proc Natl Acad Sci USA* 109

Barcons-Simon A, Cordon-Obras C, Guizetti J, Bryant JM & Scherf A (2020) CRISPR
Interference of a Clonally Variant GC-Rich Noncoding RNA Family Leads to General
Repression of var Genes in Plasmodium falciparum. *mBio* 11

Birnbaum J, Flemming S & Reichard N (2017) A genetic system to study Plasmodium
falciparum protein function. *Nat Methods* 14: 450–456

Bozdech Z, Llinás M, Pulliam BL, Wong ED, Zhu J & DeRisi JL (2003) The Transcriptome
of the Intraerythrocytic Developmental Cycle of Plasmodium falciparum. *PLoS Biol* 1:
e5

Collins KA, Ceesay S, Drammeh S, Jaiteh FK, Guery MA, Lanke K, Grignard L, Stone W,
Conway DJ, D'Alessandro U, *et al* (2022) A Cohort Study on the Duration of
Plasmodium falciparum Infections During the Dry Season in The Gambia. *The*
Journal of Infectious Diseases 226: 128–137

Crabb BS, Cooke BM, Reeder JC, Waller RF, Caruana SR, Davern KM, Wickham ME,
Brown GV, Coppel RL & Cowman AF (1997) Targeted Gene Disruption Shows That
Knobs Enable Malaria-Infected Red Cells to Cytoadhere under Physiological Shear
Stress. *Cell* 89: 287–296

van Dam TJP, Zwartkruis FJT, Bos JL & Snel B (2011) Evolution of the TOR Pathway. *J*
Mol Evol 73: 209–220

Dawn A, Singh S, More KR, Siddiqui FA, Pachikara N, Ramdani G, Langsley G & Chitnis
CE (2014) The Central Role of cAMP in Regulating Plasmodium falciparum
Merozoite Invasion of Human Erythrocytes. *PLoS Pathog* 10: e1004520

Diffendall GM, Barcons-Simon A, Baumgarten S, Dingli F, Loew D & Scherf A (2022)
Discovery of RUF6 ncRNA–interacting proteins involved in *P. falciparum* immune
evasion. *Life Sci Alliance* 6: e202201577

Fan Y, Shen S, Wei G, Tang J, Zhao Y, Wang F, He X, Guo G, Shang X, Yu X, *et al* (2020)
Rrp6 Regulates Heterochromatic Gene Silencing via ncRNA RUF6 Decay in Malaria
Parasites. *mBio* 11: e01110-20

Fang J, Sullivan M & McCutchan TF (2004) The Effects of Glucose Concentration on the
Reciprocal Regulation of rRNA Promoters in Plasmodium falciparum. *Journal of*
Biological Chemistry 279: 720–725

Garba IH & Ubom GA (2006) Potential Role of Serum Magnesium Measurement as a
Biomarker of Acute Falciparum Malaria Infection in Adult Patients. *BTER* 114: 115–
120

Gardner MJ, Hall N, Fung E, White O, Berriman M, Hyman RW, Carlton JM, Pain A, Nelson
KE, Bowman S, *et al* (2002) Genome sequence of the human malaria parasite
Plasmodium falciparum. *Nature* 419: 498–511

Graczyk D, White RJ & Ryan KM (2015) Involvement of RNA Polymerase III in Immune
Responses. *Molecular and Cellular Biology* 35: 1848–1859

Guillochon E, Fraering J, Joste V, Kamaliddin C, Vianou B, Houzé L, Baudrin LG, Faucher
JF, Aubouy A, Houzé S, *et al* (2022) Transcriptome Analysis of *Plasmodium*
falciparum Isolates From Benin Reveals Specific Gene Expression Associated With
Cerebral Malaria. *The Journal of Infectious Diseases* 225: 2187–2196

Guizetti J, Barcons-Simon A & Scherf A (2016) Trans-acting GC-rich non-coding RNA at
var expression site modulates gene counting in malaria parasite. *Nucleic Acids Res* 44:
9710–9718

Guizetti J & Scherf A (2013) Silence, activate, poise and switch! Mechanisms of antigenic
variation in Plasmodium falciparum. *Cell Microbiology* 15: 718–726

Hasenkamp S, Merrick CJ & Horrocks P (2013) A quantitative analysis of Plasmodium
falciparum transfection using DNA-loaded erythrocytes. *Mol Biochem Parasitol* 187:
117–120

Hess FI, Kilian A, Söllner W, Nothdurft HD, Pröll S & Löscher T (1995) Plasmodium
falciparum and Plasmodium berghei: Effect of Magnesium on the Development of
Parasitemia. *Experimental Parasitology* 80: 196–193

Jahnen-Dechent W & Ketteler M (2012) Magnesium basics. *Clinical Kidney Journal* 5: i3–
i14

Jensen MD, Conley M & Helstowski LD (1983) Culture of Plasmodium falciparum: The Role
of pH, Glucose, and Lactate. *The Journal of Parasitology* 69: 1060

Kanellopoulou C, George AB, Masutani E, Cannons JL, Ravell JC, Yamamoto TN,
Smelkinson MG, Jiang PD, Matsuda-Lennikov M, Reilley J, *et al* (2019) Mg2+
regulation of kinase signaling and immune function. *Journal of Experimental*
Medicine 216: 1828–1842

Kimenyi KM, Wamae K & Ochola-Oyier LI (2019) Understanding P. falciparum
Asymptomatic Infections: A Proposition for a Transcriptomic Approach. *Front*
Immunol 10: 2398

Kumar M, Skillman K & Duraisingh MT (2021) Linking nutrient sensing and gene
expression in *Plasmodium falciparum* blood-stage parasites. *Mol Microbiol* 115: 891–
900

Leech JH, Barnwell JW, Miller LH & Howard RJ (1984) Identification of a strain-specific
malarial antigen exposed on the surface of plasmodium falciparum-infected
erythrocytes. *J Exp Med* 159: 1567–1575

Lemieux JE, Gomez-Escobar N, Feller A, Carret C, Amambua-Ngwa A, Pinches R, Day F,
Kyes SA, Conway DJ, Holmes CC, *et al* (2009) Statistical estimation of cell-cycle
progression and lineage commitment in *Plasmodium falciparum* reveals a
homogeneous pattern of transcription in ex vivo culture. *Proc Natl Acad Sci USA* 106:
7559–7564

Leopold SJ, Apinan S, Ghose A, Kingston HW, Plewes KA, Hossain A, Dutta AK, Paul S,
Barua A, Sattar A, *et al* (2019) Amino acid derangements in adults with severe
falciparum malaria. *Sci Rep* 9: 6602

Li H & Durbin R (2009) Fast and accurate short read alignment with BurrowsWheeler
transform. *Bioinformatics* 25: 1754–1760

Lopez-Rubio JJ, Mancio-Silva L & Scherf A (2009) Genome-wide analysis of
heterochromatin associates clonally variant gene regulation with perinuclear
repressive centers in malaria parasites. *Cell Host Microbe* 5: 179–190

Mancio-Silva L, Slavic K, Grilo Ruivo MT, Grosso AR, Modrzynska KK, Vera IM, Sales-
Dias J, Gomes AR, MacPherson CR, Crozet P, *et al* (2017a) Nutrient sensing
modulates malaria parasite virulence. *Nature* 547: 213–216

Mancio-Silva L, Slavic K, Grilo Ruivo MT, Grosso AR, Modrzynska KK, Vera IM, Sales-
Dias J, Gomes AR, MacPherson CR, Crozet P, *et al* (2017b) Nutrient sensing
modulates malaria parasite virulence. *Nature* 547: 213–216

Mancio-Silva L, Zhang Q, Scheidig-Benatar C & Scherf A (2010) Clustering of dispersed
ribosomal DNA and its role in gene regulation and chromosome-end associations in
malaria parasites. *Proc Natl Acad Sci USA* 107: 15117–15122

Marreiros IM, Marques S, Parreira A, Mastrodomenico V, Mounce BC, Harris CT, Kafsack
BF, Billker O, Zuzarte-Luís V & Mota MM (2023) A non-canonical sensing pathway
mediates Plasmodium adaptation to amino acid deficiency. *Commun Biol* 6: 205

McLean KJ & Jacobs-Lorena M (2017) Plasmodium falciparum Maf1 Confers Survival upon
Amino Acid Starvation. *mBio* 8: 02317–16

Michels AA, Robitaille AM, Buczynski-Ruchonnet D, Hodroj W, Reina JH, Hall MN &
Hernandez N (2010) mTORC1 Directly Phosphorylates and Regulates Human MAF1.
Mol Cell Biol 30: 3749–3757

Miller LH, Ackerman HC, Su X & Wellems TE (2013) Malaria biology and disease
pathogenesis: insights for new treatments. *Nat Med* 19: 156–167

Miller LH, Baruch DI, Marsh K & Doumbo OK (2002) The pathogenic basis of malaria.
Nature 415: 673–679

M'Kaibi FK, Steyn NP, Ochola S & Du Plessis L (2015) Effects of agricultural biodiversity
and seasonal rain on dietary adequacy and household food security in rural areas of
Kenya. *BMC Public Health* 15: 422

Murray L, Stewart LB, Tarr SJ, Ahouidi AD, Diakite M, Amambua-Ngwa A & Conway DJ
(2017) Multiplication rate variation in the human malaria parasite Plasmodium
falciparum. *Sci Rep* 7: 6436

Nacer A, Claes A, Roberts A, Scheidig-Benatar C, Sakamoto H, Ghorbal M, Lopez-Rubio J
& Mattei D (2015) Discovery of a novel and conserved Plasmodium falciparum
exported protein that is important for adhesion of PfEMP1 at the surface of infected
erythrocytes. *Cell Microbiol* 17: 1205–1216

Nyarko PB & Claessens A (2021) Understanding Host–Pathogen–Vector Interactions with
Chronic Asymptomatic Malaria Infections. *Trends in Parasitology* 37: 195–204

Oluboyo AO, Fakologbon OD, Oluboyo BO, Odewusi OO & Ajayi FO (2018) Variations in
levels of selected micronutrients during malaria infection. *J Biomed Sci* 5: 4–9

Perez-Riverol Y, Bai J, Bandla C, García-Seisdedos D, Hewapathirana S, Kamatchinathan S,
Kundu DJ, Prakash A, Frericks-Zipper A, Eisenacher M, *et al* (2022) The PRIDE
database resources in 2022: a hub for mass spectrometry-based proteomics evidences.
Nucleic Acids Research 50: D543–D552

Pluta K, Lefebvre O, Martin NC, Smagowicz WJ, Stanford DR, Ellis H SR, AK S, A B, & M.
(2001) Maf1p, a negative effector of RNA polymerase III in Saccharomyces
cerevisiae. *Mol Cell Biol* 21: 5031–5040

Poullet P, Carpentier S & Barillot E (2007) myProMS, a web server for management and
validation of mass spectrometry-based proteomic data. *Proteomics* 7: 2553–2556

Rana T, Misra S, Mittal MK, Farrow AL, Wilson KT, Linton MF, Fazio S, Willis IM &
Chaudhuri G (2011) Mechanism of Down-regulation of RNA Polymerase III-
transcribed Non-coding RNA Genes in Macrophages by Leishmania. *Journal of
Biological Chemistry* 286: 6614–6626

Rollins J, Veras I, Cabarcas S, Willis I & Schramm L (2007) Human Maf1 negatively
regulates RNA Polymerase III transcription via the TFIIB family members Brf1 and
Brf2. *Int J Biol Sci*: 292–302

Scherf A, Hernandez-Rivas R, Buffet P, Bottius E, Benatar C, Pouvelle B, Gysin J & Lanzer
M (1998) Antigenic variation in malaria: in situ switching, relaxed and mutually
exclusive transcription of var genes during intra-erythrocytic development in
Plasmodium falciparum. *EMBO J* 17: 5418–5426

Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T, Preibisch S,
Rueden C, Saalfeld S, Schmid B, *et al* (2012) Fiji: an open-source platform for
biological-image analysis. *Nat Methods* 9: 676–682

Schneider VM, Visone JE, Harris CT, Florini F, Hadjimichael E, Zhang X, Gross MR, Rhee
974   KY, Ben Mamoun C, Kafsack BFC, *et al* (2023) The human malaria parasite
975   *Plasmodium falciparum* can sense environmental changes and respond by antigenic
976   switching. *Proc Natl Acad Sci USA* 120: e2302152120

977   Serfontein J, Nisbet RER, Howe CJ & de Vries PJ (2010) Evolution of the TSC1/TSC2-TOR
978   Signaling Pathway. *Sci Signal* 3

979   Smith JD, Chitnis CE, Craig AG, Roberts DJ, Hudson-Taylor DE, Peterson DSP, R. N, I. C &
980   Miller LH (1995) Switches in expression of plasmodium falciparum var genes
981   correlate with changes in antigenic and cytoadherent phenotypes of infected
982   erythrocytes. *Cell* 82: 101–110

983   Smith JD, Rowe JA, Higgins MK & Lavstsen T (2013) Malaria's deadly grip: cytoadhesion
984   of Plasmodium falciparum-infected erythrocytes: Plasmodium falciparum
985   cytoadhesion and var genes. *Cell Microbiol* 15: 1976–1983

986   The M, MacCoss MJ, Noble WS & Käll L (2016) Fast and Accurate Protein False Discovery
987   Rates on Large-Scale Proteomics Data Sets with Percolator 3.0. *J Am Soc Mass*
988   *Spectrom* 27: 1719–1727

989   Thomson-Luque R, Votborg-Novél L, Ndovie W, Andrade CM, Niangaly M, Attipa C, Lima
990   NF, Coulibaly D, Doumtabe D, Guindo B, *et al* (2021) Plasmodium falciparum
991   transcription in different clinical presentations of malaria associates with circulation
992   time of infected erythrocytes. *Nat Commun* 12: 4711

993   Valot B, Langella O, Nano E & Zivy M (2011) MassChroQ: a versatile tool for mass
994   spectrometry quantification. *Proteomics* 11: 3572–3577

995   Vannini A, Ringel R, Kusser AG, Berninghausen O, Kassavetis GA & Cramer P (2010)
996   Molecular basis of RNA polymerase III transcription repression by Maf1. *Cell* 143: 70

997   Wahlgren M, Goel S & Akhouri RR (2017) Variant surface antigens of Plasmodium
998   falciparum and their roles in severe malaria. *Nat Rev Microbiol*

999   W.H.O. (2022) World Malaria Report 2022.

1000   Wullschleger S, Loewith R & Hall MN (2006) TOR Signaling in Growth and Metabolism.
1001   *Cell* 124: 471–484

1002   Zhang M, Wang C, Otto TD, Oberstaller J, Liao X, Adapa U SR, K B, IF C, D M, M B, *et al*
1003   (2018) Uncovering the essential genes of the human malaria parasite Plasmodium
1004   falciparum by saturation mutagenesis. *Science*

1005   Zhang Q, Siegel TN, Martins RM, Wang F, Cao J, Gao Q, Cheng X, Jiang L, Hon CC,
1006   Scheidig-Benatar C, *et al* (2014) Exonuclease-mediated degradation of nascent RNA
1007   silences genes linked to severe malaria. *Nature* 513: 431–435

1008   Zhang X & Deitsch KW (2022) The mystery of persistent, asymptomatic Plasmodium
1009   falciparum infections. *Current Opinion in Microbiology* 70: 102231

1010

1011
1012