Scalable querying of human cell atlases via a foundational model reveals

commonalities across fibrosis-associated macrophages

Graham Heimberg[1,2,*,✉], Tony Kuo[3,*], Daryle DePianto[4], Tobias Heigl[5], Nathaniel Diamant[2], Omar

Salem[3], Gabriele Scalia[2], Tommaso Biancalani[2], Shannon Turley[4,5], Jason Rock[5], Héctor Corrada Bravo[6],

Josh Kaminker[1,§,✉], Jason A. Vander Heiden[1,4,§,✉], Aviv Regev[7,§,✉]

Affiliations:

[1] Department of OMNI Bioinformatics, Genentech, South San Francisco, CA 94080, USA.

[2] Department of Machine Learning for Biology, Genentech, South San Francisco, CA 94080,

USA.

[3] Roche Informatics, F. Hoffmann-La Roche Ltd., Mississauga, Canada

[4] Department of Immunology Discovery, Genentech, South San Francisco, CA 94080, USA.

[5] Department of Regenerative Medicine, Genentech, South San Francisco, CA 94080, USA.

[6] Data Science and Statistical Computing, Genentech, South San Francisco, CA 94080, USA.

[7] Research and Early Development, Genentech, South San Francisco, CA 94080, USA.

[*] These authors contributed equally

[§] These authors contributed equally

[✉] email: heimberg@gene.com, regev.aviv@gene.com, vanderheiden.jason@gene.com,

kaminker@gene.com

Abstract

Single-cell RNA-seq (scRNA-seq) studies have profiled over 100 million human cells across diseases, developmental stages, and perturbations to date. A singular view of this vast and growing expression landscape could help reveal novel associations between cell states and diseases, discover cell states in unexpected tissue contexts, and relate *in vivo* cells to *in vitro* models. However, these require a common, scalable representation of cell profiles from across the body, a general measure of their similarity, and an efficient way to query these data. Here, we present SCimilarity, a metric learning framework to learn and search a unified and interpretable representation that annotates cell types and instantaneously queries for a cell state across tens of millions of profiles. We demonstrate SCimilarity on a 22.7 million cell corpus assembled across 399 published scRNA-seq studies, showing accurate integration, annotation and querying. We experimentally validated SCimilarity by querying across tissues for a macrophage subset originally identified in interstitial lung disease, and showing that cells with similar profiles are found in other fibrotic diseases, tissues, and a 3D hydrogel system, which we then repurposed to yield this cell state *in vitro*. SCimilarity serves as a foundational model for single cell gene expression data and enables researchers to query for similar cellular states across the entire human body, providing a powerful tool for generating novel biological insights from the growing Human Cell Atlas.

INTRODUCTION

Characterizing the contexts in which cells employ different expression programs is critical for deciphering their functional role in health and disease. To date, well over 100 million individual cells have been profiled using single-cell or single-nucleus RNA-seq (sc/snRNA-seq) across homeostatic, disease, and perturbed conditions[1]. Individually, even the largest multi-tissue scRNA-seq atlases[2–4] capture only a relatively small portion of cell states across human tissues; collectively, these atlases provide a vast, pan-human view of cell and disease biology that has the potential to address fundamental questions about human biology[1]. By aggregating across atlases, we may uncover biological insights and enable investigations into cell states that are common across multiple studies of the same organ and conditions (*e.g.*, similar neural progenitor populations across independent studies of brain development), different organs and conditions (*e.g.*, inflammatory fibroblasts in both ulcerative colitis and cancer); or between the human body and *in vitro* lab models (*e.g.*, regulatory T cells genetically perturbed to recapitulate *in vivo* cells from diseased tissue).

Despite this promise and the rapid growth in data, our ability to realize the potential of cross-datasets, pan-body, analyses remains limited and hampered by the need for laborious manual curation, harmonization, and dataset aggregation by expert analysts, as well as the painstaking process of selecting datasets, standardizing cell type annotations, and finding a common low-dimensional representation. As a result, most aggregation efforts have been limited in their biological scope and number of datasets, with some recent notable exceptions focused on genes rather than cell representation[5–8].

To leverage and query the massive scale and richness of available single-cell atlases, we need both (1) a foundational model of cell states with an effective representation for single-cell profiles across different cell types and conditions that can be used across many applications without retraining; and (2) a measure of cell similarity that is robust to technical noise, scales to hundreds of millions of cells, and accurately generalizes to datasets and cell states not observed in the training. Established unsupervised methods to learn low dimensional representations of scRNA-seq profiles, such as Principal Component Analysis (PCA) or autoencoders[3,9–11], faithfully preserve information from the input[3,9–11] and may even eliminate technical variation for explicitly defined batches. However, they do not learn general features that encode relationships between cells needed to represent and query new data sets in the context of cross-study, pan-tissue biological variation.

Machine learning methods, metric learning in particular, have successfully learned representations for diverse entities and a measure of similarity between them, especially in image analysis. For example, metric learning models for facial recognition are explicitly trained to embed images of the same person closer together than images of different people, by exploiting visual features that are critical to distinguish individuals[12]. Once trained, images are embedded into a low-dimensional space, where distances between images represent a measure of similarity based on the learned features. Users can then query with an image not in the training set to find additional similar images that are nearby in the latent space and depict the same person. We reasoned that, analogously, metric learning could provide a meaningful representation of and similarity metric for cell profiles. By training a model using annotated scRNA-seq data, we can learn a low dimensional representation that places similar cells near each other and dissimilar cells farther apart. If learned

from a sufficient diversity of cell profiles, such a representation would, in turn, provide a

foundational model of cells and would allow efficient searches for cells with similar expression

states (**Fig. 1a**).

Here, we introduce SCimilarity, a class of deep metric learning models that quantify similarity

between single-cell expression profiles (SCimilarity score) and provide a single-cell gene

expression foundational reference model to systematically query for comparable cell states across

tissues and diseases. SCimilarity uses a training set of diverse author-annotated cell profiles to

learn a universal representation and distance metric that facilitates efficient searches across a

massive reference meta-atlas for the most expression-similar cells. To train a foundational model

that can be broadly applied to many applications across tissues and studies, we built a

programmatic pipeline for massive data import and automated standardized curation and used it to

assemble a corpus of 22,699,774 cells from 399 datasets spanning a broad range of organs, systems

and conditions across the human body. After training and testing on a subset of 66 studies and

7.9M single-cell profiles, the learned models generalize well, representing and quantifying

similarities between 14.9M cells from another 347 studies excluded from training. By tuning a

single parameter during SCimilarity's training, we yield models optimized for either data

integration and visualization of millions of cells across hundreds of studies, or for fast and efficient

(millisecond) queries of a new cell state across tens of millions of cells. Finally, we illustrate the

power of SCimilarity by querying for a fibrosis-associated macrophage (FMΦ) subset previously

identified in interstitial lung disease (ILD), finding comparable cell populations (but with different

annotated names and signatures) in other ILD studies, as well as in new contexts, including

COVID-19, different tumors including pancreatic ductal adenocarcinoma (PDAC), and even

healthy lung (at low abundance). Surprisingly, SCimilarity recovered FMΦ-like cells among PMBCs cultured and stimulated in a 3D hydrogel system *in vitro*, which we experimentally validated, producing an FMΦ-like *in vitro* cell system for future functional studies of a tissue-resident cell state. Overall, SCimilarity preserves expression diversity across cells in an integrated foundational model of a human cell atlas, and allows a novel scaled cell search across organs, systems, and conditions, as a powerful framework for generating biological insights and experimentally testable hypotheses.

RESULTS

SCimilarity: novel similarity metrics and representations for single-cell expression profiles

SCimilarity is a family of models that blend unsupervised representation learning and supervised metric learning, through simultaneously optimizing two objectives : (1) a supervised triplet loss function, which is used to embed expression profiles from matching cell types close together, effectively integrating cells of the same type across studies[13–15], and (2) an unsupervised mean squared error (MSE) reconstruction loss function, which encourages the model to preserve variation from the input expression profiles, capturing subtler differences in expression patterns within cells of the same type, such as those related to tissue residency of immune cells (**Fig. 1b, Methods**). The balance of these two objectives, set by a single hyperparameter, β, determines the properties of the representation (**Methods**). Increasing the relative weight of the triplet loss function improves dataset integration, while increasing the relative weight of the reconstruction loss improves querying performance. Therefore, different loss function weightings within the same model architecture can address different applications.

We train SCimilarity with tens of millions of cell triplets sampled from data with author-provided

standardized cell type annotations from the Cell Ontology[16] (**Fig. 1b**, **Methods**). Specifically, each

training triplet consists of similar anchor and positive cells (*i.e.*, same cell type) from different

studies, while anchor and negative cells are dissimilar (*i.e.*, distinct cell types; from the same or a

different study). However, even with standardized Cell Ontology terms, some cell type

comparisons are ambiguous due to arbitrary differences in annotation granularity across studies

(*e.g.*, it is ambiguous if cells annotated as "T cell" in one study and "CD4$^+$ T cell" in another are

similar or dissimilar). To address this, SCimilarity excludes cell pairings with such vertical

ancestor-descendant Cell Ontology relationships from training triplets, and learns only from cells

that are either explicitly similar or unambiguously dissimilar (**Fig. 1b, Methods**). By sampling

only unambiguous triplets we eliminate the need to manually flatten or harmonize every cell type

annotation and are able to seamlessly scale the training set across dozens of studies.

A learned SCimilarity representation of 22.7M cells across dozens of tissues and disease

datasets collated by an automated curation and processing pipeline

To test SCimilarity models, we assembled a compendium of sc/snRNA-seq datasets across human

biology. We focused on studies generated with one experimental platform (10x Genomics

Chromium droplet-based scRNA-seq) and data publicly available on the Gene Expression

Omnibus (GEO)[17] or CELLxGENE[18]. These data capture much of the published scRNA-seq data,

and were generated with similar library preparation protocols and computational pipelines[19]. There

were 753 human sc/snRNA-seq datasets matching our search criteria and keywords as of March

23rd, 2021 (with Biopython Entrez[20], **Methods**). The number of samples and cells matching our

criteria has at least doubled every 6 months between December 2018 and March 2021; (**Extended**

Data Fig. 1a,b). We programmatically downloaded 13,401,599 cell profiles from 333 of the

identified studies with their respective GEO metadata and unnormalized gene count matrices

(**Methods, Extended Data Table 1**). We manually ingested another 66 well-annotated studies

from either the CELLxGENE portal[18] or from large studies and consortia not available through

GEO that passed the same dataset filtering criteria (**Methods**). Overall, we assembled a corpus of

399 studies comprising 22,699,774 cells from 33,815 tissue samples with 184 unique Tissue

Ontology terms[21], 132 Disease Ontology terms[22], and 204 Cell Ontology cell type terms[16], with

each Cell Ontology term appearing in at least two separate datasets (**Fig. 2a, Extended Data Fig.**

1c, Extended Data Table 1).

We trained SCimilarity models with a training set of 7,913,892 single-cell profiles from 52 studies

with Cell Ontology author annotations that reflected a diversity of conditions and tissues

(**Extended Data Fig. 1d**, **Extended Data Table 1**), sampling 50,000,000 of the most informative

triplets (**Methods**). We withheld 14 studies comprising 1,384,283 cells with Cell Ontology

annotations for testing the learned representation and metric (**Fig. 2a**). We excluded tumor, cell

lines, and iPSC-derived samples from the training and test sets, because cell identity of tumor cells

and cell lines can be ambiguous.

Tuning of SCimilarity's reconstruction and triplet loss functions yields models optimized for

integration *vs*. cell search tasks

We examined six different blends for SCimilarity's objective function, varying the relative

weighting of the reconstruction and triplet loss functions, and finding that the two loss function

components gave rise to different behaviors in a trained model. Briefly, we assessed the models

on two tasks – data integration and searching for cells similar to a query profile – using studies

entirely held out from training. To evaluate data integration, we quantified how coherently cells of

each type are clustered and how distinct each cell type cluster is from other clusters. To this end,

we created an ontology-aware variation of average silhouette width[23] to quantify integration

capabilities across datasets without harmonizing cell type annotations (**Methods**). To evaluate our

cell search distance metric, we compared searches with SCimilarity to gene signature scoring

(**Methods**). The higher the correlation between these two quantities, the more our similarity metric

corresponds to traditional signature-based similarity to represent a cell state of interest.

Models with higher triplet loss weighting scored higher on integration benchmarks, while models

with higher reconstruction loss weighting encoded distances between cells in a manner better

correlated with differences in representative expression signature scores (**Extended Data Fig.**

2a,b). Pure triplet loss, which is calculated at the level of cell type labels, does not reliably preserve

subtle cell state differences, such as tissue specificity or disease response within cells of the same

type. Mean squared error reconstruction loss complements this by preserving more subtle gene

expression patterns, while the triplet loss ensures that cells of the same type are embedded closely

together. Based on the biological question, a user can tune this balance to yield the highest utility.

We thus pursued two SCimilarity models: an integration model, optimized for the task of learning

a low dimensional representation that groups cells by type rather than by study; and a cell search

model that is optimized for the task of retrieving cells with an expression state similar to that of a

query cell across hundreds or thousands of scRNA-seq datasets (**Extended Data Fig. 2a**,

Methods).

SCimilarity's latent space representation filters outlier cells and integrates test datasets

without batch correction

We next benchmarked if SCimilarity's latent space representation from the integration model generalizes well to cells from entire datasets held out of training compared to other methods. In a low-dimensional embedding, unannotated cell profiles from nine lung studies (7 training set, 2 test set) visually intermix well when embedded into SCimilarity's learned 128-dimensional space (**Extended Data Fig. 2a**). SCimilarity's data integration model scored higher than Harmony, scVI, and scArches on integration tasks by the ontology-adjusted ASW measure of cluster coherence, but scored lower for normalized mutual information (NMI) and adjusted rand index (ARI), which measure the extent of study mixing within each cluster (**Extended Data Fig. 2b**). Thus, without directly training on the full data set or performing additional batch correction, the integration model clusters cells by type rather than study at a level that is competitive with existing methods trained directly on the data. This demonstrates that the triplet loss learns features that capture meaningful biology, while reducing technical sources of noise and avoiding overfitting to the training set.

SCimilarity quantifies a confidence level for each cell's representation, providing both outlier detection and an assessment of the representation's relevance in the context of new data. When computing the representation of a new cell, the further outside the scope of model training it is, the harder it is for the model to accurately represent it. Using SCimilarity's score to quantify how distant a query cell is from the training data distribution provides a heuristic about the quality and scope of the representation – a cell scoring as highly similar to cells seen during training can be confidently represented by the model. Overall, 79.5% of *in vivo* holdout cells had high

representation confidence. Tissue samples with particularly low representation confidence, such

as stomach (n = 0 training studies), fetal gut (n=1), and bladder (n=0) were either absent or poorly

represented in training (**Methods, Extended Data Fig. 2c**), suggesting that more labeled training

datasets from those tissues could improve the model's representation. Similarly, 43.8% of *in vitro*

cell profiles were considered low confidence due to poor matching to the training set (which

excluded *in vitro* samples).

We combined SCimilarity's ability to generalize to new datasets and its confidence-based filtering

to systematically generate meta-atlases for 21 different human tissues without labor-intensive

dataset harmonization and no additional training (**Fig. 2b**). If datasets have already been embedded

using SCimilarity, this task only requires concatenation of cells of interest and standard

visualization workflows.

SCimilarity assigns an unannotated query cell to a cell type by finding similar cells in a

labeled reference

We next used SCimilarity to find the cells in the annotated reference that are most similar to an

unannotated query cell profile, and then annotate the query cell accordingly (**Fig. 3a, Methods**).

This approach is distinct from established annotation methods in that it (1) relies on a large, pan-

human annotated cell repository, (2) employs a measure of expression similarity, and (3) classifies

at the single cell rather than cluster level, providing greater transparency into the classification

itself. Thus, users can see which individual cells, studies, and tissues are driving the classification

decision. Moreover, since each cell is annotated independently, no clustering or associated

parameter selection, such as the number and resolution of clusters, are required. A user can choose

to annotate a cell's profile by comparing it either to a desired subset of cell types (*e.g.*, for a tissue-

specific query) or to the entire annotated cell reference. Because SCimilarity is built using metric

learning, finding the most similar cells is the same as retrieving the query cell's nearest neighbors.

This operation is extremely efficient with the hnswlib algorithm[24], where searching a precomputed

approximate nearest neighbor index of all the annotated reference cells in SCimilarity's latent

space takes just 20 milliseconds (**Methods**). Low SCimilarity scores to reference cells flag an

outlier query cell, which may be either a cell type that is not within the reference or a query cell of

low quality.

SCimilarity quickly and accurately assigned cell types for entire datasets held out from training,

as well as for the rest of the 22.7M cell corpus. When limiting potential cell types to author-selected

labels, 94.5% of SCimilarity's predicted labels from healthy kidney samples[25] match the author-

provided cell type annotations (**Fig. 3b-d**, **Methods**). In some cases, where SCimilarity's

predictions did not match author-provided annotations, SCimilarity's predictions were more

accurate or granular. For example, 94% of the cells that the authors[25] annotated as CD4$^+$ T cells

but SCimilarity annotated as CD8$^+$ T cells express *CD8A* or *CD8B* (and none express *CD4*),

supporting SCimilarity's annotation (**Fig. 3e-h**). Separately, when allowing cells to be annotated

as any cell type in the repository, 6.3% of the author-annotated CD4$^+$ T cells were reannotated by

SCimilarity as regulatory T cells (T$_{regs}$) (**Extended Data Fig. 3a**), most of which (85.2%)

expressed at least one T$_{regs}$ marker (FOXP3, IL2RA, or IKZF2, **Extended Data Fig. 3b-d**).

Similarly, 1.8% of author-annotated mesenchymal stem cells (**Fig. 3b**) were reassigned by

SCimilarity as myofibroblasts (**Extended Data Fig. 3a**) and 93% of those express the

myofibroblast-associated gene *ACTA2* (**Extended Data Fig. 3e**). Cell type prediction was rapid,

taking 3-5 seconds to embed and annotate 10,000 cells from a dataset. Overall, across all 14 test

datasets spanning 78 Cell Ontology terms, 71% of the cell populations had high agreement (>85%

of the cell population) between author and SCimilarity annotations (**Fig. 3i**). SCimilarity

performed poorly on one dataset (Cano-Gomez et al.[26]), due to fine granularity and redundancy of

author labels (*e.g.*, CD4$^+$ αβ T cells, helper T cell, memory T cell, naive T cell, and regulatory T

cell).

We used SCimilarity's cell type assignment to rapidly annotate all 22.7M cell profiles in one

common model, newly-annotating 13,401,599 profiles and reannotating 9,298,175 author-

annotated profiles (**Methods**) to a single set spanning 74 cell type labels (21 coarser lineages) from

25 simplified tissue categories (**Fig. 4a,b, Extended Data Fig. 3f**). A consistent annotation across

datasets facilitates cross-study and cross-tissue analyses of one cell type or lineage, as SCimilarity

can extract cells from hundreds of studies, aggregating vast biological diversity across one cell

type. For example, we readily aggregated 1,172,325 fibroblasts and myofibroblasts (**Extended**

Data Fig. 3g) and 2,507,879 monocytes and macrophages (**Extended Data Fig. 3h**) from

hundreds of studies profiling different primary tissue samples.

SCimilarity's representations comprise of interpretable biological features

To interpret SCimilarity's annotations, we quantified the importance of each gene for cell type

annotations assigned by the foundational query model using Integrated Gradients, a method that

identifies the impact on model predictions of small disturbances to the input expression profiles

(**Methods**). For example, the top gene attributions that distinguish lung alveolar type 2 (AT2) cells

are surfactant genes *SFTPA2*, *SFTPA1*, *SFTPB*, and *SFTPC*, consistent with known AT2 cell

function[27]. SCimilarity learned these without prior knowledge of cell type specific genes,

signatures, or highly variable genes. Overall, SCimilarity's top importance genes agreed well with

differentially expressed marker genes for 17 different matched types[3] with the exception of rare

neuroendocrine cells (average AUC=0.84, **Extended Data Table 2, Extended Data Fig. 3i**).

Thus, SCimilarity's representation captured known and validated biological markers within its

features.

Cell search identifies fibrosis-associated macrophages across tissues and diseases

With a single representation and common definition of cell types, we hypothesized that

SCimilarity could help elucidate the role of tissue-resident immune cells. As a case study, we

focused on macrophages, given their remarkable plasticity in cell states and their important

specialized roles in tissue repair, regeneration, and fibrosis[28,29]. Recent scRNA-seq studies in

fibrotic diseases, including lung fibrosis, cancer, obesity, and COVID-19 have reported seemingly-

similar *SPP1*⁺ fibrosis-associated macrophage (FMΦ) populations[30–38]. However, because each

study identified them independently, using different nomenclatures and marker gene signatures to

define subsets, it is unclear how similar these cell states are. Moreover, it is unknown how broadly

associated such cell states are with other diseases, especially those with prominent fibrosis. We

reasoned that SCimilarity's cell search should allow us to query our corpus with an FMΦ cell

profile from one study to identify similar cells across other tissues and conditions, thereby

clarifying the cell identity of similarly-described cells and the conditions in which FMΦ arise (**Fig.**

5a).

We queried our model with the FMΦ cell profile, searching for similar cells across 2,578,221 cells

annotated by SCimilarity as monocytes or macrophages in the 22.7M cell corpus (**Fig. 5a**).

SCimilarity queries can use either an individual cell profile or a centroid of multiple cell profiles.

Here, we input the centroid profile of a macrophage cell subset from Adams *et al.*[30] that we defined

using a gene signature consisting of the extracellular matrix remodeling and fibrosis-associated

genes *SPP1*, *TREM2*, *GPNMB*, *MMP9*, *CHIT1*, and *CHI3L1* (**Methods**). In two seconds,

SCimilarity exhaustively computed the pairwise similarity of our query profile to each of the 2.6M

in vivo profiles of the cells it annotated as monocytes or macrophages in our corpus (**Fig. 5b** and

Extended Data Fig. 4a). Alternatively, simply identifying the 10,000 cells with the highest

SCimilarity score takes 0.05 seconds (**Methods**). By comparison, a more conventional approach

that scores each cell in the corpus with a literature-defined FMΦ gene signature took 2 hours and

46 minutes (**Extended Data Fig. 4b**). The gene signature and SCimilarity scores are broadly

correlated ($r = 0.50$, $p < 10^{-300}$, **Extended Data Fig. 4a-c**), showing that the granular cell state,

not just the cell type, is well-represented in SCimilarity query score and embedding.

The SCimilarity search showed that FMΦs are common in ILD lung samples in our compendium,

as well as present in some cancers, including uveal melanoma, pancreatic ductal adenocarcinoma

(PDAC), and colon cancer (**Fig. 5c-e, Extended Data Table 3**). Of the top 1% of monocytes and

macrophages most similar to our query, 99.1% were from lung tissue and 87.2% from ILD and

COVID-19 lung samples. The prevalence of FMΦ-like cells in the lung varied by disease: the

proportion of monocytes and macrophages that were FMΦ-like was 20% and 4% in two systemic

sclerosis (SSc) studies, 6.1% on average (SE = 1.4%) across 13 ILD studies (excluding SSc), 1.2%

on average across seven COVID-19 lung studies (SE = 0.5%, 0% in non-lung COVID-19 data)

and 0.4% in 19 studies annotated as "healthy", "normal" or with no disease annotation (SE =

0.2%). While abundant in SSc lung, FMΦ-like cells were much rarer (0.14% of myeloid cells) in

SSc skin[39]. There were some FMΦ-like cells in other fibrotic diseases and tissues, such as one

primary pancreatic ductal adenocarcinoma (PDAC) tumor[40] (0.85% of 1,171 myeloid cells) and

one liver metastasis[41] of PDAC (0.5% of 1,199 cells). Thus, while our query FMΦ profile was

derived from IPF samples, it uncovered FMΦ-like cells in many contexts, including SSc-ILD,

COVID-19 lung and PDAC. These results confirm previous observations of FMΦs in lung

injury[38,42] and suggest a role for FMΦ-like cells across other organs and diseases.

Integrated gradients analysis reveals commonalities between SCimilarity score and

established gene signatures

Because FMΦ-like cells are detected by SCimilarity across many ILD studies, we hypothesized

that the cells captured by different marker genes and nomenclature in different studies refer to the

same biological cell state. To test this, we applied integrated gradients to quantify each gene's

importance when SCimilarity distinguishes FMΦs from randomly sampled monocytes and

macrophages (**Methods**). The genes identified as important for distinguishing FMΦs are enriched

in key fibrotic processes, including extracellular matrix remodeling (*MMP7*, *MMP9*, *FN1*, *SDC2*,

SPARC, *SPP1*), lipid metabolism and lipoprotein clearance (*APOC1*, *APOE*, *LPL*, *LIPA*), and

damage-associated molecular pattern recognition (*MARCO*, *MSR1*) (**Fig. 5f**, **Extended Data Fig.**

4d,e, **Extended Data Table 4**). While SCimilarity found many FMΦ marker genes that were

already discussed in the literature, such as TREM2 (**Extended Data Fig. 4f**), it also identified

novel genes elevated in FMΦs such as HLA-DQA1 and RGS1 (**Extended Data Fig. 4g,h**).

The genes with the highest importance scores in the SCimilarity embedding of FMΦs significantly

overlap (p<6.7x10^{-13}) with published gene signatures describing similar macrophage populations

or with genes whose differential expression defined each study's macrophage population of

interest (**Extended Data Table 5**). While cell signatures from IPF lung had a high signature match

(AUC ≥ 0.95), the negative control signatures of M1 and M2 macrophages[43] had lower ones at the

bottom three (AUC = 0.85 (2.65x10^{-2}) and AUC=0.92 (p<4.92x10^{-6}), respectively; **Fig. 5f**,

Extended Data Fig. 4d).

FMΦ-like cells identified among *ex vivo* stimulated peripheral blood mononuclear cells

(PBMCs) help establish a novel human cell model

Research to understand the role of a novel cell state or subtype in disease, such as FMΦs, benefits

greatly from the ability to model, perturb, and study the cells *in vitro*. However, there is currently

no systematic way to identify *in vitro* culture conditions that generate cells that match cells

identified *in vivo*. To accelerate development of an *in vitro* FMΦ system, we used SCimilarity to

search for FMΦ-like cells across *in vitro* stimulated samples with the goal of identifying previously

employed experimental conditions that might resemble the tissue cell state. We filtered our full

reference cell collection for *in vitro* and *ex vivo* studies containing at least 50 monocytes or

macrophages, resulting in 41,926 monocytes and macrophages across 40 samples from 17 such

studies. These span diverse and complex conditions, such as lung organoids infected with SARS-

CoV-2[44], *ex vivo* treated acute myeloid leukemia samples[45], or PBMCs stimulated with morphine

and lipopolysaccharide[46].

The cells most similar to our query FMΦ expression profile were monocytes grown as part of a heterogenous PBMC culture for 5 days in a 3D hydrogel culture system that was designed for expansion of hematopoietic stem cells (HSCs) from PBMCs[47] (**Fig. 6a, Extended Data Table 6**). This study is unrelated to lung biology and its authors did not report any results for myeloid cells. Nevertheless, while no FMΦ-like cells were present among myeloid cells on day 0, 15% of cells grown for five or more days in this system were highly similar to FMΦs (SCimilarity score >25) and expressed *TREM2*, *GPNMB*, *CCL18* and *MMP9* (**Fig. 6b-e**). This was a surprising result, because of the seeming irrelevance of the study to fibrosis or macrophage biology and the rarity of FMΦ-like cells in PBMC samples *in vivo*.

To validate SCimilarity's prediction of an FMΦ-like cell culture condition, we used a similar protocol to replicate the 3D hydrogel system[47], followed by scRNA-seq to assess the yield of FMΦ-like cells (**Fig. 6b,c,f**). While relative cellular abundances differed between the original day 5 data (Xu et al, 2022) and our day 8 replication of the same conditions (**Methods**), 10.1% of all cells in the Day 8 experiment were predicted as HSCs by SCimilarity (**Fig. 6g**). Moreover, 41.5% of the myeloid cells in day 8 validation experiments from three donors were predicted as FMΦ-like macrophages (**Fig. 6b,f**, 37.1%, 42.5%, and 44.9%; SCimilarity score > 25). Furthermore, FMΦ hallmark genes, such as *CCL18*, *GPNMB*, *SPP1*, and *TREM2*, were enriched in the myeloid compartment of our replicate experiment compared to day 0 conditions (**Fig. 6c**). This experiment validates that an FMΦ-like population can be generated from PBMCs in culture conditions. Taken together, these results demonstrate SCimilarity's ability to interrogate publicly available data at scale, query a reference of *in vivo* and *in vitro* data for biologically similar conditions, and help identify experimental conditions to reproduce those results in laboratory settings.

DISCUSSION

To date, more than a hundred million human cells have been profiled across tissues in health and

disease, and such data continue to grow exponentially. This growing human cell atlas should be

the starting point for researchers aiming to readily search, query and compare cell states of interest

across different protocols, treatments, tissues, and diseases.

SCimilarity systematically annotates and repurposes tens of millions of expression profiles from

hundreds of studies, to create an integrated, searchable and queryable foundational model of pan-

human cellular diversity. SCimilarity is comprised of three key features: (1) a 22.7M cell human

scRNA-seq data repository (at present), (2) a foundational model for single cell gene expression

with a generalizable embedding and similarity metric (which could readily be retrained for larger

datasets), and (3) methods to efficiently query across this entire pan-body human cell atlas.

Together, these provide new context, capabilities, and workflows for extracting insights from new

and existing scRNA-seq datasets in the human cell atlas and other atlases. SCimilarity's

framework architecture can easily accommodate quick updates as data continue to grow.

Because SCimilarity can generalize to cells and datasets not seen in the training, cell profiles can

be added as entirely new studies or removed by applying new cell filters without recomputing the

low dimensional representations. This flexibility allows us to change the analysis' scope at any

point without redoing work, enabling modularized workflows for scRNA-seq analysis.

Downstream tasks, such as cell type annotation, cell queries, and gene signature derivation all are

simplified using SCimilarity's generalized low dimensional representation and can be applied to

cells not seen during training without informing the model about the importance or variability of

specific genes during training. Outlier detection helps both filter out technical errors and highlight

potentially novel cell subsets. Although generalized models that do not require recomputing low

dimensional representations would alleviate time and expertise barriers that currently impede

researchers, to the best of our knowledge, generalization has rarely been optimized in single-cell

expression analysis.

There is no single objective measure of similarity, or dissimilarity, between cellular profiles.

Curated gene signatures are useful when a small number of explanatory genes are sufficient to

define a cell state. SCimilarity uses the full expression profile of a cell as its query, defined by

either a single representative cell or the centroid of a set of cell profiles. Thus, SCimilarity's cell-

based search bypasses the manual curation requirements and biases inherent in defining a gene

signature. In cases where such a gene signature is desired, SCimilarity can compute a robust

signature for a cell state across studies.

Exploration of transcriptionally-similar populations across a vast atlas of human scRNA-seq data

provides critical context to a cell population of interest. First, observing a query population across

many similar studies shows that the original observation was reproducible, a key for subsequent

scientific research[48]. Second, SCimilarity queries can connect results from independent studies.

While one study may find a cell population in a disease, another may show similar cells with

functional characterization, allowing us to formulate a new hypothesis on the functional properties

of disease-associated cells.

This is illustrated by how SCimilarity allowed us to search for and identify FMΦ-like cells across

tissues and disease states, construct a cross-study set of explanatory marker genes, and uncover a

cell culture system that elicits a similar FMΦ-like state *in vitro*. Modeling FMΦs from readily

available PBMCs is exceptionally valuable, because isolation of cells from human lung explants

is prohibitive for many functional assays. Surprisingly, in addition to fibrotic lung, FMΦs were

present in multiple tumor types, particularly PDAC, a heavily fibrotic cancer, where macrophages

play an important role in mediating the associated fibrosis and have been linked to tumor

progression[49]. The identification of a common FMΦ state across fibrosis, cancer, and infection

suggests a broader role for these cells in the damage response and tissue remodeling processes

across diseases. Moreover, SCimilarity's search identified FMΦ-like cells in an *in vitro* study– an

observation that could not have been gleaned by reviewing the paper or based on the description

of the culture system – but that we validated in the lab. The variations we observed between the

original and replicate *in vitro* experiment may be attributed to differences in culture duration, cell

extraction from the hydrogel, lymphocyte proportions, or other batch effects. Furthermore, these

results invite new hypotheses, such as whether the 3D hydrogel provides key ECM-like

environmental cues that promote an FMΦ-like state and induction of remodeling genes, such as

MMP9 and *SPP1*, and which factors can be added to drive an even stronger FMΦ phenotype. Thus,

SCimilarity provides a powerful framework to iteratively generate and validate such experimental

hypotheses.

SCimilarity is not appropriate for all applications and will need further improvements to continue

to scale with exponential data growth and to more comprehensively span human biology as the

Human Cell Atlas continues to grow. Training SCimilarity requires Cell Ontology labels.

Fortunately, scRNA-seq data sharing practices are increasingly relying on using the Cell Ontology

for standardization. However, the Cell Ontology itself is a large, yet incomplete, effort. Cell states

are only considered in training if they are recognized in the Cell Ontology, and the number of these

states is growing rapidly. Furthermore, while we trained SCimilarity on vast amounts of data,

cancer cells and cell lines were deliberately withheld from training due to lack of clear cell type

identity and therefore may not be well represented. In addition, in our experience, we see poor

performance on fetal samples, likely due to most of the training data being sourced from adult

tissues.

The current data integration and cell search models provide generalizable representations of 22.7M

single-cell profiles across the human body, and include a Python API for querying cell profiles of

interest. Future improvements to SCimilarity could include pre-training on the massive amounts

of unlabeled data, effectively exposing the model to more cell states and more technical variability

during training. With effective representations we can more easily combine embeddings to include

other species or data modalities. We believe that SCimilarity brings a new framework to single-

cell genomics, enabling re-use of rich public data resources through instantaneous queries and

demonstrates how this can be used to provide novel biological insights.

Contributions:

GH conceived of the method with input from AR, JVH, HCB, and JK. GH and TK performed

data ingest and model implementation with input from JVH, ND, GS, TB, JK and AR. Python

API was developed by TK with help from JVH, OS, and GH. Interpretability was developed by

ND and GS with input from HCB and TB. JVH conceived of biological application of method

with input from GH, ST, JR and DD. DD and TH performed experimental validation with

guidance from JR and ST. GH wrote the manuscript with input from JVH, JK, AR and HCB.

Acknowledgements:

We thank Anupriya Tripathi for coming up with the name "SCimilarity" and Jenna Collier,

Gokcen Eraslan, John Marioni, and Jake Freimer for their suggestions that strengthened the

manuscript.

Competing interests:

All authors are employees of Genentech or Roche. A.R. is a co-founder and equity holder of

Celsius Therapeutics, an equity holder in Immunitas, and until July 31, 2020 was an S.A.B.

member of Thermo Fisher Scientific, Syros Pharmaceuticals, Neogene Therapeutics and

Asimov.

References

1. Rood, J.E., Maartens, A., Hupalowska, A., Teichmann, S.A., and Regev, A. (2022). Impact
of the Human Cell Atlas on medicine. Nat. Med. *28*, 2486–2496.

2. Domínguez Conde, C., Xu, C., Jarvis, L.B., Rainbow, D.B., Wells, S.B., Gomes, T., Howlett,
S.K., Suchanek, O., Polanski, K., King, H.W., et al. (2022). Cross-tissue immune cell analysis
reveals tissue-specific features in humans. Science *376*, eabl5197.

3. Eraslan, G., Drokhlyansky, E., Anand, S., Fiskin, E., Subramanian, A., Slyper, M., Wang, J.,
Van Wittenberghe, N., Rouhana, J.M., Waldman, J., et al. (2022). Single-nucleus cross-tissue
molecular reference maps toward understanding disease gene function. Science *376*,
eabl4290.

4. Tabula Sapiens Consortium*, Jones, R.C., Karkanias, J., Krasnow, M.A., Pisco, A.O., Quake, S.R., Salzman, J., Yosef, N., Bulthaup, B., Brown, P., et al. (2022). The Tabula Sapiens: A multiple-organ, single-cell transcriptomic atlas of humans. Science *376*, eabl4896.

5. Rosen, Y., Brbić, M., Roohani, Y., Swanson, K., Li, Z., and Leskovec, J. (2023). Towards Universal Cell Embeddings: Integrating Single-cell RNA-seq Datasets across Species with SATURN. bioRxiv. 10.1101/2023.02.03.526939.

6. Cui, H., Wang, C., Maan, H., Pang, K., Luo, F., and Wang, B. (2023). scGPT: Towards Building a Foundation Model for Single-Cell Multi-omics Using Generative AI. bioRxiv, 2023.04.30.538439. 10.1101/2023.04.30.538439.

7. Theodoris, C.V., Xiao, L., Chopra, A., Chaffin, M.D., Al Sayed, Z.R., Hill, M.C., Mantineo, H., Brydon, E.M., Zeng, Z., Liu, X.S., et al. (2023). Transfer learning enables predictions in network biology. Nature. 10.1038/s41586-023-06139-9.

8. Shen, H., Shen, X., Hu, J., Liu, J., Zhang, C., Wu, D., Feng, M., Yang, M., Li, Y., Yang, Y., et al. (2022). Generative pretraining from large-scale transcriptomes: Implications for single-cell deciphering and clinical translation. bioRxiv, 2022.01.31.478596. 10.1101/2022.01.31.478596.

9. Heimberg, G., Bhatnagar, R., El-Samad, H., and Thomson, M. (2016). Low Dimensionality in Gene Expression Data Enables the Accurate Extraction of Transcriptional Programs from Shallow Sequencing. Cell Systems *2*, 239–250.

10. Lopez, R., Regier, J., Cole, M.B., Jordan, M.I., and Yosef, N. (2018). Deep generative modeling for single-cell transcriptomics. Nat. Methods *15*, 1053–1058.

11. Lotfollahi, M., Naghipourfar, M., Luecken, M.D., Khajavi, M., Büttner, M., Wagenstetter, M., Avsec, Ž., Gayoso, A., Yosef, N., Interlandi, M., et al. (2022). Mapping single-cell data to reference atlases by transfer learning. Nat. Biotechnol. *40*, 121–130.

12. Schroff, F., Kalenichenko, D., and Philbin, J. (6/2015). FaceNet: A Unified Embedding for Face Recognition and Clustering. 2015 IEEE Conference on Computer Vision and Pattern Recognition (CVPR), 815–823.

13. Simon, L., Wang, Y.-Y., and Zhao, Z. (2021). Integration of millions of transcriptomes using batch-aware triplet neural networks. Nature Machine Intelligence *3*, 1–11.

14. Yang, M., Yang, Y., Xie, C., Ni, M., Liu, J., Yang, H., Mu, F., and Wang, J. (2022). Contrastive learning enables rapid mapping to multimodal single-cell atlas of multimillion scale. Nature Machine Intelligence *4*, 696–709.

15. Yu, X., Xu, X., Zhang, J., and Li, X. (2023). Batch alignment of single-cell transcriptomics data using deep metric learning. Nat. Commun. *14*, 960.

16. Diehl, A.D., Meehan, T.F., Bradford, Y.M., Brush, M.H., Dahdul, W.M., Dougall, D.S., He, Y., Osumi-Sutherland, D., Ruttenberg, A., Sarntivijai, S., et al. (2016). The Cell Ontology

2016: enhanced content, modularization, and ontology interoperability. J. Biomed. Semantics
7, 44.

17. Edgar, R., Domrachev, M., and Lash, A.E. (2002). Gene Expression Omnibus: NCBI gene
expression and hybridization array data repository. Nucleic Acids Res. *30*, 207–210.

18. Chan Zuckerberg CELLxGENE Discover (2022). Cellxgene Data Portal.

19. Zheng, G.X.Y., Terry, J.M., Belgrader, P., Ryvkin, P., Bent, Z.W., Wilson, R., Ziraldo, S.B.,
Wheeler, T.D., McDermott, G.P., Zhu, J., et al. (2017). Massively parallel digital
transcriptional profiling of single cells. Nat. Commun. *8*, 14049.

20. Cock, P.J.A., Antao, T., Chang, J.T., Chapman, B.A., Cox, C.J., Dalke, A., Friedberg, I.,
Hamelryck, T., and Kauff, F. (2009). Biopython: freely available Python tools for
computational molecular biology and bioinformatics. Bioinformatics.

21. Gremse, M., Chang, A., Schomburg, I., Grote, A., Scheer, M., Ebeling, C., and Schomburg,
D. (2011). The BRENDA Tissue Ontology (BTO): the first all-integrating ontology of all
organisms for enzyme sources. Nucleic Acids Res. *39*, D507-13.

22. Schriml, L.M., Mitraka, E., Munro, J., Tauber, B., Schor, M., Nickle, L., Felix, V., Jeng, L.,
Bearer, C., Lichenstein, R., et al. (2019). Human Disease Ontology 2018 update:
classification, content and workflow expansion. Nucleic Acids Res. *47*, D955–D962.

23. Luecken, M.D., Büttner, M., Chaichoompu, K., Danese, A., Interlandi, M., Mueller, M.F.,
Strobl, D.C., Zappia, L., Dugas, M., Colomé-Tatché, M., et al. (2022). Benchmarking atlas-
level data integration in single-cell genomics. Nat. Methods *19*, 41–50.

24. Malkov, Y.A., and Yashunin, D.A. (2020). Efficient and Robust Approximate Nearest
Neighbor Search Using Hierarchical Navigable Small World Graphs. IEEE Trans. Pattern
Anal. Mach. Intell. *42*, 824–836.

25. Young, M.D., Mitchell, T.J., Custers, L., Margaritis, T., Morales-Rodriguez, F., Kwakwa, K.,
Khabirova, E., Kildisiute, G., Oliver, T.R.W., de Krijger, R.R., et al. (2021). Single cell
derived mRNA signals across human kidney tumors. Nat. Commun. *12*, 3896.

26. Cano-Gamez, E., Soskic, B., Roumeliotis, T.I., So, E., Smyth, D.J., Baldrighi, M., Willé, D.,
Nakic, N., Esparza-Gordillo, J., Larminie, C.G.C., et al. (2020). Single-cell transcriptomics
identifies an effectorness gradient shaping the response of CD4+ T cells to cytokines. Nat.
Commun. *11*, 1801.

27. Beers, M.F., and Moodley, Y. (2017). When Is an Alveolar Type 2 Cell an Alveolar Type 2
Cell? A Conundrum for Lung Stem Cell Biology and Regenerative Medicine. Am. J. Respir.
Cell Mol. Biol. *57*, 18–27.

28. Wynn, T.A., and Vannella, K.M. (2016). Macrophages in Tissue Repair, Regeneration, and
Fibrosis. Immunity *44*, 450–462.

29. Lis-López, L., Bauset, C., Seco-Cervera, M., and Cosín-Roger, J. (2021). Is the Macrophage Phenotype Determinant for Fibrosis Development? Biomedicines *9*. 10.3390/biomedicines9121747.

30. Adams, T.S., Schupp, J.C., Poli, S., Ayaub, E.A., Neumark, N., Ahangari, F., Chu, S.G., Raby, B.A., DeIuliis, G., Januszyk, M., et al. (2020). Single-cell RNA-seq reveals ectopic and aberrant lung-resident cell populations in idiopathic pulmonary fibrosis. Science Advances *6*, eaba1983.

31. Ayaub, E.A., Poli, S., Ng, J., Adams, T., Schupp, J., Quesada-Arias, L., Poli, F., Cosme, C., Robertson, M., Martinez-Manzano, J., et al. (2021). Single Cell RNA-seq and Mass Cytometry Reveals a Novel and a Targetable Population of Macrophages in Idiopathic Pulmonary Fibrosis. 10.1101/2021.01.04.425268.

32. Jaitin, D.A., Adlung, L., Thaiss, C.A., Weiner, A., Li, B., Descamps, H., Lundgren, P., Bleriot, C., Liu, Z., Deczkowska, A., et al. (2019). Lipid-Associated Macrophages Control Metabolic Homeostasis in a Trem2-Dependent Manner. Cell *178*, 686-698.e14.

33. Morse, C., Tabib, T., Sembrat, J., Buschur, K.L., Bittar, H.T., Valenzi, E., Jiang, Y., Kass, D.J., Gibson, K., Chen, W., et al. (2019). Proliferating SPP1/MERTK-expressing macrophages in idiopathic pulmonary fibrosis. Eur. Respir. J. *54*. 10.1183/13993003.02441-2018.

34. Mulder, K., Patel, A.A., Kong, W.T., Piot, C., Halitzki, E., Dunsmore, G., Khalilnezhad, S., Irac, S.E., Dubuisson, A., Chevrier, M., et al. (2021). Cross-tissue single-cell landscape of human monocytes and macrophages in health and disease. Immunity *54*, 1883-1900.e5.

35. Ramachandran, P., Dobie, R., Wilson-Kanamori, J.R., Dora, E.F., Henderson, B.E.P., Luu, N.T., Portman, J.R., Matchett, K.P., Brice, M., Marwick, J.A., et al. (2019). Resolving the fibrotic niche of human liver cirrhosis at single-cell level. Nature *575*, 512–518.

36. Reyfman, P.A., Walter, J.M., Joshi, N., Anekalla, K.R., McQuattie-Pimentel, A.C., Chiu, S., Fernandez, R., Akbarpour, M., Chen, C.-I., Ren, Z., et al. (2019). Single-Cell Transcriptomic Analysis of Human Lung Provides Insights into the Pathobiology of Pulmonary Fibrosis. Am. J. Respir. Crit. Care Med. *199*, 1517–1536.

37. Wendisch, D., Dietrich, O., Mari, T., von Stillfried, S., Ibarra, I.L., Mittermaier, M., Mache, C., Chua, R.L., Knoll, R., Timm, S., et al. (2021). SARS-CoV-2 infection triggers profibrotic macrophage responses and lung fibrosis. Cell *184*, 6243-6261.e27.

38. Gao, X., Jia, G., Guttman, A., DePianto, D.J., Morshead, K.B., Sun, K.-H., Ramamoorthi, N., Vander Heiden, J.A., Modrusan, Z., Wolters, P.J., et al. (2020). Osteopontin Links Myeloid Activation and Disease Progression in Systemic Sclerosis. Cell Reports Medicine *1*, 100140.

39. Mirizio, E., Tabib, T., Wang, X., Chen, W., Liu, C., Lafyatis, R., Jacobe, H., and Torok, K.S. (2020). Single-cell transcriptome conservation in a comparative analysis of fresh and cryopreserved human skin tissue: pilot in localized scleroderma. Arthritis Res. Ther. *22*, 263.

40. Lin, W., Noel, P., Borazanci, E.H., Lee, J., Amini, A., Han, I.W., Heo, J.S., Jameson, G.S.,
Fraser, C., Steinbach, M., et al. (2020). Single-cell transcriptome analysis of tumor and
stromal compartments of pancreatic ductal adenocarcinoma primary tumors and metastatic
lesions. Genome Med. *12*, 80.

41. Kemp, S.B., Steele, N.G., Carpenter, E.S., Donahue, K.L., Bushnell, G.G., Morris, A.H., The,
S., Orbach, S.M., Sirihorachai, V.R., Nwosu, Z.C., et al. (2021). Pancreatic cancer is marked
by complement-high blood monocytes and tumor-associated macrophages. Life Science
Alliance *4*, e202000935.

42. Bhattacharya, M. (2022). Insights from Transcriptomics: CD163+ Profibrotic Lung
Macrophages in COVID-19. Am. J. Respir. Cell Mol. Biol. *67*, 520–527.

43. Martinez, F.O., Gordon, S., Locati, M., and Mantovani, A. (2006). Transcriptional Profiling
of the Human Monocyte-to-Macrophage Differentiation and Polarization: New Molecules
and Patterns of Gene Expression1. The Journal of Immunology *177*, 7303–7311.

44. Salahudeen, A.A., Choi, S.S., Rustagi, A., Zhu, J., van Unen, V., de la O, S.M., Flynn, R.A.,
Margalef-Català, M., Santos, A.J.M., Ju, J., et al. (2020). Progenitor identification and SARS-
CoV-2 infection in human distal lung organoids. Nature *588*, 670–675.

45. Duy, C., Li, M., Teater, M., Meydan, C., Garrett-Bakelman, F.E., Lee, T.C., Chin, C.R.,
Durmaz, C., Kawabata, K.C., Dhimolea, E., et al. (2021). Chemotherapy Induces Senescence-
Like Resilient Cells Capable of Initiating AML Recurrence. Cancer Discov. *11*, 1542–1561.

46. Karagiannis, T.T., Cleary, J.P., Gok, B., Henderson, A.J., Martin, N.G., Yajima, M., Nelson,
E.C., and Cheng, C.S. (2020). Single cell transcriptomics reveals opioid usage evokes
widespread suppression of antiviral gene program. Nat. Commun. *11*, 2611.

47. Xu, Y., Zeng, X., Zhang, M., Wang, B., Guo, X., Shan, W., Cai, S., Luo, Q., Li, H., Li, X., et
al. (2022). Efficient expansion of rare human circulating hematopoietic stem/progenitor cells
in steady-state blood using a polypeptide-forming 3D culture. Protein Cell. 10.1007/s13238-
021-00900-4.

48. Regev, A., Teichmann, S.A., Lander, E.S., Amit, I., Benoist, C., Birney, E., Bodenmiller, B.,
Campbell, P., Carninci, P., Clatworthy, M., et al. (2017). The Human Cell Atlas. Elife *6*,
e27041.

49. Liou, G.-Y. (2017). Inflammatory Cytokine Signaling during Development of Pancreatic and
Prostate Cancers. J Immunol Res *2017*, 7979637.

Methods

SCimilarity model architecture and loss function

Model architecture

The SCimilarity model consists of one fully connected encoder and one decoder stage and reuses the same encoding network three times per training triplet, such that updates to the model after each batch are shared equally for each subsequent batch of training triplets. The decoder stage is not part of the conventional triplet loss architecture, but is included to compute a mean squared error (MSE) reconstruction loss.

Expression profiles are reduced through an encoder network, starting from 28,231 genes through three hidden layers with dimensions 1,024, 1,024, and 128. The 128-dimensional outputs are unit length normalized, forcing all low dimensional cell representations to lie on the surface of a hypersphere. During training, the input layer is subjected to 40% dropout, zeroing out many gene expression values at random, and each hidden layer is subjected to 50% dropout rates for maximum regularization [1].

While hyperspheric spaces have been infrequently used for representation of single-cell profiles [2], the triplet loss model often uses hypersphere embeddings to ensure consistency between the model hyperparameters [3]. During triplet loss training, the objective is to place cells of different types sufficiently far apart. The minimum desired distance between cells of different types is called the margin. By fixing the volume of the embedding space to the surface of a unit length

64-dimensional hypersphere, the margin is interpreted consistently between model runs. Without normalization, cells can be placed up to an infinite distance apart, rendering the margin meaningless.

Triplet loss training

To learn features that place data points considered similar near each other, the loss function depends on distances between data points embedded in a learned low dimensional latent space, described with:

$$d(x, y) = ||f(x) - f(y)||_2^2$$

where x and y are two high dimensional vectors (here, cell profiles), passed through a neural network encoder $f()$.

The triplet loss model learns from three vectors at a time: the anchor (x_i^a), positive (x_i^p) and negative (x_i^n). The anchor and positive vectors are considered similar, whereas the anchor and negative are dissimilar.

The model parameters are iteratively updated to decrease the number of triplets where the distance between the anchor and negative data vectors is insufficiently large relative to the distance between the anchor and the positive points, thus minimizing the triplet loss function:

$$L_{triplet} = \frac{\sum_i^N max\left(d\left(x_i^a, x_i^p\right) - d\left(x_i^a, x_i^n\right) + \alpha, 0\right)}{N}$$

where α is the margin, which denotes how much further the negatives should be from the anchor than the positives, and i is the index of the triplet.

Reconstruction loss training

The reconstruction loss is computed on the anchor cell only, because each anchor cell is used only once as an anchor within a batch. The reconstruction loss is defined as:

$$L_{MSE} = \frac{\sum_{i}^{N} \left\| x_i^a - g(f(x_i^a)) \right\|_2^2}{N}$$

where N is the number of anchor cells in a batch, set to $N=1000$ in SCimilarity, and $g()$ is the function learned by the neural network decoder stage.

Combined loss function

Adding a reconstruction loss to classification models has been shown to improve generalization [4] through a regularization effect. The SCimilarity loss function combines the triplet loss and reconstruction loss functions as follows:

$$L = (1 - \beta) * L_{MSE} + \beta * L_{triplet}$$

where β is a weighting term in $[0, 1]$. $\beta = 0$ corresponds to a conventional autoencoder, and $\beta = 1$ corresponds to a pure triplet loss model. Empirically, $\beta = 0.001$ performed best on the cell search task (query model) and $\beta = 1$ performed best on batch integration (integration model) (**Extended Data Fig. 2a**).

Use of Cell Ontology terms and relationships

Authors may annotate cell types at different granularities, which confounds triplet sampling by introducing cell type annotations with hierarchical relationships that cannot be unambiguously defined as either similar or dissimilar. As such, cell type annotations used for training are defined using standardized Cell Ontology terms and valid triplets are restricted to cells without vertical

Cell Ontology relationships between members of the triplet. A vertical relationship is defined as any directed path of one or more ancestor-descendant relationships in the Cell Ontology network. Thus, there are three binary relations defined for annotation: (1) similar pairs with identical annotations (*e.g.*, "T cell" and "T cell"), (2) dissimilar pairs with non-vertical ontology relationships (*e.g.*, "CD4-positive, alpha-beta T cell" and "CD8-positive, alpha-beta T cell"), and (3) ambiguous pairs with vertical relationships (*e.g.*, "T cell" and "CD4-positive, alpha-beta T cell"). Positives are drawn from cells similar to the anchor, negatives are drawn from cells dissimilar to the anchor, and cells that are ambiguous to the anchor are excluded from sampling.

GEO data aggregation

334 human scRNA-seq datasets were obtained from the Gene Expression Omnibus (GEO)[5]. Multiple filtering steps were used to restrict the datasets analyzed to samples from human tissue, that were generated using the 10x Chromium platform, and which reported unnormalized gene count data that could be automatically processed. To select appropriate datasets, search criteria were designed for the Biopython Entrez search tool (Cock et al., 2019) to find GEO studies that had specific properties, such as metadata keywords, file formats, and species. Then, using GEOparse[6], the GEO text metadata was downloaded for each sample and searched for blacklisted words in the metadata or download URLs (*e.g.*, "smartseq", "trizol", and "fasta") to further filter out samples that were not generated using 10x Chromium. Data for samples and corresponding download links that passed the metadata filter stage were automatically downloaded. No datasets were realigned. 753 studies were identified for download. A set of import functions was designed for the most common file type formats (.mtx, .h5ad, and gene

expression matrices in .tsv or .csv). Any dataset that could not be successfully downloaded or read in was discarded. Once read in, each sample was automatically tested for count data and gene names that match a reference gene list or gene name mapper before saving each file in a uniform h5ad format for later processing. This resulted in a total of 334 published studies that were not duplicates of studies found in CELLxGENE [7] for use in our analysis.

Data preprocessing

All UMI count data were natural log normalized per-cell with a scaling factor of 10,000 using the scanpy.pp.normalize_to_target(adata, 10000) and scanpy.pp.log1p(adata) functions from scanpy[8].

Manual data aggregation, normalization and filtering

Datasets with author-provided cell type annotations used for training were obtained from Tabula Sapiens[9], 10x Genomics[10], the single nucleus cross-tissue atlas[11], and the human lung cell atlas[12] and subjected to the same preprocessing procedures as programmatically-downloaded datasets. Cell type annotations were manually converted into terms contained within the Cell Ontology. Cells that with annotations that did not clearly map to the Cell Ontology were not included in training.

Cell profiles previously annotated as doublets, scored as doublets by infer_doublets from Pegasus[13], had >20% total UMI counts aligned to mitochondrial genes, or had <500 total genes detected were removed.

Preparation of training and test data

Training and test sets were chosen such that entire studies were held out of training (rather than holding out a subset of cells from each dataset) (**Extended Data Table 1**); there were 52 and 14 datasets in the training and test sets, respectively. This presents a harder generalization challenge and reflects how users are likely to use SCimilarity. Test datasets were selected to reflect the tissue diversity within the training sets.

Selection of Cell Ontology terms for training

Cell Ontology terms were selected for training if they were observed in at least two separate studies in the training set. Terms that appeared in only one study were not used because SCimilarity is trained by comparing cells across studies. To rescue single-study terms, the immediate parent terms were inspected across studies. If a single-study term's parent was observed in at least two other datasets then the original cell type annotation was replaced with the coarser parent term (**Extended Data Table 1**). Otherwise, all cells with this annotation were removed from training. As the size or annotation quality of training data grows, the number of Cell Ontology terms meeting the inclusion criteria are expected to increase.

Triplet sampling and semi-hard triplet mining

During training, batches of 1,024 cells are sampled from the training datasets. This sampling is weighted by study and cell type to have a similar number of observations per cell type from each study per batch.

Because of the *maximum* operation within the loss function, not all viable triplets contribute to the gradient, and are categorized as easy, semi-hard or hard, based on their contribution to the gradient.

Easy negatives are defined as:

$$\left\| f(x_i^a) - f(x_i^p) \right\|_2^2 < \left\| f(x_i^a) - f(x_i^n)) \right\|_2^2 + \alpha$$

Easy negatives provide no information to the gradient because the distances between the cells in the low dimensional embedding already satisfy the objective, such that the *maximum* operation returns 0 to the triplet loss sum. Because there are many easy triplets after training a small number of batches, randomly sampling triplets does not train models effectively. To accelerate training, triplets are mined to search for training triplets that are especially informative for model training[3].

Hard negatives are defined as:

$$\left\| f(x_i^a) - f(x_i^p) \right\|_2^2 > \left\| f(x_i^a) - f(x_i^n)) \right\|_2^2 + \alpha$$

Hard negatives contribute the largest quantity to the loss function, because they do not fit and are far from fitting the desired latent relationships. In practice, hard triplets are rarely useful for training, because they contribute to model collapse during training[3,14]. Hard negatives may be enriched for incorrectly annotated cells.

Semi-hard negatives are defined as:

$$\left\| f(x_i^a) - f(x_i^n)) \right\|_2^2 - \left\| f(x_i^a) - f(x_i^p) \right\|_2^2 < \alpha$$

Semi-hard negatives contribute small amounts to the loss function because they nearly satisfy the desired distances between cells in low dimensional space. Meaning, the negative cell profile is further from the anchor cell than the positive cell, but by a less than desired distance α. Semihard negatives are often used in triplet loss models[3].

Overall, we chose to train SCimilarity using only semi-hard negative triplets.

Explainability framework and marker gene identification

An explainability framework was used to identify genes whose variation leads to the most significant variations of the learned features and, in turn, affects the relative distance between different cells.

An explanation for a pair of cells is defined as those genes which have the greatest impact on the relative distance between those cells in latent space. Given $d(x, y) = ||f(x) - f(y)||_2^2$, the distance between two cell profiles x and y in latent space f, the integrated gradient approach (Sundararajan et al. 2017) was extended to compute the importance of each gene i in the comparison between cell profiles x and y as:

$$Importance_i(x) = \left| \max((x_i - y_i), 0) \times \int_{\alpha=0}^{1} \frac{\partial d(y + \alpha \times (x - y), y)}{\partial x_i} \right|$$

High values of $Importance_i(x)$ correspond to genes that are highly expressed in x, and their modification (i.e., gradient) affects $d(x, y)$ more. Intuitively, the expression of each gene in y is gradually increased to match x along the trajectory from x to y. Through this trajectory, the rate

of change of $d(x, y)$ is computed for each gene, aggregating the results. The score is scaled by $(x_i - y_i)$. In order to identify genes that are up regulated in a subset of interest, genes i with expression $x_i < y_i$ are ignored.

This approach differs in several key ways from the standard integrated gradient approach, because: (1) gradients are computed with respect to a learned distance instead of output features, (2) attributions where $x_i < y_i$ are ignored and (3) the sign of the integral is ignored due to the complex interactions between features.

To identify important genes for a cell type t, a set of cells $T \in \{t_1, ..., t_N\}$ with cell type t and a set of cells $B \in \{b_1, ..., b_N\}$ with cell types different from t are randomly sampled. Pairwise importances are computed for each pair of cells t_i in T and b_j in B and aggregated to obtain a signature that characterizes cell type t as:

$$Signature_i(t) = \frac{1}{N} \sum_{c=1}^{N} Importance_i(t_c, b_c)$$

Since the pairwise comparisons are averaging relative comparisons, the sampling of $\{b_1, ..., b_N\}$ impacts the signature scoring. To obtain general cell type markers, a background of all cell types is sampled. To obtain a cell state specific signature, a background of cells in other states of the same type are sampled. Confidence intervals for each gene i are computed as the standard error of the mean. This results in an attribution score for each gene.

Training and evaluation metrics

SCimilarity score

The SCimilarity score is defined as the inverse of the cosine distance of two embedded cell profiles:

$$SCimilarity\ score\ =\frac{1}{1-c_i \bullet c_j}$$

where c_i and c_j are the embeddings of the ith and jth cell profiles with unit length, respectively and $i \neq j$. The threshold for similarity varies in practice by question and cell types.

Ontology-aware modified average silhouette width

Average silhouette width (ASW) has been used to assess the performance of data integration tasks on multiple scRNA-seq studies[15] by quantifying how coherently grouped each cell type is across studies. The silhouette width of cell profile i of cell type t typically compares the average intra-cell type distances $a(i)$ and the average inter-cell type distances $b(i)$ between cells of type t and cells of the nearest cell type, defined as:

$$a(i) = \frac{1}{|C_I|-1} \sum_{j \in C_I, i \neq j} d(i,j)$$

$$b(i) = \min_{J \neq I} \frac{1}{|C_J|-1} \sum_{j \in C_J} d(i,j)$$

where, typically, C_I is the set of cells of author-annotated type t and C_J are the cells of all other cell types.

However, the ASW as typically formulated does not account for differences in granularity of cell type annotations across studies. To address those, a modified formulation is used where C_I contains cell type label t and all of its ontological descendants and C_J is the set of all other cell types, except cells of type t and any of its ontological descendants or ancestors. For example, if computing $a(i)$ for a T cell, distances between all types of T cell terms ("CD4-positive", "alpha-beta T cell", "CD8-positive", "alpha-beta T cell and CD4-positive", "CD25-positive", "alpha-beta regulatory T cell", etc) are members of the "T cell" term. Ancestor terms of T cells, such as the term "Lymphocytes", are not members of the T cell class (nor a T cell subset) but are excluded from the summation indices in the calculations of $a(i)$ and $b(i)$.

Correlation to predefined gene signatures

To test how the SCimilarity distance represents distance between predefined cell states, a signature-based definition of cell state was correlated with the SCimilarity score (above).

For each cell in the test set, both the signature score[16] and a SCimilarity score *vs*. the cell query are calculated, yielding two vectors, and Pearson's correlation coefficient is calculated between the vectors.

Selection of models for downstream analysis

Models were run in triplicate along 6 different β parameters ranging from [0,1] and one query model and one integration model were selected based on two criteria. First, query performance was tested by how well cell similarities to a query FMΦ profile correlated with a signature defining that same state (*TREM2, GPNMB, SPP1, CCL18, MMP9, CTSK, APOE, CHIT1, LIPA,*

CHI3L1, *CD14*, *APOC1*). Second, ontology aware ASW was used to quantify how well the cells of the same type from different studies intermixed in SCimilarity's representation. The query model was selected as the model with the highest query test performance. The integration model was selected from the β=1 models. Since the three replicates had nearly identical integration scores, we picked the model with the highest query test score as it performed much better on the query task than the other high integration models. (**Extended Data Fig. 2a**). The selected integration model had more study mixing than the query model according to the study (NMI) and study adjusted rand index (ARI)[15].

Benchmarking vs. integration methods

SCimilarity's integration and cell search models were each compared to three batch integration methods: Harmony[17], scVI[18], and scArches[19]. A test dataset of 34,713 cells was created by sampling cells from lung tissue studies with uniform probability across studies. The modified ASW (above), adjusted Rand index (ARI) and normalized mutual information (NMI) were calculated as integration benchmark metrics. Harmony was run using the wrapper in Pegasus[13] following the workflow described in https://pegasus-tutorials.readthedocs.io/en/latest/_static/tutorials/batch_correction.html. scVI and scArches were run using the scvi-tools workflow described in https://docs.scvi-tools.org/en/stable/tutorials/notebooks/harmonization.html and https://docs.scvi-tools.org/en/stable/tutorials/notebooks/scarches_scvi_tools.html, respectively. As the scArches workflow requires a reference dataset, 101,133 cell profiles were sampled across all training datasets with uniform probability across studies for use as the reference.

Cell type annotation

Cell type assignments were performed by k-nearest neighbors (k-NN) classification combined with an annotated reference set. SCimilarity's reduced dimensionality latent space was used to determine $k=50$ nearest neighbors in the reference data set to a query cell t, and the query cell was annotated either by tallying votes based each cell's annotation with either equal weights,

$$\text{Celltype } (t) = \arg \max_{t} \left(\sum_{i \in t} \frac{1}{n} \right)$$

or with weights by distance in SCimilarity's reduced dimensionality latent space:

$$\text{Celltype } (t) = \arg \max_{t} \left(\sum_{y \in t} \frac{1}{d(x,y)} \right)$$

To allow users to annotate new datasets from a restricted list of cell types of interest, excluding (blocklisting) or limiting to (safelisting) specified cell type annotations is used, and is recommended when feasible to improve interpretability and reduce spurious annotations. However, extensive blocklisting or safelisting can slow the annotation process significantly, because the pre-built k-NN indices are not optimized for a modified target cell type list.

kNN parameters for annotation and querying

Two separate kNN indices were used for efficient and accurate queries. For cell type annotation, a 7.9M cell k-NN index was built using hnswlib[20] with ef_construction = 1000 and M = 80. Searching this k-NN found the 50 nearest neighbors (default behavior) for cell type annotation (k=50) and ef=100.

Cell query relied on a separate 22.7M cell k-NN index also built using hnswlib. This index was constructed with the following parameters: ef_construction=400 and M=50. The search parameters are set by the user's request for how many similar cells to return. Default behavior is set to k=1000 and ef=k, but in practice k can vary widely depending on the use case.

Outlier filtering

To filter outlier cells prior to visualization and downstream analysis, SCimilarity's score is used to flag cells that are out of distribution. Cells with a SCimilarity score < 50 from the nearest cell in the training set were removed prior to further analysis. Many of these cells were from immortalized cell lines, and reflect their difference from primary cells (and absence in the training). Note that if out of distribution cells are not removed, these cells won't be accurately annotated and can confound visualization.

Macrophage query preprocessing

To prepare a cell query for FMΦ cells, a public dataset[21] (GSE136831 and https://www.ipfcellatlas.com) was preprocessed with the same steps for all ingested data and scored use Scanpy's scanpy.tl.score_genes function with a gene signature of *SPP1*, *TREM2*, *GPNMB*, *MMP9, CHIT1*, and *CHI3L1* Scanpy[8]. The average profile of the top 50 scoring cell was embedded using SCimilarity and used as the input query to SCimilarity's cell search model and used throughout analyses in **Fig. 5** and **6**.

Important genes and pathway enrichment

Important genes were identified using SCimilarity's attribution score method. This method requires two cell groups to compare, identifying which genes differ between them. Here we used 1,000 cells that were considered similar to the average FMΦ profile calculated from Adams et al. as the FMΦ-like group. This query excluded any cells from the Adams et al. dataset. To compare to the FMΦ-like group comparison, 1,000 dissimilar monocytes and macrophages were randomly sampled (any monocyte or macrophage that was not within the top 10,000 most FMΦ similar results).

Reactome pathways enriched for the 100 genes with the top importance scores for FMΦ were determined using the method provided in the ReactomePA[22] R package, with multiple hypothesis correction using the Benjamini-Hochberg method and the background gene universe restricted to the ~28,000 genes included in SCimilarity. Pathways were considered significant if they met the criteria of adjusted p-value (Q) ≤ 0.05 and gene count ≥ 5.

3DCS culture of PBMC

Peripheral blood was sourced from healthy volunteers at Genentech that were consented as per IRB. Samples were collected in heparin collection tubes and subsequently diluted 1:1 with a solution of PBS containing 2% FBS and 1mM EDTA. 30 ml of diluted blood was overlayed onto 15 ml of Lymphoprep (STEMCELL Technologies) in a 50ml tube and centrifuged at 3,000 rpm for 20 minutes at 4°C. PBMCs were isolated from the interphase after centrifugation and diluted

with PBS containing 2% FBS and 1 mM EDTA and centrifuged at 300 x g for 10 minutes at 4°C. Cell pellet was washed again with PBS containing 2% FBS and 1mM EDTA. Red blood cell lysis was performed on the cell pellet by resuspending in RBC Lysis Buffer (Cell Signaling Technology) for 5 minutes at room temperature, followed by inactivation with addition of RPMI media containing 10% FBS. Cells were pelleted by centrifugation at 300 x g for 10 minutes at 4°C and subsequently washed with PBS containing 2% FBS and 1 mM EDTA. Cells were then resuspended in a 10% sucrose solution at a concentration of 2 x 106 cells/ml right before plating into 3D hydrogel culture. Puramatrix hydrogel (Corning) was vortexed for 30 seconds and diluted 1:1 with a 20% sucrose solution. 250 µl of diluted Puramatrix hydrogel was mixed with 250 µl of resuspended PBMCs and plated in a 24-well tissue culture plate. To induce gelation, RPMI media was overlaid onto the hydrogel/PBMC mixture and incubated for 5 minutes in a 37°C incubator with 5% CO_2. Overlayed media was aspirated off of the 3D hydrogel and washed twice with RPMI media, after which 600 µl of 3DCS media, formulated as previously described (Xu, Y. et al., Protein & Cell 2022, 13:808-824) was overlaid onto the hydrogel. Cells were cultured in a 37°C incubator with 5% CO2 for 8 days, with media exchanges every other day. On day 8, culture cells were recovered from the 3D hydrogel for scRNA-seq.

Single cell RNA-Seq from 3DCS cultures

Wells of 3D hydrogel culture were washed with PBS, followed by recovery of the hydrogel and cells by gentle pipetting in PBS buffer. This solution was centrifuged for 5 minutes at 750 x g and the hydrogel/PBMC pellet was resuspended in TrypLE solution (ThermoFisher Scientific) and incubated at 37°C for 10 minutes. RPMI media with 10% FBS was added and the solution was centrifuged for 5 minutes at 750 x g. The resultant pellet was washed twice with PBS to

remove hydrogel matrix debris. PBMCs were resuspended in PBS and passed through a 40 µM filter, pelleted by centrifugation at 300 x g for 5 minutes, and resuspended in RPMI media with 10% FBS. The cell solution was subjected to FACS to isolate cells from any remaining hydrogel debris and recovered cells were concentrated to 1,000 cells/µl in RPMI media with 10% FBS for downstream profiling by scRNA-seq.

ScRNA-seq was performed using the Chromium Single Cell 3' Library and Gel bead kit v3 (10x Genomics), following manufacturer's user guide. Briefly, cell density and viability of single-cell suspension were determined by Vi-CELL XR cell counter (Beckman Coulter). Cell density was used to impute the volume of single cell suspension needed in the reverse transcription (RT) master mix, aiming to achieve ~10,000 cells per sample. cDNAs and libraries were prepared following the manufacturer's user guide (10x Genomics). Libraries were profiled by Bioanalyzer High Sensitivity DNA kit (Agilent Technologies) and quantified using Kapa Library Quantification Kit (Kapa Biosystems). Libraries were sequenced on a NovaSeq 6000 (Illumina) following the manufacturer's specifications with 28+90 bp paired-end reads at a depth of 101M mate-pair reads. Sequencing reads were aligned to the GENCODE 27 Basic gene model on the human genome assembly GRCh38 using Cell Ranger v6.0 (10x Genomics, Pleasanton, CA, USA).

Individual samples were genetically demultiplexing using the singularity container provided with Souporcell 2.0 [23]. No genotype information was provided to the pipeline. Since PBMCs were provided from 3 donors, a k of 3 was used to cluster the samples into 3 genotypes. These samples

were pre-processed consistently with the previously ingested samples and then embedded using SCimilarity to enable direct comparisons to Xu et al as well as the rest of the public datasets.

SCimilarity cell type classification was applied to both public and validation cells using SCimilarity with the following safelist: B cell, CD4-positive, alpha-beta T cell, CD8-positive, alpha-beta T cell, conventional dendritic cell, hematopoietic stem cell, macrophage, monocyte, natural killer cell, plasma cell, plasmacytoid dendritic cell.

Code performance benchmarking

Benchmarks were run on servers with 8 Intel Xeon E5-2650 v4 CPUs with 2.20GHz cores and a total of 128 GB of RAM.

Query runtimes, using the pre-built approximate k-NN index[20] to find the top n most similar cells, had an average runtime of 50 milliseconds. Some API functions use the query and summarize the metadata within one function call. That function timing is dominated by summarizing metadata and computing statistics from the query results, which requires an additional 3.3 seconds. This performance differs from an exhaustive comparison (**Fig. 5b**), where the query was directly compared against 2.58M monocytes and macrophages with a runtime of 2 seconds.

Cell signatures were calculated using scanpy.tl.score_genes. The scanpy score_genes function was applied to the already normalized data. This runtime totalled 2 hours, 46 minutes and 20

seconds when it was applied across each h5ad file (one file per tissue sample). Even though h5ad files were not stored with any compression, file reading was a dominant factor in runtime.

Code availability

Code and tutorials are available at https://github.com/Genentech/scimilarity.

Licensing

- Code license: Apache 2.0

- Pretrained model weights, kNN and pre-built indices license: CC-BY-SA 4.0

References

1. Baldi, P., and Sadowski, P. (2013). Understanding dropout. In Proceedings of the 26th International Conference on Neural Information Processing Systems - Volume 2 NIPS'13. (Curran Associates Inc.), pp. 2814–2822.

2. Ding, J., and Regev, A. (2021). Deep generative model embedding of single-cell RNA-Seq profiles on hyperspheres and hyperbolic spaces. Nat. Commun. 12, 2554.

3. Schroff, F., Kalenichenko, D., and Philbin, J. (2015). FaceNet: A Unified Embedding for Face Recognition and Clustering. 2015 IEEE Conference on Computer Vision and Pattern Recognition (CVPR), 815–823.

4. Le, L., Patterson, A., and White, M. (2018). Supervised autoencoders: improving generalization performance with unsupervised regularizers. In Proceedings of the 32nd International Conference on Neural Information Processing Systems NIPS'18. (Curran Associates Inc.), pp. 107–117.

5. Edgar, R., Domrachev, M., and Lash, A.E. (2002). Gene Expression Omnibus: NCBI gene expression and hybridization array data repository. Nucleic Acids Res. 30, 207–210.

6. Gumienny, R. GEOparse: Python library to access Gene Expression Omnibus Database (GEO).

7. Chan Zuckerberg CELLxGENE Discover (2022). Cellxgene Data Portal.

8. Wolf, F.A., Angerer, P., and Theis, F.J. (2018). SCANPY: large-scale single-cell gene expression data analysis. Genome Biol. 19, 15.

9. Tabula Sapiens Consortium*, Jones, R.C., Karkanias, J., Krasnow, M.A., Pisco, A.O., Quake, S.R., Salzman, J., Yosef, N., Bulthaup, B., Brown, P., et al. (2022). The Tabula Sapiens: A multiple-organ, single-cell transcriptomic atlas of humans. Science 376, eabl4896.

10. Zheng, G.X.Y., Terry, J.M., Belgrader, P., Ryvkin, P., Bent, Z.W., Wilson, R., Ziraldo, S.B., Wheeler, T.D., McDermott, G.P., Zhu, J., et al. (2017). Massively parallel digital transcriptional profiling of single cells. Nat. Commun. 8, 14049.

11. Eraslan, G., Drokhlyansky, E., Anand, S., Fiskin, E., Subramanian, A., Slyper, M., Wang, J., Van Wittenberghe, N., Rouhana, J.M., Waldman, J., et al. (2022). Single-nucleus cross-tissue molecular reference maps toward understanding disease gene function. Science 376, eabl4290.

12. Travaglini, K.J., Nabhan, A.N., Penland, L., Sinha, R., Gillich, A., Sit, R.V., Chang, S., Conley, S.D., Mori, Y., Seita, J., et al. (2020). A molecular cell atlas of the human lung from single-cell RNA sequencing. Nature 587, 619–625.

13. Li, B., Gould, J., Yang, Y., Sarkizova, S., Tabaka, M., Ashenberg, O., Rosen, Y., Slyper, M., Kowalczyk, M.S., Villani, A.-C., et al. (2020). Cumulus provides cloud-based data analysis for large-scale single-cell and single-nucleus RNA-seq. Nat. Methods 17, 793–798.

14. Wu, C.-Y., Manmatha, R., Smola, A.J., and Krähenbühl, P. (2017). Sampling Matters in Deep Embedding Learning. arXiv [cs.CV].

15. Luecken, M.D., Büttner, M., Chaichoompu, K., Danese, A., Interlandi, M., Mueller, M.F., Strobl, D.C., Zappia, L., Dugas, M., Colomé-Tatché, M., et al. (2022). Benchmarking atlas-level data integration in single-cell genomics. Nat. Methods 19, 41–50.

16. Satija, R., Farrell, J.A., Gennert, D., Schier, A.F., and Regev, A. (2015). Spatial reconstruction of single-cell gene expression data. Nat. Biotechnol. 33, 495–502.

17. Korsunsky, I., Millard, N., Fan, J., Slowikowski, K., Zhang, F., Wei, K., Baglaenko, Y., Brenner, M., Loh, P.-R., and Raychaudhuri, S. (2019). Fast, sensitive and accurate integration of single-cell data with Harmony. Nat. Methods 16, 1289–1296.

18. Lopez, R., Regier, J., Cole, M.B., Jordan, M.I., and Yosef, N. (2018). Deep generative modeling for single-cell transcriptomics. Nat. Methods 15, 1053–1058.

19. Lotfollahi, M., Naghipourfar, M., Luecken, M.D., Khajavi, M., Büttner, M., Wagenstetter, M., Avsec, Ž., Gayoso, A., Yosef, N., Interlandi, M., et al. (2022). Mapping single-cell data to reference atlases by transfer learning. Nat. Biotechnol. 40, 121–130.

20. Malkov, Y.A., and Yashunin, D.A. (2020). Efficient and Robust Approximate Nearest Neighbor Search Using Hierarchical Navigable Small World Graphs. IEEE Trans. Pattern Anal. Mach. Intell. 42, 824–836.

21. Adams, T.S., Schupp, J.C., Poli, S., Ayaub, E.A., Neumark, N., Ahangari, F., Chu, S.G., Raby, B.A., DeIuliis, G., Januszyk, M., et al. (2020). Single-cell RNA-seq reveals ectopic and aberrant lung-resident cell populations in idiopathic pulmonary fibrosis. Science Advances 6, eaba1983.

22. Yu, G., and He, Q.-Y. (2016). ReactomePA: an R/Bioconductor package for reactome pathway analysis and visualization. Mol. Biosyst. 12, 477–479.

23. Heaton, H., Talman, A.M., Knights, A., Imaz, M., Gaffney, D.J., Durbin, R., Hemberg, M., and Lawniczak, M.K.N. (2020). Souporcell: robust clustering of single-cell RNA-seq data by genotype without reference genotypes. Nat. Methods 17, 615–620.

a

Query cell

Searchable datatsets
– 22.7M cells
– 399 studies
– pan tissue & disease
– perturbations

SCimilarity search

SCimilarity score

Tissue | Study

1. Intestine — 337 similar cells: Kim et al. 2022
Sample: Small intestine bioposy, donor 7

2. Intestine — 293 similar cells: Torres et al. 2022
Sample: Crohns bioposy, treated, patient 4

3. *In vitro* PBMCs — **Novel cell model**
277 similar cells: Smith et al. 2021
Sample: Peripheral blood, gene knockouts

7. COVID-19 lung — **Novel disease association**
104 similar cells: Johnson et al. 2023
Sample: COVID-19 lung, donor 02

10,000 similar cells found in 0.05 seconds

low   high

b

Training data

From across the body
Healthy tissue
Immune
Infectious disease

Datasets | Annotations
T cell
CD4+
CD8+
Treg

– 52 training studies
– 14 test labeled studies

Triplet Sampling

Cell ontology
T cell
CD4+ T cell   CD8+ T cell
Treg

Triplet sampling
Ancestor-Descendent   Cousin

Anchor
Positive
Negative

Ambiguous Comparisons ✗   Valid Comparisons ✓

Metric Learning

$$Loss=(1-\beta)\cdot L_{MSE}(\mathbf{A},\hat{\mathbf{A}}) + \beta\cdot L_{triplet}(\mathbf{A},\mathbf{P},\mathbf{N})$$

Input layer   Output layer

$$\textbf{SCimilarity score} = \frac{1}{\text{Embedded distance}}$$

Distance

Fig. 1. SCimilarity metric learning enables cell search in large human scale atlases.

a, Cell Querying with SCimilarity. Left: A query cell profile is compared to a searchable reference

collection of 22.7M profiles from 399 studies. Center: Sample with similar cells are identified and

returned with information about the original sample conditions, including unexpected tissue, *in*

vitro or diseases contexts. Right: A SCimilarity score is computed between the query cell and each

cell within a tissue sample. **b,** Triplet loss training. From left: 52 training and 14 test annotated (by

the Cell Ontology) datasets from across the body are sampled for cell triplets (an anchor, a

"positive" (anchor-similar), and a "negative" (anchor-dissimilar) cell; based on Cell Ontology

annotations) to train a neural network that embeds similar cells closer than dissimilar ones

(**Methods**).

Fig. 2. SCimilarity learns a universal representation that generalizes to new datasets.

a, A large-scale reference database of public gene expression datasets across tissues and diseases. Number of cells (circle size) across tissues (outermost light blue circles) and disease states (middle green circles) across individual studies (innermost circles) in the training (gold), test (pink) or unannotated (purple) datasets. **b,** Integration between studies without feature selection or batch correction. Uniform manifold approximation and projection (UMAP) embedding of cell profiles (dots) generated on a 128-dimensional latent space from SCimilarity's integration model (**Methods**) for cells from 21 tissues (panels) and 239 unique studies (color code). For tissues with more than a million cell profiles, the UMAP embedding was computed on a random uniform subsampling of 1 million cells from the studies for that tissue.

Fig. 3. SCimilarity accurately annotates cell types across the human body

a, SCimilarity cell annotation. A new unannotated cell (red, bottom left) is embedded in

SCimilarity's common low-dimensional space and compared against the precomputed reference

for cell type annotation (0.02 seconds per cell). **b-d**, SCimilarity annotation of a kidney scRNA-

seq dataset. (**b,c**) UMAP embedding of cell profiles (dots) from SCimilarity's latent representation

of a held out kidney dataset[1] colored by author provided (**b**) or SCimilarity predicted (**c**) cell type

annotations. (**d**) Percentage (color bar and number) of author-annotated cells (columns) with each

SCimilarity annotation (rows). **e-h,** SCimilarity-corrected author annotations. UMAP embedding

of cell profiles (dots) from SCimilarity's latent representation for cells either author-annotated or

predicted as CD4$^+$ or CD8$^+$ T cells, colored by author (**e**) or SCimilarity (**f**) annotations, and by

CD8A (**g**) or *CD4* (**h**) expression. **i,** Classification performance. F1 score (y axis) for SCimilarity

vs. author annotation for each cell population (dot) from each of 14 held out datasets (x axis).

Right: Distribution of F1 scores.

Fig. 4. SCimilarity annotations scale to tens of millions of cells from hundreds of datasets

a,b Predicted cell types group by biologically-accurate lineages. **a,** Hierarchical clustering

dendrogram of centroids of predicted cell types (leaves) in SCimilarity latent space, colored by

lineage. Clustering was performed using cosine distance and average linkage. **b,** UMAP of

2,000,000 embedded cells uniformly sampled from the 22.7M reference, colored by clusters (as

labeled in **a**).

Fig. 5. SCimilarity cell search reveals FMΦs across ILD and other diseases

a, SCimilarity cell search. A query cell profile (bottom left) is embedded into the learned

SCimilarity representation along with the reference of 22.7M cells, and its nearest neighbors,

determined by distance from the embedded query in the low dimensional space, are tabulated by

study, tissue and disease. **b-e,** Identification of FMΦs across tissues by SCimilarity cell search. (**b**)

SCimilarity scores (y axis, $\log_{10}$ scale, and color bar) against a FMΦ query profile for each

annotated monocyte or macrophage (dot) from 1,041 *in vivo* tissue samples from 143 studies (x

axis), ordered by mean SCimilarity score. (**c**) Number of cells (circle size) across tissues

(outermost light blue circles), disease states (middle green circles), and individual studies

(innermost circles, colored by fraction of monocytes and macrophages with SCimilarity scores

>95th percentile of all FMΦ SCimilarity scores (log scaled color bar)). Circle size for disease and

individual study are scaled relative to other diseases in the same tissue, or studies in the same

disease. (**d,e**) UMAP embeddings of cell profiles (dots) from the SCimilarity representation (query

model) from an ILD[2] (**d**) and PDAC[3] (**e**) studies, colored by FMΦ query SCimilarity scores (color

bar). **f,** Identification of FMΦs associated genes by importance. Integrated gradients attribution

scores (y axis, top) for the genes (x axis top, and columns, bottom) with top 50 scores for FMΦs

vs. lung macrophages (**Methods**), and their membership (red: presence; grey: absence) in

published macrophage signatures (bottom, rows). Left color bar: AUC of the ranking of each

published signature in SCimilarity attribution scores (AUC=1: all *n* signature genes are listed as

the top *n* genes by SCimilarity attribution scores for distinguishing FMΦ). Martinez-M1 and -M2:

macrophage states expected to be different from FMΦs. P-value: hypergeometric test

Fig. 6. SCimilarity cell search identifies *in vitro* cells matching an *in vivo* FMΦ state and a novel *in vitro* disease model.

a, Identification of FMΦs-like cells across *in vitro* samples by SCimilarity cell search. SCimilarity scores (y axis, $\log_{10}$ scale, and color bar) against a FMΦ query profile for each annotated myeloid cell (dot) from 40 *in vitro* samples (x axis) (from 17 studies), ordered by mean SCimilarity score. Gray boxes: Day 0 and Day 5 samples from a 3D-hydrogel culture system[4]. **b-f,** 3D conditions yield FMΦ-like cells *in vitro* in validation experiments. **b,** SCimilarity scores (y axis, $\log_{10}$ scale, and color bar) against a FMΦ query profile for each annotated myeloid cell (dot) from the original 3D-hydrogel culture system dataset[4] and from 3 donors in the validation experiment (x axis). **c,** Mean expression (dot color) and proportion of expressing cells (dot size) of genes (rows) with high SCimilarity attribution score for distinguishing FMΦs *in vivo* (as in **Fig. 5f**) for myeloid cells in the original 3D-hydrogel culture system[4] and the validation experiment (columns). **d-f,** UMAP embedding from SCimilarity's query model latent space of cell profiles (dots) from day 0 (**d**) or day 5 (**e**) of the original 3D-hydrogel culture system[4] or from day 8 of the replication experiment (**f**), colored by FMΦ SCimilarity score (color bar). **g,** replication of Xu et al.'s original finding of HSC expansion. Proportion of HSCs between Xu et al.'s day 0, day 5 and our validation day 8 time points.

Extended Data figure legends

Extended Data Fig. 1. Data compendium to assemble a pan-human reference.

a,b, Cumulative number of cells (**a**, y axis) and samples (**b**, y axis) profiled by sc/snRNA-seq (and

matching our filters; **Methods**) over time (x axis). Doubling time is calculated based on the

publication date from the most recent 150 data points (dashed red line). **c,** Author-annotated cell

types used in training. Number of author-annotated cells (color bar) from each Cell Ontology type

(rows) and study (columns) used for SCimilarity model training. **d,** Tissues and diseases used in

training. Number of studies (heatmap tiles, text and color bar) and cells (margins, y or x axis) used

for model training from each tissue (rows, right y-axis) and disease (columns, top x-axis).

Extended Data Fig. 2. SCimilarity training details and hyperparameter search.

a, Impact of triplet and autoencoder loss mixing on model performance, where the leftmost column is a traditional autoencoder and the rightmost column is exclusively triplet loss. The FMΦ retrieval test quantifies how much correlation there is between signature scoring of FMΦs and SCimilarity score to FMΦ. The multiple tissue integration test quantifies an ontology-aware average silhouette width where a higher score denotes more coherent clusters for each cell type. The bottom row shows UMAPs for each loss function mix for nine lung scRNA-seq datasets, colored by study. **b,** Benchmarking SCimilarity to established data integration models. Ontology-aware average silhouette width (ASW, y axis, top), normalized mutual information (NMI, y axis, bottom left) and adjusted Rand index (ARI, y axis, bottom right) for SCimilarity's integration and search models and for scVI, scArches, and Harmony (x axis), each applied to nine lung datasets. **c,** Outlier cells from different types and tissues. Fraction (x axis) of cells from different disease (left) or healthy (right) tissue samples with low similarity (SCimilarity score <50) to training data.

Extended Data Fig. 3. Validation of large-scale integration and annotation.

a, Unconstrained cell annotation. UMAP embedding of single cell profiles (dots) from SCimilarity's latent representation of the held out scRNA-Seq kidney data[1] (as in **Fig. 3b,c,** colored by cell annotation without constraining target labels to the scope of author-provided labels in this studyor by expression of select marker genes of regulatory T cells (**b-d)** or myofibroblasts (**e**). **f,** UMAP of 2,000,000 embedded cells uniformly sampled from the 22.7M reference, colored by tissue (as in **Fig. 4b**). **g,h,** UMAP embedding of cell profiles predicted by SCimilarity as fibroblast/myofibroblast (**g**) or monocytes/macrophages (**h**), colored by study (for the 60 studies contributing most cells). **i,** SCimilarity cell-type important genes match cell-type specific signatures. Fraction of cell type-specific differentially expressed genes (from Eraslan *et al.*[5]) (y axis) captured by top-n important genes (x axis) for that cell type by SCimilarity's integrated gradients attribution analysis.

Extended Data Fig. 4. FMΦs among monocytes and macrophages.

a-c, Agreement between SCimilarity and traditional FMΦ cell scores. **a,b** UMAP embedding of

2,578,221 monocyte and macrophage cell profiles (dots) from SCimilarity's latent space

representation colored by SCimilarity score using a prototypical FMΦ cellular profile defined from

Adams *at el.*[6] (**a**) or Scanpy's signature score for FMΦ associated genes (**b**). (**c**) Scanpy FMΦ gene

signature score (x axis) and FMΦ SCimilarity score (y axis) for each cell (shown as density). **d,**

Agreement between SCimilarity FMΦ important genes and published FMΦ signatures. ROC curve

of the fraction of each study's gene sets (y axis) captured within the top genes by SCimilarity

attribution ranking (x axis). **e,** FMΦ important genes are enriched for relevant pathways. False

discovery rate ($-\log_{10}$(q value), hypergeometric test, x axis) for enrichment of Reactome pathways

(y axis, $Q \leq 0.05$ and gene count ≥ 5) with the 100 genes with the top integrated gradients

attribution scores for the FMΦ query (ranked by score). Color: ratio of important genes within a

Reactome pathway to the total size of the pathway. **f-h,** Expression of known and novel genes

associated with FMΦs. Pseudobulked gene expression values for ILD tissue samples for known

marker TREM2 (**f**) and enriched genes not previously described (**g,h**).

1. Young, M. D. *et al.* Single cell derived mRNA signals across human kidney tumors. *Nat. Commun.* **12**, 3896 (2021).

2. Morse, C. *et al.* Proliferating SPP1/MERTK-expressing macrophages in idiopathic pulmonary fibrosis. *Eur. Respir. J.* **54**, (2019).

3. Lin, W. *et al.* Single-cell transcriptome analysis of tumor and stromal compartments of pancreatic ductal adenocarcinoma primary tumors and metastatic lesions. *Genome Med.* **12**, 80 (2020).

4. Xu, Y. *et al.* Efficient expansion of rare human circulating hematopoietic stem/progenitor cells in steady-state blood using a polypeptide-forming 3D culture. *Protein Cell* (2022) doi:10.1007/s13238-021-00900-4.

5. Eraslan, G. *et al.* Single-nucleus cross-tissue molecular reference maps toward understanding disease gene function. *Science* **376**, eabl4290 (2022).

6. Adams, T. S. *et al.* Single-cell RNA-seq reveals ectopic and aberrant lung-resident cell populations in idiopathic pulmonary fibrosis. *Science Advances* **6**, eaba1983 (2020).