Habitat structural complexity increases age-class coexistence and population growth rate through relaxed cannibalism in medaka fish

Eric Edeline[1*], Yoann Bennevault[2], David Rozen-Rechels[3]

1: DECOD (Ecosystem Dynamics and Sustainability), INRAE, IFREMER, Institut Agro, Rennes, France.

2: U3E, INRAE, Agrocampus Ouest, Rennes, France.

3: Sorbonne Université, CNRS, IRD, INRAE, Université Paris Est Créteil, Université Paris Cité, Institute of Ecology and Environmental Sciences of Paris (iEES- Paris), Paris, France.

* corresponding author: eric.edeline@inrae.fr

Key words: Anthropogenic change, Biodiversity, Body size, Cannibalism, Competition, Conservation, Density-dependent population dynamics, Habitat degradation.

ABSTRACT

Structurally-complex habitats harbour more taxonomically-diverse and more productive
20   communities, a phenomenon generally ascribed to habitat complexity relaxing the strength of inter-
specific predation and competition. Here, we challenge this classical, community-centred view by
showing that positive complexity-diversity and complexity-productivity relationships may also
emerge from between-age-class, *intra*-specific interactions at a single-population level. In the
laboratory, we show that medaka fish (*Oryzias latipes*) are strongly cannibalistic in complexity-free
25   habitats, and that cannibalism may occur over a wide range of victim/cannibal body size ratios. In
replicated outdoor pond populations, we manipulated habitat structural complexity using floating
artificial structures, which selectively hampered movements of large-bodied medaka. Habitat
structural complexity relaxed the strength of cannibalism, resulting in (1) increased survival of age-
0+ individuals, (2) elevated age-class diversity through more balanced age-class proportions, (3) a
30   80 % increase in population growth rate, (4) dampened negative density-dependence in the effects
of age-1+ medaka on the recruitment of age-0+ medaka, resulting in elevated habitat carrying
capacity. The resultant higher population density in complex habitats was associated with increased
competition for food among both age-0+ and age-1+ individuals. Our results highlight that positive
complexity-diversity and complexity-productivity relationships may be considered as a generally-
35   emergent property of size-structured populations and communities in which a larger body size
brings a predation or interference advantage. Hence, enhancement of habitat structural complexity
may be seen as a pivotal management strategy not only in favour of taxonomic diversity, but also to
increase the productivity of exploited populations and to improve the conservation status of
endangered species.

40 **INTRODUCTION**

Anthropogenic changes and biodiversity loss are often congruent with a simplification in the physical structure of habitats. For instance, agriculture transforms complex, mature forests into pastures or grain fields that, eventually, may return to secondary forests which are structurally simpler than mature forests (Colorado Zuluaga and Rodewald 2015). In aquatic or terrestrial

45 habitats, top-predator extirpation may have cascading effects resulting in a loss of the large autotrophic organisms that structure space (Estes et al. 2011). In oceans, eutrophication, acidification, bottom trawling and dredging all tend to destroy the complex structure provided by benthic habitats such as coral reefs (National Research Council 2002, Rogers et al. 2014). In freshwaters, channelisation for hydropower, shipping, urban development or log-driving

50 tremendously simplifies river habitats, and eutrophication may drive a loss of submerged macrophytes which are key to structuring rivers, ponds or lakes (Scheffer 2004).

In parallel, many studies report that structurally-complex habitats favour increased species coexistence and higher numerical abundances and biomass in animal communities. This positive

55 effect is generally ascribed to habitat complexity relaxing the strengths of both predation and competition, and thus allowing more species to coexist through increasing surface area, niche diversity and the spatial partitioning of limiting resources (Smith 1972, Crowder and Cooper 1982, Diehl 1988, 1992, Heck and Crowder 1991, Hixon and Menge 1991, Janssen et al. 2007, Kovalenko et al. 2012, Reichstein et al. 2013, Rogers et al. 2014, Soukup et al. 2022).

This is because highly-structured habitats make prey less detectable visually and hinder movements and encounters (Stenseth 1980, Bartholomew et al. 2000, St Pierre and Kovalenko 2014, Soukup et al. 2022), such that predators may be restricted to ambush hunting only (Schultz et al. 2009, Soukup et al. 2022). In particular, in complex habitats larger-bodied individuals selectively suffer from a

65   reduced agility and from a reduced accessibility to microhabitats and crevices (Rogers et al. 2014,

Soukup et al. 2022). Hence, structurally-complex habitats decrease predator attack rates, resulting in

a reduced risk of prey overexploitation at all trophic levels and in relaxed apparent competition

among prey, i.e., relaxed indirect competition through supporting a common predator (Holt 1987).

70   Historically, research on the ecological effects of habitat structural complexity has focused on

interspecific interactions with, to our knowledge, a poor consideration for intraspecific interactions.

This is despite that many populations are size-structured due to both overlapping age classes and

due to heterogeneous somatic growth rates within age classes (Ebenman and Persson 1988). In such

size-structured populations, apparent competition is by definition not possible, but large-bodied

75   individuals often dominate in interference competition (Post et al. 1999, Le Bourlot et al. 2014), and

may even cannibalize smaller-bodied conspecifics (Fox 1975, Smith and Reay 1991, Wise 2006). It

is therefore likely that, in size-structured populations, just like in size-structured communities, a

high habitat structural complexity impedes the dominance of large-bodied individuals.

80   Accordingly, a few studies suggest that habitat structural complexity increases age-class diversity in

the form of enhanced juvenile-adult coexistence, supports larger population sizes, and favours

population persistence in laboratory populations of the guppy *Poecilia reticulata* (Yamagishi 1976,

Nilsson and Persson 2013) and of the mosquitofish *Gambusia affinis* (Benoît et al. 2000). However,

given the paucity of studies dealing with habitat complexity in size-structured populations, the

85   generality of positive complexity-diversity and complexity-productivity relationships at the

population level remains questionable. Additionally, the respective contributions of competition and

cannibalism to the emergence of complexity-productivity relationships at the population level

remain largely unexplored.

In this paper, we test the prediction that habitat structural complexity increases age-class diversity and increases population productivity in the Japanese medaka fish (*Oryzias latipes*), and that the underlying mechanisms involve a relaxation in the strengths of both interference competition and cannibalism. Our approach was to first check that medaka express size-dependent cannibalism in the laboratory, and then to quantify the effects of structural complexity on medaka population dynamics in outdoor-pond experiments. Our results suggest that habitat complexity increases age-class diversity and population productivity, but point to cannibalism as the only underlying mechanism.

MATERIALS AND METHODS

Medaka fish

The medaka is an oviparous fish belonging to the group of Beloniformes, a sister group of Cyprinodontiformes which includes killifishes (Kinoshita et al. 2009). Medaka naturally inhabit slow-moving fresh- and brackish-waters of South-East Asia. Juveniles and adults have a highly-overlapping diet of zooplankton, small benthic invertebrates and filamentous algae (Terao 1985, Edeline et al. 2016). Due to its high thermal tolerance and ease of manipulation, the medaka can be used for parallel experiments in the laboratory, where generation time is 2-3 months under optimal light, food and temperature conditions, and in outdoor ponds, where perennial populations maintain themselves over years under natural conditions without any artificial feeding (Bouffet-Halle et al. 2021). For these reasons, the medaka is a good model species for studies in genetics, developmental biology, ecology and evolution (Kinoshita et al. 2009, Renneville et al. 2016, 2020, Diaz Pauli et al. 2019, 2020, Le Rouzic et al. 2020, Evangelista et al. 2020a, b, 2021, Bouffet-Halle et al. 2021).

Cannibalistic behavioural assays and predation window

115 Trophic interactions are often size-dependent (Woodward et al. 2005), such that the prey/predator body-length ratio is constrained to a specific range called the "predation window" (Claessen et al. 2000). The lower and upper limits of the predation window have far-reaching consequences for population dynamics and individual life histories, respectively (Claessen et al. 2002). We performed cannibalistic behavioural assays to determine whether medaka are cannibalistic under no-

120 complexity conditions and, if so, to estimate the limits of their predation window.

Cannibalistic assays were performed at the Centre de recherche en Ecologie Expérimentale et Prédictive (CEREEP, https://www.cereep.bio.ens.psl.eu/) using the Kiyosu medaka strain (Toyohashi, Aichi Prefecture, Japan, see Bouffet-Halle et al. 2021). Temperature was 21°C (±1.5

125 SD) and the light regime followed the natural photoperiod at the time of experiment (12-13[th] March 2013), i.e., dark from 17:00 to 9:00. Adult medaka were measured individually for standard body length (Sdl, from the tip of the snout to the base of the caudal peduncle, n = 45 adults, mean ± SD = 22.8 mm ± 4.2), determined for sex according to their secondary sexual characters (Yamamoto 1975, Kinoshita et al. 2009), and were kept separately in a 0.5 L complexity-free aquarium filled

130 with dechlorinated tap water. After 72h of adult fastening (i.e., at 15:30 on March 12[th]), two medaka newborns where chosen randomly from a pool of 2-10 days old larvae, were measured individually for Sdl (n = 90 larvae, mean ± SD = 5.6 mm ± 0.7), and were randomly-deposited with an adult in one of the 45 aquariums. It was not possible to follow larvae individually, and we therefore counted the number of surviving larvae in each aquarium after 2.5 hours, and then every 3 hours during 17.5

135 hours (yielding n = 270 observations). This approach made it possible to study the kinetics of larvae survival in each aquarium.

We used the relationship between larvae survival probability and victim/cannibal Sdl ratios (taking the average Sdl for the two larvae) to estimate the lower (δ) and higher (ε) limits of medaka predation window, as well as the optimal victim/cannibal Sdl ratio (ϕ), as defined by Claessen et al. (2002). Specifically, using the probability of a cannibalistic attack estimated at first census (2.5 hours of cannibalistic assays), i.e., when most of size-dependency in cannibalism was expressed (see Methods and Results), we arbitrarily defined δ and ε as the lower- and higher end Sdl ratios, respectively, at which cannibalistic probability becomes less than 0.05, and ϕ as the Sdl ratio at which cannibalistic probability is maximal.

Fig. 1. Geotextile manipulation of habitat structural complexity in pond medaka populations. A: Detailed structure of the geotextile used as artificial refuges to manipulate habitat complexity. The 2 euro-cent coin has a diameter of 22.25mm. The inset shows a juvenile medaka swimming through the geotextile. For a detailed analysis of geotextile structure, see Appendix 1. **B:** A high-complexity pond during the 2021 experiment, hosting single-layered tiles of geotextile, as well as two floating plastic brushes that ensured that spawning substrates for medaka were non-limiting. In 2022, high-complexity habitats were created using five-layered geotextile mats (see Appendix 1). NB: blue ropes ensured that the insect-proof net that covered ponds did not touch water surface.

Geotextile to manipulate habitat complexity

During outdoor-pond experiments (see below), we manipulated habitat structural complexity using tiles of Maccaferri MacMat®, a high-porosity geotextile used in soil-erosion control. This floating geotextile forms a 3D surface with regularly-spaced, 12-13 mm-deep hills and valleys (Fig. 1A). The 3D surface is made from an irregular mesh of polyamide threads, which present a multiplicity

170   of holes of various sizes through which fish can swim (Fig. 1A, Appendix 1). Such a floating

artificial structure is well adapted to providing shelters to medaka, which is a surface-dwelling fish

species (Yamamoto 1975, Kinoshita et al. 2009).

A detailed comparison of geotextile structure with the structure provided by various types of natural

175   vegetation is beyond the scope of this study. However, we note that near-surface or floating

vegetation mats may be formed by riverbank-tree roots, accumulations of dead wood or other

floating debris, by floating macro-algae (*Spirogyra* sp. or *Oedogonium* sp)*,* or by macrophytes, be

they ground-rooted (e.g., *Myriophylumm spp.*) or strictly-floating (e.g., *Utricularia vulgaris* or

Ceratophyllum demersum) in natural water bodies.

To gain some insights into the mechanisms through which the geotextile could affect medaka

behaviour, we measured the distribution of geotextile's hole sizes (in 2D, as seen from a top view),

and compared it with the distributions of medaka body width and height, which are the largest

horizontal and vertical linear distances, respectively, perpendicular to the the fish normal direction

185   of motion (Bartholomew et al. 2000). The distributions of medaka body width and height were fully

included in geotextile's hole-size distribution, suggesting that even the largest-bodied medaka can

swim through the geotextile (Appendix 1). However, small-bodied medaka had many more holes

available to swim through, suggesting that the geotextile would selectively hamper the movement of

the largest medaka.

To test this hypothesis, we performed behavioural assays aiming at measuring the size-dependency

of medaka ability to swim through a single layer of geotextile. The results confirmed that medaka of

all sizes can swim through the geotextile, but that medaka larger than ca. 15 mm in Sdl need much

more time to find their way through the geotextile (Appendix 1). Hence, the geotextile selectively

195   hampered the movements of larger-bodied medaka, exactly as natural habitat structures selectively

hamper the movement of large-bodied predators (Crowder and Cooper 1982, Diehl 1988, 1992, Heck and Crowder 1991, Bartholomew et al. 2000, Kovalenko et al. 2012, Rogers et al. 2014). Although we did not specifically test for visual effects, we hypothesize that the geotextile further interfered with medaka ability to see or sense small-bodied conspecifics (Bartholomew et al. 2000).

We quantified the effects of geotextile-manipulated habitat complexity on medaka population dynamics during two separate outdoor-pond experiments (16, 2 m^2 circular ponds, see below). In 2021, we used single-layered geotextile tiles (12-13 mm deep) of varying surface to induce a complexity contrast. Specifically, single-layered tiles of geotextile covered either a low proportion

205   of pond surface (from 3.1 to 9.9 %, mean 8.6 %, hereafter low-complexity treatment, n = 8 ponds) or a large proportion of pond surface (from 36.2 to 50 %, mean 39.4 %, n = 8 ponds, hereafter high-complexity treatment, Fig. 1B).

In 2022, to induce a more drastic complexity contrast, we used single-layered geotextile tiles in

210   low-complexity habitats and five-layered geotextile mats (90-100 mm deep) in high-complexity habitats (Appendix 1). Specifically, the low-complexity treatment consisted of 0.063 m^2 of single-layered geotextile tiles (3.1 % of pond surface), while the high-complexity treatment consisted of 0.750 m^2 of five-layered geotextile tiles (amounting to 187.5 % of pond surface). Note that ponds were 500-600 mm deep (see below), such that most of the water column was unstructured in both

215   the low- and high-complexity treatments.

In both 2021 and 2022, we added two floating plastic brushes to the ponds to avoid a limitation by the spawning substrate (Fig. 1B).

220   **Pond ecosystems and medaka populations**

On 31st March 2021, 12 circular, 2 m^2 ponds (H = 0.60 m, diam. = 1.65 m, vol. = 1 m^3) were installed at the U3E experimental research unit (https://www6.rennes.inrae.fr/u3e_eng/), filled with dechlorinated tap water, and seeded with 2 L of a mixture of benthic detritus (decaying leaves and phytoplankton), benthic organisms and plankton collected in mature, fishless ponds using a kick net

225   (0.3 mm mesh). Benthos and plankton additions were repeated on April 2nd, 9th and 13th. Additionally, on three occasions (April 2nd, 9th and 15th) each pond received a solution of KH$_2$PO$_4$ amounting to 10 µg L^{-1} of phosphorus. Enrichment was complemented on April 13th, 19th 27th and June 4th with additions of 1 L of an algal mixture (*Chlorella* sp. and *Desmodesmus* sp). All ponds were covered with an insect-proof net (1 mm mesh size). On July 1st 2021, 4 more ponds were

230   installed and submitted to the same seeding and enrichment treatments. Hence, the experiment included 16 ponds in total.

On March 31st 2021 (July 8th for the four extra ponds), from 18 to 60 adult medaka (mean = 37) born in 2020 (hereafter "age-1+") were randomly introduced in ponds. Stocking numbers were n =

235   35 ± 12 fish at low complexity and n = 39 ± 14 fish at high complexity (estimate = 0.107, SE = 0.174, t-value = 0.617, p-value = 0.547). To vary the genetic background of the fish populations, we used medaka originating from the Kiyosu population, as well as five other strains corresponding to different locations in Japan and provided by the Japanese National Bioresource Project[1]: Hamochi, Higashidori, Inawashiro, Tokamachi, and Kushima. The six strains were not mixed in a pond, but

240   seeded each in two separate ponds (one low-complexity pond and one high-complexity pond), except the Kiyosu strain that was seeded in 6 ponds (three low-complexity ponds and three high-complexity ponds). Preliminary analyses of the 2021 data showed that medaka strains had no

---

1   https://shigen.nig.ac.jp/medaka/

significant effect on population growth rate or body sizes (Appendix 2), and we chose to discard the strain effect from our subsequent analyses and experimental design (see below).

From November 15^{th} to 17^{th} 2021, i.e., after the reproductive period, all fish from each pond were sampled using both hand nets and a seine net. To reduce fish handling and to provide fast and reproducible body-size measurements, fish were live-photographed in batches in a transparent tray set above a light source, and were released in their pond. These fish included a mixture of both age-

250 1+ adults that were survivors from initial introductions and age-0+ recruits that were born in the ponds. Each photograph was analysed with ImageJ to automatically fit an ellipse to each fish shape using the "fit-ellipse" tool (Schneider et al. 2012). Sdl was equated to ellipse major distance.

On March 22^{nd} 2022, the 16 ponds were dried, and water and sediments were mixed and randomly

255 redistributed among the 16 ponds. This was the start of a long-term experiment requiring identical initial conditions among ponds. Therefore, we homogeneously mixed medaka strains in each pond by seeding 3, 8, 7, 15, 8 and 25 fish from the Hamochi, Higashidori, Inawashiro, Tokamachi, Kushima and Kiyosu strains, respectively (66 fish per pond). Fish were batch-photographed for Sdl measurement using ImageJ on November 8^{th} and 9^{th} 2022, and were released in their pond to pursue

260 the long-term experiment.

Filamentous algae

Filamentous algae represent an important food source for medaka, both in the wild (Terao 1985, Edeline et al. 2016) and in mesocosms (Bouffet-Halle et al. 2021). In November 2022, before

265 medaka fishing, two independent observers visually estimated the percentage of pond surface covered by filamentous algae in each pond.

Statistical analyses

Cannibalistic behavioural assays

From cannibalistic assays in the laboratory, we estimated how the probability for an average-sized larvae to be cannibalized changed with (i) the victim-to-cannibal Sdl ratio, (ii) the time elapsed since the start of the assays, and (iii) the sex of the cannibal. To that aim, we used an overdispersed binomial (logit link) generalized additive model (GAM) of the form:

$$C_{i,T} \sim \text{Binomial}(2, \pi_{i,T})$$
$$\ln\left(\frac{\pi_{i,T}}{1-\pi_{i,T}}\right) = \beta_{Sex[i]} + f(SdlR_i, T) \quad \text{(Model 1)},$$
$$var(C_{i,T}) = \theta \, 2 \, \pi_{i,T}(1-\pi_{i,T})$$

where ln is the natural logarithm, C is the number of larvae eaten, i indexes an aquarium (and cannibal individual), T indexes the time of census ($i \times T$ = 270 observations), π is cannibalism probability, $\beta_{Sex[i]}$ is a sex effect of cannibal individual i, *SdlR* is mean victim/cannibal Sdl ratio , f is a tensor product of natural cubic splines with 3 knots, which was parsimonious in terms of smoother wigliness and, at the same time, captured the essential features of the cannibalism-size relationship. Finally, θ is a positive parameter accounting for a slight overdispersion in the data. Estimating a different f for each sex increased model's generalised cross-validation score, and we thus preferred the simpler model including *Sex* as a fixed effect on the intercept.

We fitted Model 1 using quasi-likelihood (`quasibinomial` family) in the `mgcv` library of the R software version 4.2.1 (Wood 2017, R Core Team 2024). This model explained 24.5 % of the deviance in the number of larvae eaten. The significance of the effects included in the model was assessed using a standard *t*-test (as provided by the `summary` function in R).

Pond populations

We analysed the effects of habitat structural complexity on pond medaka population dynamics in two steps. Step one aimed a gaining a general understanding of unstructured population dynamics using a classical Ricker model of total population numbers (which were measured with no error). As a second step, we tried to gain a more detailed understanding of the mechanisms behind population dynamics by deciphering age structure in the populations. We did so using a Gaussian mixture model that estimated individual age from body-size distributions.

At step one, we specifically explored unstructured medaka population dynamics using the Ricker logistic equation (Case 2000):

$$\mathrm{N}(t+1) = \mathrm{N}(t)\,e^{\mathrm{r}\left(1 - \frac{\mathrm{N(t)}}{\mathrm{K}}\right)}$$
$$\ln\left(N(t+1)\right) = \ln\left(N(t)\right) + \mathrm{r} - \gamma\, N(t)$$
(Eq. 1),

where $N(t+1)$ is total fish number after reproduction (recruitment), $N(t)$ is the number of age-1+ medaka parents initially introduced before reproduction (stock), r is maximum population growth rate, and $\gamma = \mathrm{r}/\mathrm{K}$ measures the strength of negative density-dependence in the population.

We tested for an effect of habitat structural complexity on the density-*in*dependent r parameter, and on the density-dependent γ parameter using a Poisson GLM:

$$N(t+1)_i \sim \mathrm{Poisson}(\lambda_i)$$
$$\ln(\lambda_i) = \ln\left(N(t)_i\right) + \alpha_y + \beta_{k[i]} - \gamma_{k[i]} N_{st}(t)_i + \epsilon_i$$
(Model 2),
$$\epsilon_i \sim \mathrm{Normal}(0, \sigma_\epsilon^2)$$

where *i* indexes pond-by-year combinations (n = 32), *y* indexes the year of experiment (2021 and 2022), α_y and $\beta_{k[i]}$ are year-of-experiment and habitat-complexity effects on population growth rate, respectively, $\gamma_{k[i]}$ captures a habitat complexity-by-density interaction, and σ_ϵ^2 is

an overdispersion parameter. To avoid any intercept-slope correlation, we standardized $N(t)$ to

zero mean (hence the $N_{st}(t)$ notation in Model 2 equation). As a consequence of this

315   standardisation, the α and β parameters in Model 2 do not estimate the Ricker stock-

recruitment relationship at origin, but at an average level of the stock.

As stressed above, the Ricker-logistic approach in Model 2 is unstructured by lumping together age-

1+ parents and their age-0+ progeny in $N(t+1)$, and is therefore unable to adult survival from

320   the *per capita* production rate of age-0+ recruits (i.e., number of age-0+ at time t+1 / number of

introduced age-1+ at time t). To gain a deeper understanding of medaka population dynamics, we

inferred individual ages at t+1 from the information present in the body-size distributions. At this

step two of our analysis, we deciphered mixtures of size distributions for age-1+ and age-0+ fish,

and their response to habitat complexity, using a log-normal mixture model fitted to standard body

325   lengths Sdl:

$$\ln(\text{Sdl}_i) \sim \sum_{j=1}^{J} \sum_{k=1}^{K} \sum_{p=1}^{P} \pi_{j[i],p[i],y[i]} \text{Normal}(\mu_{j[i],i}, \sigma^2_{j[i],k[i],y[i]})$$
$$\mu_{0+,i} = \alpha 1_{y[i]} + \alpha 2_{k[i]} + \alpha 3_{y[i],k[i]} + \xi_{p[i]}$$
$$\mu_{1+,i} = \beta 1_{y[i]} + \beta 2_{k[i]} + \beta 3_{y[i],k[i]} + \zeta_{p[i]} \qquad \text{(Model 3),}$$
$$\xi_{p[i]} \sim \text{Normal}(0, \sigma^2_\xi)$$
$$\zeta_{p[i]} \sim \text{Normal}(0, \sigma^2_\zeta)$$

where i indexes individual fish (n = 3423), j indexes age groups (age 0+ *vs*. age 1+, J = 2),

k indexes the complexity treatment (low vs. high, K = 2), and p indexes ponds (P = 16).

$\pi_{j[i],p[i],y[i]}$ is the proportion of age j fish in pond p and year of experiment y such that,

330   for each p and y, $\pi_j \geq 0, \sum_{j=1}^{J} \pi_j = 1$. The $\alpha 1_{y[i]}$ and $\beta 1_{y[i]}$ parameters capture year-

of-experiment effects (2021 vs. 2022) on mean body lengths at age-0+ and age-1+, respectively.

The $\alpha 2_{k[i]}$ and $\beta 2_{k[i]}$ parameters capture the effects of habitat structural complexity on mean

body lengths at age 0+ and age 1+, respectively. The $\alpha3_{y[i],k[i]}$ and $\beta3_{y[i],k[i]}$ parameters capture complexity-by-year interactions on mean body lengths at age 0+ and age 1+, respectively.

335   The $\xi_{p[i]}$ and $\zeta_{p[i]}$ parameters capture random pond effects on mean body lengths at age 0+ and age 1+, respectively. Finally, Model 3 included heteroscedasticity in the form of a triple age class-by-complexity-by-year interaction on residual variance $\sigma^2_{j[i],k[i],y[i]}$ (i.e., eight variances were estimated separately).

340   Estimates for the number of individuals in each age class in each pond and each year were computed from Bayesian posterior estimates (see below) of π_j as $\widetilde{N_{j,p,y}}(t+1)=\pi_{j,p,y}N_{p,y}(t+1)$, where $N_{p,y}$ is total fish number. To separate the effects of habitat complexity on *per capita* production rate of age-0+ recruits and on age-1+ parent survival, we used two separate Ricker-logistic models similar to Model 2. We modelled the median posterior

345   number of age-0+ recruits $\widetilde{N}_{0+}(t+1)$ as:

$$\ln(\widetilde{N}_{0+}(t+1)_i)\sim\text{Normal}(\mu_i,\sigma^2_{\widetilde{N}_{0+}(t+1)_i}) \quad \text{(Model 4)},$$
$$\mu_i=\ln(N(t)_i)+\alpha_y+\beta_{k[i]}-\gamma_{k[i]}N_{st}(t)_i$$

where $\sigma^2_{\widetilde{N}_{0+}(t+1)_i}$ is the posterior variance in $\widetilde{N}_{0+}(t+1)$, supplied here as data so as to propagate

350   estimation uncertainty, α_y and $\beta_{k[i]}$ capture the effects of year-of-experiment and habitat complexity, respectively, on *per capita* production rate of age-0+ recruits, evaluated at an average level of the parental stock (i.e., at $N_{st}(t)=0$). The $\gamma_{k[i]}$ captured the density-by-habitat complexity interaction on *per capita* production rate of age-0+ recruits.

355   Following a similar rationale, we modelled the median posterior number of surviving age-1+ fish

after reproduction $\widetilde{N}_{1+}(t+1)$ as:

$$\ln\left(\widetilde{N}_{1+}(t+1)_i\right)\sim \text{Normal}\left(\mu_i, \sigma^2_{\widetilde{N}_{1+}(t+1)_i}\right)$$
$$\mu_i = \ln\left(N(t)_i\right) + \alpha_y + \beta_{k[i]} - \gamma_{k[i]} N_{st}(t)_i$$

(Model 5),

360   where $\sigma^2_{\widetilde{N}_{1+}(t+1)_i}$ is posterior variance in $\widetilde{N}_{1+}(t+1)$, intercepts α and β capture the year-

of-experiment and habitat-complexity effects on mean, ln-transformed survival probabilities of age-

1+ fish through the reproductive period, and slopes $\gamma_{k[i]}$ capture the habitat complexity-by-

density interaction on age-1+ survival probabilities. We preferred Model 5 to a Binomial or a Beta

models, which are more naturally adapted to modelling probabilities, but can not straightforwardly

365   incorporate estimation uncertainty $\sigma^2_{\widetilde{N}_{1+}(t+1)_i}$. However, as a complement to Model 5, we provide a

Binomial analysis of age-1+ survival probabilities in Appendix 3.

We fitted Models 2-5 using Markov Chain Monte Carlo (MCMC) in JAGS 4.3.0 (Plummer 2003)

through the `jagsUI` package (Kellner 2019). Priors were chosen to be weakly informative, except

370   in mixture Model 3, where we imposed the following constraints: (i) mean standard body length

was smaller at age 0+ than at age 1+ (i.e., $\alpha_1 < \beta_1$ and $\alpha_2 < \beta_2$), (ii) body-length variance was

larger at age 0+ than at age 1+, as evidenced by a visual inspection of size distributions (i.e.,

$\sigma^2_{\bar{\xi}} > \sigma^2_{\xi}$), and (iii) the number of age-1+ individuals after reproduction at time t+1 was not larger

than before reproduction at time t (i.e., $\pi_{1+} \leq min\left(1, \dfrac{N(t)}{N(t+1)}\right)$). We further prevented label

375   switching by assigning age-class 0+ to fish shorter than 10 mm and age-class 1+ to fish longer than

30 mm (Edeline et al. 2016, Bouffet-Halle et al. 2021). We ran 3 parallel MCMC chains until

parameter convergence was reached, as assessed using the Gelman–Rubin statistic (Gelman and

Rubin 1992).

380   We assessed goodness of fit of Models 2-5 by using a Bayesian P-value (Gelman et al. 1996).

Briefly, we computed residuals for the actual data as well as for synthetic data simulated from

estimated model parameters (i.e., residuals from fitting the model to ''ideal'' data). The Bayesian P-

value is the proportion of simulations in which ideal residuals are larger than true residuals. If the

model fits the data well, the Bayesian P-value is close to 0.5. Bayesian P values for Models 2-5

385   were 0.49, 0.52, 0.12 and 0.07, respectively, indicating fits ranging from excellent to fair only

(Gelman et al. 1996). The lower fit of Models 4 and 5 is explained by the fact that error variance

was not freely estimated but supplied as data in these models.

The significance of the effects included in Models 2-5 was assessed using MCMC p-values (not to

390   be confounded with the Bayesian P-value above), which quantify posterior overlap with zero.

Specifically, at each MCMC iteration and for each model parameter, we computed Δ as the

difference between posterior parameters under low and high habitat complexity. We then computed

MCMC p-values as twice the proportion of Δ for which the sign of Δ was opposite to that of its

mean value.

Finally, the effect of habitat complexity on the percentage of pond surface covered by filamentous

algae Pr was modelled using a Beta GLMM:

$$
\begin{aligned}
Pr_i &\sim Beta(p_i, q_i) \\
p_i &= \mu_i * \tau_i \\
q_i &= \tau_i - \mu_i * \tau_i \\
\ln\left(\frac{\mu_i}{1-\mu_i}\right) &= \alpha_{k[i]} + \beta_{p[i]} \\
\beta_{p[i]} &\sim Normal(0, \sigma_\beta^2)
\end{aligned}
\qquad \text{(Model 6),}
$$

where *i* indexes an observer-by-pond combination (n = 32 observations), $\alpha_{k[i]}$ captures the effect

of habitat structural complexity (low vs. high), and $\beta_{p[i]}$ is a random pond effect (n = 16 ponds).

We fitted Model 6 in the `glmmTMB` library of R (Brooks et al. 2017), and evaluated significance of

the complexity effect using the `summary` function of R.

<h1 style="text-align:center">RESULTS</h1>

Cannibalistic behavioural assays and predation window

The relationship between medaka cannibalistic behaviour and the victim/cannibal standard body

length (Sdl) ratio predicted by GAM Model 1 was nonlinear, and changed during the course of

cannibalistic assays (Fig. 2A). At first census, after 2.5 hours of exposure, freely-adjusting splines

showed that the relationship followed a bell-shaped curve, thus validating the hypotheses of the

predation window (Fig. 2B).

Fig. 2. Cannibalistic behaviour in the laboratory with no habitat structure. A: Surface plot for the nonlinear interaction between victim/cannibal standard body length (Sdl) ratio and time of exposure on the probability for cannibalism to occur. Blue solid lines: male cannibals; Pink dashed lines: female cannibals. The surfaces were predicted from Model 1. Dots show the fitted values: Circles: male cannibals; Triangles: female cannibals. **B:** Defining parameters of the predation window from predicted cannibalism probability surfaces. The cannibalistic curves are the same as in Fig. 2A for the fist census (2.5 hours of cannibalistic assays). δ and ε are the lower and upper limits of the predation window, arbitrarily defined as the lower- and upper-end Sdl ratios, respectively, at which cannibalistic probability becomes less than 0.05 (horizontal dashed line). ϕ is the optimal victim-to-cannibal Sdl ratio. The resultant predation window is plotted in Fig. 3. Note that the higher cannibalistic voracity in males results in a wider predation window.

As time of victim to cannibal exposure was increasing, the relationship progressively became body-

435   size independent. By the end of the assays (17.5 hours of exposure), victim survival probability was

very low under almost all Sdl ratios (Fig. 2A). GAM Model 1 further showed that overall mean

cannibalism probability was lower in female than male cannibals (Fig. 2A; β_{Male} = 1.01, SE =

0.288, t-value = 3.52, p-value < 0.001).

Fig. 3. Predation window for victim *vs*. cannibal standard body length (Sdl) in medaka. The black solid line shows the optimal victim length while ribbons show the upper and lower predation limits for males (blue solid lines) and females (pink dashed lines), as defined by the 5 % cannibalism probabilities displayed in Fig. 2B. The top marginal distribution shows the body sizes typically present in pond medaka populations in France before reproduction in March (Bouffet-Halle et al. 2021), i.e., the size distribution of potential cannibals. The right-hand marginal distribution shows larval body sizes at hatch (Renneville et al. 2020), i.e., the size distribution of potential victims. Vertical and horizontal dashed lines show the 2.5 and 97.5 % quantiles of Sdl distributions.

Due to this sex effect on cannibalistic voracity, the upper and lower limits of the predation window

were sex-dependent (Fig. 2B). Specifically, the predation window was slightly wider in male (δ

= 0.09, ϕ = 0.26, ε = 0.41) than in female cannibals (δ = 0.13, ϕ = 0.26, ε = 0.38).

460   These parameters show that newly-hatched medaka larvae are under strong cannibalistic risk (Fig.

3). An average-sized hatchling (3.8 mm Sdl) may be cannibalised by any 9.3-to-42.2 mm Sdl male

(10.0 to 29.2 mm female) conspecific. This body-size range encompasses not only the whole range

of parental body sizes, but also late-juvenile body sizes (Fig. 3), indicating that medaka larvae are

exposed to both inter- and intra-cohort cannibalism.

Table 1. Parameter estimates from Bayesian Models 2-5. The models were fitted using an "effect" parametrization, i.e., intercepts and slopes are evaluated at a reference level (low complexity and year-of-experiment 2021), coded with a "0" in the "Parameter column", and effects of high habitat complexity and year-of experiment 2022 are estimated as deviations from this reference level. MCMC P-values are twice the proportion of the posterior which sign was opposite to the sign of the posterior mode. P-values significant at the 5 % risk are bold-faced. MCMC P-values are not relevant, and thus not supplied, for variance parameters.

Response	Model	Distribution	Link	Effect	Parameter	mean	sd	Rhat	MCMC P-value
Recruitment Rate $\frac{N(t+1)}{N(t)}$	2	Poisson	Log	Model intercept (year 2021, Low complexity)	$\alpha 0$	0.636	0.115	1.001	**0.000**
				Year 2022 effect	α	-0.478	0.176	1.001	**0.013**
				High complexity effect	β	0.596	0.094	1.000	**0.000**
				Nst(t) at Low complexity	$\gamma 0$	-0.021	0.006	1.001	**0.001**
				Nst(t), High complexity effect	γ	0.015	0.006	1.001	**0.010**
				Overdispersion	σ^2_ε	0.236	0.040	1.001	
Standard Body Length Log(Sdl)	3	Gaussian	Identity	Intercept on age-0+ (year 2021, Low complexity)	$\alpha 0$	2.551	0.046	1.003	**0.000**
				Year 2022 effect on age-0+	$\alpha 1$	0.286	0.127	1.000	**0.045**
				High complexity effect on age-0+	$\alpha 2$	-0.179	0.041	1.000	**0.000**
				Year 2022-by-High complexity interaction on age-0+	$\alpha 3$	-0.147	0.136	1.000	0.260
				Intercept on age-1+ (year 2021, Low complexity)	$\beta 0$	3.152	0.024	1.000	**0.000**
				Year 2022 effect on age-1+	$\beta 1$	-0.071	0.021	1.000	**0.001**
				High complexity effect on age-1+	$\beta 2$	-0.014	0.024	1.000	0.553
				Year 2022-by-High complexity interaction on age-1+	$\beta 3$	-0.099	0.031	1.000	**0.005**
				Pond random effect on age-0+	σ^2_γ	0.159	0.034	1.001	
				Pond random effect on age-1+	σ^2_ζ	0.059	0.015	1.001	
Age-0+ Production Rate $\ln(\widetilde{N}_{0+}(t+1))$	4	Gaussian	Identity	Model intercept (year 2021, Low complexity)	$\alpha 0$	-0.181	0.036	1.000	**0.000**
				Year 2022 effect	α	-0.619	0.034	1.001	**0.000**
				High complexity effect	β	1.324	0.037	1.000	**0.000**
				Nst(t) at Low complexity	$\gamma 0$	-0.049	0.002	1.000	**0.000**
				Nst(t), High complexity effect	γ	0.039	0.002	1.000	**0.000**
Age-1+ Survival Rate $\ln(\widetilde{N}_{1+}(t+1))$	5	Gaussian	Identity	Model intercept (year 2021, Low complexity)	$\alpha 0$	-0.052	0.032	1.000	0.104
				Year 2022 effect	α	-0.230	0.075	1.001	**0.003**
				High complexity effect	β	-0.043	0.045	1.001	0.347
				Nst(t) at Low complexity	$\gamma 0$	0.008	0.004	1.000	**0.039**
				Nst(t), High complexity effect	γ	0.012	0.003	1.001	**0.002**

Pond populations

In replicated pond populations, complex habitats increased the population growth rate (Table 1-

475   Model 2, positive β effect), as indicated by green lines being higher than red lines in Fig. 4A.

Specifically, population growth rate increased on average from 1.5 year^{-1} under a low complexity to

2.7 year^{-1} under a high complexity (80 % increase).

Fig. 4. Effect of habitat structural complexity on medaka population dynamics. A. Relationship between the number of medaka after ($N(t+1)$) and before reproduction ($N(t)$) in replicated pond populations. Dots show the raw data (squares: low complexity; diamonds: high complexity), and curves show median posterior predictions from Model 2 (violet-red: low-complexity habitats; Green: high-complexity habitats). Ribbons show 95 % credible intervals. The solid, straight y = x black line intersects coloured curves at habitat carrying capacity K. **B.** Body-length distributions after reproduction at *t*+1. Light bars show the raw data and curves show median posterior distributions predicted from mixture Model 3. There are 8 curves, one per age-class per complexity treatment per year of experiment.

Additionally, medaka population dynamics were negatively density-dependent under a low habitat

complexity (Table 1-Model 2, significantly negative γ_0 effect), but a high habitat complexity

500   dampened this negative density dependence (Table 1-Model 2, positive γ effect). Accordingly,

the stock-recruitment relationship changed from flat at low habitat complexity (red lines) to positive

at high habitat complexity (green lines), reflecting increased habitat carrying capacity (Fig. 4A).

Fitting Model 2 without centring $N(t)$, so as to preserve the original biological interpretation of

 r (see Methods), yielded a median posterior for K that increased from 82 under low habitat

505   complexity to 194 under high habitat complexity in 2021, and from 59 to 143 in 2022.

Mixture Model 3 allowed us to quantify the effects of habitat structural complexity on age-specific, natural log-transformed body-length distributions. Increased habitat complexity decreased mean age-0+ ln(Sdl) in both years of experiment (Table 1-Model 3, α_2 and α_3 parameters), and

510 decreased mean age-1+ ln(Sdl) in 2022, when the complexity contrast was more extreme, but not significantly so in 2021 (Table 1-Model 3, β_3 *vs.* β_2 parameters).

In 2021, more complex habitats increased variability in age-0+ ln(Sdl) (Model 3, $\sigma_{\text{age-0+, Low-complexity, 2021}} = 0.269$ *vs.* $\sigma_{\text{age-0+, High-complexity, 2021}} = 0.303$, MCMC P-value = 0.022), but

515 decreased variability in age-1+ ln(Sdl) (Model 3, $\sigma_{\text{age-1+, Low-complexity, 2021}} = 0.140$ *vs.* $\sigma_{\text{age-1+, High-complexity, 2021}} = 0.113$, MCMC P-value = 0.038). In 2022, age-0+ fish were too few in low-complexity ponds to provide any robust inference on Sdl variability (MCMC P-value = 0.652), while more complex habitats increased age-1+ ln(Sdl) (Model 3, $\sigma_{\text{age-1+, Low-complexity, 2022}} = 0.102$ *vs.* $\sigma_{\text{age-1+, High-complexity, 2022}} = 0.136$, MCMC P-value = 0).

Model 2 is a model of unstructured population dynamics, where age-0+ and age-1+ individuals are lumped in $N(t+1)$. Hence, from Model 2, it is not possible to tell whether the increase in $N(t+1)/N(t)$ under a high habitat complexity reflected increased *per capita* production rate of age-0+ recruits and/or increased survival probability of age-1+ fish. To answer this question,

525 mixture Model 3 provided us with separate estimates for the numbers of age-0+ and age-1+ fish at *t*+1, and thus allowed us to separate these two effects using Models 4 and and 5 (see Methods).

Model 4, which was fitted to the posterior number of age-0+ medaka, perfectly parallel results from Model 2 above. Specifically, *per capita* production of age-0+ fish strongly increased from on

530 average 0.8 at low complexity to 3.1 under a high habitat complexity (Table 1-Model 4, positive β

effect). Additionally, the habitat complexity-by-density interaction reveals a strong negative density dependence under a low habitat complexity (Table 1-Model 4, negative γ_0 effect) that was dampened under a high habitat complexity (Table 1-Model 4, positive γ effect).

535   Patterns of age-1+ survival probability, as inferred from Model 5, were opposite to patterns of *per capita* production rates of age-0+ recruits, suggesting opposite effects of habitat complexity on age-0+ and age-1+ fish. Specifically, a high habitat complexity *decreased* age-1+ survival probability in both the Gaussian and binomial models, but significantly-so in the binomial model only (Table 1-Model 5 *vs*. Appendix 3: negative β effects). The binomial model predicted that age-1+ survival

540   probability decreased from 0.91 at low complexity to 0.77 under a high habitat complexity.

Surprisingly, survival probability of age-1+ fish was *positively* density-dependent under low habitat complexity in both the Gaussian and binomial models (Table 1-Model 5 and Appendix 3: positive γ_0 effects), and a high habitat complexity amplified this positive density dependence, but

545   significantly-so in the Gaussian model only (Table 1-Model 5 *vs*. Appendix 3: positive γ effects), suggesting a weak amplification.

Taken together, results from Models 4 and 5 indicate that the positive effect of habitat complexity on medaka population growth rate resulted from increased *per capita* production of age-0+ fish, and

550   was weakly opposed by depressed survival of age-1+fish.

Filamentous algae

On average, filamentous algae covered 7.1 % of pond surface in high-complexity ponds, and 13.8 % in low-complexity ponds (Fig. 5), suggesting reduced food availability in more complex habitats.

555 Analysis of Model 6 showed that this difference was statistically significant (α_{Low} = 0.728, SE = 0.259, z-value = 2.81, p-value < 0.005).

560 **Fig. 5. Filamentous algae.** Algal cover on each pond was estimated independently by two observers before medaka fishing in November 2022. Symbols show the raw data (n = 32).

DISCUSSION

In multispecies assemblages, habitat structural complexity increases species coexistence and productivity through a relaxation in the strengths of both interspecific competition and predation 570 (Smith 1972, Crowder and Cooper 1982, Holt 1987, Diehl 1988, 1992, Heck and Crowder 1991, Hixon and Menge 1991, Rogers et al. 2014). Our present results in medaka fish show that habitat structural complexity may further increase age-class coexistence, population growth rate and habitat carrying capacity at a single-population level. Hence, intraspecific processes may provide a substantial, but currently neglected, contribution to the positive effects of habitat complexity on 575 biodiversity and productivity at a community level.

Our results may further be used to gain some understanding of the underlying ecological mechanisms that drive positive habitat complexity-productivity relationships. We show that increased age-class coexistence and productivity in medaka populations were mainly achieved

580   through an increased *per capita* production of age-0+ recruits by their age-1+ parents. In theory, such an increase in recruitment may result from increases in both parental fecundity (i.e., increased egg production) and in larval survival to the recruit stage. However, several lines of evidence point to increased larval survival as the only mechanism at play.

585   Filamentous algae, one of the main food sources for medaka in mesocosms, were less abundant in high-complexity than in low-complexity ponds. As a result of this food shortage, both age-0+ and age-1+ medaka had smaller and more variable body sizes in more complex habitats, indicating increased competition for food (Ohlberger et al. 2013). Age-1+ survival probability further decreased in more complex habitats, consistent with negative effects of food competition on fitness

590   in age-1+ medaka, which are dominated by their age-0+ progeny in exploitative competition for food (Edeline et al. 2016). Hence, although more complex habitats increase substrate area, leading to potentially higher production of primary producers and sessile animals (Soukup et al. 2022), we conclude that food stress was stronger in more complex habitats in our experiments due to higher fish densities. Therefore, fecundity of age-1+ medaka most likely *decreased* under a higher habitat

595   complexity, and a higher recruitment in more complex habitats is most likely explained by increased survival of age-0+ medaka due to relaxed cannibalism.

Interestingly, habitat complexity did not only relax overall age-0+ vulnerability to cannibalism, but also interfered with density-dependence in cannibalism-mediated processes. Specifically, habitat

600   complexity relaxed both (i) the *negative* effect of age-1+ density on age-0+ survival, and (ii) the *positive* effect of age-1+ density on their own survival probability, the later being explained by decreasing densities of dominant age-0+ medaka relaxing the competition for food experienced by competitively-inferior age-1+ medaka (Dionne 1985, Claessen et al. 2000). Therefore, our study suggests that habitat structural complexity may reduce both the rate of dangerous encounters

605 between conspecifics and the likelihood that such encounters will lead to successful cannibalistic

attacks (Rosenheim and Schreiber 2022).

In the wild in Japan, medaka starve to death when they turn age-1+, while their newly-hatched

progeny grow very rapidly (Terao 1985, Awaji and Hanyu 1987, Egami et al. 1988). This

610 "semelparous" life history pattern seems to reflect a competitive exclusion of age-1+ parents by

their age-0+ progeny (Edeline et al. 2016). In contrast, in experimental ponds in France medaka

survive to age 2+ and contribute to two reproductive bouts in their lives, i.e., are "iteroparous" (this

study, Bouffet-Halle et al. 2021). In multiple taxa, cannibalism allows large-bodied individuals to

overturn the competitive superiority of smaller-bodied conspecifics (this study, Dionne 1985, Polis

615 1988, Claessen et al. 2000, Wise 2006), and increased cannibalism, as resulting from a lower habitat

complexity, is probably an important driver of the extended lifespan of age-1+ medaka in

experimental ponds.

Accordingly, compared to experimental ponds where complexity was restricted to top water layers,

620 wild habitats harbour rooted water plants or wood debris that provide refuges across the whole

water column down to benthic habitats, which provide substantial amounts of food to medaka when

their preferred planktonic prey are scarce (Terao 1985). Additionally, wild habitats often harbour

shallow and warm areas that are particularly sought after by medaka fish (Awaji and Hanyu 1987),

but which expose large-bodied medaka to stranding and thus protect small-bodied juveniles against

625 cannibalism. Such shallow areas were absent from our experimental ponds.

If the higher complexity of wild habitats explains relaxed cannibalism and the resultant dominance

of age-0+ medaka in the wild, then we could expect a hump-shaped habitat complexity-productivity

relationship in medaka populations, with a shift along an increasing habitat-complexity gradient

630   from a cannibal-mediated, age-1+ dominance at low complexity to an exploitative competition-mediated, age-0+ dominance at high complexity, resulting in maximal age-class coexistence and population productivity at intermediate complexity levels where the ontogenetic asymmetry in dominance is most relaxed. Maximal diversity and productivity at intermediate levels of habitat structural complexity are indeed observed in complex, multispecies communities (Crowder and

635   Cooper 1982, Rogers et al. 2018), supporting the contention that complexity-diversity and complexity-productivity relationships emerge through similar, size-dependent predation processes operating at both the community and population levels.

Size-dependent cannibalism is a very common form of interaction in the animal kingdom (Fox

640   1975, Polis 1981, Smith and Reay 1991, Wise 2006), and cannibalism plays a key role in determining life histories and population dynamics in wild populations of annelids (Elliott 2004), insects (Baskauf 2009), crustaceans (Grosholz et al. 2021), acarids (Walde et al. 1992), fish (Claessen et al. 2000, 2002, Persson et al. 2003), or amphibians (Wissinger et al. 2010). We thus expect habitat complexity, through relaxing cannibalism, to enhance age-class coexistence and

645   population growth rate across a broad taxonomic range. Around the globe, habitat structural complexity becomes increasingly threatened by anthropogenic perturbations such as deforestation, eutrophication, bottom trawling and dredging, river channelization, by trophic cascades resulting from top-predator extinction, or by ocean acidification, thus making global change convergent with a general habitat simplification. Ample evidence already demonstrates that habitat complexity

650   supports species diversity, such that habitat simplification may entail a cascade of species extinctions in multispecies communities (Smith 1972, Crowder and Cooper 1982, Diehl 1988, 1992, Heck and Crowder 1991, Hixon and Menge 1991, Janssen et al. 2007, Kovalenko et al. 2012, Reichstein et al. 2013, Rogers et al. 2014). Our study further suggests that, within the surviving

species, overwhelming dominance of large-bodied individuals may also reduce juvenile survival,
655 and ultimately impair the growth and long-term persistence of populations.

Acknowledgement. We are grateful to the NBRP medaka (https://shigen.nig.ac.jp/medaka/) and to Prof. Kiyoshi Naruse (NIBB Okazaki) for providing us with the medaka strains. We thank André De Roos for insightful comments on an earlier manuscript version. Alexandre Kempf, Solène
660 Moulin, Louisiane Perrin and Morgan Verdeil provided invaluable help during behavioural assays. This experiment was approved by the Charles Darwin Ethical Committee (Ce5/2010/041).

Author contributions: EE designed and contributed to performing experiments, analysed data and wrote the first draft version. EE, YB and DRR performed experiments. All authors contributed to
665 result interpretation and draft improvements.

Funding: This work has benefited from technical and human resources provided by CEREEP-Ecotron IleDeFrance (CNRS/ENS UMS 3194) as well as from financial support from the Regional Council of Ile-de-France under the DIM Program R2DS bearing the references I-05-098/R and
670 2015-1657. It has received a support under the program 'Investissements d'Avenir' launched by the French government and implemented by ANR with the references ANR-10-EQPX-13-01 Planaqua and ANR-11-INBS-0001 AnaEE France, and from Pépinière interdisciplinaire CNRS de site PSL (Paris-Sciences et Lettres) "Eco-Evo-Devo". EE was further supported by grants from Sorbonne Université (program Convergences, project C14234) and from Rennes Métropole (AIS 18C0356).
675

Conflict of interest: The authors declare that they have no conflict of interest relating to the content of this article.

Data, script and code availability: All data and codes needed to run the analyses and to reproduce

680   figures may be downloaded from Zenodo: https://doi.org/10.5281/zenodo.14044171

References

Awaji, M. and Hanyu, I. 1987. Annual reproductive cycle of the wild type medaka. - Nippon Suisan Gakkai Shi Bull. Jpn. Soc. Sci. Fish. 53: 959–965.

Bartholomew, A., Diaz, R. J. and Cicchetti, G. 2000. New dimensionless indices of structural habitat complexity: predicted and actual effects on a predator's foraging success. - Mar. Ecol. Prog. Ser. 206: 45–58.

Baskauf, S. 2009. Factors influencing population dynamics of the southwestern corn borer (Lepidoptera: Crambidae): a reassessment. - Environ. Entomol. 32: 915–928.

Benoît, H. P., Post, J. R., Barbet, A. D. and Ross, S. T. 2000. Recruitment dynamics and size structure in experimental populations of the mosquitofish, *Gambusia affinis*. - Copeia 2000: 216–221.

Bouffet-Halle, A., Mériguet, J., Carmignac, D., Agostini, S., Millot, A., Perret, S., Motard, E., Decenciere, B. and Edeline, E. 2021. Density-dependent selection mediates harvest-induced trait changes. - Ecol. Lett. 24: 648–657.

Brooks, M. E., Kristensen, K., Benthem, K. J. van, Magnusson, A., Berg, C. W., Nielsen, A., Skaug, H. J., Mächler, M. and Bolker, B. M. 2017. glmmTMB balances speed and flexibility among packages for zero-inflated generalized linear mixed modeling. - R J. 9: 378–400.

Case, T. J. 2000. An illustrated guide to theoretical ecology. - Oxford University Press.

Claessen, D., de Roos, A. M. and Persson, L. 2000. Dwarfs and giants: cannibalism and competition in size-structured populations. - Am. Nat. 155: 219–237.

Claessen, D., van Oss, C., de Roos, A. M. and Persson, L. 2002. The impact of size-dependent predation on population dynamics and individual life history. - Ecology 83: 1660–1675.

Colorado Zuluaga, G. J. and Rodewald, A. D. 2015. Response of mixed-species flocks to habitat alteration and deforestation in the Andes. - Biol. Conserv. 188: 72–81.

Crowder, L. B. and Cooper, W. E. 1982. Habitat structural complexity and the interaction between bluegills and their prey. - Ecology 63: 1802–1813.

Diaz Pauli, B., Garric, S., Evangelista, C., Vøllestad, L. A. and Edeline, E. 2019. Selection for small body size favours contrasting sex-specific life histories, boldness and feeding in medaka, *Oryzias latipes*. - BMC Evol. Biol. 19: 127.

Diaz Pauli, B., Edeline, E. and Evangelista, C. 2020. Ecosystem consequences of multi-trait response to environmental changes in Japanese medaka, *Oryzias latipes*. - Conserv. Physiol. 8: coaa011.

Diehl, S. 1988. Foraging efficiency of three freshwater fishes: effects of structural complexity and light. - Oikos 53: 207–214.

Diehl, S. 1992. Fish predation and benthic community structure: the role of omnivory and habitat complexity. - Ecology 73: 1646–1661.

Dionne, M. 1985. Cannibalism, food availability, and reproduction in the mosquito fish (*Gambusia affinis*): a laboratory experiment. - Am. Nat. 126: 16–23.

1988. Size structured populations, ecology and evolution (B Ebenman and L Persson, Eds.). - Springer.

Edeline, E., Terao, O. and Naruse, K. 2016. Empirical evidence for competition-driven semelparity in wild medaka. - Popul. Ecol. 58: 371–383.

Egami, N., Terao, O. and Iwao, Y. 1988. The life span of wild populations of the fish *Oryzias latipes* under natural conditions. - Zoolog. Sci. 5: 1149–1152.

Elliott, J. M. 2004. Contrasting dynamics in two subpopulations of a leech metapopulation over 25 year-classes in a small stream. - J. Anim. Ecol. 73: 272–282.

Estes, J. A., Terborgh, J., Brashares, J. S., Power, M. E., Berger, J., Bond, W. J., Carpenter, S. R., Essington, T. E., Holt, R. D., Jackson, J. B. C., Marquis, R. J., Oksanen, L., Oksanen, T., Paine, R. T., Pikitch, E. K., Ripple, W. J., Sandin, S. A., Scheffer, M., Schoener, T. W., Shurin, J. B., Sinclair, A. R. E., Soulé, M. E., Virtanen, R. and Wardle, D. A. 2011. Trophic downgrading of planet earth. - Science 333: 301–306.

Evangelista, C., Diaz Pauli, B., Vøllestad, L. A. and Edeline, E. 2020a. Stoichiometric consequences of size-selective mortality: an experimental test using the Japanese medaka (*Oryzias latipes*). - Sci. Total Environ. 724: 138193.

Evangelista, C., Vøllestad, L. A., Diaz Pauli, B. and Edeline, E. 2020b. Density-dependent consequences of size-selective induced life-history changes to population fitness in medaka (*Oryzias latipes*). - Can. J. Fish. Aquat. Sci. 77: 1741–1748.

Evangelista, C., Dupeu, J., Sandkjenn, J., Pauli, B. D., Herland, A., Meriguet, J., Vøllestad, L. A. and Edeline, E. 2021. Ecological ramifications of adaptation to size-selective mortality. - R. Soc. Open Sci. 8: 210842.

Fox, L. R. 1975. Cannibalism in natural populations. - Annu. Rev. Ecol. Syst. 6: 87–106.

Gelman, A. and Rubin, D. B. 1992. Inference from iterative simulation using multiple sequences. - Stat. Sci: 457–472.

Gelman, A., Meng, X. L. and Stern, H. 1996. Posterior predictive assessment of model fitness via realized discrepancies. - Stat. Sin. 6: 733–807.

Grosholz, E., Ashton, G., Bradley, M., Brown, C., Ceballos-Osuna, L., Chang, A., de Rivera, C., Gonzalez, J., Heineke, M., Marraffini, M., McCann, L., Pollard, E., Pritchard, I., Ruiz, G., Turner, B. and Tepolt, C. 2021. Stage-specific overcompensation, the hydra effect, and the failure to eradicate an invasive predator. - Proc. Natl. Acad. Sci. 118: e2003955118.

Heck, K. L. and Crowder, L. B. 1991. Habitat structure and predator—prey interactions in vegetated aquatic systems. - In: Bell, S. S. et al. (eds), Habitat structure: the physical arrangement of objects in space. Springer Netherlands, pp. 281–299.

Hixon, M. A. and Menge, B. A. 1991. Species diversity: prey refuges modify the interactive effects of predation and competition. - Theor. Popul. Biol. 39: 178–200.

Holt, R. D. 1987. Prey communities in patchy environments. - Oikos 50: 276–290.

Janssen, A., Sabelis, M. W., Magalhaes, S., Montserrat, M. and van der Hammen, T. 2007. Habitat structure affects intraguild predation. - Ecology 88: 2713–2719.

Kellner, K. 2019. jagsUI: a wrapper around "rjags" to streamline "JAGS" analyses.

Kinoshita, M., Murata, K., Naruse, K. and Tanaka, M. 2009. Medaka. Biology, management and experimental protocols. - Wiley.

Kobayashi, S. 2002. Growth and movement of medaka (Japanese killifish), *Oryzias latipes* in Nagano prefecture rice paddies. - Bull. Nagano Nat. Conserv. Res. Inst. 5: 9–14.

Kovalenko, K. E., Thomaz, S. M. and Warfe, D. M. 2012. Habitat complexity: approaches and future directions. - Hydrobiologia 685: 1–17.

Le Bourlot, V., Tully, T. and Claessen, D. 2014. Interference versus exploitative competition in the regulation of size-structured populations. - Am. Nat. 184: 609–623.

Le Rouzic, A., Renneville, C., Millot, A., Agostini, S., Carmignac, D. and Edeline, E. 2020. Unidirectional response to bidirectional selection on body size. II Quantitative genetics. - Ecol. Evol. 10: 11453–11466.

Meng, Y., Lin, B.-L., Tominaga, M. and Nakanishi, J. 2006. Simulation of the population-level effects of 4-nonylphenol on wild Japanese medaka (*Oryzias latipes*). - Ecol. Model. 197: 350–360.

National Research Council 2002. Effects of trawling and dredging on seafloor habitat. - The National Academies Press.

Nilsson, K. A. and Persson, L. 2013. Refuge availability and within-species differences in cannibalism determine population variability and dynamics. - Ecosphere 4: art100.

Ohlberger, J., Otero, J., Edeline, E., Winfield, I. J., Stenseth, N. Chr. and Vøllestad, L. A. 2013. Biotic and abiotic effects on cohort size distributions in fish. - Oikos 122: 835–844.

Persson, L., De Roos, A. M., Claessen, D., Byström, P., Lövgren, J., Sjögren, S., Svanbäck, R., Wahlström, E. and Westman, E. 2003. Gigantic cannibals driving a whole-lake trophic cascade. - Proc. Natl. Acad. Sci. 100: 4035–4039.

Pinheiro, J., Bates, D., DebRoy, S., Sarkar, D., and R Core Team 2018. nlme: linear and nonlinear mixed effects models.

Plummer, M. 2003. JAGS: a program for analysis of Bayesian graphical models using Gibbs sampling.

Polis, G. A. 1981. The evolution and dynamics of intraspecific predation. - Annu. Rev. Ecol. Syst. 12: 225–251.

Polis, G. A. 1988. Exploitation competition and the evolution of interference, cannibalism, and intraguild predation in age/size-structured populations (B Ebenman and L Persson, Eds.). - Size-Struct. Popul.: 185–202.

Post, J. R., Parkinson, E. A. and Johnston, N. T. 1999. Density-dependent processes in structured fish populations: interaction strengths in whole-lake experiments. - Ecol. Monogr. 69: 155–175.

R Core Team 2024. R: a language and environment for statistical computing. - R Foundation for Statistical Computing.

Reichstein, B., Schröder, A., Persson, L. and De Roos, A. M. 2013. Habitat complexity does not promote coexistence in a size-structured intraguild predation system. - J. Anim. Ecol. 82: 55–63.

Renneville, C., Le Rouzic, A., Baylac, M., Millot, A., Loisel, S. and Edeline, E. 2016. Morphological drivers of trophic cascades. - Oikos 125: 1193–1202.

Renneville, C., Millot, A., Agostini, S., Carmignac, D., Maugars, G., Dufour, S., Le Rouzic, A. and Edeline, E. 2020. Unidirectional response to bidirectional selection on body size. I. Phenotypic, life-history, and endocrine responses. - Ecol. Evol. 10: 10571–10592.

Rogers, A., Blanchard, J. L. and Mumby, P. J. 2014. Vulnerability of coral reef fisheries to a loss of structural complexity. - Curr. Biol. 24: 1000–1005.

Rogers, A., Blanchard, J. L. and Mumby, P. J. 2018. Fisheries productivity under progressive coral reef degradation. - J. Appl. Ecol. 55: 1041–1049.

Rosenheim, J. A. and Schreiber, S. J. 2022. Pathways to the density-dependent expression of cannibalism, and consequences for regulated population dynamics. - Ecology 103: e3785.

Scheffer, M. 2004. Ecology of shallow lakes. - Springer.

Schneider, C. A., Rasband, W. S. and Eliceiri, K. W. 2012. NIH Image to ImageJ: 25 years of image analysis. - Nat. Methods 9: 671–675.

Schultz, S., Kruschel, C. and Bakran-Petricioli, T. 2009. Influence of seagrass meadows on predator–prey habitat segregation in an Adriatic lagoon. - Mar. Ecol. Prog. Ser. 374: 85–99.

Smith, F. E. 1972. Spatial heterogeneity, stability and diversity in ecosystems. - Trans. Conn. Acad. Arts Sci. 44: 309–335.

Smith, C. and Reay, P. 1991. Cannibalism in teleost fish. - Rev. Fish Biol. Fish. 1: 41–64.

Soukup, P. R., Näslund, J., Höjesjö, J. and Boukal, D. S. 2022. From individuals to communities: habitat complexity affects all levels of organization in aquatic environments. - WIREs Water 9: e1575.

St Pierre, J. I. and Kovalenko, K. E. 2014. Effect of habitat complexity attributes on species richness. - Ecosphere 5: art22.

Stenseth, N. Chr. 1980. Spatial heterogeneity and population stability: some evolutionary consequences. - Oikos 35: 165–184.

Terao, O. 1985. Contribution to the study of the ecology of the medaka, *Oryzias latipes*, under natural conditions: life span, reproduction, food habits and its seasonal changes.

Walde, S. J., Nyrop, J. P. and Hardman, J. M. 1992. Dynamics of *Panonychus ulmi* and *Typhlodromus pyri*: factors contributing to persistence. - Exp. Appl. Acarol. 14: 261–291.

Wise, D. H. 2006. Cannibalism, food limitation, intraspecific competition, and the regulation of spider populations. - Annu. Rev. Entomol. 51: 441–465.

Wissinger, S. A., Whiteman, H. H., Denoël, M., Mumford, M. L. and Aubee, C. B. 2010. Consumptive and nonconsumptive effects of cannibalism in fluctuating age-structured populations. - Ecology 91: 549–559.

Wood, S. N. 2017. Generalized additive models: an introduction with R. - Chapman & Hall/CRC.

Woodward, G., Ebenman, B., Emmerson, M., Montoya, J. M., Olesen, J. M., Valido, A. and Warren, P. H. 2005. Body size in ecological networks. - Trends Ecol. Evol. 20: 402–409.

Yamagishi, H. 1976. Experimental study on population dynamics in the guppy, *Poeclia reticulata* (Peters). Effect of shelters on the increase of population density. - J. Fish Biol. 9: 51–65.

Yamamoto, T. 1975. Medaka (killifish): biology and strains. - Keigaku Pub. Co.

Appendix 1. Artificial habitat structure.

Geotextile 2D structure

685   To gain some insights about geotextile's 2D structure, we performed an ImageJ analysis from a top view of a 27 x 42 cm geotextile sample. We found that holes in the mesh represented 38 % of the whole geotextile surface (Fig. S1A). We plotted hole-size distribution, and compared it with the distributions of medaka body width and height, which determine medaka ability to pass through geotextile mesh (Fig. 1B).

690   **Fig. S1. 2D structure of a 27 x 42 cm sample of a single layer of geotextile. A.** Top view of the mesh structure of the geotextile, with holes appearing in black and the geotextile mesh appearing in white. **B. Grey distribution:** distribution of hole size (Feret's diameter) in the geotextile mesh, as measured using ImageJ from Fig. S1A. Feret's diameter, also known as maximum caliper, is the longest distance between any two points along the hole boundary. Hole Feret's diameters ranged from 0.11 to 15.00 mm (mean ± SD = 1.54 ± 1.42 mm). **Red and green distributions:** distributions
695   of medaka body width (red) and height (green) during the 2021 and 2022 mesocosm experiments, as predicted from standard body lengths (Sdl) using allometric relationships estimated by Renneville et al. (2016).

The distributions of medaka body width and height were fully included in hole-size distribution (Fig. S1B), indicating that a floating single-layered geotextile tile could not provide any absolute
700   refuge against cannibalism to small-bodied medaka. However, small-bodied medaka had many more holes available to pass through the geotextile than large-bodied medaka, and we hypothesized that the geotextile should selectively hamper the movements of large-bodied medaka.

Behavioural assays

705   To test this hypothesis, we evaluated the effect of the geotextile on medaka movements in a series

of four behavioural assays performed on July 1st 2024. We used a 55 x 67 cm, single-layered tile of

geotextile affixed onto a wooden frame and placed in a 100 x 100 cm tank filled with 30 cm of

dechlorined tap water (Fig. S2). The frame was put on wedges, such that the geotextile was

maintained 3-4 cm below the water surface, and such that the fish put inside the frame would have

710   to swim through the geotextile to escape to the tank.

Fig. S2. Single-layered geotextile tile affixed onto a wooden frame. A: Bottom view of the 55 x 67 cm wooden frame. **B**. Top view of the wooden frame placed in the tank before the start of behavioural assays.

The medaka used for assays were from a stock of the Kiyosu strain maintained permanently in large

715   outdoor mesocosms with mild artificial feeding. Four batches of 210, 256, 251, and 220 naive fish,

respectively, were gently captured using an aquarium net. With no further habituation, each batch of

fish was deposited in the wooden frame and left undisturbed under room light during 1, 5, 10 or 20

minutes, respectively. At the end of each assay, the frame was removed from the tank and fish were

separated among those that had swum through the geotextile and those that did not. Each individual

720   fish was assayed only once. All fish were photographed and measured for standard body length

(Sdl) using ImageJ as described in the main text, and were returned to their outdoor mesocosm.

We modelled the effect of medaka Sdl (range 8-32 mm, mean $\pm$ SD = 20.4 $\pm$ 4.8 mm) and duration

of assay on medaka probability to swim through the geotextile using a Bernouilli GLM:

$$S_i \sim Bernoulli(p_i)$$
$$\ln\left(\frac{p_i}{1-p_i}\right) = \beta_0 + \beta_1 Sdl_i + \beta_2 D_{batch[i]} + \beta_3 Sdl_i * D_{batch[i]} \qquad \text{(Model A1)},$$

where S_i is the status of individual fish i at the end of the behavioural assay, i.e., $S_i = 1$ if the

fish swam through the geotextile and 0 otherwise, and $D_{batch[i]}$ is the duration of assay associated

730   with the batch to which fish i belonged. The results of maximum-likelihood estimation of the β

parameters using the `glm` function of R are shown in Table A1.

Table A1. Results of maximum-likelihood estimation of parameters from model A1. Bolded-faced p-values are significant at the 5 % risk.

Response	Distribution	Link	Effect	Parameter	Estimate	sd	t-value	p-value
			Intercept	β0	6.303	0.949	6.643	**5.20E-11**
Individual state (passed through geotextile or not)	Bernouilli	Logit	Standard length Sdl	β1	-0.240	0.041	-5.804	**8.86E-09**
			Assay duration D	β2	-0.123	0.086	-1.437	1.51E-01
			Sdl-by-D interaction	β3	0.009	0.004	2.304	**2.14E-02**

Parameter estimation shows that the probability for a medaka to swim through the single-layered

geotextile tile was negatively body-length dependent, and that there was a significant length-by-

duration interaction (Table A1). To further visualize this interaction, we plotted model predictions in

Fig. S3.

Fig. S3. Interaction between medaka standard body length (Sdl) and behavioural-assay duration on medka probability to swim through a single layer of geotextile. The surface shows predictions from Model A1 for the whole range of assayed body sizes. Symbols show the location of fitted values.

Fig. S3 shows that the geotextile strongly reduced passage of the largest-bodied medaka during short-duration assays (1-5 minutes), but almost not during the 20-min assay. This result confirms that a single-layered geotextile tile slowed down, but did not prevent, the movements of large-bodied, potentially cannibalistic medaka.

755   *Multiple-layered geotextile mats.*

During the 2021 pond experiment, we used single-layered geotextile tiles and varied habitat complexity through contrasted tile surfaces. In 2022, we aimed at increasing the complexity contrast, and we varied complexity through contrasting both geotextile surface and the number of geotextile layers. Specifically, we created high-complexity habitats using geotextile mats comprised

760   of five stacked tiles, which structure is shown in Fig. S4.

Fig. S4. Geotextile mat made from a 5-layered stack of geotextile tiles. A. Close view showing the 5 layers of geotextile, separated by a 2 cm x 1 cm black plastic mesh used to prevent tiles to interlock. This mesh was large enough not to restrict fish movements. The whole mat was about 10 cm thick. **B.** Distant view showing mat positioning at the top of the water column. Mat layers were bound together by a pair of plastic collars, here visible on the right-hand side of the mat.

Appendix 2. Strain effects on population growth rate and body sizes.

In the 2021 pond experiment, medaka strains were not mixed in a pond (see Methods), and it was possible to test for strain effects on population dynamics. We tested for a strain effect on population growth rate using a model similar to Model 2 in the main text:

$$
\begin{gathered}
N(t+1)_i \sim \text{Poisson}(\lambda_i) \\
\ln(\lambda_i) = \ln(N(t)_i) + \alpha_S + \beta \, N_{st}(t)_i + \epsilon_i \qquad \text{(Model A2)}, \\
\epsilon_i \sim \text{Normal}(0, \sigma_\epsilon^2)
\end{gathered}
$$

where i indexes ponds (n = 16), α_S is a strain effect on population growth rate, β is the density-dependent parameter, and σ_ϵ^2 is a overdispersion parameter. Model A2 was fitted in the `glmmTMB` library of R (Brooks et al. 2017). We tested for a stain effect using a Chi-squared likelihood ratio test between Model A2 and a similar model that did not include any strain effect using the `anova` function of R. The results suggest no strain effect on population growth rate (Table S2).

Table S2. Results from running the `anova` function in R to compare Model A2 with a similar model that omits a strain effect.

	Df	AIC	BIC	logLik	deviance	Chisq	Chi Df	p-value
Model without any strain effect	3	165.8	168.2	-79.9	159.8			
Model A2	8	167.6	173.8	-75.8	151.6	8.2	5	0.144

We tested for a strain effect on standard body lengths Sdl using a linear mixed-effects model fitted by restricted maximum likelihood in the `nlme` library of R (Pinheiro et al. 2018):

$$
\begin{gathered}
\ln(\text{Sdl}_i) \sim \text{Normal}(\mu_i, \sigma^2) \\
\mu_i = \alpha_S + \xi_{p[i]} \qquad \text{(Model A3)}, \\
\xi_{p[i]} \sim \text{Normal}(0, \sigma_\xi^2)
\end{gathered}
$$

where ln is the natural logarithm, i indexes individual fish ($n = 1917$), p indexes ponds ($n = 16$), and α_S captured strain effects on mean ln-transformed body lengths. The $\xi_{p[i]}$ parameter

captured random pond effects. We then tested significance of the variance explained by α_S in the

model using an F-test in the `anova` function of R. The results show that the strain effect was not

790 statistically significant (Table S3).

Table S3. Result of running the `anova` function in R to test significance of a strain effect in Model A3.

	numDF	denDF	F-value	p-value
(Intercept)	1	1901	5122.4	<.0001
Strain	5	10	1.0	0.451

795 **Appendix 3. Binomial analysis of survival probability of age-1+ medaka through the reproductive period.**

Compared to Model 5 in the main text, an alternative approach to modelling survival probability of age-1+ medaka, which does not propagate estimation uncertainty in age-1+ numbers but preserves

800 the underlying Binomial process, consists in using an overdispersed Binomial model of the form:

$$\widetilde{N}_{1+}(t+1)_i \sim \text{Binomial}\left(N(t)_i, \pi_i\right)$$
$$\ln\left(\frac{\pi_i}{1-\pi_i}\right) = \alpha_{y[i]} + \beta_{k[i]} - \gamma_{k[i]} N_{st}(t)_i + \epsilon_i \qquad \text{(Model A4),}$$
$$\epsilon_i \sim \text{Normal}\left(0, \sigma_\epsilon^2\right)$$

where $\widetilde{N}_{1+}(t+1)$ is the median posterior number of age-1+ fish after their reproductive period,

805 and ϵ is an overdispersion parameter. Other variables and parameters are as in Model 5 in the main text. Model A4 had a Bayesian P-value of 0.49, indicating an excellent fit to the data, and yielded parameter estimates presented in Table S4.

Table S4. Summary statistics for MCMC estimation of the parameters of Model A4. Bolded-faced

810 p-values are significant at the 5 % risk.

Response	Distribution	Link	Effect	Parameter	mean	sd	Rhat	MCMC P-value
Age-1+ Survival probability π	Binomial	Logit	Model intercept (year 2021, Low complexity)	$\alpha 0$	2.386	0.657	1.037	**0.000**
			Year 2022 effect	α	-1.747	1.067	1.022	0.132
			High complexity effect	β	-1.154	0.583	1.017	**0.047**
			$N_{st}(t)$ at Low complexity	$\gamma 0$	0.098	0.039	1.046	**0.026**
			$N_{st}(t)$, High complexity effect	γ	0.002	0.034	1.025	0.946
			Overdispersion	σ_ϵ^2	1.527	0.249	1.005	