Habitat structural complexity increases age-class coexistence and productivity in fish populations

Eric Edeline[1*], Yoann Bennevault[2], David Rozen-Rechels[3]

1: DECOD (Ecosystem Dynamics and Sustainability), INRAE, IFREMER, Institut Agro, Rennes, France.

2: U3E, INRAE, Agrocampus Ouest, Rennes, France.

3: Sorbonne Université, CNRS, IRD, INRAE, Université Paris Est Créteil, Université Paris Cité, Institute of Ecology and Environmental Sciences of Paris (iEES- Paris), Paris, France.

* corresponding author: eric.edeline@inrae.fr

15 **Abstract**

Structurally-complex habitats harbour more taxonomically-diverse and more productive communities, a phenomenon generally ascribed to habitat complexity relaxing the strength of interspecific predation and competition. Here, we challenge this classical, community-centred view by showing that positive habitat complexity-productivity relationships may also emerge from

20 between-age-class, *intra*-specific interactions at a single-population level. In replicated outdoor pond populations of the medaka fish (*Oryzias latipes*), structurally complex habitats provide refuges to newborns and relax the strength of cannibalism, resulting in increased survival of age-0+ individuals, in a 80 % increase in population growth rate, and in dampened negative density-dependence indicating elevated habitat carrying capacity. The resultant higher population density in

25 complex habitats was associated with increased competition for food among age-0+ and age-1+ individuals, as revealed by their smaller and more variable body sizes. Positive habitat complexity-productivity relationships may thus be considered as a generally-emergent property of both size-structured communities and populations, in which a larger body size brings a predation advantage. Our results highlight that anthropogenic habitat simplification drives biodiversity loss not only from

30 community- but also from population-level processes, and hence may further reduce population productivity in the surviving species. Enhancement of habitat structural complexity is therefore a pivotal action for biodiversity improvement, not only in the context of ecosystem management, but also for successful conservation of endangered populations.

35 **Key words:** Apparent competition, Biodiversity, Body size, Cannibalistic behaviour, Food competition, Density-dependent population dynamics, Habitat heterogeneity, Predation window.

INTRODUCTION

40   Anthropogenic changes and biodiversity loss are often congruent with a simplification in the physical structure of habitats. For instance, agriculture transforms complex mature forests into pastures or grain fields that, eventually, may return to secondary forest which are structurally simpler than mature forest (Colorado Zuluaga & Rodewald 2015). In oceans, eutrophication, acidification, bottom trawling and dredging all tend to destroy the complex structure provided by

45   benthic habitats such as coral reefs (National Research Council 2002, Rogers *et al.* 2014). In freshwaters, eutrophication may drive a loss of submerged macrophytes which are key to habitat structuring (Scheffer 2004). This ubiquitous and global simplification of habitat structure plays a central role in anthropogenic biodiversity loss and the disruption of ecosystem functions.

50   Many studies report that structurally-complex habitats favour increased species coexistence and higher numerical abundances and biomass in animal communities. This positive effect is generally ascribed to habitat complexity relaxing the strength of predation and competition, and thus allowing more species to coexist through spatial partitioning of limiting resources (F. E. Smith 1972, Crowder & Cooper 1982, Diehl 1988, Heck & Crowder 1991, Hixon & Menge 1991, Diehl 1992,

55   Janssen *et al.* 2007, Kovalenko *et al.* 2012, Reichstein *et al.* 2013, Rogers *et al.* 2014).

In essence, the emergence of positive habitat complexity-diversity relationships strongly depends on predators being larger-bodied than their prey. In highly-structured and fine-grained habitats, movements are slowed down and encounter rates are reduced (Stenseth 1980, St Pierre &

60   Kovalenko 2014). In particular, larger-bodied individuals selectively suffer from a reduced agility and from a reduced accessibility to microhabitats (Rogers *et al.* 2014). Hence, structurally-complex habitats decrease predator attack rates, resulting in a reduced risk of prey overexploitation and in relaxed apparent competition among prey (Holt 1987).

65 Historically, research on the ecological effects of habitat structural complexity has focused on interspecific interactions with, to our knowledge, a poor consideration for intraspecific interactions. This is despite that many populations are size-structured due to both overlapping age classes and heterogeneous somatic growth rates within age classes. Within such size-structured populations apparent competition by definition is not possible, but large-bodied individuals often dominate in

70 interference competition (Post *et al.* 1999, Le Bourlot *et al.* 2014), and may even cannibalize smaller-bodied conspecifics (Fox 1975, C. Smith & Reay 1991). It is therefore likely that, in size-structured populations just like in size-structured communities, a high habitat structural complexity impedes the dominance of large-bodied individuals.

75 Accordingly, a few studies suggest that habitat structural complexity increases juvenile-adult coexistence, supports larger population sizes, and favours population persistence in laboratory populations of the guppy *Poecilia reticulata* (Yamagishi 1976, Nilsson & Persson 2013) and the mosquitofish *Gambusia affinis* (Benoît et al. 2000). However, given the paucity of studies dealing with habitat complexity in size-structured populations, the generality of positive complexity-

80 productivity relationships at the population level remains questionable. Additionally, the respective contributions of competition and cannibalism to the emergence of complexity-productivity relationships at the population level remain largely unexplored. In this paper, we test the prediction that habitat structural complexity increases population productivity in the Japanese medaka (*Oryzias latipes*), and that the underlying mechanism involves a relaxation in the strengths of both

85 interference competition and cannibalism.

MATERIALS AND METHODS

90 *Medaka fish*

The medaka is an oviparous fish belonging to the group of Beloniformes, a sister group of Cyprinodontiformes which includes killifishes (Kinoshita *et al.* 2009). Medaka naturally inhabit slow-moving fresh- and brackish-waters of South-East Asia. Juveniles and adults have a highly overlapping diet of zooplankton, small benthic invertebrates and filamentous algae (Terao 1985,

95 Edeline *et al.* 2016). Due to its high thermal tolerance and ease of manipulation, the medaka can be used for parallel experiments in the laboratory, where generation time is 2-3 months under optimal light, food and temperature conditions, and in outdoor ponds, where perennial populations maintain themselves during years under natural conditions without any artificial feeding (Bouffet-Halle *et al.* 2021). For these reasons, the medaka is a good model species for studies in genetics, developmental

100 biology, ecology and evolution (Kinoshita *et al.* 2009, Renneville *et al.* 2016, 2020, Diaz Pauli *et al.* 2019, 2020, Evangelista *et al.* 2020a, 2020b, 2021, Le Rouzic *et al.* 2020, Bouffet-Halle *et al.* 2021).

Cannibalistic behavioural assays and predation window

105 Trophic interactions are often size-dependent (Woodward *et al.* 2005), such that the prey-to-predator body-length ratio is constrained to a specific range called the ''predation window'' (Claessen *et al.* 2000). The lower and upper limits of the predation window have far-reaching consequences for population dynamics and individual life histories, respectively (Claessen *et al.* 2002). We performed cannibalistic behavioural assays to determine whether medaka are

110 cannibalistic under no-complexity conditions and, if so, to estimate the limits of their predation window.

Cannibalistic assays were performed with newborns (2-10 days old) and adults under laboratory

conditions at the CEREEP (https://www.cereep.bio.ens.psl.eu/) at 21°C (±1.5 SD) in complexity-

115   free aquaria. Medaka newborns and adults were measured for standard body length (Sdl, from the

tip of the snout to the base of the caudal peduncle, mean ± SD = 5.6 mm ± 0.7 in newborns and 22.8

mm ± 4.2 in adults), and adults were further determined for sex according to their secondary sexual

characters (Yamamoto 1975, Kinoshita *et al.* 2009). The light regime followed the natural

photoperiod at the time of experiment (12[th] and 13[th] March 2013), i.e., dark from 17:00 to 9:00.

Starting at 15:30 on March 12[th] and during 17.5 hours, two newborns were presented to one adult

(72h fasting) in 45 plastic aquaria (0.5 L, no habitat structure) filled with dechlorinated tap water.

The number of surviving larvae in each aquarium were first counted after 2.5 hours and then every

3 hours, allowing us to study the kinetics of medaka cannibalistic behaviour (yielding n = 270

125   observations).

Following Claessen et al. (2002), we estimated the lower (δ), higher (ε) and optimal (φ) victim-to-

cannibal Sdl ratios from behavioural data. Specifically, we used the relationship between the

probability of a cannibalistic attack and Sdl ratio estimated at first census (2.5 hours of cannibalistic

130   assays), when most of the size-dependency in cannibalism was expressed (see Statistical analyses

for the method of estimation and Results). We arbitrarily defined δ as the lower-end ratio at which

cannibalistic probability becomes less than 0.05, ε as the higher-end ratio at which cannibalistic

probability becomes less then 0.05, and φ as the ratio at which cannibalistic probability was

maximal.

Replicated pond populations with varying habitat complexity

We performed two experiments, one in 2021 and one in 2022. In 2021, on 31st March, 12 circular

140 ponds (2m²) were installed at the U3E experimental research unit (https://www6.rennes.inrae.fr/u3e_eng/), filled with dechlorinated tap water, and seeded with 1–2 L of a mixture of benthic detritus (decaying leaves and phytoplankton), benthic organisms and plankton collected in mature, fishless ponds using a kick net (0.3 mm mesh). Benthos and plankton additions were repeated on April 2nd, 9th and 13th 2021. Additionally, on three occasions (April 2nd,

145 9th and 15th) each pond received a solution of KH_2PO_4 amounting to 10 µg L⁻¹ of phosphorus. Enrichment was complemented on April 13th, 19th 27th and June 4th with additions of 1–2 L of an algal mixture (*Chlorella* sp. and *Desmodesmus* sp). All ponds were covered with an insect-proof net (1 mm mesh size). On July 1st 2021, 4 more ponds were installed and submitted to the same seeding and enrichment treatments. Hence, the experiment included 16 ponds in total.

In ponds, we varied habitat structural complexity using single-layered, square patches of Maccaferri MacMat®. This floating polyamide geomaterial has a 3D, irregular mesh structure that provides crevices where small-bodied medaka can hide (Fig. 1a). The complexity structure covered either a low proportion of pond surface (from 3.1 to 9.9 %, mean 8.6 %, hereafter low-complexity

155 treatment, n = 8 ponds) or a large proportion of pond surface (from 36.2 to 50 %, mean 39.4 %, n = 8 ponds, hereafter high-complexity treatment, Fig. 1b). We further added two floating plastic brushes in all ponds to mimic vegetation and ensure that spawning substrates for medaka were non-limiting (Fig. 1b).

Fig. 1. Manipulation of habitat structural complexity in pond medaka populations in 2021. A: Detailed structure of the geomaterial used as artificial refuges to manipulate habitat complexity during the first experiment (2021). The 2 euro-cent coin has a diameter of 22.25mm. The inset shows a juvenile medaka hiding in the artificial refuge. B: A high-complexity pond hosting a $1m^2$ surface (50 % of pond surface) of floating artificial refuges as well as two floating plastic brushes (spawning substrate). NB: blue ropes ensured that the insect-proof net that covered ponds did not touch water surface.

On March 31st 2021 (July 8th for the four extra ponds), from 18 to 60 adult medaka (mean = 37) born in 2020 (hereafter "age-1+") were randomly introduced in ponds. In order to vary the genetic background of the fish populations, we used medaka originating from the Kiyosu population

(Toyohashi, Aichi Prefecture, Japan), as described in Renneville et al. (2020), as well as five other strains corresponding to different locations in Japan and provided by the Japanese National Bioresource Project[1]: Hamochi, Higashidori, Inawashiro, Tokamachi, and Kushima. The six strains were not mixed in a pond, but seeded each in two separate ponds (one low-complexity pond and

175 one high-complexity pond), except the Kiyosu strain that was seeded in 6 ponds (three low-complexity ponds and three high-complexity ponds). Preliminary analyses showed that medaka strains had no significant effect on population growth rate (Chisq = 6.83, DF = 5, p-value = 0.23) or body sizes ($F_{5, 10}$ = 1.26, p-value = 0.35), and we therefore discarded the strain effect from our subsequent analyses.

From November 15^{th} to 17^{th} 2021, i.e., after the reproductive period, all fish from each pond were sampled using hand nets and a seine net, batch-photographed for individual measurement of their standard body length using ImageJ, and released in their pond. These fish (n = 1917) included a mixture of age-1+ adults that were survivors from initial introductions and age-0+ recruits that were

185 born in the ponds.

In 2022, we repeated the same experiment as in 2021, except for two major differences: we mixed medaka strains within a pond, and we amplified the contrast in habitat structural complexity. On March 22^{nd}, the 16 ponds were dried, all fish were collected, and fish, water and sediments were

190 mixed and redistributed among the 16 ponds. Specifically, the medaka strain mixture in each pond was 3, 8, 7, 15, 8 and 25 fish from the Hamochi, Higashidori, Inawashiro, Tokamachi, Kushima and Kiyosu strains, respectively (66 fish per pond). The low-complexity treatment consisted in 0.063 m^2 of single-layered geomaterial (3.1 % of pond surface), while the high-complexity treatment consisted in 0.750 m^2 of five-layered geomaterial (187.5 % of pond surface). Similar to 2021, we

---

1   https://shigen.nig.ac.jp/medaka/

195 added two floating plastic brushes in all ponds to mimic vegetation. Batch photographs for standard

body length measurement using ImageJ were performed on November 8th and 9th 2022 and all fish

were released in their pond.

Table 1. Parameter estimates from Bayesian Models 2-4. The models were fitted using an "effect"
200 parametrization, i.e., intercepts and slopes are evaluated at a reference level (Low complexity and Year-of-experiment 2021), coded with a "0" in the "Parameter column", and effects of High habitat complexity and Year-of experiment 2022 are estimated as deviations from this reference level. MCMC P-values are twice the proportion of the posterior which sign was opposite to the sign of the posterior mode. P-values significant at the 5 % risk are bold-faced. MCMC P-values are not relevant, and thus not supplied, for variance parameters.

Model	Response	Distribution	Link	Effect	Parameter	mean	sd	Rhat	MCMC P-value
2	$\dfrac{N(t+1)}{N(t)}$	Poisson	Log	Model intercept (year 2021, Low complexity)	$\alpha0$	0.636	0.115	1.001	**0.000**
				Year 2022 effect	α	-0.478	0.176	1.001	**0.013**
				High complexity effect	β	0.596	0.094	1.000	**0.000**
				$N\mathrm{st}(t)$ at Low complexity	$\gamma0$	-0.021	0.006	1.001	**0.001**
				$N\mathrm{st}(t)$, High complexity effect	γ	0.015	0.006	1.001	**0.010**
				Overdispersion	σ^2_ε	0.236	0.040	1.001	
3	log(Sdl)	Gaussian	Identity	Intercept on age-0+ (year 2021, Low complexity)	$\alpha0$	2.551	0.046	1.003	**0.000**
				Year 2022 effect on age-0+	$\alpha1$	0.286	0.127	1.000	**0.045**
				High complexity effect on age-0+	$\alpha2$	-0.179	0.041	1.000	**0.000**
				Year 2022-by-High complexity interaction on age-0+	$\alpha3$	-0.147	0.136	1.000	0.260
				Intercept on age-1+ (year 2021, Low complexity)	$\beta0$	3.152	0.024	1.000	**0.000**
				Year 2022 effect on age-1+	$\beta1$	-0.071	0.021	1.000	**0.001**
				High complexity effect on age-1+	$\beta2$	-0.014	0.024	1.000	0.553
				Year 2022-by-High complexity interaction on age-1+	$\beta3$	-0.099	0.031	1.000	**0.005**
				Pond random effect on age-0+	σ^2_γ	0.159	0.034	1.001	
				Pond random effect on age-1+	σ^2_ζ	0.059	0.015	1.001	

205 Table 1 continues on the next page…

Model	Response	Distribution	Link	Effect	Parameter	mean	sd	Rhat	MCMC P-value
				Model intercept (year 2021, Low complexity)	$\alpha 0$	-0.181	0.036	1.000	**0.000**
				Year 2022 effect	α	-0.619	0.034	1.001	**0.000**
4	$\ln(\widetilde{N}_{0+}(t+1))$	Gaussian	Identity	High complexity effect	β	1.324	0.037	1.000	**0.000**
				$N\mathrm{st}(t)$ at Low complexity	$\gamma 0$	-0.049	0.002	1.000	**0.000**
				$N\mathrm{st}(t)$, High complexity effect	γ	0.039	0.002	1.000	**0.000**
				Model intercept (year 2021, Low complexity)	$\alpha 0$	-0.052	0.032	1.000	0.104
				Year 2022 effect	α	-0.230	0.075	1.001	**0.003**
5	$\ln(\widetilde{N}_{1+}(t+1))$	Gaussian	Identity	High complexity effect	β	-0.043	0.045	1.001	0.347
				$N\mathrm{st}(t)$ at Low complexity	$\gamma 0$	0.008	0.004	1.000	**0.039**
				$N\mathrm{st}(t)$, High complexity effect	γ	0.012	0.003	1.001	**0.002**

Statistical analyses

Data from cannibalistic assays in the laboratory were analysed using an overdispersed binomial (logit link) generalized additive model (GAM) of the form:

$$C_{i,T} \sim Bin(2, \pi_{i,T})$$
$$\ln\left(\frac{\pi_{i,T}}{1-\pi_{i,T}}\right) = \beta_{Sex[i]} + f(SdlR_i, T) \quad \text{(Model 1)},$$
$$var(C_{i,T}) = \theta\, 2\, \pi_{i,T}(1-\pi_{i,T})$$

where *C* is the number of larvae eaten, *i* indexes an aquarium (and cannibal individual), *T* indexes the time of census (n = 270 aquarium-by-time observations), π is cannibalism probability,

215 $\beta_{Sex[i]}$ is a sex effect of cannibal individual *i*, *SdlR* is cannibal to victim Sdl ratio (cannibal Sdl / mean of the two victims Sdl), *f* is a tensor product of natural cubic splines with 3 knots, which was parsimonious in terms of smoother wigliness and, at the time, captured the essential features of the cannibalism-size relationship. Finally, θ is a positive parameter accounting for a slight overdispersion in the data. Estimating a different *f* for each sex increased model's generalised cross-

220 validation score, and we thus preferred the simpler model including *Sex* as a fixed effect on the intercept. We fitted Model 1 using quasi-likelihood in the `mgcv` library (`quasibinomial` family)

of the R software version 4.2.1 (Wood 2017, R Core Team 2023). This model explained 24.5 % of the deviance in data.

225   In pond populations, we tested for an effect of habitat structural complexity (Low vs. High) on medaka population dynamics using the Ricker logistic equation (Case 2000):

$$N(t+1) = N(t) e^{r(1 - \frac{N(t)}{K})} \quad \text{(Eq. 1)},$$
$$\ln(N(t+1)) = \ln(N(t)) + r - \gamma N(t)$$

where $N(t+1)$ is total fish number after reproduction (recruitment), $N(t)$ is the number of age-1+ medaka parents initially introduced before reproduction (stock), r is the slope of stock-

230   recruitment relationship at origin and describes the maximum reproductive rate, and $\gamma = r/K$ measures the strength of negative density-dependence in the population. Specifically, we tested for an effect of habitat structural complexity on the density-*in*dependent r parameter, and on the density-dependent γ parameter using a Ricker stock-recruitment model:

235
$$N(t+1)_i \sim \text{Poisson}(\lambda_i)$$
$$\ln(\lambda_i) = \ln(N(t)_i) + \alpha_y + \beta_{k[i]} - \gamma_{k[i]} N_{st}(t)_i + \epsilon_i \quad \text{(Model 2)}$$
$$\epsilon_i \sim N(0, \sigma_\epsilon^2)$$

where *i* indexes ponds (n = 32), *y* indexes the year of experiment (2021 and 2022), α_y is a year-of-experiment effect on the r parameter from Eq. 1, $\beta_{k[i]}$ is habitat-complexity effect (Low vs. High) on the r parameter from Eq. 1, $\gamma_{k[i]}$ is habitat-complexity effect on the γ

240   parameters from Eq. 1, and ϵ_i is an overdispersion parameter. To avoid an intercept-slope correlation when fitting the model, we standardized $N(t)$ to zero mean in the regression (hence the $N_{st}(t)$ notation in Model 2 equation). As a consequence of this standardisation, the α and

β parameters in Model 2 do not estimate the slope of the Ricker stock-recruitment relationship at

origin, but at an average level of the stock.

We assessed the effect of habitat structural complexity on the size distributions of age-1+ medaka

parents and their age-0+ progeny in pond populations using a log-normal mixture model fitted to

standard body lengths Sdl:

$$
\begin{aligned}
\ln(\text{Sdl}_i) &\sim \sum_{j=1}^{J} \sum_{k=1}^{K} \sum_{p=1}^{P} \pi_{j[i],p[i]} N\left(\mu_{j[i],i}, \sigma^2_{j[i],k[i],y[i]}\right) \\
\mu_{0+,i} &= \alpha 1_{y[i]} + \alpha 2_{k[i]} + \alpha 3_{y[i],k[i]} + \xi_{p[i]} \\
\mu_{1+,i} &= \beta 1_{y[i]} + \beta 2_{k[i]} + \beta 3_{y[i],k[i]} + \zeta_{p[i]} \\
\xi_{p[i]} &\sim N\left(0, \sigma^2_y\right) \\
\zeta_{p[i]} &\sim N\left(0, \sigma^2_\zeta\right)
\end{aligned} \qquad \text{(Model 3)},
$$

where i indexes individual fish (n = 3423), j indexes age groups (age 0+ *vs*. age 1+, J = 2),

 k indexes the complexity treatment (low vs. high, K = 2), p indexes ponds (P = 16), and N is

the normal distribution with mean μ and variance σ^2 . $\pi_{j[i],p[i]}$ is the proportion of age j

255 fish in each pond p such that, for each p , $\pi_j \geq 0, \sum_{j=1}^{J} \pi_j = 1$. The $\alpha 1_{y[i]}$ and $\beta 1_{y[i]}$

parameters capture year-of-experiment effects (2021 vs. 2022) on mean body lengths at age-0+ and

age-1+, respectively. The $\alpha 2_{k[i]}$ and $\beta 2_{k[i]}$ parameters capture the effects of habitat structural

complexity (Low vs. High) on mean body lengths at age 0+ and age 1+, respectively. The

 $\alpha 3_{y[i],k[i]}$ and $\beta 3_{y[i],k[i]}$ parameters capture complexity-by-year interactions on mean body

260 lengths at age 0+ and age 1+, respectively. The $\xi_{p[i]}$ and $\zeta_{p[i]}$ parameters capture random pond

effects on mean body lengths at age 0+ and age 1+, respectively. Finally, Model 3 included

heteroscedasticity in the form of a triple age class-by-complexity-by-year interaction on residual variance.

265 Mixture Model 3 provided us with pond-specific estimates for the number of age-0+ *vs*. age-1+ medaka after reproduction at time t+1. We were thus able to estimate the specific effects of habitat structural complexity on the production rate of estimated age-0+ recruits per introduced spawner in a Ricker stock-recruitment model:

270
$$
\begin{aligned}
&\ln\left(\widetilde{N}_{0+}(t+1)_i\right) \sim \mathrm{Normal}\left(\mu_i, \sigma^2_{\widetilde{N}_{0+}(t+1)_i}\right) \\
&\mu_i = \ln\left(N(t)_i\right) + \alpha_y + \beta_{k[i]} - \gamma_{k[i]} N_{st}(t)_i
\end{aligned}
\qquad \text{(Model 4),}
$$

where $\widetilde{N}_{0+}(t+1)$ is the median posterior number of age-0+ fish produced, and $\sigma^2_{\widetilde{N}_{0+}(t+1)_i}$ is the posterior variance in $\widetilde{N}_{0+}(t+1)$, supplied here as data so as to propagate estimation uncertainty. Other variables and parameters are as in Model 2 and have the same biological interpretation.

Following a similar rationale, we modelled the median posterior number of surviving age-1+ fish after reproduction as:

$$
\begin{aligned}
&\ln\left(\widetilde{N}_{1+}(t+1)_i\right) \sim \mathrm{Normal}\left(\mu_i, \sigma^2_{\widetilde{N}_{1+}(t+1)_i}\right) \\
&\mu_i = \ln\left(N(t)_i\right) + \alpha_y + \beta_{k[i]} - \gamma_{k[i]} N_{st}(t)_i
\end{aligned}
\qquad \text{(Model 5),}
$$

where $\widetilde{N}_{1+}(t+1)$ and $\sigma^2_{\widetilde{N}_{1+}(t+1)_i}$ are the median and variance, respectively, in posterior number of age-1+ fish after their reproductive period. Hence, Model 5 propagates estimation uncertainty in the number of surviving age-1+ fish. Intercepts α and β capture the year-of-experiment and habitat-complexity effects on mean, log-transformed survival probabilities of age-1+ fish through

285 the reproductive period, and slopes γ capture the density-by-habitat complexity interaction on

age-1+ survival probabilities. We preferred Model 5 to a Binomial or Beta model, which are more

naturally adapted to modelling probabilities, but can not straightforwardly incorporate estimation

uncertainty $\sigma^2_{\tilde{N}_{1+}(t+1)_i}$. However, as a complement to Model 5, we provide a Binomial analysis of

age-1+ survival probabilities in Appendix 1.

We fitted Models 2-5 using Markov Chain Monte Carlo (MCMC) in JAGS 4.3.0 (Plummer 2003)

through the `jagsUI` package (Kellner 2019). Priors were chosen to be weakly informative except

in mixture Model 3, where we imposed the following constraints: (i) mean standard body length

was smaller at age 0+ than at age 1+ (i.e., $\alpha_1 < \beta_1$ and $\alpha_2 < \beta_2$), (ii) body-length variance was

295 larger at age 0+ than at age 1+, as evidenced by a visual inspection of size distributions (i.e.,

$\sigma^2_\gamma > \sigma^2_\xi$, Fig. 4b), and (iii) the number of age-1+ individuals after reproduction at time $t+1$ was

not larger than before reproduction at time t (i.e., $\pi_{1+} \leq min\left(1, \frac{N(t)}{N(t+1)}\right)$). We further prevented

label switching by assigning age-class 0+ to fish shorter than 10 mm and age-class 1+ to fish longer

than 30 mm (Edeline *et al.* 2016, Bouffet-Halle *et al.* 2021). We ran 3 parallel MCMC chains until

300 parameter convergence was reached, as assessed using the Gelman–Rubin statistic (Gelman &

Rubin 1992).

We assessed goodness of fit of Models 2-5 by using a Bayesian P-value (Gelman *et al.* 1996a).

Briefly, we computed residuals for the actual data as well as for synthetic data simulated from

305 estimated model parameters (i.e., residuals from fitting the model to ''ideal'' data). The Bayesian P-

value is the proportion of simulations in which ideal residuals are larger than true residuals. If the

model fits the data well, the Bayesian P-value is close to 0.5. Bayesian P values for Models 2-5

were equal to 0.49, 0.52, 0.12 and 0.07, respectively, indicating fits ranging from excellent to fair

only (Gelman *et al.* 1996b). The lower fit of Models 4 and 5 is explained by the fact that error

310   variance was not freely estimated but supplied as data in these models.

Finally, the significance of the effects included in our models was assessed using MCMC p-values

(not to be confounded with the Bayesian P-value above), which quantify posterior overlap with

zero. Specifically, at each MCMC iteration and for each model parameter, we computed Δ as the

315   difference between posterior parameters under Low and High habitat complexity. We then

computed MCMC p-values as twice the proportion of Δ for which the sign of Δ was opposite to that

of its mean value.

320   **RESULTS**

Cannibalistic behavioural assays and predation window

The relationship between medaka cannibalistic behaviour and the victim-to-cannibal ratio for

standard body length (Sdl) predicted by GAM Model 1 was nonlinear, and changed during the

course of cannibalistic assays (Fig. 2A). At first census, after 2.5 hours of exposure, the

325   cannibalism-Sdl ratio relationship followed a bell-shaped curve, thus validating the hypotheses of

the predation window (Fig. 2B).

As time of victim to cannibal exposure was increasing, the relationship progressively became body-

size independent. By the end of the assays (17.5 hours of exposure), victim survival probability was

330   very low under almost all Sdl ratios (Fig. 2A). GAM Model 1 further showed that overall mean

cannibalism probability increased significantly from 0.67 in female cannibals to 0.85 in male

cannibals (Fig. 2A; β_{Male} = 1.01, SE = 0.288, t-value = 3.52, p-value < 0.001).

Fig. 2. Cannibalistic behaviour in the laboratory with no habitat structure. A: Surface plot for the nonlinear interaction between a the victim-to-cannibals body length ratio and time of exposure on the probability for cannibalism to occur. Blue solid lines: male cannibals; Pink dashed lines: female cannibals. The surfaces were predicted from model (1). Dots show the fitted values: Circles: male cannibals; Triangles: female cannibals. Sdl: standard body length. **B:** Defining parameters of the predation window from predicted cannibalism probability surfaces. The cannibalistic curves are the same as in Fig. 2A for the fist census (2.5 hours of cannibalistic assays). δ and ε are the lower and upper limits of the predation window, arbitrarily defined as the lower- and upper-end ratios, respectively, at which cannibalistic probability becomes less than 0.05 (horizontal dashed line). φ is the optimal victim-to-cannibal Sdl ratio. Note that the higher cannibalistic voracity in males results in a wider predation window.

Due to this sex effect on cannibalistic voracity, the upper and lower limits of the predation window were sex-dependent (Fig. 2B). Specifically, predation window was slightly wider in male (δ = 0.09, φ = 0.26, ε = 0.41) than in female cannibals (δ = 0.13, φ = 0.26, ε = 0.38). These parameters show

that newly-hatched medaka larvae are under strong cannibalistic risk (Fig. 3). An average-sized

hatchling (3.8 mm Sdl) may be cannibalised by any 9.3-to-42.2 mm Sdl male (10.0 to 29.2 mm

female) conspecific. This body-size range encompasses not only the whole range of parental body

350   sizes, but also late-juvenile body sizes (Fig. 3), indicating that medaka larvae are exposed to both

inter- and intra-cohort cannibalism.

Fig. 3. Predation window for cannibal *vs*. victim length in medaka. The black solid line shows the

optimal victim length while ribbons show the upper and lower predation limits for males (blue solid lines)

355   and females (pink dashed lines), as defined by the 5 % cannibalism probabilities displayed in Fig. 2B. The

top marginal distribution shows the body sizes typically present in pond mesocosm medaka populations in

France before reproduction in March (Bouffet-Halle *et al.* 2021), i.e., the size distribution of potential

cannibals. The right-hand marginal distribution shows larval body sizes at hatch (Renneville *et al.* 2020), i.e.,

the size distribution of potential victims. Sdl: standard body length. Vertical and horizontal dashed lines show

360   the 95 % body-size limits of potential cannibals and larval victims, respectively.

Replicated pond populations

In replicated pond populations, complex habitats increased the total number of fish present in ponds

after the reproductive period $N(t+1)$ (Table 1-Model 2, positive β effect), as indicated by green

365 lines being higher than red lines in Fig. 4A. Specifically, population growth rate $N(t+1)/N(t)$

increased on average from 1.5 year^{-1} under a low complexity to 2.7 year^{-1} under a high complexity

(80 % increase).

Additionally, medaka population dynamics were negatively density-dependent under a low habitat

370 complexity (Table 1-Model 2, significantly negative γ_0 effect), but a high habitat complexity

dampened this negative density dependence (Table 1-Model 2, positive γ effect). Accordingly,

the stock-recruitment relationship changes from flat at low habitat complexity (red lines) to positive

at high habitat complexity (green lines), reflecting increased habitat carrying capacity (Fig. 4A).

Fitting Model 2 without centring $N(t)$, so as to preserve the original biological interpretation of

375 r (see Methods), yielded a median posterior for K that increased under High habitat complexity

from 82 to 194 fish in 2021, and from 59 to 143 fish in 2022.

Mixture Model 3 allowed us to quantify the effects of habitat structural complexity on age-specific

log-body-length distributions. Increased habitat complexity decreased mean body length in age-0+

380 medaka in both years of experiment (Table 1-Model 3, α_2 and α_3 parameters), and decreased

mean body length in age-1+ medaka in 2022, when the complexity contrast was more extreme, but

not in 2021 (Table 1-Model 3, β_3 *vs.* β_2 parameters).

In 2021 more complex habitats increased variability of log-transformed body-length in age-0+ fish

385 (Model 3, $\sigma_{\text{age-0+, Low-complexity, 2021}} = 0.269$ *vs.* $\sigma_{\text{age-0+, High-complexity, 2021}} = 0.303$, MCMC P-value =

0.022), but decreased it in age-1+ fish (Model 3, $\sigma_{\text{age-1+, Low-complexity, 2021}} = 0.140$ *vs.*

$\sigma_{\text{age-1+, High-complexity, 2021}} = 0.113$, MCMC P-value = 0.038). In 2022, age-0+ were to few in low-

complexity ponds to provide any robust inference on body-length variability (MCMC P-value =

0.652), while more complex habitats increased body-length variability in age-1+ fish (Model 3,

390 $\sigma_{\text{age-1+, Low-complexity, 2022}} = 0.102$ *vs.* $\sigma_{\text{age-1+, High-complexity, 2022}} = 0.136$, MCMC P-value = 0).

Model 2 is a model of unstructured population dynamics, where age-0+ and age-1+ individuals are

lumped in $N(t+1)$. Hence, the increase in $N(t+1)/N(t)$ under a high habitat complexity

may reflect increased *per capita* production of age-0+ fish and/or increased survival probability of

395 age-1+ fish through the reproductive period. Mixture Model 3 provided us with separate estimates

for the numbers of age-0+ and age-1+ fish at *t*+1, and thus allowed us to separate these two effects

using Ricker Models 4 and and 5 (see Methods).

Ricker Model 4, which was fitted to the number of age-0+ medaka, perfectly parallel results from

400 Model 2 above. Specifically, *per capita* production of age-0+ fish strongly increased from on

average 0.8 at low complexity to 3.1 under a high habitat complexity (Table 1-Model 4, positive β

effect). Additionally, the habitat complexity-by-density interaction reveals a strong negative density

dependence under a low habitat complexity (Table 1-Model 4, negative γ_0 effect) that was

dampened under a high habitat complexity (Table 1-Model 4, positive γ effect).

Patterns of age-1+ survival probability were different from patterns of age-0+ production. A high

habitat complexity *decreased* age-1+ survival probability, but significantly-so in the binomial model

only (Table 1-Model 5 *vs*. Appendix 1: negative β effects), suggesting a relatively weak effect. Specifically, the binomial model predicts that age-1+ survival probability decreased from 0.91 at

410 low complexity to 0.77 under a high habitat complexity (Appendix 1).

Fig. 4. Effect of habitat structural complexity on medaka population dynamics. A. Relationship between the number of medaka after ($N(t$+1)) and before reproduction ($N(t)$) in replicated pond populations. Dots show the raw data (squares: Low complexity; diamonds: High complexity), and curves show median

415 posterior predictions from of a Ricker Model 2 (violet-red: Low-complexity habitats; Green: High-complexity habitats). Ribbons show 95 % credible intervals. The straight y = x black line intersects coloured curves at habitat carrying capacity K. **B.** Body-length distributions after reproduction at t+1. Bars show the raw data and curves show median posterior distributions predicted from the log-Gaussian mixture Model 3. There are 8 curves, one per age-class per complexity treatment per year of experiment.

Age-1+ survival probability was *positively* density-dependent under low habitat complexity in both the Gaussian and binomial models (Table 1-Model 5 and Appendix 1: positive γ_0 effects), and a high habitat complexity amplified this positive density dependence, but significantly-so in the

Gaussian model only (Table 1-Model 5 *vs*. Appendix 1: positive γ effects), suggesting a weak

425 amplification.

Taken together, results from Models 4 and 5 indicate that the positive effect of habitat complexity on medaka population growth rate resulted from increased *per capita* production of age-0+ fish, and was weakly opposed by depressed survival of age-1+fish.

DISCUSSION

In multispecies assemblages, habitat structural complexity increases species coexistence and productivity through a relaxation in the strengths of interspecific competition and predation (F. E.

435 Smith 1972, Crowder & Cooper 1982, Holt 1987, 1987, Diehl 1988, 1992, Heck & Crowder 1991, Hixon & Menge 1991, Rogers *et al.* 2014). Our present results in medaka fish show that habitat structural complexity may further increase age-class coexistence and productivity at a single-population level. Hence, intraspecific processes may provide a substantial, but currently neglected, contribution to the positive effects of habitat complexity on biodiversity and productivity, which are

440 almost always measured at the community level.

Our results may further be used to gain some understanding of the underlying ecological mechanisms that drive positive habitat complexity-productivity relationships. We show that increased age-class coexistence and productivity in medaka populations was achieved through an

445 increased *per capita* production of age-0+ recruits by their age-1+ parents which, in theory, may result either from increased parental fecundity (i.e., egg production) and/or from increased survival

of newborns to the recruit stage in November. Our results, however, provide evidence against increased adult fecundity in more complex habitats.

450　In more complex habitats, both age-0+ and age-1+ medaka had smaller and more variable body sizes, indicating increasing competition for food (Ohlberger *et al.* 2013). Additionally, age-1+ survival probability decreased in more complex habitats, consistent with reinforced exploitative competition for food, in which large-bodied medaka parents are excluded by their progeny (Edeline *et al.* 2016). Increased food intake is a necessary condition for increased reproductive output

455　(Kooijman 2010). Hence, in a context of food shortage in more complex habitats, increased fecundity in age-1+ medaka was very unlikely. We therefore conclude that increased age-class coexistence and productivity of medaka populations in more complex habitats resulted primarily from increased survival of newborns.

460　Increased newborn survival resulted from a relaxed dominance of their larger-bodied parents, as evidenced by relaxed negative density-dependence of the stock-recruitment relationship in more complex pond habitats. In animal populations, dominance of large-bodied individuals typically operates through interference competition and/or cannibalism (Post *et al.* 1999, Claessen *et al.* 2004, Le Bourlot *et al.* 2014). In medaka, interference with a dominance of large-bodied individuals

465　was previously documented in complexity-free aquariums in the laboratory (Magnuson 1962). Our cannibalistic assays now further demonstrate that, in structurally-simple habitats, newborn medaka are exposed to cannibalism from a large body-size range of both age-0+ and age-1+ individuals. Hence, in principle structurally-complex habitats may potentially relax both interference competition and cannibalism in medaka populations.

However, our results in outdoor ponds show that competition for food experienced by both age-0+ and age-1+ medaka was stronger, not weaker, in more complex habitats (see above). This result is inconsistent with a relaxation of interference competition from age-1+ on age-0+ individuals, and rather highlights a general increase in exploitative competition for food in more complex ponds.

475 Hence, we safely conclude that a relaxed cannibalism was the primary driver of increased newborn survival in more complex ponds.

This conclusion is in perfect line with both experimental work (Dionne 1985) and theoretical models (Claessen *et al.* 2000) showing that a relaxed cannibalism increases the strength of

480 intraspecific competition for food. The strong positive density-dependence in age-1+ survival probability in pond medaka populations may thus be explained by cannibalism which, through relaxing competition for food in which age-0+ dominate (see above and below), allowed age-1+ medaka to indirectly limit their competitive exclusion and to increase their own survival probability. Size-dependent cannibalism is a very common form of interaction in the animal kingdom (Fox

485 1975, Polis 1981, C. Smith & Reay 1991), and we thus expect habitat complexity to enhance age-class coexistence and population growth rate across a broad taxonomic range.

While in experimental ponds in France cannibalism allow medaka to survive to age 2+ (this study, Bouffet-Halle et al. 2021), in the wild in Japan juvenile medaka exclude their age-1+ parents in

490 exploitative competition for food (Terao 1985, Edeline *et al.* 2016). Two separate and non mutually-exclusive mechanisms may explain the dominance of age-0+ medaka in the wild. First, wild habitats are probably more structurally complex than the most complex ponds in our experiments, and may effectively provide age-0+ medaka with absolute refuges against cannibalism. In particular, wild habitats often harbour shallow areas, which are key to provide juvenile fish with

495   absolute predation refuges (Radinger *et al.* 2023), but which were absent from the most complex

ponds in our experiments.

Second, in Japan summer temperatures are often warmer than in France. Yet, net energy gain by

individual ectotherms (energy intake rate minus maintenance rate) increases faster with temperature

500   for small consumers, which thus have a competitive advantage under warm conditions (Vasseur &

McCann 2005, Ohlberger *et al.* 2011, Edeline *et al.* 2013). Hence, warmer temperatures favour the

competitive exclusion of age-1+ medaka by their age-0+ progeny (Edeline *et al.* 2016).

If the higher complexity of wild habitats explains the dominance of age-0+ medaka in the wild, then

505   we could expect a hump-shaped habitat complexity-productivity relationship in medaka

populations, with a shift along an increasing habitat-complexity gradient from a cannibal-mediated,

age-1+ dominance at low complexity to an exploitative competition-mediated, age-0+ dominance at

high complexity, resulting in maximal age-class coexistence and population productivity at

intermediate complexity levels where the asymmetry in dominance is most relaxed. Maximal

510   diversity/productivity at intermediate levels of habitat structural complexity may also be observed

in complex, multispecies communities (Crowder & Cooper 1982, Rogers *et al.* 2018), supporting

the contention that complexity-diversity/productivity relationships measured at the community level

may reflect processes effectively acting at the population level.

515   Around the globe, habitat structural complexity in natural habitats is often provided by sessile

organisms such as plants or corals. These "ecosystem engineers" become increasingly threatened by

anthropogenic perturbations such as deforestation, eutrophication, bottom trawling and dredging, or

by marine trophic cascades or ocean acidification, thus making biodiversity loss convergent with a

global habitat simplification. Community theory predicts that such habitat simplification may entail

520   a cascade of species extinctions by virtue of the positive habitat complexity-diversity relationship. Our study further suggests that, within the surviving species, increased dominance of large-bodied individuals may further reduce juvenile survival, and ultimately impair population productivity and long-term persistence. These results call for an incorporation of strong actions to preserve or restore habitat structural complexity, not only into ecosystem-based management programs, but also in

525   conservation plans for endangered populations.

Acknowledgement. We are grateful to the NBRP medaka (https://shigen.nig.ac.jp/medaka/) and to Prof. Kiyoshi Naruse (NIBB Okazaki) for providing us with the medaka strains. We thank André

530   De Roos for insightful comments on an earlier manuscript version. Alexandre Kempf, Solène Moulin, Louisiane Perrin and Morgan Verdeil provided invaluable help during behavioural assays. This experiment was approved by the Charles Darwin Ethical Committee (Ce5/2010/041).

Author contributions: EE designed and contributed to performing experiments, analysed data and

535   wrote the first draft version. YB and DRR performed experiments. All authors contributed to result interpretation and draft improvements.

Funding: This work has benefited from technical and human resources provided by CEREEP-Ecotron IleDeFrance (CNRS/ENS UMS 3194) as well as from financial support from the Regional

540   Council of Ile-de-France under the DIM Program R2DS bearing the references I-05-098/R and 2015-1657. It has received a support under the program 'Investissements d'Avenir' launched by the French government and implemented by ANR with the references ANR-10-EQPX-13-01 Planaqua and ANR-11-INBS-0001 AnaEE France, and from Pépinière interdisciplinaire CNRS de site PSL

(Paris-Sciences et Lettres) "Eco-Evo-Devo". EE was further supported by grants from Sorbonne

545   Université (program Convergences, project C14234) and from Rennes Métropole (AIS 18C0356).

Data availability statement

The data and model codes will be made available online upon acceptance of the manuscript.

550   **References**

Benoît, H. P., Post, J. R., Barbet, A. D., & Ross, S. T. 2000. Recruitment dynamics and size
     structure in experimental populations of the mosquitofish, *Gambusia affinis*. Copeia,
     2000(1), 216–221. DOI: 10.1643/0045-8511(2000)2000[0216:RDASSI]2.0.CO;2

Bouffet-Halle, A., Mériguet, J., Carmignac, D., Agostini, S., Millot, A., Perret, S., Motard, E.,
     Decenciere, B., & Edeline, E. 2021. Density-dependent selection mediates harvest-induced
     trait changes. Ecology Letters, 24(4), 648–657. DOI: 10.1111/ele.13677

Case, T. J. 2000. An illustrated guide to theoretical ecology. 1 ed. New York: Oxford University
     Press.

Claessen, D., de Roos, A. M., & Persson, L. 2000. Dwarfs and giants: cannibalism and competition
     in size-structured populations. The American Naturalist, 155(2), 219–237. DOI:
     10.1086/303315

Claessen, D., de Roos, A. M., & Persson, L. 2004. Population dynamic theory of size-dependent
     cannibalism. Proceedings of the Royal Society of London. Series B: Biological Sciences,
     271(1537), 333. DOI: 10.1098/rspb.2003.2555

Claessen, D., van Oss, C., de Roos, A. M., & Persson, L. 2002. The impact of size-dependent
     predation on population dynamics and individual life history. Ecology, 83(6), 1660–1675.
     DOI: 10.2307/3071986

Colorado Zuluaga, G. J., & Rodewald, A. D. 2015. Response of mixed-species flocks to habitat
     alteration and deforestation in the Andes. Special Issue: Ecology and Conservation of Avian

Insectivores of the Rainforest Understory: A Pan-Tropical Perspective, 188, 72–81. DOI: 10.1016/j.biocon.2015.02.008

Crowder, L. B., & Cooper, W. E. 1982. Habitat structural complexity and the interaction between bluegills and their prey. Ecology, 63(6), 1802–1813. DOI: 10.2307/1940122

Diaz Pauli, B., Edeline, E., & Evangelista, C. 2020. Ecosystem consequences of multi-trait response to environmental changes in Japanese medaka, *Oryzias latipes*. Conservation Physiology, 8(1), coaa011. DOI: 10.1093/conphys/coaa011

Diaz Pauli, B., Garric, S., Evangelista, C., Vøllestad, L. A., & Edeline, E. 2019. Selection for small body size favours contrasting sex-specific life histories, boldness and feeding in medaka, *Oryzias latipes*. BMC Evolutionary Biology, 19(1), 127. DOI: 10.1186/s12862-019-1460-x

Diehl, S. 1988. Foraging efficiency of three freshwater fishes: effects of structural complexity and light. Oikos, 53(2), 207–214. DOI: 10.2307/3566064

Diehl, S. 1992. Fish predation and benthic community structure: the role of omnivory and habitat complexity. Ecology, 73(5), 1646–1661. DOI: 10.2307/1940017

Dionne, M. 1985. Cannibalism, food availability, and reproduction in the mosquito fish (*Gambusia affinis*): a laboratory experiment. The American Naturalist, 126(1), 16–23. DOI: 10.1086/284392

Edeline, E., Lacroix, G., Delire, C., Poulet, N., & Legendre, S. 2013. Ecological emergence of thermal clines in body size. Global Change Biology, 19(10), 3062–3068. DOI: 10.1111/gcb.12299

Edeline, E., Terao, O., & Naruse, K. 2016. Empirical evidence for competition-driven semelparity in wild medaka. Population Ecology, 58(3), 371–383. DOI: 10.1007/s10144-016-0551-4

Evangelista, C., Diaz Pauli, B., Vøllestad, L. A., & Edeline, E. 2020a. Stoichiometric consequences of size-selective mortality: An experimental test using the Japanese medaka (*Oryzias latipes*). Science of The Total Environment, 724, 138193. DOI: 10.1016/j.scitotenv.2020.138193

Evangelista, C., Dupeu, J., Sandkjenn, J., Pauli, B. D., Herland, A., Meriguet, J., Vøllestad, L. A., & Edeline, E. 2021. Ecological ramifications of adaptation to size-selective mortality. Royal Society Open Science, 8(10), 210842. DOI: 10.1098/rsos.210842

Evangelista, C., Vøllestad, L. A., Diaz Pauli, B., & Edeline, E. 2020b. Density-dependent consequences of size-selective induced life-history changes to population fitness in medaka (*Oryzias latipes*). Canadian Journal of Fisheries and Aquatic Sciences, 77, 1741–1748. DOI: 10.1139/cjfas-2019-0406

Fox, L. R. 1975. Cannibalism in natural populations. Annual Review of Ecology and Systematics, 6(1), 87–106. DOI: 10.1146/annurev.es.06.110175.000511

Gelman, A., Meng, X. L., & Stern, H. 1996a. Posterior predictive assessment of model fitness via realized discrepancies. Statistica Sinica, 6(4), 733–807.

Gelman, A., Meng, X. L., & Stern, H. 1996b. Posterior predictive assessment of model fitness via realized discrepancies. Statistica Sinica, 6(4), 733–807.

Gelman, A., & Rubin, D. B. 1992. Inference from iterative simulation using multiple sequences. Statist. Sci., (4), 457–472. DOI: 10.1214/ss/1177011136

Heck, K. L., & Crowder, L. B. 1991. Habitat structure and predator—prey interactions in vegetated aquatic systems. In: S. S. Bell, E. D. McCoy, & H. R. Mushinsky (Eds.), Habitat structure: the physical arrangement of objects in space. pp. 281–299. Dordrecht: Springer Netherlands.

Hixon, M. A., & Menge, B. A. 1991. Species diversity: prey refuges modify the interactive effects of predation and competition. Theoretical Population Biology, 39(2), 178–200. DOI: 10.1016/0040-5809(91)90035-E

Holt, R. D. 1987. Prey communities in patchy environments. Oikos, 50(3), 276–290. DOI: 10.2307/3565488

Janssen, A., Sabelis, M. W., Magalhaes, S., Montserrat, M., & van der Hammen, T. 2007. Habitat structure affects intraguild predation. Ecology, 88(11), 2713–2719.

Kellner, K. 2019. jagsUI: a wrapper around "rjags" to streamline "JAGS" analyses.

Kinoshita, M., Murata, K., Naruse, K., & Tanaka, M. 2009. Medaka. Biology, management and experimental protocols. 1 ed. Ames (USA): Wiley: p. 444.

Kooijman, S. A. L. M. 2010. Dynamic energy budget theory for metabolic organisation. 3 ed. Cambridge: Cambridge University Press.

Kovalenko, K. E., Thomaz, S. M., & Warfe, D. M. 2012. Habitat complexity: approaches and future directions. Hydrobiologia, 685(1), 1–17. DOI: 10.1007/s10750-011-0974-z

Le Bourlot, V., Tully, T., & Claessen, D. 2014. Interference versus exploitative competition in the regulation of size-structured populations. The American Naturalist, 184(5), 609–623. DOI: 10.1086/678083

Le Rouzic, A., Renneville, C., Millot, A., Agostini, S., Carmignac, D., & Edeline, E. 2020. Unidirectional response to bidirectional selection on body size. II Quantitative genetics. Ecology and Evolution, 10(20), 11453–11466. DOI: 10.1002/ece3.6783

Magnuson, J. J. 1962. An analysis of aggressive behavior, growth, and competition for food and space in medaka (Oryzias latipes (Pisces, Cyprinodontidae)). Canadian Journal of Zoology, 40(2), 313–363. DOI: 10.1139/z62-029

National Research Council. 2002. Effects of trawling and dredging on seafloor habitat. Washington, DC: The National Academies Press. DOI: 10.17226/10323

Nilsson, K. A., & Persson, L. 2013. Refuge availability and within-species differences in cannibalism determine population variability and dynamics. Ecosphere, 4(8), art100. DOI: 10.1890/ES13-00105.1

Ohlberger, J., Edeline, E., Vøllestad, L. A., Stenseth, N. C., & Claessen, D. 2011. Temperature-driven regime shifts in the dynamics of size-structured populations. The American Naturalist, 177(2), 211–223. DOI: 10.1086/657925

Ohlberger, J., Otero, J., Edeline, E., Winfield, I. J., Stenseth, N. Chr., & Vøllestad, L. A. 2013. Biotic and abiotic effects on cohort size distributions in fish. Oikos, 122(6), 835–844. DOI: 10.1111/j.1600-0706.2012.19858.x

Plummer, M. 2003. JAGS: a program for analysis of Bayesian graphical models using Gibbs sampling. Vienna, Austria.

Polis, G. A. 1981. The evolution and dynamics of intraspecific predation. Annual Review of Ecology and Systematics, 12, 225–251.

Post, J. R., Parkinson, E. A., & Johnston, N. T. 1999. Density-dependent processes in structured fish populations: interaction strengths in whole-lake experiments. Ecological Monographs, 69(2), 155–175. DOI: 10.1890/0012-9615(1999)069[0155:DDPISF]2.0.CO;2

R Core Team. 2023. R: a language and environment for statistical computing. Vienna, Austria: R Foundation for Statistical Computing.

Radinger, J., Matern, S., Klefoth, T., Wolter, C., Feldhege, F., Monk, C. T., & Arlinghaus, R. 2023. Ecosystem-based management outperforms species-focused stocking for enhancing fish populations. Science, 379(6635), 946–951. DOI: 10.1126/science.adf0895

Reichstein, B., Schröder, A., Persson, L., & De Roos, A. M. 2013. Habitat complexity does not promote coexistence in a size-structured intraguild predation system. Journal of Animal Ecology, 82(1), 55–63. DOI: 10.1111/j.1365-2656.2012.02032.x

Renneville, C., Le Rouzic, A., Baylac, M., Millot, A., Loisel, S., & Edeline, E. 2016. Morphological drivers of trophic cascades. Oikos, 125(8), 1193–1202. DOI: 10.1111/oik.02877

Renneville, C., Millot, A., Agostini, S., Carmignac, D., Maugars, G., Dufour, S., Le Rouzic, A., & Edeline, E. 2020. Unidirectional response to bidirectional selection on body size. I. Phenotypic, life-history, and endocrine responses. Ecology and Evolution, 10(19), 10571–10592. DOI: 10.1002/ece3.6713

Rogers, A., Blanchard, J. L., & Mumby, P. J. 2014. Vulnerability of coral reef fisheries to a loss of structural complexity. Current Biology, 24(9), 1000–1005. DOI: 10.1016/j.cub.2014.03.026

Rogers, A., Blanchard, J. L., & Mumby, P. J. 2018. Fisheries productivity under progressive coral reef degradation. Journal of Applied Ecology, 55(3), 1041–1049. DOI: 10.1111/1365-2664.13051

Scheffer, M. 2004. Ecology of shallow lakes. 1 ed. New York: Springer: p. 357.

Smith, C., & Reay, P. 1991. Cannibalism in teleost fish. Reviews in Fish Biology and Fisheries, 1(1), 41–64. DOI: 10.1007/BF00042661

Smith, F. E. 1972. Spatial heterogeneity, stability and diversity in ecosystems. Transactions of the Connecticut Academy of Arts and Sciences, 44, 309–335.

St Pierre, J. I., & Kovalenko, K. E. 2014. Effect of habitat complexity attributes on species richness. Ecosphere, 5(2), art22. DOI: 10.1890/ES13-00323.1

Stenseth, N. Chr. 1980. Spatial heterogeneity and population stability: some evolutionary consequences. Oikos, 35(2), 165–184. DOI: 10.2307/3544425

Terao, O. 1985. Contribution to the study of the ecology of the medaka, *Oryzias latipes*, under natural conditions: life span, reproduction, food habits and its seasonal changes. Master Thesis Tokyo. p. 63.

Vasseur, D. A., & McCann, K. S. 2005. A mechanistic approach for modeling temperature-dependent consumer-resource dynamics. The American Naturalist, 166(2), 184–198.

Wood, S. N. 2017. Generalized additive models: an introduction with R. 2 ed. Boca Raton: Chapman & Hall/CRC: p. 410.

Woodward, G., Ebenman, B., Emmerson, M., Montoya, J. M., Olesen, J. M., Valido, A., & Warren, P. H. 2005. Body size in ecological networks. Trends in Ecology & Evolution, 20(7), 402–409. DOI: 10.1016/j.tree.2005.04.005

Yamagishi, H. 1976. Experimental study on population dynamics in the guppy, Poeclia reticulata (Peters). Effect of shelters on the increase of population density. Journal of Fish Biology, 9(1), 51–65. DOI: 10.1111/j.1095-8649.1976.tb04661.x

Yamamoto, T. 1975. Medaka (killifish): biology and strains. 1 ed. Tokyo: Keigaku Pub. Co: p. 365.

Appendix 1. Binomial analysis of survival probability of age-1+ medaka through the reproductive period.

555 An alternative approach to modelling survival probability of age-1+ medaka, which does not propagate estimation uncertainty in age-1+ numbers but preserves the underlying Binomial process, consists in using an overdispersed Binomial model of the form:

$$\widetilde{N}_{1+}(t+1)_i \sim Bin\left(N_1(t)_i, \pi_i\right)$$
$$logit(\pi_i) = \alpha_{y[i]} + \beta_{k[i]} - \gamma_{k[i]} N_{st}(t)_i + \epsilon_i \qquad ,$$
$$\epsilon_i \sim N\left(0, \sigma_\epsilon^2\right)$$

where $\widetilde{N}_{1+}(t+1)$ is the median posterior number of age-1+ fish after their reproductive period, and ϵ is an overdispersion parameter. Other variables and parameters are as in Model 5 in the main text. This model has a Bayesian P-value of 0.49, indicating an excellent fit to the data, and yields the following results:

Response	Distribution	Link	Effect	Parameter	mean	sd	Rhat	MCMC P-value
Age-1+ Survival probability π	Binomial	Logit	Model intercept (year 2021, Low complexity)	$\alpha0$	2.386	0.657	1.037	**0.000**
			Year 2022 effect	α	-1.747	1.067	1.022	0.132
			High complexity effect	β	-1.154	0.583	1.017	**0.047**
			Nst(t) at Low complexity	$\gamma0$	0.098	0.039	1.046	**0.026**
			Nst(t), High complexity effect	γ	0.002	0.034	1.025	0.946
			Overdispersion	σ_ϵ^2	1.527	0.249	1.005	