1 **A novel machine learning algorithm picks proteome signature to specifically identify**
2 **cancer exosomes**
3

Bingrui Li[1], Fernanda G. Kugeratski[1], Raghu Kalluri[1, 2, 3]

[1]Department of Cancer Biology, University of Texas MD Anderson Cancer Center, Houston, TX

[2]Department of Molecular and Cellular Biology, Baylor College of Medicine, Houston, TX

[3]Department of Bioengineering, Rice University, Houston, TX

Corresponding author:

Raghu Kalluri, MD, PhD

Email: rkalluri@mdanderson.org
12
13

Abstract

Non-invasive early cancer diagnosis remains challenging due to the low sensitivity and

specificity of current diagnostic approaches. Exosomes are membrane-bound nanovesicles

secreted by all cells that contain DNA, RNA, and proteins that are representative of the parent

cells. This property, along with the abundance of exosomes in biological fluids makes them

compelling candidates as biomarkers. However, a rapid and flexible exosome-based diagnostic

method to distinguish human cancers across cancer types in diverse biological fluids is yet to be

defined. Here, we describe a novel machine learning-based computational method to distinguish

cancers using a panel of proteins associated with exosomes. Employing datasets of exosome

proteins from human cell lines, tissue, plasma, serum and urine samples from a variety of cancers,

we identify Clathrin Heavy Chain (CLTC), Ezrin, (EZR), Talin-1 (TLN1), Adenylyl cyclase-

associated protein 1 (CAP1) and Moesin (MSN) as highly abundant universal biomarkers for

exosomes and define three panels of pan-cancer exosome proteins that distinguish cancer

exosomes from other exosomes and aid in classifying cancer subtypes employing random forest

models. All the models using proteins from plasma, serum, or urine-derived exosomes yield

AUROC scores higher than 0.91 and demonstrate superior performance compared to Support

Vector Machine, K Nearest Neighbor Classifier and Gaussian Naive Bayes. This study provides

a reliable protein biomarker signature associated with cancer exosomes with scalable machine

learning capability for a sensitive and specific non-invasive method of cancer diagnosis.

Introduction

Tissue biopsies have traditionally been a definitive way to diagnose and stage cancer, however, a biopsy may not be easily accessible for many tumors such as those in the pancreas, lung and brain[1-5]. In addition, the small amount of biopsied tissue does not represent the entire heterogeneous pathological profile of the tumor[6]. In recent years, liquid biopsy has emerged as a plausible diagnostic and monitoring approach with the capability to detect tumor biomarkers in more accessible biological fluids such as plasma, serum and urine[7]. Detectable tumor biomarkers can include circulating tumor DNA (ctDNA), circulating tumor cells (CTCs) and exosomes[8].

Exosomes are extracellular vesicles of endosomal origin that are between ~40-180 nm in diameter and have been shown to mediate intercellular communication in health and disease[8-10]. They can contain a variety of biomolecules including DNA, RNA, proteins, lipids, metabolites and other materials representative of the parent cell[8]. Exosomes are present at high concentrations in biological fluids, which is a potential advantage as a biomarker[11, 12]. Exosomal mRNAs and miRNAs have been investigated intensively as diagnostic biomarkers, and mounting evidence suggests that exosomal proteins circulating in biological fluids could be used for cancer diagnosis and monitoring cancer progression[7, 13]. Challenges that remain include standardization of methods for consistent exosome isolation from various tissues, identification of biomarkers that distinguish cancer and normal exosomes across different cancer types, and the identification of biomarkers that are unique to specific biological fluids (e.g., plasma, serum, urine, etc.).

In addition to accurately identifying exosomal protein biomarkers, it is also challenging to successfully utilize them to diagnose cancers due to intratumor and interpatient heterogeneity[14]. Conventional diagnostic approaches predominantly rely on a single biomarker, which is often not specific or sensitive[15, 16]. However, recent advances in machine learning

60   algorithms connected to artificial intelligence provide an opportunity to construct a classifier that

61   identifies a panel of exosomal protein biomarkers that would possess a more comprehensive

62   ability to reflect the complex disease status of different patients and distinguish cancer samples

63   from normal samples with significantly improved sensitivity and specificity.

65   **Results**
66
Unbiased proteomics analysis of exosomes identifies 18 abundant plasma membrane

protein markers for various human cell lines

To identify universal exosomal protein biomarkers for differentiating cancer from non-cancer

exosomes, we analyzed protein abundance data from 228 cancer and 57 control cell line-derived

exosomes, representing various cancer types (**Figure 1A; Supplementary Table S1**). Because

studies employ distinct isolation and mass spectrometry quantification techniques, the number of

identified proteins is different among the studies. To overcome the bias caused by such technical

factors, we examined the proteins common to all studies and identified 1124 overlapping

proteins (**Figure 2A**). To determine the heterogeneity among cancer and control cell line-derived

exosomes, we performed principal component analysis (PCA) using these 1124 proteins in 285

different studies with cancer and control cell line-derived exosomes (**Figure 2B**). The PCA

indicated that the exosomes derived from cancer and control cell lines are heterogeneous and

show significant variation in protein expression across cell lines.

We next investigated the frequency of the proteins detected in the exosomes from all cell

lines, only the cancer cell lines and just the control cell lines. Commonly used exosome

biomarkers (e.g., CD9, tetraspanins) were examined first, and from the 12 traditional exosome

markers[17-19], only eight proteins were detected among the 1124 overlapping proteins from all cell

lines. Six of eight proteins were detected in at least 90% of all samples, with CD9 and HSPA4

detected with the least frequency (**Figure 2C**). In addition, the frequency of FLOT1, FLOT2 and

TSG101 proteins was higher in cancer cell line-derived exosomes when compared to control cell

line-derived exosomes (**Figure 2C**).

88   To identify biomarkers detectable at a high frequency in all cancer and control cell line-

derived exosomes, we searched for the proteins that were detected in ≥ 90% of all samples

(**Supplementary Table S2**). We annotated these proteins using IPA and found that 78.0% of

these proteins localize to the cytoplasm and 13.5% of proteins are associated with the plasma

membrane (**Figure 2D**). Gene Ontology (GO) analysis revealed the enrichment of proteins from

pathways related to vesicle-mediated transport, secretory vesicles, exocytosis, endocytosis, and

other exosome-related pathways (**Figure 2E**). To further explore the utility of the proposed

biomarkers, we examined the proteins located on the plasma membrane that met the threshold of

detection in ≥ 90% of all samples (**Figure 2F**). Clathrin Heavy Chain (CLTC) was ranked as the

top plasma membrane protein detected in 99.6% of all samples and 100% of control samples

(**Figure 2F**). In addition, the scaffolding protein, Syntenin-1(SDCBP) was detected at a high

frequency of 97.9% of all samples, corroborating previous findings[20]. Next, we sought to identify

unique markers that can identify cancer cell-derived exosomes (cancer exosomes) by filtering out

the proteins present in ≤ 10% of 57 control cell line-derived exosomes. Interestingly, Ataxin 2

Like (ATXN2L), which has been reported to promote cancer cell invasiveness and resistance to

chemotherapy, was uniquely detected in the cancer cell-derived exosomes (**Figure 3A**)[21]. In

total, we identified a set of 18 exosome protein markers that are present at a high abundance in

all the cancer exosomes examined (**Figure 2F**).

Comparison of exosomal proteins derived from cell lines and tissues identified five universal plasma membrane protein markers

We next sought to investigate common biomarkers for the tissue-derived exosomes across cancer types. We calculated the detection frequency of commonly used exosome markers in the 157 samples (101 cancers; 56 controls). Two established exosome markers, CD63 and TSG101, were only detected in 33.1% and 45.9% of all samples, respectively (**Figure 3B**). To identify high-frequency biomarkers for exosomes derived from both cell lines and tissues, we examined the overlapping proteins that met a threshold of ≥ 90% in all samples for exosomes derived from cell lines and tissues and found 31 common proteins (**Figure 3C**). Among the 31 proteins, there were five proteins that were detected in over 90% of all cell line and tissue-derived exosomes (**Figure 3D**). These include Clathrin Heavy Chain (CLTC), Ezrin, (EZR), Talin-1 (TLN1), Adenylyl cyclase-associated protein 1 (CAP1) and Moesin (MSN).

An exosome proteome signature of 18 proteins can differentiate cancer exosomes from non-cancer exosomes across multiple cancer types

Plasma or serum is the most readily accessible source for non-invasive biopsies. We next sought to identify if exosome proteins in plasma and serum could differentiate cancer exosomes from non-cancer exosomes across multiple cancer types. We pooled exosome proteomics data derived from plasma or serum for 205 cancer and 51 control samples from five different studies[22-26], which included breast cancer, colorectal cancer, glioblastoma, lung carcinoma, liver cancer, neuroblastoma, and pancreatic cancer (**Supplementary Table S1**). To account for differences in the methodologies of these studies, we first reviewed the proteins and identified 46 proteins that were detected in all the studies (**Figure 4A**). We then examined their abundances in the 205 cancer and 51 control samples (**Figure 4B**). Although we could detect differences between the

exosomes derived from cancer cell and non-cancer cells, it was difficult to classify them robustly

based on PCA (**Supplementary Figure S1A**). Therefore, we sought to employ the advanced

machine learning algorithm to differentiate cancer exosomes from non-cancer exosomes. We

first calculated the mutual information score (MI) for each protein and trained the random forest

classifier using the different numbers of top proteins according to their MI score to determine the

best set of proteins we should include in the classifier. We found that the model performed best

using 18 proteins, where the performance was evaluated by the area under the curve of the

receiver operating characteristic curve (AUROC) (**Figure 4C**). Several key cancer-associated

proteins were included among these 18 proteins. For example, apolipoprotein C1 (APOC1),

which was ranked as the top protein, is significantly decreased in samples of cancer patients

(**Supplementary Figure S1B**) and has been previously reported to be down-regulated in the

non-small lung cancer, colorectal cancer, papillary thyroid carcinoma and pediatric

nephroblastoma[27-30].

Employing all 18 proteins with 5-fold cross-validation, we constructed a random forest

classifier[31] to distinguish cancer and control samples and compared it with multiple popular

machine learning models, including Support Vector Machine (SVM)[32], K Nearest Neighbor

Classifier (K-NN)[33] and Gaussian Naive Bayes[34]. Our random forest classifier demonstrated the

highest AUROC (**Figure 4D**). More specifically, it yielded an AUROC of 0.96, with an accuracy

of 0.92, a precision of 0.94, and a recall of 0.96 (**Figure 4E& F**). When applied to the

independent test set, the model yielded an AUROC of 0.99, an accuracy of 0.95, a precision of

0.96 and a recall of 0.98 (**Figure 4G& H**). Importantly, only one sample was misclassified in the

independent test set, and 51 cancer samples were correctly classified. Taken together, our results

showed the advantage and clinical potential of applying the random forest classifier model to

plasma or serum exosome protein based liquid biopsy for cancer diagnosis.

Five plasma/serum exosomal proteins can reliably differentiate five common cancer types

We next sought to further enhance the clinical utility of the exosomes for differentiating between

cancer types. We analyzed proteomics data from plasma or serum derived exosomes from

patients with five common cancer types, including breast cancer, colorectal cancer, glioma, lung

cancer and pancreatic cancer. Initial PCA revealed differences in exosome levels among cancer

patients but failed to distinguish the five cancer types (**Figure 5A**). We determined the crucial

features for cancer-type classification by computing mutual information scores for 46 common

proteins and built a random forest classifier to determine the optimal number of features to

include in the final classifier. We ultimately selected a set of five proteins based on AUROC

scores (**Figure 5B**). We then increased the independent testing data size by utilizing 40% of the

total samples and used the remaining 60% as the training set to minimize overfitting issues.

Employing the five proteins with 5-fold cross-validation to train the random forest classifier, the

model achieved a very high accuracy of 0.99 (**Figure 5C**), and when applied to the independent

test set, the model consistently yielded a high accuracy of 0.94 (**Figure 5D**). The abundance of

the five protein features varied across the five cancer types, reflecting the potential roles of these

proteins in specific cancers (**Figure 5E**). For example, Histidine-rich glycoprotein (HRG),

which was abundant in the colorectal cancer plasma-derived exosomes, has been reported to

promote the tumor migration of colorectal cancer patients[35]. Overall, our results demonstrated

that this exosome protein-based classification model can reliably differentiate between cancer

types and further enhance the diagnostic value of our approach.

A urinary exosome proteome signature consisting of 17 proteins detects cancer exosomes across multiple cancer types

Urine is emerging as a superior non-invasive marker for urologic cancers, as its composition directly reflects the physiological changes in the urogenital system[36]. To test the use of urinary exosome proteins in cancer diagnosis, we collected data from 261 cancer patient samples and 124 control samples from four studies, including bladder cancer, prostate cancer, and renal cancer, lung cancer, cervical cancer, colorectal cancer, esophageal and gastric cancer (**Supplementary Table S1**)[36-39]. Upon examination of proteins common across all four studies, 229 proteins were identified (**Figure 6A**). PCA revealed the variance between samples but failed to differentiate between cancer and control samples (**Figure 6B**). As described earlier, we next employed the random forest classifier method to differentiate cancer and control samples based on their exosomal proteomic profiles. To reduce the feature space and select the most relevant features, we utilized the mutual information score to rank the 229 protein features and then trained the random forest classifier using varying numbers of the top-ranking proteins (**Figure 6C**). Based on the AUROC scores of including the different number of features, we selected 17 features that resulted in the highest AUROC score (**Figure 6C**). A majority of these 17 proteins displayed significant variations in abundance between cancer and control samples (**Figure 6D**). By training a random forest classifier with the 17 protein features and 5-fold cross-validation, the model achieved an AUROC of 0.96, an accuracy of 0.90, a precision of 0.92, and a recall of 0.93 (**Figure 6E& F**). When tested on an independent set, the model produced an AUROC of 0.91, an accuracy of 0.82, a precision of 0.83, and a recall of 0.92 (**Figure 6G& H**). To summarize, our findings indicated the promising clinical potential of using urinary exosome proteins for the diagnosis of urologic cancers as well as other non-urologic cancer types.

Discussion

Liquid biopsy has numerous benefits in the early detection of cancer, categorizing cancer types,

tracking cancer progression, and monitoring response to treatment[40]. Exosomes found in

biological fluids can provide a forensic view of their cells of origin. Despite previous studies

proposing common protein biomarkers to identify exosomes, a comprehensive set of exosome

biomarkers derived from different biological materials has not been established, owing to

limitations in isolation and quantification methods[41-44]. Additionally, a reliable diagnostic tool

based on proteins associated with exosomes that can be applied across all cancers is yet to be

identified.

Here, we generate a comprehensive proteomics profile of exosomes derived from cell

lines, tissues, plasma, serum, and urine from 1083 cancer and control samples. An extensive

analysis of these samples showed that several widely used exosome markers, such as CD63,

CD81, HSP70, and HSPA8, are absent in exosomes derived from a subset of cell lines. Further,

FLOT1, FLOT2, TSG101, and CD63 are present at low levels (< 60%) in all exosomes derived

from tissues, indicating a need for the identification of additional universal markers for both cell

line- and tissue-derived exosomes. Our study identifies five highly abundant universal exosome

biomarkers-CLTC, EZR, TLN1, CAP1, and MSN- that were present in over 90% of all cell line-

and tissue-derived samples. Additionally, we found that ATXN2L was only present in cancer

exosomes and absent in non-cancer exosomes. In this regard, GPC1 which was identified in

many studies identified as cancer exosomes specific marker, was not identified here as many of

databases did not pick up this protein in analysis.

Here we describe a novel computational approach using the random forest classifier

method to define exosome protein panels that serve as effective biomarkers specifically for

plasma, serum or urine across cancer types. By training the random forest model and testing with

independent datasets, our model yields excellent scores in AUROC, sensitivity, and specificity

and differentiates cancer exosomes from non-cancer exosomes. We show that this approach can

also be used to classify, with high accuracy, five common cancer types based on their exosome

protein signatures.

A majority of the protein makers identified in this study have demonstrable biological

relevance in cancer. As an example, ITIH3, which was identified among the protein features for

the plasma or serum-based classifier and highly abundant in cancer samples, was reported to be

more highly expressed in the plasma of gastric cancer samples compared to the control[45] and

increased with tumor staging of clear cell renal cell carcinoma patients [46]. Importantly, the

biomarker panels identified for cell lines, tissues, plasma/serum, and urine overcome the bias

associated with the isolation and quantification method as well as the inter-patient variability

inherent to the complex process of cancer. These panels are designed to be applied to a variety of

cancers.

Collectively, our results demonstrate that exosome protein features can be utilized as

reliable biomarkers for the early detection of cancer, classification of cancer types, and

potentially for diagnosing tumors of undetermined origin. These results have the potential to

advance the development and standardization of innovative and optimized methods for the

isolation of exosomes and the implementation of routine plasma, serum- and urine-based

exosome screening in clinical settings.

Methods

Public exosome proteomics data

We collected publicly available exosome protein data from cell lines, plasma, serum, and urine from previous studies as summarized in **Supplementary Table S1**[20, 22-26, 36-39, 43, 47]. We obtained the raw spectral count and intensity data from the original paper or directly from the authors through communication. The obtained data was log normalized accordingly.

Feature selection and machine learning algorithm

We employed mutual information scores to evaluate the importance of protein features for each prediction, quantifying the amount of uncertainty reduction for one variable given the knowledge of another variable. We calculated the mutual information score for each protein with a target label (cancer/control) using SelectKBest from the scikit-learn 1.1.1 Python library. According to the sample size and characteristics of the data collected for plasma and urine-derived exosomes, we selected a customized number of best features for each prediction. To select the optimum number of proteins to include in the prediction, we ranked the protein features based on the mutual information score and built a random forest classifier to evaluate performance on including a range of number of features. The random forest model can decrease the probability of over-fitting and enhances the resilience towards outliers and input data noise. The area under the curve of the receiver operating characteristic curve (AUROC) was employed to evaluate the performance of the classifier. We also calculated accuracy, precision and recall for comprehensive evaluation. All models were evaluated using 5-fold cross-validation with stratified train-test splits that preserved the percentage of samples for the prediction target. We also tested the performance of alternate machine learning algorithms including support vector classifier, K nearest neighbor classifier and gaussian naive bayes. Overall, the random forest

classifier achieved the best performance in our analysis (**Supplementary Figure S1B**). To

visualize high-dimensional datasets, kernel principal component analysis (PCA) from scikit-learn

1.1.1 Python library was employed to perform PCA and plots were generated using the

scatterplot function from seaborn 0.11.2 Python library.

Gene ontology and pathway enrichment analysis

The WebGestalt 2019[48] online tool was used to perform the gene ontology and pathway

enrichment analysis of the selected proteins. Pathways with FDR < 0.05 were considered

significant.

Statistical analysis

All statistical analyses were conducted using R 4.2.1 software. Significance was determined by

the Wilcoxon rank-sum test unless specified otherwise. Significance was concluded if the p-

value was <0.05, while in pathway analysis, significance was concluded if the FDR was <0.05

after correction for multiple comparisons.

Disclosure of conflict of interest

UT MD Anderson Cancer Center and R.K. hold patents in the area of exosome biology licensed

to Codiak Biosciences, Inc. UT MD Anderson Cancer Center and R.K. are stock equity holders

in Codiak Biosciences, Inc. R.K. is a consultant and scientific adviser for Codiak Biosciences,

Inc. The remaining authors declare no competing interests.

Acknowledgments

The EV work in the Kalluri lab is supported by MD Anderson Cancer Center, NIH

R35CA263815, and NIH P40OD024628 and gifts from Fifth Generation (Love, Tito's), Lyda

Hill Philanthropies, and Bosarge Family Trust.

Data availability

The public datasets used in this study were described in the Methods sections.

Contributions

B.L. and R.K. conceived and designed the study. B.L. developed the computational algorithm

and performed the analysis with supervision from R.K. F.G.K. generated the proteomics data of

the cell lines. B.L. and R.K. wrote the manuscript. All the authors read the manuscript and

discussed the results.

References

1. Distler, M., Aust, D., Weitz, J., Pilarsky, C. & Grutzmann, R. Precursor lesions for sporadic pancreatic cancer: PanIN, IPMN, and MCN. *Biomed Res Int* **2014**, 474905 (2014).
2. Morris, J.P.t., Cano, D.A., Sekine, S., Wang, S.C. & Hebrok, M. Beta-catenin blocks Kras-dependent reprogramming of acini into pancreatic cancer precursor lesions in mice. *J Clin Invest* **120**, 508-520 (2010).
3. Suram, A. et al. Oncogene-induced telomere dysfunction enforces cellular senescence in human cancer precursor lesions. *EMBO J* **31**, 2839-2851 (2012).
4. Mullerat, J., Deroide, F., Winslet, M.C. & Perrett, C.W. Proliferation and p53 expression in anal cancer precursor lesions. *Anticancer Res* **23**, 2995-2999 (2003).
5. Hammes, L.S. et al. Up-regulation of VEGF, c-fms and COX-2 expression correlates with severity of cervical cancer precursor (CIN) lesions and invasive disease. *Gynecol Oncol* **110**, 445-451 (2008).
6. Oshi, M. et al. Urine as a Source of Liquid Biopsy for Cancer. *Cancers (Basel)* **13** (2021).
7. Hu, C. et al. Potentiality of Exosomal Proteins as Novel Cancer Biomarkers for Liquid Biopsy. *Front Immunol* **13**, 792046 (2022).
8. Kalluri, R. & LeBleu, V.S. The biology, function, and biomedical applications of exosomes. *Science* **367** (2020).
9. Johnstone, R.M., Adam, M., Hammond, J.R., Orr, L. & Turbide, C. Vesicle formation during reticulocyte maturation. Association of plasma membrane activities with released vesicles (exosomes). *J Biol Chem* **262**, 9412-9420 (1987).
10. Maas, S.L.N., Breakefield, X.O. & Weaver, A.M. Extracellular Vesicles: Unique Intercellular Delivery Vehicles. *Trends Cell Biol* **27**, 172-188 (2017).
11. Colombo, M., Raposo, G. & Thery, C. Biogenesis, secretion, and intercellular interactions of exosomes and other extracellular vesicles. *Annu Rev Cell Dev Biol* **30**, 255-289 (2014).
12. Valencia, K. & Montuenga, L.M. Exosomes in Liquid Biopsy: The Nanometric World in the Pursuit of Precision Oncology. *Cancers (Basel)* **13** (2021).
13. Chen, I.H. et al. Phosphoproteins in extracellular vesicles as candidate markers for breast cancer. *Proc Natl Acad Sci U S A* **114**, 3175-3180 (2017).
14. Muluneh, M., Shang, W. & Issadore, D. Track-etched magnetic micropores for immunomagnetic isolation of pathogens. *Adv Healthc Mater* **3**, 1078-1085 (2014).
15. Ballehaninna, U.K. & Chamberlain, R.S. Serum CA 19-9 as a Biomarker for Pancreatic Cancer-A Comprehensive Review. *Indian J Surg Oncol* **2**, 88-100 (2011).
16. Singh, S., Tang, S.J., Sreenarasimhaiah, J., Lara, L.F. & Siddiqui, A. The clinical utility and limitations of serum carbohydrate antigen (CA19-9) as a diagnostic tool for pancreatic cancer and cholangiocarcinoma. *Dig Dis Sci* **56**, 2491-2496 (2011).
17. Thery, C., Amigorena, S., Raposo, G. & Clayton, A. Isolation and characterization of exosomes from cell culture supernatants and biological fluids. *Curr Protoc Cell Biol* **Chapter 3**, Unit 3 22 (2006).
18. Kowal, J. et al. Proteomic comparison defines novel markers to characterize heterogeneous populations of extracellular vesicle subtypes. *Proc Natl Acad Sci U S A* **113**, E968-977 (2016).

19.   Jeppesen, D.K. et al. Reassessment of Exosome Composition. *Cell* **177**, 428-445 e418
(2019).
20.   Kugeratski, F.G. et al. Quantitative proteomics identifies the core proteome of exosomes
with syntenin-1 as the highest abundant protein and a putative universal biomarker. *Nat*
Cell Biol **23**, 631-641 (2021).
21.   Lin, L. et al. ATXN2L upregulated by epidermal growth factor promotes gastric cancer
cell invasiveness and oxaliplatin resistance. *Cell Death Dis* **10**, 173 (2019).
22.   Hoshino, A. et al. Extracellular Vesicle and Particle Biomarkers Define Multiple Human
Cancers. *Cell* **182**, 1044-1061 e1018 (2020).
23.   Vykoukal, J. et al. Plasma-derived extracellular vesicle proteins as a source of biomarkers
for lung adenocarcinoma. *Oncotarget* **8**, 95466-95480 (2017).
24.   Li, S. et al. Proteomic Landscape of Exosomes Reveals the Functional Contributions of
CD151 in Triple-Negative Breast Cancer. *Mol Cell Proteomics* **20**, 100121 (2021).
25.   Lin, K. et al. Comprehensive proteomic profiling of serum extracellular vesicles in
patients with colorectal liver metastases identifies a signature for non-invasive risk
stratification and early-response evaluation. *Mol Cancer* **21**, 91 (2022).
26.   Hallal, S. et al. A Comprehensive Proteomic SWATH-MS Workflow for Profiling Blood
Extracellular Vesicles: A New Avenue for Glioma Tumour Surveillance. *Int J Mol Sci* **21**
(2020).
27.   Zhang, Q., Wang, J., Dong, R., Yang, S. & Zheng, S. Identification of novel serum
biomarkers in child nephroblastoma using proteomics technology. *Mol Biol Rep* **38**, 631-
638 (2011).
28.   Jin, Y. et al. Identification a novel clinical biomarker in early diagnosis of human non-
small cell lung cancer. *Glycoconj J* **36**, 57-68 (2019).
29.   Engwegen, J.Y. et al. Detection of Colorectal Cancer by Serum and Tissue Protein
Profiling: A Prospective Study in a Population at Risk. *Biomark Insights* **3**, 375-385
(2008).
30.   Fan, Y. et al. Discovery and identification of potential biomarkers of papillary thyroid
carcinoma. *Mol Cancer* **8**, 79 (2009).
31.   Ho, T.K. Random decision forests. *Proceedings of 3rd International Conference on*
Document Analysis and Recognition **1**, 278-282 vol.271 (1995).
32.   Cortes, C.a.V., Vladimir Support-vector networks. *Machine learning* **20**, 273-297 (1995).
33.   Fix, E.a.H., Joseph l. Discriminatory Analysis. Nonparametric Discrimination:
Consistency Properties. *International Statistical Review / Revue Internationale de*
Statistique **57**, 238-247 (1989).
34.   Pedregosa, F. et al. Scikit-learn: Machine Learning in Python. *J Mach Learn Res* **12**,
2825-2830 (2011).
35.   Bogoevska, V. et al. HRG/HER2/HER3 signaling promotes AhR-mediated Memo-1
expression and migration in colorectal cancer. *Oncogene* **36**, 2394-2404 (2017).
36.   Dhondt, B. et al. Unravelling the proteomic landscape of extracellular vesicles in prostate
cancer by density-based fractionation of urine. *J Extracell Vesicles* **9**, 1736935 (2020).
37.   Zhang, C. et al. Urine Proteome Profiling Predicts Lung Cancer from Control Cases and
Other Tumors. *EBioMedicine* **30**, 120-128 (2018).
38.   Overbye, A. et al. Identification of prostate cancer biomarkers in urinary exosomes.
Oncotarget **6**, 30357-30376 (2015).

39. Suh, J. et al. Next-generation Proteomics-Based Discovery, Verification, and Validation
of Urine Biomarkers for Bladder Cancer Diagnosis. *Cancer Res Treat* **54**, 882-893 (2022).
40. Nikanjam, M., Kato, S. & Kurzrock, R. Liquid biopsy: current technology and clinical
applications. *J Hematol Oncol* **15**, 131 (2022).
41. Castillo, J. et al. Surfaceome profiling enables isolation of cancer-specific exosomal
cargo in liquid biopsies from pancreatic cancer patients. *Ann Oncol* **29**, 223-229 (2018).
42. Gangoda, L. et al. Proteomic Profiling of Exosomes Secreted by Breast Cancer Cells with
Varying Metastatic Potential. *Proteomics* **17** (2017).
43. Hurwitz, S.N. et al. Proteomic profiling of NCI-60 extracellular vesicles uncovers
common protein cargo and cancer type-specific biomarkers. *Oncotarget* **7**, 86999-87015
(2016).
44. Ji, H. et al. Proteome profiling of exosomes derived from human primary and metastatic
colorectal cancer cells reveal differential expression of key metastatic factors and signal
transduction components. *Proteomics* **13**, 1672-1686 (2013).
45. Chong, P.K. et al. ITIH3 is a potential biomarker for early detection of gastric cancer. *J*
Proteome Res **9**, 3671-3679 (2010).
46. Chang, Q.H. et al. Pan-cancer analysis identifies ITIH1 as a novel prognostic indicator
for hepatocellular carcinoma. *Aging (Albany NY)* **13**, 11096-11119 (2021).
47. Rontogianni, S. et al. Proteomic profiling of extracellular vesicles allows for human
breast cancer subtyping. *Commun Biol* **2**, 325 (2019).
48. Liao, Y., Wang, J., Jaehnig, E.J., Shi, Z. & Zhang, B. WebGestalt 2019: gene set analysis
toolkit with revamped UIs and APIs. *Nucleic Acids Res* **47**, W199-W205 (2019).

415
416

417 **Figure legends**

419 **Figure 1. Overview of the study.**

420
421

Figure 2. Proteomic characterization of exosomes derived from 285 cell lines from four

studies. (A) Overlapped proteins from four different studies of cell line-derived exosomes. **(B)**

PCA plot of cancer and control cell line-derived exosomes. **(C)** Positivity for 8 commonly used

exosomal protein biomarkers in various cell lines. The percentage of samples expressing each

protein is shown in the boxes. Darker red indicates a higher percentage. **(D)** Annotation of the

proteins detected in more than 90% of all samples. **(E)** GO and KEGG pathway enrichment

analysis of the proteins detected in more than 90% of all samples. **(F)** Plasma membrane proteins

detected in more than 90 % of all samples.

Figure 3. Proteomic characterization of exosomes derived from cell lines and tissues. (A)

Proteins detected at higher frequency in cancer cell line-derived exosomes. **(B)** Positivity for 11

commonly used exosomal protein biomarkers in various tissues. **(C)** Overlapping proteins (>90%

frequency) between cell line- and tissue-derived exosomes. **(D)** Positivity of five plasma

membrane proteins detected in more than 90% of both cell line- and tissue-derived exosomes.

Figure 4. Identification of the signature proteins of plasma or serum-derived exosomes and

the evaluation of random forest classifier. (A) Overlapping exosome proteins detected in the

plasma and serum of 205 cancer and 51 control samples from five studies. **(B)** Heat map of 46

overlapping exosome proteins in cancer and control plasma or serum samples. **(C)** AUROC

score of the random forest classifier on including various numbers of protein features. **(D)**

AUROC of different models in comparison. **(E)** Classification error matrix of the 75% training

set using a random forest classifier for the 18 selected proteins. The number of samples is

indicated in each box. **(F)** AUROC score of the random forest classifier trained using 75% of the

dataset. Other metrics are indicated on right. **(G)** Classification error matrix of 25% testing set

using a random forest classifier for the 18 selected proteins. The number of samples is indicated

in each box. **(H)** AUROC score of the random forest classifier tested using 25% of the dataset.

Other metrics are indicated on right.

Figure 5. Identification of signature proteins expressed by plasma or serum-derived

exosomes for classifying five common cancer types and evaluation of random forest

classifier. (A) PCA plot of plasma or serum-derived exosomes from five cancer types. **(B)**

AUROC score of the random forest classifier by including various number of protein features.

(C-D) Classification error matrix of a 60% training set and 40% testing set to classify the five

cancer types using a random forest classifier for the 5 selected proteins. The number of samples

is indicated in each box. **(E)** Protein abundance of five selected protein features in five cancer

types.

Figure 6. Identification of signature proteins expressed by urine-derived exosomes and

evaluation of random forest classifier. (A) Overlapping exosome proteins detected in the urine

from 261 cancer and 124 control samples from four studies. **(B)** PCA plot of cancer and control

urine-derived exosomes. **(C)** AUROC score of the random forest classifier by including a various

number of protein features. **(D)** Protein abundance of 17 selected protein features in cancer- and

control urine-derived exosomes. **(E)** Classification error matrix of 75% training set using a

random forest classifier for the 17 selected proteins. The number of samples is indicated in each

box. **(F)** AUROC score of the random forest classifier trained using 75% of the dataset. Other

metrics are indicated on right. **(G)** Classification error matrix of 25% testing set using a random

forest classifier for the 17 selected proteins. The number of samples is indicated in each box. **(H)**

AUROC score of the random forest classifier tested using 25% of the dataset. Other metrics are

indicated on right.

Supplementary Figure S1. Machine learning models for plasma-derived exosomes. (A) PCA

plot of cancer and control plasma or serum-derived exosomes. **(B)** Protein abundance of 18

selected protein features in cancer- and control plasma or serum-derived exosomes.

Figure 1

A

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6