Antimicrobial activity of iron-depriving pyoverdines against human

opportunistic pathogens

Vera Vollenweider[1], Karoline Rehm[2], Clara Chepkirui[3], Manuela Pérez-Berlanga[1],

Magdalini Polymenidou[1], Jörn Piel[3], Laurent Bigler[2], Rolf Kümmerli[1]

[1]Department of Quantitative Biomedicine, University of Zurich, Zurich, Switzerland

[2]Department of Chemistry, University of Zurich, Zurich, Switzerland

[3]Institute of Microbiology, Eidgenössische Technische Hochschule (ETH) Zurich,

Zurich, Switzerland

Short title: Pyoverdines against pathogens

Corresponding author: Rolf Kümmerli, Department of Quantitative Biomedicine,

University of Zurich, Winterthurerstrasse 190, 8057 Zurich, Switzerland.

Email: rolf.kuemmerli@uzh.ch / Phone: +41 44 635 48 01

Abstract

The global rise of antibiotic resistance calls for new drugs against bacterial pathogens. A common approach is to search for natural compounds deployed by microbes to inhibit competitors. Here we show that the iron chelating pyoverdines, siderophores produced by environmental *Pseudomonas* spp., have strong antibacterial properties by inducing iron starvation and growth arrest in pathogens. A screen of 320 natural *Pseudomonas* isolates used against 12 human pathogens uncovered several pyoverdines with particularly high antibacterial properties and distinct chemical characteristics. The most potent pyoverdine effectively reduced growth of the pathogens *Acinetobacter baumannii*, *Klebsiella pneumoniae* and *Staphylococcus aureus* in a concentration- and iron-dependent manner. Pyoverdine increased survival of infected *Galleria mellonella* host larvae, and showed low toxicity for the host, mammalian cell lines, and erythrocytes. Furthermore, experimental evolution combined with whole-genome sequencing revealed reduced potentials for resistance evolution compared to an antibiotic. Thus, pyoverdines from environmental strains have the potential to become a new class of sustainable antibacterials against specific human pathogens.

Introduction

There are approximately 4.95 million fatalities associated with antibiotic-resistant bacteria worldwide each year[1]. This high mortality rate contrasts with the low development rate of new antibacterial agents for clinical applications[2,3]. Thus, the development of alternative therapies with higher sustainability to counteract the rapidly dwindling treatment options due to resistance evolution is imperative[4–6]. Numerous alternative approaches have been proposed including phage therapy, antimicrobial peptides and nanobodies, anti-virulence treatments to disarm pathogens, treatments that capitalize on ecological competition between susceptible and resistant bacteria, and the search for new antibacterial compounds produced by environmental microbes[7–12].

Mining natural microbial communities has become a promising endeavour in the hunt for novel antimicrobials because most microbes deploy bioactive compounds to contend with their competitors in the diverse assemblies they live in. Competition often involves the secretion of secondary metabolites with specific antimicrobial properties against other microbes[13]. The isolation, characterization and synthesis of such secondary metabolites can result in novel prospective antibiotics for clinical applications[14,15]. Secondary metabolites that act as toxins to kill competitors are often considered promising candidates. However, microbes compete through a variety of mechanisms other than toxins. For example, there is increasing evidence that competition for iron is a main determinant of species interactions in soil and freshwater communities[16,17]. Competition for iron involves the secretion of siderophores, a class of secondary metabolites that scavenges environmental iron with high affinity[18]. Because siderophores are often species specific, they have two opposing effects: they make iron available for related community members possessing the matching receptor but withhold iron from competitors with non-matching receptors. Thus, siderophores can be competitive agents in inter-species interactions[19–22].

Here, we apply the concept of siderophore-mediated iron competition observed in natural communities to human opportunistic pathogens. Specifically, we investigate whether siderophores from non-pathogenic environmental bacteria can induce iron starvation and growth arrest in human pathogens. We focus on pyoverdines, a class of siderophores with high iron affinity that are produced and

secreted by fluorescent *Pseudomonas* spp. under iron-limited conditions[23]. Upon

complexation with ferric iron, the iron-loaded pyoverdine is imported into the cell by

receptors with high specificity and subsequently reduced to the bio-available ferrous

iron[24,25]. Pyoverdines show an extraordinary structural diversity with more than 70

described variants that differ in their peptide backbone[26,27], and have an extremely

high affinity for iron (K_a = 10^{32} M^{-1}, pyoverdine produced by *Pseudomonas*

fluorescens biotype B)[28]. For these reasons, we propose that pyoverdines from non-

pathogenic *Pseudomonas* spp. could be potent agents to reduce the growth of

opportunistic human pathogens by intensifying iron competition.

To test our hypothesis, we screened pyoverdines from a library of 320

environmental *Pseudomonas* strains, isolated from soil and freshwater habitats, for

their activity against 12 strains of human opportunistic bacteria. For the most

promising pyoverdine candidates, we elucidated the chemical structure to assess

diversity and to understand the chemical properties important for iron competition.

Subsequently, we assessed the efficacy of the three top pyoverdine candidates

against four human opportunistic pathogens (*Acinetobacter baumannii*, *Klebsiella*

pneumoniae, *Pseudomonas aeruginosa* and *Staphylococcus aureus*) *in vitro* and/or

in vivo via infections of the Greater wax moth larvae (*Galleria mellonella*). We further

assessed their toxicity in the host, towards two mammalian cell lines and red blood

cells. Finally, we used experimental evolution combined with whole-genome

sequencing to study the potential of pathogens evolving resistance against

pyoverdine treatment.

92　Results

93　Pyoverdines can inhibit the growth of human opportunistic pathogens

To assess the antibacterial properties of pyoverdines, we performed a growth-

inhibition screen using a well-characterized collection of 320 natural *Pseudomonas*

isolates[16,29–31]. The isolates stem from freshwater and soil communities and belong

to the group of fluorescent pseudomonads (e.g, *P. fluorescens*, *P. putida*, *P.*

syringae, *P. chlororaphis*) that are non-pathogenic to humans and can produce and

secrete pyoverdines, a group of high iron-affinity siderophores[32]. Important to note is

that several isolates are likely to produce secondary siderophores in addition to

pyoverdine[33]. Given that secondary siderophores generally have lower iron affinity

than pyoverdine, we anticipate treatment effects to be primarily driven by
pyoverdines.

In a first screen, we evaluated how the sterile pyoverdine-containing
supernatants of the 320 isolates affect the growth of 12 human opportunistic
pathogens (Table 1). Specifically, we grew the pathogens in 70% fresh casamino
acid medium (CAA) supplemented with 30% spent supernatant. We found high
variation in the supernatant effect, ranging from complete pathogen growth inhibition
to growth promotion compared to the control treatment (70% CAA supplemented
with 30% sodium chloride 0.8% solution) (Figure 1A and Figure S1A). The mean
supernatant effect across all pathogens showed a bimodal distribution with two
peaks at 0.47 – 0.58 (approximately 50% growth inhibition) and 1.00 – 1.10 (no
growth inhibition) (Figure S1B). Our observations are consistent with previous
studies, in which supernatants containing siderophores can have both growth
inhibitory and stimulatory effects[20,29]. For each supernatant, we quantified the
number of pathogens that experienced a growth reduction of at least 20% compared
to the control and found that 25 supernatants inhibited all 12 pathogens (Figure 1A
and S1C).

Table 1 | Species and strains of human opportunistic pathogens used for the
pyoverdine growth inhibition assay

Species and strain name	Class	Gram-staining	Frequency of inhibiting supernatants
Acinetobacter johnsonii	Gammaproteobacteria	negative	75.0
Acinetobacter junii	Gammaproteobacteria	negative	73.1
Burkholderia cenocepacia H111	Betaproteobacteria	negative	52.2
Burkholderia cenocepacia K56-2	Betaproteobacteria	negative	54.7
Cronobacter sakazakii	Gammaproteobacteria	negative	48.8
Escherichia coli K12	Gammaproteobacteria	negative	31.6
Klebsiella michiganensis	Gammaproteobacteria	negative	30.9
Pseudomonas aeruginosa PA14	Gammaproteobacteria	negative	9.7
Pseudomonas aeruginosa PAO1	Gammaproteobacteria	negative	20.9
Shigella sp.	Gammaproteobacteria	negative	57.8
Staphylococcus aureus Cowan	Bacilli	positive	47.8
Staphylococcus aureus JE2	Bacilli	positive	41.6

To confirm that the growth-inhibiting properties of the 25 most potent
supernatants are indeed caused by pyoverdines and not by other agents in the
supernatant, we repeated the above assay, but this time supplemented the
supernatant with 40 µM FeCl$_3$. Under such iron-replete conditions, pyoverdines (and
secondary siderophores) should not be able to substantially sequester environmental
iron, and pathogen growth should therefore be restored. We identified seven top
candidate supernatants for which a relative growth recovery of at least 0.2 occurred
for all 12 pathogens (Figure 1B). Among the remaining 18 supernatants, the overall
rate of growth recovery was also high, indicating that pyoverdines are responsible for
pathogen growth inhibition under iron-limited conditions in most cases.

Finally, we assessed the relationship between growth inhibitory effects of
supernatants and the phylogenetic affiliation of the producing strains (Figure S2). We
found that growth inhibitory effects were not restricted to a few closely related strains
but were scattered across the phylogenetic tree. Nonetheless, there was a certain
level of clustering with two *Pseudomonas* spp. clades showing an accumulation of
strains with growth-inhibitory effects on pathogens. This analysis suggests that
closely as well as distantly related strains can produce inhibitory pyoverdines.

141
Figure 1. Effect of supernatants from environmental *Pseudomonas* isolates on the
growth of 12 human opportunistic pathogens. (A) Screen to assess the extent to which
pyoverdine-containing supernatants from 320 natural *Pseudomonas* isolates inhibit the
growth of 12 human opportunistic pathogens. The heatmap depicts relative pathogen growth
in the supernatant treatment (70% CAA + 30% spent supernatant) relative to the control
(70% CAA + 30% sodium chloride solution), with values ranging from stimulation (yellow) to
inhibition (blue) based on four independent replicates. Grey bars above the heatmap show
the number of pathogens that were at least 20% inhibited in their growth by a given

supernatant. The screen returned 25 supernatant candidates that inhibited the growth of all

pathogens. (B) Control screen with the 25 top supernatant candidates to check whether

pyoverdine causes the observed growth inhibition. The heatmap depicts the level of growth

recovery in the 12 pathogens when iron was added to the supernatant, with values ranging

from no recovery (red) to full recovery (blue). Growth recovery in iron-rich medium indicates

that pyoverdines are involved in growth inhibition in iron-limited medium. Grey bars above

the heatmap show the number of pathogens that experienced a relative growth recovery of

at least 0.2. The screen returned 7 supernatant top candidates for which growth recovery

occurred for all pathogens under iron-rich conditions.

Growth-inhibiting pyoverdines show distinct chemical features

In previous studies, we developed a new method for pyoverdine structure elucidation

from low-volume crude extracts using ultra-high-performance liquid chromatography

high-resolution tandem mass spectrometry (UHPLC-HR-MS/MS)[27,34]. These studies

included the pyoverdines extracted from the seven top candidate supernatants and

the analysis therein yielded five structurally different pyoverdines (Figure 2A). We

found one novel pyoverdine type (**1**, from isolate 3A06) and four pyoverdine types

already known from the literature (**2**-**5**). The pyoverdine from isolate 3G07 can occur

in either a linear or cyclic conformation (**2a** and **2b**). Three isolates (s3b09, s3b10,

s3b12), all originating from the same soil sample, had an identical pyoverdine type

(**3**).

Here, we investigated whether the chemical properties of growth-inhibitory

pyoverdines differ from pyoverdines that do not inhibit pathogen growth. To this end,

we further elucidated the chemical structure of a set of six pyoverdines extracted

from supernatants that inhibited none of the 12 pathogens, and one additional

pyoverdine inhibiting the growth of 10 pathogens[27]. We found that inhibitory

pyoverdines had significantly larger collision-cross sections (CCS in Å^2) than non-

inhibitory ones ($F_{1,22} = 10.63$, $p = 0.0036$, measured by ion mobility spectrometry

(IMS)) (Figure 2B). As CCS values correlate with the volume of a molecule, our

results suggest that inhibitory pyoverdines are larger and have more complex

molecule structures than non-inhibitory pyoverdines. In contrast, there was no

significant difference between the two classes of pyoverdines regarding molecule

polarity ($F_{1,22} = 2.00$, $p = 0.1711$, assessed by the chromatographic retention time on

a reverse phase column) (Figure 2C), but molecule polarity was significantly higher

for ferri- than for apo-pyoverdines ($F_{1,22}$ = 12.78, p = 0.0017). Finally, we assessed

the stability of ferri-pyoverdines to derive a proxy for iron affinity. We first measured

the propensity of ferri-pyoverdines to undergo collision-induced unfolding (CIU) by

IMS measurements at different Δ6 voltages and found that inhibitory pyoverdines are

significantly more stable than non-inhibitory pyoverdines ($F_{1,11}$ = 7.55, p = 0.019)

(Figure 2D). The same trend was observed for our second measure of stability

(CE50, normalised collision energy necessary to fragment 50% of ferri-pyoverdines),

but in this case, the difference between the two pyoverdine classes was not

significant ($F_{1,11}$ = 1.87, p = 0.198) (Figure 2E). Taken together, the chemical

analyses suggest that inhibitory pyoverdines are larger and have a higher affinity for

iron than non-inhibitory pyoverdines.

Figure 2. Chemical structure of growth-inhibitory pyoverdines and their properties compared to non-inhibitory pyoverdines. (A) The chemical structures of the seven top candidate pyoverdines were elucidated using UHPLC-HR-MS/MS and revealed 5 unique pyoverdine structures (**1**-**5**) differing in their peptide backbone (see[27] for an in-depth chemical analysis). Pyoverdine **1** is a novel structure (from isolate 3A06). Pyoverdine **2** can occur in either a linear **2a** or a cyclic **2b** form (from isolate 3G07). Pyoverdine **3** was found in three different isolates, originating from the same soil sample (from isolates s3b09, s3b10 and s3b12). Pyoverdine **4** and **5** are from isolate s3c13 and s3e20, respectively. (B) The CCS values of inhibitory pyoverdines are higher than for non-inhibitory pyoverdines. (C) Molecule polarity, measured as the chromatographic retention time, is higher for ferri-pyoverdines (iron loaded) than for apo-pyoverdines (iron free). (D) Iron complex stability, assessed by the collision induced unfolding of ferri-pyoverdines, was significantly higher for inhibitory than non-inhibitory pyoverdines. (E) The normalised collision energy (CE50) necessary to fragment 50% of ferri-pyoverdines was not different between inhibitory and non-inhibitory pyoverdines. Box plots show the median and the first and third quartiles across the 7 inhibitory and 6 non-inhibitory pyoverdines. Whiskers represent the 1.5x interquartile range.

Pyoverdines inhibit human pathogens in a concentration-dependent manner

We purified three out of the five most potent pyoverdines and tested their efficacy against the four human pathogens *A. baumannii*, *K. pneumoniae, P. aeruginosa* PAO1 and *S. aureus* JE2. For these experiments, we decided to include the most common (s3b09), the novel (3A06), and the cyclical-linear (3G07) pyoverdines. We crude-purified all three pyoverdines using established protocols[16,35] and further purified the most common pyoverdine (s3b09) using preparative and analytical HPLC. During HPLC purification, we observed that ferribactin (the final pyoverdine precursor) was also highly abundant in the supernatant[36,37], presumably released by cells during the centrifugation process. This precursor is typically not secreted but matures into pyoverdine in the periplasma[38]. Due to its compromised iron-binding capacity, we used it as a negative control for which no activity against pathogens is expected[37]. As positive treatment control, we used the antibiotic ciprofloxacin. We subjected the four pathogens to the three pyoverdine variants, ferribactin, and ciprofloxacin across a concentration gradient and measured their effects on pathogen growth (Figure 3, Figure S3).

For *A. baumannii* and *S. aureus,* we observed classical dose-response curves
in all pyoverdine assays, characterized by a decrease in bacterial growth at low and
intermediate concentrations, followed by a complete halt of bacterial growth at higher
pyoverdine concentrations. There was a good qualitative match between the dose-
response curves obtained with the HPLC vs. the crude-purified pyoverdines. Note
that pyoverdine concentration is unknown for the crude extracts and is thus
expressed as relative concentrations. In contrast, the pyoverdine concentration is
absolute for the HPLC-purified version, allowing the determination of an IC50 value
for pyoverdine s3b09, which is 5.037 µg/mL and 12.163 µg/mL for *A. baumannii* and
S. aureus, respectively.

For *K. pneumoniae*, the pyoverdines were less effective in reducing pathogen
growth. The dose-response curves for the pyoverdines s3b09 (HPLC and crude) and
3A06 showed similar trajectories and the model fits showed a growth reduction
between 9-12%. The strongest inhibitory effects were found for pyoverdine 3G07
with an estimated 54% growth reduction. One reason why pyoverdines might be less
effective against *K. pneumoniae* is that this species produces enterobactin, a
siderophore that has higher iron affinity than pyoverdine[39].

For *P. aeruginosa*, we observed that all pyoverdines were growth inhibitory at
intermediate concentrations. The initial inhibition converted into growth promotions at
high pyoverdine concentrations in three out of four cases. The only exception was
pyoverdine 3G07, which yielded consistent growth inhibitions of about 66%. One
reason for these non-standard dose-response curves could be that *P. aeruginosa*
possesses two receptors (FpvA and FpvB) for pyoverdine uptake[40]. It might thus
capitalize on the supplemented pyoverdines, especially at high concentrations via
the up-regulation of the heterologous promiscuous FpvB receptor[41,42]. Our findings
for *P. aeruginosa* are not unexpected and show that pyoverdine treatment should not
be applied against this pathogen.

Taken together, we found that all pyoverdines are highly potent against *A.*
baumannii and *S. aureus,* and one pyoverdine (3G07) is potent against *K.*
pneumoniae. Especially in the case of *A. baumannii* and *S. aureus,* the pyoverdine
dose-response curves follow the standard trajectory observed for antibiotics (Figure
S3 for ciprofloxacin). In support of our hypothesis that the inhibitory effects of
pyoverdines operate via withholding iron from pathogens, we found no growth

inhibition when the pathogens were exposed to purified ferribactin (Figure S3), the
non-iron chelating precursor of pyoverdine.

To further validate that the antibacterial effect of pyoverdine is due to iron
sequestration, we saturated pyoverdine s3b09 with varying concentrations of iron
and repeated the dose-response experiment with *A. baumannii* (Figure S4). Here,
we expect pyoverdine potency to decrease because iron is no longer a growth
limiting factor. Indeed, we observed that the inhibition of *A. baumannii* was greatly
reduced with intermediate levels of iron supplementation (40 µm) and completely
stalled with high levels of iron supplementation (200 µM) (Figure S4).

Figure 3. Pyoverdine dose-response curves for *A. baumannii, K. pneumoniae, P.
aeruginosa* and *S. aureus. We exposed the four human opportunistic pathogens to three
pyoverdines (3A06, 3G07, s3b09) that were among the most potent ones. We used crude-
purified extracts of all three pyoverdines and a HPLC-purified variant for pyoverdine s3b09.
The absolute concentrations of the crude-purified extracts are unknown and therefore
expressed relative to the weighed amount of 6 mg. The absolute concentration of the HPLC-
purified variant is given in µg/mL. Growth values are scaled relative to the untreated control
in CAA medium. Dots and error bars show mean values and standard errors, respectively,

across a minimum of three replicates per concentration. Dose-response curves were fitted
using 4- or 5-parameter logistic regressions.

Pyoverdines have low toxicity for mammalian cell lines, erythrocytes, and
hosts

We investigated the potential toxicity of crude-purified pyoverdines against mouse
neuroblastoma-spinal cord (NSC-34) hybrid cells and human embryonic kidney 293
(HEK-293) cells. NSC-34 are motor neuron-like cells, which do not divide in their
differentiated state, thus mimicking established tissue. For these cells, we observed
no adverse effects on cell viability at low and intermediate pyoverdine concentrations
(Figure 4A), representing the doses that strongly inhibited the pathogens (Figure 3).
A moderate decrease of cell viability was only observed at the highest pyoverdine
concentrations (Figure 4A). Conversely, HEK-293 cells divide rapidly and inform us
on whether pyoverdines interfere with cell proliferation. We indeed noticed a more
pronounced effect on cell viability already at intermediate pyoverdine concentrations
(Figure 4A), which suggests that pyoverdine chelates the iron required for cell
proliferation.

To assess the haemolytic activity of pyoverdines, we exposed sheep
erythrocytes to a concentration gradient of the crude-purified pyoverdines. We found
that haemolysis was extremely low (<1%) relative to the positive control Triton X-100
surfactant for pyoverdine concentrations up to 0.5 (Figure 4B) and remained low
even for the highest concentration (haemolysis rates for pyoverdines 3A06, 3G07
and s3b09 were 2.01%, 1.89% and 5.92%, respectively). This result shows that
pyoverdines are unable to retrieve iron from haemoglobin.

Finally, we injected pyoverdine into the larvae of *G. mellonella* and followed
host survival over time. We selected three pyoverdine concentrations (0.01, 0.05,
and 0.1), which we later used for the infection experiments, and which were
minimally toxic to cell lines (coloured dots in Figure 4A). We found that pyoverdine
injections did not significantly reduce host survival at low and intermediate
concentrations (Figure 4C and Table S1). In contrast, we observed a tendency
towards higher larval mortality levels with the highest pyoverdine concentration
administered, especially for 3G07, suggesting that pyoverdines can be mildly toxic
for the host at high concentrations.

Figure 4. Toxicity assays for pyoverdines from environmental *Pseudomonas* spp.
against human cell lines, sheep erythrocytes and the host larvae of *G. mellonella*. (A)
We exposed mouse neuroblastoma-spinal cord (NSC-34) and human embryonic kidney 293
(HEK-293) cells to three crude-purified pyoverdines (3A06, 3G07, s3b09) that were among
the most potent ones to inhibit bacterial growth. An MTT assay was used to assess the
metabolic activity of cells as an indicator of cell viability and proliferation. Cell viability data
are scaled relative to the pyoverdine-free treatment, whereby dots and error bars show
means and standard error across three replicates, respectively. The absolute concentrations
of the crude-purified pyoverdines are unknown and concentrations are therefore expressed
relative to the highest one used. Coloured dots indicate pyoverdine dosages used for the *in*
vivo experiments (pyoverdines/10 µL). Dose-response curves were fitted using 5-parameter
logistic regressions. (B) We evaluated the haemolytic activity of the pyoverdines by adding
them to sheep erythrocytes along a concentration gradient (range 0.002 – 1). Triton X-100
and PBS served as positive and negative control, respectively. Haemolytic activity is scaled

relative to the positive control, dots and error bars show means and standard error across 6

replicates from two independent experiments, respectively. (C) To assess the toxic effects of

pyoverdine on the host, we injected pyoverdines (three relative concentrations, 0.01, 0.05,

0.1) into larvae 4 hours after a mock infection with PBS. The percentage of larval survival

was tracked over 48 hours post-treatment. Data stem from three independent experiments

with each 10 larvae per infection and treatment.

Pyoverdine treatment increases the survival of infected hosts

We then investigated whether pyoverdines can be used to treat infections in *G.*

mellonella. We used *A. baumannii*, *K. pneumoniae,* and *P. aeruginosa* PAO1 as

pathogens. We did not include *S. aureus* for these experiments as it is literally

avirulent for *G. mellonella* given the infection loads and time frames used[43]. We first

determined the bacterial infection load needed to reduce larval survival by at least

50% (Figure S5) over 48 hours. We found this to be the case with $1.8*10^5$

CFU/larvae of *A. baumannii* and $8.9*10^5$ CFU/larvae of *K. pneumoniae*. For *P.*

aeruginosa, all larvae died within 48 hours even with the lowest infection dose and

we decided to use 56 CFU/larvae. We infected the larvae with one of the three

pathogens, and subsequently treated them with either pyoverdine (relative

concentrations 0.01, 0.05 and 0.1) or a PBS control four hours post infection,

followed by monitoring larval survival over time (Figure 5 and Figure S6).

352   We found that the two pyoverdines 3A06 and 3G07 were effective treatments

against *A. baumannii* and *K. pneumoniae* and significantly decreased the risk of

death compared to untreated infections (Figure 5, Table S2). Specifically, *A.*

baumannii treated with 3A06 at a concentration of 0.05 and 3G07 at a concentration

of 0.01 reduced the risk of death (hazard) by 57.9% (z = -1.973, P = 0.0485) and

59.4% (z = -2.057, p = 0.0397), respectively. For s3b09, the same tendency was

observed (z = -1.496, p = 0.135) albeit not significant. For *K. pneumoniae*, the

pyoverdine treatments 3A06 and 3G07 (concentration of 0.01) reduced the risk of

death by 51.5% (z = -2.027, p = 0.0426) and 57.6% (z = -2.341, p = 0.0192),

respectively. These results suggest that low and intermediate doses of pyoverdine

are more effective than the highest concentration. This notion was further supported

when analysing the time-to-death of larvae (Figure S6). We found hump-shaped

relationships between pyoverdine concentrations and time-to-death in four out of six

cases with two of them being significant (quadratic regression models for *K.*

pneumoniae treated with pyoverdine 3G07: linear term: $F_{1,71}$ = 0.88, p = 0.3506,

quadratic term: $F_{1,71}$ = 6.11, p = 0.0159 and s3b09: linear term: $F_{1,73}$ = 0.31, p =

0.5806, quadratic term: $F_{1,73}$ = 6.16, p = 0.0153). In contrast, infections with *P.*

aeruginosa led to near 100% larval mortality regardless of pyoverdine treatment and

dosage so that no significant treatment effect arose (Figure 5) and no effect on the

time-to-death of larvae could be observed (Figure S6).

Altogether, our infection experiments show that pyoverdines are effective

against the moderately virulent pathogens *A. baumannii* and *K. pneumoniae* by

decreasing the risk of death by more than 50%, while they are ineffective against the

fast-killing highly virulent *P. aeruginosa*. Moreover, intermediate pyoverdine

concentration are most effective, reinforcing our observation that pyoverdine can

exhibit some toxicity at higher concentrations (Figure 4C).

Figure 5. Pyoverdine treatments significantly increase survival of the host *G.

***mellonella* when infected with *A. baumannii* and *K. pneumoniae*.** Larvae of the greater

wax moth were first infected with either *A. baumannii*, *K. pneumoniae*, or *P. aeruginosa* and

then treated with one of three pyoverdines (3A06, 3G07, or s3b09) at four relative

pyoverdine concentrations (0, 0.01, 0.05, 0.1). All panels show the larvae survival over 48

hours post-treatment. Data stem from three independent experiments with each 10 larvae per infection and treatment. Asterisks indicate significant differences in larval survival between treated and untreated infections based on Cox proportional hazard regressions ($p < 0.05$).

Pathogens show low levels of resistance evolution against pyoverdine treatment

We assessed the ability of *A. baumannii*, *K. pneumoniae*, *P. aeruginosa* and *S. aureus* to evolve resistance against the pyoverdine treatment by exposing all pathogens to both single pyoverdine treatments (3A06, 3G07, and s3b09) and combination treatments (double and triple combinations) over 16 days of experimental evolution. One exception was *K. pneumoniae*, which was only exposed to 3G07 as only this pyoverdine was inhibitory. We had six independent lineages (populations) per pathogen and treatment combination (see Table S2 for treatment concentrations). We further subjected all pathogens to the antibiotic ciprofloxacin as positive control in which resistance evolution is expected. Additionally, we let the pathogens grow in untreated growth medium to control for adaptation to the growth medium. Overall, we had 180 lineages that were transferred daily to fresh medium.

After experimental evolution, we observed that all four pathogens grew significantly better under ciprofloxacin treatment than the ancestor and beyond the level observed for growth medium adaptation (Table S3; Figure 6A and S9). This provides strong evidence for pervasive resistance evolution against this conventional antibiotic. In contrast, we found that none of the four pathogens treated with pyoverdines experienced a significant increase in growth relative to the observed level of growth medium adaptation (Table S3, Figure 6A), suggesting low potentials of resistance evolution at the population level.

However, population screens can hide the presence of resistant clones within populations. We thus picked eight random clones from two populations for each of the pyoverdine single treatments and one combo treatment for each pathogen. With a total of 208 clones, we repeated the above growth assay and expressed clonal growth as fold-change relative to the growth of populations evolved in the growth medium alone (Figure 6B and S10). For *A. baumannii* and *K. pneumoniae*, the average growth fold-change across clones was 1.02 ± 0.004 and 0.84 ± 0.02,

respectively, and there was no sign of resistant clones being present in populations.
This was different for *P. aeruginosa* and *S. aureus*, for which we observed high
variation in growth fold-changes across clones, with many clones growing
moderately better than the growth medium controls. The growth increase across
clones was 1.11 ± 0.02 and 1.10 ± 0.02, and significant in three and two population
of *P. aeruginosa* and *S. aureus*, respectively. Thus, the clonal screen indicates some
level of adaptation to pyoverdine treatment for these two pathogens.
425

Mutational and functional patterns in response to pyoverdine treatment are
species specific
To obtain insights into the genetic basis of putative resistance evolution, we
sequenced the whole genomes of 52 evolved clones (16 per pathogen, except 4 for
K. pneumoniae), 16 medium-evolved control populations and the 4 ancestors. We
first mapped reads to the reference genomes and then to our ancestors. Among the
clones evolved under pyoverdine treatment, we identified 97 synonymous, 96
nonsynonymous, 59 intergenic and 9 nonsense single-nucleotide polymorphisms
(SNPs). Additionally, there were 48 insertions, 59 deletions and 45 substitutions
(Figure 6B). We observed that growth peaked in clones with a single mutation and
decreased in clones with multiple mutations (Figure 6C, linear fit: $F_{1,50} = 6.57$, p =
0.0135), suggesting positive selection for a few specific advantageous mutations.
Next, we excluded all mutations that were present at least twice in the no-
treatment controls, and all intergenic mutations. We identified seven mutations in *A.*
baumannii, one in *K. pneumoniae*, ten in *P. aeruginosa*, and 20 in *S. aureus* that
exclusively arose under pyoverdine treatment and could thus be associated with
resistance evolution. We then classified the SNPs and small deletions (<600 bp) into
functional categories (Figure 6D, Table S4). We found that mutations in genes
related to regulation and metabolism were consistently associated with increased
growth in *S. aureus*, whereas mutations in genes related to biofilm and motility were
associated with increased growth in *P. aeruginosa* (Figure 6D).
Furthermore, we identified six large deletions (≥600 bp; Table S5). One *K.*
pneumoniae clone lost a ~42 kb plasmid encoding various functional proteins, such
as methyltransferases and toxin-antitoxin systems[44]. Four *S. aureus* clones from two
independent populations showed parallel loss of the ~14 kb pathogenicity island
SaPI5, containing the virulence genes *sek* and *seq*[45,46]. Additionally, one *S. aureus*

clone exhibited a ~55 kb deletion containing the staphylococcal cassette
chromosome mec type IV (SCCmecIV), which harbours the *mecA* gene responsible
for β-lactam antibiotic resistance, and the arginine catabolic mobile element
(ACME)[45].

Taken together, we found that the observed weak resistance evolution against
pyoverdine in *P. aeruginosa* and *S. aureus* was not linked to mechanisms directly
enhancing the pathogen's iron acquisition but with mutations in global regulators,
metabolism, biofilm and motility genes. In *A. baumanii* and *K. pneumoniae*, no
evidence for resistance evolution was observed. Moreover, certain pathogens lost
mobile genetic elements containing virulence and antibiotic resistance genes,
suggesting that pyoverdine treatment could selects for less virulent and antibiotic
sensitive strains.

Figure 6. Phenotypic and genotypic analysis of experimentally evolved pathogens
reveal weak levels of resistance evolution against pyoverdine treatment. (A) We
exposed evolved and ancestral pathogen populations to the treatment in which they evolved
in and quantified their growth (area under the curve). Growth values were scaled relative to
the ancestor in untreated medium, and the panels show the scaled growth differences
between evolved and ancestral populations. The shaded areas show the scaled growth
difference between the ancestor and the populations evolved in untreated medium and is
representative of medium adaptation. The blue and green dots represent the pathogen
populations with the lowest (population 1) and highest (population 2) scaled growth
difference, respectively, which were subsequently used to pick clones. The dots show mean
values across two independent replicates and asterisks show significant growth increases
relative to the medium-adapted control. Box plots show the median and the first and third

quartiles across the 6 independently evolved populations. Whiskers represent the 1.5x
interquartile range. (B) We repeated the above growth assays with 208 individual clones
evolved under the pyoverdine treatments (picked from population 1 and population 2) and 24
populations evolved in growth medium alone (control). Each square represents a clone or a
control population, and the number indicates the number of mutations identified based on
whole-genome sequencing. The heatmap shows the fold-change in growth relative to the
evolved control populations. Asterisks depict significant fold-increases in growth compared
to control populations. (C) The relationship between the number of mutations and fold-
change in growth across all sequenced clones (n = 52) The red line and shaded area are the
regression line and 95% confidence interval, respectively. (D) Heatmap showing the number
of mutations per pathogen and per functional gene categories across clones together with
the respective fold-change in growth (n = 32). For this analysis, we excluded intergenic
mutations and deletions larger than 600 bp. Triangle, square and cross depict growth fold-
change of *A. baumannii*, *P. aeruginosa* and *S. aureus*, respectively.
492

Discussion

We investigated whether pyoverdines produced by environmental *Pseudomonas*
spp. have antibacterial activities against human opportunistic pathogens through iron
sequestration and withholding this trace element from pathogens. Our screen
involving 320 environmental isolates and 12 pathogen strains revealed five top
pyoverdine candidates with broad-spectrum activity. These candidates could be
distinguished from non-inhibitory pyoverdines by their high CCS values (standing for
larger and more complex molecules) and their higher iron complexation stability.
Experiments with crude- and HPLC-purified pyoverdines showed that pyoverdines
completely stall the growth of *A. baumannii* and *S. aureus*, while showing
intermediate activity against *K. pneumoniae* and *P. aeruginosa*. When administered
as treatment to infected *G. mellonella* larvae, we observed significantly increased
host survival rates in infections with *A. baumannii* and *K. pneumoniae*,
demonstrating that pyoverdine exhibits antibacterial activity in hosts. Furthermore,
we found low toxicity of pyoverdines at effective concentrations and observed low
potentials for resistance evolution. Overall, our results reveal pyoverdines from non-
pathogenic *Pseudomonas* spp. as potent antibacterials against several human
opportunistic pathogens.

Our novel treatment approach assumes that pyoverdines sequester iron and
thereby induce iron starvation and growth arrest in pathogens. Several of our
findings support this view. First, the growth-inhibiting effect of our top pyoverdine
candidates exclusively occurred under iron-limited conditions, while pathogen growth
was restored under iron-replete conditions (Figure 1, Figure S4). Second, non-
inhibitory ferri-pyoverdines were more prone towards collision induced unfolding than
inhibitory ferric-pyoverdines (Figure 2D), suggesting that inhibitory pyoverdines have
a higher iron binding affinity. Third, the pyoverdine precursor ferribactin did not inhibit
pathogen growth (Figure S3). The two molecules are identical apart from one ring in
the chromophore core that is not cyclized and not oxidized in ferribactin, which
compromises the binding of iron. These findings suggest that pyoverdine does not
target the pathogens directly, but rather indirectly via the induction of iron starvation.

We observed that pyoverdines showed pathogen-specific efficacies in curbing
bacterial growth (Figure 3). We propose that differences can be explained by the
various strategies of these pathogens to cope with iron stress. *A. baumannii* and *S.*
aureus produce siderophores that have a much simpler chemical structures than
pyoverdines and are expected to bind iron with lower affinity (*A. baumannii*:
acinetobactin (association constant: 10^{26} M^{-1}), fimsbactin (10^{27} M^{-1}) and
baumannoferrin (unknown)[47–49], *S. aureus*: staphyloferrin A+B (unknown)[50]) than
pyoverdine (10^{32} M^{-1})[28]. Hence, the growth suppression of *A. baumannii* and *S.*
aureus most likely occurs because their siderophores are too weak to successfully
compete with pyoverdine. The situation is different for *K. pneumoniae*, which
produces enterobactin, the siderophore with the highest known iron affinity (10^{52} M^{-1}
)[39]. Consistent with the concept of iron competition, we observed that pyoverdine
treatment is less effective against *K. pneumoniae*, and was only potent for one out of
three pyoverdines (3G07, Figure. 3). Finally, pyoverdine treatment is not expected to
work well against *P. aeruginosa*, which itself produces a version of this siderophore.
The two competing pyoverdines likely have similar iron affinities and the presence of
pyoverdine receptors in the pathogen might foster the uptake of the supplemented
pyoverdine. Our results indeed support the view that pyoverdine treatment is not
effective against *P. aeruginosa*. Taken together, we predict that pyoverdine
treatment could be highly potent against certain pathogens like *A. baumannii* and *S.*
aureus that have less efficient siderophores and are unable to use the pyoverdine as
iron source.

For every new treatment, it is important to consider whether it has
unintentional consequences for the targeted pathogen. Because our approach does
not kill pathogens directly, there are multiple ways of how pathogens may react to
pyoverdine treatment. For example, pathogens could increase the production of their
own siderophores in response to the severe iron limitation induced by pyoverdines.
Alternatively, siderophores can serve as signalling molecules in certain species like
P. aeruginosa, regulating the expression of additional virulence factors like proteases
and toxins[51]. While pyoverdine treatment may affect such signalling cascades in
other pathogens as well, and thus potentially increase virulence factor production,
we observed the opposite in *S. aureus*, which showed a loss of virulence genes.
Similarly, iron limitation can increase[52] or decrease[53–55] the formation of biofilms,
depending on the species or even strains[56,57] and it is well possible that pyoverdine
treatment thus affects the propensity of pathogens to form biofilms. Finally,
pyoverdines may have differential effects on the various members of the microbiota
or the different pathogens in polymicrobial infections and thereby induces shifts in
species composition due to the unequal suppression of community members. In
complex communities, the inhibitory effect of pyoverdine on the pathogen may get
diluted as the burden of increased iron limitation is shared across community
members. All these considerations show that it will be crucial to assess the
consequences of pyoverdine treatments in more complex settings including hosts.

Not only bacteria but also host cells need iron. Consequently, pyoverdine
treatment could negatively affect host iron homeostasis. Our toxicity assays reveal
that there is a range of pyoverdine concentrations for which only mild adverse effects
against cell lines and *G. mellonella* larvae are observed and no haemolysis occurs.
Especially the finding that pyoverdines have no negative effects on the survival of
non-replicating NSC-34 cells is encouraging, as this cell line is representative of
intact host tissue, in which iron is bound to strong chelators such as transferrin,
lactoferrin or ferritin[58,59]. Additionally, recent research has demonstrated that purified
pyoverdine remains non-toxic even after 72 hours of treatment at concentrations up
to 200 μM[60]. The result that pyoverdines are unable to obtain iron from haemoglobin
is encouraging, too. Previous work showed that the pyoverdine of *P. aeruginosa* can
retrieve iron from human transferrin when working in concert with proteases
(proteolytically degrading transferrin)[61] or phenazines (spontaneously reducing ferric
to ferrous iron)[62]. However, these *P. aeruginosa* specific mechanisms require the

presence of this pathogen[63], which is not the case when purified pyoverdines are
used as a treatment against other pathogens like *A. baumannii*. Nonetheless, we
found that pyoverdines can have certain adverse effects, especially for the rapidly
proliferating HEK293 cells (Figure 4). These cells represent a regenerating tissue
and thus have high iron demands. Similarly, we observed mild adverse effects for
some pyoverdines (3G07) when administered at high concentrations to *G. mellonella*
larvae (Figure 5). These findings corroborate the results from previous studies on
Caenorhabditis elegans, where pyoverdines were found to interfere with host iron
homeostasis[64,65]. These considerations highlight that finding the right pyoverdine
concentration will be key to meet the fine line between maximizing pathogen
inhibition and minimizing tissue damage. By contrast, the adverse host effects of
pyoverdines might come with additional opportunities, for instance in cancer therapy,
where siderophores reduce iron levels in tumours and slow tumour progression[66].

We propose that pyoverdine treatment could synergistically interact with the
host innate immune system. Mammalian neutrophils produce siderocalin to
sequester bacterial siderophore-iron complexes, rendering them non-functional[58].
While this strategy is efficient against many siderophores, some bacteria have
evolved so-called stealth siderophores that no longer bind to siderocalin.
Pyoverdines are such stealth siderophores[67]. Regarding pyoverdine treatment, this
means that pathogens with regular siderophores will hit a double wall: their
siderophores are immobilized by siderocalins, while pyoverdines lock away the
remaining iron.

Another promising finding of our work is that the potential for resistance
evolution against pyoverdines seems low. Conceptually, one reason for low
resistance evolution could be that pyoverdines, unlike many antibiotics, are not
internalized into bacterial cells. Thus, classic resistance mechanisms such as
reduced drug influx, increased drug efflux, and intra-cellular target modification
cannot operate[68]. We initially expected that resistance evolution would involve
mechanisms that improve the pathogen's iron acquisition. This could include the up-
regulation of the pathogen's own siderophores or switching from siderophore-based
ferric iron acquisition to the use of reductases that foster ferrous iron uptake[68].
However, our sequencing analysis did not reveal mutations in genes associated with
iron uptake systems. One explanation is that such mutations (albeit beneficial) might
not reach high frequencies in populations because increased siderophore production

and secretion would increase the iron acquisition of resistant and non-resistant cells
alike[68]. We instead observed an overrepresentation of mutations in regulatory,
biofilm and motility, and metabolic genes. While we do not know the phenotypes of
these mutations, one possibility is that increased biofilm formation could offer some
protection against pyoverdine treatment, for example by restricting pyoverdine from
chelating free iron. Important to note is also that patterns of resistance evolution may
differ between *in vitro* and *in vivo* settings[4,69]. For example, our experimental
evolution setup did not allow for horizontal transfer of siderophore receptors, a
scenario that might occur in the context of polymicrobial infections and confer
resistance to pyoverdine therapy.

In conclusion, our treatment approach using pyoverdine is based on the
principle of paying like with like. Pathogens secrete siderophores to scavenge iron
for their growth. We add a strong heterologous siderophore as treatment to withhold
iron from pathogens to curb their growth. The concept of targeting bacterial
infections with both natural and synthetic iron-chelators has been highlighted in the
literature for quite some time[70–74]. For instance, the siderophore desferoxamine
(produced by *Streptomyces* species) is used in clinical settings to treat iron
overload[75]. However, its efficacy as antibacterial is limited because of its low iron
affinity (10^{27} M^{-1}, compared to pyoverdine and enterobactin)[76] and several pathogens
have receptors for deferoxamine uptake and can thus use it as an iron source[77].
Synthetic iron chelators like deferiprone and deferasirox have been proposed as
alternatives to overcome these limitations. However, the minimum inhibitory
concentration (MIC) of deferiprone needed to inhibit *P. aeruginosa* was found to be
up to 36.7 times higher than the plasma levels in human therapy, raising concerns
about its cytotoxicity and clinical viability for treating bacterial infections[78,79].
Furthermore, at sub-MIC levels, deferiprone only partially suppressed the growth of
P. aeruginosa and *A. baumannii,* indicating insufficient iron sequestration from these
pathogens[79,80]. Our approach offers a distinct advantage by utilizing naturally
evolved iron chelators from non-pathogenic species, which possess unique chemical
structures and exceptionally high iron affinities. These properties reduce the
likelihood of their exploitation as iron sources by pathogens and provide a
competitive advantage over siderophores with lower iron affinities. Combined with
low toxicity to hosts at therapeutically effective concentrations and minimal potential
for resistance evolution, these natural chelators hold promise as a new class of

effective antibacterials, either as standalone treatments or in combination with
conventional antibiotics.

Material and methods

Bacterial strains

We used a strain collection of 320 natural isolates, originating from soil and pond
habitats. The isolates originate from 16 independent soil and water samples (20
isolates per sample) collected on the Campus Irchel Park of the University of Zurich.
In previous work, we have extensively studied and characterized all isolates[16,29–31].
Details on the isolates can be found in the Supporting Information.

For the supernatant screening assay, we used a collection of 12 strains of
opportunistic human pathogens (Table 1), including *A. johnsonii*, *A. junii*, *C.*
sakazakii, *K. michiganensis* (belonging to the *K. oxytoca* complex)[81] and *Shigella* sp.
(all provided by the laboratory of Leo Eberl, University of Zurich), and *B.*
cenocepacia strains H111 and K56-2, *P. aeruginosa* strains PA14 and PAO1, *E. coli*
K12 and *S. aureus* strains Cowan and JE2 from our own strain collection. For dose-
response and infection experiments, we used *A. baumannii* (DSM 30007) and *K.*
pneumoniae (DSM 30104), both purchased from the German Collection of
Microorganisms and Cell Cultures GmbH, in addition to *P. aeruginosa* PAO1 and *S.*
aureus JE2.

Culturing conditions

Pathogen overnight cultures were grown in either 8 mL tryptic soy broth (TSB; for the
two *S. aureus* strains) or 8 mL lysogeny broth (LB; all other pathogens) in 50 mL
tubes at 37 °C and 220 rpm agitation. Following two washing steps with 0.8% NaCl,
we adjusted the overnight cultures to an optical density at 600 nm (OD_{600}) of 1. For
overnights of environmental isolates, we followed the same procedure when working
with low sample sizes, with the only difference being that culturing occurred at 28 °C.
For the large-scale screening experiments, we grew environmental isolates in 200
μL LB in 96-well plates shaken at 170 rpm. Cultures from plates were then directly
used for experiments. For all main experiments (screen for bioactive pyoverdines
and dose-response curves), we used casamino acid (CAA) medium (1% casamino

acids, 5 mM $K_2HPO_4 * 3H_2O$, 1 mM $MgSO_4 * 7 H_2O$, 25 mM HEPES buffer). This

medium contains low levels of iron, triggering pyoverdine production[82].

Supernatant screening assays

We created sterile supernatants from all the 320 *Pseudomonas* isolates. Specifically,

we transferred 2 μL of overnight cultures (grown in 96-well plates in LB in four-fold

replication for 24 hours as described above) to new plates containing 200 μL of CAA

medium supplemented with 250 μM 2,2'-bipyridyl. Cultures were incubated for 24

hours and subsequently centrifuged at 2250*g* for 10 minutes. The supernatants were

transferred to 0.2 μm membrane filter plates (PALL AcroPrep Advance), centrifuged

a second time, and frozen at -20 °C.

For the supernatant screening assay, the pathogens were cultured and diluted

as described above and added to 140 μL CAA medium in 96-well plates. We added

60 μL of thawed *Pseudomonas* supernatants (30%) or 0.8% NaCl as control and

incubated the plates at 37 °C and 170 rpm. Growth was measured as OD_{600} after 24

hours using a Tecan Infinite M-200 plate reader (Tecan Group, Männedorf,

Switzerland). We then scaled the growth of pathogens in the supernatant relative to

their growth in the control treatment. Relative growth values smaller and larger than

one indicate growth inhibition and promotion by the supernatant, respectively.

To test whether pyoverdine is responsible for pathogen growth reduction, we

repeated the screening assay but supplemented the CAA medium containing the

Pseudomonas supernatants with 40 μM $FeCl_3$. In this iron-rich condition, pyoverdine

should not be able to limit iron availability and thus pathogen growth should be

restored.

Structure elucidation of pyoverdine and analyses of chemical properties

From the seven top supernatant candidates that inhibited all twelve pathogens, we

extracted the pyoverdines and elucidated their structure. For this purpose, we have

developed a new protocol that allows structure elucidation from low volume

supernatants using UHPLC-HR-MS/MS. The methodological details are described in

Rehm *et al.*[27,34]. Afterwards, we assessed whether growth inhibiting pyoverdines can

be distinguished from non-inhibiting pyoverdines by their chemical properties. We

focussed on polarity, CCS values, and stability upon collision induced unfolding or

fragmentation. Detailed protocols and instrument settings can be found in the Supporting Information.

Crude pyoverdine purification

To test the efficacy of the pyoverdines against the four focal human opportunistic pathogens (*A. baumannii*, *K. pneumoniae*, *P. aeruginosa* (PAO1), and *S. aureus* (JE2)), and to assess their cytotoxicity against human cell lines, we crude-purified three pyoverdines (3A06, 3G07, s3b09). The three pyoverdines were among the five most potent ones identified in our supernatant screening assay. The adapted method for pyoverdine purification from the previous studies[16,35] are further described in the Supporting Information.

HPLC purification of pyoverdine and ferribactin

To further validate that pyoverdines are the compounds that induce pathogen growth inhibition, we purified the most common pyoverdine (s3b09) among the most potent pyoverdines from our screens using reversed-phase high-performance liquid chromatography (HPLC). Methodological details thereof are described in the Supporting Information.

Dose-response curves

To determine the inhibition potential of crude- and HPLC-purified pyoverdines, as well as ferribactin, we subjected *A. baumannii*. *K. pneumoniae*, *P. aeruginosa* and *S. aureus* to serially diluted extracts. For this, the extracts were dissolved in CAA medium, filter sterilized and diluted in the same medium. Bacterial overnight cultures were grown and prepared as described above and added to a total of 200 µL medium on a 96-well plate with 4 replicates per extract dilution. As positive control where growth inhibition is expected, we subjected all three pathogens to a dilution series of the antibiotic ciprofloxacin (highest concentration 4 µg/mL), in the exact same way as for the extracts. We incubated the plates at 37 °C statically in the plate reader and measured the OD_{600} every 15 minutes for 24 hours. Before each measurement, plates were shaken for 15 seconds. We repeated the experiment up to five times with different initial extract starting concentrations. This was done to ensure that the full dose response to pyoverdine was captured. Thus, sample size

per individual extract concentration varied between four and 20 replicates. We
subtracted the blank values and the background values caused by the treatment of
the respective extract concentration from the measured growth data and calculated
the integral (area under the growth curve) using the R package Growthcurver[83]. We
then expressed growth relative to the control medium, where bacteria grew in CAA
without extract or antibiotic addition. Finally, we plotted extract/antibiotic
concentration vs. relative growth and fitted dose-response curves (see below).

Cytotoxicity assays

To determine the cytotoxicity of pyoverdine treatments against mammalian cells, we
subjected human embryonic kidney 293 (HEK-293) and mouse motor neuron-like
neuroblastoma-spinal cord hybrid (NSC-34) cells to the crude extract of the three
most potent pyoverdines (3A06, 3G07, s3b09). Detailed culturing conditions can be
found in the Supporting Information. Briefly, approximately 10^4 cells/well were plated
on 96-well plates (Greiner Bio-One, 655090) and, after 48 hours, the culture medium
was replaced and the pyoverdines were added covering a range of concentrations.
After another 48 hours, the pyoverdine-containing culture medium was removed, and
cells were incubated with 1 µg/mL thiazolyl blue tetrazolium bromide (Sigma, D5655)
in 100 µL of fresh medium for 1.5 hours (NSC-34) or 40 minutes (HEK-293). The
reaction was stopped, and formazan crystals simultaneously solubilized with the
addition of 100 µl of a solution containing 10% sodium dodecyl sulfate (SDS; Sigma-
Aldrich, 05030) and 0.03% HCl (Supelco, 100319). Finally, the absorbance of cell
debris and other contaminants at 630 nm were subtracted from the absorbance of
the solubilized formazan at 570 nm.

Haemolysis assay

To determine the haemolytic activity of pyoverdines, we quantified sheep erythrocyte
haemolysis at different pyoverdine concentrations. We first centrifuged 25 mL of
fresh sheep blood (Chemie Brunschwig AG, Switzerland) in a 50 mL tube at 1500g
for 5 minutes at 4 °C. Plasma was then gently aspirated, replaced by PBS and tubes
were gently inverted to mix. This washing procedure was repeated three times.
Pyoverdines were diluted in PBS to final relative concentrations of 1, 0.5, 0.25, 0.2,
0.02, and 0.002. PBS and 0.1% Triton X-100 (Sigma Aldrich, Switzerland) served as

negative and positive control, respectively. 100 µL erythrocyte solution was mixed
with either 100 µL pyoverdine solution or the controls in a 96-well plate in triplicates.
Following an incubation at 37 °C and 120 rpm for 1 hour, the plate was centrifuged
at 1036g for 5 minutes at 4 °C. Subsequently, 100 µL of the supernatant were
transferred to a new 96-well plate. Haemoglobin release was measured at 570 nm in
the plate reader and the percent haemolysis was calculated relative to the positive
control (% haemolysis = (OD570 of sample − negative control) / (positive control −
negative control)).
787

Infection model

We tested the efficacy of pyoverdine to treat infections in larvae of the host model *G.*
mellonella. Detailed information on the larvae handling and infection procedure are
described in the Supporting Information. On the pathogen side, we determined the
LD$_{50}$ (lethal dose, killing 50% of the larvae) for *A. baumannii* (1.8 * 10^5 CFU/larvae)
and *K. pneumoniae* (8.9*10^5 CFU/larvae) (Figure S5). For *P. aeruginosa*, we use 56
CFU/larvae because all doses tested killed 100% of the larvae (Figure S5). As
negative control, we administered 10 µL/larvae of sterile PBS. Following infections,
larvae were individually placed in wells of 24-well plates and incubated at 37 °C in
the dark for 4 hours.
On the treatment side, we used pyoverdines 3A06, 3G07, and s3b09 at three
concentrations that were non-toxic for the NSC-34 cells. For this, we prepared stocks
of 60 mg/mL, 30 mg/mL, and 6 mg/mL of crude pyoverdines. Since the injected
treatment volume was 10 µL per larvae, the final pyoverdine concentrations were 0.6
mg, 0.3 mg, and 0.06 mg per larvae, respectively. When scaling theses
concentrations to the doses used for the *in vivo* dose-response curves (Figure 4), the
relative pyoverdine concentrations were 0.1, 0.05 and 0.01. Pyoverdine treatments
were administered 4 hours after the infection through injections (see Supporting
Information). Prior to treatment, larvae were immobilized on ice for 15 minutes. We
also applied treatments to larvae infected with sterile PBS to quantify potential
negative effects of the treatment in the absence of an infection. Following treatment,
larvae were put back to their allocated well of the 24-well plate and kept at 37 °C in
the dark.

From the 12th hour post infection (hpi) onwards, we regularly checked the

survival of all larvae for a total of 48 hpi. Specifically, larvae were poked with a

pipette tip and individuals not responding to the physical stimulation were considered

dead. We conducted three independent experiments per pathogen-treatment

combination with 10 larvae per experiment, resulting in a total of 30 larvae per

pathogen-treatment combination. Larvae in their last instar stage were purchased

from a local vendor (Bait Express GmbH, Basel, Switzerland) and different batches

of larvae were used for the three independent experiments.

Experimental evolution experiment

We first determined the IC50 concentrations of (crude) pyoverdines 3A06, 3G07, and

s3b09, and the antibiotic ciprofloxacin against all four pathogens. For this, we fitted

logistic regressions to the dose-response curves. We used these IC50 values as

treatment concentrations during experimental evolution (Table S2). We also mixed

different pyoverdines in double and triple combo treatments. In combo treatments,

pyoverdine concentrations were equivalent to those of the single treatments,

meaning that we divided the single treatment concentrations by two or three. The

experimental evolution experiment included (i) three single pyoverdine treatments

(3A06, 3G07, s3b09), (ii) three double pyoverdine treatments (3A06 and 3G07, 3A06

and s3b09, 3G07 and s3b09), (iii) one triple pyoverdine treatment (3A06, 3G07 and

s3b09), and (iv) one ciprofloxacin treatment for *A. baumannii*, *P. aeruginosa* and *S.*

aureus. For *K. pneumoniae*, which was only affected by one pyoverdine (3G07), the

experiment consisted of one single pyoverdine treatment and one ciprofloxacin

treatment.

A detailed description of the experimental evolution setup is provided in the

Supporting Information. Briefly, for each pathogen-treatment combination, we

included six independently replicated lineages (populations) on 96-well plates. Each

plate further contained control populations that evolved in plain CAA medium. At the

start, we grew the pathogens overnight, washed and diluted them as described

above, and transferred 2 µL of the diluted pathogens into 200 µL medium in 96-well

plates. Subsequently, the plates were incubated at 37 °C for 22 hours, shaken at

170 rpm. After incubation, we measured growth (OD_{600}) of all evolving lineages with

the plate reader. We then diluted the evolving cultures 1:1,000 in CAA medium and

added them to a fresh treatment plate. After the transfer, the new plates were again
incubated at 37 °C and 170 rpm agitation for 22 hours. We added glycerol to the old
plates to a final concentration of 25% and stored them at -80 °C.

Phenotypic screening of evolved populations
We screened evolved populations for improved growth performance relative to the
ancestor under pyoverdine treatment, which we took as a proxy for resistance
evolution. For this purpose, we thawed the final plates (day 16) of the experimental
evolution experiment and transferred 2 µl of all populations to 200 µl of LB or TSB
(for *S aureus*) to create overnight cultures in 96-well plates. We further included
ancestor cultures of all four species. After incubation for 18 hours at 37 °C and 170
rpm, we diluted overnights 1:1,000 with 0.8% NaCl and subjected all evolved
populations to the conditions they had evolved in. Ancestors were simultaneously
exposed to the respective treatment conditions. The culturing volume and conditions
matched those used during experimental evolution, but this time we incubated the
plates in a plate reader, measuring OD_{600} every 15 minutes to obtain growth kinetics.
The population screen was repeated twice independently. From the growth kinetics,
we first calculated the area under the curve (AUC) for each growth curve. AUC
values were then scaled relative to the growth of the ancestor in the untreated CAA
control medium. For each evolved population, we then calculated the difference in
growth in comparison to the mean growth of the ancestor subjected to the same
treatment.

Phenotypic screening of evolved clones
To screen for resistant clones within populations, we selected the pathogen
populations with the lowest and highest growth difference for each single treatment
and for one combination treatment and picked 8 random colonies. We transferred
them to 200 µL LB or TSB (*S. aureus*) in 96-well plates and incubated plates for 18
hours at 37 °C (170 rpm) to create overnight cultures. We followed the same
protocol as above for quantifying the difference in growth of evolved and ancestral
clones.

Genomic analyses of evolved populations and clones

We sequenced the genomes of the four ancestors, four control populations that had evolved in medium without pyoverdine treatment, and two random clones (out of the eight picked) per population that had received pyoverdine treatment and were used for clonal growth analysis (Figure 6B). The Supporting Information features a detailed description of the sequencing and data analysis protocols used. Variants were predicted relative to the reference genomes ATCC 19606 (*A. baumannii*), ATCC 13883 (*K. pneumonia*), ATCC 15692 (*P. aeruginosa*) and CP020619.1 (*S. aureus*) using the breseq pipeline[84]. In the pipeline, we specified that medium-adapted controls were treated as populations, whereas the treatment-adapted colonies picked for sequencing were treated as clones.

Data analyses

All statistical analyses were performed in R 4.0.2 and RStudio version 1.3.1056. We conducted Shapiro-Wilk tests and consulted diagnostic plots to check whether our data (residuals) were normally distributed. For normally distributed data sets, we used linear models and two-sided tests for all analyses. For non-normally distributed data sets, we used non-parametric rank tests. We used ANOVA to test whether chemical properties differ between growth inhibiting and non-inhibiting pyoverdines, in their ferri- versus apo-pyoverdine state. For the *in vitro* dose-response and cytotoxicity experiments, we fitted dose-response curves with either 4- or 5-parameter logistic regressions using the nplr package[85]. Survival analyses were performed using the Cox proportional hazards regression model or the log-rank test (adjusted for multiple comparisons using the Holm method) from the survival package[86]. We used two-sided t-tests, Welch's t-tests, or Wilcoxon tests to compare growth differences between the ancestor and the evolved populations under treatment. To compare growth differences between the ancestor and the evolved clones under treatment, we used one- and two-way ANOVAs. To construct the cladogram in Figure S2, we followed the procedure described in Kramer et al[31] using partial *rpoD* gene sequences with lengths ≥600 bp. We used iTOL tool[87] for the graphic presentation of the cladogram and to link phylogeny to the supernatant effects.

Acknowledgements

We thank Richard Allen, Dominik Bär, Johanna Giger, Kevin Schiefelbein and Julien Weber for laboratory assistance, and Markus Seeger for insightful discussion. This work was supported by funding from the Swiss National Science Foundation (grant numbers: 31003A_182499 to RK, PP00P3_144862 and PP00P3_176966 to MP) and a Candoc Grant (Forschungskredit) from the University of Zurich (to MPB).

Authors contributions

VV and RK conceived the project. VV, KR, CC and MPB performed the experiments and analysed experimental results. All authored interpreted the data. VV and RK wrote the manuscript with contributions from all authors.

Competing interests

Authors declare no competing interests.

Data availability

The raw data underlying the figures are available from the Figshare depository (https://doi.org/10.6084/m9.figshare.26388565). The sequencing data for this study have been deposited in the European Nucleotide Archive (ENA) at EMBL-EBI under accession number PRJEB78468 (https://www.ebi.ac.uk/ena/browser/view/PRJEB78468).

References

1. Murray, C. J., Ikuta, K. S., Sharara, F., Swetschinski, L., Aguilar, G. R., Gray, A., Han, C., Bisignano, C., Rao, P., Wool, E., Johnson, S. C., Browne, A. J., Chipeta, M. G., Fell, F., Hackett, S., Haines-Woodhouse, G., Hamadani, B. H. K., Kumaran, E. A. P., McManigal, B., Agarwal, R., Akech, S., Albertson, S., Amuasi, J., Andrews, J., Aravkin, A., Ashley, E., Bailey, F., Baker, S., Basnyat, B., Bekker, A., Bender, R., Bethou, A., Bielicki, J., Boonkasidecha, S., Bukosia, J., Carvalheiro, C., Castañeda-Orjuela, C., Chansamouth, V., Chaurasia, S., Chiurchiù, S., Chowdhury, F., Cook, A. J., Cooper, B., Cressey, T. R., Criollo-Mora, E., Cunningham, M., Darboe, S., Day, N. P. J., Luca, M. D., Dokova, K.,

Dramowski, A., Dunachie, S. J., Eckmanns, T., Eibach, D., Emami, A., Feasey,
N., Fisher-Pearson, N., Forrest, K., Garrett, D., Gastmeier, P., Giref, A. Z., Greer,
R. C., Gupta, V., Haller, S., Haselbeck, A., Hay, S. I., Holm, M., Hopkins, S.,
Iregbu, K. C., Jacobs, J., Jarovsky, D., Javanmardi, F., Khorana, M., Kissoon, N.,
Kobeissi, E., Kostyanev, T., Krapp, F., Krumkamp, R., Kumar, A., Kyu, H. H.,
Lim, C., Limmathurotsakul, D., Loftus, M. J., Lunn, M., Ma, J., Mturi, N., Munera-
Huertas, T., Musicha, P., Mussi-Pinhata, M. M., Nakamura, T., Nanavati, R.,
Nangia, S., Newton, P., Ngoun, C., Novotney, A., Nwakanma, D., Obiero, C. W.,
Olivas-Martinez, A., Olliaro, P., Ooko, E., Ortiz-Brizuela, E., Peleg, A. Y.,
Perrone, C., Plakkal, N., Ponce-de-Leon, A., Raad, M., Ramdin, T., Riddell, A.,
Roberts, T., Robotham, J. V., Roca, A., Rudd, K. E., Russell, N., Schnall, J.,
Scott, J. A. G., Shivamallappa, M., Sifuentes-Osornio, J., Steenkeste, N.,
Stewardson, A. J., Stoeva, T., Tasak, N., Thaiprakong, A., Thwaites, G., Turner,
C., Turner, P., Doorn, H. R. van, Velaphi, S., Vongpradith, A., Vu, H., Walsh, T.,
Waner, S., Wangrangsimakul, T., Wozniak, T., Zheng, P., Sartorius, B., Lopez,
A. D., Stergachis, A., Moore, C., Dolecek, C. & Naghavi, M. Global burden of
bacterial antimicrobial resistance in 2019: a systematic analysis. *The Lancet* **399,**
629–655 (2022).

2. WHO. 2021 antibacterial agents in clinical and preclinical development: an
overview and analysis. (2022). at https://www.who.int/publications-detail-
redirect/9789240047655

3. Beyer, P. & Paulin, S. The antibacterial research and development pipeline
needs urgent solutions. *ACS Infect. Dis.* **6,** 1289–1291 (2020).

4. Bell, G. & MacLean, C. The search for 'evolution-proof' antibiotics. *Trends*
Microbiol. **26,** 471–483 (2018).

5. Monserrat-Martinez, A., Gambin, Y. & Sierecki, E. Thinking outside the bug:
molecular targets and strategies to overcome antibiotic resistance. *Int. J. Mol.*
Sci. **20,** 1255 (2019).

6. Rezzoagli, C., Granato, E. T. & Kümmerli, R. Harnessing bacterial interactions to
manage infections: a review on the opportunistic pathogen Pseudomonas
aeruginosa as a case example. *J. Med. Microbiol.* **69,** 147–161 (2020).

7. Bhushan, A., Peters, E. E. & Piel, J. *Entotheonella bacteria as source of sponge-*
derived natural products: Opportunities for biotechnological production. (Springer
International Publishing, 2017).

8. Ghosh, C., Sarkar, P., Issa, R. & Haldar, J. Alternatives to conventional
antibiotics in the era of antimicrobial resistance. *Trends Microbiol.* **27,** 323–338
(2019).

9. Mei, Y., Chen, Y., Sivaccumar, J. P., An, Z., Xia, N. & Luo, W. Research
progress and applications of nanobody in human infectious diseases. *Front.*
Pharmacol. **13,** (2022).

10. Rezzoagli, C., Archetti, M., Mignot, I., Baumgartner, M. & Kümmerli, R.
Combining antibiotics with antivirulence compounds can have synergistic effects
and reverse selection for antibiotic resistance in Pseudomonas aeruginosa.
PLOS Biol. **18,** (2020).

11. Saeki, E. K., Kobayashi, R. K. T. & Nakazato, G. Quorum sensing system: target
to control the spread of bacterial infections. *Microb. Pathog.* **142,** (2020).

12. Wale, N., Sim, D. G., Jones, M. J., Salathe, R., Day, T. & Read, A. F. Resource
limitation prevents the emergence of drug resistance by intensifying within-host
competition. *Proc. Natl. Acad. Sci.* **114,** 13774–13779 (2017).

13. Hug, J. J., Krug, D. & Müller, R. Bacteria as genetically programmable producers
of bioactive natural products. *Nat. Rev. Chem.* **4,** 172–193 (2020).

14. Imai, Y., Meyer, K. J., Iinishi, A., Favre-Godal, Q., Green, R., Manuse, S.,
Caboni, M., Mori, M., Niles, S., Ghiglieri, M., Honrao, C., Ma, X., Guo, J. J.,
Makriyannis, A., Linares-Otoya, L., Böhringer, N., Wuisan, Z. G., Kaur, H., Wu,
R., Mateus, A., Typas, A., Savitski, M. M., Espinoza, J. L., O'Rourke, A., Nelson,
K. E., Hiller, S., Noinaj, N., Schäberle, T. F., D'Onofrio, A. & Lewis, K. A new
antibiotic selectively kills Gram-negative pathogens. *Nature* **576,** 459–464
(2019).

15. Ling, L. L., Schneider, T., Peoples, A. J., Spoering, A. L., Engels, I., Conlon, B.
P., Mueller, A., Schäberle, T. F., Hughes, D. E., Epstein, S., Jones, M.,
Lazarides, L., Steadman, V. A., Cohen, D. R., Felix, C. R., Fetterman, K. A.,
Millett, W. P., Nitti, A. G., Zullo, A. M., Chen, C. & Lewis, K. A new antibiotic kills
pathogens without detectable resistance. *Nature* **517,** 455–459 (2015).

16. Butaitė, E., Baumgartner, M., Wyder, S. & Kümmerli, R. Siderophore cheating
and cheating resistance shape competition for iron in soil and freshwater
Pseudomonas communities. *Nat. Commun.* **8,** 414 (2017).

17. Deveau, A., Gross, H., Palin, B., Mehnaz, S., Schnepf, M., Leblond, P.,
Dorrestein, P. C. & Aigle, B. Role of secondary metabolites in the interaction

between Pseudomonas fluorescens and soil microorganisms under iron-limited
conditions. *FEMS Microbiol. Ecol.* **92,** (2016).

18. Hider, R. C. & Kong, X. Chemistry and biology of siderophores. *Nat. Prod. Rep.*
27, 637–657 (2010).

19. Figueiredo, A. R. T., Özkaya, Ö., Kümmerli, R. & Kramer, J. Siderophores drive
invasion dynamics in bacterial communities through their dual role as public good
versus public bad. *Ecol. Lett.* **25,** 138–150 (2022).

20. Gu, S., Wei, Z., Shao, Z., Friman, V.-P., Cao, K., Yang, T., Kramer, J., Wang, X.,
Li, M., Mei, X., Xu, Y., Shen, Q., Kümmerli, R. & Jousset, A. Competition for iron
drives phytopathogen control by natural rhizosphere microbiomes. *Nat. Microbiol.*
5, 1002–1010 (2020).

21. Lee, W., van Baalen, M. & Jansen, V. A. A. An evolutionary mechanism for
diversity in siderophore-producing bacteria. *Ecol. Lett.* **15,** 119–125 (2012).

22. Niehus, R., Picot, A., Oliveira, N. M., Mitri, S. & Foster, K. R. The evolution of
siderophore production as a competitive trait. *Evolution* **71,** 1443–1455 (2017).

23. Visca, P., Imperi, F. & Lamont, I. L. Pyoverdine siderophores: from biogenesis to
biosignificance. *Trends Microbiol.* **15,** 22–30 (2007).

24. Bonneau, A., Roche, B. & Schalk, I. J. Iron acquisition in Pseudomonas
aeruginosa by the siderophore pyoverdine: an intricate interacting network
including periplasmic and membrane proteins. *Sci. Rep.* **10,** 120 (2020).

25. Cornelis, P., Tahrioui, A., Lesouhaitier, O., Bouffartigues, E., Feuilloley, M.,
Baysse, C. & Chevalier, S. High affinity iron uptake by pyoverdine in
Pseudomonas aeruginosa involves multiple regulators besides Fur, PvdS, and
FpvI. *BioMetals* (2022).

26. Meyer, J.-M., Gruffaz, C., Raharinosy, V., Bezverbnaya, I., Schäfer, M. &
Budzikiewicz, H. Siderotyping of fluorescent Pseudomonas: molecular mass
determination by mass spectrometry as a powerful pyoverdine siderotyping
method. *BioMetals* **21,** 259–271 (2008).

27. Rehm, K., Vollenweider, V., Kümmerli, R. & Bigler, L. A comprehensive method
to elucidate pyoverdines produced by fluorescent Pseudomonas spp. by UHPLC-
HR-MS/MS. *Anal. Bioanal. Chem.* **414,** 2671–2685 (2022).

28. Meyer & Abdallah, M. A. The fluorescent pigment of Pseudomonas fluorescens:
biosynthesis, purification and physicochemical properties. *Microbiology* **107,**
319–328 (1978).

29. Butaitė, E., Kramer, J., Wyder, S. & Kümmerli, R. Environmental determinants of
pyoverdine production, exploitation and competition in natural Pseudomonas
communities. *Environ. Microbiol.* **20,** 3629–3642 (2018).

30. Butaitė, E., Kramer, J. & Kümmerli, R. Local adaptation, geographical distance
and phylogenetic relatedness: assessing the drivers of siderophore-mediated
social interactions in natural bacterial communities. *J. Evol. Biol.* **34,** 1266–1278
(2021).

31. Kramer, J., López Carrasco, M. Á. & Kümmerli, R. Positive linkage between
bacterial social traits reveals that homogeneous rather than specialised
behavioral repertoires prevail in natural Pseudomonas communities. *FEMS
Microbiol. Ecol.* **96,** (2020).

32. Meyer, J.-M. Pyoverdines: pigments, siderophores and potential taxonomic
markers of fluorescent Pseudomonas species. *Arch. Microbiol.* **174,** 135–142
(2000).

33. Cornelis, P. Iron uptake and metabolism in pseudomonads. *Appl. Microbiol.
Biotechnol.* **86,** 1637–1645 (2010).

34. Rehm, K., Vollenweider, V., Kümmerli, R. & Bigler, L. Rapid identification of
pyoverdines of fluorescent Pseudomonas spp. by UHPLC-IM-MS. *BioMetals* **36,**
19–34 (2023).

35. Meyer, J.-M., Stintzi, A., De Vos, D., Cornelis, P., Tappe, R., Taraz, K. &
Budzikiewicz, H. 1997. Use of siderophores to type pseudomonads: the three
Pseudomonas aeruginosa pyoverdine systems. *Microbiology* **143,** 35–43 (1997).

36. Budzikiewicz, H., Schäfer, M., Fernández, D. U., Matthijs, S. & Cornelis, P.
Characterization of the chromophores of pyoverdins and related siderophores by
electrospray tandem mass spectrometry. *BioMetals* **20,** 135–144 (2007).

37. Nadal-Jimenez, P., Koch, G., Reis, C. R., Muntendam, R., Raj, H., Jeronimus-
Stratingh, C. M., Cool, R. H. & Quax, W. J. PvdP Is a tyrosinase that drives
maturation of the pyoverdine chromophore in Pseudomonas aeruginosa. *J.
Bacteriol.* **196,** 2681–2690 (2014).

38. Sugue, M.-F., Burdur, A. N., Ringel, M. T., Dräger, G. & Brüser, T. PvdM of
fluorescent pseudomonads is required for the oxidation of ferribactin by PvdP in
periplasmic pyoverdine maturation. *J. Biol. Chem.* **298,** (2022).

39. Harris, W. R., Carrano, C. J., Cooper, S. R., Sofen, S. R., Avdeef, A. E., McArdle,
J. V. & Raymond, K. N. Coordination chemistry of microbial iron transport

compounds. 19. Stability constants and electrochemical behavior of ferric
enterobactin and model complexes. *J. Am. Chem. Soc.* **101,** 6097–6104 (1979).

40. Ghysels, B., Dieu, B. T. M., Beatson, S. A., Pirnay, J.-P., Ochsner, U. A., Vasil,
M. L. & Cornelis, P. 2004. FpvB, an alternative type I ferripyoverdine receptor of
Pseudomonas aeruginosa. *Microbiology* **150,** 1671–1680 (2004).

41. González, J., Salvador, M., Özkaya, Ö., Spick, M., Reid, K., Costa, C., Bailey, M.
J., Avignone Rossa, C., Kümmerli, R. & Jiménez, J. I. Loss of a pyoverdine
secondary receptor in Pseudomonas aeruginosa results in a fitter strain suitable
for population invasion. *ISME J.* **15,** 1330–1343 (2021).

42. Sexton, D. J., Glover, R. C., Loper, J. E. & Schuster, M. Pseudomonas protegens
Pf-5 favours self-produced siderophore over free-loading in interspecies
competition for iron. *Environ. Microbiol.* **19,** 3514–3525 (2017).

43. Ménard, G., Rouillon, A., Ghukasyan, G., Emily, M., Felden, B. & Donnio, P.-Y.
Galleria mellonella larvae as an infection model to investigate sRNA-mediated
pathogenesis in Staphylococcus aureus. *Front. Cell. Infect. Microbiol.* **11,** (2021).

44. Singh, G., Yadav, M., Ghosh, C. & Rathore, J. S. Bacterial toxin-antitoxin
modules: classification, functions, and association with persistence. *Curr. Res.*
Microb. Sci. **2,** 100047 (2021).

45. Diep, B. A., Gill, S. R., Chang, R. F., Phan, T. H., Chen, J. H., Davidson, M. G.,
Lin, F., Lin, J., Carleton, H. A., Mongodin, E. F., Sensabaugh, G. F. & Perdreau-
Remington, F. Complete genome sequence of USA300, an epidemic clone of
community-acquired meticillin-resistant Staphylococcus aureus. *The Lancet* **367,**
731–739 (2006).

46. Novick, R. P. & Ram, G. Staphylococcal pathogenicity islands - movers and
shakers in the genomic firmament. *Curr. Opin. Microbiol.* **38,** 197–204 (2017).

47. Bohac, T. J., Fang, L., Giblin, D. E. & Wencewicz, T. A. Fimsbactin and
acinetobactin compete for the periplasmic siderophore binding protein BauB in
pathogenic Acinetobacter baumannii. *ACS Chem. Biol.* **14,** 674–687 (2019).

48. Shapiro, J. A. & Wencewicz, T. A. Acinetobactin isomerization enables adaptive
iron acquisition in Acinetobacter baumannii through pH-triggered siderophore
swapping. *ACS Infect. Dis.* **2,** 157–168 (2016).

49. Sheldon, J. R. & Skaar, E. P. Acinetobacter baumannii can use multiple
siderophores for iron acquisition, but only acinetobactin is required for virulence.
PLOS Pathog. **16,** (2020).

50. Hammer, N. D. & Skaar, E. P. Molecular mechanisms of Staphylococcus aureus
iron acquisition. *Annu. Rev. Microbiol.* **65,** 129–147 (2011).

51. Lamont, I. L., Beare, P. A., Ochsner, U., Vasil, A. I. & Vasil, M. L. Siderophore-
mediated signaling regulates virulence factor production in Pseudomonas
aeruginosa. *Proc. Natl. Acad. Sci.* **99,** 7072–7077 (2002).

52. Johnson, M., Cockayne, A. & Morrissey, J. A. Iron-regulated biofilm formation in
Staphylococcus aureus Newman requires ica and the secreted protein Emp.
Infect. Immun. **76,** 1756–1765 (2008).

53. Banin, E., Vasil, M. L. & Greenberg, E. P. Iron and Pseudomonas aeruginosa
biofilm formation. *Proc. Natl. Acad. Sci.* **102,** 11076–11081 (2005).

54. Lin, M.-H., Shu, J.-C., Huang, H.-Y. & Cheng, Y.-C. Involvement of iron in biofilm
formation by Staphylococcus aureus. *PLOS ONE* **7,** e34388 (2012).

55. Kang, D. & Kirienko, N. V. Interdependence between iron acquisition and biofilm
formation in Pseudomonas aeruginosa. *J. Microbiol.* **56,** 449–457 (2018).

56. Gentile, V., Frangipani, E., Bonchi, C., Minandri, F., Runci, F. & Visca, P. Iron
and Acinetobacter baumannii biofilm formation. *Pathogens* **3,** 704–719 (2014).

57. Oliveira, F., Rohde, H., Vilanova, M. & Cerca, N. The emerging role of iron
acquisition in biofilm-associated infections. *Trends Microbiol.* **29,** 772–775
(2021).

58. Cassat, J. E. & Skaar, E. P. Iron in infection and immunity. *Cell Host Microbe* **13,**
509–519 (2013).

59. Skaar, E. P. The Battle for iron between bacterial pathogens and their vertebrate
hosts. *PLOS Pathog.* **6,** (2010).

60. Kang, D., Xu, Q. & Kirienko, N. V. In vitro lung epithelial cell model reveals novel
roles for Pseudomonas aeruginosa siderophores. *Microbiol. Spectr.* **12,** e03693-
23 (2024).

61. Döring, G., Pfestorf, M., Botzenhart, K. & Abdallah, M. A. Impact of proteases on
iron uptake of Pseudomonas aeruginosa pyoverdin from transferrin and
lactoferrin. *Infect. Immun.* **56,** 291–293 (1988).

62. Cox, C. D. Role of pyocyanin in the acquisition of iron from transferrin. *Infect.*
Immun. **52,** 263–270 (1986).

63. Wolz, C., Hohloch, K., Ocaktan, A., Poole, K., Evans, R. W., Rochel, N.,
Albrecht-Gary, A. M., Abdallah, M. A. & Döring, G. Iron release from transferrin

by pyoverdin and elastase from Pseudomonas aeruginosa. *Infect. Immun.* **62,** 4021–4027 (1994).

64. Kang, D. & Kirienko, N. V. An in vitro cell culture model for pyoverdine-mediated Virulence. *Pathogens* **10,** 9 (2021).

65. Kirienko, N. V., Kirienko, D. R., Larkins-Ford, J., Wählby, C., Ruvkun, G. & Ausubel, F. M. Pseudomonas aeruginosa disrupts caenorhabditis elegans iron homeostasis, causing a hypoxic response and death. *Cell Host Microbe* **13,** 406–416 (2013).

66. Pita-Grisanti, V., Chasser, K., Sobol, T. & Cruz-Monserrate, Z. Understanding the potential and risk of bacterial siderophores in cancer. *Front. Oncol.* **12,** (2022).

67. Peek, M. E., Bhatnagar, A., McCarty, N. A. & Zughaier, S. M. Pyoverdine, the major siderophore in Pseudomonas aeruginosa, evades NGAL recognition. *Interdiscip. Perspect. Infect. Dis.* **2012,** 843509 (2012).

68. Rezzoagli, C., Wilson, D., Weigert, M., Wyder, S. & Kümmerli, R. Probing the evolutionary robustness of two repurposed drugs targeting iron uptake in Pseudomonas aeruginosa. *Evol. Med. Public Health* **2018,** 246–259 (2018).

69. Sommer, M. O. A., Munck, C., Toft-Kehler, R. V. & Andersson, D. I. Prediction of antibiotic resistance: time for a new preclinical paradigm? *Nat. Rev. Microbiol.* **15,** 689–696 (2017).

70. Qiu, D.-H., Huang, Z.-L., Zhou, T., Shen, C. & Hider, R. C. In vitro inhibition of bacterial growth by iron chelators. *FEMS Microbiol. Lett.* **314,** 107–111 (2011).

71. Kontoghiorghes, G. J., Kolnagou, A., Skiada, A. & Petrikkos, G. The role of iron and chelators on infections in iron overload and non iron loaded conditions: prospects for the design of new antimicrobial therapies. *Hemoglobin* **34,** 227–239 (2010).

72. Ribeiro, M., Sousa, C. A. & Simões, M. Harnessing microbial iron chelators to develop innovative therapeutic agents. *J. Adv. Res.* **39,** 89–101 (2022).

73. Coraça-Huber, D. C., Dichtl, S., Steixner, S., Nogler, M. & Weiss, G. Iron chelation destabilizes bacterial biofilms and potentiates the antimicrobial activity of antibiotics against coagulase-negative Staphylococci. *Pathog. Dis.* **76,** fty052 (2018).

74. Hatcher, H. C., Singh, R. N., Torti, F. M. & Torti, S. V. Synthetic and natural iron chelators: therapeutic potential and clinical use. *Future Med. Chem.* **1,** 1643–1670 (2009).

1180 75. Mobarra, N., Shanaki, M., Ehteram, H., Nasiri, H., Sahmani, M., Saeidi, M., Goudarzi, M., Pourkarim, H. & Azad, M. A review on iron chelators in treatment of iron overload syndromes. *Int. J. Hematol.-Oncol. Stem Cell Res.* **10,** 239–247 (2016).

1184 76. Elalfy, M. S., Hamdy, M., Adly, A., Ebeid, F. S. E., Temin, N. T., Rozova, A., Lee, D., Fradette, C. & Tricta, F. Efficacy and safety of early-start deferiprone in infants and young children with transfusion-dependent beta thalassemia: evidence for iron shuttling to transferrin in a randomized, double-blind, placebo-controlled, clinical trial (START). *Am. J. Hematol.* **98,** 1415–1424 (2023).

77. Eto, D., Watanabe, K., Saeki, H., Oinuma, K., Otani, K., Nobukuni, M., Shiratori-Takano, H., Takano, H., Beppu, T. & Ueda, K. Divergent effects of desferrioxamine on bacterial growth and characteristics. *J. Antibiot.* **66,** 199–203 (2013).

78. Thompson, M. G., Corey, B. W., Si, Y., Craft, D. W. & Zurawski, D. V. Antibacterial activities of iron chelators against common nosocomial pathogens. *Antimicrob. Agents Chemother.* **56,** 5419–5421 (2012).

79. Visca, P., Bonchi, C., Minandri, F., Frangipani, E. & Imperi, F. The dual personality of iron chelators: growth inhibitors or promoters? *Antimicrob. Agents Chemother.* **57,** 2432–2433 (2013).

80. de Léséleuc, L., Harris, G., KuoLee, R. & Chen, W. In vitro and in vivo biological activities of iron chelators and gallium nitrate against Acinetobacter baumannii. *Antimicrob. Agents Chemother.* **56,** 5397–5400 (2012).

81. Yang, J., Long, H., Hu, Y., Feng, Y., McNally, A. & Zong, Z. Klebsiella oxytoca complex: update on taxonomy, antimicrobial resistance, and virulence. *Clin. Microbiol. Rev.* **35,** (2021).

82. Cunrath, O., Geoffroy, V. A. & Schalk, I. J. Metallome of Pseudomonas aeruginosa: a role for siderophores. *Environ. Microbiol.* **18,** 3258–3267 (2016).

83. Sprouffske, K. & Wagner, A. Growthcurver: An R package for obtaining interpretable metrics from microbial growth curves. *BMC Bioinformatics* **17,** 172 (2016).

84. Deatherage, D. E. & Barrick, J. E. Identification of mutations in laboratory-evolved microbes from next-generation sequencing data using breseq. *Eng. Anal. Multicell. Syst. Methods Protoc.* 165–188 (2014). doi:10.1007/978-1-4939-0554-6_12

85. Commo, F. & Bot, B. nplr: n-parameter logistic regression. R package version
0.1-7. (2016).
86. Therneau, T. A package for survival analysis in R. (2020). at https://CRAN.R-
project.org/package=survival
87. Letunic, I. & Bork, P. Interactive tree of life (iTOL) v3: an online tool for the
display and annotation of phylogenetic and other trees. *Nucleic Acids Res.* **44,**
W242–W245 (2016).
1221