1
2
3 **Sperm induction of somatic cell-cell fusion as a novel functional test**
4
5
6
7

8 Nicolas G. Brukman[1#], Clari Valansi[1], Benjamin Podbilewicz[1#]
9
10
11 [1]Department of Biology, Technion- Israel Institute of Technology, Haifa 32000, Israel
12
[#]Correspondence should be addressed to N.G.B. (nbrukman@gmail.com) and B.P.
(podbilew@technion.ac.il).

Summary

The fusion of mammalian gametes requires the interaction between IZUMO1 on the
sperm and JUNO on the oocyte. We have recently shown that ectopic expression of
mouse IZUMO1 induces cell-cell fusion and that sperm can fuse to fibroblasts
expressing JUNO. Here, we found that the incubation of mouse sperm with hamster
fibroblasts or human epithelial cells in culture induces the fusion between these
somatic cells and formation of syncytia, a pattern previously observed with some
animal viruses. This sperm-induced cell-cell fusion requires a species-matching JUNO
on both fusing cells, can be blocked by an antibody against IZUMO1, and does not
rely on the synthesis of new proteins. The fusion is dependent on the sperm's
fusogenic capacity, making this a reliable, fast and simple method for predicting sperm
function during the diagnosis of male infertility.

Highlights

- Sperm induces viral-like fusion of somatic cells expressing JUNO
- We developed a new technique to determine the fertilization potential of sperm
- The test measures the capacity of sperm to induce somatic cell fusion
- The degree of somatic cell fusion correlates with the fertilizing ability of sperm

Graphical abstract

Keywords

Introduction

Infertility is estimated to affect approximately 15% of the population (World Health Organization 2023), and many couples dealing with this burden turn to assisted reproductive techniques (ARTs) as a possible solution (Steptoe and Edwards 1978; Palermo et al. 1992). During fertility evaluation of a male patient, a basic semen examination is performed, where different sperm parameters are determined: concentration, motility, morphology and vitality (World Health Organization, HRP 2021). This basic analysis can be complemented with additional tests to determine the fertilizing potential of the sperm and choose the best treatment for the couple. The hamster oocyte penetration (HOP) test has been proposed as a quantitative method for analyzing the fusogenic potential of human spermatozoa (Aitken and Elton 1986; Yanagimachi, Yanagimachi, and Rogers 1976; Zainul Rashid et al. 1998), however, it was excluded from the newest WHO laboratory manual for the examination and processing of human semen (World Health Organization, HRP 2021) for being considered obsolete. Therefore, to date there is no other standardized methodology to analyze specifically the ability of sperm to fuse to oocytes.

In mammals, the adhesion of the sperm to the oocyte plasma membranes is mediated by the species-specific interaction of two membrane proteins: IZUMO1 and JUNO (Bianchi and Wright 2015; Aydin et al. 2016; Ohto et al. 2016; Matsumura et al. 2021). The transmembrane protein IZUMO1 is expressed during spermatogenesis and localizes to the fusogenic region of the sperm head after an exocytic process named acrosome reaction (Inoue et al. 2005; Satouh et al. 2012; Inoue et al. 2015; Nishimura et al. 2016). On the other hand, the IZUMO1 receptor, JUNO, is an oocyte protein bound to the plasma membrane by a GPI lipid anchor (Bianchi et al. 2014). The subsequent fusion of the two gametes relies on the action of IZUMO1 in a unilateral manner (Brukman et al. 2023). The IZUMO1-JUNO interaction was characterized also in humans (Aydin et al. 2016; Ohto et al. 2016) and its relevance for human fertility is supported by the presence of antibodies against IZUMO1 in the sera of immuno-infertile women (Clark and Naz 2013), the mutations on JUNO in cases of fertilization failure and polyspermy (Yu et al. 2018), and the lower levels of IZUMO1 observed in sperm that failed to fertilize during *in vitro* fertilization (IVF) treatments (Enoiu et al. 2022).

Recently, we found that mouse sperm can fuse to fibroblasts ectopically expressing the sperm-receptor JUNO (Brukman et al. 2023). Here, we show that after sperm fuses with a somatic cell, this cell can fuse with additional cells inducing syncytia formation - a single cell with several nuclei (multinucleated cell). This is likely mediated by the transfer of the fusion machinery from the sperm to the target somatic cell. We call this process "SPerm-Induced CEll-cell fusion Requiring JUNO" (SPICER). We found that sperm with higher fertilizing ability can induce the fusion of somatic cells more efficiently, judged by the increased levels of multinucleation. Therefore, the number and size of these syncytia can indicate the fertilizing potential of the sperm

and can help embryologists to better predict the success of the different ARTs for each
individual. Based on this, we propose a new method for diagnosis of male fertility
based on the ability of sperm to induce cell-cell fusion *in vitro*.

Results

Sperm fusion to fibroblasts promotes syncytia formation

There are reports showing that mammalian sperm can fuse to somatic cells (Rival
et al. 2019; Mattioli et al. 2009; Bendich, Borenfreund, and Sternberg 1974). In our
experimental conditions, mouse sperm cells only fuse to Baby Hamster Kidney (BHK)
cells if they are induced to express the sperm-receptor, JUNO (Brukman et al. 2023).
This fusion was demonstrated by detecting the transfer of the DNA binding GFP (GFP-
MBD) from the BHK cells to the sperm heads (Brukman et al. 2023). To further study
the mechanisms of mammalian sperm-oocyte fusion, we incubated mouse sperm with
BHK cells expressing JUNO (functioning as pseudo-oocytes) and determined the
efficiency of the sperm-BHK cell interactions. Surprisingly, we found that sperm
induces the formation of multinucleated BHK cells (syncytia; Fig. 1A-C). The induction
of multinucleation was dependent on the presence of JUNO (Fig. 1B-C), however,
JUNO expression alone is not sufficient to induce this process. Only cells with sperm
fused to them form syncytia (Fig. 1C). Thus, the fusion of sperm with JUNO-expressing
BHK cells is required for inducing subsequent multinucleation of these somatic cells.
This phenomenon was unexpected, leading us to further study the mechanisms of this
process.

Sperm induce syncytia formation using a viral-like mechanism

To our knowledge, sperm-induced cell-cell fusion has not been described in any
species. However, decades ago, it was first described that some somatic cells can
fuse following viral infection (Okada 1962; Kohn 1965) and later confirmed for diverse
viruses, including SARS-CoV2 (Buchrieser et al. 2020). This process may require the
synthesis of new viral proteins and therefore induces Fusion From Within (FFWI) while
other viruses induce fusion independently of protein synthesis in a process called
Fusion From Without (FFWO) (Bratt and Gallaher 1969; Duelli and Lazebnik 2007).
For instance, orthoreoviruses induce the expression of fusion-associated small
transmembrane (FAST) proteins upon infection, promoting FFWI (Ciechonska, Key,
and Duncan 2014; Theuerkauf et al. 2021). We hypothesized that in a way reminiscent
of viral induced cell-cell fusion, sperm could induce syncytia formation following its
merger via FFWI or FFWO. To distinguish between these alternative mechanisms, we
performed the experiment in the presence of cycloheximide to inhibit protein synthesis.
We found no differences in the levels of multinucleation between control and
cycloheximide treatments (Fig 1B and D). This shows that *de novo* protein synthesis
is not required for the induction of multinucleation by sperm and suggests a
mechanism of FFWO. Additionally, live imaging of the process showed that fusion
between cells occurs efficiently within 10-15 minutes after the addition of the sperm

(Fig. 1E, Movie S1), suggesting that protein synthesis is not involved. We have previously shown that IZUMO1 fades from the sperm after fusion to the fibroblast (Brukman et al. 2023) indicating that this transmembrane protein is transferred to the target cell plasma membrane. Consistent with these findings, IZUMO1 was observed diffusing to the oocyte plasma membrane during in vitro fertilization assays (Satouh et al. 2012). Here we detected IZUMO1 by immunostaining and by a fluorescent reporter and confirmed that it diffuses from the sperm head after its fusion with the BHK cell (Fig. S1A-B), suggesting a critical function for the fusogenic machinery carried by the sperm during the FFWO process. Thus, sperm-BHK fusion appears to trigger BHK-BHK cell fusion, observed as multinucleation, without the requirement of *de novo* synthesis of proteins, likely by passing the sperm's fusogenic machinery onto the BHK cells. In this sense, two models may explain FFWO induced by sperm (Fig. S1C), similarly to what was reported for viruses (Tang et al. 2021). The first model involves an initial fusion of the sperm to a BHK cell expressing JUNO, the transfer of IZUMO1 to the target cell (forming a pseudo-sperm from a pseudo-oocyte) and a subsequent fusion to a new JUNO-expressing BHK (Fig. S1Ci). On the other hand, the second model suggests the fusion of one sperm cell with two JUNO-positive BHK cells, bridging both and forming a continuous syncytium (Fig. S1Cii). At the moment, we cannot exclude either model, or their coexistence. Even though it is not known whether sperm can induce fusion of somatic cells *in vivo*, we show that sperm use a viral-like mechanism of FFWO when the fibroblasts express the oocyte JUNO *in vitro*.

Syncytia formation results from the fusion of cells expressing JUNO

Then we asked whether the sperm-induced multinucleation was a consequence of BHK-BHK fusion and if so, whether it requires JUNO to be present on both fusing cells. For that, we employed a content mixing experiment where two populations of cells expressing different fluorescent markers were mixed and then exposed to sperm (Fig. 2A). We observed the formation of multinucleated hybrid cells containing both fluorescent markers, confirming the fusion of BHK cells expressing JUNO (Fig. 2B). This BHK-BHK fusion was not observed when only one or neither of the cell populations expressed JUNO (Fig. 2B-C), indicating that sperm-induced multinucleation is indeed BHK-BHK fusion which relies on bilateral expression of JUNO. An alternative explanation for sperm-induced multinucleation is a failure in cytokinesis. However, the presence of an inhibitor of the cell cycle (FdUdr, 5-fluoro-2′-deoxyuridine, (Valansi et al. 2017)) does not inhibit multinucleation (Fig. 3A-B), ruling out that syncytia formation is as a consequence of failure in the division of the BHKs and confirming the occurrence of fusion between them. Following these findings, we named this process SPICER that stands for "SPerm-Induced CEll-cell fusion Requiring JUNO".

Sperm-induced multinucleation as a readout of sperm fertilizing ability

Following the discovery of the ability of sperm to induce cell-cell fusion, we decided to evaluate whether BHK multinucleation could be used as a readout of sperm

fusogenic potential. For this purpose, we incubated the sperm in media lacking BSA or Ca^{2+} that do not support capacitation (i.e. a process by which sperm acquires its fusogenic activity) (Visconti et al. 1995; Yanagimachi 1988). We found that sperm cells incubated under these conditions failed to fuse to BHK cells, as well as to induce syncytia formation (Fig 3A-B). Thus, BHK-BHK cell fusion requires fully capacitated sperm. A sperm which is unable to fuse with BHK cells, will not induce subsequent multinucleation.

SPICER depends on the activity of IZUMO1

The sperm IZUMO1 protein is essential for sperm-egg interactions via binding to JUNO (Bianchi et al. 2014). More recently, IZUMO1 was shown to induce fusion of cells in a JUNO-independent way using a different domain that is not required for binding to the egg receptor (Brukman et al. 2023). To study whether IZUMO1 is required for the fusion of sperm to BHK cells and for the induction of multinucleation of the fibroblasts, we decided to inhibit IZUMO1 activity using a blocking antibody. For this purpose, we used a monoclonal antibody against IZUMO1 which has been shown to inhibit sperm-egg fusion (Inoue et al. 2013) in our sperm-induced multinucleation assay. We found that the anti-IZUMO1 antibody had the ability to block both sperm-BHK and sperm-induced BHK-BHK fusions even when sperm cells were fully capacitated (Fig 3A-B). Altogether, our results show that the SPICER assay is able to determine sperm fertilization potential that depends on the IZUMO-JUNO interactions.

Cell-cell fusion requires species-specific JUNO

Elegant work on IZUMO1-JUNO interactions among different mammalian species has shown species-specific selectivity between these crucial interactions (Bianchi and Wright 2015). To evaluate the effect of species-specificity of the SPICER assay, we incubated the mouse sperm with BHK cells expressing human JUNO. Under these conditions, we observed lower levels of sperm-BHK fusion and a reduction in BHK multinucleation compared to cells expressing mouse JUNO (Fig 4A-C). Interestingly, the low activity mediated by human JUNO was still significant compared to the control without JUNO. This confirms the requirement of a species-matching JUNO for the assay; however, it puts in evidence a residual cross-interaction between mouse sperm and human JUNO.

Sperm induce the fusion of human epithelial cells

To exclude any BHK cell-specific requirement for SPICER, we tested the epithelial human embryonic kidney 293T (HEK) cells. Due to the smaller size and more rounded shape of these cells that hinder the quantification of multinucleation, we opted to analyze fusion by content mixing. For this, we utilized the Dual Split Protein system (Wang et al. 2014; Ishikawa et al. 2012) which employs two GFP halves expressed in different populations of cells. Upon fusion of these two populations, the reporter self-assembles and the cells become fluorescent (Nakane and Matsuda 2015). The

number of green fluorescent cells relative to the number of transfected cells was
determined as an indication of fusion levels (Fig. 5A). As observed for the BHK cells,
HEK cells bearing JUNO fused between them when they were co-incubated with
sperm (Fig. 5B-C). On the contrary, neither HEK cells without JUNO nor HEK
expressing JUNO but without sperm showed content mixing (Fig. 5B-C). These results
show that SPICER is not restricted to BHK cells or fibroblasts and opens the possibility
for adapting this assay to different cell lines and approaches to evaluate cell-cell
fusion.

Discussion

SPICER as a diagnostic tool

Determining sperm's fusogenic potential is of great interest for infertility diagnosis
of both human and stud animals. Considering that the hamster oocyte penetration
(HOP) assay became obsolete, an assay that evaluates the fusion of somatic cells
with ectopic expression of JUNO induced by competent sperm (SPICER, Fig. 6)
represents a powerful diagnostic tool. SPICER could predict the success chances of
ARTs like intrauterine insemination (IUI) or conventional IVF, which require a fusion-
competent sperm and if it is necessary to proceed with a more complex technique
such as intracytoplasmic sperm injection (ICSI). This is particularly important
considering the routine use of the ICSI is under debate as it is associated with a slightly
higher risk of adverse outcomes in the progeny (Practice Committees of the American
Society for Reproductive Medicine and the Society for Assisted Reproductive
Technology 2020).

Considering our previous work showing that IZUMO1 can mediate cell-cell fusion
(Brukman et al. 2023) and that significantly lower levels of this protein were detected
in the sperm cells from patients with total fertilization failure (Enoiu et al. 2022), we
expect that the SPICER assay will be able to resolve some unexplained cases of male
infertility that result from loss of function of components of the fusion machinery. In this
context, other proteins that are essential for gamete fusion, such as SPACA6,
TMEM95, FIMP, SOF1 and DCST1/2 (Barbaux et al. 2020; Noda et al. 2020;
Lorenzetti et al. 2014; Lamas-Toranzo et al. 2020; Fujihara et al. 2020; Inoue,
Hagihara, and Wada 2021; Noda et al. 2022), might be transferred to the somatic cells
together with IZUMO1 during sperm-induced cell-cell fusion. Finally, considering that
antibodies against IZUMO1 were detected on the sera of infertile women (Clark and
Naz 2013) and here we observed that a monoclonal antibody against this protein was
able to block sperm-induced syncytia, SPICER could serve as tool to readily analyze
cases of immuno-infertility.

SPICER occurrence *in vivo*

Apart from being expressed in oocytes, JUNO has been described in lymphoid
tissues (Spiegelstein, Eudy, and Finnell 2000), more specifically in regulatory T cells

(Yamaguchi et al. 2007). Therefore, we cannot exclude that sperm could bind and fuse
to other cell types *in vivo*, and whether it has any physiological or pathological
relevance. This fusion cannot occur before acrosomal exocytosis, when IZUMO1 is
enclosed into the acrosome (Inoue et al. 2005; Satouh et al. 2012), and therefore, it
will be relevant to study sperm fusion to somatic cells within the oviduct of the female,
close to the fertilization site (Muro et al. 2016; Hino et al. 2016; La Spina et al. 2016).
Furthermore, JUNO was detected in the first polar body (Suzuki et al. 2017) and
whether sperm can fuse to it or if sperm fusion to the oocyte can induce the polar body-
egg fusion remains unknown.

Expanding SPICER across species

The assay that we described here can be a potent tool to study cross-fertilization
between different species and mechanisms of speciation and evolution, presenting a
step beyond biochemical approaches to study IZUMO1-JUNO interactions (Bianchi
and Wright 2015; Nakajima et al. 2022). On the other hand, by simply exchanging the
JUNO sequence used, this assay could be easily adapted to allow the analysis of the
sperm fertilizing ability in different mammalian species, such as humans or cattle,
having implications not only for reproductive health but also for biotechnological and
agricultural uses. Furthermore, even though IZUMO-JUNO interaction is a specific
requirement during sperm-egg fusion of mammals (Grayson 2015), we cannot exclude
that alternative configurations, using the right egg receptors (Lu and Ikawa 2022), will
be possible in the future to induce somatic cell-cell fusion by sperm of other sexually
reproducing organisms. In species where *Izumo1* and *Juno* genes are absent, such
as plants and protists, the gamete fusion is promoted by the fusogen HAPLESS
2/GENERATIVE CELL-SPECIFIC 1 (HAP2/GCS1) (Brukman, Li, and Podbilewicz
2022). In other organisms like zebrafish or nematodes, IZUMO1 orthologs were
described (Binner et al. 2021; Nishimura et al. 2015; Takayama et al. 2021) but, since
JUNO is present only in mammals (Vance and Lee 2020; Grayson 2015), it is unclear
whether male gametes in other species interact with different receptors on the female
gamete. Therefore, it can be speculated, for instance, that the expression of a species-
specific Bouncer will be required to activate the fusion of fish sperm (Gert et al. 2023;
Herberg et al. 2018) or *Chlamydomonas* FUS1 to mediate the fusion of the algae
minus gamete (Pinello, Liu, and Snell 2021; Ferris, Woessner, and Goodenough
1996).

Further applications for SPICER

In addition to the potential uses of SPICER for diagnosis of infertility, this assay
could be used to evaluate potential sperm donors and animal studs. Moreover, our
SPICER tool has the potential to aid screening for compounds that enhance
fertilization (new fertility treatments) or that block gamete interactions (Stepanenko et
al. 2022), or to easily determine the effect of genetic variations of JUNO (Allingham
and Floriano 2021; Takaiso et al. 2016). SPICER could also facilitate studies aimed at
resolving standing enigmas of sperm-egg fusion (Deneke and Pauli 2021;

Yanagimachi 2022). This includes the role of IZUMO1 and other potential fusogens as
well as additional cellular pathways that are known to influence fusion in different
systems, including the action of the cytoskeleton and molecular motors (Shilagardi et
al. 2013; Yang et al. 2017; Zhang et al. 2017), specific lipids (e.g. phosphatidylserine)
(Rival et al. 2019; van den Eijnde et al. 2001; Verma et al. 2018; Abay et al. 2017),
calcium signaling (Tsuchiya et al. 2018; Earles et al. 2001), phosphorylation cascades
(Eigler et al. 2021) and more (Brukman et al. 2019). Finally, whether other cell types
rather than sperm, such as muscle cells or osteoclasts, can induce syncytia formation
after ectopic fusion remains an open question, with important biological implications.

Limitations of the study

The main caveat of our assay is that it cannot discriminate between defects in
sperm docking or gamete fusion, as a poor performance in SPICER could be explained
by the altered interaction of IZUMO1 and JUNO or by a reduced capacity of IZUMO1
(and IZUMO1 partners in the sperm) to induce membrane merger. Moreover,
simultaneous measurements of the levels of spontaneous acrosome reaction will be
needed to exclude a faulty capacitation. Alternatively, capacitation could be further
stimulated pharmacologically using a Ca^{2+} ionophore (Tateno et al. 2013). In this
context, further analysis may be required to refine the characterization of defective
sperm. Lastly, the exact pathway for the FFWO mediated by the sperm will be a
subject of future investigations.

MATERIALS AND METHODS
305
306 **KEY RESOURCES TABLE**
307

REAGENT or RESOURCE	SOURCE	IDENTIFIER
Antibodies		
Anti-IZUMO1 Mab120 monoclonal antibody	Merck Millipore	Cat# MABT1357
Anti-goat IgG secondary antibody, Alexa Fluor 647	Thermo Fisher Scientific	Cat# A-21247, RRID: AB_141778
Bacterial and virus strains		
DH5α competent cells	Thermo Fisher Scientific	Cat# 18265017
Chemicals, peptides, and recombinant proteins		
jetPRIME® transfection reagent	Polyplus Transfection	Cat# 101000046

FdUdr, 5-Fluoro-2'-deoxyuridine	Sigma	Cat# F0503, CAS: 50-91-9
BSA, bovine serum albumin	Sigma	Cat# A7906, CAS: 9048-46-8
Hyaluronidase	Sigma	Cat# H3506, CAS: 37326-33-3
Cycloheximide	Sigma	Cat# C7698, CAS: 66-81-9
Puromycin Dihydrochloride	GoldBio	Cat# P-600-100, CAS: 58-58-2
Poly-L-lysine hydrobromide	Sigma	Cat# P2636, CAS: 25988-63-0
Experimental models: Cell lines		
BHK-21, clone 13	ATCC	Cat# CCL-10, RRID: CVCL_1915
HEK293T	ATCC	Cat# CRL-3216, RRID:CVCL_0063
HEK293T-DSP1-7	This paper	N/A
HEK293T-DSP8-11	This paper	N/A
Experimental models: Organisms/strains		
B6D2.C57BL/6-Tg(Zp3-EGFP/Cd9)1Osb mouse line	(Satouh et al. 2012)	N/A
FVB/129sv/CF1	This paper	N/A
Oligonucleotides		
IZUMO1-mCherry F (Genotyping): CCTTCCTGCGGCTTGTTCTCT	(Satouh et al. 2012)	N/A

IZUMO1-mCherry R (Genotyping): ATCAAGGTCTCAGAACTGTTCTCCCAAACC	(Satouh et al. 2012)	N/A
NheI Human Juno (Cloning into pCI::H2B-RFP): TTATCGCTAGCATGGCATGCTGGTGGCCGCTC	This paper	N/A
EcoRV Human Juno R (Cloning into pCI::H2B-RFP): CCAGGATATCTCAGGAAAGGAACGGCAGGAAC	This paper	N/A
Recombinant DNA		
pCI::H2B-RFP	(Williams et al. 2018)	Addgene #92398
pCI::GFPnes	(Moi et al. 2022)	N/A
pCI::mJUNO::H2B-RFP	(Brukman et al. 2023)	N/A
pCI::mJUNO::GFPnes	(Brukman et al. 2023)	N/A
pcDNA3.1-EGFP-MBD-nls	(Yamagata et al. 2005)	N/A
pCMV6-humanJUNO	(Wood and Wright 2019)	N/A
pCI::humanJUNO::H2B-RFP	This paper	N/A
pIRESpuro3-DSP1–7	(Wang et al. 2014)	N/A
pIRESpuro3-DSP8-11	(Wang et al. 2014)	N/A
Software and algorithms		
GraphPad Prism 9	GraphPad Prism	RRID:SCR_002798
FIJI (ImageJ 1.53c)	Image J	RRID:SCR_002285
Photoshop CS6	Adobe	RRID:SCR_014199
Illustrator CS6	Adobe	RRID:SCR_010279
ZEN microscopy software 7.0.4.0	ZEISS	RRID:SCR_013672

MetaMorph image analysis software 7.8.1.0	Molecular Devices	RRID:SCR_002368
Biorender	Biorender	RRID:SCR_018361

RESOURCE AVAILABILITY

Lead contact

312 Further information and requests for resources and reagents should be directed to and
313 will be fulfilled by the lead contact, Benjamin Podbilewicz (podbilew@technion.ac.il).

Materials availability

316 Plasmids or cell lines generated in this study are available upon request. Signing a
317 materials transfer agreement (MTA) may be required.

Data and code availability

Any additional information required to reanalyze the data reported in this paper is
available from the lead contact upon request.

EXPERIMENTAL MODEL AND SUBJECT DETAILS

Animals

All animal studies were approved by the Committee on the Ethics of Animal
Experiments of the Technion - Israel Institute of Technology. In this study, we used
wild-type male mice from a FVB/129sv/CF1 mixed background and B6D2-Tg(Izumo1-
mCherry) mouse line (Satouh et al. 2012). Animals were bred and housed in the
Technion animal facility under specific pathogen–free conditions with *ad libitum*
access to food and water. The primers used for genotyping Izumo1-mCherry are
outlined in the Key Resources Table. In all cases, male mice between 3 and 6 months
old were used for the experiments.

Cell lines and DNA transfection

BHK cells (Cat# CCL-10; ATCC, RRID: CVCL_1915) and HEK293T cells (Cat#
CRL-3216; ATCC, RRID: CVCL_0063) were grown and maintained in DMEM containing
10% FBS. Cells were cultured at 37°C in 5% CO_2. Plasmids were transfected into cells
using 2 µl jetPRIME (PolyPlus-transfection) per µg of DNA in 100 µl of reaction buffer
for every ml of medium. HEK293T cells stable lines for Dual Split Proteins (DSP) 1-7
and 8-11 were prepared by transfecting pIRESpuro3-DSP1–7 and pIRESpuro3-
DSP8-11 respectively and selecting with 2 µg/ml of puromycin for 10–13 days as
previously described (Wang et al. 2014).

METHOD DETAILS

Sperm collection and capacitation

Sperm were recovered by incising the cauda epididymis, obtained from adult male mice, in 300 µl of mHTF medium (Kito et al. 2004) supplemented with 4 mg/ml of BSA. The sperm were diluted in fresh medium to a concentration of 5×10^6 cells/ml and incubated for 90 min at 37°C and 5% CO_2 to induce capacitation.

Sperm-to-BHK fusion and multinucleation

BHK cells were grown on 24-well glass bottom tissue-culture plates. 24 h after plating, cells were transfected with 0.25 µg pcDNA3.1-EGFP-MBD-nls plasmids and 0.5 µg of either pCI::H2B-RFP or pCI::JUNO::H2B-RFP. 24 h after transfection, 2×10^6 capacitated wild-type sperm cells in mHTF were added to each well and co-incubated with the BHK cells for 4 h at 37°C and 5% CO_2. After one wash with PBS, the cells were fixed with 4% PFA in PBS and stained with 1 µg/ml DAPI. Micrographs were obtained using wide-field illumination using an ELYRA system S.1 microscope (Plan-Apochromat 20× NA 0.8; Zeiss). Multinucleation percentage was determined as the ratio between the number of nuclei in multinucleated cells (NuM) and the total number of nuclei in fluorescent cells (NuF), as follows: % of multinucleation = (NuM/NuF) × 100. 500 nuclei (NuF) were counted in each independent repetition (experimental point). In some cases, the number of sperm fused was determined by evaluating the transfer of EGFP-MBD-nls signal from the BHK cell to the sperm nuclei (number of fused sperm/500 BHK cells). In some experiments, sperm cells obtained from transgenic mice expressing IZUMO1-mCherry were employed to analyze IZUMO1 localization.

Content mixing assay using different colors

BHK cells at 70% confluence in 35-mm plates were transfected with 1 µg pCI::H2B-RFP, pCI::GFPnes, pCI::JUNO::H2B-RFP or pCI::JUNO::GFPnes. 4 h after transfection, the cells were washed four times with DMEM with 10% serum, four times with PBS, and detached using Trypsin (Biological Industries). The cells were collected, resuspended in DMEM with 10% serum, and counted. Equal amount of H2B-RFP and GFPnes cells (1.25×10^5 each) were mixed and seeded on glass-bottom plates (12-well black, glass-bottom #1.5H; Cellvis) and incubated at 37°C and 5% CO_2. 18 h after mixing, 4×10^6 capacitated wild-type sperm cells in mHTF were added to the BHK cells and co-incubated for 4 h after which they were washed with PBS, fixed with 4% PFA in PBS and stained with 1 µg/ml DAPI. Micrographs were obtained using wide-field illumination using an ELYRA system S.1 microscope (Plan-Apochromat 20× NA 0.8; Zeiss). The percentage of mixing was defined as the ratio between the nuclei in mixed cells (NuM) and the total number of nuclei in mixed cells and fluorescent cells in contact that did not fuse (NuC), as follows: % of mixing = (NuM/[NuM + NuC]) × 100. 1,000 nuclei (NuM + NuC) were counted in each independent repetition (experimental point).

Content mixing assay using the Dual Split Protein

Content mixing assay using the Dual Split Protein

Equal amount of HEK293T cells stably expressing DSP1-7 or DSP8-11 were
mixed (1.25×10^5 each) and seeded on glass-bottom plates (12-well black, glass-
bottom #1.5H; Cellvis) pre-treated with 20 µg/ml of Poly-L-lysine. 24 h later, the cells
were transfected with 1 µg pCI::H2B-RFP or pCI::JUNO::H2B-RFP. 18 h after
transfection, 4×10^6 capacitated wild-type sperm cells in mHTF were added to the
HEK293T cells and co-incubated for 4 h after which they were washed with PBS, fixed
with 4% PFA in PBS and stained with 1 µg/ml DAPI. Micrographs were obtained as
above, using wide-field illumination using an ELYRA system S.1 microscope (Plan-
Apochromat 20× NA 0.8; Zeiss). The number of GFP positive cells per 1,000 nuclei
was determined. Between 1,000 to 2,000 red nuclei were counted in each independent
repetition (experimental point).

Inhibition of sperm-induced cell-cell fusion

Different conditions were tested to evaluate their effect on sperm-induced cell-cell
fusion. Non-capacitated sperm were incubated in medium mHTF lacking Ca^{2+} or BSA
for 90 min at 37°C and 5% CO_2. In other cases, capacitated sperm were mixed before
their addition to the BHK cells with 1 µg/µl of anti-IZUMO1 antibody (Inoue et al. 2013),
20 µM of the cell cycle inhibitor FdUdr (Valansi et al. 2017) or 200 µg/ml of the inhibitor
of protein synthesis cycloheximide (Wengler 1975).

Time-lapse Imaging of sperm induced cell-cell fusion

BHK cells were grown on 35 mm glass bottom tissue-culture plates (Greiner Bio-
one) and, 24 h after plating, cells were transfected with 1.5 µg pCI::JUNO::GFPnes. 24
h after transfection, time-lapse images of the cells were imaged before and after
adding 5×10^6 capacitated wild-type sperm cells in mHTF. Images of the cells were
acquired every 3 min for 1 h to record cell-to-cell fusion, using a spinning disc confocal
microscope (CSU-X; Yokogawa Electric Corporation) with an Eclipse Ti inverted
microscope and a Plan-Apochromat ×20 (NA, 0.75; Nikon) objective. Images were
obtained using an iXon3 EMCCD camera (ANDOR) through MetaMorph (Molecular
Devices, version 7.8.1.0). Images in differential interference contrast and green
channels were recorded.

Immunofluorescence

The localization of IZUMO1 was determined by immunostaining. Briefly, after
fixation cell were permeabilized with 0.1% Triton X-100 in PBS and incubated with
anti-IZUMO1, clone Mab120 (1:500, Cat# MABT1357; Merck Millipore) followed by the
secondary antibody Alexa Fluor 647 goat anti-rat (1:500, Cat# A-21247; Thermo
Fisher Scientific, RRID: AB_141778). Later, the nuclei were stained with 1 µg/ml DAPI
and micrographs were obtained using wide-field illumination using an ELYRA system
S.1 microscope (Plan-Apochromat 20× NA 0.8; Zeiss) with an EMCCD iXon camera

(Andor) through ZEN microscopy software 7.0.4.0 (RRID: SCR_013672; Zeiss).

QUANTIFICATION AND STATISTICAL ANALYSIS

Statistics and data analysis

Results are shown as means ± SEM. For each experiment, at least three
independent biological repetitions were performed. The significance of differences
between the averages were analyzed using one-way ANOVA as described in the
legends (GraphPad Prism 9, RRID: SCR_002798). Figures were prepared with
Photoshop CS6 (Adobe, RRID: SCR_014199), Illustrator CS6 (Adobe, RRID:
SCR_010279), BioRender.com (RRID: SCR_018361), and FIJI (ImageJ 1.53c, RRID:
SCR_002285).

Acknowledgements

We thank Gavin Wright (University of York, York, UK) for the pCMV6-hJuno
plasmid, Kazuo Yamagata (Kindai University, Higashiosaka City, Osaka, Japan) for
the EGFP-MBD-NLS plasmid, Masahito Ikawa (Osaka University, Osaka, Japan) for
the IZUMO1-mCherry transgenic mouse line, and Zene Matsuda (University of Tokyo,
Japan) for the DSP plasmids. Finally, we thank Dan Cassel (Technion- Israel Institute
of Technology), Leonid Chernomordik (NICHD/DIR, NIH, US) and the members of
Podbilewicz Lab for critically reading the manuscript.

This project has received funding from the Israel Science Foundation (257/17,
2462/18, 2327/19, and 178/20 to B. Podbilewicz) and the European Union's Horizon
2020 research and innovation program under the Marie Skłodowska-Curie Actions
(844807 to N.G. Brukman).

Author contributions

B.P. conceived the study. N.G.B. and C.V. designed and conducted experiments,
and processed and analyzed the data supervised by B.P.; N.G.B. wrote the initial draft
of the manuscript. All authors participated in discussions of results and manuscript
editing.

Declaration of interests

N.G.B., C.V. and B.P. are inventors on a patent application filed by the
Technion- Israel Institute of Technology, based on this work.

Figures

Figure 1: Sperm induce syncytia formation of fibroblasts

(A) Scheme of experimental design: Baby Hamster Kidney (BHK) cells were
transfected with either pCI::H2B-RFP or pCI::JUNO::H2B-RFP vectors and
pcDNA3.1-EGFP-MBD-nls. Mouse sperm were obtained from the epididymis of adult
mice and capacitated in HTF capacitating medium. Sperm cells were co-incubated
with the BHK cells for 4 h and then the cells were fixed and stained with DAPI to detect
the DNA. Grey lollipops represent JUNO molecules **(B)** Representative images
showing H2B-RFP (magenta), GFP-MBD (green), DAPI (blue) channels and the
merge. Dotted lines contour relevant cells. Green arrowheads point to fused sperm
(GFP positive). The presence of 200 µg/ml of cycloheximide is indicated (CHX). Scale
bars, 20 µm. **(C)** The percentage of multinucleation was defined as the ratio between
the nuclei in multinucleated cells (NuM) and the total number of nuclei in fluorescent
cells (NuF), as follows: % of multinucleation = (NuM/NuF) × 100. We show individual
data and means ± SEM of six independent experiments. The number of nuclei counted
per experiment and per treatment was 500. For JUNO transfected cells,
multinucleation was counted separately for cells with and without sperm fused to them.

Comparisons were made with one-way ANOVA followed by Tukey's test. ns = non-
significant, ****P < 0.0001. **(D)** In another set of experiments JUNO-transfected cells
were treated with 200 µg/ml CXH to inhibit *de novo* synthesis of proteins. We show
individual data and means ± SEM of five independent experiments. ns = non-
significant, **P < 0.01. **(E)** Time-lapse images from a movie showing sperm-induced
cell-cell fusion. BHK cells were transfected with the pCI::JUNO::GFPnes plasmid and
sperm were added at time=0 min. Arrows and arrowheads indicate contacting and
fused cells, respectively. The green channel (GFPnes) and the DIC images are shown
(see also Movie S1). Scale bars, 20 µm.

<u>Figure 2: Sperm-induced fusion is dependent on JUNO</u>

(A) Scheme of experimental design. BHK cells transfected with pCI::GFPnes or pCI::H2B-RFP empty vectors or containing the coding sequence for the expression of JUNO were mixed as indicated. Later, the cells were co-incubated with sperm for 4 h, fixed and stained with DAPI. (B) Representative images for each treatment. Each separate channel for GFPnes (green cytoplasm), H2B-RFP (magenta nuclei) and DAPI staining (blue) are shown. Scale bars, 20 µm. Fused, mixed cells contain both GFPnes and H2B-RFP staining. (C) Quantification of content mixing experiments. The percentage of mixing was defined as the ratio between the nuclei in mixed cells (NuM) and the total number of nuclei in mixed cells and fluorescent cells in contact that did not fuse (NuC), as follows: % of mixing = (NuM/[NuM + NuC]) × 100. Bar chart showing individual experiment values (each corresponding to 1,000 nuclei) and means ± SEM of six independent experiments. Comparisons by one-way ANOVA followed by Tukey's test. ns = non-significant, ***P < 0.001, ****P < 0.0001.

Figure 3: Syncytia formation requires a functional sperm, is dependent on IZUMO1
and is not affected by an inhibitor of cell division

BHK cells were transfected with pcDNA3.1-EGFP-MBD-nls together with pCI::H2B-
RFP (empty vector) or pCI::JUNO::H2B-RFP and co-incubated with control sperm in
the presence of 20 µM of the inhibitor of cell division FdUdr or of 1 µg/µl of anti-IZUMO1
antibody (clone Mab120). Alternatively, cells were mixed with sperm incubated in a
medium lacking calcium (-Ca^{2+}) or bovine serum albumin (-BSA); both conditions fail
to support sperm capacitation. The number of sperm fused per 500 fluorescent cells
(A) and the percentage of multinucleation **(B)** were determined. Bar charts showing
individual experiment values and means ± SEM of five independent experiments.
Comparisons by one-way ANOVA followed by Tukey's test. In red are the comparisons
against the empty vector. ns = non-significant, ***P < 0.001, ****P < 0.0001.

<u>Figure 4: Sperm induce BHK-BHK fusion in a species-specific manner.</u>

BHK cells were transfected with pcDNA3.1-EGFP-MBD-nls together pCI::H2B-RFP
(empty vector) or pCI::JUNO::H2B-RFP encoding for human JUNO (hJUNO) or mouse
JUNO (mJUNO), and co-incubated with capacitated mouse sperm. **(A)** Representative
images showing H2B-RFP (magenta), GFP-MBD (green), DAPI (blue) channels and
the merge. Dotted lines contour a cell transfected with human JUNO with two mouse
sperm fused (arrowheads). Scale bar, 20 µm. The number of sperm fused per 500
fluorescent cells **(B)** and the percentage of multinucleation **(C)** were determined. Bar
charts showing individual experiment values and means ± SEM of five independent
experiments. Comparisons by one-way ANOVA followed by Tukey's test. *P < 0.05,
**P < 0.01, ***P < 0.001.

Figure 5: Sperm-induced fusion of human cells evaluated by Dual Split Proteins (DSP)

(A) Scheme of experimental design. HEK293T cells stably expressing the split GFP (DSP1-7 or DSP8-11) were mixed and transfected with pCI::H2B-RFP or pCI::JUNO::H2B-RFP vectors. When indicated, the cells were later co-incubated with sperm for 4 h, fixed and stained with DAPI. (B) Representative images for each treatment. Each separate channel for GFP (assembled GFP in fused cells), H2B-RFP (magenta nuclei) and DAPI staining (blue) are shown. Scale bars, 20 μm. (C) Quantification of content mixing experiments. The extent of fusion was determined by counting the number of GFP positive cells per 1,000 nuclei. Bar chart showing individual experiment values (each corresponding to 1,000 to 2,000 nuclei) and means ± SEM of at least four independent experiments. Comparisons by one-way ANOVA followed by Tukey's test. ns = non-significant, ****P < 0.0001.

<u>Figure 6: Summary of the SPICER method</u>

Schematic representation of the multinucleation assay employed to determine the
sperm fusogenic potential. (1) BHK are seeded on a plate. (2) The cells are transfected
with the plasmids encoding for JUNO and EGFP-MBD. (3) Mouse sperm cells are
collected and capacitated. (4) The BHK and sperm cells are co-incubated. (5) The
cells are washed, fixed and stained with DAPI. (6) Multinucleation levels are quantified
as indicated.

 Supplementary Information

Figure S1: IZUMO1 diffuses into the target cell upon sperm fusion

(A) BHK were transfected with pcDNA3.1-EGFP-MBD-nls alone or together with pCI::JUNO::H2B-RFP and co-incubated with sperm obtained from transgenic mice expressing IZUMO1-mcherry. Representative images showing H2B-RFP and Izumo1-mCherry (magenta), GFP-MBD (green), DAPI (blue) channels and the merge with and without DAPI. Dotted lines contour relevant cells. Scale bar, 20 µm. Magenta arrowheads point to unfused sperm (GFP negative) presenting IZUMO1-mCherry signal while green arrowheads point to fused sperm (GFP positive) without Izumo1-mCherry signal. **(B)** BHK were transfected with pcDNA3.1-EGFP-MBD-nls together with pCI::JUNO::H2B-RFP and co-incubated with wild-type mouse sperm. After fixation, an immunostaining against IZUMO1 was performed. Representative images showing H2B-RFP (magenta), GFP-MBD (green), IZUMO1 staining (yellow), DAPI (blue) channels and the merge with and without DAPI. Dotted lines contour relevant cells. Scale bar, 20 µm. Yellow arrows point to unfused sperm (GFP negative)

presenting IZUMO1 signal while green arrowheads point to fused sperm (GFP positive) without IZUMO1 signal. **(C)** Models for sperm-induced cell-cell fusion. In general, this phenomenon resembles the so-called "fusion from without" in which viruses can induce the fusion of the target cells (Falke, Knoblich, and Müller 1985; Tang et al. 2021; Edwards and Brown 1986). i- One sperm fuses with a BHK, IZUMO1 is transferred by diffusion and mediates the fusion to another cell. ii- One sperm fuses to two BHK bridging them and inducing the formation of a syncytium.

Movie S1: Time-lapse experiment using spinning disk confocal microscopy showing the fusion of two cells expressing JUNO and GFPnes after the addition of sperm at t=0 min. Time in hours:minutes. Green and DIC channels are shown. Scale bars, 20 µm.

References

Abay, Zehra C., Michelle Yu-Ying Wong, Jean-Sébastien Teoh, Tarika Vijayaraghavan, Massimo A. Hilliard, and Brent Neumann. 2017. "Phosphatidylserine Save-Me Signals Drive Functional Recovery of Severed Axons in Caenorhabditis Elegans." *Proceedings of the National Academy of Sciences of the United States of America* 114 (47): E10196–205.

Aitken, R. J., and R. A. Elton. 1986. "Quantitative Analysis of Sperm-Egg Interaction in the Zona-Free Hamster Egg Penetration Test." *International Journal of Andrology* 9 (s6): 14–30.

Allingham, Jessica, and Wely B. Floriano. 2021. "Genetic Diversity in the IZUMO1-JUNO Protein-Receptor Pair Involved in Human Reproduction." *PloS One* 16 (12): e0260692.

Aydin, Halil, Azmiri Sultana, Sheng Li, Annoj Thavalingam, and Jeffrey E. Lee. 2016. "Molecular Architecture of the Human Sperm IZUMO1 and Egg JUNO Fertilization Complex." *Nature* 534 (7608): 562–65.

Barbaux, Sandrine, Côme Ialy-Radio, Myriam Chalbi, Elisa Dybal, Méline Homps-Legrand, Marcio Do Cruzeiro, Daniel Vaiman, Jean-Philippe Wolf, and Ahmed Ziyyat. 2020. "Sperm SPACA6 Protein Is Required for Mammalian Sperm-Egg Adhesion/Fusion." *Scientific Reports* 10 (1): 5335.

Bendich, A., E. Borenfreund, and S. S. Sternberg. 1974. "Penetration of Somatic Mammalian Cells by Sperm." *Science* 183 (4127): 857–59.

Bianchi, Enrica, Brendan Doe, David Goulding, and Gavin J. Wright. 2014. "Juno Is the Egg Izumo Receptor and Is Essential for Mammalian Fertilization." *Nature* 508: 483–87.

Bianchi, Enrica, and Gavin J. Wright. 2015. "Cross-Species Fertilization: The Hamster Egg Receptor, Juno, Binds the Human Sperm Ligand, Izumo1." *Philosophical Transactions of the Royal Society of London. Series B, Biological Sciences* 370(1661): 20140101.

Binner, Mirjam I., Anna Kogan, Karin Panser, Alexander Schleiffer, Victoria E. Deneke, and Andrea Pauli. 2021. "The Sperm Protein Spaca6 Is Essential for Fertilization in Zebrafish." *Frontiers in Cell and Developmental Biology* 9: 806982.

Bratt, Michael A., and William R. Gallaher. 1969. "Preliminary analysis of the requirements for fusion from within and fusion from without by newcastle disease virus*." *Proceedings of the National Academy of Sciences* 64 (2): 536–43.

Brukman, Nicolas G., Xiaohui Li, and Benjamin Podbilewicz. 2022. "Fusexins, HAP2/GCS1 and Evolution of Gamete Fusion." *Frontiers in Cell and Developmental Biology* 9: 824024.

Brukman, Nicolas G., Kohdai P. Nakajima, Clari Valansi, Kateryna Flyak, Xiaohui Li, Tetsuya Higashiyama, and Benjamin Podbilewicz. 2023. "A Novel Function for the Sperm Adhesion Protein IZUMO1 in Cell-Cell Fusion." *J Cell Biol. 2023 Feb 6;222(2):e202207147.*

Brukman, Nicolas G., Berna Uygur, Benjamin Podbilewicz, and Leonid V. Chernomordik. 2019. "How Cells Fuse." *J Cell Biol.* 218 (5): 1436–51.

Buchrieser, Julian, Jérémy Dufloo, Mathieu Hubert, Blandine Monel, Delphine Planas, Maaran Michael Rajah, Cyril Planchais, et al. 2020. "Syncytia Formation by SARS-CoV-2-Infected Cells." *The EMBO Journal* 39 (23): e106267.

Ciechonska, Marta, Tim Key, and Roy Duncan. 2014. "Efficient Reovirus- and Measles Virus-Mediated Pore Expansion during Syncytium Formation Is Dependent on Annexin A1 and Intracellular Calcium." *Journal of Virology* 88 (11): 6137–47.

Clark, Sydney, and Rajesh K. Naz. 2013. "Presence and Incidence of Izumo Antibodies in Sera of Immunoinfertile Women and Men." *Am J Reprod Immunol. 2013 Mar;69(3):256-63.*

Deneke, Victoria E., and Andrea Pauli. 2021. "The Fertilization Enigma: How Sperm and Egg Fuse." *Annual Review of Cell and Developmental Biology* 37 (October): 391–414.

Duelli, Dominik, and Yuri Lazebnik. 2007. "Cell-to-Cell Fusion as a Link between Viruses and Cancer." *Nature Reviews. Cancer* 7 (12): 968–76.

Earles, C. A., J. Bai, P. Wang, and E. R. Chapman. 2001. "The Tandem C2 Domains of Synaptotagmin Contain Redundant Ca2+ Binding Sites That Cooperate to Engage T-SNAREs and Trigger Exocytosis." *The Journal of Cell Biology* 154 (6): 1117–23.

Edwards, J., and D. T. Brown. 1986. "Sindbis Virus-Mediated Cell Fusion from without Is a Two-Step Event." *The Journal of General Virology* 67 (Pt 2) (February): 377–80.

Eigler, Tamar, Giulia Zarfati, Emmanuel Amzallag, Sansrity Sinha, Nadav Segev, Yishaia Zabary, Assaf Zaritsky, et al. 2021. "ERK1/2 Inhibition Promotes Robust Myotube Growth via CaMKII Activation Resulting in Myoblast-to-Myotube Fusion." *Developmental Cell* 56 (24): 3349–63.e6.

Eijnde, S. M. van den, M. J. van den Hoff, C. P. Reutelingsperger, W. L. van Heerde, M. E. Henfling, C. Vermeij-Keers, B. Schutte, M. Borgers, and F. C. Ramaekers. 2001. "Transient Expression of Phosphatidylserine at Cell-Cell Contact Areas Is Required for Myotube Formation." *Journal of Cell Science* 114 (Pt 20): 3631–42.

Enoiu, Simona Ioana, Marie Berg Nygaard, Mona Bungum, Søren Ziebe, Morten R. Petersen, and Kristian Almstrup. 2022. "Expression of Membrane Fusion Proteins in Spermatozoa and Total Fertilisation Failure during in Vitro Fertilisation." *Andrology.* 2022 Oct;10(7):1317-1327.

Falke, D., A. Knoblich, and S. Müller. 1985. "Fusion from without Induced by Herpes Simplex Virus Type 1." *Intervirology* 24 (4): 211–19.

Ferris, P. J., J. P. Woessner, and U. W. Goodenough. 1996. "A Sex Recognition Glycoprotein Is Encoded by the plus Mating-Type Gene fus1 of Chlamydomonas Reinhardtii." *Molecular Biology of the Cell* 7 (8): 1235.

Fujihara, Yoshitaka, Yonggang Lu, Taichi Noda, Asami Oji, Tamara Larasati, Kanako Kojima-Kita, Zhifeng Yu, Ryan M. Matzuk, Martin M. Matzuk, and Masahito Ikawa. 2020. "Spermatozoa Lacking Fertilization Influencing Membrane Protein (FIMP) Fail to Fuse with Oocytes in Mice." *Proceedings of the National Academy of Sciences of the United States of America* 117 (17): 9393–9400.

Gert, Krista R. B., Karin Panser, Joachim Surm, Benjamin S. Steinmetz, Alexander Schleiffer, Luca Jovine, Yehu Moran, Fyodor Kondrashov, and Andrea Pauli. 2023. "Divergent Molecular Signatures in Fish Bouncer Proteins Define Cross-Fertilization Boundaries." *Nature Communications* 14 (1): 3506.

Grayson, Phil. 2015. "Izumo1 and Juno: The Evolutionary Origins and Coevolution of Essential Sperm–egg Binding Partners." *R Soc Open Sci.* 2015 Dec 16;2(12):150296.

Herberg, Sarah, Krista R. Gert, Alexander Schleiffer, and Andrea Pauli. 2018. "The Ly6/uPAR Protein Bouncer Is Necessary and Sufficient for Species-Specific Fertilization." *Science* 361 (6406): 1029.

Hino, Toshiaki, Yuko Muro, Miwa Tamura-Nakano, Masaru Okabe, Hiroyuki Tateno, and Ryuzo Yanagimachi. 2016. "The Behavior and Acrosomal Status of Mouse Spermatozoa In Vitro, and Within the Oviduct During Fertilization after Natural

Mating." *Biology of Reproduction* 95 (3): 50.

Inoue, Naokazu, Yoshihisa Hagihara, and Ikuo Wada. 2021. "Evolutionarily Conserved Sperm Factors, DCST1 and DCST2, Are Required for Gamete Fusion." *Elife.* 2021 Apr 19;10:e66313.

Inoue, Naokazu, Yoshihisa Hagihara, Danelle Wright, Takahisa Suzuki, and Ikuo Wada. 2015. "Oocyte-Triggered Dimerization of Sperm IZUMO1 Promotes Sperm-Egg Fusion in Mice." *Nature Communications* 6: 8858.

Inoue, Naokazu, Daizo Hamada, Hironari Kamikubo, Kunio Hirata, Mikio Kataoka, Masaki Yamamoto, Masahito Ikawa, Masaru Okabe, and Yoshihisa Hagihara. 2013. "Molecular Dissection of IZUMO1, a Sperm Protein Essential for Sperm-Egg Fusion." *Development* 140 (15): 3221–29.

Inoue, Naokazu, Masahito Ikawa, Ayako Isotani, and Masaru Okabe. 2005. "The Immunoglobulin Superfamily Protein Izumo Is Required for Sperm to Fuse with Eggs." *Nature* 434 (7030): 234–38.

Ishikawa, Hirohito, Fanxia Meng, Naoyuki Kondo, Aikichi Iwamoto, and Zene Matsuda. 2012. "Generation of a Dual-Functional Split-Reporter Protein for Monitoring Membrane Fusion Using Self-Associating Split GFP." *Protein Engineering, Design & Selection: PEDS* 25 (12): 813–20.

Kito, Seiji, Tatsuo Hayao, Yoshiko Noguchi-Kawasaki, Yuki Ohta, Uhara Hideki, and Shintaro Tateno. 2004. "Improved in Vitro Fertilization and Development by Use of Modified Human Tubal Fluid and Applicability of Pronucleate Embryos for Cryopreservation by Rapid Freezing in Inbred Mice." *Comparative Medicine* 54 (5): 564–70.

Kohn, A. 1965. "Polykaryocytosis induced by newcastle disease virus in monolayers of animal cells." *Virology* 26 (June): 228–45.

Lamas-Toranzo, Ismael, Julieta G. Hamze, Enrica Bianchi, Beatriz Fernández-Fuertes, Serafín Pérez-Cerezales, Ricardo Laguna-Barraza, Raúl Fernández-González, et al. 2020. "TMEM95 Is a Sperm Membrane Protein Essential for Mammalian Fertilization." *Elife.* 2020 Jun 2;9:e53913.

La Spina, Florenza A., Lis C. Puga Molina, Ana Romarowski, Alejandra M. Vitale, Tomas L. Falzone, Dario Krapf, Noritaka Hirohashi, and Mariano G. Buffone. 2016. "Mouse Sperm Begin to Undergo Acrosomal Exocytosis in the Upper Isthmus of the Oviduct." *Developmental Biology* 411 (2): 172–82.

Lorenzetti, Diego, Christophe Poirier, Ming Zhao, Paul A. Overbeek, Wilbur Harrison, and Colin E. Bishop. 2014. "A Transgenic Insertion on Mouse Chromosome 17 Inactivates a Novel Immunoglobulin Superfamily Gene Potentially Involved in Sperm-Egg Fusion." *Mammalian Genome: Official Journal of the International Mammalian Genome Society* 25: 141–48.

Lu, Yonggang, and Masahito Ikawa. 2022. "Eukaryotic Fertilization and Gamete Fusion at a Glance." *Journal of Cell Science* 135 (22). https://doi.org/10.1242/jcs.260296.

Matsumura, Takafumi, Taichi Noda, Yuhkoh Satouh, Akane Morohoshi, Shunsuke Yuri, Masaki Ogawa, Yonggang Lu, Ayako Isotani, and Masahito Ikawa. 2021. "Sperm IZUMO1 Is Required for Binding Preceding Fusion with Oolemma in Mice and Rats." *Frontiers in Cell and Developmental Biology* 9: 3849.

Mattioli, M., A. Gloria, A. Mauro, L. Gioia, and B. Barboni. 2009. "Fusion as the Result of Sperm-Somatic Cell Interaction." *Reproduction* 138 (4): 679–87.

Moi, David, Shunsuke Nishio, Xiaohui Li, Clari Valansi, Mauricio Langleib, Nicolas G. Brukman, Kateryna Flyak, et al. 2022. "Discovery of Archaeal Fusexins Homologous to Eukaryotic HAP2/GCS1 Gamete Fusion Proteins." *Nature*

Communications 13 (1): 3880.

Muro, Yuko, Hidetoshi Hasuwa, Ayako Isotani, Haruhiko Miyata, Kazuo Yamagata,
Masahito Ikawa, Ryuzo Yanagimachi, and Masaru Okabe. 2016. "Behavior of
Mouse Spermatozoa in the Female Reproductive Tract from Soon after Mating to
the Beginning of Fertilization." *Biology of Reproduction* 94 (4): 80.

Nakane, Shuhei, and Zene Matsuda. 2015. "Dual Split Protein (DSP) Assay to Monitor
Cell-Cell Membrane Fusion." In *Cell Fusion: Overviews and Methods: Second*
Edition, Second, 229–36. Springer New York.

Nakajima KP, Valansi C, Kurihara D, Sasaki N, Podbilewicz B, Higashiyama T. Live
imaging-based assay for visualising species-specific interactions in gamete
adhesion molecules. Sci Rep. 2022 Jun 10;12(1):9609.

Nishimura, Hitoshi, Tatsuya Tajima, Heather Skye Comstra, Elizabeth J. Gleason, and
Steven W. L'Hernault. 2015. "The Immunoglobulin-like Gene Spe-45 Acts during
Fertilization in Caenorhabditis Elegans like the Mouse Izumo1 Gene." *Current*
Biology: CB 25 (24): 3225.

Nishimura, Kaoru, Ling Han, Enrica Bianchi, Gavin J. Wright, Daniele de Sanctis, and
Luca Jovine. 2016. "The Structure of Sperm Izumo1 Reveals Unexpected
Similarities with Plasmodium Invasion Proteins." *Current Biology: CB* 26 (14):
R661–62.

Noda, Taichi, Andreas Blaha, Yoshitaka Fujihara, Krista R. Gert, Chihiro Emori,
Victoria E. Deneke, Seiya Oura, et al. 2022. "Sperm Membrane Proteins DCST1
and DCST2 Are Required for Sperm-Egg Interaction in Mice and Fish."
Communications Biology 5 (1): 332.

Noda, Taichi, Yonggang Lu, Yoshitaka Fujihara, Seiya Oura, Takayuki Koyano,
Sumire Kobayashi, Martin M. Matzuk, and Masahito Ikawa. 2020. "Sperm
Proteins SOF1, TMEM95, and SPACA6 Are Required for Sperm-Oocyte Fusion
in Mice." *Proceedings of the National Academy of Sciences of the United States*
of America 117 (21): 11493–502.

Ohto, Umeharu, Hanako Ishida, Elena Krayukhina, Susumu Uchiyama, Naokazu
Inoue, and Toshiyuki Shimizu. 2016. "Structure of IZUMO1–JUNO Reveals
Sperm–oocyte Recognition during Mammalian Fertilization." *Nature* 534 (7608):
566–69.

Okada, Y. 1962. "Analysis of Giant Polynuclear Cell Formation Caused by HVJ Virus
from Ehrlich's Ascites Tumor Cells. I. Microscopic Observation of Giant
Polynuclear Cell Formation." *Experimental Cell Research* 26 (February): 98–107.

Palermo, G., H. Joris, P. Devroey, and A. C. Van Steirteghem. 1992. "Pregnancies
after Intracytoplasmic Injection of Single Spermatozoon into an Oocyte." *The*
Lancet 340 (8810): 17–18.

Pinello, Jennifer F., Yanjie Liu, and William J. Snell. 2021. "MAR1 Links Membrane
Adhesion to Membrane Merger during Cell-Cell Fusion in Chlamydomonas."
Dev Cell. 2021 Dec 20;56(24):3380-3392.e9.

Practice Committees of the American Society for Reproductive Medicine and the
Society for Assisted Reproductive Technology. 2020. "Intracytoplasmic Sperm
Injection (ICSI) for Non-Male Factor Indications: A Committee Opinion." *Fertility*
and Sterility 114 (2): 239–45.

Rival, Claudia M., Wenhao Xu, Laura S. Shankman, Sho Morioka, Sanja Arandjelovic,
Chang Sup Lee, Karen M. Wheeler, et al. 2019. "Phosphatidylserine on Viable
Sperm and Phagocytic Machinery in Oocytes Regulate Mammalian Fertilization."
Nature Communications 10 (1): 4456.

Satouh, Yuhkoh, Naokazu Inoue, Masahito Ikawa, and Masaru Okabe. 2012.

"Visualization of the Moment of Mouse Sperm-Egg Fusion and Dynamic Localization of IZUMO1." *Journal of Cell Science* 125 (Pt 21): 4985–90.

Shilagardi, Khurts, Shuo Li, Fengbao Luo, Faiz Marikar, Rui Duan, Peng Jin, Ji Hoon Kim, Katherine Murnen, and Elizabeth H. Chen. 2013. "Actin-Propelled Invasive Membrane Protrusions Promote Fusogenic Protein Engagement during Cell-Cell Fusion." *Science* 340 (6130): 359–63.

Spiegelstein, O., J. D. Eudy, and R. H. Finnell. 2000. "Identification of Two Putative Novel Folate Receptor Genes in Humans and Mouse." *Gene* 258 (1-2): 117–25.

Stepanenko, Nataliia, Omri Wolk, Enrica Bianchi, Gavin James Wright, Natali Schachter-Safrai, Kiril Makedonski, Alberto Ouro, Assaf Ben-Meir, Yosef Buganim, and Amiram Goldblum. 2022. "In Silico Docking Analysis for Blocking JUNO-IZUMO1 Interaction Identifies Two Small Molecules That Block in Vitro Fertilization." *Frontiers in Cell and Developmental Biology* 10 (April): 824629.

Steptoe, P. C., and R. G. Edwards. 1978. "Birth after the Reimplantation of a Human Embryo." *The Lancet* 2 (8085): 366.

Suzuki, Banri, Yukou Sugano, Jun Ito, Haruka Saito, Sueo Niimura, and Hideaki Yamashiro. 2017. "Location and Expression of Juno in Mice Oocytes during Maturation." *JBRA Assisted Reproduction* 21 (4): 321–26.

Takaiso, Nobue, Haruki Nishizawa, Sachie Nishiyama, Tomio Sawada, Eriko Hosoba, Tamae Ohye, Tsutomu Sato, Hidehito Inagaki, and Hiroki Kurahashi. 2016. "Mutation Analysis of the *JUNO* Gene in Female Infertility of Unknown Etiology." *Fujita Medical Journal* 2 (3): 59–61.

Takayama, Jun, Tatsuya Tajima, Shuichi Onami, and Hitoshi Nishimura. 2021. "C. Elegans Spermatozoa Lacking Spe-45 Are Incapable of Fusing with the Oocyte Plasma Membrane." *MicroPubl Biol.* 2021: 10.17912/micropub.biology.000372.

Tang, Jiajia, Giada Frascaroli, Xuan Zhou, Jan Knickmann, and Wolfram Brune. 2021. "Cell Fusion and Syncytium Formation in Betaherpesvirus Infection." *Viruses.* 2021 Oct; 13(10): 1973.

Tateno, Hiroyuki, Dario Krapf, Toshiaki Hino, Claudia Sánchez-Cárdenas, Alberto Darszon, Ryuzo Yanagimachi, and Pablo E. Visconti. 2013. "Ca2+ Ionophore A23187 Can Make Mouse Spermatozoa Capable of Fertilizing in Vitro without Activation of cAMP-Dependent Phosphorylation Pathways." *Proceedings of the National Academy of Sciences of the United States of America* 110 (46): 18543–48.

Theuerkauf, Samuel A., Alexander Michels, Vanessa Riechert, Thorsten J. Maier, Egbert Flory, Klaus Cichutek, and Christian J. Buchholz. 2021. "Quantitative Assays Reveal Cell Fusion at Minimal Levels of SARS-CoV-2 Spike Protein and Fusion from without." *iScience* 24 (3): 102170.

Tsuchiya, Masaki, Yuji Hara, Masaki Okuda, Karin Itoh, Ryotaro Nishioka, Akifumi Shiomi, Kohjiro Nagao, et al. 2018. "Cell Surface Flip-Flop of Phosphatidylserine Is Critical for PIEZO1-Mediated Myotube Formation." *Nature Communications* 9 (1): 2049.

Valansi, Clari, David Moi, Evgenia Leikina, Elena Matveev, Martín Graña, Leonid V. Chernomordik, Héctor Romero, Pablo S. Aguilar, and Benjamin Podbilewicz. 2017. "Arabidopsis HAP2/GCS1 Is a Gamete Fusion Protein Homologous to Somatic and Viral Fusogens." *The Journal of Cell Biology* 216: 571–81.

Vance, Tyler D. R., and Jeffrey E. Lee. 2020. "Virus and Eukaryote Fusogen Superfamilies." *Current Biology: CB* 30 (13): R750–54.

Verma, Santosh K., Evgenia Leikina, Kamran Melikov, Claudia Gebert, Vardit Kram, Marian F. Young, Berna Uygur, and Leonid V. Chernomordik. 2018. "Cell-Surface

Phosphatidylserine Regulates Osteoclast Precursor Fusion." *The Journal of Biological Chemistry* 293 (1): 254–70.

Visconti, Pablo E., Janice L. Bailey, Grace D. Moore, Dieyun Pan, Patricia Olds-Clarke, and Gregory S. Kopf. 1995. "Capacitation of Mouse Spermatozoa. I. Correlation between the Capacitation State and Protein Tyrosine Phosphorylation." *Development* 121: 1129–37.

Wang, Hongyun, Xiao Li, Shuhei Nakane, Shujun Liu, Hirohito Ishikawa, Aikichi Iwamoto, and Zene Matsuda. 2014. "Co-Expression of Foreign Proteins Tethered to HIV-1 Envelope Glycoprotein on the Cell Surface by Introducing an Intervening Second Membrane-Spanning Domain." *PloS One* 9 (5): e96790.

Wengler, G. 1975. "Studies on the Synthesis of Viral RNA-Polymerase-Template Complexes in BHK 21 Cells Infected with Semliki Forest Virus." *Virology* 66 (1): 322–26.

Williams, Ruth M., Upeka Senanayake, Mara Artibani, Gunes Taylor, Daniel Wells, Ahmed Ashour Ahmed, and Tatjana Sauka-Spengler. 2018. "Genome and Epigenome Engineering CRISPR Toolkit for in Vivo Modulation of Cis-Regulatory Interactions and Gene Expression in the Chicken Embryo." *Development* 145 (4). https://doi.org/10.1242/dev.160333.

Wood, Laura, and Gavin J. Wright. 2019. "High-Content Imaging for Large-Scale Detection of Low-Affinity Extracellular Protein Interactions." *SLAS Discovery : Advancing Life Sciences R & D* 24 (10): 987–99.

World Health Organization. 2023. *Infertility Prevalence Estimates, 1990–2021*. Edited by World Health Organization. Geneva.

World Health Organization, HRP. 2021. "WHO Laboratory Manual for the Examination and Processing of Human Semen." Sixth Edition. World Health Organization.

Yamagata, Kazuo, Taiga Yamazaki, Misuzu Yamashita, Yuki Hara, Narumi Ogonuki, and Atsuo Ogura. 2005. "Noninvasive Visualization of Molecular Events in the Mammalian Zygote." *Genesis* 43 (2): 71–79.

Yamaguchi, Tomoyuki, Keiji Hirota, Kanji Nagahama, Katsuya Ohkawa, Takeshi Takahashi, Takeshi Nomura, and Shimon Sakaguchi. 2007. "Control of Immune Responses by Antigen-Specific Regulatory T Cells Expressing the Folate Receptor." *Immunity* 27 (1): 145–59.

Yanagimachi, R., H. Yanagimachi, and B. J. Rogers. 1976. "The Use of Zona-Free Animal Ova as a Test-System for the Assessment of the Fertilizing Capacity of Human Spermatozoa." *Biology of Reproduction* 15 (4): 471–76.

Yanagimachi, Ryuzo. 1988. "Chapter 1 Sperm–Egg Fusion." In *Current Topics in Membranes and Transport*, edited by Felix Bronner, 32:3–43. Academic Press.

Yanagimachi, Ryuzo. 2022. "Mysteries and Unsolved Problems of Mammalian Fertilization and Related Topics." *Biology of Reproduction* 106 (4): 644–75.

Yang, Yihong, Yan Zhang, Wen Jun Li, Yuxiang Jiang, Zhiwen Zhu, Huifang Hu, Wei Li, et al. 2017. "Spectraplakin Induces Positive Feedback between Fusogens and the Actin Cytoskeleton to Promote Cell-Cell Fusion." *Developmental Cell* 41 (1): 107–20.e4.

Yu, Mengru, Han Zhao, Tailai Chen, Ye Tian, Mei Li, Keliang Wu, Yuehong Bian, et al. 2018. "Mutational Analysis of IZUMO1R in Women with Fertilization Failure and Polyspermy after in Vitro Fertilization." *Journal of Assisted Reproduction and Genetics* 35 (3): 539–44.

Zainul Rashid, M. R., S. B. Fishel, S. Thornton, J. A. Hall, G. Ndukwe, M. Aloum, and S. D. Fleming. 1998. "The Predictive Value of the Zona-Free Hamster Egg Penetration Test in Relation to in-Vitro Fertilization at Various Insemination

875   Concentrations." *Human Reproduction* 13 (3): 624–29.

876 Zhang, Yan, Yihong Yang, Zhiwen Zhu, and Guangshuo Ou. 2017. "WASP-Arp2/3-

877   Dependent Actin Polymerization Influences Fusogen Localization during Cell-Cell

878   Fusion in Caenorhabditis elegans Embryos." *Biology Open* 6 (9): 1324–28.