1    **Oxygen isotopes in orangutan teeth reveal recent and ancient climate variation**
2

3    Tanya M. Smith[1,2], Manish Arora[3], Christine Austin[3], Janaína N. Ávila[1,4], Mathieu Duval[2,5,6],
4    Tze Tshen Lim[7], Philip J. Piper[8], Petra Vaiglova[1,2,8], John de Vos[9], Ian S. Williams[10], Jian-
5    xin Zhao[4,11], Daniel R. Green[2,12]
6
1 Griffith Centre for Social and Cultural Research, Griffith University, Australia
2 Australian Research Centre for Human Evolution, Griffith University, Australia
3 Department of Environmental Medicine and Public Health, Icahn School of Medicine at
Mount Sinai, USA
4 School of Earth and Environmental Sciences, University of Queensland, Australia
5 Centro Nacional de Investigación sobre la Evolución Humana (CENIEH), Spain
6 Palaeoscience Labs, Dept. Archaeology and History, La Trobe University, Australia
7 Department of Geology, Universiti Malaya, Malaysia
8 School of Archaeology and Anthropology, The Australian National University, Australia
9 Department of Geology, Naturalis Biodiversity Center, The Netherlands
10 Research School of Earth Sciences, The Australian National University, Australia
11 Radiogenic Isotope Facility, School of Earth and Environmental Sciences, University of
Queensland, Australia
12 Lamont-Doherty Earth Observatory, Earth Institute and Climate School, Columbia
University, USA
22

**Abstract**

Studies of climate variation commonly rely on chemical and isotopic changes

recorded in sequentially-produced growth layers, such as in corals, shells and tree rings, as

well as in accretionary deposits—ice and sediment cores, and speleothems. Oxygen isotopic

compositions ($\delta^{18}O$) of tooth enamel are a direct method of reconstructing environmental

variation experienced by an individual animal. Here we utilize long-forming orangutan

dentitions (*Pongo* spp.) to probe recent and ancient rainfall trends on a weekly basis over ~

3–11 years per individual. We first demonstrate the lack of any consistent isotopic

enrichment effect during exclusive nursing, supporting the use of primate first molar teeth as

environmental proxies. Comparisons of $\delta^{18}O$ values (n = 2016) in six modern Bornean and

Sumatran orangutans reveal a high degree of overlap, with more consistent annual and

bimodal rainfall patterns in the Sumatran individuals. Comparisons with fossil orangutan

$\delta^{18}O$ values (n = 955) reveal similarities between modern and late Pleistocene fossil Sumatran

individuals, but differences between modern and late Pleistocene/early Holocene Bornean

orangutans. These suggest drier and more open environments with reduced monsoon intensity

during this earlier period in northern Borneo, consistent with other Niah Caves studies and

long-term speleothem $\delta^{18}O$ records in the broader region. This approach can be extended to

test hypotheses about the paleoenvironments that early humans encountered in southeast

Asia.

**Introduction**

Present-day rainfall patterns in Indonesia are controlled by the Asian and Australian monsoon systems, yielding annual trends that vary considerably with geography, topography, and the direction of monsoonal winds (Aldrian and Susanto, 2003; Moron et al., 2009; Qian et al., 2013; Belgaman et al., 2017). Northern Sumatra and western Borneo experience high annual rainfall and relatively stable annual temperatures, with a bimodal distribution of rainfall governed by the Intertropical Convergence Zone (van Schaik, 1986; Aldrian and Susanto, 2003; Belgaman et al., 2017). These islands are also under the influence of inter-annual climate fluctuations driven by the El-Niño Southern Oscillation (ENSO); a periodic coupling of atmospheric and oceanic temperature gradients that initiates in the tropical Pacific, and influences global temperature and precipitation trends (McPhaden et al., 2006).

It is well understood that variation in rainfall patterns influences the fundamental structure of primate habitats (Brockman and van Schaik, 2005; Wessling et al., 2018). Dense tropical forests are sustained by fairly consistent rainfall and short, irregular dry seasons, while woodland communities in more arid environments have smaller trees, less dense canopies, and more deciduous trees (Vico et al., 2017; Archibald et al., 2019). In regions with prolonged dry seasons, low annual rainfall and savannah landscapes abound, in addition to disturbances such as wildfires (Pletcher et al., 2022).

Open woodland and savannah environments are unfavorable for slow-moving orangutans, the largest mammal with an arboreal lifestyle, particularly in regions with predators such as tigers or humans (Thorpe and Crompton, 2009; Ashbury et al., 2015; Spehar et al., 2018). Supra-annual ENSO events may also impact orangutan energy balance, reproduction, and social organization through the inducement of mast-fruiting, or dramatic seed production events in dipterocarp forests (Knott 1998; Curran et al., 1999; Marshall et al., 2009). Such climate fluctuations over the past several hundred years have been documented in coral isotopes and tree-ring analyses, revealing especially marked changes during the past few decades (Cole et al., 1993; Stahle et al., 1998; Hughen et al., 1999; Urban et al., 2000; Tudhope et al., 2001; Pumijumnong et al., 2020).

Detailed climate records prior to the era of human-induced climate change are somewhat limited for island southeast Asia, but they are directly relevant to understanding the recent distribution of orangutans, and the arrival and dispersal of modern humans in the region during the Late Pleistocene (e.g., Piper 2016; Bae et al., 2017; Spehar et al., 2018). A small number of studies of fossil corals, molluscs, marine sediments, and speleothems has

provided insights into the last interglacial and glacial periods (e.g., Hughen et al., 1999;

Tudhope et al., 2001; Stephens et al., 2016; Yang et al., 2016; Buckingham et al. 2022). For

example, oxygen isotopes in fossil corals from seven periods during the last 130,000 years

suggest that ENSO activity in the western Pacific over that time was comparable to modern

records, although there was variation in the intensity of such activity at different timepoints

(Tudhope et al., 2001). This study was also able to resolve bimodal annual rainfall peaks in

modern corals, yet such detailed subannual records are extremely uncommon, particularly

from terrestrial environments where early humans once lived alongside orangutans and other

mammals.

*Oxygen isotope studies for paleoenvironmental reconstruction*

Oxygen isotope values ($\delta^{18}O$) in water vary with latitude, altitude, temperature and

precipitation cycles, and are also impacted by precipitation sources. In tropical regions the

primary determinant of rainfall isotope compositions is rainfall amount (Dansgaard, 1964;

Rozanski et al., 1993; Belgaman et al., 2017). During wet seasons, rainfall $\delta^{18}O$ values are

relatively low, while the opposite pattern is evident in periods with less rain, although other

meteorological factors can influence isotope values as well (Belgaman et al., 2016). This

primary tropical pattern influences isotopic variation in meteoric, surface, and leaf waters,

which may show further elevations in $\delta^{18}O$ values during dryer periods due to preferential

evaporative loss of the lighter isotope, $^{16}O$ (da Silveira et al., 1989; Bowen, 2010; Roberts et

al., 2017).

In addition to $\delta^{18}O$ values in fossil corals, tree rings and speleothems, other fine-

scaled oxygen isotopic climate proxies include otoliths (fish ear bones) and mollusc shells

(e.g., Aubert et al., 2012; Stephens et al., 2016; Prendergast et al., 2018)—although these are

rarely preserved in rainforest environments. Records of $\delta^{18}O$ values in mammalian tooth

enamel are a more direct means of studying seasonality (reviewed in Green et al., 2018,

2022), providing insight into the actual climates experienced by individuals, in contrast to

indirect proxies for which it can be difficult to establish concurrence. Unlike bone, teeth do

not remodel during life, and the phosphate component of the enamel mineral (hydroxyapatite)

is especially resistant to modification after burial (reviewed in Smith et al., 2018a; Pederzani

and Britton, 2019).

Tooth enamel is most commonly sampled with hand-held drills to recover the isotopic

composition of oxygen inputs from water and food preserved in the hydroxyapatite (e.g.,

Janssen et al., 2016; Roberts et al., 2020; Kubat et al., 2023). This coarse drilling method

yields spatially and temporally blurred powdered samples formed over a substantial and

unknown period of time, however, precluding the identification of precise seasonal

environmental patterns. To circumvent this limitation, we have employed the stable isotope

sensitive high-resolution ion microprobe (SHRIMP SI) to measure $\delta^{18}O$ values sequentially

from thin sections of teeth, relating these to daily increments and birth lines to determine

enamel formation times, and in some instances, calendar ages (Smith et al., 2018a, 2022;

Green et al., 2022; Vaiglova et al., in review).

It is well established that $\delta^{18}O$ values in tooth enamel are closely related to local water

oxygen isotope compositions (reviewed in Green et al., 2018, 2022).  For teeth that form after

birth and during periods of milk consumption, $\delta^{18}O$ values are expected to be higher, as a

result of infant evaporative water loss while consuming $^{18}O$-enriched mother's milk (Bryant

et al., 1996; Wright and Schwartz 1999; Britton et al., 2015). Studies of large-bodied

mammals report that milk $\delta^{18}O$ values are elevated by ~ 1–6 ‰ relative to local drinking

water $\delta^{18}O$ (Kornexl et al., 1997; Lin et al., 2003; Chesson et al., 2010; Green et al., 2018; but

see Cherney et al., 2010). Comparable data on human or nonhuman primate milk enrichment

appear to be lacking, save for a study of 44 British infants aged 5–16 weeks (Roberts et al.,

1988). The urine of infants who were breast-fed showed isotopic enrichment of 1–3 ‰

compared to infants who were fed formula prepared from sterile local tap water.

While such studies point to potential changes in infant body water during nursing, it is

unclear whether such differences prohibit the use of early-formed enamel in studies of

climate variation (Blumenthal et al., 2017; Luyt and Sealey 2018). Two studies of $\delta^{18}O$

values in the dentitions of modern sheep, horses, and zebras reported higher bulk values (~ 1–

2 ‰) in five molars (M1) compared to the rest of the permanent dentition (Bryant et al.,

1996; Fricke and O'Neil, 1996). This led Fricke and O'Neil (1996) to suggest that M1s are

unlikely to reflect the values of local meteoric water due to the influence of maternal inputs in

utero and through lactation. However, near-weekly $\delta^{18}O$ values over the first 2.75 years of

life in a Neanderthal M1 measured with SHRIMP SI showed clear annual trends and

maximum $\delta^{18}O$ values corresponding to a period after nursing has ceased (Smith et al.,

2018a). An examination of longer continuous periods of enamel formation within and

between teeth will help to clarify whether early-formed primate teeth should be avoided for

studies of climate seasonality.

Here we first assess whether wild orangutans show elevated $\delta^{18}O$ values in early-

formed enamel, testing the suggestion that M1s are significantly affected by nursing $^{18}O$-

enrichment, thereby precluding their use in climatological reconstructions. We then explore

approximately 30 years of weekly $\delta^{18}O$ values (n = 2016 measurements) to compare

orangutan individuals from the islands of Sumatra and Borneo. Finally, we contrast $\delta^{18}O$

values between modern and Pleistocene orangutans, including those from key regions of

early human occupation: Lida Ajer, Sumatra (Hooijer, 1948; Westaway et al., 2017) and Niah

Caves, Malaysia (Hooijer, 1961; Barker et al., 2007) (Figure 1, Table 1). Novel

understanding of climate patterns in these fossil assemblages may inform debates about the

likelihood of modern humans living in dense Asian rainforests, and the conditions that would

support savannah corridors for human dispersals throughout the region (e.g., de Vos, 1983;

Bird et al., 2005; Westaway et al., 2017; Louys and Roberts, 2020).

**Results**

*Modern orangutans*

The $\delta^{18}O$ ranges of twelve modern and six fossil orangutan molars, representing 2971

near-weekly measurements spanning 57.6 years of tooth formation, are listed in Table 2.

Prior to making comparisons between individuals, geographic regions, or time periods, we

first consider the potential intra-individual effect of isotopic enrichment from maternal milk

on $\delta^{18}O$ values. Comparisons of $\delta^{18}O$ values during the first, second, and third years of life in

five modern orangutan first molars (M1) do not show consistently elevated values during

their first year (Figure 2). Mean yearly $\delta^{18}O$ values in the first year are elevated by only 0.3

‰ compared to the second year. While three of the five M1s showed first year $\delta^{18}O$ values

higher than second year values (p ≤ 0.05), only two individuals showed mean values that

were ~1 – 2 ‰ higher during year one; one individual showed no difference from the first to

the second year, and one individual showed lower values during the first year than during the

second year (p ≤ 0.05) (Supplemental Table 1). A sixth individual was only sampled from

193 days of age, but maximum values from this point onwards were similar across more than

three years of life. Similarly variable patterns were observed for the six putative fossil

orangutan M1s (Supplemental Figure 1).

Comparisons across serial molars in four modern orangutans show no consistent trend

of elevated $\delta^{18}O$ values in M1s relative to successive molars (Figure 3). Only two individuals

showed maximum $\delta^{18}O$ values in their M1s relative to M2s; in both instances M3s were

unavailable due to their lack of development prior to death. The other two individuals showed

higher $\delta^{18}O$ values in M2s or M3s than in their respective M1s. In the case of the oldest

individual (ZSM 1981/248), the highest $\delta^{18}O$ values appeared at approximately 5.8 years of

age, well past the age when exclusive nursing ends.

Comparison of the $\delta^{18}$O values in the full datasets of modern Bornean and Sumatran

orangutans reveals a high degree of overlap. Values from the three Bornean individuals

ranged from 12.7 to 20.0 ‰ (n = 955 near weekly measurements), while the three Sumatran

individuals ranged from 11.3 to 20.6 ‰ (n = 1061 measurements). Comparisons of periodic

trends via spectral power distribution analysis revealed more consistent bimodal patterns in

the Sumatran individuals; three of the six Bornean molars were aperiodic (statistical power of

0.1 or less), while all six of the Sumatran molars revealed annual or semiannual cycles with

greater power (Supplemental Figure 2). Rapid oxygen isotopic shifts on the order of ~ 6–8 ‰

are evident in the single Bornean and Sumatran individuals with $\delta^{18}$O measurements

spanning M1 to M3, which may represent one or more supra-annual ENSO events captured

during the ~ 9–11 years these molars were forming.

*Fossil orangutans – oxygen isotopes*

Concurrently forming teeth (molar specimens 11594.12 and 11595.105) from same

individual at Lida Ajer, Sumatra are nearly isotopically identical; $\delta^{18}$O values range from

15.1 to 19.9 ‰ and 15.7 to 20.0 ‰, respectively, supporting the biogenic fidelity of these

records. The $\delta^{18}$O values of two individuals from the nearby Sibrambang site (15.3–20.4 ‰,

14.7–20.8 ‰) are very similar to those of the Lida Ajer individual. These Sumatran fossils all

fall at the upper end of the range of modern Sumatran orangutans (Figure 4), and reveal

approximately annual $\delta^{18}$O periodicities (0.9–1.3 years), as well as strong bimodal

distribution patterns in one instance (11565.162).

The two fossils from the Niah Caves were excavated from different regions and

stratigraphic depths; $\delta^{18}$O in the tooth from grid US/22 ranged from 15.9 to 24.8 ‰ and,

unlike the three modern Bornean individuals, yielded an annual periodicity (1.0 years). The

$\delta^{18}$O in the tooth from grid Y/F4 ranged from 14.2 to 22.9 ‰ and showed a stronger bimodal

trend than an annual one, although its short formation time may have prohibited identification

of longer trends. The range of values from these two fossil molars (14.2–24.8 ‰) markedly

exceeds the range of modern Bornean orangutans (12.7–20.0 ‰) (Figure 4), with the mean

$\delta^{18}$O value at least 2‰ heavier, suggesting possibly drier conditions with greater seasonality

during their formation.

*Fossil orangutans – U-series age estimates*

The six fossil teeth have very low uranium concentrations in their enamel (<0.5 ppm),

regardless of their origin (Supplementary Table 2). These enamel values are very close to the

detection limit of the Nu Plasma II MC-ICP-MS, and thus are not useful for estimating

minimum ages. The dentine of Lida Ajer specimen 11595.105 shows a spatial gradient of

increasing uranium concentration from ~ 41 to 66 ppm, and decreasing age estimates from ~

51 to 40 ka (Supplementary Table 2). This trend might result from a preferential uranium

leaching overprint near the end of the root. Spot DE10, positioned near the EDJ, is less likely

to be impacted (Supplemental Figure 3), and is thus assumed to provide the most reliable

minimum age for the tooth, ~ 40 ka. Uranium values from Lida Ajer specimen 11594.12

show a similar trend of concentrations decreasing from ~ 31 to 24 ppm towards the root tip.

However, the U-series age estimates remain constant within the range 31–34 ka across the

dentine (Supplemental Table 2; Supplemental Figure 3). No evidence for a recent overprint is

observed, supporting a minimum age of 33 ka. In summary, this individual's minimum age is

~ 33–40 ka.

U-series analysis of the dentine of Sibrambang specimen 11565.162 shows a slight

decreasing trend of uranium concentration from the EDJ to the root tip (from > 60 ppm to <

60 ppm), and corresponding increasing age estimates (56–62 ka) (Supplemental Table 2;

Supplemental Figure 4). This might result from a slight uranium leaching overprint; a

minimum age of 60 ka is likely for this tooth. The U-series age estimates obtained for

Sibrambang specimen 11564.5 show a decreasing trend from the EDJ toward the

circumpulpal dentine from 75 to 65 ka (Supplemental Table 2; Supplemental Figure 4.)

However, given the associated uncertainties, this trend might not be meaningful. An average

dentine U-series age of 70.3 ± 5.5 ka (2σ) may be regarded as a minimum age for the fossil,

which is broadly consistent with the single age estimate obtained from the enamel (64 ka). In

summary, the two teeth from Sibrambang yield U-series apparent ages of ~ 60–70 ka.

The uranium concentration measured across the dentine of the Niah Caves specimen

from grid Y/F4 shows little variability, 4.2–4.9 ppm. The U-series age estimates are between

6.0 and 8.7 ka (Supplemental Table 2; Supplemental Figure 5). The average dentine U-series

minimum age is 7.6 ± 1.3 ka. Similarly, the Niah Cave specimen from grid US/22 shows a

consistent uranium concentration through the dentine (1.3–1.4 ppm), with relatively large

uncertainties that nonetheless bracket individual age estimates (Supplemental Table 2;

Supplemental Figure 5). The average dentine U-series minimum age is 8.8 ± 3.0 ka. In

summary, the two teeth from Niah Cave yield consistent apparent ages of ~ 8–9 ka.

**Discussion**

*Primate oxygen isotope compositions do not reveal a clear milk enrichment effect*

Half of our modern sample, and potentially all of our fossil sample, are composed of

M1s. These begin forming around birth and continue growing for three or more years (Smith,

2016). Orangutan infants rely exclusively on maternal milk during their first year of life,

supplementing this with solid foods in the second year, which are increased until suckling

ceases prior to nine years of age (van Noordwijk et al., 2013; Smith et al., 2017). Our

developmentally-guided sampling approach allows us to examine fine-scaled trends in $\delta^{18}O$

values during birth, exclusive nursing, supplemental feeding, and also after nursing ends (in

those individuals with available serial molar teeth).

We find that the five modern orangutans show only minor and inconsistently elevated

$\delta^{18}O$ values during the first year of life when compared to the subsequent year. These data do

not support the hypothesis that infant orangutans had markedly elevated body water $\delta^{18}O$

values during exclusive nursing. Data from the majority of 12 human M1s studied by

Vaiglova et al. (in review) similarly revealed maximum $\delta^{18}O$ values after the first year of

tooth formation, well beyond the duration of exclusive milk intake. This was also evident in

the M1 of a Neanderthal born in the spring (Smith et al., 2018a); $\delta^{18}O$ values mostly rose for

the first 3.5 months of life, but did not reach a maximum for another two years — long after

the infant would have begun consuming supplemental foods and liquids. This final dataset

points to the influence of season of birth on initial postnatal $\delta^{18}O$ values, as inferred in other

mammals (Bryant et al., 1996; Frick and O'Neil, 1996).

Comparisons of serially-forming teeth in four wild orangutans also failed to show a

consistent elevation of $\delta^{18}O$ values in M1s versus M2s (or M3s in two cases). Comparisons of

M1 $\delta^{18}O$ values with subsequent-forming teeth in four baboons, two tantalus monkeys, and

one mona monkey (from Green et al., 2022: SI Dataset S1) also largely fail to support the

enriched "Pattern 1" trend modelled by Bryant et al. (1996: Figure 4, p. 401). This is also the

case in comparisons of $\delta^{18}O$ values from bulk samples of human teeth—Wright and Schwartz

(1999) demonstrated that M1s have higher $\delta^{18}O$ values than later-forming teeth in only four

of seven individuals. In summary, the data from a range of primates, humans included, do not

support the exclusion of early-forming primate teeth from the assessment of environmental

seasonality.

*Modern orangutans show similar isotopic values across the islands of Borneo and Sumatra*

The two Bornean juveniles from the Munich collection (ZSM 1981/48, ZSM 1981/87)

reflect the environmental conditions of the late 1880s and early 1890s in Skalau—a region

where orangutans might now be locally extinct. Similarly, the teeth from the two Sumatran

individuals from the Munich collection (ZSM 1981/246, ZSM 1981/248) were collected prior

to 1939 in northern Aceh, from where orangutans also have since disappeared (Spehar et al.,

2018). While the individuals from northernmost Sumatra might have inhabited somewhat

higher elevations than those from western Borneo, there does not appear to be an evident

altitude effect (lower isotopic values at higher altitudes), as these four individuals show

similar isotopic values, save for a single brief excursion below 12 ‰ in ZSM 1981/248

(Table 1, Figure 3). It is unknown to what extent local rainfall may have been isotopically

distinct at the time the teeth were forming.

The $\delta^{18}$O values shown in Figure 1 reflect estimates of monthly and annual average

precipitation from the Online Isotopes in Precipitation Calculator (3.0) compiled for

www.waterisotopes.org. Actual measurements of precipitation $\delta^{18}$O from the islands of

Borneo and Sumatra are extremely limited. The closest observation facilities to the ZMS

orangutan locations yield similar patterns of modern annual rainfall $\delta^{18}$O variability

(Belgaman et al., 2017), yet specific measurements from the six facilities that make up

"Cluster 3" in this reference are not available for comparison.

Other studies underscore the complexity of water transport in this region—multiple

factors such as the oceanic origin of water vapor, cloud cover and type, and the post-

condensation process influence the short-term variability of $\delta^{18}$O values in rainfall (Moerman

et al., 2013; Suwarman et al., 2013; Belgaman et al., 2016). For example, Moerman et al.

(2013) provided five years of daily rainfall $\delta^{18}$O measurements from Northern Borneo

(Gunung Mulu National Park, Malaysia); daily rainfall $\delta^{18}$O values ranged from + 0.7 ‰ to

−18.5 ‰ and showed 1 – 3 month, annual, and supra-annual cycle frequencies. Interannual

rainfall $\delta^{18}$O fluctuations of 6–8 ‰ were significantly correlated with ENSO events; these are

similar in scale to the large fluctuations in our serial tooth datasets (Figure 3).

Another potential source of isotopic variability derives from dietary variation, as

orangutans obtain the majority of their body water from plants. Oxygen isotope values can be

stratified within tropical forest canopies (Sternberg et al., 1989; Roberts et al., 2017)—

potentially leading to offset values amongst various animals, including primates, that

consume different resources in the same forest (Nelson, 2013). While differences in $\delta^{18}$O

values are apparent in comparisons of diverse arboreal and terrestrial mammals (reviewed in

Green et al., 2022), it remains to be seen whether primates with broadly similar diets and

habitats show meaningful differences in tooth $\delta^{18}$O values, and to what extent plant-derived

oxygen influences the pattern and amplitude of seasonality relative to rainfall.

Oxygen isotope compositions in the six modern individuals from the islands of
Borneo and Sumatra are very similar. Orangutans from both islands prefer ripe fruit when
available, with some differences in the consumption of bark, leaves, unripe fruits, and
insects—which varies between sites and across seasons (reviewed in Smith et al., 2012).
Orangutan $\delta^{18}O$ values are also quite similar to the $\delta^{18}O$ values from five humans from
Flores, Indonesia (14.8–21.0 ‰) dated at ~ 2.2–3.0 ka (Vaiglova et al., in review). This is
remarkable given the major dietary differences between frugivorous orangutans and coastal-
dwelling humans, and suggests that their enamel $\delta^{18}O$ values may be influenced
predominantly by regional precipitation.
322
*Fossil orangutan isotope values suggest different ancient climates in Sumatra and Borneo*
Dating studies at Lida Ajer have established the presence of the oldest human remains
in insular Southeast Asia, ~ 63–73 ka (Westaway et al., 2017), and a broad survey of the cave
has reconfirmed an age of MIS 4 (59–76 ka) for the mammalian fauna (Louys et al., 2022).
This is consistent with the minimum age of ~ 33–40 ka estimated for the two molars
examined in the current study. The Sumatran Sibrambang Cave is regarded as roughly
contemporaneous to Lida Ajer given broad faunal similarities (de Vos, 1983). Unfortunately,
it has yet to be relocated since Dubois' excavations over a century ago (Bacon et al., 2015).
Although no independent age controls are available, the Sibrambang fossil assemblage
appears similar to, or slightly older than, that from Lida Ajer, given the minimum U-series
age estimates for teeth from both sites. Our analysis of $\delta^{18}O$ values in Sumatran orangutan
fossil molars reveals a close similarity across sites and with modern Sumatran individuals,
although the fossil compositions fall at the upper end of the modern range. This may indicate
a slightly dryer and less variable climate during the late Pleistocene.
Pollen records from the Niah Caves archaeological site indicate that there were a
number of local ecological shifts from lowland rainforest to more open environments during
the Late Pleistocene and into the Holocene (Hunt et al., 2012), where humans may have
begun hunting orangutans at ~ 45 ka (Spehar et al., 2018). While it is not possible to locate
the two fossil orangutan molars in these pollen records, Piper and Rabett (2016) considered
that the large animal bone assemblages accumulated within the Lobang Hangus entrance and
defined by the Harrisson spit depths of 12"- 42" were of terminal Pleistocene age. More
broadly, the orangutan specimen from grid US/22 (32"-36") is stratigraphically positioned
between radiocarbon ages of 14,206−15,061 cal. BP (OxA-13936) and 36,583−38,059 cal.
BP (OxA-13938), and this provides plausible minimum and maximum age constraints that

are not incompatible with the apparent minimum U-series age of ~ 9 ka. Based on these

results, the tooth is likely to date from the latest part of the Late Pleistocene. The specimen

from grid Y/F4 might date from the latest part of the Late Pleistocene to the early Holocene,

by comparison with the shell and fauna assemblage from other excavated areas (Piper et al.,

2016.)

Both orangutan molars from the Niah Caves yield wide ranges of $\delta^{18}O$, which is

particularly notable given the short periods of time sampled compared to the other fossils and

most modern orangutan molars. Given the similar offsets in $\delta^{18}O$ values between modern and

prehistoric Bornean orangutans, and baboons living in Ugandan forests and the Ethiopian rift

region (Green et al., 2022), we regard the higher $\delta^{18}O$ values in the Niah Cave orangutans as

possibly indicative of reduced rainfall when compared to recent conditions. This is consistent

with paleoclimate reconstructions for Borneo and Flores during the late Pleistocene and early

Holocene (Griffith et al. 2009; Buckingham et al., 2022), when the environment around the

Niah Caves is believed to have been a drier, more open seasonal forest (Harrison, 1996; Hunt

et al., 2012). A study of $\delta^{18}O$ values in Niah Caves shell middens dating from the early to

mid- Holocene indicates a shift to periods of high rainfall with less variation than modern

conditions (Stephens et al., 2016). The transition from a drier environment to moist tropical

rainforest is also reflected in the increasing number and higher frequencies of canopy-adapted

mammalian taxa in excavated layers of the Pleistocene-Holocene transition (Piper and Lim,

2021).

Our approach has the potential to contribute to reconstructions of ancient

paleoenvironments in SE Asia based on studies of pollen, molluscs, faunal community

compositions, guano records, and stable isotopes of teeth (e.g., Jablonski et al., 2000; Bird et

al., 2005; Louys and Meijaard, 2010; Wurster et al., 2010; Hunt et al., 2012; Janssen et al.,

2016; Stephens et al., 2016; Louys and Roberts, 2020; Bacon et al., 2021; Louys et al., 2022).

This may be especially timely given that recent work examining modern fauna compositions

in African landscapes has cautioned that fossil herbivore assemblages tend to overestimate

the extent of ancient grasslands in comparison to woodlands (Negash and Barr, 2023). Fine-

scaled tooth sampling may also allow an expansion of inferences from $\delta^{18}O$ values of bulk-

sampled Asian hominin remains (Janssen et al., 2016; Roberts et al., 2020; Kubat et al.,

2023), which are difficult to interpret for understanding seasonal rainfall dynamics in tropic

environments (Green et al., 2022). Such information could better inform debates about

whether humans employed arid savannah corridors to avoid dense tropical forests, or whether

humans were adept at colonizing such environments during their consequential migration

throughout island Southeast Asia.

**Materials and Methods**

*Orangutan samples*

Thin (histological) sections of twelve molar teeth from six modern orangutans and six

molar teeth from five fossil orangutans were employed (Table 1). These sections were

previously prepared for studies of incremental tooth development, enamel thickness,

elemental chemistry, and Asian hominoid taxonomy (Smith, 2016; Smith et al., 2011, 2012,

2017, 2018b). Four modern individuals were sourced from the Munich State Anthropological

Collection (ZSM); two were collected in 1893-1894 from Skalau (north of the Kapuas River

and south of the Klingkang Mountains in eastern West Borneo), and two were collected prior

to 1939 from Aceh (northwest Sumatra) (Röhrer-Ertl, 1988: Figure 3, p. 14) (Figure 1). It

was not possible to determine from which specific regions or time periods the two other

modern individuals derive—collection notes were not available for these specimens from the

Harvard Museum of Natural History (MCZ) or the Humboldt Museum (ZMB). Ages at death

were determined for five of six individuals from assessments of incremental features and

elemental registration of serially forming molars (detailed in Smith, 2016; Smith et al., 2017).

We also studied four Sumatran fossil orangutan teeth that were collected more than a

century ago from the Lida Ajer and Sibrambang Caves in the Padang Highlands by Eugene

Dubois (de Vos, 1983). Right and left lower molars from Lida Ajer (11594.12, 11595.105)

show identical trace element patterns in their dentine (Supplemental Figure 6), as well as

similar occlusal fissure patterns and light wear, consistent with their attribution to the same

individual. Two Bornean fossil orangutan teeth from Niah Caves (Malaysia) were also

included in this study. The caves have yielded significant late Pleistocene and early Holocene

human remains since the Harrissons began excavations in the 1950s (Barker et al., 2007).

These lower molars were derived from two different entrances to the cave system, Gan Kira

(grid square Y/F4) and Lobang Angus/Hangus (grid square US/22), with burial depths of 12–

18 inches and 30–36 inches, respectively (Hooijer, 1961). Although Hooijer (1948, 1961)

identified all six of these fossil teeth as M1s, we regard this as tentative, given that isolated

orangutan molars are notoriously difficult to seriate (Grine and Franzen, 1994).

*Dating of fossil samples*

413           Preliminary assessments at the Australian National University Radiocarbon Dating

Laboratory confirmed that collagen preservation in the six fossil orangutans was insufficient

for radiocarbon dating, a common limitation in tropical environments (e.g., Wood et al.,

2016). Laser ablation uranium series (U-series) analyses were carried out at the Radiogenic

Isotope Facility of the University of Queensland using an ASI RESOlution SE laser ablation

system connected to a Nu Plasma II MC-ICP-MS. A succession of several rasters (< 2-minute

linear ablations) were made in a transect across the dentine and enamel of each tooth

(Supplemental Figures 4-6) following Grün et al. (2014). The $^{230}Th/^{238}U$ and $^{234}U/^{238}U$

activity ratios of the samples were normalized to bracketing analyses of a homogeneous rhino

tooth standard that has been precisely calibrated by isotope dilution (Grün et al., 2014).

Importantly, dental tissues are known to behave as open systems for U-series elements;

provided there is no occurrence of uranium leaching, age estimates should therefore be

regarded as minimum age constraints since uranium uptake into dental tissues may be

significantly delayed after death.

*Tooth formation and oxygen isotope analyses*

429           Thin sections were first imaged with transmitted light microscopy. Enamel daily

secretion rates were measured between sequential accentuated growth lines to yield the time

of formation (see Smith, 2016: Fig. 1, p. 94), and enamel extension rates were calculated

between accentuated lines to guide placement of the analysed spots at approximately weekly

intervals of growth from the dentine horn tip to the enamel cervix (Smith et al., 2018a; Green

et al., 2022). Following the removal of cover slips by immersion in xylene, each thin section

was analysed for $\delta^{18}O$ at the SHRIMP Laboratory at the Australian National University

according to methods detailed in Vaiglova et al. (in review).

437           In brief, a 15 kV Cs primary ion beam focused to a spot ~ 15 x 20 µm diameter was

used to sequentially sample the enamel as close as possible to the enamel-dentine junction

(EDJ). Oxygen secondary ions were extracted at 10 kV and analysed isotopically by a

multiple collector equipped with dual electrometers operated in resistor mode. The $\delta^{18}O$

values were calculated relative to reference apatite (Durango 3) measured every 10–15

sample analyses. Distances of SHRIMP $\delta^{18}O$ measurements along the innermost enamel from

the cusp to cervix were converted to time in days following Green et al. (2022). A polynomial

regression relating distances to days was created using the enamel extension rates, and this

regression was applied to estimate the timing of secretory deposition at every SHRIMP spot

location. The Lomb–Scargle periodogram was used to assess time-dependent patterns of $\delta^{18}O$

values, which estimates the power of sine wave periods within a given range to produce the

temporal patterns present within those measurements.

The probability that differences between first and second year $\delta^{18}O$ values in modern

first molars might have arisen by chance was assessed by one-way paired t-tests, with alpha =

0.05 adjusted by Bonferroni correction due to repeated comparisons across multiple teeth.

Figure and data plotting using Python 3 in the Google Colab environment were aided by

ChatGPT, a language model based on the GPT-3.5 architecture developed by OpenAI.

**Acknowledgements**

Manfred Ade, Steffen Bock, Judy Chupasko, Mark Omura, Carina Phillips, and Olav Rohrer-

Ertl assisted with access to living orangutan material. Graeme Barker, Ipoi Datan, and

Natasja den Ouden assisted with access to fossil orangutan material. The following museums

and universities provided access to material: Humboldt Museum (Berlin), Harvard University

Museum of Comparative Zoology (Cambridge), Naturalis Museum (Leiden), State

Anthropological Collection (Munich), and the Sarawak Museum (Kuching). Technical

assistance was provided by Kamil Sokolowski and Brian Tse at the Preclinical Imaging Core

Facility at the Translational Research Institute, funding support for which came from

Therapeutic Innovation Australia, under the National Collaborative Research Infrastructure

Strategy. Kate Britton, Luca Fiorenza, Gathorne Gathorne-Hardy (Earl of Cranbrook), Terry

Harrison, Jessica Rippengal, Pam Walton, and Rachel Wood provided research assistance.

Sample preparation and oxygen isotope analysis was funded by the Australian Academy of

Science Regional Collaborations Programme, the Australian National University, the

Australian Research Council (DP210101913), and Griffith University. The U-series dating

was funded by the Australian Research Council (FT150100215) and the Spanish Ramón y

Cajal Fellowship (RYC2018-025221-I). The latter is funded by MCIN/AEI/

10.13039/501100011033 and by 'ESF Investing in your future.' U-series dating analyses

were carried out within the framework of the existing Brisbane Geochronology Alliance

(Griffith University, University of Queensland, and Queensland University of Technology);

Y. Feng helped with the U-series data acquisition.

**References**

Aldrian ED, Susanto R. 2003. Identification of three dominant rainfall regions within
Indonesia and their relationship to sea surface temperature. International Journal of
Climatology **23**: 1435-1452. DOI: https://doi.org/10.1002/joc.950
484

485    Archibald S, Bond WJ, Hoffmann W, Lehmann C, Staver C, Stevens N. 2019. Distribution
486    and determinants of Savannas. *In: Savanna Woody Plants and Large Herbivores*, (eds P.F.
487    Scogings and M. Sankaran). DOI: https://doi.org/10.1002/9781119081111.ch1
488

489    Ashbury AM, Posa MRC, Dunkel LP, Spillmann B, Atmoko SSU, van Schaik CP, van
490    Noordwijk MA. 2015. Why do orangutans leave the trees? Terrestrial behavior among wild
491    Bornean orangutans (*Pongo pygmaeus wurmbii*) at Tuanan, Central Kalimantan. *American*
492    *Journal of Primatology* **77**: 1216-1229. DOI: https://doi.org/10.1002/ajp.22460
493

494    Aubert M, Williams IS, Boljkovac K, Moffat I, Moncel MH, Dufour E, Grün R. 2012. In situ
495    oxygen isotope micro-analysis of faunal material and human teeth using a SHRIMP II: A new
496    tool for palaeo-ecology and archaeology. *Journal of Archaeological Science* **39**: 3184-3194.
497    DOI: https://doi.org/10.1016/j.jas.2012.05.002
498

Bacon A-M, Bourgon N, Welker F, Cappellini E, Fiorillo D, Tombret O, Huong NTM, Tuan
NA, Sayavonkhamdy T, Souksavatdy V, Sichanthongtip P, Antoine P-O, Duringer P, Ponche
J-L, Westaway K, Joannes-Boyau R, Boesch Q, Suzzoni E, Frangeul S, Patole-Edoumba E,
Zachwieja A, Shackelford L, Demeter F, Hublin J-J, Dufour É. 2021. A multi-proxy
approach to exploring Homo sapiens' arrival, environments and adaptations in Southeast
Asia. *Scientific Reports* **11**: 21080. DOI: https://doi.org/10.1038/s41598-021-99931-4
505

Bacon A-M, Westaway K, Antoine P-O, Duringer P, Blin A, Demeter F, Ponche J-L, Zhao J-
X, Barnes LM, Sayavonkhamdy T, Thuy NTK, Long VT, Patole-Edoumba E, Shackelford L.
2015. Late Pleistocene mammalian assemblages of Southeast Asia: New dating, mortality
profiles and evolution of the predator-prey relationships in an environmental context.
*Palaeogeography, Palaeoclimatology, Palaeoecology* **422**: 101-127. DOI:
https://doi.org/10.1016/j.palaeo.2015.01.011
512

513    Bae CJ, Douka K, Petraglia MD. 2017. On the origin of modern humans: Asian perspectives.
514    *Science* **358**: eaai9067. DOI: https://doi.org/10.1126/science.aai9067
515

Barker G, Barton H, Bird M, Daly P, Datan I, Dykes A, Farr L, Gilbertson D, Harrisson B,
Hunt C, Higham T, Kealhofer L, Krigbaum J, Lewis H, McLaren S, Paz V, Pike A, Piper P,
Pyatt B, Rabett R, Reynolds T, Rose J, Rushworth G, Stephens M, Stringer C, Thompson J,
Turney C. 2007. The 'human revolution' in lowland tropical Southeast Asia: the antiquity and
behavior of anatomically modern humans at Niah Cave (Sarawak, Borneo). *Journal of*
*Human Evolution* **52**: 243-261. DOI: https://doi.org/10.1016/j.jhevol.2006.08.011
522

523    Belgaman HA, Ichiyanagi K, Tanoue M, Suwarman R. 2016. Observational research on
524    stable isotopes in precipitation over Indonesian maritime continent. *Journal of Japanese*
525    *Association of Hydrological Sciences* **46**: 7–28. DOI: https://10.4145/jahs.46.7
526

527    Belgaman HA, Ichiyanagi K, Suwarman R, Tanoue M, Aldrian E, Utami AID,
528    Kusumaningtyas SDA. 2017. Characteristics of seasonal precipitation isotope variability in
529    Indonesia. *Hydrological Research Letters* **11**: 92-98. DOI: https://doi.org/10.3178/hrl.11.92
530

Bird MI, Taylor D, Hunt C. 2005. Palaeoenvironments of insular Southeast Asia during the
Last Glacial Period: A savanna corridor in Sundaland? *Quaternary Science Reviews* **24**:
2228-2242. DOI: https://doi.org/10.1016/j.quascirev.2005.04.004

Blumenthal SA, Levin NE, Brown FH, Brugal JP, Chritz KL, Harris JM, Jehle GE, Cerling
TE. 2017. Aridity and hominin environments. *Proceedings of the National Academy of
Sciences of the United States of America* **114**: 7331-7336. DOI:
https://doi.org/10.1073/pnas.1700597114

Bowen GJ. 2010. Isoscapes: Spatial pattern in isotopic biogeochemistry. *Annual Review of
Earth and Planetary Sciences* **38**: 161-187. DOI: https://doi.org/10.1146/annurev-earth-
040809-152429

Britton K, Fuller BT, Tütken T, Mays S, Richards MP. 2015. Oxygen isotope analysis of
human bone phosphate evidences weaning age in archaeological populations. *American
Journal of Physical Anthropology* **157**: 226-241. DOI: https://doi.org/10.1002/ajpa.22704

Brockman D, vain Schaik C. (Eds.). 2005. *Seasonality in Primates: Studies of Living and
Extinct Human and Non-Human Primates* (Cambridge Studies in Biological and
Evolutionary Anthropology). Cambridge: Cambridge University Press. DOI:
https://doi.org/10.1017/CBO9780511542343

Bryant JD, Froelich PN, Showers WJ, Genna BJ. 1996. A tale of two quarries: Biologic and
taphonomic signatures in the oxygen isotope composition of tooth enamel phosphate from
modern and miocene equids. *Palaios* **11**: 397-408. DOI: https://doi.org/10.2307/3515249

Buckingham FL, Carolin SA, Partin JW, Adkins JF, Cobb KM, Day CC, Ding Q, He C, Liu
Z, Otto-Bliesner B, Roberts WHG, Lejau S, Malang J. 2022. Termination 1 millennial-scale
rainfall events over the Sunda Shelf. *Geophysical Research Letters* **49**: e2021GL096937.
DOI: https://doi.org/10.1029/2021GL096937

Cherney MD, Fisher DC, Hren MT, Shirley EA. 2021. Stable isotope records of nursing and
weaning: A case study in elephants with implications for paleobiological investigations.
*Palaeogeography, Palaeoclimatology, Palaeoecology* **567**: 110223. DOI:
https://doi.org/10.1016/j.palaeo.2021.110223

Chesson LA, Valenzuela LO, O'Grady SP, Cerling TE, Ehleringer JR. 2010. Hydrogen and
oxygen stable isotope ratios of milk in the United States. *Journal of Agricultural and Food
Chemistry* **58**: 2358-2363. DOI: https://doi.org/10.1021/jf904151c

Cole JE, Fairbanks RG, Shen GT. 1993. Recent variability in the southern oscillation:
Isotopic results from a Tarawa Atoll coral. *Science* **260**: 1790-1793. DOI:
https://doi.org/10.1126/science.260.5115.1790

Curran LM, Caniago I, Paoli GD, Astianti D, Kusneti M, Leighton M, Nirarita CE,
Haeruman H. 1999. Impact of El Niño and logging on canopy tree recruitment in Borneo.
*Science* **286**: 2184-2188. DOI: https://doi.org/10.1126/science.286.5447.2184

Dansgaard W. 1964. Stable isotopes in precipitation. *Tellus* **16**: 436-468. DOI:
https://doi.org/10.1111/j.2153-3490.1964.tb00181.x

de Vos J. 1983. The *Pongo* faunas from Java and Sumatra and their significance for
biostratigraphical and paleo-ecological interpretations. *Palaeontology Proceedings B*. **86**:
417-425.
584
585  da Silveira L, Sternberg L, Mulkey SS, Joseph Wright S. 1989. Oxygen isotope ratio
586  stratification in a tropical moist forest. *Oecologia* **81**: 51-56. DOI:
587  https://doi.org/10.1007/BF00377009
588
589  Fricke HC, O'Neil JR. 1996. Inter- and intra-tooth variation in the oxygen isotope
590  composition of mammalian tooth enamel phosphate: Implications for palaeoclimatological
591  and palaeobiological research. *Palaeogeography, Palaeoclimatology, Palaeoecology* **126**: 91-
592  99. https://doi.org/10.1016/S0031-0182(96)00072-7
593
594  Green DR, Ávila JN, Cote S, Dirks W, Lee D, Poulsen CJ, Williams IS, Smith TM. 2022.
595  Fine-scaled climate variation in equatorial Africa revealed by modern and fossil primate
596  teeth. *Proceedings of the National Academy of Sciences of the United States of America* **119**:
597  e212336611. DOI: https://doi.org/10.1073/pnas.2123366119
598
599  Green DR, Olack G, Colman AS. 2018. Determinants of blood water δ18O variation in a
600  population of experimental sheep: Implications for paleoclimate reconstruction. *Chemical
601  Geology* **485**: 32-43. DOI: https://doi.org/10.1016/j.chemgeo.2018.03.034
602
603  Grine F, Franzen JL. 1994. Fossil hominid teeth from the Sangiran Dome (Java, Indonesia).
604  *Courier Forschungsinstitut Senckenberg* **171**: 75–103.
605
606  Griffiths ML, Drysdale RN, Gagan MK, Zhao J-X, Ayliffe LK, Hellstrom JC, Hantoro WS,
607  Frisia S, Feng Y-X, Cartwright I, Pierre ES, Fischer MJ, Suwargadi BW. 2009. Increasing
608  Australian-Indonesian monsoon rainfall linked to early Holocene sea-level rise. *Nature
609  Geoscience* **2**: 636-639. DOI: https://doi.org/10.1038/ngeo605
610
611  Grün R, Eggins S, Kinsley L, Moseley H, Sambridge M. 2014. Laser ablation U-series
612  analysis of fossil bones and teeth. *Palaeogeography, Palaeoclimatology, Palaeoecology* **416**:
613  150-167. DOI: https://doi.org/10.1016/j.palaeo.2014.07.023
614
615  Harrison T. 1996. The paleoecological context at Niah Cave Sarawak: Evidence from the
616  primate fauna. *Bulletin of the Indo-Pacific Prehistory Association* **14**, 90–100.
617
618  Hooijer DA. 1948. Prehistoric teeth of man and of the orang-utan from central Sumatra, with
619  notes on the fossil orang-utan from Java and Southern China. *Zoologische Mededelingen* **29**:
620  175–301.
621
622  Hooijer DA. 1961. The orang-utan in Niah Cave prehistory. *Sarawak Museum Journal* **9**:
623  408-421.
624
625  Hughen KA, Schrag DP, Jacobsen SB, Hantoro W. 1999. El Niño during the last interglacial
626  period recorded by a fossil coral from Indonesia. *Geophysical Research Letters* **26**: 3129-
627  3132. DOI: https://doi.org/10.1029/1999GL006062
628

629     Hunt CO, Gilbertson DD, Rushworth G. 2012. A 50,000-year record of late Pleistocene
630     tropical vegetation and human impact in lowland Borneo. *Quaternary Science Reviews* **37**:
631     61-80. DOI: https://doi.org/10.1016/j.quascirev.2012.01.014
632
633     Jablonski NG, Whitfort MJ, Roberts-Smith N, Qinqi X. 2000. The influence of life history
634     and diet on the distribution of catarrhine primates during the Pleistocene in eastern Asia.
635     *Journal of Human Evolution* **39**: 131-157. DOI: https://doi.org/10.1006/jhev.2000.0405
636
637     Janssen R, Joordens JCA, Koutamanis DS, Puspaningrum MR, de Vos J, van der Lubbe
638     JHJL, Reijmer JJG, Hampe O, Vonhof HB. 2016. Tooth enamel stable isotopes of Holocene
639     and Pleistocene fossil fauna reveal glacial and interglacial paleoenvironments of hominins in
640     Indonesia. *Quaternary Science Reviews* **144**: 145-154. DOI:
641     https://doi.org/10.1016/j.quascirev.2016.02.028
642
643     Knott CD. 1998. Changes in orangutan caloric intake, energy balance, and ketones in
644     response to fluctuating fruit availability. *International Journal of Primatology* **19**: 1061-1079.
645     DOI: https://doi.org/10.1023/A:1020330404983
646
647     Kornexl BE, Werner T, Roßmann A, Schmidt H-L. 1997. Measurement of stable isotope
648     abundances in milk and milk ingredients - A possible tool for origin assignment and quality
649     control. *Zeitschrift fur Lebensmittel -Untersuchung und -Forschung* **205**: 19-24. DOI:
650     https://doi.org/10.1007/s002170050117
651
652     Kubat J, Nava A, Bondioli L, Dean MC, Zanolli C, Bourgon N, Bacon A-M, Demeter F,
653     Peripoli B, Albert R, Lüdecke T, Hertler C, Mahoney P, Kullmer O, Schrenk F, Müller W.
654     2023. Dietary strategies of Pleistocene *Pongo sp.* and *Homo erectus* on Java (Indonesia).
655     *Nature Ecology and Evolution* **7**: 279-289. DOI: https://doi.org/10.1038/s41559-022-01947-0
656
657     Lin GP, Rau YH, Chen YF, Chou CC, Fu WG. 2006. Measurements of $\delta$D and $\delta^{18}$O stable
658     isotope ratios in milk. *Journal of Food Science* **68**: 2192-2195. DOI:
659     https://doi.org/10.1111/j.1365-2621.2003.tb05745.x
660
661     Louys J, Duval M, Price GJ, Westaway K, Zaim Y, Rizal Y, Aswan A, Puspaningrum M,
662     Trihascaryo A, Breitenbach SFM, Kwiecien O, Cai Y, Higgins P, Albers PCH, de Vos J,
663     Roberts P. 2022. Speleological and environmental history of Lida Ajer cave, western
664     Sumatra. *Philosophical Transactions of the Royal Society B: Biological Sciences* **377**:
665     20200494. DOI: https://doi.org/10.1098/rstb.2020.0494
666
667     Louys J, Meijaard E. 2010. Palaeoecology of Southeast Asian megafauna-bearing sites from
668     the Pleistocene and a review of environmental changes in the region. *Journal of*
669     *Biogeography* **37**: 1432-1449. DOI: https://doi.org/10.1111/j.1365-2699.2010.02297.x
670
671     Louys J, Roberts P. 2020. Environmental drivers of megafauna and hominin extinction in
672     Southeast Asia. *Nature* **586**: 402-406. DOI: https://doi.org/10.1038/s41586-020-2810-y
673
674     Luyt J, Sealy J. 2018. Inter-tooth comparison of $\delta^{13}$C and $\delta^{18}$O in ungulate tooth enamel
675     from south-western Africa. *Quaternary International* **495**:144-152. DOI:
676     https://doi.org/10.1016/j.quaint.2018.02.009
677

Marshall AJ, Ancrenaz M, Brearley FQ, Fredriksson GM, Ghaffar N, Heydon M, Husson SJ,
Leighton M, McConkey KR, Morrogh-Bernard HC, Proctor J, van Schaik CP, Yeager CP,
Wich SA. 2009. The effects of forest phenology and floristics on populations of Bornean and
Sumatran orangutans: Are Sumatran forests better orangutan habitat than Bornean
forests?', in Wich SA, Utami Atmoko SS, Setia TM, van Schaik CP (eds), *Orangutans:*
*Geographic Variation in Behavioral Ecology and Conservation* (Oxford, 2008; online
edn, Oxford Academic, 1 May 2009) 97-117. DOI:
https://doi.org/10.1093/acprof:oso/9780199213276.003.0007,
686
687    McPhaden MJ, Zebiak SE, Glantz MH. 2006. ENSO as an integrating concept in earth
688    science. *Science* **314**: 1740-1745. DOI: https://doi.org/10.1126/science.1132588
689
690    Moerman JW, Cobb KM, Adkins JF, Sodemann H, Clark B, Tuen AA. 2013. Diurnal to
691    interannual rainfall $\delta^{18}O$ variations in northern Borneo driven by regional hydrology. *Earth*
692    *and Planetary Science Letters* **369-370**: 108-119. DOI:
693    https://doi.org/10.1016/j.epsl.2013.03.014
694
695    Moron V, Robertson AW, Boer R. 2009. Spatial coherence and seasonal predictability of
696    monsoon onset over Indonesia. *Journal of Climate* **22**: 840-850. DOI:
697    https://doi.org/10.1175/2008JCLI2435.1
698
699    Negash EW, Barr WA. 2023. Relative abundance of grazing and browsing herbivores is not a
700    direct reflection of vegetation structure: Implications for hominin paleoenvironmental
701    reconstruction. *Journal of Human Evolution* **177**: 103328. DOI:
702    https://doi.org/10.1016/j.jhevol.2023.103328
703
704    Nelson SV. 2013. Chimpanzee fauna isotopes provide new interpretations of fossil ape and
705    hominin ecologies. *Proceedings of the Royal Society B* **280**: 20132324. DOI:
706    https://doi.org/10.1098/rspb.2013.2324
707
708    Pederzani S, Britton K. 2019. Oxygen isotopes in bioarchaeology: Principles and
709    applications, challenges and opportunities. *Earth-Science Reviews* **188**: 77-107. DOI:
710    https://doi.org/10.1016/j.earscirev.2018.11.005
711
712    Piper PJ. 2016. Human cultural, technological and adaptive changes from the end of the
713    Pleistocene to the mid-Holocene in Southeast Asia. in Oxenham, M. & Buckley, H. R. (eds.)
714    *The Routledge Handbook of Bioarchaeology in Southeast Asia and the Pacific Islands*,
715    Routledge, 24-44.
716
717    Piper P, Lewis H, Reynolds T, McLaren S, Rabett R, Cole F, Szábo K, Farr L, Barker G.
718    2016. The NCP Excavations in Lobang Hangus, Lobang Tulang, Gan Kira, and Kain Hitam,
719    in G. Barker, L. Farr (eds.), *Archaeological Investigations in the Niah Caves, Sarawak: The*
720    *Archaeology of the Niah Caves, Sarawak*, **2**: 65-80. Cambridge, United Kingdom: McDonald
721    Institute for Archaeological Research.
722
723    Piper PJ, Lim TT. 2021. Zooarchaeology at Niah Cave: contributions to our understanding of
724    southeast Asian prehistory. Malayan Nat J 81[st] Anniversary Special Issue, 195-208.
725
726    Piper P, Rabett R. 2016. Vertebrate Fauna from the Niah Caves, in G. Barker, L. Farr (eds.),
727    *Archaeological Investigations in the Niah Caves, Sarawak: The Archaeology of the Niah*

728    *Caves, Sarawak*, **2**: 401-438. Cambridge, United Kingdom: McDonald Institute for
729    Archaeological Research
730
731    Pletcher E, Staver C, Schwartz NB. 2022. The environmental drivers of tree cover and forest–
732    savanna mosaics in Southeast Asia. *Ecography* **2022**: e06280. DOI:
733    https://doi.org/10.1111/ecog.06280
734
735    Prendergast AL, Pryor AJE, Reade H, Stevens RE. 2018. Seasonal records of
736    palaeoenvironmental change and resource use from archaeological assemblages. *Journal of*
737    *Archaeological Science: Reports* **21**: 1191–1197. DOI:
738    https://doi.org/10.1016/j.jasrep.2018.08.006
739
740    Pumijumnong N, Bräuning A, Sano M, Nakatsuka T, Muangsong C, Buajan S. 2020. A 338-
741    year tree-ring oxygen isotope record from Thai teak captures the variations in the Asian
742    summer monsoon system. *Scientific Reports* **10**: 8966. DOI: https://doi.org/10.1038/s41598-
743    020-66001-0
744
745    Qian J-H, Robertson AW, Moron V. 2013. Diurnal cycle in different weather regimes and
746    rainfall variability over Borneo associated with ENSO. *Journal of Climate* **26**: 1772-1790.
747    DOI: https://doi.org/10.1175/JCLI-D-12-00178.1
748
749    Roberts SB, Coward WA, Ewing G, Savage J, Cole TJ, Lucas, A. 1988. Effect of weaning on
750    accuracy of doubly labled water method in infants. *American Journal of Physiology-*
751    *Regulatory, Integrative and Comparative Physiology* **254**: R622-R627. DOI:
752    https://doi.org/10.1152/ajpregu.1988.254.4.R622
753
754    Roberts P, Blumenthal SA, Dittus W, Wedage O, Lee-Thorp JA. 2017. Stable carbon,
755    oxygen, and nitrogen, isotope analysis of plants from a South Asian tropical forest:
756    Implications for primatology. *American Journal of Primatology* **79**: e22656. DOI:
757    https://doi.org/10.1002/ajp.22656
758
759    Roberts P, Louys J, Zech J, Shipton C, Kealy S, Carro SS, Hawkins S, Boulanger C, Marzo
760    S, Fiedler B, Boivin N, Mahirta, Aplin K, O'Connor S. 2020. Isotopic evidence for initial
761    coastal colonization and subsequent diversification in the human occupation of Wallacea.
762    *Nature Communications* **11**: 2068. DOI: https://doi.org/10.1038/s41467-020-15969-4
763
764    Röhrer-Ertl O. 1988 Research history, nomenclature, and taxonomy of the orang-utan, in
765    Schwartz JH (ed) *Orang-utan Biology*, Oxford University Press, New York, 7-18.
766
767    Rozanski K, Araguás-Araguás L, Gonfiantini R. 1993. Isotopic patterns in modern global
768    precipitation, in Swart PK, Lohmann KC, Mckenzie J, Savin S (eds), Climate change in
769    continental isotopic records. American Geophysical Union, Washington D.C., USA; 1–36.
770    DOI: https://doi.org/10.1029/GM078p0001
771
772    Smith TM. 2016. Dental development in living and fossil orangutans. *Journal of Human*
773    *Evolution* **94**: 92-105. DOI: https://doi.org/10.1016/j.jhevol.2016.02.008
774
Smith TM, Austin C, Ávila JN, Dirks W, Green DR, Williams IS, Arora M. 2022. Permanent
signatures of birth and nursing initiation are chemically recorded in teeth. *Journal of*
*Archaeological Science* **140**: 105564. DOI: https://doi.org/10.1016/j.jas.2022.105564

Smith TM, Austin C, Green DR, Joannes-Boyau R, Bailey S, Dumitriu D, Fallon S, Grün R,
James HF, Moncel MH, Williams IS, Wood R, Arora M. 2018a. Wintertime stress, nursing,
and lead exposure in Neanderthal children. *Science Advances* **4**: eaau9483. DOI:
https://doi.org/10.1126/sciadv.aau9483
783

784  Smith TM, Austin C, Hinde K, Vogel ER, Arora M. 2017. Cyclical nursing patterns in wild
785  orangutans. *Science Advances* **3**: e1601517. DOI: https://doi.org/10.1126/sciadv.1601517
786

787  Smith, TM., Bacon A-M, Demeter F, Kullmer O, Nguyen KT, de Vos J, Wei W, Zermeno JP,
788  Zhao L. 2011. Dental tissue proportions in fossil orangutans from mainland Asia and
789  Indonesia. *Human Origins Research* **1**: e1-e6.
790

791  Smith TM, Houssaye A, Kullmer O, Le Cabec A, Olejniczak AJ, Schrenk F, de Vos J,
792  Tafforeau P. 2018b. Disentangling isolated dental remains of asian pleistocene hominins and
793  pongines. *PLoS ONE* **13**: e0204737. DOI: https://doi.org/10.1371/journal.pone.0204737
794

795  Smith TM, Kupczik K, MacHanda Z, Skinner MM, Zermeno JP. 2012. Enamel thickness in
796  Bornean and Sumatran orangutan dentitions. *American Journal of Physical Anthropology*
797  **147**: 417-426. DOI: https://doi.org/10.1002/ajpa.22009
798

799  Spehar SN, Sheil D, Harrison T, Louys J, Ancrenaz M, Marshall AJ, Wich SA, Bruford MW,
800  Meijaard E. 2018. Orangutans venture out of the rainforest and into the anthropocene.
801  *Science Advances* **4**: e1701422. DOI: https://doi.org/10.1126/sciadv.1701422
802

803  Stahle DS, D'Arrigo RD, Krusic PJ, Cleaveland MK, Cook ER, Allan RJ, Cole JE, Dunbar
804  RB, Therrell MD, Gay DA, Moore MD, Stokes MA, Burns BT, Villanueva-Diaz J,
805  Thompson LG. 1998. Experimental Dendroclimatic Reconstruction of the Southern
806  Oscillation. *Bulletin of the American Meteorological Society* **79**: 2137-2152. DOI:
807  https://doi.org/10.1175/1520-0477(1998)079<2137:EDROTS>2.0.CO;2
808

809  Stephens M, Rose J, Mattey D, Gilbertson D. D. 2016. Stable isotope analysis of shells from
810  the West Mouth: palaeoenvironments, seasonality and harvesting, in G. Barker, L. Farr (eds.),
811  *Archaeological Investigations in the Niah Caves, Sarawak: The Archaeology of the Niah*
812  *Caves, Sarawak*, **2**: 177-200. Cambridge, United Kingdom: McDonald Institute for
813  Archaeological Research.
814

815  Suwarman R, Ichiyanagi K, Tanoue M, Yoshimura K, Mori S, Yamanaka MD, Kurita N,
816  Syamsudin F. 2013. The variability of stable isotopes and water origin of precipitation over
817  the maritime continent. *Scientific Online Letters on the Atmosphere* **9**: 74-78. DOI:
818  https://doi.org/10.2151/sola.2013-017
819

820  Thorpe SKS, Crompton RH. 2009. Orangutan positional behavior: Interspecific variation and
821  ecological correlates, in Wich SA, Utami Atmoko SS, Setia TM, van Schaik CP
822  (eds), *Orangutans: Geographic Variation in Behavioral Ecology and*
823  *Conservation* (Oxford, 2008; online edn, Oxford Academic, 1 May 2009) 33-48. DOI:
824  https://doi.org/10.1093/acprof:oso/9780199213276.003.0003
825

826    Tshen LT. 2016. Biogeographic distribution and metric dental variation of fossil and living
827    orangutans (*Pongo spp.*). *Primates* **57**: 39-50. DOI: https://doi.org/10.1007/s10329-015-
828    0493-z
829

830    Tudhope AW, Chilcott CP, McCulloch MT, Cook ER, Chappell J, Ellam RM, Lea DW,
831    Lough JM, Shimmield GB. 2001. Variability in the El Niño-Southern Oscillation through a
832    glacial-interglacial cycle. *Science* **291**: 1511-1517. DOI:
833    https://doi.org/10.1126/science.1057969
834

835    Urban FE, Cole JE, Overpeck JT. 2000. Influence of mean climate change on climate
836    variability from a 155-year tropical Pacific coral record. *Nature* **407**: 989-993. DOI:
837    https://doi.org/10.1038/35039597
838

839    Vaiglova P, Ávila JN, Buckley H, Galipaud JC, Green DR, Halcrow S, James HF, Kinaston
840    R, Oxenham M, Paz V, Simanjukntak T, Snoeck C, Trinh HH, Williams IS, Smith TM. (in
841    review) Holocene rainfall patterns in Southeast Asia revealed by microanalysis of $\delta^{18}O$
842    values in human teeth. *eLife*
843

844    van Noordwijk MA, Willems EP, Utami Atmoko SS, Kuzawa CW, van Schaik CP. 2013.
845    Multi-year lactation and its consequences in Bornean orangutans (*Pongo pygmaeus wurmbii*).
846    *Behavioral Ecology and Sociobiology* **67**: 805-814. DOI: https://doi.org/10.1007/s00265-013-
847    1504-y
848

849    Van Schaik CP. 1986. Phenological changes in a Sumatran rain forest. *Journal of Tropical*
850    *Ecology* **2**: 327-347. DOI: https://doi.org/10.1017/S0266467400000973
851

852    Vico G, Dralle D, Feng X, Thompson S, Manzoni S. 2017. How competitive is drought
853    deciduousness in tropical forests? A combined eco-hydrological and eco-evolutionary
854    approach. *Environmental Research Letters* **12**: 065006. DOI: https://doi.org/10.1088/1748-
855    9326/aa6f1b
856

857    Wessling EG, Kühl HS, Mundry R, Deschner T, Pruetz JD. 2018. The costs of living at the
858    edge: Seasonal stress in wild savanna-dwelling chimpanzees. *Journal of Human Evolution*
859    **121**: 1-11. DOI: https://doi.org/10.1016/j.jhevol.2018.03.001
860

861    Westaway KE, Louys J, Awe RD, Morwood MJ, Price GJ, Zhao J-X, Aubert M, Joannes-
862    Boyau R, Smith TM, Skinner MM, Compton T, Bailey RM, Van Den Bergh GD, de Vos J,
863    Pike AWG, Stringer C, Saptomo EW, Rizal Y, Zaim J, Santoso WD, Trihascaryo A, Kinsley
864    L, Sulistyanto B. 2017. An early modern human presence in Sumatra 73,000-63,000 years
865    ago. *Nature* **548**: 322-325. DOI: https://doi.org/10.1038/nature23452
866

867    Wood R, Duval M, Mai Huong NT, Tuan NA, Bacon AM, Demeter F, Duringer P, Oxenham
868    M, Piper P. 2016. The effect of grain size on carbonate contaminant removal from tooth
869    enamel: Towards an improved pretreatment for radiocarbon dating. *Quaternary*
870    *Geochronology* **36**: 174-187. DOI: https://doi.org/10.1016/j.quageo.2016.08.010
871

872    Wright LE, Schwarcz HP. 1999. Correspondence between stable carbon, oxygen and nitrogen
873    isotopes in human tooth enamel and dentine: Infant diets at Kaminaljuyu. *Journal of*
874    *Archaeological Science* **26**: 1159-1170. DOI: https://doi.org/10.1006/jasc.1998.0351
875

876    Wurster CM, Bird MI, Bull ID, Creed F, Bryant C, Dungait JAJ, Paz V. 2010. Forest
877    contraction in north equatorial Southeast Asia during the Last Glacial Period. *Proceedings of*
878    *the National Academy of Sciences of the United States of America* **107**: 15508-15511. DOI:
879    https://doi.org/10.1073/pnas.1005507107
880
881    Yang H, Johnson KR, Griffiths ML, Yoshimura K. 2016. Interannual controls on oxygen
882    isotope variability in Asian monsoon precipitation and implications for paleoclimate
883    reconstructions. *Journal of Geophysical Research* **121**: 8410-8428. DOI:
884    https://doi.org/10.1002/2015JD024683
885

886    **Figures**
887
888    Figure 1. Approximate location of select modern and fossil orangutans superimposed on
889    modeled isotopic variation.
890

891
892
893
894    Figure modified from www.waterisotopes.org based on data from the Online Isotopes in
895    Precipitation Calculator (3.0). See Table 1 for the location of particular individuals.
896

897    Figure 2. Comparison of sequential $\delta^{18}O$ values across multiple years of first molar formation
898    in five modern orangutans from Borneo and Sumatra.
899

900
901
902    Bornean individuals: MCZ 5290, ZSM 1981/48 ZSM 1981/87; Sumatran individuals: ZMB
903    83508, ZSM 1981/248.  The width of each curve is a kernel density estimate (KDE)
904    corresponding to the distribution of $\delta^{18}O$ values. First year data (Y1) is shown with a purple
905    violin plot, second year data (Y2) with a green plot, and third year data (Y3) with a yellow
906    plot where complete/available. Actual data are plotted as black circles.
907

908    Figure 3. Comparison of sequential δ¹⁸O values across multiple years of serial molar
909    formation in two modern orangutans from Borneo (top) and two from Sumatra (bottom).
910

911
912
913
914
915    Individual in upper left: ZSM 1981/48; upper right: ZSM 1981/87; lower left: ZSM
916    1981/246; lower right: ZSM 1981/248. Developmental overlap was determined through
917    registration of trace elements as in Smith et al. (2017).

923    Figure 4. Comparison of $\delta^{18}O$ values in fossil and modern orangutans from Borneo (blue) and
924    Sumatra (green).
925

926
927    Violin plots show kernel density estimates representing the distribution of $\delta^{18}O$ values in
928    modern individuals (left plot), and in fossil individuals (right plot). Actual $\delta^{18}O$
929    measurements are shown as black circles.
930

931    Table 1. Modern and fossil orangutan teeth employed in the current study.
932

| Taxon | Accession | Origin | Sex | Age (Years) | Teeth |
|---|---|---|---|---|---|
| *Pongo pygmaeus* | ZSM 1981/48 | Skalau, Borneo | F | ~8.4 | RUM1, LLM2 |
| | ZSM 1981/87 | Skalau, Borneo | F | > 9 | LUM1, RUM2, RLM3 |
| | MCZ 5290 | Borneo (location unspecified) | n/a | 4.5 | RUM1 |
| *Pongo abelii* | ZSM 1981/246 | Aceh, Sumatra | M | ~8.5 | LLM1, LUM2 |
| | ZSM 1981/248 | Aceh, Sumatra | F | adult | LUM1, LUM2, LLM3 |
| | ZMB 83508 | Sumatra (location unspecified) | n/a | 8.8 | RLM1 |
| Fossil *Pongo* spp. | 11564.5 | Sibrambang, Sumatra | n/a | n/a | RUM |
| | 11565.162 | Sibrambang, Sumatra | n/a | n/a | LUM |
| | 11594.12 | Lida Ajer, Sumatra | n/a | n/a | RLM |
| | 11595.105 | Lida Ajer, Sumatra | n/a | n/a | LLM |
| | US/22 | Niah Caves, Malaysia | n/a | n/a | RLM |
| | Y/F4 | Niah Caves, Malaysia | n/a | n/a | LLM |

933
934
935
936    Numerous taxonomic assignments have been made for fossil orangutans (*Pongo* spp.), some
937    of which have not been based on clear morphological characteristics (Lim, 2016), and are not
938    relevant for the focus of this paper.
939

940    Table 2. Modern and fossil orangutan molar $\delta^{18}O$ values.
941

| Taxon | Accession | Tooth | Cusp | Spots | Time (Days) | dO18 Range |
|---|---|---|---|---|---|---|
| *Pongo pygmaeus* | ZSM 1981/48 | RUM1 | dl | 151 | 1241 | 13.6-19.9 |
| | ZSM 1981/48 | LLM2 | mb | 107 | 804 | 13.0-18.8 |
| | ZSM 1981/87 | LUM1 | ml | 131 | 869 | 13.7-18.2 |
| | ZSM 1981/87 | RUM2 | ml | 196 | 1195 | 12.7-20.0 |
| | ZSM 1981/87 | RLM3 | mb | 220 | 1350 | 13.7-19.2 |
| | MCZ 5290 | RUM1 | ml | 150 | 1002 | 13.8-18.1 |
| *Pongo abelii* | ZSM 1981/246 | LLM1 | mb | 136 | 1425 | 12.3-18.3 |
| | ZSM 1981/246 | LUM2 | ml | 229 | 1376 | 12.6-18.0 |
| | ZSM 1981/248 | LUM1 | db | 177 | 1072 | 11.3-19.3 |
| | ZSM 1981/248 | LUM2 | db | 193 | 1374 | 13.5-20.6 |
| | ZSM 1981/248 | LLM3 | db | 191 | 1461 | 15.2-21.2 |
| | ZMB 83508 | RLM1 | db | 135 | 1029 | 13.4-20.4 |
| Fossil *Pongo* spp. | 11564.5 | RUM | mb | 178 | 1387 | 15.3-20.4 |
| | 11565.162 | LUM | ml | 143 | 1144 | 14.7-20.8 |
| | 11594.12 | RLM | ml | 154 | 1081 | 15.1-19.9 |
| | 11595.105 | LLM | mb | 197 | 1312 | 15.7-20.0 |
| | US/22 | RLM | mb | 149 | 1023 | 15.9-24.8 |
| | Y/F4 | LLM | db | 134 | 869 | 14.2-22.9 |

942
943

944    Supplemental Figure 1. Comparison of sequential $\delta^{18}O$ values across multiple years of molar
945    formation in six putative fossil orangutan M1s from Borneo and Sumatra.
946

947
948
The width of each curve is a kernel density estimate (KDE) corresponding to the distribution
of oxygen isotope values measured from different teeth. From each tooth, first year data (Y1)
is shown with a purple violin plot, second year data (Y2) with a green plot, and third year
data (Y3, if present) with a yellow plot. Actual data are plotted as black circles.

Supplemental Figure 2. Inferred seasonality of $\delta^{18}O$ values from a Bornean (top) and
Sumatran (bottom) M1.

955
956
Lomb-Scargle periodograms show potential periods in days (x-axis) against period power (y-
axis), where higher values on the y-axis indicate underlying sine-wave periods that produce,
contribute to, or explain $\delta^{18}O$ value oscillations. Best-fit periodicities are shown as light gray
vertical lines, whereas annual periodicities are indicated by blue dashed horizontal lines. We
regard the Bornean M1 (MCZ 5290) as largely aperiodic; a minor peak is observed at c. 6
months, and increasing powers at very high periods are an artifact of limited sampling length
within teeth relative to the model. The Sumatran M1 (ZMB 83508) has a 1.1 year inferred
frequency, as well as an approximately 6-month peak, likely reflecting semiannual monsoons.
965

Supplemental Figure 3. Laser ablation profiles performed across the two teeth from Lida
Ajer.

#576 (left) refers to specimen 11595.105; #578 (right) refers to specimen 11594.12. The red
dots represent the position of the rasters, and arrows indicate the sequence of the analyses.
EN = enamel. DE = dentine.

Supplemental Figure 4. Laser ablation profiles across the two teeth from Sibrambang Cave.

#577 (left) refers to specimen 11565.162; #581 (right) refers to specimen 11564.5. The red
dots represent the position of the rasters, and arrows indicate the sequence of the analyses.
EN = enamel. DE = dentine.

Supplemental Figure 5. Laser ablation profiles across the two teeth from Niah Caves.

#579 (left) refers to specimen from grid Y/F4; #580 (right) refers to specimen from grid
US/22. The red dots represent the position of the rasters, and arrows indicate the sequence of
the analyses. EN = enamel. DE = dentine.

984
985

986    Supplementary Figure 6. Matching trace element patterns in cross-sections of two isolated
987    molars from the Dubois collection of fossil orangutan teeth from Lida Ajer.
988

989
990
991    High concentrations are shown in warm colors, low concentration are in cool colors; here
992    Li/Ca is shown, but identical corresponding patterns were also observed for Ba/Ca and Sr/Ca
993    (not shown). The enamel cap of each tooth is to the left, and root dentine is to the right. Trace
994    elements were measured according to LA-ICP-MS methods detailed in Smith et al. (2017).
995

996    Supplementary Table 1. Comparisons of first and second year $\delta^{18}O$ values in five first molars.
997

| Specimen | Adjusted p-values | Higher $\delta^{18}O$ values |
|---|---|---|
| MCZ 5290 | p = 0.010 | Year 1 |
| ZMB 83508 | p = 0.006 | Year 1 |
| ZSM 1981/48 | p = 0.161 (N.S.) | Year 1 |
| ZSM 1981/87 | p < 0.001 | Year 2 |
| ZSM 1981/248 | p < 0.001 | Year 1 |

998
999

1000      Supplementary Table 2. U-series dates for fossil orangutan material.

1001

| LA raster | Tissue | U(ppm) | $^{230}$Th/$^{238}$U | ± | $^{234}$U/$^{238}$U | ± | Age (ka) | ± |
|---|---|---|---|---|---|---|---|---|
| 11595.105-EN11 | enamel | *0.02* | *-1.230* | *0.090* | *0.459* | *0.037* | n.c. | n.c. |
| 11595.105-EN12 | enamel | *0.00* | *-12.82* | *0.084* | *-4.590* | *0.018* | n.c. | n.c. |
| 11595.105-EN13 | enamel | *0.00* | *-31.04* | *0.082* | *-17.11* | *0.018* | n.c. | n.c. |
| 11595.105-EN14 | enamel | *0.10* | *0.623* | *0.168* | *1.155* | *0.105* | n.c. | n.c. |
| 11595.105-EN15 | enamel | *0.04* | *-0.826* | *0.082* | *0.752* | *0.051* | n.c. | n.c. |
| 11595.105-DE1 | dentine | *41.4* | *0.516* | *0.020* | *1.360* | *0.047* | 50.6 | 3.40 |
| 11595.105-DE2 | dentine | *44.2* | *0.470* | *0.020* | *1.359* | *0.055* | 45.2 | 3.33 |
| 11595.105-DE3 | dentine | *48.6* | *0.471* | *0.015* | *1.344* | *0.045* | 46.0 | 2.65 |
| 11595.105-DE4 | dentine | *58.9* | *0.481* | *0.015* | *1.356* | *0.038* | 46.6 | 2.50 |
| 11595.105-DE5 | dentine | *61.1* | *0.482* | *0.019* | *1.341* | *0.045* | 47.5 | 3.10 |
| 11595.105-DE6 | dentine | *65.8* | *0.446* | *0.019* | *1.346* | *0.051* | 43.0 | 3.04 |
| 11595.105-DE7 | dentine | *64.0* | *0.440* | *0.016* | *1.346* | *0.044* | 42.3 | 2.55 |
| 11595.105-DE8 | dentine | *60.0* | *0.430* | *0.016* | *1.338* | *0.046* | 41.4 | 2.56 |
| 11595.105-DE9 | dentine | *64.6* | *0.433* | *0.018* | *1.336* | *0.049* | 41.8 | 2.81 |
| 11595.105-DE10 | dentine | *65.9* | *0.411* | *0.016* | *1.325* | *0.047* | 39.7 | 2.53 |

1002

| LA raster | Tissue | U(ppm) | $^{230}$Th/$^{238}$U | ± | $^{234}$U/$^{238}$U | ± | Age (ka) | ± |
|---|---|---|---|---|---|---|---|---|
| 11594.12-EN1 | enamel | *0.00* | *-13.08* | *0.082* | *-9.394* | *0.019* | n.c. | n.c. |
| 11594.12-EN2 | enamel | *0.00* | *-23.91* | *0.104* | *-11.05* | *0.016* | n.c. | n.c. |
| 11594.12-EN3 | enamel | *0.00* | *-22.35* | *0.078* | *-4.621* | *0.027* | n.c. | n.c. |
| 11594.12-EN4 | enamel | *0.00* | *-16.49* | *0.089* | *-6.110* | *0.015* | n.c. | n.c. |
| 11594.12-EN5 | enamel | *0.01* | *17.58* | *0.103* | *16.21* | *0.031* | *224* | *3.12* |
| 11594.12-DE6 | dentine | 30.9 | 0.337 | 0.016 | 1.335 | 0.041 | 31.2 | 2.02 |
| 11594.12-DE7 | dentine | 30.6 | 0.347 | 0.013 | 1.337 | 0.036 | 32.2 | 1.68 |
| 11594.12-DE8 | dentine | 31.9 | 0.355 | 0.017 | 1.332 | 0.049 | 33.3 | 2.31 |
| 11594.12-DE9 | dentine | 31.4 | 0.366 | 0.013 | 1.335 | 0.035 | 34.4 | 1.78 |
| 11594.12-DE10 | dentine | 32.1 | 0.359 | 0.011 | 1.329 | 0.038 | 33.8 | 1.71 |
| 11594.12-DE11 | dentine | 24.6 | 0.344 | 0.014 | 1.357 | 0.031 | 31.4 | 1.67 |
| 11594.12-DE12 | dentine | 24.9 | 0.352 | 0.015 | 1.351 | 0.032 | 32.3 | 1.85 |

1003

| LA raster | tissue | U(ppm) | $^{230}$Th/$^{238}$U | ± | $^{234}$U/$^{238}$U | ± | Age (ka) | ± |
|---|---|---|---|---|---|---|---|---|
| 11565.162-EN1 | enamel | 0.58 | 0.434 | 2.630 | 1.025 | 1.495 | 59.7 | 496 |
| 11565.162-EN2 | enamel | *0.00* | *-4.899* | *0.132* | *-81.42* | *0.020* | n.c. | n.c. |
| 11565.162-EN3 | enamel | *0.00* | *-23.35* | *0.075* | *-16.84* | *0.018* | n.c. | n.c. |
| 11565.162-EN4 | enamel | *0.00* | *-37.09* | *0.082* | *-22.74* | *0.021* | n.c. | n.c. |
| 11565.162-EN5 | enamel | *0.01* | *-5.283* | *0.068* | *-0.022* | *0.029* | n.c. | n.c. |
| 11565.162-DE6 | dentine | 62.2 | 0.438 | 0.287 | 1.073 | 0.470 | 56.5 | 59.4 |
| 11565.162-DE7 | dentine | 63.6 | 0.461 | 0.023 | 1.066 | 0.048 | 61.2 | 5.73 |
| 11565.162-DE8 | dentine | 61.4 | 0.467 | 0.021 | 1.072 | 0.044 | 61.7 | 5.23 |
| 11565.162-DE9 | dentine | 59.8 | 0.470 | 0.028 | 1.073 | 0.060 | 62.2 | 7.01 |
| 11565.162-DE10 | dentine | 57.4 | 0.480 | 0.026 | 1.072 | 0.051 | 64.2 | 6.51 |

| LA raster | tissue | U(ppm) | $^{230}\text{Th}/^{238}\text{U}$ | ± | $^{234}\text{U}/^{238}\text{U}$ | ± | Age (ka) | ± |
|---|---|---|---|---|---|---|---|---|
| 11564.5-EN1 | enamel | 0.11 | 0.304 | 0.181 | 0.959 | 0.086 | 41.6 | 30.6 |
| 11564.5-EN2 | enamel | *0.00* | *-2215* | *0.086* | *-486.53* | *0.018* | n.c. | n.c. |
| 11564.5-EN3 | enamel | *0.00* | *-97.95* | *0.065* | *-4.236* | *0.031* | n.c. | n.c. |
| 11564.5-EN4 | enamel | 9.86 | 0.502 | 0.211 | 1.115 | 0.162 | 64.3 | 38.8 |
| 11564.5-DE5 | dentine | 51.6 | 0.564 | 0.055 | 1.116 | 0.052 | 75.4 | 11.8 |
| 11564.5-DE6 | dentine | 45.8 | 0.541 | 0.057 | 1.111 | 0.047 | 71.6 | 11.6 |
| 11564.5-DE7 | dentine | 49.1 | 0.530 | 0.039 | 1.117 | 0.079 | 69.0 | 10.2 |
| 11564.5-DE8 | dentine | 47.8 | 0.512 | 0.010 | 1.128 | 0.016 | 65.0 | 2.13 |

1004

| LA raster | tissue | U(ppm) | $^{230}\text{Th}/^{238}\text{U}$ | ± | $^{234}\text{U}/^{238}\text{U}$ | ± | Age (ka) | ± |
|---|---|---|---|---|---|---|---|---|
| Y/F4-EN1 | enamel | *0.01* | *-4.737* | *0.084* | *-0.283* | *0.025* | n.c. | n.c. |
| Y/F4-EN2 | enamel | *0.00* | *-42.98* | *0.070* | *-2.119* | *0.024* | n.c. | n.c. |
| Y/F4-EN3 | enamel | *0.00* | *-137.8* | *0.078* | *-18.50* | *0.017* | n.c. | n.c. |
| Y/F4-EN4 | enamel | *0.00* | *-49.66* | *0.079* | *3.398* | *0.033* | n.c. | n.c. |
| Y/F4-DE5 | dentine | 4.41 | 0.058 | 0.016 | 1.082 | 0.031 | 5.97 | 1.74 |
| Y/F4-DE6 | dentine | 4.93 | 0.072 | 0.018 | 1.057 | 0.034 | 7.69 | 1.99 |
| Y/F4-DE7 | dentine | 4.80 | 0.077 | 0.018 | 1.088 | 0.031 | 7.95 | 1.93 |
| Y/F4-DE8 | dentine | 4.15 | 0.085 | 0.041 | 1.102 | 0.065 | 8.74 | 4.43 |

1005

| LA raster | tissue | U(ppm) | $^{230}\text{Th}/^{238}\text{U}$ | ± | $^{234}\text{U}/^{238}\text{U}$ | ± | Age (ka) | ± |
|---|---|---|---|---|---|---|---|---|
| US/22-EN1 | enamel | *0.00* | *-7.442* | *0.102* | *-6.047* | *0.023* | n.c. | n.c. |
| US/22-EN2 | enamel | *0.00* | *-53.99* | *0.082* | *-14.98* | *0.020* | n.c. | n.c. |
| US/22-EN3 | enamel | *0.00* | *215.0* | *0.103* | *78.83* | *0.019* | n.c. | n.c. |
| US/22-EN4 | enamel | *0.00* | *2.054* | *0.109* | *-12.69* | *0.020* | n.c. | n.c. |
| US/22-EN5 | enamel | *0.02* | *-2.955* | *0.109* | *1.044* | *0.052* | n.c. | n.c. |
| US/22-DE6 | dentine | 1.28 | 0.066 | 0.070 | 1.205 | 0.100 | 6.09 | 6.66 |
| US/22-DE7 | dentine | 1.36 | 0.085 | 0.073 | 1.179 | 0.086 | 8.10 | 7.22 |
| US/22-DE8 | dentine | 1.38 | 0.094 | 0.060 | 1.183 | 0.071 | 8.94 | 5.98 |
| US/22-DE9 | dentine | 1.41 | 0.126 | 0.062 | 1.183 | 0.087 | 12.27 | 6.38 |
| US/22-DE10 | dentine | 1.41 | 0.090 | 0.050 | 1.234 | 0.080 | 8.17 | 4.77 |

1006

$^{230}\text{Th}/^{238}\text{U}$ and $^{234}\text{U}/^{238}\text{U}$ are activity ratios. It is worth noting that, for most transect analyses,
the $^{232}\text{Th}$ signal, which was measured on a Faraday collector, was indistinguishable from
background noise. In this regard, the corresponding $^{230}\text{Th}/^{232}\text{Th}$ activity ratio of each transect
should be >>100, and thus non-radiogenic or detrital $^{230}\text{Th}$ correction would have negligible
impact on the age. U-series data in italics should be viewed with caution due to U
concentrations of ≤ 0.5 ppm. All errors are 2-σ. Key: EN= enamel; DE = dentine; n.c. = not
calculable. Negative values were caused by background extraction from their measured peaks
with intensities at detection levels.