1 **Oxygen isotopes in orangutan teeth reveal recent and ancient climate variation**
2

3 Tanya M. Smith[1,2], Manish Arora[3], Christine Austin[3], Janaína N. Ávila[1,4], Mathieu Duval[2,5,6],
4 Tze Tshen Lim[7], Philip J. Piper[8], Petra Vaiglova[1,2,8], John de Vos[9], Ian S. Williams[10], Jian-
5 xin Zhao[11], Daniel R. Green[2,12]
6

1 Griffith Centre for Social and Cultural Research, Griffith University, Australia
2 Australian Research Centre for Human Evolution, Griffith University, Australia
3 Department of Environmental Medicine and Public Health, Icahn School of Medicine at
Mount Sinai, USA
4 School of the Environment, The University of Queensland, Australia
5 Centro Nacional de Investigación sobre la Evolución Humana (CENIEH), Spain
6 Palaeoscience Labs, Department of Archaeology and History, La Trobe University,
Australia
7 Department of Geology, Universiti Malaya, Malaysia
8 School of Archaeology and Anthropology, The Australian National University, Australia
9 Department of Geology, Naturalis Biodiversity Center, The Netherlands
10 Research School of Earth Sciences, The Australian National University, Australia
11 Radiogenic Isotope Facility, School of the Environment, The University of Queensland,
Australia
12 Department of Human Evolutionary Biology, Harvard University, USA
22

Abstract

Studies of climate variation commonly rely on chemical and isotopic changes

recorded in sequentially-produced growth layers, such as in corals, shells and tree rings, as

well as in accretionary deposits—ice and sediment cores, and speleothems. Oxygen isotopic

compositions ($\delta^{18}O$) of tooth enamel are a direct method of reconstructing environmental

variation experienced by an individual animal. Here we utilize long-forming orangutan

dentitions (*Pongo* spp.) to probe recent and ancient rainfall trends on a weekly basis over ~

3–11 years per individual. We first demonstrate the lack of any consistent isotopic

enrichment effect during exclusive nursing, supporting the use of primate first molar teeth as

environmental proxies. Comparisons of $\delta^{18}O$ values (n = 2016) in twelve molars from six

modern Bornean and Sumatran orangutans reveal a high degree of overlap, with more

consistent annual and bimodal rainfall patterns in the Sumatran individuals. Comparisons

with fossil orangutan $\delta^{18}O$ values (n = 955 measurements from six molars) reveal similarities

between modern and late Pleistocene fossil Sumatran individuals, but differences between

modern and late Pleistocene/early Holocene Bornean orangutans. These suggest drier and

more open environments with reduced monsoon intensity during this earlier period in

northern Borneo, consistent with other Niah Caves studies and long-term speleothem $\delta^{18}O$

records in the broader region. This approach can be extended to test hypotheses about the

paleoenvironments that early humans encountered in southeast Asia.

Introduction

Present-day rainfall patterns in Indonesia are controlled by the Asian and Australian monsoon systems, yielding annual trends that vary considerably with geography, topography, and the direction of monsoonal winds (Aldrian and Susanto, 2003; Moron et al., 2009; Qian et al., 2013; Belgaman et al., 2017). Northern Sumatra and western Borneo experience high annual rainfall and relatively stable annual temperatures, with a bimodal distribution of rainfall governed by the Intertropical Convergence Zone (van Schaik, 1986; Aldrian and Susanto, 2003; Belgaman et al., 2017). These islands are also under the influence of inter-annual climate fluctuations driven by the El-Niño Southern Oscillation (ENSO); a periodic coupling of atmospheric and oceanic temperature gradients that initiates in the tropical Pacific, and influences global temperature and precipitation trends (McPhaden et al., 2006).

It is well understood that variation in rainfall patterns influences the fundamental structure of primate habitats (Brockman and van Schaik, 2005; Wessling et al., 2018). Dense tropical forests are sustained by fairly consistent rainfall and short, irregular dry seasons, while woodland communities in more arid environments have smaller trees, less dense canopies, and more deciduous trees (Vico et al., 2017; Archibald et al., 2019). In regions with prolonged dry seasons, low annual rainfall and savannah landscapes abound, in addition to disturbances such as wildfires (Pletcher et al., 2022).

Open woodland and savannah environments are unfavorable for slow-moving orangutans, the largest mammal with an arboreal lifestyle, particularly in regions with predators such as tigers or humans (Thorpe and Crompton, 2009; Ashbury et al., 2015; Spehar et al., 2018). Supra-annual ENSO events may also impact orangutan energy balance, reproduction, and social organization through the inducement of mast-fruiting, or dramatic seed production events in dipterocarp forests (Knott 1998; Curran et al., 1999; Marshall et al., 2009). Such climate fluctuations over the past several hundred years have been documented in coral isotopes and tree-ring analyses, revealing especially marked changes during the past few decades (Cole et al., 1993; Stahle et al., 1998; Hughen et al., 1999; Urban et al., 2000; Tudhope et al., 2001; Pumijumnong et al., 2020).

Detailed climate records prior to the era of human-induced climate change are somewhat limited for island southeast Asia, but they are directly relevant to understanding the recent distribution of orangutans, and the arrival and dispersal of modern humans in the region during the Late Pleistocene (e.g., Piper 2016; Bae et al., 2017; Spehar et al., 2018). A small number of studies of fossil corals, molluscs, marine sediments, and speleothems has

provided insights into the last interglacial and glacial periods (e.g., Hughen et al., 1999;

Tudhope et al., 2001; Stephens et al., 2016; Yang et al., 2016; Buckingham et al. 2022). For

example, oxygen isotopes in fossil corals from seven periods during the last 130,000 years

suggest that ENSO activity in the western Pacific over that time was comparable to modern

records, although there was variation in the intensity of such activity at different timepoints

(Tudhope et al., 2001). This study was also able to resolve bimodal annual rainfall peaks in

modern corals, yet such detailed subannual records are extremely uncommon, particularly

from terrestrial environments where early humans once lived alongside orangutans and other

mammals.

Oxygen isotope studies for paleoenvironmental reconstruction

Oxygen isotope values ($\delta^{18}O$) in water vary with latitude, altitude, temperature and

precipitation cycles, and are also impacted by precipitation sources. In tropical regions the

primary determinant of rainfall isotope compositions is rainfall amount (Dansgaard, 1964;

Rozanski et al., 1993; Belgaman et al., 2017). During wet seasons, rainfall $\delta^{18}O$ values are

relatively low, while the opposite pattern is evident in periods with less rain, although other

meteorological factors can influence isotope values as well (Belgaman et al., 2016). This

primary tropical pattern influences isotopic variation in meteoric, surface, and leaf waters,

which may show further elevations in $\delta^{18}O$ values during dryer periods due to preferential

evaporative loss of the lighter isotope, ^{16}O (da Silveira et al., 1989; Bowen, 2010; Roberts et

al., 2017).

In addition to $\delta^{18}O$ values in fossil corals, tree rings and speleothems, other fine-

scaled oxygen isotopic climate proxies include otoliths (fish ear bones) and mollusc shells

(e.g., Aubert et al., 2012; Stephens et al., 2016; Prendergast et al., 2018)—although these are

rarely preserved in rainforest environments. Records of $\delta^{18}O$ values in mammalian tooth

enamel are a more direct means of studying seasonality (reviewed in Green et al., 2018,

2022), providing insight into the actual climates experienced by individuals, in contrast to

indirect proxies for which it can be difficult to establish concurrence. Unlike bone, teeth do

not remodel during life, and the phosphate component of the enamel mineral (hydroxyapatite)

is especially resistant to modification after burial (reviewed in Smith et al., 2018a; Pederzani

and Britton, 2019).

Tooth enamel is most commonly sampled with hand-held drills to recover the isotopic

composition of oxygen inputs from water and food preserved in the hydroxyapatite (e.g.,

Janssen et al., 2016; Roberts et al., 2020; Kubat et al., 2023). This coarse drilling method

yields spatially and temporally blurred powdered samples formed over a substantial and

unknown period of time, however, precluding the identification of precise seasonal

environmental patterns. To circumvent this limitation, we have employed the stable isotope

sensitive high-resolution ion microprobe (SHRIMP SI) to measure $\delta^{18}O$ values sequentially

from thin sections of teeth, relating these to daily increments and birth lines to determine

enamel formation times, and in some instances, calendar ages (Smith et al., 2018a, 2022;

Green et al., 2022; Vaiglova et al., in review).

It is well established that $\delta^{18}O$ values in tooth enamel are closely related to local water

oxygen isotope compositions (reviewed in Green et al., 2018, 2022). For teeth that form after

birth and during periods of milk consumption, $\delta^{18}O$ values are expected to be higher, as a

result of infant evaporative water loss while consuming [18]O-enriched mother's milk (Bryant

et al., 1996; Wright and Schwartz 1999; Britton et al., 2015). Studies of large-bodied

mammals report that milk $\delta^{18}O$ values are elevated by ~ 1–6 ‰ relative to local drinking

water $\delta^{18}O$ (Kornexl et al., 1997; Lin et al., 2003; Chesson et al., 2010; Green et al., 2018; but

see Cherney et al., 2010). Comparable data on human or nonhuman primate milk enrichment

appear to be lacking, save for a study of 44 British infants aged 5–16 weeks (Roberts et al.,

1988). The urine of infants who were breast-fed showed isotopic enrichment of 1–3 ‰

compared to infants who were fed formula prepared from sterile local tap water.

While such studies point to potential changes in infant body water during nursing, it is

unclear whether such differences prohibit the use of early-formed enamel in studies of

climate variation (Blumenthal et al., 2017; Luyt and Sealey 2018). Two studies of $\delta^{18}O$

values in the dentitions of modern sheep, horses, and zebras reported higher bulk values (~ 1–

2 ‰) in five molars (M1) compared to the rest of the permanent dentition (Bryant et al.,

1996; Fricke and O'Neil, 1996). This led Fricke and O'Neil (1996) to suggest that M1s are

unlikely to reflect the values of local meteoric water due to the influence of maternal inputs in

utero and through lactation. However, near-weekly $\delta^{18}O$ values over the first 2.75 years of

life in a Neanderthal M1 measured with SHRIMP SI showed clear annual trends and

maximum $\delta^{18}O$ values corresponding to a period after nursing has ceased (Smith et al.,

2018a). An examination of longer continuous periods of enamel formation within and

between teeth will help to clarify whether early-formed primate teeth should be avoided for

studies of climate seasonality.

Here we first assess whether wild orangutans show elevated $\delta^{18}O$ values in early-

formed enamel, testing the suggestion that M1s are significantly affected by nursing [18]O-

enrichment, thereby precluding their use in climatological reconstructions. We then explore

approximately 30 years of weekly $\delta^{18}O$ values (n = 2016 measurements) to compare

orangutan individuals from the islands of Sumatra and Borneo. Finally, we contrast $\delta^{18}O$

values between modern and Pleistocene orangutans, including those from key regions of

early human occupation: Lida Ajer, Sumatra (Hooijer, 1948; Westaway et al., 2017) and Niah

Caves, Malaysia (Hooijer, 1961; Barker et al., 2007) (Figure 1, Table 1). Novel

understanding of climate patterns in these fossil assemblages may inform debates about the

likelihood of modern humans living in dense Asian rainforests, and the conditions that would

support savannah corridors for human dispersals throughout the region (e.g., de Vos, 1983;

Bird et al., 2005; Westaway et al., 2017; Louys and Roberts, 2020).

154   Insert Figure 1 and Table 1 here

Results

Modern orangutans

The $\delta^{18}O$ ranges of twelve modern and six fossil orangutan molars, representing 2971

near-weekly measurements spanning 57.6 years of tooth formation, are listed in Table 2.

Prior to making comparisons between individuals, geographic regions, or time periods, we

first consider the potential intra-individual effect of isotopic enrichment from maternal milk

on $\delta^{18}O$ values. Comparisons of $\delta^{18}O$ values during the first, second, and third years of life in

five modern orangutan first molars (M1) do not show consistently elevated values during

their first year (Figure 2). Mean yearly $\delta^{18}O$ values in the first year are elevated by only 0.3

‰ compared to the second year. While three of the five M1s showed first year $\delta^{18}O$ values

higher than second year values ($p \leq 0.05$), only two individuals showed mean values that

were ~1 – 2 ‰ higher during year one; one individual showed no difference from the first to

the second year, and one individual showed lower values during the first year than during the

second year ($p \leq 0.05$) (Supplementary Table 1). A sixth individual was only sampled from

193 days of age, but maximum values from this point onwards were similar across more than

three years of life. Similarly variable patterns were observed for the six putative fossil

orangutan M1s (Supplementary Figure 1).

174   Insert Figure 2 and Table 2 here

Comparisons across serial molars in four modern orangutans show no consistent trend

of elevated $\delta^{18}O$ values in M1s relative to successive molars (Figure 3). Only two individuals

showed maximum $\delta^{18}O$ values in their M1s relative to M2s; in both instances M3s were

unavailable due to their lack of development prior to death. The other two individuals showed

higher $\delta^{18}O$ values in M2s or M3s than in their respective M1s. In the case of the oldest

individual (ZSM 1981/248), the highest $\delta^{18}O$ values appeared at approximately 5.8 years of

age, well past the age when exclusive nursing ends.

184   Insert Figure 3 here

Comparison of the $\delta^{18}O$ values in the full datasets of modern Bornean and Sumatran

orangutans reveals a high degree of overlap. Values from the three Bornean individuals

ranged from 12.7 to 20.0 ‰ (n = 955 near weekly measurements), while the three Sumatran

individuals ranged from 11.3 to 20.6 ‰ (n = 1061 measurements). Comparisons of periodic

trends via spectral power distribution analysis revealed more consistent bimodal patterns in

the Sumatran individuals; three of the six Bornean molars were aperiodic (statistical power of

0.1 or less), while all six of the Sumatran molars revealed annual or semiannual cycles with

greater power (Supplementary Figure 2). Rapid oxygen isotopic shifts on the order of ~ 6–8

‰ are evident in the single Bornean and Sumatran individuals with $\delta^{18}O$ measurements

spanning M1 to M3, which may represent one or more supra-annual ENSO events captured

during the ~ 9–11 years these molars were forming.

Fossil orangutans – oxygen isotopes

Concurrently forming teeth (molar specimens 11594.12 and 11595.105) from same

individual at Lida Ajer, Sumatra are nearly isotopically identical; $\delta^{18}O$ values range from

15.1 to 19.9 ‰ and 15.7 to 20.0 ‰, respectively, supporting the biogenic fidelity of these

records. The $\delta^{18}O$ values of two individuals from the nearby Sibrambang site (15.3–20.4 ‰,

14.7–20.8 ‰) are very similar to those of the Lida Ajer individual. These Sumatran fossils all

fall at the upper end of the range of modern Sumatran orangutans (Figure 4), and reveal

approximately annual $\delta^{18}O$ periodicities (0.9–1.3 years), as well as strong bimodal

distribution patterns in one instance (11565.162).

208   Insert Figure 4 here

The two fossils from the Niah Caves were excavated from different regions and

stratigraphic depths; $\delta^{18}O$ values in the tooth from grid US/22 ranged from 15.9 to 24.8 ‰

and, unlike the three modern Bornean individuals, yielded an annual periodicity (1.0 years).

The $\delta^{18}O$ in the tooth from grid Y/F4 ranged from 14.2 to 22.9 ‰ and showed a stronger

bimodal trend than an annual one, although its short formation time may have prohibited

identification of longer trends. The range of values from these two fossil molars (14.2–24.8

‰) markedly exceeds the range of modern Bornean orangutans (12.7–20.0 ‰) (Figure 4),

with the mean $\delta^{18}O$ value at least 2 ‰ heavier. This suggests possibly drier conditions with

greater seasonality during fossil molar formation (Supplementary Figure 3).

Fossil orangutans – U-series age estimates

The six fossil teeth have very low uranium concentrations in their enamel (<0.5 ppm),

regardless of their origin (Supplementary Table 2). These enamel values are very close to the

detection limit of the Nu Plasma II MC-ICP-MS, and thus are not useful for estimating

minimum ages. The dentine of Lida Ajer specimen 11595.105 shows a spatial gradient of

increasing uranium concentration from ~ 41 to 66 ppm, and decreasing age estimates from ~

51 to 40 ka (Supplementary Table 2). This trend might result from a preferential uranium

leaching overprint near the end of the root. Spot DE10, positioned near the EDJ, is less likely

to be impacted (Supplementary Figure 4), and is thus assumed to provide the most reliable

minimum age for the tooth, ~ 40 ka. Uranium values from Lida Ajer specimen 11594.12

show a similar trend of concentrations decreasing from ~ 31 to 24 ppm towards the root tip.

However, the U-series age estimates remain constant within the range 31–34 ka across the

dentine (Supplementary Table 2; Supplementary Figure 4). No evidence for a recent

overprint is observed, supporting a minimum age of 33 ka. In summary, this individual's age

is at least 33 ka, and possibly > 40 ka.

U-series analysis of the dentine of Sibrambang specimen 11565.162 shows a slight

decreasing trend of uranium concentration from the EDJ to the root tip (from > 60 ppm to <

60 ppm), and corresponding increasing age estimates (56–62 ka) (Supplementary Table 2;

Supplementary Figure 5). This might result from a slight uranium leaching overprint; a

minimum age of 60 ka is likely for this tooth. The U-series age estimates obtained for

Sibrambang specimen 11564.5 show a decreasing trend from the EDJ toward the

circumpulpal dentine from 75 to 65 ka (Supplementary Table 2; Supplementary Figure 5.)

However, given the associated uncertainties, this trend might not be meaningful. An average

dentine U-series age of 70.3 ± 5.5 ka (2σ) may be regarded as a minimum age for the fossil,

which is broadly consistent with the single age estimate obtained from the enamel (64 ka). In

summary, the two teeth from Sibrambang yield U-series apparent ages of ~ 60–70 ka.

The uranium concentration measured across the dentine of the Niah Caves specimen

from grid Y/F4 shows little variability, 4.2–4.9 ppm. The U-series age estimates are between

6.0 and 8.7 ka (Supplementary Table 2; Supplementary Figure 6). The average dentine U-

series minimum age is 7.6 ± 1.3 ka. Similarly, the Niah Cave specimen from grid US/22

shows a consistent uranium concentration through the dentine (1.3–1.4 ppm), with relatively

large uncertainties that nonetheless bracket individual age estimates (Supplementary Table 2;

Supplementary Figure 6). The average dentine U-series age is 8.8 ± 3.0 ka. In summary, the

two teeth from Niah Cave yield consistent apparent ages of ~ 8–9 ka, which should be

regarded as a minimum age constraint for the fossils.

Discussion

Primate oxygen isotope compositions do not reveal a clear milk enrichment effect

Half of our modern sample, and potentially all of our fossil sample, are composed of

M1s. These begin forming around birth and continue growing for three or more years (Smith,

2016). Orangutan infants rely exclusively on maternal milk during their first year of life,

supplementing this with solid foods in the second year, which are increased until suckling

ceases prior to eight or nine years of age (van Noordwijk et al., 2013; Smith et al., 2017). Our

developmentally-guided sampling approach allows us to examine fine-scaled trends in δ^{18}O

values during birth, exclusive nursing, supplemental feeding, and also after nursing ends (in

those individuals with available serial molar teeth).

We find that five modern orangutans show only minor and inconsistently elevated

δ^{18}O values during the first year of life when compared to the subsequent year. These data do

not support the hypothesis that primate infants have markedly elevated body water δ^{18}O

values during exclusive nursing. Data from the majority of 12 human M1s studied by

Vaiglova et al. (in review) similarly reveal maximum δ^{18}O values after the first year of tooth

formation, well beyond the duration of exclusive milk intake. This is also evident in the M1

of a Neanderthal born in the spring (Smith et al., 2018a); δ^{18}O values mostly rose for the first

3.5 months of life, but did not reach a maximum for another two years — long after the infant

would have begun consuming supplemental foods and liquids. This final dataset points to the

influence of season of birth on initial postnatal δ^{18}O values, as inferred in other mammals

(Bryant et al., 1996; Frick and O'Neil, 1996).

Comparisons of serially-forming teeth in four wild orangutans also fails to show a

consistent elevation of δ^{18}O values in M1s versus M2s (or M3s in two cases). Comparisons of

M1 δ^{18}O values with subsequent-forming teeth in four baboons, two tantalus monkeys, and

one mona monkey (from Green et al., 2022: SI Dataset S1) also largely fail to support the

enriched "Pattern 1" trend modelled by Bryant et al. (1996: Figure 4, p. 401). This is also the

case in comparisons of δ^{18}O values from bulk samples of human teeth—Wright and Schwartz

(1999) demonstrated that M1s have higher δ^{18}O values than later-forming teeth in only four

of seven individuals. In summary, the data from a range of primates, humans included, do not

support the exclusion of early-forming primate teeth from the assessment of environmental

seasonality.

Modern orangutans show similar isotopic values across the islands of Borneo and Sumatra

The two Bornean juveniles from the Munich collection (ZSM 1981/48, ZSM 1981/87)

reflect the environmental conditions of the late 1880s and early 1890s in Skalau—a region

where orangutans might now be locally extinct. Similarly, the teeth from the two Sumatran

individuals from the Munich collection (ZSM 1981/246, ZSM 1981/248) were collected prior

to 1939 in northern Aceh, from where orangutans also have since disappeared (Spehar et al.,

2018). While the individuals from northernmost Sumatra might have inhabited somewhat

higher elevations than those from western Borneo, there does not appear to be an evident

altitude effect (lower isotopic values at higher altitudes), as these four individuals show

similar isotopic values, save for a single brief excursion below 12 ‰ in ZSM 1981/248

(Table 1, Figure 3). It is unknown to what extent local rainfall may have been isotopically

distinct at the time the teeth were forming.

The δ^{18}O values shown in Figure 1 reflect estimates of monthly and annual average

precipitation from the Online Isotopes in Precipitation Calculator (3.0) compiled for

www.waterisotopes.org. Actual measurements of precipitation δ^{18}O from the islands of

Borneo and Sumatra are extremely limited. The closest observation facilities to the ZMS

orangutan locations yield similar patterns of modern annual rainfall δ^{18}O variability

(Belgaman et al., 2017), yet specific measurements from the six facilities that make up

"Cluster 3" in this reference are not available for comparison.

Other studies underscore the complexity of water transport in this region—multiple

factors such as the oceanic origin of water vapor, cloud cover and type, and the post-

condensation process influence the short-term variability of δ^{18}O values in rainfall (Moerman

et al., 2013; Suwarman et al., 2013; Belgaman et al., 2016). For example, Moerman et al.

(2013) provided five years of daily rainfall δ^{18}O measurements from Northern Borneo

(Gunung Mulu National Park, Malaysia); daily rainfall δ^{18}O values ranged from + 0.7 ‰ to

−18.5 ‰ and showed 1 – 3 month, annual, and supra-annual cycle frequencies. Interannual

rainfall $\delta^{18}O$ fluctuations of 6–8 ‰ were significantly correlated with ENSO events; these are

similar in scale to the large fluctuations in our serial tooth datasets (Figure 3).

Another potential source of isotopic variability derives from dietary variation, as

orangutans obtain the majority of their body water from plants (MacKinnon, 1974). Plant

oxygen isotope compositions can be stratified within tropical forest canopies (Sternberg et al.,

1989; Roberts et al., 2017; Lowry et al. 2021)—potentially leading to offset values amongst

various animals, including primates, that consume different resources in the same forest

(Krigbaum et al., 2013; Nelson, 2013; Fannin and McGraw, 2020). Orangutans forage at

different canopy heights ranging from the ground to high in the canopy (MacKinnon, 1974;

Ungar, 1996; Thorpe and Crompton, 2005; Ashbury et al. 2015). MacKinnon (1974) reported

that Bornean and Sumatran orangutans obtain 95 % of their food from the middle and upper

levels of the canopy, where preferred foods are most abundant. In contrast, Ungar (1996)

reported that Sumatran orangutans were quite variable in feeding heights, with a mean of

approximately 19 meters; lower than gibbons who fed preferentially in the high canopy.

Thorpe and Crompton (2005) reported stratification in Sumatran orangutans, with immature

individuals feeding below 20 meters, females feeding both below and above this height, and

adult/subadult males preferring to feed high in the canopy.

While differences in enamel $\delta^{18}O$ values are apparent in comparisons of sympatric

arboreal and terrestrial mammals (reviewed in Lowry et al. 2021; Green et al., 2022), it

remains to be seen whether primates with broadly similar diets and habitats show meaningful

differences in $\delta^{18}O$ values, and to what degree plant physiology influences the pattern and

amplitude of seasonality relative to rainfall. Oxygen isotope compositions in the six modern

individuals from the islands of Borneo and Sumatra are very similar. Orangutans from both

islands prefer ripe fruit when available, with some differences in the consumption of bark,

leaves, unripe fruits, and insects—which varies between sites and across seasons (reviewed in

Smith et al., 2012). Seasonal variation in diets and the stratification of food within the canopy

may also contribute to enamel oxygen isotope variation within individuals, in addition to the

seasonal rainfall trends we observe in our datasets. Orangutan $\delta^{18}O$ values are also quite

similar to the $\delta^{18}O$ values from five humans from Flores, Indonesia (14.8–21.0 ‰) dated at ~

2.2–3.0 ka (Vaiglova et al., in review). This is remarkable given the major dietary differences

between frugivorous orangutans and omnivorous coastal-dwelling humans, and suggests that

their enamel $\delta^{18}O$ values are predominantly influenced by regional precipitation.

Fossil orangutan isotope values suggest different ancient climates in Sumatra and Borneo

Dating studies at Lida Ajer have established the presence of the oldest human remains

in insular Southeast Asia, ~ 63–73 ka (Westaway et al., 2017), and a broad survey of the cave

has reconfirmed an age of MIS 4 (59–76 ka) for the mammalian fauna (Louys et al., 2022).

This is consistent with the minimum age of ~ 33–40 ka estimated for the two molars

examined in the current study. The Sumatran Sibrambang Cave has been regarded as roughly

contemporaneous to Lida Ajer given broad faunal similarities (de Vos, 1983). Recent U-

series dating of two fossil orangutans from the Sibrambang assemblage yielded minimum

ages of >56 ka and >85 ka (Louys et al. in press), which bracket the apparent U-series

minimum ages of ~ 60–70 ka in the current study. Sibrambang primates appear similar to, or

slightly older than, those from Lida Ajer, given the minimum U-series age estimates for teeth

from both sites, but this is not definitive given the absence of finite numerical ages for the

fossils. Our analysis of $\delta^{18}O$ values in Sumatran orangutan fossil molars reveals a close

similarity across sites and with modern Sumatran individuals, although the fossil

compositions fall at the upper end of the modern range. This may indicate a slightly dryer and

less variable climate during the late Pleistocene; elevated tooth $\delta^{18}O$ values are also indicative

of elevated values in hydrological systems globally, resulting from increased ice volumes at

in glaciers and at the poles.

Pollen records from the Niah Caves archaeological site indicate that there were a

number of local ecological shifts from lowland rainforest to more open environments during

the Late Pleistocene and into the Holocene (Hunt et al., 2012), where humans may have

begun hunting orangutans at ~ 45 ka (Spehar et al., 2018). While it is not possible to locate

the two fossil orangutan molars in these pollen records, Piper and Rabett (2016) considered

that the large animal bone assemblages accumulated within the Lobang Hangus entrance and

defined by the Harrisson spit depths of 12"- 42" were of terminal Pleistocene age. More

broadly, the orangutan specimen from grid US/22 (32"-36") is stratigraphically positioned

between radiocarbon ages of 14,206−15,061 cal. BP (OxA-13936) and 36,583−38,059 cal.

BP (OxA-13938), and this provides plausible minimum and maximum age constraints that

are not incompatible with the apparent minimum U-series age of ~ 9 ka. Based on these

results, the tooth is likely to date from the latest part of the Late Pleistocene. The specimen

from grid Y/F4 might date from the latest part of the Late Pleistocene to the early Holocene,

by comparison with the shell and fauna assemblage from other excavated areas (Piper et al.,

2016.)

Both orangutan molars from the Niah Caves yield wide ranges of $\delta^{18}O$, which is

particularly notable given the short periods of time sampled compared to the other fossils and

most modern orangutan molars. Given the similar offsets in $\delta^{18}O$ values between modern

baboons living in Ugandan forests and the Ethiopian rift region (Green et al., 2022) and

modern and prehistoric Bornean orangutans, we regard the higher $\delta^{18}O$ values in the Niah

Cave orangutans as possibly indicative of reduced rainfall when compared to recent

conditions. This is consistent with paleoclimate reconstructions for Borneo and Flores during

the late Pleistocene and early Holocene (Griffith et al. 2009; Buckingham et al., 2022), when

the environment around the Niah Caves is believed to have been a drier, more open seasonal

forest (Harrison, 1996; Hunt et al., 2012). A study of $\delta^{18}O$ values in Niah Caves shell

middens dating from the early to mid- Holocene indicates a shift to periods of high rainfall

with less variation than modern conditions (Stephens et al., 2016). The transition from a drier

environment to moist tropical rainforest is also reflected in the increasing number and higher

frequencies of canopy-adapted mammalian taxa in excavated layers of the Pleistocene-

Holocene transition (Piper and Lim, 2021).

Our approach has the potential to contribute to reconstructions of ancient

paleoenvironments in SE Asia based on studies of pollen, molluscs, faunal community

compositions, guano records, and stable isotopes of teeth (e.g., Jablonski et al., 2000; Bird et

al., 2005; Louys and Meijaard, 2010; Wurster et al., 2010; Hunt et al., 2012; Janssen et al.,

2016; Stephens et al., 2016; Louys and Roberts, 2020; Bacon et al., 2021; Louys et al., 2022,

in press). This may be especially timely given that recent work examining modern fauna

compositions in African landscapes has cautioned that fossil herbivore assemblages tend to

overestimate the extent of ancient grasslands in comparison to woodlands (Negash and Barr,

2023; also see Sokolowski et al. 2023). Fine-scaled tooth sampling may also allow an

expansion of inferences from $\delta^{18}O$ values of bulk-sampled Asian hominin remains (Janssen et

al., 2016; Roberts et al., 2020; Kubat et al., 2023), which are difficult to interpret for

understanding seasonal rainfall dynamics in tropic environments (Green et al., 2022). Such

information could better inform debates about whether humans employed arid savannah

corridors to avoid dense tropical forests, or whether humans were adept at colonizing such

environments during their consequential migration throughout island Southeast Asia.

Materials and Methods

Orangutan samples

Thin (histological) sections of twelve molar teeth from six modern orangutans and six
molar teeth from five fossil orangutans were employed (Table 1). These sections were
previously prepared for studies of incremental tooth development, enamel thickness,
elemental chemistry, and Asian hominoid taxonomy (Smith, 2016; Smith et al., 2011, 2012,
2017, 2018b). Four modern individuals were sourced from the Munich State Anthropological
Collection (ZSM); two were collected in 1893-1894 from Skalau (north of the Kapuas River
and south of the Klingkang Mountains in eastern West Borneo), and two were collected prior
to 1939 from Aceh (northwest Sumatra) (Röhrer-Ertl, 1988: Figure 3, p. 14) (Figure 1). It
was not possible to determine from which specific regions or time periods the two other
modern individuals derive—collection notes were not available for these specimens from the
Harvard Museum of Natural History (MCZ) or the Humboldt Museum (ZMB). Ages at death
were determined for five of six individuals from assessments of incremental features and
elemental registration of serially forming molars (detailed in Smith, 2016; Smith et al., 2017).
We also studied four Sumatran fossil orangutan teeth that were collected more than a
century ago from the Lida Ajer and Sibrambang Caves in the Padang Highlands by Eugene
Dubois (de Vos, 1983). Right and left lower molars from Lida Ajer (11594.12, 11595.105)
show identical trace element patterns in their dentine (Supplementary Figure 7), as well as
similar occlusal fissure patterns and light wear, consistent with their attribution to the same
individual. Two Bornean fossil orangutan teeth from Niah Caves (Malaysia) were also
included in this study. The caves have yielded significant late Pleistocene and early Holocene
human remains since the Harrissons began excavations in the 1950s (Barker et al., 2007).
These lower molars were derived from two different entrances to the cave system, Gan Kira
(grid square Y/F4) and Lobang Angus/Hangus (grid square US/22), with burial depths of 12–
18 inches and 30–36 inches, respectively (Hooijer, 1961). Although Hooijer (1948, 1961)
identified all six of these fossil teeth as M1s, we regard this as tentative, given that isolated
orangutan molars are notoriously difficult to seriate (Grine and Franzen, 1994).
439

Dating of fossil samples
Preliminary assessments at the Australian National University Radiocarbon Dating
Laboratory confirmed that collagen preservation in the six fossil orangutans was insufficient
for radiocarbon dating, a common limitation in tropical environments (e.g., Wood et al.,
2016). Laser ablation uranium series (U-series) analyses were carried out on longitudinal
sections of teeth at the Radiogenic Isotope Facility of the University of Queensland using an
ASI RESOlution SE laser ablation system connected to a Nu Plasma II MC-ICP-MS. A

succession of several rasters (< 2-minute linear ablations) was made in a transect across the

dentine and enamel of each tooth (Supplementary Figures 4-6) following Grün et al. (2014).

The $^{230}Th/^{238}U$ and $^{234}U/^{238}U$ activity ratios of the samples were normalized to bracketing

analyses of a homogeneous rhino tooth standard that has been precisely calibrated by isotope

dilution (Grün et al., 2014). Importantly, dental tissues are known to behave as open systems

for U-series elements; provided there is no occurrence of uranium leaching, age estimates

should therefore be regarded as minimum age constraints since uranium uptake into dental

tissues may be significantly delayed after death.

Tooth formation and oxygen isotope analyses

Thin sections were first imaged with transmitted light microscopy. Enamel daily

secretion rates were measured between sequential accentuated growth lines to yield the time

of formation (see Smith, 2016: Fig. 1, p. 94), and enamel extension rates were calculated

between accentuated lines to guide placement of the analysed spots at approximately weekly

intervals of growth from the dentine horn tip to the enamel cervix (Smith et al., 2018a; Green

et al., 2022). Following the removal of cover slips by immersion in xylene, each thin section

was analysed for $\delta^{18}O$ at the SHRIMP Laboratory at the Australian National University

according to methods detailed in Vaiglova et al. (in review).

In brief, a 15 kV Cs primary ion beam focused to a spot ~ 15 x 20 μm diameter was

used to sequentially sample the enamel as close as possible to the enamel-dentine junction

(EDJ). Oxygen secondary ions were extracted at 10 kV and analysed isotopically by a

multiple collector equipped with dual electrometers operated in resistor mode. The $\delta^{18}O$

values were calculated relative to reference apatite (Durango 3) measured every 10–15

sample analyses. Distances of SHRIMP $\delta^{18}O$ measurements along the innermost enamel from

the cusp to cervix were converted to secretory time in days following Green et al. (2022). A

polynomial regression relating distances to days was created using the enamel extension

rates, and this regression was applied to estimate the timing of secretory deposition at every

SHRIMP spot location. The Lomb–Scargle periodogram was used to assess time-dependent

patterns of $\delta^{18}O$ values, which estimates the power of sine wave periods within a given range

to produce the temporal patterns present within those measurements.

The probability that differences between first and second year $\delta^{18}O$ values in modern

first molars might have arisen by chance was assessed by one-way paired t-tests, with alpha =

0.05 adjusted by Bonferroni correction due to repeated comparisons across multiple teeth.

Figure and data plotting using Python 3 in the Google Colab environment were aided by ChatGPT, a language model based on the GPT-3.5 architecture developed by OpenAI.

Acknowledgements

Manfred Ade, Steffen Bock, Judy Chupasko, Mark Omura, Carina Phillips, and Olav Rohrer-Ertl assisted with access to living orangutan material. Graeme Barker, Ipoi Datan, and Natasja den Ouden assisted with access to fossil orangutan material. The following museums and universities provided access to material: Humboldt Museum (Berlin), Harvard University Museum of Comparative Zoology (Cambridge), Naturalis Museum (Leiden), State Anthropological Collection (Munich), and the Sarawak Museum (Kuching). Technical assistance was provided by Kamil Sokolowski and Brian Tse at the Preclinical Imaging Core Facility at the Translational Research Institute, funding support for which came from Therapeutic Innovation Australia, under the National Collaborative Research Infrastructure Strategy. Kate Britton, Luca Fiorenza, Gathorne Gathorne-Hardy (Earl of Cranbrook), Terry Harrison, Jessica Rippengal, Pam Walton, and Rachel Wood provided research assistance. Sample preparation and oxygen isotope analysis was funded by the Australian Academy of Science Regional Collaborations Programme, the Australian National University, the Australian Research Council (DP210101913), and Griffith University. The U-series dating was funded by the Australian Research Council (FT150100215) and the Spanish Ramón y Cajal Fellowship (RYC2018-025221-I). The latter is funded by MCIN/AEI/ 10.13039/501100011033 and by 'ESF Investing in your future.' U-series dating analyses were carried out within the framework of the existing Brisbane Geochronology Alliance (Griffith University, University of Queensland, and Queensland University of Technology); Y. Feng helped with the U-series data acquisition.

References

Aldrian ED, Susanto R. 2003. Identification of three dominant rainfall regions within Indonesia and their relationship to sea surface temperature. International Journal of Climatology **23**: 1435-1452. DOI: https://doi.org/10.1002/joc.950

Archibald S, Bond WJ, Hoffmann W, Lehmann C, Staver C, Stevens N. 2019. Distribution and determinants of Savannas. *In: Savanna Woody Plants and Large Herbivores*, (eds P.F. Scogings and M. Sankaran). DOI: https://doi.org/10.1002/9781119081111.ch1

517 Ashbury AM, Posa MRC, Dunkel LP, Spillmann B, Atmoko SSU, van Schaik CP, van
518 Noordwijk MA. 2015. Why do orangutans leave the trees? Terrestrial behavior among wild
519 Bornean orangutans (*Pongo pygmaeus wurmbii*) at Tuanan, Central Kalimantan. *American*
520 *Journal of Primatology* **77**: 1216-1229. DOI: https://doi.org/10.1002/ajp.22460
521
522 Aubert M, Williams IS, Boljkovac K, Moffat I, Moncel MH, Dufour E, Grün R. 2012. In situ
523 oxygen isotope micro-analysis of faunal material and human teeth using a SHRIMP II: A new
524 tool for palaeo-ecology and archaeology. *Journal of Archaeological Science* **39**: 3184-3194.
525 DOI: https://doi.org/10.1016/j.jas.2012.05.002
526
Bacon A-M, Bourgon N, Welker F, Cappellini E, Fiorillo D, Tombret O, Huong NTM, Tuan
NA, Sayavonkhamdy T, Souksavatdy V, Sichanthongtip P, Antoine P-O, Duringer P, Ponche
J-L, Westaway K, Joannes-Boyau R, Boesch Q, Suzzoni E, Frangeul S, Patole-Edoumba E,
Zachwieja A, Shackelford L, Demeter F, Hublin J-J, Dufour É. 2021. A multi-proxy
approach to exploring Homo sapiens' arrival, environments and adaptations in Southeast
Asia. *Scientific Reports* **11**: 21080. DOI: https://doi.org/10.1038/s41598-021-99931-4
533
534 Bae CJ, Douka K, Petraglia MD. 2017. On the origin of modern humans: Asian perspectives.
535 *Science* **358**: eaai9067. DOI: https://doi.org/10.1126/science.aai9067
536
Barker G, Barton H, Bird M, Daly P, Datan I, Dykes A, Farr L, Gilbertson D, Harrisson B,
Hunt C, Higham T, Kealhofer L, Krigbaum J, Lewis H, McLaren S, Paz V, Pike A, Piper P,
Pyatt B, Rabett R, Reynolds T, Rose J, Rushworth G, Stephens M, Stringer C, Thompson J,
Turney C. 2007. The 'human revolution' in lowland tropical Southeast Asia: the antiquity and
behavior of anatomically modern humans at Niah Cave (Sarawak, Borneo). *Journal of*
Human Evolution **52**: 243-261. DOI: https://doi.org/10.1016/j.jhevol.2006.08.011
543
544 Belgaman HA, Ichiyanagi K, Tanoue M, Suwarman R. 2016. Observational research on
545 stable isotopes in precipitation over Indonesian maritime continent. *Journal of Japanese*
546 *Association of Hydrological Sciences* **46**: 7–28. DOI: https://10.4145/jahs.46.7
547
548 Belgaman HA, Ichiyanagi K, Suwarman R, Tanoue M, Aldrian E, Utami AID,
549 Kusumaningtyas SDA. 2017. Characteristics of seasonal precipitation isotope variability in
550 Indonesia. *Hydrological Research Letters* **11**: 92-98. DOI: https://doi.org/10.3178/hrl.11.92
551
552 Bird MI, Taylor D, Hunt C. 2005. Palaeoenvironments of insular Southeast Asia during the
553 Last Glacial Period: A savanna corridor in Sundaland? *Quaternary Science Reviews* **24**:
554 2228-2242. DOI: https://doi.org/10.1016/j.quascirev.2005.04.004
555
556 Blumenthal SA, Levin NE, Brown FH, Brugal JP, Chritz KL, Harris JM, Jehle GE, Cerling
557 TE. 2017. Aridity and hominin environments. *Proceedings of the National Academy of*
558 *Sciences of the United States of America* **114**: 7331-7336. DOI:
559 https://doi.org/10.1073/pnas.1700597114
560
561 Bowen GJ. 2010. Isoscapes: Spatial pattern in isotopic biogeochemistry. *Annual Review of*
562 *Earth and Planetary Sciences* **38**: 161-187. DOI: https://doi.org/10.1146/annurev-earth-
563 040809-152429
564

565 Britton K, Fuller BT, Tütken T, Mays S, Richards MP. 2015. Oxygen isotope analysis of
566 human bone phosphate evidences weaning age in archaeological populations. *American*
567 *Journal of Physical Anthropology* **157**: 226-241. DOI: https://doi.org/10.1002/ajpa.22704
568
569 Brockman D, vain Schaik C. (Eds.). 2005. *Seasonality in Primates: Studies of Living and*
570 *Extinct Human and Non-Human Primates* (Cambridge Studies in Biological and
571 Evolutionary Anthropology). Cambridge: Cambridge University Press. DOI:
572 https://doi.org/10.1017/CBO9780511542343
573
574 Bryant JD, Froelich PN, Showers WJ, Genna BJ. 1996. A tale of two quarries: Biologic and
575 taphonomic signatures in the oxygen isotope composition of tooth enamel phosphate from
576 modern and miocene equids. *Palaios* **11**: 397-408. DOI: https://doi.org/10.2307/3515249
577
578 Buckingham FL, Carolin SA, Partin JW, Adkins JF, Cobb KM, Day CC, Ding Q, He C, Liu
579 Z, Otto-Bliesner B, Roberts WHG, Lejau S, Malang J. 2022. Termination 1 millennial-scale
580 rainfall events over the Sunda Shelf. *Geophysical Research Letters* **49**: e2021GL096937.
581 DOI: https://doi.org/10.1029/2021GL096937
582
583 Cherney MD, Fisher DC, Hren MT, Shirley EA. 2021. Stable isotope records of nursing and
584 weaning: A case study in elephants with implications for paleobiological investigations.
585 *Palaeogeography, Palaeoclimatology, Palaeoecology* **567**: 110223. DOI:
586 https://doi.org/10.1016/j.palaeo.2021.110223
587
588 Chesson LA, Valenzuela LO, O'Grady SP, Cerling TE, Ehleringer JR. 2010. Hydrogen and
589 oxygen stable isotope ratios of milk in the United States. *Journal of Agricultural and Food*
590 *Chemistry* **58**: 2358-2363. DOI: https://doi.org/10.1021/jf904151c
591
592 Cole JE, Fairbanks RG, Shen GT. 1993. Recent variability in the southern oscillation:
593 Isotopic results from a Tarawa Atoll coral. *Science* **260**: 1790-1793. DOI:
594 https://doi.org/10.1126/science.260.5115.1790
595
596 Curran LM, Caniago I, Paoli GD, Astianti D, Kusneti M, Leighton M, Nirarita CE,
597 Haeruman H. 1999. Impact of El Niño and logging on canopy tree recruitment in Borneo.
598 *Science* **286**: 2184-2188. DOI: https://doi.org/10.1126/science.286.5447.2184
599
600 Dansgaard W. 1964. Stable isotopes in precipitation. *Tellus* **16**: 436-468. DOI:
601 https://doi.org/10.1111/j.2153-3490.1964.tb00181.x
602
603 de Vos J. 1983. The *Pongo* faunas from Java and Sumatra and their significance for
604 biostratigraphical and paleo-ecological interpretations. *Palaeontology Proceedings B*. **86**:
605 417-425.
606
607 da Silveira L, Sternberg L, Mulkey SS, Joseph Wright S. 1989. Oxygen isotope ratio
608 stratification in a tropical moist forest. *Oecologia* **81**: 51-56. DOI:
609 https://doi.org/10.1007/BF00377009
610
611 Fannin LD, McGraw WS. 2020. Does oxygen stable isotope composition in primates vary as
612 a function of vertical stratification or folivorous behaviour? *Folia Primatologica* **91**: 219–
613 227.
614

Fricke HC, O'Neil JR. 1996. Inter- and intra-tooth variation in the oxygen isotope
composition of mammalian tooth enamel phosphate: Implications for palaeoclimatological
and palaeobiological research. *Palaeogeography, Palaeoclimatology, Palaeoecology* **126**: 91-
99. https://doi.org/10.1016/S0031-0182(96)00072-7

Green DR, Ávila JN, Cote S, Dirks W, Lee D, Poulsen CJ, Williams IS, Smith TM. 2022.
Fine-scaled climate variation in equatorial Africa revealed by modern and fossil primate
teeth. *Proceedings of the National Academy of Sciences of the United States of America* **119**:
e212336611. DOI: https://doi.org/10.1073/pnas.2123366119

Green DR, Olack G, Colman AS. 2018. Determinants of blood water δ18O variation in a
population of experimental sheep: Implications for paleoclimate reconstruction. *Chemical*
Geology **485**: 32-43. DOI: https://doi.org/10.1016/j.chemgeo.2018.03.034

Grine F, Franzen JL. 1994. Fossil hominid teeth from the Sangiran Dome (Java, Indonesia).
Courier Forschungsinstitut Senckenberg **171**: 75–103.

Griffiths ML, Drysdale RN, Gagan MK, Zhao J-X, Ayliffe LK, Hellstrom JC, Hantoro WS,
Frisia S, Feng Y-X, Cartwright I, Pierre ES, Fischer MJ, Suwargadi BW. 2009. Increasing
Australian-Indonesian monsoon rainfall linked to early Holocene sea-level rise. *Nature*
Geoscience **2**: 636-639. DOI: https://doi.org/10.1038/ngeo605

Grün R, Eggins S, Kinsley L, Moseley H, Sambridge M. 2014. Laser ablation U-series
analysis of fossil bones and teeth. *Palaeogeography, Palaeoclimatology, Palaeoecology* **416**:
150-167. DOI: https://doi.org/10.1016/j.palaeo.2014.07.023

Harrison T. 1996. The paleoecological context at Niah Cave Sarawak: Evidence from the
primate fauna. *Bulletin of the Indo-Pacific Prehistory Association* **14**, 90–100.

Hooijer DA. 1948. Prehistoric teeth of man and of the orang-utan from central Sumatra, with
notes on the fossil orang-utan from Java and Southern China. *Zoologische Mededelingen* **29**:
175–301.

Hooijer DA. 1961. The orang-utan in Niah Cave prehistory. *Sarawak Museum Journal* **9**:
408-421.

Hughen KA, Schrag DP, Jacobsen SB, Hantoro W. 1999. El Niño during the last interglacial
period recorded by a fossil coral from Indonesia. *Geophysical Research Letters* **26**: 3129-
3132. DOI: https://doi.org/10.1029/1999GL006062

Hunt CO, Gilbertson DD, Rushworth G. 2012. A 50,000-year record of late Pleistocene
tropical vegetation and human impact in lowland Borneo. *Quaternary Science Reviews* **37**:
61-80. DOI: https://doi.org/10.1016/j.quascirev.2012.01.014

Jablonski NG, Whitfort MJ, Roberts-Smith N, Qinqi X. 2000. The influence of life history
and diet on the distribution of catarrhine primates during the Pleistocene in eastern Asia.
Journal of Human Evolution **39**: 131-157. DOI: https://doi.org/10.1006/jhev.2000.0405

Janssen R, Joordens JCA, Koutamanis DS, Puspaningrum MR, de Vos J, van der Lubbe
JHJL, Reijmer JJG, Hampe O, Vonhof HB. 2016. Tooth enamel stable isotopes of Holocene

and Pleistocene fossil fauna reveal glacial and interglacial paleoenvironments of hominins in Indonesia. *Quaternary Science Reviews* **144**: 145-154. DOI: https://doi.org/10.1016/j.quascirev.2016.02.028

Knott CD. 1998. Changes in orangutan caloric intake, energy balance, and ketones in response to fluctuating fruit availability. *International Journal of Primatology* **19**: 1061-1079. DOI: https://doi.org/10.1023/A:1020330404983

Kornexl BE, Werner T, Roßmann A, Schmidt H-L. 1997. Measurement of stable isotope abundances in milk and milk ingredients - A possible tool for origin assignment and quality control. *Zeitschrift fur Lebensmittel -Untersuchung und -Forschung* **205**: 19-24. DOI: https://doi.org/10.1007/s002170050117

Krigbaum J, Berger MH, Daegling DJ, McGraw WS. 2013. Stable isotope canopy effects for sympatric monkeys at Taï Forest, Côte d'Ivoire. *Biology Letters* **9**: 20130466.

Kubat J, Nava A, Bondioli L, Dean MC, Zanolli C, Bourgon N, Bacon A-M, Demeter F, Peripoli B, Albert R, Lüdecke T, Hertler C, Mahoney P, Kullmer O, Schrenk F, Müller W. 2023. Dietary strategies of Pleistocene *Pongo sp.* and *Homo erectus* on Java (Indonesia). *Nature Ecology and Evolution* **7**: 279-289. DOI: https://doi.org/10.1038/s41559-022-01947-0

Lin GP, Rau YH, Chen YF, Chou CC, Fu WG. 2006. Measurements of δD and δ^{18}O stable isotope ratios in milk. *Journal of Food Science* **68**: 2192-2195. DOI: https://doi.org/10.1111/j.1365-2621.2003.tb05745.x

Louys J, Duval M, Price GJ, Westaway K, Zaim Y, Rizal Y, Aswan A, Puspaningrum M, Trihascaryo A, Breitenbach SFM, Kwiecien O, Cai Y, Higgins P, Albers PCH, de Vos J, Roberts P. 2022. Speleological and environmental history of Lida Ajer cave, western Sumatra. *Philosophical Transactions of the Royal Society B: Biological Sciences* **377**: 20200494. DOI: https://doi.org/10.1098/rstb.2020.0494

Louys J, Meijaard E. 2010. Palaeoecology of Southeast Asian megafauna-bearing sites from the Pleistocene and a review of environmental changes in the region. *Journal of Biogeography* **37**: 1432-1449. DOI: https://doi.org/10.1111/j.1365-2699.2010.02297.x

Louys J, Price GJ, Higgins P, de Vos J, Zaim J, Rizal Y, Aswan, Puspaningrum MR, Tri Hascaryo A, Drawhorn GM, Albers PCH. in press. Geochronology and palaeoenvironments of Sibrambang and Djambu caves, western Sumatra, in Louys J, Albers PCH, van der Geer AAE (eds), *Quaternary Palaeontology and Archaeology of Sumatra*. (Terra Australis, Australian National University Press, Canberra)

Louys J, Roberts P. 2020. Environmental drivers of megafauna and hominin extinction in Southeast Asia. *Nature* **586**: 402-406. DOI: https://doi.org/10.1038/s41586-020-2810-y

Lowry BE, Wittig RM, Pittermann J, Oelze VM. 2021. Stratigraphy of stable isotope ratios and leaf structure within an African rainforest canopy with implications for primate isotope ecology. *Scientific Reports* **11**:14222. DOI: https://doi.org/10.1038/s41598-021-93589-8

Luyt J, Sealy J. 2018. Inter-tooth comparison of δ^{13}C and δ^{18}O in ungulate tooth enamel

from south-western Africa. *Quaternary International* **495**:144-152. DOI: https://doi.org/10.1016/j.quaint.2018.02.009

MacKinnon J. 1974. The behaviour and ecology of wild orang-utans (*Pongo pygmaeus*). *Animal Behaviour* 22: 3-74.

Marshall AJ, Ancrenaz M, Brearley FQ, Fredriksson GM, Ghaffar N, Heydon M, Husson SJ, Leighton M, McConkey KR, Morrogh-Bernard HC, Proctor J, van Schaik CP, Yeager CP, Wich SA. 2009. The effects of forest phenology and floristics on populations of Bornean and Sumatran orangutans: Are Sumatran forests better orangutan habitat than Bornean forests?, in Wich SA, Utami Atmoko SS, Setia TM, van Schaik CP (eds), *Orangutans: Geographic Variation in Behavioral Ecology and Conservation* (Oxford, 2008; online edn, Oxford Academic, 1 May 2009) 97-117. DOI: https://doi.org/10.1093/acprof:oso/9780199213276.003.0007,

McPhaden MJ, Zebiak SE, Glantz MH. 2006. ENSO as an integrating concept in earth science. *Science* **314**: 1740-1745. DOI: https://doi.org/10.1126/science.1132588

Moerman JW, Cobb KM, Adkins JF, Sodemann H, Clark B, Tuen AA. 2013. Diurnal to interannual rainfall δ^{18}O variations in northern Borneo driven by regional hydrology. *Earth and Planetary Science Letters* **369-370**: 108-119. DOI: https://doi.org/10.1016/j.epsl.2013.03.014

Moron V, Robertson AW, Boer R. 2009. Spatial coherence and seasonal predictability of monsoon onset over Indonesia. *Journal of Climate* **22**: 840-850. DOI: https://doi.org/10.1175/2008JCLI2435.1

Negash EW, Barr WA. 2023. Relative abundance of grazing and browsing herbivores is not a direct reflection of vegetation structure: Implications for hominin paleoenvironmental reconstruction. *Journal of Human Evolution* **177**: 103328. DOI: https://doi.org/10.1016/j.jhevol.2023.103328

Nelson SV. 2013. Chimpanzee fauna isotopes provide new interpretations of fossil ape and hominin ecologies. *Proceedings of the Royal Society B* **280**: 20132324. DOI: https://doi.org/10.1098/rspb.2013.2324

Pederzani S, Britton K. 2019. Oxygen isotopes in bioarchaeology: Principles and applications, challenges and opportunities. *Earth-Science Reviews* **188**: 77-107. DOI: https://doi.org/10.1016/j.earscirev.2018.11.005

Piper PJ. 2016. Human cultural, technological and adaptive changes from the end of the Pleistocene to the mid-Holocene in Southeast Asia. in Oxenham, M. & Buckley, H. R. (eds.) *The Routledge Handbook of Bioarchaeology in Southeast Asia and the Pacific Islands*, Routledge, 24-44.

Piper P, Lewis H, Reynolds T, McLaren S, Rabett R, Cole F, Szábo K, Farr L, Barker G. 2016. The NCP Excavations in Lobang Hangus, Lobang Tulang, Gan Kira, and Kain Hitam, in G. Barker, L. Farr (eds.), *Archaeological Investigations in the Niah Caves, Sarawak: The Archaeology of the Niah Caves, Sarawak*, **2**: 65-80. Cambridge, United Kingdom: McDonald Institute for Archaeological Research.

765
766 Piper PJ, Lim TT. 2021. Zooarchaeology at Niah Cave: contributions to our understanding of
767 southeast Asian prehistory. Malayan Nat J 81[st] Anniversary Special Issue, 195-208.
768
769 Piper P, Rabett R. 2016. Vertebrate Fauna from the Niah Caves, in G. Barker, L. Farr (eds.),
770 *Archaeological Investigations in the Niah Caves, Sarawak: The Archaeology of the Niah*
771 *Caves, Sarawak*, **2**: 401-438. Cambridge, United Kingdom: McDonald Institute for
772 Archaeological Research
773
774 Pletcher E, Staver C, Schwartz NB. 2022. The environmental drivers of tree cover and forest–
775 savanna mosaics in Southeast Asia. *Ecography* **2022**: e06280. DOI:
776 https://doi.org/10.1111/ecog.06280
777
778 Prendergast AL, Pryor AJE, Reade H, Stevens RE. 2018. Seasonal records of
779 palaeoenvironmental change and resource use from archaeological assemblages. *Journal of*
780 *Archaeological Science: Reports* **21**: 1191–1197. DOI:
781 https://doi.org/10.1016/j.jasrep.2018.08.006
782
783 Pumijumnong N, Bräuning A, Sano M, Nakatsuka T, Muangsong C, Buajan S. 2020. A 338-
784 year tree-ring oxygen isotope record from Thai teak captures the variations in the Asian
785 summer monsoon system. *Scientific Reports* **10**: 8966. DOI: https://doi.org/10.1038/s41598-
786 020-66001-0
787
788 Qian J-H, Robertson AW, Moron V. 2013. Diurnal cycle in different weather regimes and
789 rainfall variability over Borneo associated with ENSO. *Journal of Climate* **26**: 1772-1790.
790 DOI: https://doi.org/10.1175/JCLI-D-12-00178.1
791
792 Roberts SB, Coward WA, Ewing G, Savage J, Cole TJ, Lucas, A. 1988. Effect of weaning on
793 accuracy of doubly labled water method in infants. *American Journal of Physiology-*
794 *Regulatory, Integrative and Comparative Physiology* **254**: R622-R627. DOI:
795 https://doi.org/10.1152/ajpregu.1988.254.4.R622
796
797 Roberts P, Blumenthal SA, Dittus W, Wedage O, Lee-Thorp JA. 2017. Stable carbon,
798 oxygen, and nitrogen, isotope analysis of plants from a South Asian tropical forest:
799 Implications for primatology. *American Journal of Primatology* **79**: e22656. DOI:
800 https://doi.org/10.1002/ajp.22656
801
802 Roberts P, Louys J, Zech J, Shipton C, Kealy S, Carro SS, Hawkins S, Boulanger C, Marzo
803 S, Fiedler B, Boivin N, Mahirta, Aplin K, O'Connor S. 2020. Isotopic evidence for initial
804 coastal colonization and subsequent diversification in the human occupation of Wallacea.
805 *Nature Communications* **11**: 2068. DOI: https://doi.org/10.1038/s41467-020-15969-4
806
807 Röhrer-Ertl O. 1988 Research history, nomenclature, and taxonomy of the orang-utan, in
808 Schwartz JH (ed) *Orang-utan Biology*, Oxford University Press, New York, 7-18.
809
810 Rozanski K, Araguás-Araguás L, Gonfiantini R. 1993. Isotopic patterns in modern global
811 precipitation, in Swart PK, Lohmann KC, Mckenzie J, Savin S (eds), Climate change in
812 continental isotopic records. American Geophysical Union, Washington D.C., USA; 1–36.
813 DOI: https://doi.org/10.1029/GM078p0001
814

815   Smith TM. 2016. Dental development in living and fossil orangutans. *Journal of Human*
816   *Evolution* **94**: 92-105. DOI: https://doi.org/10.1016/j.jhevol.2016.02.008
817
818   Smith TM, Austin C, Ávila JN, Dirks W, Green DR, Williams IS, Arora M. 2022. Permanent
819   signatures of birth and nursing initiation are chemically recorded in teeth. *Journal of*
820   *Archaeological Science* **140**: 105564. DOI: https://doi.org/10.1016/j.jas.2022.105564
821
822   Smith TM, Austin C, Green DR, Joannes-Boyau R, Bailey S, Dumitriu D, Fallon S, Grün R,
823   James HF, Moncel MH, Williams IS, Wood R, Arora M. 2018a. Wintertime stress, nursing,
824   and lead exposure in Neanderthal children. *Science Advances* **4**: eaau9483. DOI:
825   https://doi.org/10.1126/sciadv.aau9483
826
827   Smith TM, Austin C, Hinde K, Vogel ER, Arora M. 2017. Cyclical nursing patterns in wild
828   orangutans. *Science Advances* **3**: e1601517. DOI: https://doi.org/10.1126/sciadv.1601517
829
830   Smith, TM., Bacon A-M, Demeter F, Kullmer O, Nguyen KT, de Vos J, Wei W, Zermeno JP,
831   Zhao L. 2011. Dental tissue proportions in fossil orangutans from mainland Asia and
832   Indonesia. *Human Origins Research* **1**: e1-e6.
833
834   Smith TM, Houssaye A, Kullmer O, Le Cabec A, Olejniczak AJ, Schrenk F, de Vos J,
835   Tafforeau P. 2018b. Disentangling isolated dental remains of asian pleistocene hominins and
836   pongines. *PLoS ONE* **13**: e0204737. DOI: https://doi.org/10.1371/journal.pone.0204737
837
838   Smith TM, Kupczik K, MacHanda Z, Skinner MM, Zermeno JP. 2012. Enamel thickness in
839   Bornean and Sumatran orangutan dentitions. *American Journal of Physical Anthropology*
840   **147**: 417-426. DOI: https://doi.org/10.1002/ajpa.22009
841
842   Sokolowski KG, Codding BF, Duc A, Faith JT (2023) Do grazers equal grasslands?
843   Strengthening paleoenvironmental inferences through analysis of present-day African
844   mammals. *Palaeogeography, Palaeoclimatology, Palaeoecology* **629**: 111786. DOI:
845   https://doi.org/10.1016/j.palaeo.2023.111786
846
847   Spehar SN, Sheil D, Harrison T, Louys J, Ancrenaz M, Marshall AJ, Wich SA, Bruford MW,
848   Meijaard E. 2018. Orangutans venture out of the rainforest and into the anthropocene.
849   *Science Advances* **4**: e1701422. DOI: https://doi.org/10.1126/sciadv.1701422
850
851   Stahle DS, D'Arrigo RD, Krusic PJ, Cleaveland MK, Cook ER, Allan RJ, Cole JE, Dunbar
852   RB, Therrell MD, Gay DA, Moore MD, Stokes MA, Burns BT, Villanueva-Diaz J,
853   Thompson LG. 1998. Experimental Dendroclimatic Reconstruction of the Southern
854   Oscillation. *Bulletin of the American Meteorological Society* **79**: 2137-2152. DOI:
855   https://doi.org/10.1175/1520-0477(1998)079<2137:EDROTS>2.0.CO;2
856
857   Stephens M, Rose J, Mattey D, Gilbertson D. D. 2016. Stable isotope analysis of shells from
858   the West Mouth: palaeoenvironments, seasonality and harvesting, in G. Barker, L. Farr (eds.),
859   *Archaeological Investigations in the Niah Caves, Sarawak: The Archaeology of the Niah*
860   *Caves, Sarawak*, **2**: 177-200. Cambridge, United Kingdom: McDonald Institute for
861   Archaeological Research.
862
863   Suwarman R, Ichiyanagi K, Tanoue M, Yoshimura K, Mori S, Yamanaka MD, Kurita N,
864   Syamsudin F. 2013. The variability of stable isotopes and water origin of precipitation over

the maritime continent. *Scientific Online Letters on the Atmosphere* **9**: 74-78. DOI: https://doi.org/10.2151/sola.2013-017

Thorpe SKS, Crompton RH. 2005. Locomotor ecology of wild orangutans (*Pongo pygmaeus abelii*) in the Gunung Leuser Ecosystem, Sumatra, Indonesia: a multivariate analysis using log-linear modeling. *American Journal of Physical Anthropology* **127**: 58–78.

Thorpe SKS, Crompton RH. 2009. Orangutan positional behavior: Interspecific variation and ecological correlates, in Wich SA, Utami Atmoko SS, Setia TM, van Schaik CP (eds), *Orangutans: Geographic Variation in Behavioral Ecology and Conservation* (Oxford, 2008; online edn, Oxford Academic, 1 May 2009) 33-48. DOI: https://doi.org/10.1093/acprof:oso/9780199213276.003.0003

Tshen LT. 2016. Biogeographic distribution and metric dental variation of fossil and living orangutans (*Pongo spp.*). *Primates* **57**: 39-50. DOI: https://doi.org/10.1007/s10329-015-0493-z

Tudhope AW, Chilcott CP, McCulloch MT, Cook ER, Chappell J, Ellam RM, Lea DW, Lough JM, Shimmield GB. 2001. Variability in the El Niño-Southern Oscillation through a glacial-interglacial cycle. *Science* **291**: 1511-1517. DOI: https://doi.org/10.1126/science.1057969

Ungar P. 1996. Feeding height and niche separation in sympatric Sumatran monkeys and apes. *Folia Primatologica* **67**: 163-168.

Urban FE, Cole JE, Overpeck JT. 2000. Influence of mean climate change on climate variability from a 155-year tropical Pacific coral record. *Nature* **407**: 989-993. DOI: https://doi.org/10.1038/35039597

Vaiglova P, Ávila JN, Buckley H, Galipaud JC, Green DR, Halcrow S, James HF, Kinaston R, Oxenham M, Paz V, Simanjukntak T, Snoeck C, Trinh HH, Williams IS, Smith TM. in review. Holocene rainfall patterns in Southeast Asia revealed by microanalysis of δ^{18}O values in human teeth. *Journal of Archaeological Science.* Preprint: http://dx.doi.org/10.2139/ssrn.4542042

van Noordwijk MA, Willems EP, Utami Atmoko SS, Kuzawa CW, van Schaik CP. 2013. Multi-year lactation and its consequences in Bornean orangutans (*Pongo pygmaeus wurmbii*). *Behavioral Ecology and Sociobiology* **67**: 805-814. DOI: https://doi.org/10.1007/s00265-013-1504-y

Van Schaik CP. 1986. Phenological changes in a Sumatran rain forest. *Journal of Tropical Ecology* **2**: 327-347. DOI: https://doi.org/10.1017/S0266467400000973

Vico G, Dralle D, Feng X, Thompson S, Manzoni S. 2017. How competitive is drought deciduousness in tropical forests? A combined eco-hydrological and eco-evolutionary approach. *Environmental Research Letters* **12**: 065006. DOI: https://doi.org/10.1088/1748-9326/aa6f1b

913   Wessling EG, Kühl HS, Mundry R, Deschner T, Pruetz JD. 2018. The costs of living at the
914   edge: Seasonal stress in wild savanna-dwelling chimpanzees. *Journal of Human Evolution*
915   **121**: 1-11. DOI: https://doi.org/10.1016/j.jhevol.2018.03.001
916
917   Westaway KE, Louys J, Awe RD, Morwood MJ, Price GJ, Zhao J-X, Aubert M, Joannes-
918   Boyau R, Smith TM, Skinner MM, Compton T, Bailey RM, Van Den Bergh GD, de Vos J,
919   Pike AWG, Stringer C, Saptomo EW, Rizal Y, Zaim J, Santoso WD, Trihascaryo A, Kinsley
920   L, Sulistyanto B. 2017. An early modern human presence in Sumatra 73,000-63,000 years
921   ago. *Nature* **548**: 322-325. DOI: https://doi.org/10.1038/nature23452
922
923   Wood R, Duval M, Mai Huong NT, Tuan NA, Bacon AM, Demeter F, Duringer P, Oxenham
924   M, Piper P. 2016. The effect of grain size on carbonate contaminant removal from tooth
925   enamel: Towards an improved pretreatment for radiocarbon dating. *Quaternary*
926   *Geochronology* **36**: 174-187. DOI: https://doi.org/10.1016/j.quageo.2016.08.010
927
928   Wright LE, Schwarcz HP. 1999. Correspondence between stable carbon, oxygen and nitrogen
929   isotopes in human tooth enamel and dentine: Infant diets at Kaminaljuyu. *Journal of*
930   *Archaeological Science* **26**: 1159-1170. DOI: https://doi.org/10.1006/jasc.1998.0351
931
932   Wurster CM, Bird MI, Bull ID, Creed F, Bryant C, Dungait JAJ, Paz V. 2010. Forest
933   contraction in north equatorial Southeast Asia during the Last Glacial Period. *Proceedings of*
934   *the National Academy of Sciences of the United States of America* **107**: 15508-15511. DOI:
935   https://doi.org/10.1073/pnas.1005507107
936
937   Yang H, Johnson KR, Griffiths ML, Yoshimura K. 2016. Interannual controls on oxygen
938   isotope variability in Asian monsoon precipitation and implications for paleoclimate
939   reconstructions. *Journal of Geophysical Research* **121**: 8410-8428. DOI:
940   https://doi.org/10.1002/2015JD024683
941

942   **Figures**
943
944   Figure 1. Approximate location of select modern and fossil orangutans superimposed on
945   modeled isotopic variation.
946

947
948
949
950   Figure modified from www.waterisotopes.org based on data from the Online Isotopes in
951   Precipitation Calculator (3.0). See Table 1 for the location of particular individuals.
952   Sibrambang Cave has yet to be relocated since Eugene Dubois' original excavations, but it is
953   known to be in the general vicinity of Lida Ajer in the Padang Highlands, possibly near to the
954   modern village of a similar name (Louys et al. in press).
955

956 Figure 2. Comparison of sequential $\delta^{18}O$ values across multiple years of first molar formation
957 in five modern orangutans from Borneo and Sumatra.
958

959
960
961 Bornean individuals: MCZ 5290, ZSM 1981/48 ZSM 1981/87; Sumatran individuals: ZMB
962 83508, ZSM 1981/248. The width of each curve is a kernel density estimate (KDE)
963 corresponding to the distribution of $\delta^{18}O$ values. First year data (Y1) is shown with a purple
964 violin plot, second year data (Y2) with a green plot, and third year data (Y3) with a yellow
965 plot where complete/available. Actual data are plotted as black circles.
966

967 Figure 3. Comparison of sequential δ¹⁸O values across multiple years of serial molar
968 formation in two modern orangutans from Borneo (top) and two from Sumatra (bottom).
969

970
971
972
973
974 Individual in upper left: ZSM 1981/48; upper right: ZSM 1981/87; lower left: ZSM
975 1981/246; lower right: ZSM 1981/248. Developmental overlap was determined through
976 registration of trace elements as in Smith et al. (2017).

982   Figure 4. Comparison of $\delta^{18}O$ values in fossil and modern orangutans from Borneo (blue) and
983   Sumatra (green).
984

985
986   Violin plots show kernel density estimates representing the distribution of $\delta^{18}O$ values in
987   modern individuals (left plot), and in fossil individuals (right plot). Actual $\delta^{18}O$
988   measurements are shown as black circles.
989

990   Table 1. Modern and fossil orangutan teeth employed in the current study.
991

Taxon	Accession	Origin	Sex	Age (Years)	Teeth
Pongo pygmaeus	ZSM 1981/48	Skalau, Borneo	F	~8.4	RUM1, LLM2
	ZSM 1981/87	Skalau, Borneo	F	> 9	LUM1, RUM2, RLM3
	MCZ 5290	Borneo (location unspecified)	n/a	4.5	RUM1
Pongo abelii	ZSM 1981/246	Aceh, Sumatra	M	~8.5	LLM1, LUM2
	ZSM 1981/248	Aceh, Sumatra	F	adult	LUM1, LUM2, LLM3
	ZMB 83508	Sumatra (location unspecified)	n/a	8.8	RLM1
Fossil *Pongo* spp.	11564.5	Sibrambang, Sumatra	n/a	n/a	RUM
	11565.162	Sibrambang, Sumatra	n/a	n/a	LUM
	11594.12	Lida Ajer, Sumatra	n/a	n/a	RLM
	11595.105	Lida Ajer, Sumatra	n/a	n/a	LLM
	US/22	Niah Caves, Malaysia	n/a	n/a	RLM
	Y/F4	Niah Caves, Malaysia	n/a	n/a	LLM

992
993
994
995   Numerous taxonomic assignments have been made for fossil orangutans (*Pongo* spp.), some
996   of which have not been based on clear morphological characteristics (Lim, 2016), and are not
997   relevant for the focus of this paper.
998

999   Table 2. Modern and fossil orangutan molar δ^{18}O values.
1000

Taxon	Accession	Tooth	Cusp	Spots	Time (Days)	dO18 Range
Pongo pygmaeus	ZSM 1981/48	RUM1	dl	151	1241	13.6-19.9
	ZSM 1981/48	LLM2	mb	107	804	13.0-18.8
	ZSM 1981/87	LUM1	ml	131	869	13.7-18.2
	ZSM 1981/87	RUM2	ml	196	1195	12.7-20.0
	ZSM 1981/87	RLM3	mb	220	1350	13.7-19.2
	MCZ 5290	RUM1	ml	150	1002	13.8-18.1
Pongo abelii	ZSM 1981/246	LLM1	mb	136	1425	12.3-18.3
	ZSM 1981/246	LUM2	ml	229	1376	12.6-18.0
	ZSM 1981/248	LUM1	db	177	1072	11.3-19.3
	ZSM 1981/248	LUM2	db	193	1374	13.5-20.6
	ZSM 1981/248	LLM3	db	191	1461	15.2-21.2
	ZMB 83508	RLM1	db	135	1029	13.4-20.4
Fossil *Pongo* spp.	11564.5	RUM	mb	178	1387	15.3-20.4
	11565.162	LUM	ml	143	1144	14.7-20.8
	11594.12	RLM	ml	154	1081	15.1-19.9
	11595.105	LLM	mb	197	1312	15.7-20.0
	US/22	RLM	mb	149	1023	15.9-24.8
	Y/F4	LLM	db	134	869	14.2-22.9

1001
1002

1003 Supplementary Figure 1. Comparison of sequential $\delta^{18}O$ values across multiple years of
1004 molar formation in six putative fossil orangutan M1s from Borneo and Sumatra.
1005

1006
1007
The width of each curve is a kernel density estimate (KDE) corresponding to the distribution
of oxygen isotope values measured from different teeth. From each tooth, first year data (Y1)
is shown with a purple violin plot, second year data (Y2) with a green plot, and third year
data (Y3, if present) with a yellow plot. Actual data are plotted as black circles.

Supplementary Figure 2. Inferred seasonality of δ^{18}O values from a Bornean (top) and
Sumatran (bottom) M1.

1014
1015
Lomb-Scargle periodograms show potential periods in days (x-axis) against period power (y-
axis), where higher values on the y-axis indicate underlying sine-wave periods that produce,
contribute to, or explain δ^{18}O value oscillations. Best-fit periodicities are shown as light gray
vertical lines, whereas annual periodicities are indicated by blue dashed horizontal lines. We
regard the Bornean M1 (MCZ 5290) as largely aperiodic; a minor peak is observed at c. 6
months, and increasing powers at very high periods are an artifact of limited sampling length
within teeth relative to the model. The Sumatran M1 (ZMB 83508) has a 1.1 year inferred
frequency, as well as an approximately 6-month peak, likely reflecting semiannual monsoons.
1024

1025 Supplementary Figure 3. Comparison of δ¹⁸O values in fossil and modern orangutans from
1026 Borneo and Sumatra.

1027
1028

1029 Supplementary Figure 4. Laser ablation profiles performed across the two teeth from Lida
1030 Ajer.

1031

1032 #576 (left) refers to specimen 11595.105; #578 (right) refers to specimen 11594.12. The red
1033 dots represent the position of the rasters, and arrows indicate the sequence of the analyses.
1034 EN = enamel. DE = dentine.

1035

1036   Supplementary Figure 5. Laser ablation profiles across the two teeth from Sibrambang Cave.

1037

1038   #577 (left) refers to specimen 11565.162; #581 (right) refers to specimen 11564.5. The red
1039   dots represent the position of the rasters, and arrows indicate the sequence of the analyses.
1040   EN = enamel. DE = dentine.

1041

1042   Supplementary Figure 6. Laser ablation profiles across the two teeth from Niah Caves.

1043

1044   #579 (left) refers to specimen from grid Y/F4; #580 (right) refers to specimen from grid
1045   US/22. The red dots represent the position of the rasters, and arrows indicate the sequence of
1046   the analyses. EN = enamel. DE = dentine.

1047
1048

1049 Supplementary Figure 7. Matching trace element patterns in cross-sections of two isolated
1050 molars from the Dubois collection of fossil orangutan teeth from Lida Ajer.
1051

1052
1053
1054 High concentrations are shown in warm colors, low concentration are in cool colors; here
1055 Li/Ca is shown, but identical corresponding patterns were also observed for Ba/Ca and Sr/Ca
1056 (not shown). The enamel cap of each tooth is to the left, and root dentine is to the right. Trace
1057 elements were measured according to LA-ICP-MS methods detailed in Smith et al. (2017).
1058

1059   Supplementary Table 1. Comparisons of first and second year δ^{18}O values in five first molars.
1060

Specimen	Adjusted p-values	Higher δ^{18}O values
MCZ 5290	p = 0.010	Year 1
ZMB 83508	p = 0.006	Year 1
ZSM 1981/48	p = 0.161 (N.S.)	Year 1
ZSM 1981/87	p < 0.001	Year 2
ZSM 1981/248	p < 0.001	Year 1

1061
1062

1063   Supplementary Table 2. U-series dates for fossil orangutan material.

1064

LA raster	Tissue	U(ppm)	^{230}Th/^{238}U	±	^{234}U/^{238}U	±	Age (ka)	±
11595.105-EN11	enamel	*0.02*	*-1.230*	*0.090*	*0.459*	*0.037*	n.c.	n.c.
11595.105-EN12	enamel	*0.00*	*-12.82*	*0.084*	*-4.590*	*0.018*	n.c.	n.c.
11595.105-EN13	enamel	*0.00*	*-31.04*	*0.082*	*-17.11*	*0.018*	n.c.	n.c.
11595.105-EN14	enamel	*0.10*	*0.623*	*0.168*	*1.155*	*0.105*	n.c.	n.c.
11595.105-EN15	enamel	*0.04*	*-0.826*	*0.082*	*0.752*	*0.051*	n.c.	n.c.
11595.105-DE1	dentine	*41.4*	*0.516*	*0.020*	*1.360*	*0.047*	50.6	3.40
11595.105-DE2	dentine	*44.2*	*0.470*	*0.020*	*1.359*	*0.055*	45.2	3.33
11595.105-DE3	dentine	*48.6*	*0.471*	*0.015*	*1.344*	*0.045*	46.0	2.65
11595.105-DE4	dentine	*58.9*	*0.481*	*0.015*	*1.356*	*0.038*	46.6	2.50
11595.105-DE5	dentine	*61.1*	*0.482*	*0.019*	*1.341*	*0.045*	47.5	3.10
11595.105-DE6	dentine	*65.8*	*0.446*	*0.019*	*1.346*	*0.051*	43.0	3.04
11595.105-DE7	dentine	*64.0*	*0.440*	*0.016*	*1.346*	*0.044*	42.3	2.55
11595.105-DE8	dentine	*60.0*	*0.430*	*0.016*	*1.338*	*0.046*	41.4	2.56
11595.105-DE9	dentine	*64.6*	*0.433*	*0.018*	*1.336*	*0.049*	41.8	2.81
11595.105-DE10	dentine	*65.9*	*0.411*	*0.016*	*1.325*	*0.047*	39.7	2.53

1065

LA raster	Tissue	U(ppm)	^{230}Th/^{238}U	±	^{234}U/^{238}U	±	Age (ka)	±
11594.12-EN1	enamel	*0.00*	*-13.08*	*0.082*	*-9.394*	*0.019*	n.c.	n.c.
11594.12-EN2	enamel	*0.00*	*-23.91*	*0.104*	*-11.05*	*0.016*	n.c.	n.c.
11594.12-EN3	enamel	*0.00*	*-22.35*	*0.078*	*-4.621*	*0.027*	n.c.	n.c.
11594.12-EN4	enamel	*0.00*	*-16.49*	*0.089*	*-6.110*	*0.015*	n.c.	n.c.
11594.12-EN5	enamel	*0.01*	*17.58*	*0.103*	*16.21*	*0.031*	*224*	*3.12*
11594.12-DE6	dentine	30.9	0.337	0.016	1.335	0.041	31.2	2.02
11594.12-DE7	dentine	30.6	0.347	0.013	1.337	0.036	32.2	1.68
11594.12-DE8	dentine	31.9	0.355	0.017	1.332	0.049	33.3	2.31
11594.12-DE9	dentine	31.4	0.366	0.013	1.335	0.035	34.4	1.78
11594.12-DE10	dentine	32.1	0.359	0.011	1.329	0.038	33.8	1.71
11594.12-DE11	dentine	24.6	0.344	0.014	1.357	0.031	31.4	1.67
11594.12-DE12	dentine	24.9	0.352	0.015	1.351	0.032	32.3	1.85

1066

LA raster	tissue	U(ppm)	^{230}Th/^{238}U	±	^{234}U/^{238}U	±	Age (ka)	±
11565.162-EN1	enamel	0.58	0.434	2.630	1.025	1.495	59.7	496
11565.162-EN2	enamel	*0.00*	*-4.899*	*0.132*	*-81.42*	*0.020*	n.c.	n.c.
11565.162-EN3	enamel	*0.00*	*-23.35*	*0.075*	*-16.84*	*0.018*	n.c.	n.c.
11565.162-EN4	enamel	*0.00*	*-37.09*	*0.082*	*-22.74*	*0.021*	n.c.	n.c.
11565.162-EN5	enamel	*0.01*	*-5.283*	*0.068*	*-0.022*	*0.029*	n.c.	n.c.
11565.162-DE6	dentine	62.2	0.438	0.287	1.073	0.470	56.5	59.4
11565.162-DE7	dentine	63.6	0.461	0.023	1.066	0.048	61.2	5.73
11565.162-DE8	dentine	61.4	0.467	0.021	1.072	0.044	61.7	5.23
11565.162-DE9	dentine	59.8	0.470	0.028	1.073	0.060	62.2	7.01
11565.162-DE10	dentine	57.4	0.480	0.026	1.072	0.051	64.2	6.51

LA raster	tissue	U(ppm)	^{230}Th/^{238}U	±	^{234}U/^{238}U	±	Age (ka)	±
11564.5-EN1	enamel	0.11	0.304	0.181	0.959	0.086	41.6	30.6
11564.5-EN2	enamel	*0.00*	*-2215*	*0.086*	*-486.53*	*0.018*	n.c.	n.c.
11564.5-EN3	enamel	*0.00*	*-97.95*	*0.065*	*-4.236*	*0.031*	n.c.	n.c.
11564.5-EN4	enamel	9.86	0.502	0.211	1.115	0.162	64.3	38.8
11564.5-DE5	dentine	51.6	0.564	0.055	1.116	0.052	75.4	11.8
11564.5-DE6	dentine	45.8	0.541	0.057	1.111	0.047	71.6	11.6
11564.5-DE7	dentine	49.1	0.530	0.039	1.117	0.079	69.0	10.2
11564.5-DE8	dentine	47.8	0.512	0.010	1.128	0.016	65.0	2.13

1067

LA raster	tissue	U(ppm)	^{230}Th/^{238}U	±	^{234}U/^{238}U	±	Age (ka)	±
Y/F4-EN1	enamel	*0.01*	*-4.737*	*0.084*	*-0.283*	*0.025*	n.c.	n.c.
Y/F4-EN2	enamel	*0.00*	*-42.98*	*0.070*	*-2.119*	*0.024*	n.c.	n.c.
Y/F4-EN3	enamel	*0.00*	*-137.8*	*0.078*	*-18.50*	*0.017*	n.c.	n.c.
Y/F4-EN4	enamel	*0.00*	*-49.66*	*0.079*	*3.398*	*0.033*	n.c.	n.c.
Y/F4-DE5	dentine	4.41	0.058	0.016	1.082	0.031	5.97	1.74
Y/F4-DE6	dentine	4.93	0.072	0.018	1.057	0.034	7.69	1.99
Y/F4-DE7	dentine	4.80	0.077	0.018	1.088	0.031	7.95	1.93
Y/F4-DE8	dentine	4.15	0.085	0.041	1.102	0.065	8.74	4.43

1068

LA raster	tissue	U(ppm)	^{230}Th/^{238}U	±	^{234}U/^{238}U	±	Age (ka)	±
US/22-EN1	enamel	*0.00*	*-7.442*	*0.102*	*-6.047*	*0.023*	n.c.	n.c.
US/22-EN2	enamel	*0.00*	*-53.99*	*0.082*	*-14.98*	*0.020*	n.c.	n.c.
US/22-EN3	enamel	*0.00*	*215.0*	*0.103*	*78.83*	*0.019*	n.c.	n.c.
US/22-EN4	enamel	*0.00*	*2.054*	*0.109*	*-12.69*	*0.020*	n.c.	n.c.
US/22-EN5	enamel	*0.02*	*-2.955*	*0.109*	*1.044*	*0.052*	n.c.	n.c.
US/22-DE6	dentine	1.28	0.066	0.070	1.205	0.100	6.09	6.66
US/22-DE7	dentine	1.36	0.085	0.073	1.179	0.086	8.10	7.22
US/22-DE8	dentine	1.38	0.094	0.060	1.183	0.071	8.94	5.98
US/22-DE9	dentine	1.41	0.126	0.062	1.183	0.087	12.27	6.38
US/22-DE10	dentine	1.41	0.090	0.050	1.234	0.080	8.17	4.77

1069

^{230}Th/^{238}U and ^{234}U/^{238}U are activity ratios. It is worth noting that, for most transect analyses,
the ^{232}Th signal, which was measured on a Faraday collector, was indistinguishable from
background noise. In this regard, the corresponding ^{230}Th/^{232}Th activity ratio of each transect
should be >>100, and thus non-radiogenic or detrital ^{230}Th correction would have negligible
impact on the age. U-series data in italics should be viewed with caution due to U
concentrations of ≤ 0.5 ppm. All errors are 2-σ. Key: EN= enamel; DE = dentine; n.c. = not
calculable. Negative values were caused by background extraction from their measured peaks
with intensities at detection levels.