Proteomic analysis revealed the function of PoElp3 in development, pathogenicity and autophagythrough the tRNA-mediated translation efficiency in the rice blast fungus

Ting Sun[1,†], Yuyong Li[1,†], Yuanhao Liu[1,†], Xianjun Wang[2], Jiahui Xie[2], Huimin Bai[2], Jiayi Hu[2], Zifan Zhang[1], Xiuxiu Li[2], Zonghua Wang[4,*], Huakun Zheng[1,*], Guifang Lin[3,*]

[1] Fujian Universities Key Laboratory for Plant Microbe Interaction, College of Life Sciences, Fujian Agriculture and Forestry University, Fuzhou 350002, China.

[2] State Key Laboratory of Ecological Pest Control for Fujian and Taiwan Crops, College of Plant Protection, Fujian Agriculture and Forestry University, Fuzhou 350002, China.

[3] Basic Forestry and Proteomics Research Center, Fujian Agriculture and Forestry University, Fuzhou, 350002, China.

[4] Fuzhou Institute of Oceanography, Minjiang University, Fuzhou 350108, China.

* Correspondence: glin2150@163.com (G.L.); huakunz2017@fafu.edu.cn (H.Z.); wangzh@fafu.edu.cn (Z.W.).

[†] These authors contributed equally to this work.

Abstract

The Elongator complex is conserved in a wide range of species and plays crucial roles in diverse cellular processes. We have previously shown that the Elongator protein PoELp3 was involved in the asexual development, pathogenicity, and autophagy of the rice blast fungus. In this study, we further revealed that PoElp3 functions via tRNA-mediated protein integrity. Based on the phenotypic results that overexpression of two of the tRNAs, tK(UUU) and tQ(UUG), could rescue the defects in fungal growth, pathogenicity and response to rapamycin of ΔPoelp3 strain, we performed TMT-based proteomic analysis, and showed that 386 proteins were down-regulated in ΔPoelp3 strain compared with wild type strain Guy11. Transcriptional analysis demonstrated that regulation of the 386 proteins by PoElp3 was independent of transcription. Codon usage assays revealed an enrichment of Glutamine CAA-biased mRNA in the 386 proteins compared with the 70-15 genome. Functional assessment of these proteins suggested that some of them were involved in either fungal growth, conidial development, pathogenicity, or autophagy. Taken together, our results suggested that PoElp3 acts through the tRNA-mediated translational efficiency to regulate asexual development, pathogenicity, and autophagy in the rice blast fungus.

Key words: Elongator, tRNA, proteomics, transcription, translational efficiency, rice blast fungus

1. Introduction

Transfer RNAs (tRNAs) are central players in protein biosynthesis that decipher the genetic code of mRNA and add appropriate amino acids into the growing amino acid chain (Holley *et al.* 1965; Ma *et al.* 2021). The simple primary sequence of tRNAs determines that it could fold into a diverse structure in addition to the one with biological function. Therefore, the posttranscriptional modifications of tRNA are essential for the maturing, folding, degradation, and decoding efficiency (Sokołowski *et al.* 2018). The 5-methoxycarbonylmethyl- 2-thiouridine (mcm^5s^2) modification of tRNA at U_{34} is one of the best-studied modifications and is conserved in many species (Leidel *et al.* 2009; Sokołowski *et al.* 2018). The mcm^5s^2-modified tRNAs decipher the NAA type codons and wobble onto NAG. For example, the lysine, glutamine, and glutamic acid are encoded by either AAA, CAA, and GAA, or AAG, CAG, and GAG, respective. And a single eukaryotic tRNA ($tRNA^{Lys}$ $_{UUU}$ (tK^{UUU}), $tRNA^{Glu}$ $_{UUC}$ (tE^{UUC}), or $tRNA^{Gln}$ $_{UUG}$ (tQ^{UUG})) with a U at position 34 can recognize both the NAA and NAG (wobble) codons (Dewez *et al.* 2008; Goffena *et al.* 2018). The Elongator complex is involved in the first step of mcm^5s^2 modification by transferring a carboxymethyl group to U_{34} at the C_5 position to form the 5-carboxymethyluridine (cm^5U) precursor (Selvadurai *et al.* 2014). Although Elongator complex was found to modify 11 U_{34}-carrying tRNA species, only tK^{UUU}, tE^{UUC}, and tQ^{UUG} were subsequently thiolated (S2) (Johansson *et al.* 2008).

Elp3 is the catalytic subunit of the Elongator complex, which was originally identified in *Saccharomyces cerevisiae* as a component of elongating RNA polymerase II (RNAPII) holoenzyme (Otero *et al.* 1999). Elp3 contains a Gcn5-related N-terminal histone acetyltransferase (HAT) domain (Winkler *et al.* 2002), which was reported to have the acetyltransferase activity on histone proteins and other proteins (Creppe *et al.* 2009; Miśkiewicz *et al.* 2011; Laguesse *et al.* 2017; Lin *et al.* 2019; Planelles-Herrero *et al.* 2022). However, the effect of Elp3 on histone acetylation could be altered in different organisms,

in suggestive of diverse regulatory mechanisms in different backgrounds. For example, the H3K14ac level was significantly reduced in *S. cerevisiae* (Wittschieben *et al.* 2000), *Fusarium graminearum* (Lee *et al.* 2014) and *Beauveria bassiana* (Cai *et al.* 2022), but not in *Aspergillus fumigatus* (Zhang *et al.* 2022) and *Pyricularia oryzae* (Zhang *et al.* 2021). Elp3 also possesses a radical S-adenosylmethionine (SAM) domain, which is responsible for the modification of wobble uridine residues in eukaryotic tRNA (Dauden *et al.* 2019; Lin *et al.* 2019; Abbassi *et al.* 2020). Increasing evidence demonstrated that overexpression of corresponding hypomodified tRNA could rescue the defects caused by Elongator complex-associated s^2U_{34} deficiency in many organisms, such as *S. cerevisiae* (Esberg *et al.* 2006; Nedialkova and Leidel 2015; Tükenmez *et al.* 2015; Jaciuk *et al.* 2023), *Caenorhabditis elegans* (Nedialkova and Leidel 2015), and *Aspergillus fumigatus* (Zhang *et al.* 2022). In line with this, studies in different species demonstrated that both the SAM and KAT domains are required for the modification of tRNA by Elp3 proteins (Selvadurai *et al.* 2014; Zhang *et al.* 2022). These results further support a key role of Elongator complex in maintaining the wobbling function of tRNA.

In *P. oryzae* (syn. *Magnaporthe orzyae*), causal agent of the destructive blast disease, autophagy is crutial for diverse developmental processes, particularly, the appressorial development (Liu *et al.* 2007; Wilson and Talbot 2009; He *et al.* 2018; Sun *et al.* 2018; Yin *et al.* 2019). We had previously shown that the Elp3 homolog was required for the development and pathogenicity in the rice blast fungus *P. oryzae*, probably by maintaining the autophagy homeostasis (Zhang *et al.* 2021). In this study, we aim to investigate whether PoElp3 is involved in the wobbling function of tRNA and subsequently the proteomic integrity.

2. Materials and methods

2.1. Manipulation of fungal strains

The fungal strains were cultivated as described previously (Anjago *et al.* 2022).

In brief, all strains were grown at 28 ºC on complete medium (CM, containing

yeast extract 0.6%, casamino acid 0.6%, sucrose 1%, agar 2%). For

rapamycin inhibition assay, the strains were grown at 28 ºC on MM medium

supplemented with or without 1 μg/mL rapamycin (He *et al.* 2018). For all the

fungal growth assays, images were taken at 7 days after cultivation. All the

assays were performed in three independent biological experiments. For RNA

and DNA extraction, the strains were grown in CM broth for 3 days before

collection.

2.2 Spray inoculation assay

The spray inoculation assays were performed as previously described with

minor modification (Zhang *et al.* 2021). Conidial suspensions (5×10^4

conidia/mL in 0.025% Tween-20 solution) of each strain were used for spray

inoculation assay on 14-day-old TP309 rice seedlings. The challenged rice

seedlings were then cultivated under the conditions described previously

(Giraldo *et al.* 2013). The symptoms were recorded and scored as previously

described (Valent *et al.* 1991).

2.3. Construction of tRNA overexpression strains

For the identification of tRNA^{Lys} $_{UUU}$ (tK^{UUU}) and tRNA^{Gln} $_{UUG}$ (tQ^{UUG}), we search

the rice blast genome for the homologs of SPBTRNALYS.06 and

SPBTRNAGLN.02, respectively, from *Schizosaccharomyces pombe* (fission

yeast). The identified coding sequences of each tRNA were amplified using the

primers listed in Appendix A and inserted into the *Xho*I and *Hind*III sites of

pKNT-RP27, which was then introduced into the ΔPoelp3 mutant strain

through the PEG3350-meidiated protoplast transformation (Sweigard *et al.*

1998). The resulting transformants were selected on TB3 medium with 600

μg/mL G418 and further tested through PCR-based genotyping.

2.4. Proteomics analysis

The proteomic analysis was performed as previously described with some

modifications (Zhang *et al.* 2022). Tandem Mass Tags (TMT)-based

proteomics analysis was conducted by Jingjie PTM Bio (Hangzhou, China). 50

µg total protein of each sample was digested with trypsin for TMT labeling and fraction. The LC-MS/MS analysis was performed using an Orbitrap Exploris™ 480 mass spectrometer (Thermo Scientific™, USA). The MS raw data were processed with Proteome Discoverer (v2.4.1.15) and searched against 70-15 reference genome (Dean *et al.* 2005).

2.5. RNA-sequencing and data analysis

The RNA-sequencing and data analysis were performed as described previously (Xie *et al.* 2023). The total RNA of each sample was used for the construction of the RNA-seq library as described previously (Zhong *et al.* 2019). The libraries were sequenced using the Illumina NovaSeq 6000 system. Quality assessment of raw reads (RNA-seq) was conducted using FASTQC v0.11.3 (www.bioinformatics. babraham.ac.uk/bugzilla/). Adapters were removed from the raw reads using Trim_galore v3.1 (https://github.com/FelixKrueger/TrimGalore/). Then, the RNA-seq reads were aligned to 70-15 reference genome (Dean *et al.* 2005) using RSEM v1.3.3 (Li and Dewey 2011). The quantification of expression counts were conducted with transcript per million (TPM).

2.6. Statistical analysis

Statistical analysis was conducted by R packages in the R studio (Version 1.4.1106) environment (Xie *et al.* 2023).

3. Results

3.1. Overexpression of tRNA could rescue the defects in ∆*Poelp3* strain

To test whether the overexpression of tRNA could complement the defects showed in ∆*Poelp3* strain, we search in the rice blast fungal genome for genes encoding tK^{UUU} and tQ^{UUG}, the two tRNA that could rescue the transcription and histone acetylation defects, as well as other phenotypes caused by the absence of Elongator when overexpressed (Esberg *et al.* 2006). We found that both tRNAs have two coding genes, with tK^{UUU}-4 and tK^{UUU}-6 localized at chr4 and chr6, tQ^{UUG}-2 and tQ^{UUG}-4 localized at chr2 and chr4, respectively. The

coding sequence of the two copies of each tRNAs was highly conserved, while the franking sequences were highly polymorphic (Appendix B). The four tRNA coding genes were fused with the ribosomal protein promoter (*RP27p*) and introduced into the Δ*Poelp3* strain, respectively. At 7 days post inoculation, the diameter of transfromants carrying the *RP27p* :: *tK^{UUU}* (herein after referred to as '*tK(UUU)*') or *RP27p* :: *tQ^{UUG}* (herein after referred to as '*tQ(UUG)*') was about 95%-100% of that of wild type, while the diameter of Δ*Poelp3* strain was about 85% of that of the wild type (Fig. 1-A and B). Upon the treatment of rapamycin, inhibitor of the Target-of-Rapamycin (TOR) signaling pathway, the inhibition rate of fungal growth in the *tK^{UUU}* and *tQ^{UUG}* strains was reduced compared to Δ*Poelp3* strain, but was still larger than that of Guy11 (Fig. 1-A and C). Spray inoculation results showed that expression of *tK^{UUU}* and *tQ^{UUG}* could also partially rescue the defects in pathogencity caused by the absence of *PoELP3* (Fig. 1-D and E). These results suggested that the defects caused by the deletion of *PoELP3* were at least partially caused by the inefficiency in protein biosynthesis.

3.2. PoElp3 is involved in the proper expression of protein

To further investigate how PoElp3 regulates fungal development, pathogenicity, and autophagy by affecting protein biosynthesis efficiency, we compared the protein abundance between Δ*Poelp3* strain and the Guy11 wild type strain through the tandem mass tags (TMT)-based quantitative proteomic analysis. Among the 5,165 proteins identified and quantified in our proteomic data, 386 proteins were significantly down-regulated [fold change (FC) > 1/1.3 (*p-value* < 0.05)], and 403 proteins were up-regulated [FC < 1.3 (*p-value* < 0.05)] (Fig. 2-A and B; Appendix C and D). The volcano plot result showed that most of the differentially accumulated proteins (DAPs) with a FC ranged from 1.3 to 2.0, and only a few of them with a FC >2.0 (Fig. 2-B). The clustering result showed that the DAPs could be divided into different groups according to their differential abundance between Guy11 and Δ*Poelp3* strain (Fig. 2-C).

3.3. The DAPs involved in different pathways

To perform functional classification of the DAPs, we divided the DAPs into 4 Categories: Q1 with a FC<1/1.5, Q2 with a FC between 1/1.5 to 1/1.3, Q3 with a FC between 1.3 to 1.5, and Q4 with a FC>1.5 (Fig. 3-A). KEGG enrichment results showed that the Q1 DAPs were enriched in pathways such as Glyoxylate and dicarboxylate metabolism, biosynthesis of unsaturated fatty acids, various types of N-glcan biosynthesis, sulfur metabolism and sphingolipid metabolism. The Q2 DAPs were mainly enriched in steroid biosynthesis, tryptophan metabolism, and terpenoid backbone biosynthesis. While the Q3 and Q4 DAPs were mainly enriched in pathways such as aminoacyl-tRNA biosynthesis, biosynthesis, and metabolism of amino acids (Fig. 3-B). Protein domain enrichment showed that the Q1 DAPs were mainly enriched in pathways such as adenylysulphate kinase, fatty acid desaturase, and pyridine nucleotide-disulphide oxidoreductase. The Q2 DAPs were mainly enriched in pathways such as dynamin, flavodoxin, enoyl-(axyl carrier protein) reductase, and alpha amylase. While the Q3 and Q4 DAPs were mainly enriched in pathways such as tRNA synthetases, pyridoxamine 5'-phosphate oxidase, NADH oxidase and oxidoreductase NAD-binding domain (Fig. 3-C).

3.4. PoElp3 regulated the accumulation of proteins independent of transcription

To further assess whether the DAPs were associated with the differential transcription level of their coding genes, we performed a comparative transcriptome analysis between the wild type and ΔPoelp3 strains. We showed that only 535 genes were down-regulated, and 1,049 genes were up-regulated in ΔPoelp3 compared with the wild type strain Guy11 (Fig. 4-A). We further investigated the transcription level of DAPs down-regulated in ΔPoelp3 strain, and found that, among the 386 proteins down-regulated in ΔPoelp3 strain, only 19 and 25 of them showing either higher or lower transcription level in the mutant strain (Fig. 4-B; Appendix E). While among the 403 DAPs up-regulated

in Δ*Poelp3* strain, only 22 showing lower transcription level in the mutant strain, but 138 showing higher transcription level in the mutant strain (Appendix F), which could be affected indirectly. Taken together, these results suggested that the protein level of most of the DAPs in Δ*Poelp3* strain was independent of the transcription level.

3.5. Enrichment of Glutamine CAA-biased mRNA in the DAPs down-regulated in Δ*Poelp3* strain

To test whether there is a bias toward the use of NAA codons, we first surveyed the coding sequence of 70-15 genome (whole genome, WG), and found that 65.0%, 52.5% and 58.0% of the coding sequences showed an NAA:NAG ratio > 1 for Lysine, Glutamine, and Glutamic acid, respectively (Table 1). We then surveyed the coding sequence of DAPs down-regulated in Δ*Poelp3* strain. Since the DAPs down-regulated in Δ*Poelp3* strain were supposed to be caused by the absence of PoElp3, and are therefore referred to as PoElp3 regulated protein candidates (Erps) hereafter. We found that 58.8.0%, 64.8% and 50.5.0% of the coding sequences showed an NAA:NAG ratio > 1 for Lysine, Glutamine, and Glutamic acid, respectively (Table 1; Appendix G). These results suggested an enrichment of Glutamine CAA-biased mRNA in the Erps compared with 70-15 genome.

3.6. The Elp3-regulated proteins involved in asexual development and pathogenicity

To assess whether any of the DAPs down-regulated in Δ*Poelp3* strain functions similarly to PoElp3, we searched in the PubMed database for publications associated with the 386 DAPs and found six DAPs involved in vegetative growth, conidiation, pathogenicity and/or autophagy. These DAPs presented a Fold Change (FC) ranging from 0.49~0.76, and their transcription level in Δ*Poelp3* strain was no less than that in Wild type strain Guy11 (Table 2 and Appendix E). Among them, *ERP1* encodes a core protein of ESCRT-III

subcomplex (Snf7) involved in asexual development, pathogenicity and autophagy (Cheng *et al.* 2018). *ERP2* encoding a Rho3 homolog is involved in appressorial development and pathogenicity (Zheng *et al.* 2007). *ERP3* encodes an ADP-ribosylation factor 6 protein (Arf6), which is involved in endocytosis and polarity during the asexual development (Zhu *et al.* 2016). *ERP4* encodes a tetrahydroxynaphthalene reductase (Buf1) and is required for melanin biosynthesis and full virulence (Zhu *et al.* 2021). *ERP5* encodes an amidophosphoribosyltransferase (Ade4), which catalyzes the conversion of 5-phosphoribosyl-1-pyrophosphate into 5-phosphoribosyl-1-amine in the de novo purine biosynthetic pathway and is required for conidiation and pathogenicity (Aron *et al.* 2022). *ERP6* encoding a glyoxylate aminotransferase 1 (Agt1) is required for mobilization and utilization of triglycerides during infection (Bhadauria *et al.* 2012a; Bhadauria *et al.* 2012b). Besides, we also found DAPs with their homologs in other fungal species associated with development or pathogenicity. For example, the MGG_03393 encodes a secondary metabolism regulator (Lae1) and is involved in asexual development and mycoparasitism of *Trichoderma atroviride* (Karimi Aghcheh *et al.* 2013). MGG_00413T0 encoding a putative β-1,3-glucanosyltransferase (Gel7) is required for the cell wall integrity under stress conditions (Zhao *et al.* 2014).

4. Discussion

The Elongator complex is conserved among different species and increasing evidence demonstrated that they function mainly through the modification of tRNA (Bauer *et al.* 2012; Candiracci *et al.* 2019; Dauden *et al.* 2019). In supporting this, overexpression of unmodified tRNA for glutamine, glutamic acid and lysine was efficient to restore the defects caused by the absence of Elongator in several studies (Esberg *et al.* 2006; Bauer *et al.* 2012; Zhang *et al.* 2022), probably by increasing the intracellular concentration of tRNAs to compensate for decreased binding affinity in the Elongator mutants (Esberg *et*

al. 2006; Nedialkova and Leidel 2015; Tükenmez *et al.* 2015; Sokołowski *et al.*

2018). In this study, the phenotypic analysis showed that overexpression of

tK(*UUU*) or *tQ*(*UUG*) could rescue fully the defects in fungal growth, and

partially the defects in pathogenicity and response to rapamycin caused by the

deletion of *PoELP3*, which indicated that PoElp3 may act through the

tRNA-mediated protein integrity to regulate the fungal growth in rice blast

fungus (Zhang *et al.* 2021).

Elp3 proteins had been proved to catalyze the modification of tRNA at

wobbling U34 and consequently maintain the translation efficiency of proteins

encoded by tK(UUU), tE(UUC), or tQ(UUG) (Bauer *et al.* 2012; Goffena *et al.*

2018). Previous studies suggested that NAA-biased codon usage is not

always necessary for the Elongator regulated protein integraty via the tRNA

modification. For example, the biased codon usage was exhibited in in yeast

(Bauer *et al.* 2012), but not in *A. fumigatus* (Zhang *et al.* 2022). In this study,

we showed that the Erps displayed an enrichment of Glutamine CAA-biased

-biased mRNA, which suggested that different regulatory mechanisms might

be applied in the Elp3 regulated translational efficiency.

The Elongator proteins could act by regulating the translational

efficiency of target with important functions. For example, in fission yeast, the

Elongator controls cell division by regulating the translational efficiency of the

Cdr2 kinase (Bauer *et al.* 2012). Our functional assessment of the Erps

showed that some of them were associated with asexual development, such

as Erp1/Snf7, Erp2/Rho3, Erp3/ArF6, Erp4/Ade4, Erp5/Buf1, Erp6/Agt1, Lae1

and Gel7. Recent study in *A. fumigatus* showed that defects in development

and pathogenicity in Δ*elp3* strain could be caused by the amino acid

homeostasis imbalance (Zhang *et al.* 2022). We also found that many Erps

were involved in the biosynthesis and metabolism of amino acids, suggesting a

role of PoElp3 in development and pathogenicity by modulating amino acid

homeostasis. Moreover, we also found Erps that may be involved in autophagy

through different mechanisms. For example, Erp1/Snf7 acts in vesicle

trafficking and autophagy (Cheng *et al.* 2018). Erp3/Arf6 is involved in autophagy homeostasis through the generation of phosphatidylinositol 4,5-bisphosphate (PIP(2)) and induction of phospholipase D (PLD) activity (Moreau *et al.* 2012). The absence of Erp4/Ade4 caused delayed glycogen mobilization and reduced turgor pressure in the appressoria (Aron *et al.* 2022), the representative cellular processes of autophagy during appressorial development in rice blast fungus (Deng *et al.* 2008; Kershaw and Talbot 2009). Besides, we also found that some Erps were involved in the sphingolipid metabolism pathway. Although the role of sphingolipid was not well studied in filamentous fungus (Liu *et al.* 2019), it was shown to be involved in autophagy homeostasis in mammals (Li *et al.* 2014). These results together suggested that PoElp3 is required for the development, pathogenicity and autophagy by regulating the translational efficiency of proteins involved in diverse pathways. And detailed analysis of the Erps may further support the function of PoElp3 in autophagy via the tRNA-mediated protein integrity.

Although the role of Elongator in tRNA-mediated protein integrity is well established, there are several lines of evidence suggesting that Elp3 also acts by acetylation of other proteins, such as microtubules (Planelles-Herrero *et al.* 2022), α-tubulin (Creppe *et al.* 2009), bruchpilot (Laguesse *et al.* 2017), and connexin-43 (Miśkiewicz *et al.* 2011) in animal cells. In this study, we also found that overexpression of the two tRNAs could only partially rescue the response of Δ*Poelp3* to rapamycin treatment. Similar results were also reported in previous studies performed in fission yeast (Bauer *et al.* 2012) and *A. fumigatus* (Zhang *et al.* 2022). And these results together suggested that Elp3 may also act through other pathway(s).

5. Conclusion

In conclusion, our results demonstrated that PoElp3 was involved in the asexual development, pathogenicity, and autophagy by regulating the tRNA-mediated protein integrity.

350   **Competing interests**

351   The authors declare that they have no competing interests.

353   **References**

354   Abbassi NE, Biela A, Glatt S and Lin TY. 2020. How Elongator Acetylates tRNA Bases. *Int J Mol Sci,* **21,**

355   Anjago WM, Biregeya J, Shi M, Chen Y, Wang Y, Wang Z, Hong Y and Chen M. 2022. The calcium
356   chloride responsive type 2C protein phosphatases play synergistic roles in regulating MAPK
357   pathways in *Magnaporthe oryzae*. *J Fungi (Basel),* **8,**

358   Aron O, Otieno FJ, Tijjani I, Yang Z, Xu H, Weng S, Guo J, Lu S, Wang Z and Tang W. 2022. De novo
359   purine nucleotide biosynthesis mediated by MoAde4 is required for conidiation, host
360   colonization and pathogenicity in *Magnaporthe oryzae*. *Appl Microbiol Biotechnol,* **106,**
361   5587-5602.

362   Bauer F, Matsuyama A, Candiracci J, Dieu M, Scheliga J, Wolf DA, Yoshida M and Hermand D. 2012.
363   Translational control of cell division by Elongator. *Cell Rep,* **1,** 424-433.

364   Bhadauria V, Banniza S, Vandenberg A, Selvaraj G and Wei Y. 2012a. Alanine: Glyoxylate
365   aminotransferase 1 is required for mobilization and utilization of triglycerides during infection
366   process of the rice blast pathogen, *Magnaporthe oryzae*. *Plant Signal Behav,* **7,** 1206-1208.

367   Bhadauria V, Banniza S, Vandenberg A, Selvaraj G and Wei Y. 2012b. Peroxisomal alanine: glyoxylate
368   aminotransferase AGT1 is indispensable for appressorium function of the rice blast pathogen,
369   *Magnaporthe oryzae*. *PLoS One,* **7,** e36266.

370   Cai Q, Wang J, Xie J, Jiang D and Keyhani NO. 2022. The Elongator subunit Elp3 regulates development,
371   stress tolerance, cell cycle, and virulence in the entomopathogenic fungus *Beauveria*
372   *bassiana*. *J Fungi (Basel),* **8,**

373   Candiracci J, Migeot V, Chionh YH, Bauer F, Brochier T, Russell B, Shiozaki K, Dedon P and Hermand D.
374   2019. Reciprocal regulation of TORC signaling and tRNA modifications by Elongator enforces
375   nutrient-dependent cell fate. *Sci Adv,* **5,** eaav0184.

376   Cheng J, Yin Z, Zhang Z and Liang Y. 2018. Functional analysis of MoSnf7 in *Magnaporthe oryzae*.
377   *Fungal Genet Biol,* **121,** 29-45.

378   Creppe C, Malinouskaya L, Volvert ML, Gillard M, Close P, Malaise O, Laguesse S, Cornez I, Rahmouni S,
379   Ormenese S, Belachew S, Malgrange B, Chapelle JP, Siebenlist U, Moonen G, Chariot A and
380   Nguyen L. 2009. Elongator controls the migration and differentiation of cortical neurons

through acetylation of alpha-tubulin. *Cell,* **136,** 551-564.

Dauden MI, Jaciuk M, Weis F, Lin TY, Kleindienst C, Abbassi NEH, Khatter H, Krutyhołowa R, Breunig KD,

Kosinski J, Müller CW and Glatt S. 2019. Molecular basis of tRNA recognition by the Elongator

complex. *Sci Adv,* **5,** eaaw2326.

Dean RA, Talbot NJ, Ebbole DJ, Farman ML, Mitchell TK, Orbach MJ, Thon M, Kulkarni R, Xu JR, Pan H,

Read ND, Lee YH, Carbone I, Brown D, Oh YY, Donofrio N, Jeong JS, Soanes DM, Djonovic S,

Kolomiets E, Rehmeyer C, Li W, Harding M, Kim S, Lebrun MH, Bohnert H, Coughlan S, Butler J,

Calvo S, Ma LJ, Nicol R, Purcell S, Nusbaum C, Galagan JE and Birren BW. 2005. The genome

sequence of the rice blast fungus *Magnaporthe grisea*. *Nature,* **434,** 980-986.

Deng YZ, Ramos-Pamplona M and Naqvi NI. 2008. Methods for functional analysis of macroautophagy

in filamentous fungi. *Methods Enzymol,* **451,** 295-310.

Dewez M, Bauer F, Dieu M, Raes M, Vandenhaute J and Hermand D. 2008. The conserved Wobble

uridine tRNA thiolase Ctu1-Ctu2 is required to maintain genome integrity. *Proc Natl Acad Sci*

U S A, **105,** 5459-5464.

Esberg A, Huang B, Johansson MJ and Byström AS. 2006. Elevated levels of two tRNA species bypass

the requirement for elongator complex in transcription and exocytosis. *Mol Cell,* **24,** 139-148.

Giraldo MC, Dagdas YF, Gupta YK, Mentlak TA, Yi M, Martinez-Rocha AL, Saitoh H, Terauchi R, Talbot NJ

and Valent B. 2013. Two distinct secretion systems facilitate tissue invasion by the rice blast

fungus *Magnaporthe oryzae*. *Nat Commun,* **4,** 1996.

Goffena J, Lefcort F, Zhang Y, Lehrmann E, Chaverra M, Felig J, Walters J, Buksch R, Becker KG and

George L. 2018. Elongator and codon bias regulate protein levels in mammalian peripheral

neurons. *Nat Commun,* **9,** 889.

He M, Xu Y, Chen J, Luo Y, Lv Y, Su J, Kershaw MJ, Li W, Wang J, Yin J, Zhu X, Liu X, Chern M, Ma B, Wang

J, Qin P, Chen W, Wang Y, Wang W, Ren Z, Wu X, Li P, Li S, Peng Y, Lin F, Talbot NJ and Chen X.

2018. MoSnt2-dependent deacetylation of histone H3 mediates MoTor-dependent

autophagy and plant infection by the rice blast fungus *Magnaporthe oryzae*. *Autophagy,* **14,**

1543-1561.

Holley RW, Apgar J, Everett GA, Madison JT, Marquisee M, Merrill SH, Penswick JR and Zamir A. 1965.

STRUCTURE OF A RIBONUCLEIC ACID. *Science,* **147,** 1462-1465.

Jaciuk M, Scherf D, Kaszuba K, Gaik M, Rau A, Kościelniak A, Krutyhołowa R, Rawski M, Indyka P,

Graziadei A, Chramiec-Głąbik A, Biela A, Dobosz D, Lin TY, Abbassi NE, Hammermeister A,

Rappsilber J, Kosinski J, Schaffrath R and Glatt S. 2023. Cryo-EM structure of the fully

assembled Elongator complex. *Nucleic Acids Res,* **51,** 2011-2032.

Johansson MJ, Esberg A, Huang B, Björk GR and Byström AS. 2008. Eukaryotic wobble uridine

modifications promote a functionally redundant decoding system. *Mol Cell Biol,* **28,**
3301-3312.

Karimi Aghcheh R, Druzhinina IS and Kubicek CP. 2013. The putative protein methyltransferase LAE1 of
Trichoderma atroviride is a key regulator of asexual development and mycoparasitism. *PLoS*
One, 8, e67144.

Kershaw MJ and Talbot NJ. 2009. Genome-wide functional analysis reveals that infection-associated
fungal autophagy is necessary for rice blast disease. *Proc Natl Acad Sci U S A,* **106,**
15967-15972.

Laguesse S, Close P, Van Hees L, Chariot A, Malgrange B and Nguyen L. 2017. Loss of Elp3 impairs the
acetylation and distribution of connexin-43 in the developing cerebral cortex. *Front Cell*
Neurosci, **11,** 122.

Lee Y, Min K, Son H, Park AR, Kim JC, Choi GJ and Lee YW. 2014. ELP3 is involved in sexual and asexual
development, virulence, and the oxidative stress response in *Fusarium graminearum*. *Mol*
Plant Microbe Interact, **27,** 1344-1355.

Leidel S, Pedrioli PG, Bucher T, Brost R, Costanzo M, Schmidt A, Aebersold R, Boone C, Hofmann K and
Peter M. 2009. Ubiquitin-related modifier Urm1 acts as a sulphur carrier in thiolation of
eukaryotic transfer RNA. *Nature,* **458,** 228-232.

Li B and Dewey CN. 2011. RSEM: accurate transcript quantification from RNA-Seq data with or without
a reference genome. *BMC Bioinformatics,* **12,** 323.

Li Y, Li S, Qin X, Hou W, Dong H, Yao L and Xiong L. 2014. The pleiotropic roles of sphingolipid signaling
in autophagy. *Cell Death Dis,* **5,** e1245.

Lin TY, Abbassi NEH, Zakrzewski K, Chramiec-Głąbik A, Jemioła-Rzemińska M, Różycki J and Glatt S.
2019. The Elongator subunit Elp3 is a non-canonical tRNA acetyltransferase. *Nat Commun,* **10,**
625.

Liu XH, Liang S, Wei YY, Zhu XM, Li L, Liu PP, Zheng QX, Zhou HN, Zhang Y, Mao LJ, Fernandes CM, Del
Poeta M, Naqvi NI and Lin FC. 2019. Metabolomics analysis identifies sphingolipids as key
signaling moieties in appressorium morphogenesis and function in *Magnaporthe oryzae*.
mBio, **10,**

Liu XH, Lu JP, Zhang L, Dong B, Min H and Lin FC. 2007. Involvement of a *Magnaporthe grisea*
serine/threonine kinase gene, MgATG1, in appressorium turgor and pathogenesis. *Eukaryot*
Cell, 6, 997-1005.

Ma X, Liu C and Cao X. 2021. Plant transfer RNA-derived fragments: Biogenesis and functions. *J Integr*
Plant Biol, **63,** 1399-1409.

Miśkiewicz K, Jose LE, Bento-Abreu A, Fislage M, Taes I, Kasprowicz J, Swerts J, Sigrist S, Versées W,

Robberecht W and Verstreken P. 2011. ELP3 controls active zone morphology by acetylating

the ELKS family member Bruchpilot. *Neuron,* **72,** 776-788.

Moreau K, Ravikumar B, Puri C and Rubinsztein DC. 2012. Arf6 promotes autophagosome formation

via effects on phosphatidylinositol 4,5-bisphosphate and phospholipase D. *J Cell Biol,* **196,**

483-496.

Nedialkova DD and Leidel SA. 2015. Optimization of codon translation rates via tRNA modifications

maintains proteome integrity. *Cell,* **161,** 1606-1618.

Otero G, Fellows J, Li Y, de Bizemont T, Dirac AM, Gustafsson CM, Erdjument-Bromage H, Tempst P and

Svejstrup JQ. 1999. Elongator, a multisubunit component of a novel RNA polymerase II

holoenzyme for transcriptional elongation. *Mol Cell,* **3,** 109-118.

Planelles-Herrero VJ, Bittleston A, Seum C, Daeden A, Gaitan MG and Derivery E. 2022. Elongator

stabilizes microtubules to control central spindle asymmetry and polarized trafficking of cell

fate determinants. *Nat Cell Biol,* **24,** 1606-1616.

Selvadurai K, Wang P, Seimetz J and Huang RH. 2014. Archaeal Elp3 catalyzes tRNA wobble uridine

modification at C5 via a radical mechanism. *Nat Chem Biol,* **10,** 810-812.

Sokołowski M, Klassen R, Bruch A, Schaffrath R and Glatt S. 2018. Cooperativity between different

tRNA modifications and their modification pathways. *Biochim Biophys Acta Gene Regul Mech,*

1861, 409-418.

Sun G, Elowsky C, Li G and Wilson RA. 2018. TOR-autophagy branch signaling via Imp1 dictates

plant-microbe biotrophic interface longevity. *PLoS Genet,* **14,** e1007814.

Sweigard JA, Carroll AM, Farrall L, Chumley FG and Valent B. 1998. *Magnaporthe grisea* pathogenicity

genes obtained through insertional mutagenesis. *Mol Plant Microbe Interact,* **11,** 404-412.

Tükenmez H, Xu H, Esberg A and Byström AS. 2015. The role of wobble uridine modifications in +1

translational frameshifting in eukaryotes. *Nucleic Acids Res,* **43,** 9489-9499.

Valent B, Farrall L and Chumley FG. 1991. *Magnaporthe grisea* genes for pathogenicity and virulence

identified through a series of backcrosses. *Genetics,* **127,** 87-101.

Wilson RA and Talbot NJ. 2009. Under pressure: investigating the biology of plant infection by

Magnaporthe oryzae. *Nat Rev Microbiol,* **7,** 185-195.

Winkler GS, Kristjuhan A, Erdjument-Bromage H, Tempst P and Svejstrup JQ. 2002. Elongator is a

histone H3 and H4 acetyltransferase important for normal histone acetylation levels *in vivo*.

Proc Natl Acad Sci U S A, **99,** 3517-3522.

Wittschieben BO, Fellows J, Du W, Stillman DJ and Svejstrup JQ. 2000. Overlapping roles for the

histone acetyltransferase activities of SAGA and elongator *in vivo*. *Embo j,* **19,** 3060-3068.

Xie J, Tang W, Lu G, Hong Y, Zhong Z, Wang Z and Zheng H. 2023. Histone H3K27me3 methylation

regulates the expression of secreted proteins distributed at fast-evolving regions through

transcriptional repression of transposable elements. *Journal of Integrative Agriculture,*

Yin Z, Chen C, Yang J, Feng W, Liu X, Zuo R, Wang J, Yang L, Zhong K, Gao C, Zhang H, Zheng X, Wang P

and Zhang Z. 2019. Histone acetyltransferase MoHat1 acetylates autophagy-related proteins

MoAtg3 and MoAtg9 to orchestrate functional appressorium formation and pathogenicity in

Magnaporthe oryzae. *Autophagy,* **15,** 1234-1257.

Zhang L, Chen S, Qi M, Cao X, Liang N, Li Q, Tang W, Lu G, Zhou J, Yu W, Wang Z and Zheng H. 2021.

The putative elongator complex protein Elp3 is involved in asexual development and

pathogenicity by regulating autophagy in the rice blast fungus. *Journal of Integrative*

Agriculture, **20,** 2944-2956.

Zhang Y, Wang Y, Fan J, Zhu G and Lu L. 2022. *Aspergillus fumigatus* Elongator complex subunit 3

affects hyphal growth, adhesion and virulence through wobble uridine tRNA modification.

PLoS Pathog, **18,**

Zhao W, Li C, Liang J and Sun S. 2014. The *Aspergillus fumigatus* β-1,3-glucanosyltransferase Gel7 plays

a compensatory role in maintaining cell wall integrity under stress conditions. *Glycobiology,*

24, 418-427.

Zheng W, Chen J, Liu W, Zheng S, Zhou J, Lu G and Wang Z. 2007. A Rho3 homolog is essential for

appressorium development and pathogenicity of *Magnaporthe grisea*. *Eukaryot Cell,* **6,**

2240-2250.

Zhong Z, Lin L, Chen M, Lin L, Chen X, Lin Y, Chen X, Wang Z, Norvienyeku J and Zheng H. 2019.

Expression Divergence as an evolutionary alternative mechanism adopted by two rice

subspecies against rice blast infection. *Rice (N Y),* **12,** 12.

Zhu S, Yan Y, Qu Y, Wang J, Feng X, Liu X, Lin F and Lu J. 2021. Role refinement of melanin synthesis

genes by gene knockout reveals their functional diversity in *Pyricularia oryzae strains*.

Microbiol Res, **242,** 126620.

Zhu X, Zhou T, Chen L, Zheng S, Chen S, Zhang D, Li G and Wang Z. 2016. Arf6 controls endocytosis and

polarity during asexual development of *Magnaporthe oryzae*. *FEMS Microbiol Lett,* **363,**

Figure legends

Fig. 1 Overexpression of *tK*(*UUU*) or *tQ*(*UUG*) could partially rescue the defects in Δ*Poelp3* strain. A, the indicated strains grown on CM, MM and MM supplemented with 1 μg/uL rapamycin (MM+Rap). B, colony diameter of indicated strains grown on CM. C, the inhibition rate of indicated strains upon the treatment of 1 μg/uL rapamycin. The letters indicate statistically significant differences ($p<0.01$). D, leaves of rice cultivar TP309 were sprayed with conidial suspensions ($5×10^4$ conidia mL^{-1}) of strains indicated. E, quantification of different lesion types. Data were shown as the mean±SE (n=3).

Fig. 2 The differentially accumulated proteins (DAPs) between Δ*Poelp3* and Guy11 strains. A, quantitative identification of proteins upregulated and downregulated in Δ*Poelp3* strain though TMT. B, volcano plot showing the fold change of the differentially accumulated proteins. C, cluster analysis of the DAPs.

Fig. 3 Functional classification of the DAPs. A, the four Q Categories of the DAPs. B, classification of DAPs according to the KEGG pathway. C, classification of DAPs according to protein domain.

Fig. 4 Comparative transcriptional analysis between wild type and Δ*Poelp3* strains. A, differentially expressed genes between wild type and Δ*Poelp3* strains. B, comparative analysis of the transcription level of DAPs down-regulated in Δ*Poelp3* strain.

Table 1 Codon usage of lysine, glutamine and glutamic acid in the Erps.

Table 2 The *ERPs* functionally studied in *P. oryzae*

Supporting information

Appendix A. Primers used in this study.

Appendix B. Alignment of tRNAs from *Schizosaccharomyces pombe* (fission yeast) and *P. oryzae*.

Appendix C. Proteins up-regulated in Δ*Poelp3* strain.

Appendix D. Proteins down-regulated in Δ*Poelp3* strain.

Appendix E. The transcription level of DAPs down-regulated in Δ*Poelp3*

542 strain.

543 **Appendix F.** Comparative analysis of the transcription level of DAPs

544 up-regulated in ΔPoelp3 strain.

545 **Appendix G.** Codons used in Erps.

Table 1 Codon usage of lysine, glutamine and glutamic acid in the Erps.

Amino acid	Biased[a]		Non-Biased[b]		Percentage of Biased (%)	
	Erps	WG	Erps	WG	Erps	WG
Lysine	227	8,337	159	4,489	58.8	65.0
Glutamine	250	6,735	136	6,091	64.8	52.5
Glutamic acid	195	7,443	194	5,383	50.5	58.0

(a: NAA:NAG ratio > 1; b: NAA:NAG ratio ≤ 1.)

Table 2 The *ERPs* functionally studied in *P. oryzae*

Code	Gene ID	Product	Fold Change	Function	Reference
ERP1	MGG_04174	SNF7 family protein	0.76	Vegetative growth, conidiation and pathogenicity, autophagy	(Cheng *et al.* 2018)
ERP2	MGG_10323	GTP-binding protein RHO3	0.73	Appressorial development and pathogenicity	(Zheng *et al.* 2007)
ERP3	MGG_10676	ADP-ribosylation factor 6	0.71	Vegetative growth and conidial morphology	(Zhu *et al.* 2016)
ERP4	MGG_02252	Tetrahydroxynaphthalene reductase MoBuf1	0.64	Conidiation, germination and pathogenicity	(Zhu *et al.* 2021)
ERP5	MGG_04618	Amidophosphoribosyltransferase	0.71	Conidiation and pathogenicity	(Aron *et al.* 2022)
ERP6	MGG_02525	Alanine-glyoxylate aminotransferase	0.49	Appressorial development	(Bhadauria *et al.* 2012a; Bhadauria *et al.* 2012b)