NAPE-PLD in the ventral tegmental area regulates reward events, feeding and energy homeostasis

Julien Castel[1], Guangping Li[1], Oriane Onimus[1], Emma Leishman[2], Patrice D. Cani[3,4,5], Heather Bradshaw[2], Ken Mackie[2,6], Amandine Everard[1,3,4,5], Serge Luquet[1,*], Giuseppe Gangarossa[1,7,*]

[1] Université Paris Cité, CNRS, Unité de Biologie Fonctionnelle et Adaptative, F-75013 Paris, France

[2] Department of Psychological and Brain Sciences, Indiana University Bloomington, Bloomington, IN, United States

[3] Metabolism and Nutrition Research group, Louvain Drug Research Institute (LDRI), UCLouvain, Université catholique de Louvain, Brussels, Belgium

[4] WELBIO-Walloon Excellence in Life Sciences and Biotechnology, WELBIO department, WEL Research Institute, avenue Pasteur, 6, 1300 Wavre, Belgium

[5] Institute of Experimental and Clinical Research (IREC), UCLouvain, Université catholique de Louvain, Brussels, Belgium.

[6] Gill Center for Biomolecular Science, Indiana University Bloomington, Bloomington, IN, United States

[7] Institut universitaire de France (IUF)

* Senior authors

Correspondence should be addressed to: Giuseppe Gangarossa (giuseppe.gangarossa@u-paris.fr, @PeppeGanga) and Serge Luquet (serge.luquet@u-paris.fr)

Keywords: NAPE-PLD, bioactive lipids, reward, dopamine, energy balance

31 **Highlights**
32
33 - NAPE-PLD and NAEs are enriched in the VTA and regulate food-reinforced behaviors and
34 reward processes.
35
36 - NAPE-PLD scales *in vivo* VTA→NAc dopamine dynamics.
37
38 - NAPE-PLD in the VTA contributes to whole-body energy balance and metabolic efficiency.
39
- Downregulation of VTA NAPE-PLD ameliorates obesity-associated metabolic features.

Abstract

The *N*-acyl phosphatidylethanolamine-specific phospholipase D (NAPE-PLD) catalyzes the

production of *N*-acylethanolamines (NAEs), a family of endogenous bioactive lipids, which

are involved in various biological processes ranging from neuronal functions to energy

homeostasis and feeding behaviors. Reward-dependent behaviors depend on the dopamine

(DA) transmission between the ventral tegmental area (VTA) and the nucleus accumbens

(NAc) which conveys reward-values and scales reinforced behaviors. However, whether and

how NAPE-PLD may contribute to the regulation of feeding and reward-dependent behaviors

has not been investigated yet. This biological question is of paramount importance since

NAEs are altered in obesity and metabolic disorders.

Here, we show that transcriptomic meta-analysis highlights a potential role for NAPE-PLD

within the VTA→NAc circuit. Using brain-specific invalidation approaches, we report that

the integrity of NAPE-PLD is required for the proper homeostasis of NAEs within the

midbrain VTA and it affects food-reward behaviors. Moreover, region-specific knock-down

of NAPE-PLD in the VTA resulted in enhanced food-reward seeking and reinforced

behaviors which were associated with increased *in vivo* DA release dynamics in response to

both food and non-food-related rewards together with heightened tropism towards food

consumption. Furthermore, midbrain knock-down of NAPE-PLD, which led to increased

energy expenditure and adapted nutrients partitioning, elicited a relative protection against

high-fat diet-mediated body fat gain and obesity-associated metabolic features.

In conclusion, these findings unravel a new key role of VTA NAPE-PLD in shaping DA-

dependent events, feeding behaviors and energy homeostasis, thus providing new insights on

the regulation of body metabolism.

Introduction

The regulation of feeding behaviors and energy homeostasis is a cardinal and evolutionarily conserved physiological feature in mammals. By mobilizing several and functionally distinct brain circuits (1,2), such regulation tightly depends on metabolic and nutritional demands as well as on the reinforcing and hedonic properties of foods. Among the different regulatory pathways which signal homeostatic states and scale feeding behaviors, lipids, either nutritional and/or endogenous species, represent key mediators in gating the functional adaptability of complex neuronal networks that synchronize food intake and energy expenditure (3,4).

Endogenous bioactive lipids are a major class of biologically active mediators with a critical role in the regulation of several functions spanning from homeostasis to cognition. Among these, the membrane phospholipid-derived long-chain fatty acids *N*-acylethanolamines (NAEs) are potent signaling molecules in peripheral tissues as well as in the central nervous system (CNS). Given their wide distribution and multiple biological functions, the modulation of NAEs tone represents an interesting target for the development of new therapeutic approaches (5).

Among NAEs, the *N*-arachidonoylethanolamine (AEA or anandamide), a *bona fide* endocannabinoid (eCB) agonist of the cannabinoid receptors CB1R and CB2R, is synthetized by the Ca^{2+}-dependent enzyme *N*-acyl phosphatidylethanolamine-specific phospholipase D (NAPE-PLD) (6,7) from the membrane *N*-acylphosphatidylethanolamine (NAPE). However, the enzyme NAPE-PLD is also important for the synthesis of other NAEs that do not bind to CBRs, including *N*-oleoylethanolamine (OEA), *N*-palmitoylethanolamine (PEA) and *N*-stearoylethanolamine (SEA) (5). In fact, while AEA binds to the cannabinoid receptors CB1R, CB2R and the transient receptor potential vanilloid 1 (TRPV1) (8), OEA, PEA and SEA activate several non-cannabinoid receptors, including peroxisome proliferator-activated receptors (PPARα), GPR55 and GPR119 (9,10). Importantly, these NAEs are involved in the regulation of appetite and energy homeostasis through different mechanisms (11–15), notably by acting within the gastrointestinal tract (13,16), at the gut-brain interface (9,17,18) and/or directly in the brain (19–21). While NAPE-PLD expression, enzymatic activity and related byproducts have been detected in the mouse, rat and human brains (7,22–24), the key functions of this enzyme within the brain remain still elusive and not well-elucidated. In fact, most of the current literature has focused on distinct NAEs based on their transducing effectors (receptors, transcription factors) and not on the enzyme itself. Only recently, a few studies have shown that pharmacological blockade of NAPE-PLD activity (25) or selective

ablation of NAPE-PLD in stress-activated neurons (21) impaired limbic functions (fear

extinction, anxiety) mainly through the regulation of the hypothalamus-pituitary-adrenal

(HPA) axis, thus highlighting the importance of NAPE-PLD and its NAEs in brain functions.

Furthermore, NAPE-PLD silencing and consequent increased availability of its NAPE

substrate are neuroprotective in response to 6-OHDA-induced loss of dopamine (DA)-neurons

(20), suggesting an important role of NAPE-PLD in regulating physiological and cellular

functions of DA-neurons.

Moreover, the developmental invalidation of the Napepld gene alters very-long chain fatty

acids composition in the brain suggesting a more complex role for NAPE-PLD than

previously appreciated and the existence of alternative biosynthetic pathways (26,27).

Furthermore, the non-CB1R-related signaling engaged by fatty acid ethanolamines (FAEs)

produced by the specific enzymatic activity encoded by the Napepld gene (ID: 242864) points

toward a role for this enzyme in the generation of bioactive FAEs with a wide array of

functions including the regulation of energy balance (OEA), inflammation (PEA) or pain

sensitivity (SEA).

Interestingly, the presence of a single-nucleotide polymorphism (SNP) on the coding region

of the Napepld gene (rs17605251) has been associated with severe obesity (BMI $\geq$ 35 kg/m^2)

(28,29), suggesting a potential role for NAPE-PLD in the regulation of energy homeostasis,

food-motivated behaviors and metabolic disorders.

In the present study, we took advantage of several integrative *in vivo* approaches following

genetic and/or viral induction of tissues-specific deletion of NAPE-PLD in basal, food-

motivated and obese conditions. Within the mouse mesolimbic reward system, notably the

ventral tegmental area (VTA), we found that the enzyme NAPE-PLD functions as a fine-

tuning gatekeeper of reward events and dopamine dynamics as well as an important regulator

of energy homeostasis and metabolic efficiency in both physiological and pathological

(obesity) contexts.

Materials and methods

Animals

All experimental procedures were approved by the Animal Care Committee of the Université Paris Cité (CEB-06-2017, APAFiS #11033) and carried out following the 2010/63/EU directive. 8-20 weeks old transgenic male or female mice were used and housed in a room maintained at 22 +/-1 °C, with a light period from 7h00 to 19h00. Regular chow diet (3.24 kcal/g, reference SAFE® A04, Augy, France) and water were provided ad libitum unless otherwise stated. Diet-induced obesity (DIO) was achieved by exposing the mice to a 3-4 months period of high-fat diet (HFD, #D12492, Research Diets Inc., 5.24 kcal/g). The following transgenic mouse lines were used: Napepld$^{f/f}$ (18,30), Nes-Cre (31,32) [JAX003771, B6.Cg-Tg(Nes-cre)1Kln/J], Villin-Cre^{ERT2} (33) [JAX020282, B6.Cg-Tg(Vil1-cre/ERT2)23Syr/J] and Pgk-Cre (34) [JAX020811, B6.C-Tg(Pgk1-cre)1Lni/CrsJ]. All mouse lines were backcrossed and maintained as C57Bl6/J. Breeding-derived lines were generated and genotyped in house.

Drugs

GBR12909 (10 mg/kg, Tocris) and cocaine hydrochloride (15 mg/kg, Sigma-Aldrich) were dissolved in saline. Tamoxifen (MP Biomedicals) was suspended in ethanol (100 mg/ml, stock solution). A ready-to-use 10 mg/ml tamoxifen solution was prepared by adding filtered sunflower oil, followed by 30 min sonication and stored at 4 °C for up to 1 week. Tamoxifen solution was sonicated 5 min before administration and injected for 5 consecutive days (100 µl/day, i.p.).

Viral constructs

pAAV-hsyn-GRAB_DA2m was a gift from Yulong Li (Addgene plasmid #140553; http://n2t.net/addgene:140553; RRID:Addgene_140553).

pAAV.CMV.HI.eGFP-Cre.WPRE.SV40 was a gift from James M. Wilson (Addgene plasmid #105545; http://n2t.net/addgene:105545; RRID:Addgene_105545).

pAAV.CMV.PI.eGFP.WPRE.bGH was a gift from James M. Wilson (Addgene plasmid #105530; http://n2t.net/addgene:105530; RRID:Addgene_105530)

Stereotaxic surgery

For all surgical procedures, mice were rapidly anesthetized with isoflurane (3%, induction),

injected (ip) with the analgesic buprenorphine (Buprecare, 0.3 mg/kg, Recipharm, Lancashire,

UK) and ketoprofen (Ketofen, 10 mg/kg, France), and maintained under isoflurane anesthesia

(1.5%) throughout the surgery.

Stereotaxic surgery. Mice were placed on a stereotactic frame (David Kopf Instruments,

California, USA). Bilateral (AAV-GFP or AAV-Cre-GFP in the VTA, 0.2 µl/side) or

unilateral (GRAB-DA2m in the NAc, 0.3 µl) micro-injections were performed at the

following coordinates (in mm from bregma): VTA (L= +/-0.45; AP= -3.4, V= -4.3) and NAc

(L= -0.9; AP= +1.18, V= -4.4). All viruses were injected at the rate of 0.05 µl/min. Mice

recovered for at least 3-4 weeks after the surgery before being involved in experimental

procedures.

In vivo fiber photometry

For *in vivo* dopamine imaging (GRAB-DA2m, (35)), a chronically implantable cannula

(Doric Lenses, Québec, Canada) composed of a bare optical fiber (400 µm core, 0.48 N.A.)

and a fiber ferrule was implanted 100 µm above the location of the viral injection site in the

NAc. The fiber was fixed onto the skull using dental cement (Super-Bond C&B, Sun

Medical). Real time fluorescence was recorded using fiber photometry as described in

(36,37). Fluorescence was collected in the NAc using a single optical fiber for both delivery

of excitation light streams and collection of emitted fluorescence. The fiber photometry setup

used 2 light emitting LEDs: 405 nm LED sinusoidally modulated at 330 Hz and a 465 nm

LED sinusoidally modulated at 533 Hz (Doric Lenses) merged in a FMC4 MiniCube (Doric

Lenses) that combines the 2 wavelengths excitation light streams and separate them from the

emission light. The MiniCube was connected to a fiber optic rotary joint (Doric Lenses)

connected to the cannula. A RZ5P lock-in digital processor controlled by the Synapse

software (Tucker-Davis Technologies, TDT, USA), commanded the voltage signal sent to the

emitting LEDs via the LED driver (Doric Lenses). The light power before entering the

implanted cannula was measured with a power meter (PM100USB, Thorlabs) before the

beginning of each recording session. The light intensity to capture fluorescence emitted by

465 nm excitation was between 25-40 µW, for the 405 nm excitation this was between 10-20

µW at the tip of the fiber. The fluorescence emitted by the GRAB was collected by a

femtowatt photoreceiver module (Doric Lenses) through the same fiber patch cord. The signal

was then received by the RZ5P processor (TDT). On-line real time demodulation of the

fluorescence due to the 405 nm and 465 nm excitations was performed by the Synapse

software (TDT). Signals were exported to Python 3.0 and analyzed offline as previously

described (36). Data are presented as z-score of ΔF/F.

Metabolic efficiency analysis

Metabolic efficiency was measured as previously described (36). Briefly, mice were

monitored for whole energy expenditure (EE), O_2 consumption, CO_2 production, respiratory

exchange rate (RER=VCO_2/VO_2, V=volume), and locomotor activity using calorimetric cages

(Labmaster, TSE Systems GmbH, Bad Homburg, Germany). Gases ratio was determined

through an indirect open circuit calorimeter. This system monitors O_2 and CO_2 at the inlet

ports of a tide cage through which a known flow of air is ventilated (0.4 L/min) and regularly

compared to an empty reference cage. O_2 and CO_2 were recorded every 15 min during the

entire experiment. EE was calculated using the Weir equation for respiratory gas exchange

measurements. Food intake was measured with sensitive sensors for automated online

measurements. Calorimetric studies to investigate voluntary exercise-induced metabolic

adaptions were performed in metabolic cages equipped with running wheels (Promethion,

Sable Systems, Nevada, USA). Mice were monitored for body weight and composition at the

entry and exit of the experiments using an EchoMRI (Whole Body Composition Analyzers,

EchoMRI, Houston, USA). Data analysis was performed on Excel XP using extracted raw

values of VO_2 (ml/h), VCO_2 (ml/h), and EE (kcal/h).

Oral glucose tolerance test (OGTT)

Animals were fasted 6 hours before oral gavage of glucose (2 g/kg). Blood glucose was

measured from the vein blood tail using a glucometer (Menarini Diagnotics, Rungis, France)

at 0, 5, 10, 15, 30, 45, 60, 90, and 120 min. Blood samples were taken at 0, 15, 30 and 60 min

to measure insulin levels (mouse ultrasensitive insulin ELISA kit, ALPCO, Salem, USA).

Behaviors

Operant conditioning – Mice were food-restricted and maintained at 90% of their initial body

weight to facilitate learning and performance during the whole operant conditioning.

Computer-controlled operant conditioning was conducted in 12 identical conditioning

chambers (Phenomaster, TSE Systems GmbH, Bad Homburg, Germany) during the light

phase, at the same hour every day until the end of the procedure. Each operant wall had two

levers (one active and one inactive) located 3 cm lateral to a central pellet dispenser. The

reinforcer was a single 20-mg peanut butter flavored sucrose tablet (TestDiet, Richmond,

USA). Operant training was carried out daily with no interruption for 1h under a fixed-ratio 1

(FR1, 1 lever press=1 pellet). When the discrimination score between active and inactive

lever press (active lever presses/inactive lever presses) exceeded chance level, mice were

shifted to sessions under a FR5 (5 lever presses=1 pellet) and/or a progressive ratio (PR) [3

lever presses more for each subsequent reinforcer (r=3N+3; N=reinforcer number)].

Whenever of interest, PR was conducted in both food-restricted and sated mice.

Conditioned-place preference (CPP) – The CPP paradigm was performed during the light

phase either in food-restricted (maintenance at 90% of initial body weight) or normally fed

mice. All the compartments were cleaned before each conditioning session. Locomotor

activity was recorded with an infrared beam-based activity monitoring system and analyzed

with the provided software (Phenomaster, TSE Systems GmbH, Bad Homburg, Germany).

The least preferred compartment during the exploration phase was designated as the reward

(HFD)-baited compartment whereas the more preferred compartment as the chow-baited

compartment (biased protocol). Animals with more than 65% of preference for a

compartment on the pre-test day were removed. To reduce anxiety, during the first two days,

animals were carefully put in the middle of the apparatus and allowed to freely explore the

two compartments for 1h. The subsequent days included alternating conditioning sessions of

1h. After 8 days of conditioning [4 sessions in each compartment (chow and HFD)], animals

freely explored the two compartments for 30 minutes. The time spent in the reward-paired

compartment before *vs* after conditioning was the primary outcome variable (preference

score).

T-Maze – Mice were food-restricted (90 % of initial body weight) during the whole paradigm

and tested for learning and cognitive flexibility in a T-maze apparatus (arm 35-cm length, 25-

cm height, 15-cm width) (38). First, they were habituated to the apparatus (15 min of

exploration) for two consecutive days. Then, mice underwent a 5-days training protocol with

one arm reinforced with a palatable food pellet (HFD, cat #D12492, 5.24 kcal/g). Each mouse

was placed at the starting point and allowed to explore the maze by choosing one of the two

arms (reinforced and non-reinforced arms). The chosen arm was then blocked for 20 seconds

and the mouse replaced again in the starting arm. This process was repeated for 10 sessions

per day. At the end of this training period, cognitive flexibility and relearning processes were

assessed in a reversed learning task which consisted in exchanging the reinforced with the

non-reinforced arm. Again, mice underwent a 5-days training protocol (10 sessions/day).

Time-locked wheel running – Mice had access to a running wheel connected to an automatic revolution counter (Intellibio Innovation) during a limited amount of time (30 min per session, one session per day) during 5 consecutive days.

Food preference and choice – Mice were tested for food choice and preference by using non-caloric and/or caloric solutions. Notably, they were exposed to graduated small bottles containing either water, sucralose (2 mM), sucrose (10% w/v) or intralipids 20%. During different days of exposure (1h session with free choice between 2 bottles), preference was measured by comparing the consumption of sucralose *vs* water, sucrose *vs* water and lipids *vs* water.

GBR-induced locomotor activity – Locomotor activity induced by GBR12909 (10 mg/kg, i.p.) was recorded in an automated online measurement system using an infrared beam-based activity monitoring system (Phenomaster, TSE Systems GmbH, Bad Homburg, Germany).

Tissue preparation and immunofluorescence

Mice were anaesthetized with pentobarbital (500 mg/kg, i.p., Sanofi-Aventis, France) and transcardially perfused with cold (4°C) PFA 4% for 5 minutes. Sections were processed and confocal imaging acquisitions were performed as previously described (37,39). GFP staining was not antibody-amplified (AAV-Cre-GFP, AAV-GFP and GRAB-DA2m). The following primary antibodies were used: rabbit anti-TH (1:1000, Merck Millipore, #AB152) and rabbit anti-cFos (1:500, Synaptic Systems, #226003). Quantification of cFos-immunopositive cells was performed using the cell counter plugin of ImageJ taking a fixed threshold of fluorescence as standard reference.

Lipidomics

Tissue extracts and HPLC/MS/MS were performed as previously described (40). In brief, samples were placed in 50 volumes of HPLC-grade methanol then spiked with 500 pmols deuterium-labeled N-arachidonoyl glycine (d8NAGly; Cayman Chemical, Ann Arbor, MI) as an internal standard to determine extraction efficiency. Samples were placed on ice in darkness for 2 hours then individually homogenized. Homogenates were then centrifuged at 19,000g for 20 minutes at 20°C. Supernatants were decanted and diluted with HPLC H_2O to make a 75:25 water to supernatant solution. Partial purification was achieved using C-18 solid

293   phase extraction columns (Agilent Technologies, Lake Forest, CA). A series of 4 elutions
294   with 1.5 mL of 60%, 75%, 85%, and 100% methanol were collected for analysis. Samples
295   were analyzed using an Applied Biosystems API 3000 triple quadrupole mass spectrometer
296   with electrospray ionization. 20μL from each elution were chromatographed using an XDB-
297   C18 reversed phase HPLC analytical column (Agilent) and optimized mobile phase gradients.
298

299   **Reclustering and transcriptomics meta-analysis**
300   Publicly available transcriptomic data were downloaded from Gene Expression Omnibus
301   (https://www.ncbi.nlm.nih.gov/geo/, GSE137763, GSE168156, GSE64526, GSE114918) and
302   analyzed using a Python 3.0 pipeline generated in line with the original publications.
303

Statistics
All data are presented as mean ± SEM. Statistical tests were performed with Prism 7
(GraphPad Software, La Jolla, CA, USA). Detailed statistical analyses are listed in the **Suppl.**
Table 1. Normality was assessed by the D'Agostino-Pearson test. Depending on the
experimental design, data were analyzed using either Student's t-test (paired or unpaired) with
equal variances, one-way ANOVA or two-way ANOVA. The significance threshold was
automatically set at $p<0.05$. ANOVA analyses were followed by Bonferroni post hoc test for
specific comparisons only when overall ANOVA revealed a significant difference (at least
$p<0.05$).

Results

NAPE-PLD is functionally expressed in the brain and mediates motivational food-responses.

In order to explore the role of the NAEs-synthetizing enzyme NAPE-PLD in food-motivated behaviors, we first addressed the consequence of whole-body NAPE-PLD developmental knock-out in a food-reward seeking behavioral paradigm. Here, we used the Napepld$^{f/f}$ mouse line in which the two LoxP sites span the exon 3 (24,30,41), the gene sequence that encodes for the catalytic activity of the enzyme and that efficiently leads to a reduction of NAPE-PLD-derived bioproducts (18,24,30,41). Napepld$^{f/f}$ mice were bred with mice expressing Cre under the pan-promoter phosphoglycerate kinase 1 (Pgk-Cre) which result in whole-body full knock-out (KO, (34)) of NAPE-PLD in subsequent generations. To study the reinforcing and motivational properties of food, we performed an operant conditioning paradigm where animals were trained, under different schedules, to press a lever to obtain a palatable sugar pellet. In both a fixed ratio 1 schedule (FR1, 1 lever press for 1 sugar pellet during 4 daily sessions) or a progressive ratio schedule (PR), which assesses the motivational component of reinforcement behaviors, Napepld$^{+/+}$ (controls) and Napepld^{KO} mice displayed similar performances (**Suppl. Fig. 1A, B**). These results suggest that either physiological compensations occurred during development, as reported by previous genetic invalidations (27), and/or that, despite the expression of NAPE-PLD in the brain (23), brain NAPE-PLD plays a marginal role in reward-seeking behavior. To disentangle these two hypotheses, we moved to brain-restricted ablation of NAPE-PLD. Ablation of NAPE-PLD in the central nervous system (CNS) was achieved by crossing Napepld$^{f/f}$ mice with mice expressing Cre under the control of the promoter Nestin (Nes$^{Cre+/-}$ mice) (31,32). We observed that CNS genetic deletion of NAPE-PLD was associated to an enhanced response to operant behavior. In fact, under FR1 schedule, Napepld$^{\Delta CNS}$ mice (males and females) collected a higher number of pellets and had a higher number of active lever presses (**Fig. 1A, A^1**). However, this enhanced reward-like phenotype was not related to differences in learning (% of active lever over inactive lever) as both genotypes were characterized by very similar discrimination scores (**Fig. 1A^2**). Once the operant conditioning established, mice were moved to the PR schedule. Again, we noticed that, despite similar learning scores, Napepld$^{\Delta CNS}$ mice showed enhanced performances (number of rewards and active lever presses) compared to control mice (**Fig. 1B, B^1, B^2**). To exclude that such phenotype was driven by the presence of Cre (Nes$^{Cre+/-}$ (32,42)) rather than the proper deletion of NAPE-PLD, we performed the same

behavioral battery in Nes$^{Cre-/-}$ (controls) and Nes$^{Cre+/-}$ mice which both displayed a very

similar phenotype on this paradigm (**Suppl. Fig. 1C, D**), thus indicating that genetic deletion

of neuronal NAPE-PLD is responsible for the enhanced reward-behavior observed in

Napepld$^{\Delta CNS}$ mice. This result revealed that tissue-specific ablation of NAPE-PLD generates

different outcomes than whole-body gene deletion. This finding, which points to an effective

role of brain NAPE-PLD in food-reward seeking behaviors, also raises the possibility that the

contribution of NAPE-PLD in multiple organs (full KO mice) might lead to physiological

adjustments eventually driving opposite consequences on a particular behavioral output with

an overall mitigated consequence.

Among the main organs that might contribute to food-dependent reward processes, the gut has

emerged as a critical modulator of reinforced behaviors (3,36,43,44). It has been previously

shown that mice with a specific and inducible deletion of NAPE-PLD in the intestinal

epithelial cells (IEC) (Napepld$^{\Delta IEC}$) exhibited a phenotype associated with specific changes in

the homeostatic regulation of food intake and altered metabolic adaptations to high-fat diet

(18,45). Therefore, we explored the potential contribution of intestinal NAPE-PLD in reward-

seeking behavior. Interestingly, Napepld$^{\Delta IEC}$ and control mice showed comparable

performances in the operant conditioning paradigm (**Suppl. Fig. 1E, F**), indicating that,

while intestinal NAPE-PLD is critical for metabolic control (18) and short-term regulation of

food intake (45), brain NAPE-PLD might represent a more direct target as acute regulator of

food-reward behaviors.

Reinforced behaviors tightly depend on key brain regions that constitute the reward system,

notably the midbrain dopamine (DA)-producing ventral tegmental area (VTA) and its

dopaminoceptive structures, including the dorsal striatum (DS)/nucleus accumbens (NAc), the

prefrontal cortex (PFC) and the hippocampus (Hippo) (46). Therefore, we first investigated

whether NAPE-PLD-produced NAEs were altered within these structures in Napepld$^{\Delta CNS}$

mice. Lipidomic analyses revealed a significant decrease of several NAEs species [AEA,

OEA, PEA, *N*-stearoylethanolamine (SEA), *N*-linoleoylethanolamine (LEA) and *N*-

docosahexaenoylethanolamine (DEA)] in the midbrain VTA following CNS deletion of

NAPE-PLD (**Fig. 1C-H**). Interestingly, either no major differences (DS/NAc and PFC),

specific significant reductions (SEA, LEA and DEA for the hippocampus) or trends of

decrease were detected in the levels of NAEs in the other reward-associated brain regions

(**Fig. 1C-H**). Of note, no alterations were detected for the endocannabinoid (eCB) 2-AG (**Fig.

1I**). Moreover, within the VTA we did not detect alterations in the levels of other fatty acids

(linoleic, arachidonic and oleic acids) (**Suppl. Fig. 2A-C**) or *N*-acylamides (**Suppl. Table 2**),

thus indicating that lack of NAPE-PLD specifically affects a subset of endogenous bioactive

lipids.

In addition, lipidomic analyses also revealed that NAEs, but not 2-AG (**Fig. 1I**), levels were

higher in the VTA compared to the DS/NAc, PFC and hippocampus (**Fig. 1C-H**).

These biochemical results, associated to the enhanced phenotype of Napepld$^{\Delta CNS}$ mice in the

operant reward-driven behavior (**Fig. 1A, B**), prompted us to investigate the structure- and/or

cell type-specific expression of NAPE-PLD in both rodents (rat and mouse) and human

brains. First, by taking advantage of single-nucleus RNA transcriptomics (snRNA-seq) in the

rat NAc (47) (GSE137763) and VTA (48) (GSE168156), we performed a clustering meta-

analyses of transcripts encoding for eCBs- and NAEs-producing enzymes (*Napepld*, *Dagla*,

Daglb) as well as for eCBs- and NAEs-related transducing effectors (*Cnr1*, *Trpv1*, *Gpr119*,

Ppara, *Pparg*). Compared to *Dagla* and *Daglb*, we observed that accumbal *Drd1*- and *Drd2*-

medium spiny neurons (MSNs) expressed low levels of *Napepld* (6% of 2819 *Drd1*-MSNs

and 5% of 1993 *Drd2*-MSNs, respectively) (**Fig. 2A, B**). On the contrary, we detected a

higher expression of *Napepld* in VTA-neurons (**Fig. 2C-E**). Since the VTA harbors different

neuronal cell types (49), we restricted our meta-analysis to VTA DA-, GABA- and glutamate

(Glut)-neurons. In the VTA, we again detected higher levels of *Dagla* and *Daglb* but,

interestingly, we observed that *Napepld* was mainly present in VTA DA- and Glut-neurons

[12% of 399 DA-neurons (**Fig. 2C**) and 20% of 698 Glut-neurons (**Fig. 2E**), respectively],

whereas 9% of GABA-neurons (896 cells) were positive for *Napepld* (**Fig. 2D**). Interestingly,

this pattern of *Napepld* expression mirrors the higher levels of NAEs observed in the VTA

compared to the DS/NAc (**Fig. 1C**). Moreover, the transcriptomic profiling of eCBs- and

NAEs-producing enzymes was in line with the meta-analysis of bulk VTA transcriptomics in

the murine (50) (*Slc6a3*-bacTRAP mice, GSE64526, **Fig. 2F**) and human midbrains (51)

(GSE114918, **Fig. 2G**).

Altogether these observations underline the potential role for NAPE-PLD in the midbrain

VTA as a regulator of food-associated reward processes.

VTA NAPE-PLD scales food-motivated behaviors and dopamine releasing dynamics

To precisely interrogate the structure-specific functions of NAPE-PLD in driving food-

motivated behaviors, we knocked-down the Napepld gene in the VTA using a local and

virally mediated delivery of Cre in the VTA of Napepld$^{f/f}$ mice (**Fig. 3A**). Next, we tested the

reinforcing and motivational properties of palatable food using a food-dependent operant

conditioning paradigm. In line with the results obtained with Napepld$^{\Delta CNS}$ mice (**Fig. 1A, B**),

we observed that viral deletion of NAPE-PLD in the VTA promoted food-operant conditioning (increased number of rewards and active lever presses) during both FR1 and PR schedules (**Fig. 3B, C**), with no differences in learning performances as both groups showed similar active/inactive discrimination index (**Fig. 3B, C**).

This enhanced reward phenotype was also present following a FR1→FR5→PR training schedule (**Suppl. Fig. 3A-C**, food restriction) and even in sated conditions (**Suppl. Fig. 3D**), therefore excluding the potentially confounding effect of hunger onto motivational drive. Importantly, this phenotype was also confirmed in female mice (**Suppl. Fig. 3E-H**), again in both food-restricted and sated conditions. Of note, in both males and females, no significant differences in initial body weight and body weight loss (food restriction) were observed between experimental groups (**Suppl. Fig. 4A, B**).

Aside from the motivational component, the liking and learning components of feeding are an integral part of food-reward processes (46). These components can be assessed through behavioral measurements of the positive valence assigned to palatable food in the conditioned-place preference (CPP, **Fig. 3D**) and T-maze (**Fig. 3E**) paradigms, which both rely on the association between reward value and context. In food-restricted conditions, we observed an increased and similar CPP score in both $Napepld^{VTA-GFP}$ and $Napepld^{\Delta VTA}$ mice (**Fig. 3F**). However, in sated conditions, only $Napepld^{\Delta VTA}$ mice showed an HFD-induced increase in CPP score (**Fig. 3G**), indicating enhanced susceptibility to the reinforcing properties of palatable foods. Using the T-maze paradigm, we next assessed the ability and flexibility of mice to actively learn in discriminating between a rewarded (HFD) and a non-rewarded arm. During the learning phase (first 5 days), we observed that both groups showed a progressive increase in correct responses (%) over training days, with $Napepld^{\Delta VTA}$ mice performing significantly better than $Napepld^{VTA-GFP}$ control mice (**Fig. 3H**). Then, mice were tested for their flexibility to relearn the task under a reversal learning schedule (in which the food reinforcer was switched to the previously unreinforced arm of the T-maze). While both groups displayed good performance in learning/flexibility, VTA-specific deletion of NAPE-PLD resulted in a better performance with a more rapid acquisition of the correct entry into the reinforced arm as compared to $Napepld^{VTA-GFP}$ control mice (**Fig. 3H**).

Next, we investigated the role of VTA NAPE-PLD in driving palatable food preference during a time-locked window (1h of exposure). First, we tested the reinforcing properties of the non-caloric sweetener sucralose (2 mM). As shown in **Fig. 3I**, $Napepld^{\Delta VTA}$ mice consumed more sucralose than $Napepld^{VTA-GFP}$ control mice. A very similar pattern of

enhanced preference was measured with the natural caloric sugar sucrose (10%, **Fig. 3J**) and

with emulsified lipids (20%, **Fig. 3K**).

We therefore decided to investigate whether the enhanced reward-like behavior observed in

Napepld$^{\Delta VTA}$ mice was associated to an increased activity of the nucleus accumbens (NAc), a

region highly innervated by VTA projections and whose activity is correlated with food-

reward processes (46). However, the enhanced neural response within the reward system

might result either from the higher tropism/consumption of palatable food of Napepld$^{\Delta VTA}$

mice or from the increased rewarding value despite a fixed amount of food-reinforcer. In

order to dissociate these two possibilities, we exposed our experimental groups (sated

conditions) to an equal amount of HFD during a time-locked window (1h during which all

mice consumed the HFD pellet) and then we analyzed the induction of cFos, a molecular

proxy of neuronal activity, in the NAc (**Fig. 3L**). Interestingly, we detected more cFos-

positive neurons in the NAc of Napepld$^{\Delta VTA}$ mice (**Fig. 3L**), thereby indicating an enhanced

responsiveness of the VTA→NAc mesolimbic axis to an equal amount of food-reward

consumption.

These results led us to hypothesize that VTA NAPE-PLD and its local NAEs bioproducts may

contribute to the regulation of DA dynamics within the VTA→NAc mesolimbic axis. To test

this hypothesis, we took advantage of *in vivo* fiber photometry coupled to virally expressed

DA biosensors (GRAB-DA2m (35)) to measure DA dynamics in the NAc of Napepld$^{VTA\text{-}GFP}$

and Napepld$^{\Delta VTA}$ mice (**Fig. 4A, B**). First, we observed that exposing both fasted (**Fig. 4C, D**)

and *ad libitum* fed mice (**Fig. 4E, F** and **Suppl. Fig. 5A**) to HFD triggered a higher DA

accumulation/release in the NAc of Napepld$^{\Delta VTA}$ mice compared to control animals.

Second, to further explore whether and how NAPE-PLD may contribute to the regulation of

DA-dependent events, we tested *in vivo* DA dynamics also in two non-food-dependent

paradigms: the administration of cocaine (**Fig. 4G, H**) and the tail suspension (TS, **Fig. 4I, J**

and **Suppl. Fig. 5B**). In both cases, we observed that Napepld$^{\Delta VTA}$ mice were characterized by

an enhanced accumulation/release of DA in the NAc than Napepld$^{VTA\text{-}GFP}$ mice. Lastly , we

administered the selective DAT blocker GBR12909 and noticed an enhanced locomotor

response in Napepld$^{\Delta VTA}$ mice compared to controls (**Suppl. Fig. 5C**), further confirming an

amplified DA release/tone as a consequence of VTA NAPE-PLD knock-down.

Overall, these results indicate that VTA NAPE-PLD tightly contributes in orchestrating the

responses of midbrain DA-neurons to both food- and non-food-related reinforcers by

promoting and boosting the release of DA at VTA→NAc synapses.

VTA NAPE-PLD contributes to the regulation of food intake and energy homeostasis

Although the regulation of energy homeostasis has been classically ascribed to the hypothalamus and the brainstem (1), new evidence indicates that the reward system also strongly contributes in scaling whole-body metabolic functions (38,52). We therefore explored the metabolic consequences of VTA NAPE-PLD knock-down in the regulation of whole-body metabolic efficiency and peripheral substrates utilization by using longitudinal measurements of indirect calorimetry. As previously observed (**Suppl. Fig. 4**), no major differences were observed in body weight and body composition between the two experimental groups (**Fig. 5A**). However, Napepld$^{\Delta VTA}$ mice displayed a spontaneous increase in locomotor activity and in cumulative food intake compared to Napepld$^{VTA-GFP}$ mice during both the light and dark circadian phases (**Fig. 5B, C**). These phenotypes were associated with an overall enhanced energy expenditure (**Fig. 5D**) and to a change in peripheral substrates utilization favoring carbohydrates over lipids-based substrates as indicated by the increase in respiratory exchange ratio (RER, 1=glucose substrate, 0.7=lipids substrate) during the light phase (**Fig. 5E**) and the consequent decrease in fatty acid oxidation (FAO) in Napepld$^{\Delta VTA}$ mice during both the light and dark phases (**Fig. 5F**). This feature was also associated with enhanced glucose tolerance during an oral glucose tolerance test at the expense of lower insulin release, suggesting enhanced whole-body glucose dynamics and insulin sensitivity (**Suppl. Fig. 6**).

We then decided to investigate how Napepld$^{\Delta VTA}$ mice adapted during manipulation of nutrients availability. We noticed that during a food deprivation period (overnight fasting) Napepld$^{\Delta VTA}$ mice were still characterized by increased locomotor activity (**Fig. 5G**) and energy expenditure (**Fig. 5I**), but with no differences in RER (**Fig. 5J**). While the capability to mobilize lipids-based substrates during the fasting-induced lipolysis was similar between the two groups, as indicated by the RER (**Fig. 5J**), the proportion of lipids used a primary source of fuel was enhanced in the fasting period as indicated by the FAO (**Fig. 5K**), thus indicating a metabolic shift toward lipids-based substrates utilization. Interestingly, upon refeeding, mice displayed similar food intake (**Fig. 5H**), locomotor activity (**Fig. 5G**) or substrates utilization (**Fig. 5J, K**), while a slight increase in energy expenditure was still detected in Napepld$^{\Delta VTA}$ mice (**Fig. 5I**). These results confirmed the hypothesis that the integrity of NAPE-PLD within the VTA was required for the proper metabolic adaptation to changes in nutrients availability.

Since fasting increases the motivational drive and responsiveness to food, we wondered

whether the expression of NAPE-PLD was required to promote DA releasing dynamics in

fasted mice exposed to a chow pellet. In contrast to the acute response to palatable HFD (**Fig**

4C-F), consumption of a chow pellet resulted in similar DA releasing dynamics in both

Napepld$^{VTA-GFP}$ and Napepld$^{\Delta VTA}$ mice (**Fig. 5L, M**). This may suggest that (*i*) the action of

VTA NAPE-PLD in the modulation of adaptive metabolic responses to nutritional

manipulations can be dissociated from DA release in the fast-refeeding transition and/or (*ii*)

VTA NAPE-PLD plays an active role in discriminating between palatable (HFD, **Fig 4C-F**)

and regular (chow, **Fig. 5L, M**) foods through the control of reward-dependent DA dynamics.

VTA NAPE-PLD does not contribute to exercise-motivated behaviors but still regulates

energy homeostasis

In mammals, exercise can function as a rewarding/motivational stimulus (53) and the eCBs

system, especially within the VTA, has been identified as a key regulator of exercise-induced

reinforced behaviors (54–56). We thus decided to extend our investigations to exercise-

motivated behaviors (*i*) to investigate whether VTA NAPE-PLD was also important in

mediating the reinforcing properties of exercise and (*ii*) to study whether metabolic

adaptations observed in Napepld$^{\Delta VTA}$ mice (**Fig. 5**) were solely dependent on enhanced

locomotor activity.

First, we performed a time-locked access (30 min session/day) to a running wheel. Despite

both Napepld$^{VTA-GFP}$ and Napepld$^{\Delta VTA}$ mice progressively spent more time wheel-running, we

surprisingly noticed a reduced performance in Napepld$^{\Delta VTA}$ mice compared to control animals

(**Fig. 6A, A^1**). This led us to investigate whole-body metabolism and metabolic efficiency in

calorimetric chambers equipped with running wheels.

Again, we observed that Napepld$^{\Delta VTA}$ mice were characterized by an enhanced spontaneous

locomotor activity (**Fig. 6B, B^1**) and reduced wheel-running activity (**Fig. 6C, C^1**). When

combining both forms of activity (spontaneous + wheel running activities), we detected no

differences in the light phase and a lower global activity in Napepld$^{\Delta VTA}$ mice during the dark

phase (**Fig. 6D**). Of interest, the peculiar metabolic signature associated with VTA NAPE-

PLD deletion also remained in this exercise-based paradigm and was characterized by

enhanced energy expenditure (**Fig. 6E, E^1**), food intake (**Fig. 6F**) and RER (**Fig. 6G**), and

lower FAO (**Fig. 6H**) in Napepld$^{\Delta VTA}$ mice.

Altogether, these results indicate that the role of VTA NAPE-PLD in regulating reward-like

processes cannot be generalized to all natural rewards (food *vs* exercise) and that the

metabolic adaptations observed in VTA NAPE-PLD-deleted mice are not solely dependent on

locomotor activity.

VTA NAPE-PLD controls metabolic adaptation to obesogenic environment.

Food-reward drive, together with changes in metabolic outputs in response to food

environment, are important contributors of the obesity pandemics. Given the above-

mentioned results showing a key role of VTA NAPE-PLD in controlling reward and

metabolic processes, we hypothesized that NAPE-PLD may influence the (mal)adaptive

responses to an obesogenic environment. Thus, Napepld$^{\text{VTA-GFP}}$ and Napepld$^{\Delta\text{VTA}}$ mice were

chronically exposed to an obesogenic diet (3 months of HFD) and then metabolically

characterized.

First, we noticed no significant differences in the body weight and lean mass composition of

both HFD-exposed experimental groups (**Fig. 7A**). However, fat body mass was significantly

lower in Napepld$^{\Delta\text{VTA}}$ mice (**Fig. 7A**). The analysis of metabolic efficiency revealed that

obese Napepld$^{\Delta\text{VTA}}$ mice displayed increased nocturnal locomotor activity (**Fig. 7B**) and

enhanced nocturnal food intake (**Fig. 7C, C^{1}**). Surprisingly, we detected a higher energy

expenditure (**Fig. 7D, D^{1}**) and FAO (**Fig. 7E, E^{1}**) in obese Napepld$^{\Delta\text{VTA}}$ mice, whereas the

RER resulted unchanged (**Fig. 7F**). This metabolic blueprint suggests that, depending on diets

(chow *vs* HFD) and metabolic profiles (lean *vs* obese), VTA NAPE-PLD readily allows the

plastic adaptation of nutrients partitioning (**Fig. 5E-F** *vs* **Fig. 7E-F**) in order to maintain a

higher energy expenditure.

These results indicate that, within the VTA, the deletion of NAPE-PLD partially protects

against diet-induced obesity.

Discussion

NAEs represent an important family of endogenous bioactive lipids involved in several biological process including adaptive stress responses and emotional states (21,57), pain (58), inflammation (59), feeding and metabolism (9,14,17,18,45). In an effort to characterize the phospholipase D (PLD)-mediated enzymatic activity that converts NAPE into NAEs, *in vitro* studies identified NAPE-PLD as able to produce NAE derivatives from NAPE (7). Genetic invalidation of the Napepld gene (27,30) revealed that, in the brain, several alternative enzymatic pathways exist for the synthesis of polyunsaturated fatty acid NAEs while NAPE-PLD activity seems more critical for saturated/monounsaturated NAEs, with a drastic decrease of these compounds escalating with carbons chain length. Later, in depth lipidomic analysis revealed a broader role for NAPE-PLD with a large spectrum and region-specific consequences in lipidome alteration in the brain (24,30). While constantly evolving, the sensitivity and limitations of quantitative methods to measure NAEs may be in part implicated in the difficulty to formerly assigned a definitive set of substrates and bioproducts to NAPE-PLD. It is now clear that NAPE-PLD bioproducts include important lipid mediators which, by acting on a variety of transcriptional and signaling cascades, control cellular and physiological responses (5).

In the present study we explored the role of NAPE-PLD in food- and reward-driven behaviors. Whole-body deletion of NAPE-PLD did not affect food-reward operant conditioning, suggesting that compensatory mechanisms, as previously reported with a model of developmental NAPE-PLD KO mice (27), may be at play. However, using neural-specific genetic deletion (Nestin-Cre) or midbrain-specific viral invalidation approaches we revealed that the integrity of NAPE-PLD within the midbrain VTA is required for NAEs synthesis (AEA, OEA, PEA, SEA, LEA and DEA) and that NAPE-PLD acts as a gatekeeper for fine-tuning food-reward behaviors and for regulating, at least as a contributor, energy balance and whole-body metabolism. In fact, viral down-regulation of NAPE-PLD in the VTA was associated with an enhanced tropism towards palatable foods and a stronger conditioning for food-related rewards (operant conditioning, conditioned-place preference and T-maze paradigm). These phenomena were associated with an enhanced activity of midbrain VTA DA-neurons and their DA release dynamics in response to food rewards and also to non-food-related stimuli. Consistent with the notion of a region-specific biosynthesis and action of NAEs, RNA-seq meta-analyses revealed low levels of *Napepld* in postsynaptic striatal

dopaminoceptive neurons (*Drd1*- and *Drd2*-MSNs), but a higher expression of *Napepld* in

VTA-neurons, notably DA- and glutamate-neurons.

Previous studies focusing on nicotine reinforcement and tobacco use disorder (TUD) have

shown that pharmacological inhibition of the fatty acid amid hydrolase (FAAH), one of the

main enzymes responsible for the degradation of NAEs, reduces nicotine-enhanced DA

transmission and nicotine reinforcement (19,60) through the activation of PPARα by

OEA/PEA and the activation of intracellular cascades leading to the reduction of nicotinic

receptors onto midbrain DA-neurons (19,60). In line with our results showing increased

dopaminergic VTA→DA transmission following downregulation of VTA NAPE-PLD, these

electrophysiological studies have clearly demonstrated that OEA/PEA inhibit DA-neurons

whereas inhibition of PPARα promotes their spontaneous activity (19,61,62). However, these

aforementioned reports, although seminal, did not formally test nor established the role of

midbrain NAPE-PLD in these processes. In our hands, specific knock-down of NAPE-PLD in

the VTA resulted in enhanced DA transmission in response to reward stimuli. In that view,

our results are perfectly in line with a putative role for NAPE-PLD-derived substrates as

negative modulators of VTA DA-neurons. While our study provides an additional mechanism

by pointing at NAPE-PLD has a potential candidate, it also extends the role of this enzyme to

the control of DA-dependent behaviors and DA releasing dynamics in response to reward

stimuli well-beyond nicotine. Although our results do not rule out whether the NAPE-

PLD/NAPE→NAEs machinery controls both the tonic and phasic DA release (63–65), they

clearly indicate that VTA NAPE-PLD activity, through the synthesis of NAEs, is an integral

component of the control of DA-neurons' activity. Whether the cellular accumulation of

NAPE and/or the decreased levels of NAEs are the primary responsible for the changes in

DA-dependent behaviors and DA releasing dynamics is still unknown. Indeed, NAPE-PLD

silencing seems to confer a protective action for increased NAPE species in 6-OHDA-induced

neural damage (20), suggesting that the regulation of DA-neurons might be linked to

NAPE/NAE membrane homeostasis. Given the multiple roles of endogenous bioactive lipids,

it is possible that the invalidation of NAPE-PLD in the VTA may lead to an imbalance in the

NAPE/NAEs ratio with ultimate consequences on a variety of processes including heightened

DA responses to reward stimuli. In addition, NAEs, either cannabinoids-like (AEA) and non-

cannabinoids-like (OEA, PEA), may be released and act through several *modus operandi*. In

fact, while the eCB AEA is retrogradely released *on demand* (66), the non-eCB NAEs may

act anterogradely as suggested by the presence of NAPE-PLD also in pre-synaptic terminals
(67), thus potentially modulating both intra-VTA microcircuits and/or VTA-projecting
circuits (*i.e.* VTA→NAc). However, the VTA does not only harbor DA-neurons (49,68).
Importantly, the relatively high expression of NAPE-PLD in VTA Glut-neurons may also
indicate a possible regulatory control of this cell type by the NAPE/NAEs ratio in the
observed behavioral and metabolic features. This regulation may be exerted either through the
local communication among all VTA-neurons or even through VTA^{Glut}→NAc projections. In
fact, these glutamatergic projections were recently shown to promote reinforcement
independently of DA release (69) and we cannot exclude that NAPE-PLD-dependent
mechanisms onto these complex neural networks may result from this additional form of
VTA→NAc communication in encoding changes in reward-driven behaviors and metabolic
efficiency. Despite this limitation, our *in vivo* imaging results clearly reveal that the
VTA→NAc dopaminergic transmission is regulated by the NAPE-PLD. Indeed, further
studies are warranted to fully flush out how and to which extent midbrain NAPE-PLD
regulates DA events by selectively focusing on the local interconnectivity and
interdependency of VTA DA-, Glut- and GABA-neurons.

While our study establishes a direct role for NAPE-PLD in the mesolimbic reward circuit in
regulating DA release and DA-dependent behaviors, it also unveils the functional connection
between VTA NAPE-PLD activity and the control of whole-body metabolism. NAPE-
PLD$^{\Delta VTA}$ mice displayed increased spontaneous locomotor activity in both fed and fasting,
but not refed, conditions, which is consistent with an enhanced activity of VTA DA-neurons
(70). These features were associated with increased cumulative food intake and whole-body
energy expenditure and, together with other recent studies (52,71,72), they point to the VTA
as an important regulator of energy balance and metabolic efficiency. In fact, on chow diet the
overall body weight was only marginally affected in NAPE-PLD$^{\Delta VTA}$ compared to control
mice, suggesting that increased energy expenditure (EE) was compensated by increased
energy intake in a closed and well-balanced homeostatic regulation (**Fig. 5**). During food
deprivation, NAPE-PLD$^{\Delta VTA}$ mice showed a drastic increase in fasting-induced foraging
pointing towards a change in adaptive strategy in response to decreased nutrients availability.
This increased activity was associated with increased EE and most likely fueled by enhanced
lipids-based metabolism (**Fig. 5**). In mice exposed to HFD, a similar increase in spontaneous
activity, EE and FAO was observed together with increased food intake. Since NAPE-

PLD$^{\Delta VTA}$ mice show increased tropism and responsiveness to HFD and palatable food it is

possible that this increase in HFD intake may be the result of enhanced palatability for

energy-dense foods. An alternative, and not mutually exclusive, hypothesis would be that

increased food intake is a homeostatic mechanism to sustain enhanced EE. Although we do

not provide a clear evidence to disentangle these two hypotheses, the fact remains that NAPE-

PLD knock-down in the VTA confers a protective phenotype against HFD-induced body fat

gain and metabolic alterations (**Fig. 7**). This result nicely echoes a study in humans that has

identified a common haplotype of the Napepld gene in severe obesity (28). In face of our

results, one should consider that VTA NAPE-PLD deletion led to a protective effect against

HFD-mediated fat mass gain and metabolic (mal)adaptations but was accompanied by

enhanced reward-driven behaviors. This is important from a translational point of view as

obesity is characterized by an alteration of peripheral and central eCBs and NAEs in humans

(73–75). Our current results, together with those depicting the role of NAPE-PLD in

peripheral tissues, culminate in the elaboration of a complex picture in which organs- and

regions-specific homeostasis of NAPE/NAEs underline the complexity by which NAPE-PLD

exerts its control onto reward-dependent behaviors, energy balance and body weight control.

Indeed, it has been previously shown that mice lacking NAPE-PLD specifically in adipocytes

displayed spontaneous obesity, higher fat mass, glucose intolerance, and lower adipocyte

browning (41). Moreover, mice lacking NAPE-PLD in the intestinal epithelial cells were

more sensitive to HFD-induced body weight gain, fat mass gain and hepatic steatosis (18), a

phenomenon partially explained by an alteration in food intake behavior (45).

In conclusion, our study provides a direct evidence for a key role of NAPE-PLD in the

control of reward-dependent behaviors, DA dynamics and energy metabolism. The main

limitation of this study primarily lies in the lack of a clear cell type-specific identification of

cellular and molecular mechanisms occurring within the heterogenous VTA. Given the

complexity of NAEs action and the variety of bioactive lipids and signaling cascades

associated with NAPE-PLD activity, further research and new investigatory tools will be

required to fully apprehend the role of NAPE-PLD and its bioproducts in promoting anti-

obesity strategies.

Acknowledgments: We thank Olja Kacanski for administrative support, Isabelle Le Parco,
Aurélie Djemat, Daniel Quintas, Magguy Boa, Ludovic Maingault and Angélique Dauvin for
animals' care, and Florianne Michel for genotyping. We acknowledge the technical platform
Functional and Physiological Exploration platform (FPE) of the Université Paris Cité, CNRS,
Unité de Biologie Fonctionnelle et Adaptative, the viral production facility of the UMR
INSERM 1089 and the animal core facility "Buffon" of the Université Paris Cité/Institut
Jacques Monod.

Funding: This work was supported by the Nutricia Research Foundation (#2022-E7), *Agence*
Nationale de la Recherche (ANR-21-CE14-0021-01, ANR-19-CE37-0020-02), *Fondation*
pour la Recherche Médicale (Équipe FRM #EQU202003010155), *Fédération pour la*
Recherche sur le Cerveau and *Association France Parkinson*, the Modern Diet and
Physiology Research Center (MDPRC), Université Paris Cité and CNRS.
G.L. is supported by a China Scholarship Council (CSC) fellowship. O.O is supported by an
FRM fellowship. We are grateful to Dr. Sylvie Robin for the gift of the Villin-Cre[ERT2] mice.
P.D.C. is recipient of grants from FNRS (FRFS-WELBIO: WELBIO-CR-2022A-02, EOS:
program no. 40007505) and La Caixa (NeuroGut). A.E. is research associate at FNRS and the
recipient of grants from FNRS (FRFS-WELBIO: WELBIO-CR-2019S-03R, FNRS:
J.0075.22).

Author contributions: Conceptualization: G.G and S.L.; **Investigation**: J.C., G.L., O.O.,
A.E., E.L., H.B., K.M., P.D.C., S.L., G.G.; **Formal Data Analysis**: J.C., S.L, G.G., H.B.,
K.M., A.E.; **Resources:** G.G., S.L., P.D.C.; **Funding acquisition**: G.G., S.L., P.D.C.;
Supervision: G.G. and S.L.; **Writing – original draft**: G.G. and S.L.; **Writing – review &**
editing: all authors.

Competing interests: P.D.C. and A.E. are inventors on patent applications dealing with the
use of specific bacteria and components in the treatment of different diseases. P.D.C. was co-
founder of The Akkermansia Company SA and Enterosys. The other authors declare no
competing interests.

     FIGURE LEGENDS

Figure 1: Deletion of NAPE-PLD in the nervous system promotes reward seeking

behaviors and reduces *N*-acylethanolamines in the VTA. The reward-like and motivational

phenotype of Napepld$^{f/f}$ and Napepld$^{\Delta CNS}$ mice was tested through an operant conditioning

paradigm (lever press). (A-A^2) Operant conditioning during 4 consecutive sessions with a

fixed ratio 1 (FR1) schedule. (A) Number of rewards, (A^1) number of active lever presses and

(A^2) percentage of discrimination between the active and inactive lever presses. (B-B^2)

Operant conditioning during 4 consecutive sessions with a progressive ratio (PR) schedule.

(B) Number of rewards, (B^1) number of active lever presses and (B^2) percentage of

discrimination between the active and inactive lever presses. (C-H) Lipidomic analysis of *N*-

acylethanolamines (NAEs) and (I) 2-AG in the midbrain ventral tegmental area (VTA), the

dorsal striatum/nucleus accumbens (DS/NAc), the prefrontal cortex (PFC) and the

hippocampus (Hippo) of Napepld$^{f/f}$ and Napepld$^{\Delta CNS}$ mice. Anandamide (AEA), *N*-

oleoylethanolamine (OEA), *N*-palmitoylethanolamine (PEA), *N*-stearoylethanolamine (SEA),

N-linoleoylethanolamine   (LEA),   *N*-docosahexaenoylethanolamine   (DEA),   2-

Arachidonoylglycerol (2-AG). Statistics: *$p<0.05$, **$p<0.01$ and ***$p<0.001$ for Napepld$^{\Delta CNS}$

vs Napepld$^{f/f}$ mice; [###]$p<0.001$ for VTA *vs* other brain structures (Napepld$^{f/f}$ mice). For

number of mice/group and statistical details see **Suppl. Table 1**.

Figure 2: *Napepld* is expressed in the VTA. (A-E) Meta-analysis and clustering of snRNA-

seq results in the rat NAc and VTA focusing of NAEs- and endocannabinoids (eCBs)-related

synthesis machineries and receptors. Percentage (%) of co-expression of *Napepld* in the NAc

Drd1-MSNs (A) and *Drd2*-MSNs (B) (extracted from (47)) as well as in the VTA dopamine

(DA)-neurons (C), VTA GABA-neurons (D) and VTA glutamate (Glut)-neurons (E)

(extracted from (48)). (F) Meta-analysis of bulk transcriptomics in the mouse (extracted from

(50)) and (G) human midbrains (extracted from (51)).

Figure 3: NAPE-PLD knock-down in the VTA promotes reward-seeking behaviors and

food preference. (A) Scheme and immunofluorescence sections indicating the viral injection

of AAV-Cre-GFP or AAV-GFP in the VTA of Napepld$^{f/f}$ mice. Scale bar: 250 μm. (B, C)

Operant conditioning during 3 consecutive sessions with a FR1 (B) and a PR (C) schedule.

For both schedules the number of rewards, the number of lever presses and the percentage of

discrimination between the active and inactive lever presses are shown. (D, E) Drawings

indicate the food-induced conditioned-place preference (CPP) and the T-Maze paradigms,
respectively. (F) CPP in Napepld$^{VTA-GFP}$ and Napepld$^{\Delta VTA}$ food-restricted mice (10% of body
weight reduction). (G) CPP in Napepld$^{VTA-GFP}$ and Napepld$^{\Delta VTA}$ sated mice. (H) Learning and
reversal learning performances of Napepld$^{VTA-GFP}$ and Napepld$^{\Delta VTA}$ food-restricted mice (10%
of body weight reduction) in the T-Maze. (I-K) Food preference in Napepld$^{VTA-GFP}$ and
Napepld$^{\Delta VTA}$ mice using three different choices: sucralose *vs* water (I), sucrose *vs* water (J),
and lipids *vs* water (K). (L) Detection and quantification of cFos-positive neurons in the NAc
of Napepld$^{VTA-GFP}$ and Napepld$^{\Delta VTA}$ mice following the exposure of a fixed amount of high-fat
diet (HFD). Scale bar: 100 μm. Statistics: *$p<0.05$, **$p<0.01$ and ***$p<0.001$ for
Napepld$^{\Delta VTA}$ *vs* Napepld$^{VTA-GFP}$ mice (B, C, H, L). Statistics: **$p<0.01$ and ***$p<0.001$ for
Napepld$^{\Delta VTA}$ (post- *vs* pre-test in CPP) or Napepld$^{VTA-GFP}$ (post- *vs* pre-test in CPP) (F, G).
Statistics: **$p<0.01$ and ***$p<0.001$ for Napepld$^{\Delta VTA}$ (sucra/sucro/lipids *vs* water) or
Napepld$^{VTA-GFP}$ mice (sucra/sucro/lipids *vs* water) (I, J, K); $^{\#\#\#}p<0.001$ for Napepld$^{\Delta VTA}$ *vs*
Napepld$^{VTA-GFP}$ mice (I, J, K). For number of mice/group and statistical details see **Suppl.**
Table 1.
781
Figure 4: NAPE-PLD knock-down in the VTA regulates *in vivo* DA release dynamics.
(A) Drawing indicates the double viral strategies to record *in vivo* DA dynamics in
Napepld$^{VTA-GFP}$ and Napepld$^{\Delta VTA}$ behaving mice by using fiber photometry coupled to the DA
biosensor GRAB-DA2m. DA dynamics were measured during food- and non-food-dependent
behaviors. (B) Immunofluorescence detection of GRAB-DA2m in the NAc and AAV-Cre-
GFP in the VTA (with TH staining). Scale bars: 250 μm. (C-J) Temporal dynamics and/or
heatmaps of DA releasing dynamics in Napepld$^{VTA-GFP}$ and Napepld$^{\Delta VTA}$ mice during
consumption of HFD in both fasted (C, D) and fed (E, F) conditions as well as during cocaine
administration (G, H) and tail suspension (I, J). Statistics: *$p<0.05$, **$p<0.01$ and ***$p<0.001$
for Napepld$^{\Delta VTA}$ *vs* Napepld$^{VTA-GFP}$ mice; $^{\#}p<0.05$ and $^{\#\#}p<0.01$ for cocaine time course in
Napepld$^{VTA-GFP}$ mice; $^{\S\S}p<0.01$ and $^{\S\S\S}p<0.01$ for cocaine time course in Napepld$^{\Delta VTA}$ mice.
For number of mice/group and statistical details see **Suppl. Table 1**.
794
Figure 5: VTA NAPE-PLD contributes to the regulation of energy balance and
metabolic efficiency. (A) Body weight and body composition of Napepld$^{VTA-GFP}$ and
Napepld$^{\Delta VTA}$ mice. (B-K) Indirect calorimetric studies in Napepld$^{VTA-GFP}$ and Napepld$^{\Delta VTA}$
mice to assess energy balance and metabolic efficiency. Longitudinal measurements in
calorimetric chambers of locomotor activity (B, B^1), food intake (C, C^1), energy expenditure

EE (D, D^1), respiratory exchange ratio RER (E, E^1) and fatty acid oxidation FAO (F, F^1) in

Napepld$^{VTA-GFP}$ and Napepld$^{\Delta VTA}$ sated mice. (G-K) Indirect calorimetric studies in

Napepld$^{VTA-GFP}$ and Napepld$^{\Delta VTA}$ mice undergoing a fasting/refeeding metabolic challenge.

(G) Locomotor activity, (H) cumulative food intake, (I) energy expenditure, (J) respiratory

exchange ratio and (K) fatty acid oxidation. (L, M) Temporal kinetics and heatmaps of *in vivo*

DA release dynamics in Napepld$^{VTA-GFP}$ and Napepld$^{\Delta VTA}$ fasted mice and exposed to a chow

pellet. Statistics: *p<0.05, **p<0.01 and ***p<0.001 for Napepld$^{\Delta VTA}$ *vs* Napepld$^{VTA-GFP}$

mice. For number of mice/group and statistical details see **Suppl. Table 1**.

Figure 6: VTA NAPE-PLD regulates energy balance independently from exercise. (A,

A^1) Time-locked (30 min/session) access to running wheels to mimic the reinforcing

properties of exercise. (B-H) Indirect calorimetric studies in calorimetric chambers equipped

with running wheels to assess energy balance and metabolic efficiency in Napepld$^{VTA-GFP}$ and

Napepld$^{\Delta VTA}$ mice. (B, B^1) Spontaneous locomotor activity, (C, C^1) running wheel activity,

(D) total locomotor activity (spontaneous + running wheel activities), (E, E^1) energy

expenditure, (F) cumulative food intake, (G) respiratory exchange ratio and (H) fatty acid

oxidation. Statistics: *p<0.05, **p<0.01 and ***p<0.001 for Napepld$^{\Delta VTA}$ *vs* Napepld$^{VTA-GFP}$

mice. For number of mice/group and statistical details see **Suppl. Table 1**.

Figure 7: VTA NAPE-PLD protects from obesity-associated metabolic features. (A)

Body weight and body composition (fat and lean mass) of Napepld$^{VTA-GFP}$ and Napepld$^{\Delta VTA}$

mice exposed to chronic high-fat diet (HFD) and consequently characterized for energy

balance and metabolic efficiency. (B) Cumulative locomotor activity, (C, C^1) food intake, (D,

D^1) energy expenditure, (E, E^1) fatty acid oxidation and (F) respiratory exchange ratio.

Statistics: *p<0.05, **p<0.01 and ***p<0.001 for Napepld$^{\Delta VTA}$ *vs* Napepld$^{VTA-GFP}$ mice. For

number of mice/group and statistical details see **Suppl. Table 1**.

Supplementary Figure 1: Tissue-specific deletion of NAPE-PLD and operant behavior.

Operant conditioning for food-seeking (lever press), under both FR1 and PR schedules, was

evaluated in different mouse lines. NAPE-PLD was genetically removed from the whole-body

(Napepld^{KO} mice, A and B) or from intestinal epithelial cells (Napepld$^{\Delta IEC}$, E and F). In

addition, operant conditioning was also performed in Nestin-Cre mice (C and D) to discard

the potential effect of Cre in the phenotype observed in Napepld$^{\Delta CNS}$ mice (**Fig. 1A, B**). For

number of mice/group and statistical details see **Suppl. Table 1**.

Supplementary Figure 2: Levels of fatty acids in the VTA of Napepld$^{\Delta CNS}$ mice.
Lipidomic detection of linoleic (A), arachidonic (B) and oleic (C) acids in the VTA of
Napepld$^{f/f}$ and Napepld$^{\Delta CNS}$ mice. For number of mice/group and statistical details see **Suppl.**
Table 1.

Supplementary Figure 3: NAPE-PLD knock-down in the VTA promotes reward-seeking
behaviors in both males and females. (A-C) Operant conditioning in Napepld$^{VTA-GFP}$ and
Napepld$^{\Delta VTA}$ male mice following a FR1→FR5→PR training schedule in food-restricted
conditions (reduction of 10% of body weight). Of note, Napepld$^{VTA-GFP}$ and Napepld$^{\Delta VTA}$ male
mice also underwent a session of PR schedule in sated conditions (D). (E-G) Operant
conditioning in Napepld$^{VTA-GFP}$ and Napepld$^{\Delta VTA}$ female mice following a FR1→FR5→PR
training schedule in food-restricted conditions (reduction of 10% of body weight). Of note,
Napepld$^{VTA-GFP}$ and Napepld$^{\Delta VTA}$ female mice also underwent a session of PR schedule in
sated conditions (H). Statistics: *p<0.05, **p<0.01 and ***p<0.001 for Napepld$^{\Delta VTA}$ *vs*
Napepld$^{VTA-GFP}$ mice. For number of mice/group and statistical details see **Suppl. Table 1**.

Supplementary Figure 4: NAPE-PLD knock-down in the VTA does not alter body
weight and food restriction-mediated weight loss. (A) Body weight of Napepld$^{VTA-GFP}$ and
Napepld$^{\Delta VTA}$ male mice undergoing the operant conditioning paradigm. (B) Body weight of
Napepld$^{VTA-GFP}$ and Napepld$^{\Delta VTA}$ female mice undergoing the operant conditioning paradigm.
Statistics: ***p<0.001 for Napepld$^{VTA-GFP}$ mice (initial *vs* food restriction); ###p<0.001 for
Napepld$^{\Delta VTA}$ mice (initial *vs* food restriction). For number of mice/group and statistical details
see **Suppl. Table 1**.

Supplementary Figure 5: VTA NAPE-PLD and *in vivo* DA dynamics. (A) Temporal
kinetics of *in vivo* DA release in Napepld$^{VTA-GFP}$ and Napepld$^{\Delta VTA}$ fed mice exposed to HFD
corresponding to **Fig. 4E-F**. (B) Temporal kinetics of *in vivo* DA release in Napepld$^{VTA-GFP}$
and Napepld$^{\Delta VTA}$ mice undergoing the tail suspension (TS) test and corresponding to **Fig. 4I-**
J. (C) Longitudinal and cumulative measurements of GBR12909-induced locomotor activity
in Napepld$^{VTA-GFP}$ and Napepld$^{\Delta VTA}$ mice. Statistics: **p<0.01 for Napepld$^{\Delta VTA}$ *vs*
Napepld$^{VTA-GFP}$ mice (GBR experiment). For number of mice/group and statistical details see
Suppl. Table 1.

Supplementary Figure 6: VTA NAPE-PLD contributes to glucose homeostasis. (A)

Blood glucose and (B) insulin levels in Napepld$^{\text{VTA-GFP}}$ and Napepld$^{\Delta\text{VTA}}$ mice after an oral

glucose tolerance test (OGTT). Statistics: **p<0.01 for Napepld$^{\Delta\text{VTA}}$ *vs* Napepld$^{\text{VTA-GFP}}$. For

number of mice/group and statistical details see **Suppl. Table 1**.

REFERENCES

1. Morton GJ, Cummings DE, Baskin DG, Barsh GS, Schwartz MW (2006): Central nervous
system control of food intake and body weight. *Nature* 443: 289–95.

2. Rossi MA, Stuber GD (2018): Overlapping Brain Circuits for Homeostatic and Hedonic
Feeding. *Cell Metab* 27: 42–56.

3. de Araujo IE, Schatzker M, Small DM (2020): Rethinking Food Reward. *Annu Rev Psychol*
71: 139–164.

4. Berland C, Small DM, Luquet S, Gangarossa G (2021): Dietary lipids as regulators of
reward processes: multimodal integration matters. *Trends Endocrinol Metab* 32: 693–705.

5. Mock ED, Gagestein B, van der Stelt M (2023): Anandamide and other N-
acylethanolamines: A class of signaling lipids with therapeutic opportunities. *Prog Lipid Res*
89: 101194.

6. Hussain Z, Uyama T, Tsuboi K, Ueda N (2017): Mammalian enzymes responsible for the
biosynthesis of N -acylethanolamines. *Biochimica et Biophysica Acta (BBA) - Molecular and*
Cell Biology of Lipids 1862: 1546–1561.

7. Okamoto Y, Morishita J, Tsuboi K, Tonai T, Ueda N (2004): Molecular characterization of
a phospholipase D generating anandamide and its congeners. *J Biol Chem* 279: 5298–305.

8. Ahern GP (2003): Activation of TRPV1 by the satiety factor oleoylethanolamide. *The*
Journal of biological chemistry 278: 30429–34.

9. Fu J, Gaetani S, Oveisi F, Lo Verme J, Serrano A, Rodriguez De Fonseca F, *et al.* (2003):
Oleylethanolamide regulates feeding and body weight through activation of the nuclear
receptor PPAR-alpha. *Nature* 425: 90–3.

10. Lo Verme J, Fu J, Astarita G, La Rana G, Russo R, Calignano A, Piomelli D (2005): The
nuclear receptor peroxisome proliferator-activated receptor-alpha mediates the anti-
inflammatory actions of palmitoylethanolamide. *Mol Pharmacol* 67: 15–19.

11. Cristino L, Becker T, Di Marzo V (2014): Endocannabinoids and energy homeostasis: an
update. *Biofactors* 40: 389–397.

12. Esposito G, Capoccia E, Turco F, Palumbo I, Lu J, Steardo A, *et al.* (2014):
Palmitoylethanolamide improves colon inflammation through an enteric glia/toll like receptor
4-dependent PPAR-α activation. *Gut* 63: 1300–1312.

13. Hansen HS, Vana V (2019): Non-endocannabinoid N-acylethanolamines and 2-
monoacylglycerols in the intestine. *Br J Pharmacol* 176: 1443–1454.

14. Rodriguez de Fonseca F, Navarro M, Gomez R, Escuredo L, Nava F, Fu J, *et al.* (2001):
An anorexic lipid mediator regulated by feeding. *Nature* 414: 209–12.

15. Terrazzino S, Berto F, Dalle Carbonare M, Fabris M, Guiotto A, Bernardini D, Leon A
(2004): Stearoylethanolamide exerts anorexic effects in mice via down-regulation of liver
stearoyl-coenzyme A desaturase-1 mRNA expression. *FASEB J* 18: 1580–1582.

16. DiPatrizio NV (2016): Endocannabinoids in the Gut. *Cannabis Cannabinoid Res* 1: 67–
77.

17. Fu J, Kim J, Oveisi F, Astarita G, Piomelli D (2008): Targeted enhancement of
oleoylethanolamide production in proximal small intestine induces across-meal satiety in rats.
Am J Physiol Regul Integr Comp Physiol 295: R45-50.

18. Everard A, Plovier H, Rastelli M, Van Hul M, de Wouters d'Oplinter A, Geurts L, *et al.*

(2019): Intestinal epithelial N-acylphosphatidylethanolamine phospholipase D links dietary
fat to metabolic adaptations in obesity and steatosis. *Nature communications* 10: 457.

19. Melis M, Pillolla G, Luchicchi A, Muntoni AL, Yasar S, Goldberg SR, Pistis M (2008):
Endogenous Fatty Acid Ethanolamides Suppress Nicotine-Induced Activation of Mesolimbic
Dopamine Neurons through Nuclear Receptors. *J Neurosci* 28: 13985–13994.

20. Palese F, Pontis S, Realini N, Piomelli D (2019): A protective role for N-
acylphosphatidylethanolamine phospholipase D in 6-OHDA-induced neurodegeneration. *Sci
Rep* 9: 15927.

21. Tevosian M, Todorov H, Lomazzo E, Bindila L, Ueda N, Bassetti D, *et al.* (2023): NAPE-
PLD deletion in stress-TRAPed neurons results in an anxiogenic phenotype. *Transl
Psychiatry* 13: 152.

22. Morishita J, Okamoto Y, Tsuboi K, Ueno M, Sakamoto H, Maekawa N, Ueda N (2005):
Regional distribution and age-dependent expression of N-acylphosphatidylethanolamine-
hydrolyzing phospholipase D in rat brain. *J Neurochem* 94: 753–762.

23. Egertova M, Simon GM, Cravatt BF, Elphick MR (2008): Localization of N-acyl
phosphatidylethanolamine phospholipase D (NAPE-PLD) expression in mouse brain: A new
perspective on N-acylethanolamines as neural signaling molecules. *The Journal of
comparative neurology* 506: 604–15.

24. Leishman E, Mackie K, Luquet S, Bradshaw HB (2016): Lipidomics profile of a NAPE-
PLD KO mouse provides evidence of a broader role of this enzyme in lipid metabolism in the
brain. *Biochim Biophys Acta* 1861: 491–500.

25. Mock ED, Mustafa M, Gunduz-Cinar O, Cinar R, Petrie GN, Kantae V, *et al.* (2020):
Discovery of a NAPE-PLD inhibitor that modulates emotional behavior in mice. *Nat Chem
Biol* 16: 667–675.

26. Muccioli GG (2010): Endocannabinoid biosynthesis and inactivation, from simple to
complex, 2010/03/23 ed. *Drug Discov Today* 15: 474–83.

27. Leung D, Saghatelian A, Simon GM, Cravatt BF (2006): Inactivation of N-acyl
phosphatidylethanolamine phospholipase D reveals multiple mechanisms for the biosynthesis
of endocannabinoids. *Biochemistry* 45: 4720–6.

28. Wangensteen T, Akselsen H, Holmen J, Undlien D, Retterstøl L (2011): A common
haplotype in NAPEPLD is associated with severe obesity in a Norwegian population-based
cohort (the HUNT study). *Obesity (Silver Spring)* 19: 612–617.

29. Doris JM, Millar SA, Idris I, O'Sullivan SE (2019): Genetic polymorphisms of the
endocannabinoid system in obesity and diabetes. *Diabetes Obes Metab* 21: 382–387.

30. Liu J, Wang L, Harvey-White J, Huang BX, Kim HY, Luquet S, *et al.* (2008): Multiple
pathways involved in the biosynthesis of anandamide. *Neuropharmacology* 54: 1–7.

31. Tronche F, Kellendonk C, Kretz O, Gass P, Anlag K, Orban PC, *et al.* (1999): Disruption
of the glucocorticoid receptor gene in the nervous system results in reduced anxiety,
1999/09/02 ed. *Nat Genet* 23: 99–103.

32. Giusti SA, Vercelli CA, Vogl AM, Kolarz AW, Pino NS, Deussing JM, Refojo D (2014):
Behavioral phenotyping of Nestin-Cre mice: implications for genetic mouse models of
psychiatric disorders. *J Psychiatr Res* 55: 87–95.

33. el Marjou F, Janssen KP, Chang BH, Li M, Hindie V, Chan L, *et al.* (2004): Tissue-
specific and inducible Cre-mediated recombination in the gut epithelium, 2004/07/30 ed.

Genesis 39: 186–93.

34. Lallemand Y, Luria V, Haffner-Krausz R, Lonai P (1998): Maternally expressed PGK-Cre transgene as a tool for early and uniform activation of the Cre site-specific recombinase. *Transgenic Res* 7: 105–112.

35. Sun F, Zhou J, Dai B, Qian T, Zeng J, Li X, *et al.* (2020): Next-generation GRAB sensors for monitoring dopaminergic activity in vivo. *Nat Methods* 17: 1156–1166.

36. Berland C, Castel J, Terrasi R, Montalban E, Foppen E, Martin C, *et al.* (2022): Identification of an endocannabinoid gut-brain vagal mechanism controlling food reward and energy homeostasis. *Mol Psychiatry* 27: 2340–2354.

37. Berland C, Montalban E, Perrin E, Di Miceli M, Nakamura Y, Martinat M, *et al.* (2020): Circulating Triglycerides Gate Dopamine-Associated Behaviors through DRD2-Expressing Neurons. *Cell Metab* 31: 773-790.e11.

38. Montalban E, Walle R, Castel J, Ansoult A, Hassouna R, Foppen E, *et al.* (2023): The addiction-susceptibility TaqIA/Ankyrin repeat and kinase domain containing 1 kinase (ANKK1) controls reward and metabolism through dopamine receptor type 2 (D2R)-expressing neurons. *Biological Psychiatry* S0006322323000847.

39. Gangarossa G, Castell L, Castro L, Tarot P, Veyrunes F, Vincent P, *et al.* (2019): Contrasting patterns of ERK activation in the tail of the striatum in response to aversive and rewarding signals. *J Neurochem* 151: 204–226.

40. Raboune S, Stuart JM, Leishman E, Takacs SM, Rhodes B, Basnet A, *et al.* (2014): Novel endogenous N-acyl amides activate TRPV1-4 receptors, BV-2 microglia, and are regulated in brain in an acute model of inflammation. *Frontiers in cellular neuroscience* 8: 195.

41. Geurts L, Everard A, Van Hul M, Essaghir A, Duparc T, Matamoros S, *et al.* (2015): Adipose tissue NAPE-PLD controls fat mass development by altering the browning process and gut microbiota. *Nat Commun* 6: 6495.

42. Declercq J, Brouwers B, Pruniau VPEG, Stijnen P, de Faudeur G, Tuand K, *et al.* (2015): Metabolic and Behavioural Phenotypes in Nestin-Cre Mice Are Caused by Hypothalamic Expression of Human Growth Hormone. *PLoS One* 10: e0135502.

43. Tellez LA, Han W, Zhang X, Ferreira TL, Perez IO, Shammah-Lagnado SJ, *et al.* (2016): Separate circuitries encode the hedonic and nutritional values of sugar. *Nat Neurosci* 19: 465–70.

44. Tellez LA, Medina S, Han W, Ferreira JG, Licona-Limon P, Ren X, *et al.* (2013): A gut lipid messenger links excess dietary fat to dopamine deficiency. *Science* 341: 800–2.

45. Rastelli M, Van Hul M, Terrasi R, Lefort C, Regnier M, Beiroa D, *et al.* (2020): Intestinal NAPE-PLD contributes to short-term regulation of food intake via gut-to-brain axis. *Am J Physiol Endocrinol Metab* 319: E647–E657.

46. Morales I, Berridge KC (2020): 'Liking' and 'wanting' in eating and food reward: Brain mechanisms and clinical implications. *Physiology & Behavior* 227: 113152.

47. Savell KE, Tuscher JJ, Zipperly ME, Duke CG, Phillips RA, Bauman AJ, *et al.* (2020): A dopamine-induced gene expression signature regulates neuronal function and cocaine response. *Sci Adv* 6: eaba4221.

48. Phillips RA, Tuscher JJ, Black SL, Andraka E, Fitzgerald ND, Ianov L, Day JJ (2022): An atlas of transcriptionally defined cell populations in the rat ventral tegmental area. *Cell Rep* 39: 110616.

49. Morales M, Margolis EB (2017): Ventral tegmental area: cellular heterogeneity, connectivity and behaviour. *Nat Rev Neurosci* 18: 73–85.

50. Brichta L, Shin W, Jackson-Lewis V, Blesa J, Yap EL, Walker Z, *et al.* (2015): Identification of neurodegenerative factors using translatome-regulatory network analysis. *Nat Neurosci* 18: 1325–33.

51. Aguila J, Cheng S, Kee N, Cao M, Wang M, Deng Q, Hedlund E (2021): Spatial RNA Sequencing Identifies Robust Markers of Vulnerable and Resistant Human Midbrain Dopamine Neurons and Their Expression in Parkinson's Disease. *Front Mol Neurosci* 14: 699562.

52. Woodworth HL, Batchelor HM, Beekly BG, Bugescu R, Brown JA, Kurt G, *et al.* (2017): Neurotensin Receptor-1 Identifies a Subset of Ventral Tegmental Dopamine Neurons that Coordinates Energy Balance. *Cell Reports* 20: 1881–1892.

53. Fernandes MFA, Matthys D, Hryhorczuk C, Sharma S, Mogra S, Alquier T, Fulton S (2015): Leptin Suppresses the Rewarding Effects of Running via STAT3 Signaling in Dopamine Neurons. *Cell Metab* 22: 741–749.

54. Medrano M-C, Hurel I, Mesguich E, Redon B, Stevens C, Georges F, *et al.* (2021): Exercise craving potentiates excitatory inputs to ventral tegmental area dopaminergic neurons. *Addict Biol* 26: e12967.

55. Muguruza C, Redon B, Fois GR, Hurel I, Scocard A, Nguyen C, *et al.* (2019): The motivation for exercise over palatable food is dictated by cannabinoid type-1 receptors. *JCI Insight* 4: e126190, 126190.

56. Dubreucq S, Durand A, Matias I, Bénard G, Richard E, Soria-Gomez E, *et al.* (2013): Ventral tegmental area cannabinoid type-1 receptors control voluntary exercise performance. *Biol Psychiatry* 73: 895–903.

57. Kathuria S, Gaetani S, Fegley D, Valiño F, Duranti A, Tontini A, *et al.* (2003): Modulation of anxiety through blockade of anandamide hydrolysis. *Nat Med* 9: 76–81.

58. Walker JM, Huang SM, Strangman NM, Tsou K, Sañudo-Peña MC (1999): Pain modulation by release of the endogenous cannabinoid anandamide. *Proc Natl Acad Sci U S A* 96: 12198–12203.

59. Lambert DM, Vandevoorde S, Jonsson K-O, Fowler CJ (2002): The palmitoylethanolamide family: a new class of anti-inflammatory agents? *Curr Med Chem* 9: 663–674.

60. Lunerti V, Li H, Benvenuti F, Shen Q, Domi A, Soverchia L, *et al.* (2022): The multitarget FAAH inhibitor/D3 partial agonist ARN15381 decreases nicotine self-administration in male rats. *European Journal of Pharmacology* 928: 175088.

61. Melis M, Scheggi S, Carta G, Madeddu C, Lecca S, Luchicchi A, *et al.* (2013): PPARα regulates cholinergic-driven activity of midbrain dopamine neurons via a novel mechanism involving α7 nicotinic acetylcholine receptors. *J Neurosci* 33: 6203–6211.

62. Melis M, Carta S, Fattore L, Tolu S, Yasar S, Goldberg SR, *et al.* (2010): Peroxisome proliferator-activated receptors-alpha modulate dopamine cell activity through nicotinic receptors. *Biol Psychiatry* 68: 256–264.

63. Grace AA (2000): The tonic/phasic model of dopamine system regulation and its implications for understanding alcohol and psychostimulant craving. *Addiction* 95 Suppl 2: S119-128.

64. Tsai H-C, Zhang F, Adamantidis A, Stuber GD, Bonci A, de Lecea L, Deisseroth K
(2009): Phasic firing in dopaminergic neurons is sufficient for behavioral conditioning.
Science 324: 1080–1084.

65. Floresco SB, West AR, Ash B, Moore H, Grace AA (2003): Afferent modulation of
dopamine neuron firing differentially regulates tonic and phasic dopamine transmission. *Nat*
Neurosci 6: 968–973.

66. Kreitzer AC, Regehr WG (2002): Retrograde signaling by endocannabinoids. *Curr Opin*
Neurobiol 12: 324–330.

67. Nyilas R, Dudok B, Urbán GM, Mackie K, Watanabe M, Cravatt BF, *et al.* (2008):
Enzymatic machinery for endocannabinoid biosynthesis associated with calcium stores in
glutamatergic axon terminals. *J Neurosci* 28: 1058–1063.

68. Pupe S, Wallén-Mackenzie Å (2015): Cre-driven optogenetics in the heterogeneous
genetic panorama of the VTA. *Trends in Neurosciences* 38: 375–386.

69. Zell V, Steinkellner T, Hollon NG, Warlow SM, Souter E, Faget L, *et al.* (2020): VTA
Glutamate Neuron Activity Drives Positive Reinforcement Absent Dopamine Co-release.
Neuron 107: 864-873.e4.

70. Boekhoudt L, Omrani A, Luijendijk MCM, Wolterink-Donselaar IG, Wijbrans EC, van
der Plasse G, Adan RAH (2016): Chemogenetic activation of dopamine neurons in the ventral
tegmental area, but not substantia nigra, induces hyperactivity in rats. *Eur*
Neuropsychopharmacol 26: 1784–1793.

71. Perez-Bonilla P, Santiago-Colon K, Matasovsky J, Ramirez-Virella J, Khan R, Garver H,
et al. (2021): Activation of ventral tegmental area neurotensin Receptor-1 neurons promotes
weight loss. *Neuropharmacology* 195: 108639.

72. Mietlicki-Baase EG, Reiner DJ, Cone JJ, Olivos DR, McGrath LE, Zimmer DJ, *et al.*
(2015): Amylin modulates the mesolimbic dopamine system to control energy balance.
Neuropsychopharmacology 40: 372–385.

73. Fanelli F, Mezzullo M, Repaci A, Belluomo I, Ibarra Gasparini D, Di Dalmazi G, *et al.*
(2018): Profiling plasma N-Acylethanolamine levels and their ratios as a biomarker of obesity
and dysmetabolism. *Molecular Metabolism* 14: 82–94.

74. Jumpertz R, Guijarro A, Pratley RE, Piomelli D, Krakoff J (2011): Central and peripheral
endocannabinoids and cognate acylethanolamides in humans: association with race, adiposity,
and energy expenditure. *J Clin Endocrinol Metab* 96: 787–791.

75. Mazier W, Saucisse N, Gatta-Cherifi B, Cota D (2015): The Endocannabinoid System:
Pivotal Orchestrator of Obesity and Metabolic Disease. *Trends Endocrinol Metab* 26: 524–
537.
1082

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7