Genome editing induces the transfer of mitochondrial DNA into the nuclear genome

Jinchun Wu[1,3], Yang Liu[1,3], Liqiong Ou[1,3], Tingting Gan[1,3], Zhengrong Zhangding[1],
Shaopeng Yuan[1], Mengzhu Liu[1], Xinyi Liu[1], Jiasheng Li[2], Jianhang Yin[1], Changchang Xin[1],
Ye Tian[2], Jiazhi Hu[1,*]

[1] The MOE Key Laboratory of Cell Proliferation and Differentiation, School of Life
Sciences, Center for Life Sciences, Genome Editing Research Center, Peking University,
Beijing 100871, China.
[2] State Key Laboratory of Molecular Developmental Biology, Institute of Genetics and
Developmental Biology, Chinese Academy of Sciences,
[3] These authors contributed equally to this work.

[*] Corresponding authors: hujz@pku.edu.cn

Abstract

Mitochondria serve as the cellular powerhouse, and their distinct DNA makes them a

prospective target for gene editing to treat genetic disorders. However, the impact of genome

editing on mitochondrial DNA (mtDNA) stability remains a mystery. Our study reveals

previously unknown risks of genome editing that both nuclear and mitochondrial editing

cause broad transfer of mitochondrial DNA segments into the nuclear genome in various cell

types including human cell lines, primary T cells, retinal cells, and mouse embryos.

Furthermore, drug-induced mitochondrial stresses and mtDNA breaks exacerbate this transfer

of mtDNA into the nuclear genome. Notably, we observe that the newly developed

mitochondrial base editor DdCBE can also cause widespread mtDNA integrations. However,

we provide a practical solution to suppress the transfer of mtDNA by co-expressing TREX1

or TREX2 exonucleases during DdCBE editing. These findings also shed light on the origins

of mitochondrial-nuclear DNA segments.

Keywords

Genome editing, DdCBE, mitochondrial base editor, CRISPR-Cas9, mitochondrial

integration into nuclear DNA, TREX1/2

Introduction

The human genome consists of two copies of nuclear DNA and a few hundred to a few thousand copies of mitochondrial DNA (mtDNA), which are separated by the nuclear and transferred to the nuclear genome during the endosymbiotic evolution (Gray et al., 1999). Consequently, the human genome contains hundreds of nuclear-mitochondrial DNA segments (NUMTs) and still undergoes the natural transfer of mtDNA to the nuclear DNA (Wei et al., 2020; Wei et al., 2022a; Ju et al., 2015). *De novo* NUMTs were detected in approximately one per 10^4 birth and 10^3 cancers (Ju et al., 2015; Wei et al., 2022a). Meanwhile, NUMTs in tumor cells tend to be embedded in gene-rich regions of nuclear DNA, suggesting that they may be the causal integration of certain tumors (Wei et al., 2022a; Kopinski et al., 2021).

Recently, the focus of gene editing has extended from the nuclear DNA (Jiang and Doudna, 2017; Hsu et al., 2014; Anzalone et al., 2020; Liu et al., 2021) to mtDNA (Silva-Pinheiro and Minczuk, 2022; Barrera-Paez and Moraes, 2022), leading to the development of a series of highly effective mitochondrial editors. These tools aim to eliminate mutated mtDNA by inducing DNA double-stranded breaks (DSBs) with mitochondrial-targeted transcription activator-like effector nucleases (mitoTALEN) (Bacman et al., 2013; Bacman et al., 2018) or correcting mutations with mitochondrial base editors (Mok et al., 2020; Cho et al., 2022). DNA insertions of genomic or viral origins have been widely observed during gene editing exerted by TALEN or clustered regularly interspaced short palindromic repeats associated system (CRISPR-Cas) (Wu et al., 2022; Yin and Hu, 2022; Kosicki et al., 2018). However, it remains to be investigated whether mitochondrial or nuclear editing can result in the integration of mtDNA segments into the nuclear genome.

Here we found that genome editing on the nuclear DNA caused mtDNA-nuclear DNA fusion at the CRISPR-Cas9 target sites *in vitro* and *in vivo*, validated by both primer-extension-mediated sequencing (PEM-seq) and target sequencing. Moreover, high-fidelity Cas9 variants and base editors were unable to eliminate the transfer of mtDNA to nuclear DNA. Mitochondrial stresses increased the level of mtDNA-nuclear DNA fusion, and mitochondrial editing tools, such as mitoTALEN and DddA-derived cytosine base editor (DdCBE), also caused mtDNA fragility and resulted in the transfer of mtDNA to nuclear DNA. Finally, we found that co-expression of either TREX1 or TREX2 exonuclease in the mitochondria could be a plausible solution to suppress mtDNA fusion to the nuclear DNA during DdCBE treatment.

Results

Mitochondrial DNA fuses to CRISPR-Cas targeting sites in the nuclear genome

To investigate the integration of mtDNA segments into nuclear CRISPR-Cas target sites, we examined potential mtDNA sequences in the CRISPR-Cas9 editing products profiled by PEM-seq (Fig. 1A). PEM-seq utilizes a primer adjacent to the "bait" target site to capture "prey" sequence fused to the target DNA breaks through a one-round primer extension, rather than PCR. Subsequently, the single-stranded DNA products were isolated and subjected to library preparation (Fig. 1A)(Wu et al., 2022; Liu et al., 2022). In order to preclude potential errors in primer annealing, library preparation and sequencing, we conducted parallel PEM-seq analysis at each locus using genomic DNA from both edited and unedited cells (Fig. 1A). By employing PEM-seq, we successfully identified fusion junctions between mtDNA and the CRISPR target sites for all the tested editing loci, including *CLIC4*, *KLHL29*, *NLRC4*, *COL8A1*, *NUDT16*, *LNX1*, *FGF18*, *VEGFA_1/2*, *HBB*, *IFNγ*, and *P2RX5-TAX1BP3* in HEK293T cells, edited with CRISPR-Cas family editors including *Lb*Cas12a, *As*Cas12a, *Un1*Cas12f, CasMINI, and CasMINI_ge4.1 (Fig. 1B) (Xin et al., 2022). Since the human nuclear genome contains multiple embedded mtDNA segments, we only counted the chimeric reads where the prey sequences were better aligned to the mtDNA as mitochondria-nuclear fusions or translocations, exemplified by two fragments fused to *VEGFA_1* or *NLRC4* baits (Fig. S1A). The frequency of mtDNA integration at CRISPR-Cas target site varied from one per 10^3 to 10^5 edited events in HEK293T cells, with the Cas12 family exhibiting slightly lower levels of mtDNA integrations in comparison to *Sp*Cas9 (Fig. 1B-E and S1B-D). In contrast, unedited samples showed no mtDNA-nuclear DNA fusions for 10 of the 12 tested loci. Only two editing events out of 625,886 for *VEGFA_1* and one editing event out of 897,959 for *HBB* loci were detected in the unedited samples (Fig. 1B, Supplementary Table 1). However, these events contained the exact same sequences as those observed in the corresponding editing samples (Supplementary Table 1), indicating that the rare events in the control libraries may originate from clustering errors during Illumina sequencing.

High-fidelity *Sp*Cas9 variants such as eSpCas9, HF1, FeCas9, EvoCas9, HiFiCas9, Hypa, LZ3, and Sniper Cas9, as well as PAM-flexible *Sp*Cas9 variants including Cas9-NG, xCas9, SpG, and SpRY, have been developed to enhance editing specificity or broaden editing scope, respectively (Schmid-Burgk et al., 2020; Zhang et al., 2021). These *Sp*Cas9 variants exhibited comparable levels of mtDNA-nuclear DNA fusions to wild-type *Sp*Cas9

(Fig. 1F). Considering the inheritability of editing products in embryonic stem cells (ESCs),

we also performed gene editing with five high-fidelity *Sp*Cas9 variants (Hypa, HF1, FeCas9,

eCas9, and SuperFi) in parallel with *Sp*Cas9 in mouse ESCs (Zhang et al., 2021; Bravo et al.,

2022). Consistently, we identified mtDNA integrations at the *c-Myc* target site in mouse

ESCs (Fig. 1G). In contrast, by avoiding DSB generation, base editors significantly reduced

mtDNA-nuclear fusion products at target loci (Fig. 1H)(Yin et al., 2022b). In total, we have

identified over 1,400 mtDNA-nuclear DNA fusion events induced by various CRISPR-Cas

editing tools. The mtDNA fusion junctions were distributed throughout the mitochondrial

genome (Fig. S1E) and over 40% of mtDNA-nuclear DNA fusion events involved

microhomology, primarily with 1-6 base pairs (Fig. S1F).

Mitochondria-nuclear fusions undergo clonal expansion

The CRISPR-Cas9 system has been utilized to generate universal chimeric antigen

receptor (CAR) T cells from human cord blood by disrupting the *TRAC*, *TRBC*, and *PDCD1*

genes (Fig. 2A) (Yin et al., 2022b). The levels of mtDNA integrations in these edited primary

immune cells were comparable to those observed in HEK293T cells, regardless of the

position of target sites (Fig. S2A). Moreover, mtDNA integrations were detected in human

CAR T cells after *ex vivo* culture for 3, 7, or 14 days (Fig. 2B and S2A) (Yin et al., 2022b).

By labeling products with random molecular barcodes (RMB), PEM-seq enables the

distinguishment of the same mtDNA-nuclear DNA fusion junction present in different

expanded cells (Fig. 2A)(Wu et al., 2022). Notably, multiple chimeric junctions occurred in

more than one cell (Fig. 2C), implying the expansion of mtDNA-nuclear DNA fusions during

T cell culture *ex vivo*.

To investigate the potential expansion of mtDNA integrations *in vivo*, we examined

CRISPR-Cas9-edited mouse TCR T cells in a chronic inflammation murine model (Wu et al.,

2022). Activated TCR T cells post Cas9 editing were infused into recipient mice for 3 weeks.

Subsequently, both activated T cells and *in vivo* expanded T cells were subjected to PEM-seq

analysis. Three different mtDNA-*c-Myc* fusions were identified in 167,611 editing events in

activated T cells (Fig. 2E). Remarkably, one fusion junction was captured 38 times after

infusion into the recipient mouse for 3 weeks (0.0132% of 286,818 editing events) (Fig. 2E).

The identified 38 reads showed diverse RMBs and prey sequences, indicating that these

fusions were originated from different cells but not PCR duplication (Fig. S2B). Moreover,

mtDNA integrations were also detected in gene editing with an age-related macular

degeneration (AMD) murine model *in vivo* (Fig. 2F), where adeno-associated virus (AAV)

was used to target the *Vegfa* gene in retinal cells (Yin et al., 2022a). One mtDNA-*Vegfa*

fusion event was found in 27,749 editing events from two different cells 4 weeks post-

injection, and one fusion was captured in 35,534 editing events 12 weeks post-treatment (Fig.

2F). Collectively, these data suggest that mtDNA integrations can be amplified multiple

times as the edited cells undergo clonal expansion.

138   The CRISPR-based base editors enable gene correction in the nuclear genome without

directly inducing DSBs (Anzalone et al., 2020). To assess the nuclear base editors, the

genome-wide off-target analysis by two-cell embryo injection (GOTI) assay was developed,

which edited the embryos at the two-cell stage and collected cells from murine E14.5

embryos for whole-genome sequencing (Zuo et al., 2019). We employed the NUMTs-

detection algorithm (Wei et al., 2022a) to analyze the discordant reads of GOTI libraries with

base editor 3 (BE3) targeting the *Tyr-D* sites (Zuo et al., 2019). For the first set, we detected

14 shared mitochondrial segments in both control and BE3-edited cells. The BE3-edited cells

had seven new mitochondrial segments and one missing segment compared to the control

cells genome-wide (Fig. 2G), suggesting a potentially higher level of mtDNA integrations

after base editing. Similar findings were obtained for the second set (Fig. S2C). Taken

together, mtDNA integrations persisted during the development of mouse embryos.

Mitochondrial perturbance enhances mtDNA transfer into nuclear DNA

152   Mitochondrial integrations into the nuclear target sites require the fusion of DNA breaks

at both the mtDNA and nuclear DNA. To explore the contributions of mtDNA breaks to

mitochondrial integrations during gene editing, we induced mitochondrial stress that can

generate mtDNA breakage and release fragmented mtDNA into the cytosol (Xian et al.,

2022; Kim et al., 2019). We employed carbonyl cyanidem-chlorophenyl hydrazone (CCCP)

to disrupt the membrane potential and used paraquat to induce oxidative damage (O'Malley

et al., 2020; Fessler et al., 2020; Riley et al., 2018). The levels of mtDNA integrations were

found to be approximately 2-fold higher after treatment with CCCP or paraquat, as

determined by PEM-seq (Fig. 3A and 3B). For further validation, we developed insertion-

enriched target sequencing (Insert-seq), which employed two rounds of targeted PCR and two

rounds of size selection to enrich fragment insertions at the nuclear editing site (Fig. 3C and

S3A). Insert-seq mainly captured DNA fragment insertions less than 500 bp with a median

length of 163 bp (Fig. S3A and S3B). We performed Insert-seq analysis on the same genomic

DNA used for PEM-seq. The mtDNA integrations at the *c-MYC* target site accounted for

approximately 0.43% and 0.42% in the presence of CCCP and paraquat, respectively, *via*

Insert-seq. In contrast, the integration level was only 0.11% in the absence of mitochondrial

stresses (Fig. 3D).

To investigate whether mtDNA breaks could be directly involved in nuclear DNA

translocations, we employed mitoTALEN to generate DSBs in the *ND4* gene of the

mitochondrial genome and performed PEM-seq by using the bait DSBs at the *c-MYC* locus

on the nuclear DNA (Fig. 4A). We found that mitoTALEN treatment resulted in over 13-fold

more mtDNA-nuclear DNA fusions than unedited cells (0.079% vs. 0.006% of total editing

events) (Fig. 4B). The fusion junctions were highly enriched at the mitoTALEN target sites

(Fig. 4C), as validated by PCR and sanger sequencing with two pairs of primers located

adjacent to the editing sites of *c-MYC* and *ND4* (Fig. S4A and S4B). We detected many

junctions over the entire mitochondrial genome beyond the *ND4* target site (Fig. 4C),

consistent with previous findings that DSBs lead mainly to rapid fragmentation and

degradation of the mtDNA (Silva-Pinheiro and Minczuk, 2022; Nissanka et al., 2018). We

also conducted Insert-seq and identified over 6-fold more mitochondrial insertions in the

presence of mitoTALEN than in untreated cells (0.39% vs. 0.06% of total insertions) (Fig.

4D and S4C), in line with PEM-seq. Of note, one mtDNA insertion at the *c-MYC* locus

captured by Insert-seq harbored the same junctions as one insertion identified by PEM-seq

(Fig. S4D). Collectively, these results suggest that fragmented mtDNA derived from

mitochondrial stresses and DSBs can integrate into nuclear breaks.

Mitochondrial editors result in mtDNA integrations into the nuclear DNA

We next sought to investigate whether mitochondrial editing tools could induce the

transfer of mtDNA into nuclear DNA. MitoTALEN has been developed to eliminate

mitochondria carrying specific mutations and thus alleviate the mitochondrial heteroplasmy

(Bacman et al., 2013; Bacman et al., 2018). MitoTALEN generated substantial DSBs within

the editing window between the two TALE binding sites. To identify *bona fide* mitochondrial

integrations, we placed the bait primer upstream of the left TALE binding site in the

mitochondrial genome and only considered PEM-seq chimeric reads harboring mtDNA

junctions within the editing window (Fig. 5A). We used mitoTALEN to edit the *ND5.1* gene

and observed 7-fold more mtDNA-nuclear DNA fusions than in untreated cells (Fig. 5B and

5C). The identified nuclear junctions were evenly distributed on the chromosomes, with

about 50.9% (57 of 112) of these junctions located in gene regions. Similarly, 5-fold more

mtDNA-nuclear DNA fusions were captured in cells post mitoTALEN editing at the *ND4*

gene (Fig. S5A and S5B).

The DdCBE enables CRISPR-free nucleotide modification in mtDNA, providing an important tool for mtDNA manipulation and disease therapy (Mok et al., 2020). We next investigated whether DdCBE could induce mtDNA integrations by cloning from *ND4* or *ND5.3* target site (Fig. 5D). At the *ND4* target site, the editing efficiency exceeded 40% (Fig. 5E). To identify *bona fide* mtDNA integrations, we only counted the chimeric junctions containing G to A mutation or missing the G nucleotide due to excision repair (Fig. 5D, 5F, and S5C). Post DdCBE treatment, we observed an 11-fold increase in mtDNA-nuclear DNA fusions at the *ND4* locus compared to untreated cells (Fig. 5F and 5G). Similarly, 4.6-fold more fusions were observed when DdCBE targeted the *ND5.3* locus (Fig. S5D-F). Besides, we also analyzed published GOTI data with DdCBE to induce the G12918A or C12336T mutation (Wei et al., 2022b). For G12918A editing, we detected eight shared mitochondrial segments in both DdCBE-edited and untreated cells, with the DdCBE-edited cells having 10 new mitochondrial segments and three missing segments compared to the untreated cells (Fig. 5H). Similar findings were obtained for C12336T editing (Fig. S5G). Altogether, these data indicated that DdCBE-editing can induce the transfer of mtDNA into the nuclear DNA.

TREX1 and TREX2 suppress the transfer of mtDNA into nuclear DNA

The transfer of mtDNA into the nuclear DNA necessitates mtDNA breakage, and thereby it is conceivable that ectopic expression of exonucleases might promote the degradation of damaged mtDNA and prevent mtDNA integration into the nuclear DNA (Fig. 6A). TREX1 and TREX2 are efficient at degrading free cytosol DNA and have been applied in nuclear gene editing (Yin et al., 2022b). We generated TREX1n by removing the C-terminal cellular trafficking domain from TREX1, which did not affect its exonuclease activity. We fused TREX1n and TREX2 with DdCBE directly or with a mitochondrial targeting sequence (MTS) to co-express with DdCBE (Fig. 6A). The fusion and co-expression of TREX1n/TREX2 had no significant impact on the editing efficiency of DdCBE or the mtDNA copy numbers (Fig. 6B and 6C). We used these mitochondrial editing tools to edit the *ND4* gene in HEK293T cells and detected 12 mtDNA-nuclear DNA fusions after DdCBE editing *via* PEM-seq. In comparison, five to seven fusions were identified when cells were treated with DdCBE either fused or co-expressed with TREX1n/TREX2 (Fig. 6D), implying a potentially lower transfer rate of mtDNA into the nuclear DNA.

To improve the throughput of mtDNA-nuclear DNA fusions for better statistical analysis, we introduced recurrent DSBs at *c-MYC* in the nuclear genome by *Sp*Cas9 and used them as bait to capture DdCBE-induced mtDNA fragmentation *via* PEM-seq (Fig. 6E). The

c-MYC bait successfully captured 47 to 160 mitochondrial integrations for each treatment

after normalization (Fig. 6F). We found 2-fold more mtDNA-nuclear DNA fusions post

DdCBE treatment at the *ND4* locus, consistent with our above findings. However, fusion or

co-expression of DdCBE with TREX1n/TREX2 resulted in a significant decrease in mtDNA-

nuclear DNA fusions, with levels dropping to those of untreated cells (Fig. 6F). Moreover,

TREX2 showed a relatively lower frequency of mtDNA-nuclear DNA fusion than TREX1n,

and the MTS-tagged TREX2 slightly outperformed DdCBE-TREX2 (Fig. 6F).

Discussion

The NUMTs with fragmented or full-length mtDNA have been widely observed in

somatic and cancer cells (Wei et al., 2022a; Wei et al., 2020; Ju et al., 2015; Dayama et al.,

2014; Mishmar et al., 2004; Woischnik and Moraes, 2002; Hazkani-Covo et al., 2010; Yuan

et al., 2020). *De novo* transfer of mtDNA to the nuclear genome can lead to altered gene

expression, such as the overactivation of *c-MYC* and *KCNMA1* oncogenes that contribute to

tumor progression (Shay et al., 1991; Ju et al., 2015), or the disruption of genes, which may

contribute to the development of Pallister-Hall syndrome and mucolipidosis (Turner et al.,

2003; Goldin et al., 2004; Borensztajn et al., 2002). In addition, mtDNA integrations lead to

mitochondrial heteroplasmy, and the level of mitochondrial heteroplasmy in individuals

could change rapidly during development (Schwartz and Vissing, 2002), similar to our

findings that mtDNA integrations can be amplified during the clonal expansion of CAR or

TCR T cells (Fig. 2A-E).

Both the nuclear and mitochondrial genomes encounter spontaneous DNA lesions

triggered by oxidative stress, ultraviolet (UV) light, chemicals, and replicative stress (Tubbs

and Nussenzweig, 2017). Furthermore, the mitochondrial genome undergoes fragmentation

during stresses, resulting in the leakage of multiple mtDNA fragments into the cytosol and

nucleus (Xian et al., 2022; Kim et al., 2019). In this study, we use the gene editing-induced

DSBs to show that any DSB in the nuclear genome might capture these mtDNA fragments,

not limited to DSBs in physiological cellular processes such as V(D)J recombination

(Lebedin et al., 2022). In this context, a 41-bp mtDNA was found to be integrated at the

junction of a reciprocal constitutional translocation (Willett-Brozick et al., 2001). Therefore,

NUMTs can occur in different types of cells and may become inheritable if they occur in ES

cells. In this context, hundreds of NUMTs have been embedded in the human genome. Since

most cancer cells are susceptible to DNA damage and possess a high oxidative metabolism,

de novo NUMTs are quite frequent in cancer cells (Wei et al., 2022a; Kopinski et al., 2021).

The CRISPR-free DdCBE system and other forms of mitochondrial base editors have

demonstrated high efficiency in modifying mtDNA and offer great potential for treating

mitochondria-related diseases (Mok et al., 2020; Mok et al., 2022; Cho et al., 2022).

However, our findings reveal a previously unknown risk that both nuclear and mitochondrial

editing systems, such as DdCBE, can cause the transfer of mtDNA into the nuclear genome

and nearly half of these mitochondrial integrations occur in gene regions. In the context of

$10^8 \sim 10^9$ edited CAR or TCR T cells being infused into patients during therapy (Stadtmauer et

al., 2020), one in $10^4 \sim 10^5$ infused cells may carry deleterious mtDNA-nuclear DNA fusions.

Therefore, caution should be exercised when using both nuclear and mitochondrial editors for

therapeutic purposes. Fortunately, we have shown that the fusion or co-expression of DdCBE

with TREX1/TREX2 can degrade mitochondrial fragments and prevent the fusion of mtDNA

with DSBs in the nuclear genome. In support, TREX1 mutations found in autoimmune

diseases were accompanied by increased amounts of escaped mtDNA fragments (Hemphill

and Perrino, 2019; Skopelja-Gardner et al., 2022). In particular, TREX2, located in the

mitochondria, outperforms DdCBE-fused TREX1n/TREX2 or co-expressed TREX1n, and

has no impact on mtDNA copy numbers, providing a feasible approach to enhance the safety

of mitochondrial base editors.

Materials and Methods

Cell culture and transfection

V6.5 mouse embryonic stem cells (a gift from Dr. Xiong Ji, School of Life Sciences,

Peking University) were cultured in Knockout^{TM} Dulbecco′s Modified Eagle′s Medium

(DMEM, Gibco) containing 15% fetal bovine serum (Gibco), MEM non-essential amino

acids solution (Sigma), nucleotides (Millipore), penicillin-streptomycin (Gibco), L-glutamine

solution (Sigma), 2-mercaptoethanol (Sigma), LIF (Millipore), CHIR99021 (Selleck), and

PD0325901 (Selleck) at 37 °C under 5% CO_2. One million cells were transfected with 2 μg

plasmid containing puromycin resistance gene, gRNA, and Cas9 variants targeting the *c-

MYC* locus by nucleofector (Lonza, 4D-Nucleofector X). The transfected cells were treated

with 1 μg/mL puromycin for 1.5 days after one day post-transfection, and then cultured with

fresh medium for another day. Genomic DNA was extracted for PEM-seq analysis.

HEK293T cells were cultured in DMEM (Gibco) supplemented with 10% fetal bovine

serum (ExCell Bio), penicillin-streptomycin (Gibco), and L-glutamine solution (Sigma) at

37 °C with 5% CO_2. HEK293T cells cultured in 10-cm dishes were transfected with *Sp*Cas9-

gRNA plasmid targeting *c-MYC* locus (15 μg) by calcium phosphate, with or without

additional mitoTALEN plasmid targeting the *ND4* locus (12 μg *ND4*-L1397C-LEFT+12 μg

ND4-L1397C-RIGHT). Regarding the mitochondrial stress treatment, 15 μg plasmid

expressing *Sp*Cas9 targeting *c-MYC* locus was transfected into HEK293T cells by calcium

phosphate. The transfected cells were then treated with 10 μM CCCP for 1 day and then

cultured with fresh medium for another day, or treated with 500 μM paraquat for 2 days.

After that, cells were harvested and the genomic DNA was extracted for PEM-seq analysis.

To determine the occurrence of mtDNA-nuclear DNA fusion caused by mtDNA editors.

HEK293T cells were transfected with mitoTALEN targeting *ND5.1* (20 μg *ND5.1*-L1397N-

LEFT + 20 μg *ND5.1*-L1397N-RIGHT) in 10-cm dishes for 3 days. Regarding DdCBEs,

HEK293T cells were also transfected with DdCBE plasmids targeting *ND4* locus (20 μg

ND4-L1397C-LEFT + 20 μg *ND4*-L1397C-RIGHT) or *ND5.3* locus (20 μg *ND5.3*-L1397C-

LEFT + 20 μg *ND5.3*-L1397C-RIGHT) for 3 days. After transfection, cells were harvested

and permeabilized with NP-40 buffer (10 mM HEPES-NaOH, pH 7.6; 10 mM KCl; 0.1 mM

EDTA; 0.3% NP-40) for 10 min, followed by centrifugation for nuclei collection and

genomic DNA extraction.

For the TREX1/2 co-expression analysis, HEK293T cells were transfected with *Sp*Cas9

plasmid targeting the *c-MYC* locus (15 μg) and DdCBE-*ND4*-L1397C plasmids (20 μg +

20 μg) containing TREX1n/TREX2 at the C-terminus of the UGI of the left arm of DdCBE.

Besides, DdCBE-*ND4*-L1397C plasmids and MTS-tagged TREX1n/TREX2 driven by a

separate promoter were also introduced into HEK293T cells. All sample cells were collected

at 3 days post-transfection.

PEM-seq library preparation and analysis

The PEM-seq libraries were prepared according to the previously reported

procedures(Liu et al., 2022). Briefly, 50 μg of genomic or nuclear DNA from HEK293T

cells, or 20 μg of genomic DNA from mES cells was generally required for each library.

PEM-seq libraries were sequenced on Illumina Hiseq platforms, 2x150 bp. Primers used in

this study were listed in the Key Resource Table.

The PEM-seq data was analyzed using the PEM-Q pipeline and aligned to the hg38

genome assembly as described(Liu et al., 2022). To identify mtDNA fusions, we only kept

chimeric reads mapped to "chrM" with a MAPQ >30. Similarly, when analyzing PEM-seq

data cloned from the mtDNA, we only kept chimeric reads mapped to the nuclear DNA with

a MAPQ >30 and then removed those reads mapped to repetitive genomic regions annotated
in the UCSC database.

Insertion-enriched target sequencing and analysis

Genomic DNA (gDNA) from cells with mitochondrial stresses or mitoTALEN
treatment were subjected to both PEM-seq and insertion-enriched target sequencing analysis.
Regarding insertion-enriched target sequencing, 2 μg gDNA was amplified by two primers
(sequences were listed in the Key Resource Table) flanking the CRISPR-Cas9 target site and
Taq DNA polymerase in 100 μL PCR mixture and then performed as follows: 95℃, 5 min;
95℃, 30 second, 59℃, 30 seconds, 72℃, 4 min, 20 cycles; 72℃, 5 min. DNA products were
size selected by AMpure beads to keep insertions. After that, another two primers (sequences
were listed in the Key Resource Table) were used to further enrich DNA products on the
Cas9-editing site by nested PCR, and the program was set as follows: 95℃, 3 min; 95℃, 30
seconds, 59℃, 30 second, 72℃, 3 min, 15 cycles; 72℃, 5 min. A second size selection was
performed to keep insertions by AMpure beads and the recovered DNA was tagged with
Illumina adapter sequences. Library DNA was sequenced on Hi-seq platforms, 2x150 bp.

Raw reads were aligned to the human reference genome, assembly hg38, by BWA-
MEM with the default parameters. The chimera reads, with MAPQ>30 for each segment,
were kept for insertion identification. Reads beginning at the primer start sites, covering at
least 40 bp of both upstream and downstream of the Cas9-editing sites, and containing
insertion fragments of less than 2000 bp were potential insertions after genome editing.
Insertions with ≥4 counts were kept for further analysis.

Data availability

All sequencing data presented in the present study have been deposited in the NCBI Gene
Expression Omnibus (GEO) under the accession number GSE226884. The PEM-Q pipeline
is available at https://github.com/JiazhiHuLab.

Contributions

J.H. conceived and supervised the project. J.W., Y.L., and J.H. designed the experiments;
J.W., Y.L., L.O., T.G., J.L., S.Y., C.X., and J.Y. performed the experiments; J.W., Y.L., Z.Z.,
X.L., M.L., and J.H. analyzed the data; and J.W., Y.L., Y.T., and J.H. wrote the paper.

Funding

This work was supported by the Ministry of Agriculture and Rural Affairs of China, the National Key R&D Program of China (2022YFC3400201), and the NSFC grants (32122018 and 31771485 to J.H.). J.H. is an investigator at the PKU-TSU Center for Life Sciences.

Conflict of Interests

The authors declare no competing interests.

Acknowledgments

We acknowledge all members of the Hu laboratory for their critical comments and helpful discussions. DdCBE plasmids were kind gifts from Dr. Chengqi Yi at Peking University. We also thank the Flow Cytometry Core at the National Center for Protein Sciences, Peking University, for the technical help.

References

Anzalone, A. V., Koblan, L. W., Liu, D. R. (2020). Genome editing with CRISPR–Cas nucleases, base editors, transposases and prime editors. Nature Biotechnology *38*, 824-844.

Bacman, S. R., Kauppila, J. H. K., Pereira, C. V., Nissanka, N., Miranda, M., Pinto, M., Williams, S. L., Larsson, N.-G., Stewart, J. B., Moraes, C. T. (2018). MitoTALEN reduces mutant mtDNA load and restores tRNAAla levels in a mouse model of heteroplasmic mtDNA mutation. Nature Medicine *24*, 1696-1700.

Bacman, S. R., Williams, S. L., Pinto, M., Peralta, S., Moraes, C. T. (2013). Specific elimination of mutant mitochondrial genomes in patient-derived cells by mitoTALENs. Nature Medicine *19*, 1111-1113.

Barrera-Paez, J. D., Moraes, C. T. (2022). Mitochondrial genome engineering coming-of-age. Trends in Genetics *38*, 869-880.

Borensztajn, K., Chafa, O., Alhenc-Gelas, M., Salha, S., Reghis, A., Fischer, A. M., Tapon-Bretaudière, J. (2002). Characterization of two novel splice site mutations in human factor VII gene causing severe plasma factor VII deficiency and bleeding diathesis. British journal of haematology *117*, 168-171.

Bravo, J. P. K., Liu, M.-S., Hibshman, G. N., Dangerfield, T. L., Jung, K., Mccool, R. S., Johnson, K. A., Taylor, D. W. (2022). Structural basis for mismatch surveillance by CRISPR–Cas9. Nature *603*, 343-347.

Cho, S.-I., Lee, S., Mok, Y. G., Lim, K., Lee, J., Lee, J. M., Chung, E., Kim, J.-S. (2022). Targeted A-to-G base editing in human mitochondrial DNA with programmable deaminases. Cell *185*, 1764-1776.e1712.

Dayama, G., Emery, S. B., Kidd, J. M., Mills, R. E. (2014). The genomic landscape of polymorphic human nuclear mitochondrial insertions. Nucleic Acids Res *42*, 12640-12649.

Fessler, E., Eckl, E.-M., Schmitt, S., Mancilla, I. A., Meyer-Bender, M. F., Hanf, M., Philippou-Massier, J., Krebs, S., Zischka, H., Jae, L. T. (2020). A pathway coordinated by DELE1 relays mitochondrial stress to the cytosol. Nature *579*, 433-437.

Goldin, E., Stahl, S., Cooney, A. M., Kaneski, C. R., Gupta, S., Brady, R. O., Ellis, J. R.,
Schiffmann, R. (2004). Transfer of a mitochondrial DNA fragment to MCOLN1
causes an inherited case of mucolipidosis IV. Hum Mutat *24*, 460-465.
Gray, M. W., Burger, G., Lang, B. F. (1999). Mitochondrial evolution. Science *283*, 1476-
1481.
Hazkani-Covo, E., Zeller, R. M., Martin, W. (2010). Molecular poltergeists: mitochondrial
DNA copies (numts) in sequenced nuclear genomes. PLoS Genet *6*, e1000834.
Hemphill, W. O., Perrino, F. W. (2019). Measuring TREX1 and TREX2 exonuclease
activities. Methods Enzymol *625*, 109-133.
Hsu, P. D., Lander, E. S., Zhang, F. (2014). Development and applications of CRISPR-Cas9
for genome engineering. Cell *157*, 1262-1278.
Jiang, F., Doudna, J. A. (2017). CRISPR-Cas9 Structures and Mechanisms. Annu Rev
Biophys *46*, 505-529.
Ju, Y. S., Tubio, J. M., Mifsud, W., Fu, B., Davies, H. R., Ramakrishna, M., Li, Y., Yates, L.,
Gundem, G., Tarpey, P. S.*, et al.* (2015). Frequent somatic transfer of mitochondrial
DNA into the nuclear genome of human cancer cells. Genome Res *25*, 814-824.
Kim, J., Gupta, R., Blanco, L. P., Yang, S., Shteinfer-Kuzmine, A., Wang, K., Zhu, J., Yoon,
H. E., Wang, X., Kerkhofs, M.*, et al.* (2019). VDAC oligomers form mitochondrial
pores to release mtDNA fragments and promote lupus-like disease. Science *366*,
1531-1536.
Kopinski, P. K., Singh, L. N., Zhang, S., Lott, M. T., Wallace, D. C. (2021). Mitochondrial
DNA variation and cancer. Nature Reviews Cancer *21*, 431-445.
Kosicki, M., Tomberg, K., Bradley, A. (2018). Repair of double-strand breaks induced by
CRISPR-Cas9 leads to large deletions and complex rearrangements. Nat Biotechnol
36, 765-771.
Lebedin, M., Foglierini, M., Khorkova, S., Vázquez García, C., Ratswohl, C., Davydov, A.
N., Turchaninova, M. A., Daubenberger, C., Chudakov, D. M., Lanzavecchia, A., De
La Rosa, K. (2022). Different classes of genomic inserts contribute to human antibody
diversity. Proceedings of the National Academy of Sciences *119*, e2205470119.
Liu, G., Lin, Q., Jin, S., Gao, C. (2021). The CRISPR-Cas toolbox and gene editing
technologies. Molecular cell.
Liu, Y., Yin, J., Gan, T., Liu, M., Xin, C., Zhang, W., Hu, J. (2022). PEM-seq
comprehensively quantifies DNA repair outcomes during gene-editing and DSB
repair. STAR Protocols *3*, 101088.
Mishmar, D., Ruiz-Pesini, E., Brandon, M., Wallace, D. C. (2004). Mitochondrial DNA-like
sequences in the nucleus (NUMTs): insights into our African origins and the
mechanism of foreign DNA integration. Hum Mutat *23*, 125-133.
Mok, B. Y., De Moraes, M. H., Zeng, J., Bosch, D. E., Kotrys, A. V., Raguram, A., Hsu, F.,
Radey, M. C., Peterson, S. B., Mootha, V. K.*, et al.* (2020). A bacterial cytidine
deaminase toxin enables CRISPR-free mitochondrial base editing. Nature *583*, 631-
637.
Mok, B. Y., Kotrys, A. V., Raguram, A., Huang, T. P., Mootha, V. K., Liu, D. R. (2022).
CRISPR-free base editors with enhanced activity and expanded targeting scope in
mitochondrial and nuclear DNA. Nature Biotechnology *40*, 1378-1387.
Nissanka, N., Bacman, S. R., Plastini, M. J., Moraes, C. T. (2018). The mitochondrial DNA
polymerase gamma degrades linear DNA fragments precluding the formation of
deletions. Nat Commun *9*, 2491.
O'malley, J., Kumar, R., Inigo, J., Yadava, N., Chandra, D. (2020). Mitochondrial Stress
Response and Cancer. Trends in Cancer *6*, 688-701.

Riley, J. S., Quarato, G., Cloix, C., Lopez, J., O'prey, J., Pearson, M., Chapman, J., Sesaki, H., Carlin, L. M., Passos, J. F.*, et al.* (2018). Mitochondrial inner membrane permeabilisation enables mtDNA release during apoptosis. The EMBO journal *37*, e99238.

Schmid-Burgk, J. L., Gao, L., Li, D., Gardner, Z., Strecker, J., Lash, B., Zhang, F. (2020). Highly Parallel Profiling of Cas9 Variant Specificity. Molecular cell *78*, 794-800.e798.

Schwartz, M., Vissing, J. (2002). Paternal Inheritance of Mitochondrial DNA. New England Journal of Medicine *347*, 576-580.

Shay, J. W., Baba, T., Zhan, Q. M., Kamimura, N., Cuthbert, J. A. (1991). HeLaTG cells have mitochondrial DNA inserted into the c-myc oncogene. Oncogene *6*, 1869-1874.

Silva-Pinheiro, P., Minczuk, M. (2022). The potential of mitochondrial genome engineering. Nature Reviews Genetics *23*, 199-214.

Skopelja-Gardner, S., An, J., Elkon, K. B. (2022). Role of the cGAS–STING pathway in systemic and organ-specific diseases. Nature Reviews Nephrology *18*, 558-572.

Stadtmauer, E. A., Fraietta, J. A., Davis, M. M., Cohen, A. D., Weber, K. L., Lancaster, E., Mangan, P. A., Kulikovskaya, I., Gupta, M., Chen, F.*, et al.* (2020). CRISPR-engineered T cells in patients with refractory cancer. Science *367*, eaba7365.

Tubbs, A., Nussenzweig, A. (2017). Endogenous DNA Damage as a Source of Genomic Instability in Cancer. Cell *168*, 644-656.

Turner, C., Killoran, C., Thomas, N. S. T., Rosenberg, M., Chuzhanova, N. A., Johnston, J., Kemel, Y., Cooper, D. N., Biesecker, L. G. (2003). Human genetic disease caused by de novo mitochondrial-nuclear DNA transfer. Human Genetics *112*, 303-309.

Wei, W., Pagnamenta, A. T., Gleadall, N., Sanchis-Juan, A., Stephens, J., Broxholme, J., Tuna, S., Odhams, C. A., Ambrose, J. C., Baple, E. L.*, et al.* (2020). Nuclear-mitochondrial DNA segments resemble paternally inherited mitochondrial DNA in humans. Nature Communications *11*, 1740.

Wei, W., Schon, K. R., Elgar, G., Orioli, A., Tanguy, M., Giess, A., Tischkowitz, M., Caulfield, M. J., Chinnery, P. F. (2022a). Nuclear-embedded mitochondrial DNA sequences in 66,083 human genomes. Nature *611*, 105-114.

Wei, Y., Li, Z., Xu, K., Feng, H., Xie, L., Li, D., Zuo, Z., Zhang, M., Xu, C., Yang, H., Zuo, E. (2022b). Mitochondrial base editor DdCBE causes substantial DNA off-target editing in nuclear genome of embryos. Cell discovery *8*, 27.

Willett-Brozick, J. E., Savul, S. A., Richey, L. E., Baysal, B. E. (2001). Germ line insertion of mtDNA at the breakpoint junction of a reciprocal constitutional translocation. Human Genetics *109*, 216-223.

Woischnik, M., Moraes, C. T. (2002). Pattern of organization of human mitochondrial pseudogenes in the nuclear genome. Genome Res *12*, 885-893.

Wu, J., Zou, Z., Liu, Y., Liu, X., Zhangding, Z., Xu, M., Hu, J. (2022). CRISPR/Cas9-induced structural variations expand in T lymphocytes in vivo. Nucleic Acids Res *50*, 11128-11137.

Xian, H., Watari, K., Sanchez-Lopez, E., Offenberger, J., Onyuru, J., Sampath, H., Ying, W., Hoffman, H. M., Shadel, G. S., Karin, M. (2022). Oxidized DNA fragments exit mitochondria via mPTP- and VDAC-dependent channels to activate NLRP3 inflammasome and interferon signaling. Immunity *55*, 1370-1385.e1378.

Xin, C., Yin, J., Yuan, S., Ou, L., Liu, M., Zhang, W., Hu, J. (2022). Comprehensive assessment of miniature CRISPR-Cas12f nucleases for gene disruption. Nature Communications *13*, 5623.

Yin, J., Fang, K., Gao, Y., Ou, L., Yuan, S., Xin, C., Wu, W., Wu, W. W., Hong, J., Yang,
H., Hu, J. (2022a). Safeguarding genome integrity during gene-editing therapy in a
mouse model of age-related macular degeneration. Nat Commun *13*, 7867.
Yin, J., Hu, J. (2022). The origin of unwanted editing byproducts in gene editing. Acta
Biochim Biophys Sin (Shanghai) *54*, 767-781.
Yin, J., Lu, R., Xin, C., Wang, Y., Ling, X., Li, D., Zhang, W., Liu, M., Xie, W., Kong, L.*, et*
al. (2022b). Cas9 exo-endonuclease eliminates chromosomal translocations during
genome editing. Nat Commun *13*, 1204.
Yuan, Y., Ju, Y. S., Kim, Y., Li, J., Wang, Y., Yoon, C. J., Yang, Y., Martincorena, I.,
Creighton, C. J., Weinstein, J. N.*, et al.* (2020). Comprehensive molecular
characterization of mitochondrial genomes in human cancers. Nat Genet *52*, 342-352.
Zhang, W., Yin, J., Zhang-Ding, Z., Xin, C., Liu, M., Wang, Y., Ai, C., Hu, J. (2021). In-
depth assessment of the PAM compatibility and editing activities of Cas9 variants.
Nucleic Acids Research.
Zuo, E., Sun, Y., Wei, W., Yuan, T., Ying, W., Sun, H., Yuan, L., Steinmetz, L. M., Li, Y.,
Yang, H. (2019). Cytosine base editor generates substantial off-target single-
nucleotide variants in mouse embryos. Science *364*, 289-292.

 Figure legends

Figure 1. Identification of mtDNA fusion to nuclear target sites of CRISPR-Cas system.

A. Schematic of mtDNA-nuclear DNA fusion captured by PEM-seq. The primer (purple

arrow) located adjacent to the CRISPR-Cas9 target site (scissor) on the nuclear DNA is

used to clone editing products (orange line), and the chimeric reads harboring nuclear

DNA around the editing site (black line) and mtDNA (red line), with MAPQ >30, were

identified as mtDNA-nuclear DNA fusion.

B. Box plot showing the frequency of mtDNA-nuclear DNA fusion events out of editing

events at CRISPR-Cas target sites (colorful dots) under editing of CRISPR-Cas enzymes.

C. Circos plot showing the fusion junctions on mtDNA (MT) and the indicated CRISPR-

Cas9 target sites (colorful triangles) on the nuclear DNA of HEK293T cells. The outer

circle shows the human genome, labeled with numbers or characters. The colorful lines

indicate the fusion between the target site and mtDNA. Annotations of mtDNA are shown

at the bottom.

D. Circos plot showing the fusion junctions on mtDNA (MT) and the indicated CRISPR-

*Lb*Cas12a target sites (colorful triangles) on the nuclear DNA of HEK293T cells.

Legends are described in C.

E. Circos plot showing the fusion junctions on mtDNA (MT) and the indicated CRISPR-

CasMINI target sites (colorful triangles) on the nuclear DNA of HEK293T cells. Legends

are described in C.

F. Box plot showing the frequency of mtDNA-nuclear DNA fusion events out of editing

events at CRISPR-Cas target sites (colorful dots) under editing of *Sp*Cas9 variants.

G. Distribution of mtDNA-nuclear DNA fusion caused by high fidelity Cas9 variants in the

mES cells. Legends are described as depicted in C.

H. Frequency of mtDNA-nuclear DNA fusion at *DNMT1*, *EMX1*, *MYC2*, *RAG1_C*, and

BCL11A loci after editing by Cas9, BE4max, and ABEmax. EMX1 and MYC2 loci were

not targetable by ABEmax. N.D., not determined.

Figure 2. Mitochondrial DNA fuses to CRISPR-Cas9 target sites of the nuclear DNA *ex*

***vivo* and *in vivo*.**

A. Schematic of ex vivo human T cell editing and PEM-seq. Briefly, human primary T cells

isolated from cord blood were activated for 3 days and subsequently transfected with

Cas9/gRNA ribonucleoprotein (RNP) complexes targeting *TRAC*, *TRBC*, and *PDCD1*

loci. Cells were collected after 3, 7, or 14 days and subjected to PEM-seq libraries. In

PEM-seq libraries, all products are labeled with random molecular barcodes (RMB) to

distinguish biological expansion and PCR duplications. mtDNA-nuclear DNA fusions

originate from the same mtDNA junction are labeled in red.

B. Distribution of mtDNA-nuclear DNA fusions after CRISPR-Cas9 editing for 3, 7, and 14

days in human CAR T cells. Legends are described in Fig. 1B.

C. Stacked bar plot showing the copy number of each mtDNA-nuclear DNA fusion type

identified from *TRAC*, *TRBC*, and *PDCD1* bait after Cas9 editing for 3, 7, and 14 days.

Numbers mark the copies of junctions with 3 copies or above.

D. Schematic showing the amplification of mtDNA-nuclear DNA fusion through clonal

expansion of TCR T cells after infusion.

E. Distribution of mtDNA-nuclear DNA fusion in CRISPR-Cas9 treated mouse TCR-T cells

before and post-infusion for 3 weeks. Legends are described as depicted in B. Percentages

show the frequency of each mtDNA-nuclear DNA fusion out of Cas9-induced editing

events.

F. mtDNA-nuclear DNA fusions between *Vegfa* and mtDNA in an AMD mouse model

treated with AAV containing CRISPR-Cas9. Legends are described as depicted in 1B.

G. Distribution of mtDNA integration in the nuclear DNA post base editor (BE3) treatment

in mouse embryos. Data were re-analyzed from GOTI libraries (SRA: SRP119022) (Zuo

et al., 2019). The grey, blue, and red lines indicate integrations detected in both control

and treatment, control-only, and treatment-only, respectively. MT, mtDNA.

Figure 3. Mitochondrial stresses exacerbate mtDNA integration into the nuclear

genome.

A. Mitochondrial stresses inducing mtDNA-nuclear DNA fusions captured by PEM-seq.

Top: cells were transfected with a plasmid containing *Sp*Cas9 and gRNA targeting *c-*

MYC for 8 hours to allow the assembly of *Sp*Cas9 and gRNA. Subsequentially, cells are

treated with CCCP or paraquat for 24 or 48 hours, respectively, and then harvested for

PEM-seq and Insert-seq analysis. Bottom: percentages of mtDNA-nuclear DNA fusions

captured by PEM-seq. Each dot represents a biological replicate. Mean±SD; *t*-test; *,

$p<0.05$.

B. Distribution of mtDNA (MT) fusion junctions captured by the *c-MYC* bait (black triangle
and grey bars) with or without mitochondrial stresses (CCCP, light purple bars; paraquat,
dark purple bars) treatment. Legends of mtDNA annotations are described as depicted in
1B. The inner circles show the number of each mtDNA-nuclear DNA fusion point on
mtDNA in a log scale.
C. Workflow of Insert-seq to enrich insertions (orange lines) at the Cas9 editing site (*c-MYC*
locus). Briefly, two rounds of targeted PCR (purple and red arrows) are used to clone the
editing events around the target site, followed by two rounds of size selection that
enriches insertions.
D. Percentages of mtDNA integrations within total insertions captured by Insert-seq at the *c-*
MYC site in the presence or absence of mitochondrial stresses. Each dot represents a
biological replicate. Mean±SD; *t*-test; *, $p < 0.05$; **, $p < 0.01$.

Figure 4. Mitochondrial editors exacerbate mtDNA integration into the nuclear target
site.

A. Illustration of mitochondrial stresses and mitoTALEN treatment induced mtDNA-nuclear
DNA fusion captured by PEM-seq and Insert-seq. CRISPR-Cas9 (scissor) targets the *c-*
MYC locus in nuclear DNA, and the primer (purple arrow) for PEM-seq is adjacent to the
target site.
B. Frequency of PEM-seq-captured mtDNA and the CRISPR-Cas9 target site fusion
junctions with or without mitoTALEN treatment. Each dot represents a biological
replicate. Mean±SD; *t*-test; **, $p < 0.01$.
C. Distribution of mtDNA (MT) fusion junctions captured by the *c-MYC* bait (black triangle
and grey bars) with or without mitoTALEN (*ND4* site, red triangle and purple bars)
treatment. Legends are described as depicted in 3B.
D. Percentages of mtDNA integrations within total insertions captured by Insert-seq at the *c-*
MYC site. Each dot represents a biological replicate. Mean±SD; *t*-test; *, $p < 0.05$.

Figure 5. Mitochondrial editors result in mtDNA integrations into nuclear DNA.

A. Schematic of mitoTALEN treatment induced mtDNA integration into the nuclear DNA,
which can be captured by a primer (purple arrow) located adjacent to the mitoTALEN
editing window (red line). mitoTALEN generated DSBs within the editing window

between two TALE target sites. Chimeric reads containing *bona fide* mitoTALEN-

induced mtDNA-nuclear DNA fusion should harbor mtDNA bait junctions within the

editing window.

B. Frequency of mtDNA-nuclear DNA fusions with mtDNA bait junctions at each

nucleotide spanning the mitoTALEN target site on *ND5.1* (green boxes), editing window

of mitoTALEN is marked with a red line. Each dot represents a biological replicate.

Mean±SD; *t*-test; *, $p < 0.05$; **, $p < 0.01$.

C. Distribution of mtDNA-nuclear DNA fusions (orange lines) with mtDNA bait junctions

within the editing window of mitoTALEN.

D. Illustration of DdCBE caused mtDNA-nuclear DNA integration. Legends are described

as depicted in A. Chimeric reads containing *bona fide* DdCBE-induced fusion should

harbor mtDNA bait junctions at the editing site.

E. Editing efficiency (C to T or G to A) of DdCBE on *ND4*. The green boxes and the red

triangle indicate the binding site and editing point of DdCBE. Each dot represents a

biological replicate; Mean±SD; *t*-test; ****, $p < 0.0001$.

F. Frequency of mtDNA-nuclear DNA fusions with mtDNA bait junctions at each

nucleotide spanning the DdCBE target sites (green boxes) on ND4. Each dot represents a

biological replicate. Mean±SD; *t*-test; **, $p < 0.01$.

G. Distribution of mtDNA-nuclear DNA fusions (orange lines) with mtDNA ending at the

editing site of DdCBE.

H. Distribution of mtDNA-nuclear DNA fusions posts DdCBE treatment in mouse embryos.

Data were re-analyzed from GOTI libraries (SRA: PRJNA786071) (Wei et al., 2022b).

Legends are described as depicted in 2G. The grey, blue, and red lines indicate fusions

detected in both control and treatment, control-only, and treatment-only, respectively.

Figure 6. TREX1 and TREX2 suppress the transfer of mtDNA into nuclear DNA.

A. Structures of DdCBE with or without TREX1n/TREX2. For the fusion form, TREX1n or

TREX2 was fused to the C-terminal domain of L-1397C-UGI. Regarding separated

TREX1n or TREX2, both nucleases were tagged with mitochondrial targeting sequence

(MTS) on the N-terminal. L-1397C-UGI, left TALE arrays fused to C-terminal DddAtox

half and UGI; R-1397N-UGI, right TALE arrays fused to N-terminal DddAtox half and

UGI; NTD, N-terminal domain; CTD, C-terminal domain.

B. Editing efficiency of DdCBE with or without TREX1n/TREX2 treatment. NC, negative

control. Mean±SD. L, L-1397C-UGI; R, R-1397N-UGI.

C. Fluorescence fold change of mtDNA to nuclear DNA detected by qPCR.

D. Distribution of mtDNA-nuclear DNA fusions (orange lines) with mtDNA bait junctions

ending at the editing site of DdCBE on *ND4*. The number (n) of fusions in each sample is

normalized to the same editing events. L, L-1397C-UGI; R, R-1397N-UGI.

E. Illustration of PEM-seq cloning from the nuclear DNA to assess DdCBE-induced

mtDNA-nuclear DNA fusion with or without TREX1n/TREX2 treatment.

F. Frequency of DdCBE-induced mtDNA fusing with the CRISPR-Cas9 target site with or

without TREX1n/TREX2 treatment. Each dot represents a biological replication.

Mean±SD; *t*-test; *, $p < 0.05$; **, $p < 0.01$; ***, $p < 0.001$. L, L-1397C-UGI; R, R-1397N-

UGI.

Supplementary Figure legends

Figure S1. Identification of mtDNA fusion to nuclear targeting site of CRISPR-Cas
systems.

A. Representative mtDNA integration sequences that are aligned to the mitochondrial

genome.

B. Circos plot showing the fusion junctions on mtDNA (MT) and the indicated *Lb*Cas12a

target sites (colorful triangles) on the nuclear DNA of HEK293T cells. Legends are

described in 1B.

C. Circos plot showing the fusion junctions on mtDNA (MT) and the indicated *Un1*Cas12f,

target sites (colorful triangles) on the nuclear DNA of HEK293T cells. Legends are

described in 1B.

D. Circos plot showing the fusion junctions on mtDNA (MT) and the indicated

CasMINI_ge4.1 target sites (colorful triangles) on the nuclear DNA of HEK293T cells.

Legends are described in 1B.

E. Distribution of mtDNA fusion points identified from CRISPR-Cas systems (1E and 1F).

F. Percentage and length of microhomology identified in chimeric reads containing mtDNA-

nuclear DNA fusion.

Figure S2. Mitochondrial DNA fuses to CRISPR-Cas9 targeting sites of the nuclear

DNA *ex vivo* and *in vivo*.

A. Percentages of mtDNA-nuclear DNA fusions at *TRAC*, *TRBC*, and *PDCD1* in human

CAR-T cells, related to 2B.

B. Sequence logo showing the frequency of nucleotides of random molecular barcodes from

reads in 38 copies of mtDNA-nuclear DNA fusion in infused mouse TCR T cells at a

single junction indicated in 2E.

C. Distribution of mtDNA-nuclear DNA fusions posts base editor treatment in mouse

embryos. Data were re-analyzed from GOTI libraries (SRA: SRP119022) (Zuo et al.,

2019). Legends are described as depicted in 2G.

Figure S3. Using target sequencing for validation of mtDNA integrations.

A. Schematic of Insert-seq captured *bona fide* insertions (orange lines) at the Cas9 editing

sites. Briefly, reads should exactly start at the primer target site and then extend at least

40 bp on the target genome. The length of insertion should be less than 2000 bp, with a

MAPQ >30. Moreover, each insertion must have more than 4 hits.

B. Length distribution of Insert-seq captured insertions. Red line, median length.

Fig. S4 Mitochondrial DNA breaks exacerbate mtDNA integration into nuclear genome.

A. PCR products showing the mtDNA-nuclear DNA fusion induced by Cas9 editing on *c-*

MYC and mitoTALEN targeting *ND4.* Primers targeting nuclear DNA (purple arrow) and

mtDNA (blue arrows) are used, and the expected products are labeled by red triangles.

B. Sanger sequencing showing the fusion between *c-MYC* (green boxes) and *ND4* (blue

boxes).

C. Distribution of Insert-seq captured mtDNA segments that were inserted into *c-MYC*.

D. Illustration of one mtDNA insertion that was captured by both PEM-seq and Insert-seq.

Reads (R1 or R2) from PEM-seq (brown) and Insert-seq (blue) harbor the same insertion

sites. Moreover, reads from Insert-seq shared an 18-bp overlap, while a 32-bp gap was found

between R1 and R2 from PEM-seq. The sequences are listed below with mtDNA sequences

marked in red.

Figure S5. Mitochondrial editors result in mtDNA integrations into nuclear DNA.

A. Frequency of mtDNA-nuclear DNA fusions with mtDNA bait junctions at each nucleotide

spanning the mitoTALEN target site on *ND4*. Each dot represents a biological replicate.

Mean±SD; *t*-test.

B. Distribution of mtDNA-nuclear DNA fusions (orange lines) with mtDNA ending at the

editing site of DdCBE. Legends are described as depicted in 5G.

C. Representative sequence of *bona fide* DdCBE-induced mtDNA-nuclear DNA fusion. The

red arrow marks the DdCBE caused G to A mutation. Light grey and green boxes mark the

sequence aligned to mitochondrial and nuclear DNA.

D. Editing efficiency of DdCBE on *ND5.3*. Three biological replicates; Mean±SD; *t*-test; ****,

p< 0.0001.

E. Distribution of mtDNA-nuclear DNA fusions (orange lines) with mtDNA bait junctions

at the editing site of DdCBE.

F. Frequency of mtDNA-nuclear DNA fusions with mtDNA bait junctions at each

nucleotide spanning the DdCBE target site on *ND5.3*. Each dot represents a biological

replicate. Mean±SD; *t*-test.

G. Distribution of mtDNA-nuclear DNA fusion post mitochondrial base editor treatment in

mouse embryos (SRA: PRJNA786071) (Wei et al., 2022b). Legends are described as

depicted in 5H.

Fig. 1 Identification of mtDNA fusion to nuclear target sites of CRISPR-Cas systems.

Fig. 2 Mitochondrial DNA fuses to CRISPR-Cas9 target sites of the nuclear DNA *ex vivo* and *in vivo*.

Fig. 3 Mitochondrial and stresses exacerbate mtDNA integration into the nuclear genome.

Fig. 4 Mitochondrial editors exacerbate mtDNA integration into the nuclear target site.

Fig. 5 Mitochondrial editors result in mtDNA integrations into the nuclear DNA.

Fig. 6 TREX1 and TREX2 suppress the transfer of mtDNA into the nuclear DNA.

A

Prey sequence from bait *VEGFA_1*:

```
TTCGTTACATGGTCCATCATAGAATTCTCACTGTGATATATAAACTCAGACCCAAACAT
---------------------------------------------------------- chrM, 12619, 100%
-------G--A--------------------A---------G--------------- chr5, 134926119, 93.3%
-----C-----A-------------------A--G------G--------------- chr5, 100048445, 91.6%
```

Prey sequence from bait *NLRC4*:

```
CTTCCACCCCCTAGCAGAAAATAGCCCACTAATCCAAACTCTAACACTATGCTTAGGCGCT
----------------------------------------------------------- chrM, 13122, 100%
-----------------G----A------------------C--------C------- chr5, 134925614, 93.5%
-----------------G-C--C-------C--C-------T--------- chr5, 100047940, 90.2%
```

Fig. S1 Identification of mtDNA fusion to nuclear target sites of CRISPR-Cas systems.

Fig. S2 Mitochondrial DNA fuses to CRISPR-Cas9 target sites of the nuclear DNA *ex vivo* and *in vivo*.

A

Ref.

① Filter 1:

Primer
Start

Insertions, ≤ 2000 bp

Primer
Start

R1

R2

Targeted region,
≥ 40 bp

Targeted region,
≥ 40 bp

② Filter 2:

Counts: ≥ 4

B

Length of insertions (bp)

2000

1000

N=4,706;
Median=163 bp

Cas9

Fig. S3 Using target sequencing for validation of mtDNA integrations.

A

Cas9: *c-MYC*

P1

nuclear DNA

+

mitoTALEN: *ND4*

P2
mtDNA
P3

P1+P2 P1+P3

Cas9 + + + +
mitoTALEN − + − +

B

Translocation1 (c-MYC+ND4-left):
c-MYC (+)
CATTCCAAATAATCAGGAAAGAACTTTAGGAGAATGTGGTTACTAGCACAGAGAGTTCTC
ND4: ChrM (−)

Translocation2 (c-MYC+ND4-left):
c-MYC (+)
CAGGAAAGAACTTTAGGGATGTGGAGCTATGTGGTTACTAGCACAGAGAGTTCTCC
ND4: ChrM (−)

Translocation3 (c-MYC+ND4-right):
c-MYC (+)
TCCAAATAATCAGGAAAGAACTTTAGGGATTCAAATATCACTCTCCTACTTACAGGACTC
ND4: ChrM (+)

C

Cas9
7 of 10,098

Cas9 + mitoTALEN-ND4
14 of 8,531

c-MYC
ND4

MT

c-MYC

Insertion fragments from mtDNA

D

c-MYC, Left, 89 bp   *ChrM,* insertion, 114 bp   *c-MYC,* Right, 90 bp

Insertion

Targeted
Insertion site
R1
18 bp
R2

PEM-seq
Junction site
R1
R2
32 bp   16 bp

Target sequencing: @E00572:537:HHY7LCCX2:3:1102:28229:61960
R1:
CCTCAGAATAGGAGAGAGTGTTCCTAGCAAGGAAAACAACTCTCC
ATTCCAAATAATCAGGAAAGAACTTTAGGGATGTGGAGCTTGGCA
ACTATGCCTAGAAGGAATAATACTATCGCTGTTCATTATAGCTACT
CTCATAACCC
R2:
AGAGCCATTCTCTGGCTCAGTAAAGGTAAGAGCGGCCTAATAGG
CACTTGGAATGGTTCCTTTCTATTCCCATAGCGGCAAAGACTAGT
ATGGCAATAGGCACAATATTGGCTAAGAGGGAGTGGGTGTTGAG
GGTTATGAGAGTAGCTA

PEM-seq: @E00572:537:HHY7LCCX2:3:2101:11749:13439
R1:
CCTCAGAATAGGAGAGAGTGTTCCTAGCAAGGAAAACAACTCTCC
ATTCCAAATAATCAGGAAAGAACTTTAGGGATGTGGAGCTTGGCA
ACTATGCCTAGAAGGAATAATACTATCGCTGTTCATTATAGCTACT
CTCATAACCC
R2:
AGAGCCATTCTCTGGCTCAGTAAAGGTAAGAGCGGCCTAATAGG
CACTTGGAATGGTTCCTTTCTATTCCCATAGCGGCAAAGACTAGT
ATGGCAATAGG

Fig. S4 Mitochondrial DNA breaks exacerbate mtDNA integration into nuclear genome.

Fig. S5 Mitochondrial editors result in mtDNA integrations into nuclear DNA.