The International Weed Genomics Consortium: Community Resources for Weed Genomics Research

Jacob S. Montgomery[1], Sarah Morran[1], Dana R. MacGregor[2], J. Scott McElroy[3], Paul Neve[4], Célia Neto[4], Martin M. Vila-Aiub[5], Maria Victoria Sandoval[5], Analia I. Menéndez[6], Julia M. Kreiner[7], Longjiang Fan[8], Ana L. Caicedo[9], Peter J. Maughan[10], Bianca Assis Barbosa Martins[11], Jagoda Mika[11], Alberto Collavo[11], Aldo Merotto Jr.[12], Nithya K. Subramanian[13] , Muthukumar V. Bagavathiannan[13], Luan Cutti[14], Md. Mazharul Islam[15], Bikram S Gill[16], Robert Cicchillo[17], Roger Gast[17], Neeta Soni[17], Terry R. Wright[18], Gina Zastrow-Hayes[18], Gregory May[18], Jenna M. Malone[19], Deepmala Sehgal[20], Shiv Shankhar Kaundun[20], Richard P. Dale[20], Barend Juan Vorster[21], Bodo Peters[11], Jens Lerchl[22], Patrick J. Tranel[23], Roland Beffa[24], Alexandre Fournier-Level[25], Mithila Jugulam[15], Kevin Fengler[18], Victor Llaca[18], Eric L. Patterson[14], Todd Gaines[1]

[1]Department of Agricultural Biology, Colorado State University, Fort Collins, Colorado, USA.

[2]Protecting Crops and the Environment, Rothamsted Research, Harpenden, Hertfordshire, UK.

[3]Department of Crop, Soil, and Environmental Sciences, Auburn University, Auburn, Alabama, USA.

[4]Department of Plant and Environmental Sciences, University of Copenhagen, Taastrup, Denmark.

[5]IFEVA-Conicet-Department of Ecology, University of Buenos Aires, Buenos Aires, Argentina.

[6]Department of Ecology, Faculty of Agronomy, University of Buenos Aires, Buenos Aires, Argentina.

[7]Department of Botany, The University of British Columbia, Vancouver, British Columbia,

Canada.

[8]Institute of Crop Sciences, Zhejiang University, Hangzhou, China.

[9]Department of Biology, University of Massachusetts Amherst, Massachusetts, USA.

[10]Department of Plant and Wildlife Sciences, Brigham Young University, Provo, Utah, USA.

[11]Bayer AG, Weed Control Research, Frankfurt, Germany.

[12]Department of Crop Sciences. Federal University of Rio Grande do Sul, Porto Alegre, Rio

Grande do Sul, Brazil.

[13]Department of Soil and Crop Sciences, Texas A&M University, College Station, Texas, USA.

[14]Department of Plant, Soil and Microbial Sciences, Michigan State University, East Lansing,

Michigan, USA.

[15]Department of Agronomy, Kansas State University, Manhattan, Kansas, USA.

[16]Department of Plant Pathology, Kansas State University, Manhattan, Kansas, USA.

[17]Crop Protection Discovery and Development, Corteva Agriscience, Indianapolis, Indiana,

USA.

[18]Genome Center of Excellence, Corteva Agriscience, Johnston, Iowa, USA.

[19]School of Agriculture, Food and Wine, University of Adelaide, Glen Osmond, South Australia,

Australia.

[20]Jealott's Hill International Research Centre, Syngenta Ltd, Bracknell, Berkshire, UK.

[21]Department of Plant and Soil Sciences, University of Pretoria, Pretoria, South Africa

[22]BASF SE, Ludwigshafen am Rhein, Germany.

[23]Department of Crop Sciences, University of Illinois, Urbana, Illinois, USA.

[24]Senior Scientist Consultant, Herbicide Resistance Action Committee / CropLife International,

46   Liederbach, Germany.

47   [25]University of Melbourne, Parkville, Victoria, Australia.

49   Author for correspondence: Todd Gaines, Associate Professor, Colorado State University, 1177

50   Campus Delivery, Fort Collins, CO 80523, USA. (Email: todd.gaines@colostate.edu)

52   Keywords: Weed science, Reference genomes, Rapid adaptation, Herbicide resistance, Public

53   resources

Abstract

The International Weed Genomics Consortium is a collaborative group of researchers focused on

developing genomic resources for the study of weedy plants. Weeds are attractive systems for

basic and applied research due to their impacts on agricultural systems and capacity to swiftly

adapt in response to anthropogenic selection pressures. Our goal is to use genomic information to

develop sustainable and effective weed control methods and to provide insights about biotic and

abiotic stress tolerance to assist crop breeding. Here, we outline resources under development by

the consortium and highlight areas of research that will be impacted by these enabling resources.

Introduction

Each year globally, agricultural producers and landscape managers spend billions of US

dollars [1, 2] and countless hours attempting to control weedy plants and reduce their adverse

effects. These management methods range from low-tech (e.g., pulling plants from the soil by

hand) to extremely high-tech (e.g., computer vision-controlled spraying of herbicides).

Regardless of technology level, effective control methods serve as strong selection pressures on

weedy plants, and often result in rapid evolution of weed populations resistant to such methods

[3-7]. Thus, humans and weeds have been locked in an arms race, where humans develop new or

improved control methods and weeds adapt and evolve to circumvent such methods.

Applying genomics to weed science will enable the development of more sustainable and

effective control methods and offer a unique opportunity to study rapid adaptation and

evolutionary rescue of diverse weedy species in the face of widespread and powerful selective

pressures. Furthermore, lessons learned from these studies may also help to improve crop

breeding efforts in the face of our ever-changing climate. While other research fields have used

genetics and genomics to uncover the basis of many biological traits [8-11] and to understand

how ecological factors affect evolution [12, 13], the field of weed science has lagged behind in

the development of genomic tools essential for such studies [14]. As research in human and crop

genetics pushes into the era of pangenomics, (i.e., multiple chromosome scale genome

assemblies for a single species [15, 16]) publicly available genomic information is still lacking or

severely limited for the majority of weed species. In fact, a recent review of current weed

genomes identified just 26 weed species with sequenced genomes [17] – many assembled to a

sub-chromosome level.

The International Weed Genomics Consortium (IWGC) is an open collaboration between

academic, government, and industry researchers focused on producing genomic tools for weedy

species from around the world. Through this collaboration, our initial aim is to provide

chromosome-level reference genome assemblies for at least 50 important weedy species from

across the globe. Each genome will include annotation of gene models and repetitive elements

and will be freely available through public databases with no intellectual property restrictions.

Species were chosen based on member input, economic impact, and global prevalence (Figure

1). Additionally, future funding of the IWGC will focus on supplementing these reference

genomes with tools that increase their utility.

The IWGC held its first conference in Kansas City, Missouri, USA in September of 2021. At

this meeting, guest speakers highlighted successful examples of using genomics to address

questions in weed science [5, 18-20]. Training workshops taught commonly used bioinformatic

pipelines, and oral and poster sessions showcased current research activities in weed genomics.

At the conclusion of this meeting, attendees participated in a forward-looking discussion about

the future of genomics in weed science and how the IWGC can help facilitate its successful

implementation. In this paper, we summarize the goals of the IWGC and how we plan to provide

support around the resources being developed to ensure they are widely accessible and utilized

by the research community. We go on to highlight areas of research where these tools can be

applied with hopes of attracting researchers from other fields to integrate weed science with the

many other research areas where genomic tools are being successfully utilized, enabling new

research towards adaptation, evolution, herbicide resistance, and genome biology.

Development of Weed Genomics Resources by the IWGC

Reference Genomes and Data Analysis Tools

The first objective of the IWGC is to provide high quality genomic resources for

agriculturally important weeds. The IWGC therefore created two main resources for information

about, access to, or analysis of weed genomic data (Figure 1). The IWGC website [21]

communicates the status and results of genome sequencing projects, information on training and

funding opportunities, upcoming events, and news in weed genomics. It also contains details of

all sequenced species including genome size, ploidy, chromosome number, herbicide resistance

status, and reference genome assembly statistics. The IWGC either compiles existing data on

genome size, ploidy, and chromosome number, or obtains the data using flow cytometry and

cytogenetics (Figure 1; Additional File 1). Through this website, users can create an account to

access our second main resource, an online genome database called WeedPedia. WeedPedia

hosts IWGC-generated and other relevant publicly accessible genomic data as well as a suite of

bioinformatic tools. Unlike what is available for other fields, weed science did not have a

centralized hub for genomics information, data, and analysis prior to the IWGC. Our intention in

creating WeedPedia is to encourage collaboration and equity of access to information across the

research community. Importantly, all genome assemblies and annotations from the IWGC (Table

1), along with the raw data used to produce them, will be made available through NCBI

GenBank. Upon completion of a one-year sponsoring member data confidentiality period for

each species, scientific teams within the IWGC produce the first genome-wide investigation to

submit for publication including whole genome level analyses on genes, gene families, and

repetitive sequences, and comparative analysis with other species. Genome assemblies and data

will be publicly available through NCBI as part of these initial publications for each species.

WeedPedia is a cloud-based omics database management platform built from the software

'CropPedia', and licensed from KeyGene (Wageningen, The Netherlands). The interface allows

users to access, visualize, and download genome assemblies along with structural and functional

annotation. The platform includes a genome browser, comparative map viewer, pangenome

tools, RNA-sequencing data visualization tools, genetic mapping and marker analysis tools, and

alignment capabilities that allow searches by keyword or sequence. Additionally, genes encoding

known target sites of herbicides have been specially annotated, allowing users to quickly identify

and compare these genes of interest. The platform is flexible, making it compatible with future

integration of other data types such as epigenetic or proteomic information. As an online

platform with a graphical user interface, WeedPedia provides user-friendly, intuitive tools that

encourage users to integrate genomics into their research. We aspire for WeedPedia to mimic the

success of other public genomic databases such as NCBI, CoGe, Phytozome, InsectBase, and

Mycocosm to name a few. WeedPedia currently hosts reference genomes for 40 species with

additional genomes in the pipeline to reach a currently planned total of reference genomes for 53

species (Table 1). These genomes include both *de novo* reference genomes generated or in

progress by the IWGC (31 species; Table 1), and publicly available genome assemblies of 22

weedy or related species (Table 2). As of October 2023, WeedPedia has over 300 registered

users representing 27 countries spread across 6 continents.

The IWGC reference genomes are generated in partnership with the Corteva Agriscience

Genome Center of Excellence (Johnston, Iowa) using a combination of single molecule long read

sequencing, optical genome maps, and chromosome conformation mapping. This strategy has

yielded highly contiguous, phased, chromosome-level assemblies for 20 weed species, with

additional assemblies currently in progress (Table 1). The IWGC assemblies have been

completed as single or haplotype-resolved double-haplotype pseudomolecules in inbreeding and

outbreeding species, respectively, with multiple genomes being near gapless. For example, the *de*

novo assemblies of the allohexaploids *Conyza sumatrensis* and *Chenopodium album* have all

chromosomes captured in single scaffolds and most chromosomes being gapless from telomere

to telomere. Complementary full-length isoform (IsoSeq) sequencing of RNA collected from

diverse tissue types and developmental stages assists in the development of gene models during

annotation. Finally, the use of PacBio Revio has enabled the re-sequencing of 80 relevant

accessions, which is enabling initial pangenomic analysis for some of the IWGC-selected

species.

As with accessibility of data, a core objective of the IWGC is to facilitate open access to

sequenced germplasm for all featured species. Historically, the weed science community has

rarely shared or adopted standard germplasm (e.g., specific weed accessions). The IWGC has

selected a specific accession of each species for reference genome assembly (typically

susceptible to herbicides). In collaboration with a parallel effort by the Herbicide Resistant Plants

committee of the Weed Science Society of America, seeds of the sequenced weed accessions

will be deposited in the United States Department of Agriculture Germplasm Resources

Information Network [22] for broad access by the scientific community. The IWGC ensures that

sequenced accessions are collected and documented to comply with the Nagoya Protocol on

access to genetic resources and the fair and equitable sharing of benefits arising from their

utilization under the Convention on Biological Diversity and related Access and Benefit Sharing

Legislation [23]. As additional accessions of weed species are sequenced (e.g., pangenomes are

obtained) the IWGC will facilitate germplasm sharing protocols to support collaboration.

Further, to simplify the investigation of herbicide resistance, the IWGC will link WeedPedia

with the International Herbicide-Resistant Weed Database [24], an already widely known and

utilized database for weed scientists.

Training and Collaboration in Weed Genomics

Beyond producing genomic tools and resources, a priority of the IWGC is to enable the

utilization of these resources across a wide range of stakeholders. A holistic approach to training

is required for weed science generally [25], and we would argue even more so for weed

genomics. To accomplish our training goals, the IWGC is developing and delivering programs

aimed at the full range of IWGC stakeholders and covering a breadth of relevant topics. We have

taken care to ensure our approaches are diverse as to provide training to researchers with all

levels of existing experience and differing reasons for engaging with these tools. Throughout, the

focus is on ensuring that our training and outreach result in impacts that benefit a wide range of

stakeholders.

Although recently developed tools are incredibly enabling and have great potential to replace

antiquated methodology [26] and to solve pressing weed science problems [14], specialized

computational skills are required to fully explore and unlock meaning from these highly complex

datasets. Collaboration with, or training of, computational biologists equipped with these skills

and resources developed by the IWGC will enable weed scientists to expand research programs

and better understand the genetic underpinnings of weed evolution and herbicide resistance. To

fill existing skill gaps, the IWGC is developing summer bootcamps and online modules directed

specifically at weed scientists that will provide training on computational skills (Figure 1).

Because successful utilization of the IWGC resources requires more than general computational

skills, we have also created three additional targeted workshops that teach practical skills related

to genomics databases, molecular biology, and population genomics (available at [27]).

Engagement opportunities during undergraduate degrees has been shown to improve

academic outcomes [28, 29]. Therefore, the IWGC sponsors opportunities for undergraduates to

undertake 10-week Research Experiences for Undergraduates (REU). These REU include an

introduction to bioinformatics, a plant genomics research project that results in a presentation,

and access to career building opportunities in diverse workplace environments. To increase

equitable access to conferences and professional communities, we supported early career

researchers to attend the first two IWGC conferences in the USA as well as workshops and

bootcamps in Europe and South America. These hybrid or in-person travel grants are

intentionally designed to remove barriers and increase participation of individuals from

backgrounds and experiences currently underrepresented within weed/plant science or genomics

[30]. Recipients of these travel awards gave presentations and gained the measurable benefits

that come from either virtual or in-person participation in conferences [31]. Moving forward,

weed scientists must amass skills associated with genomic analyses and collaborate with other

area experts to fully leverage resources developed by the IWGC.

Evolution of Weediness: Potential Research Utilizing New Weed Genomics

Tools

Weeds can evolve from non-weed progenitors through wild colonization, crop de-

domestication, or crop-wild hybridization [32]. Because the time span in which weeds have

evolved is necessarily limited by the origins of agriculture, these non-weed relatives often still

exist and can be leveraged through population genomic and comparative genomic approaches to

identify the adaptive changes that have driven the evolution of weediness. The ability to rapidly

adapt, persist, and spread in agroecosystems are defining features of weedy plants, leading many

to advocate agricultural weeds as ideal candidates for studying rapid plant adaptation [33-36].

The insights gained from applying plant ecological approaches to the study of rapid weed

adaptation will move us towards the ultimate goals of mitigating such adaptation and increasing

the efficacy of crop breeding and biotechnology [14].

Biology and Ecological Genomics of Weeds

The impressive community effort to create and maintain resources for *Arabidopsis thaliana*

ecological genomics provides a motivating example for the emerging study of weed genomics

[37-40]. *Arabidopsis thaliana* was the first flowering plant species to have its genome fully

sequenced [41] and rapidly became a model organism for plant molecular biology. As weedy

genomes become available, collection, maintenance, and resequencing of globally distributed

accessions of these species will help to replicate the success found in ecological studies of *A.*

thaliana [42-48]. Evaluation of these accessions for traits of interest to produce large phenomics

data sets (as in [49-53]) enables genome-wide association studies and population genomics

analyses aimed at dissecting the genetic basis of variation in such traits [54]. Increasingly, these

resources (e.g the 1001 genomes project [42]) have enabled *A. thaliana* to be utilized as a model

species to explore the eco-evolutionary basis of plant adaptation in a more realistic ecological

context. Weedy species should supplement lessons in eco-evolutionary genomics learned from

these experiments in *A. thaliana*.

Untargeted genomic approaches for understanding the evolutionary trajectories of

populations and the genetic basis of traits as described above rely on the collection of genotypic

information from across the genome of many individuals. While whole-genome resequencing

accomplishes this requirement and requires no custom methodology, this approach provides

more information than is necessary and is prohibitively expensive in species with large genomes.

Development and optimization of genotype-by-sequencing methods for capturing reduced

representations of newly sequence genomes like those described by [55-57] will reduce the cost

and computational requirements of genetic mapping and population genetic experiments.

Additionally, the species sequenced by the IWGC do not currently have protocols for stable

transformation, a key development in the popularity of *A. thaliana* as a model organism and a

requirement for many functional genomic approaches. Functional validation of genes/variants

believed to be responsible for traits of interest in weeds has thus far relied on transiently

manipulating endogenous gene expression [58, 59] or ectopic expression of a transgene in a

model system [60-62]. While these methods have been successful, few weed species have well-

studied viral vectors to adapt for use in virus induced gene silencing and spray induced gene

silencing is relatively ineffective without the use of nanocarriers [63], which require specialized

equipment and expertise. Furthermore, traits with complex genetic architecture divergent

between the researched and model species may not be amenable to functional genomic

approaches using transgenesis techniques in model systems. Developing protocols for reduced

representation sequencing, stable transformation, and gene editing/silencing in weeds will allow

for more thorough characterization of candidate genetic variants underlying traits of interest.

Beyond rapid adaptation, some weedy species offer an opportunity to better understand co-

evolution, like that between plants and pollinators and how their interaction leads to the spread of

weedy alleles (Additional File 2). A suite of plant-insect traits has co-evolved to maximize the

attraction of the insect pollinator community and the efficiency of pollen deposition between

flowers ensuring fruit and seed production in many weeds [64, 65]. Genetic mapping

experiments have identified genes and genetic variants responsible for many floral traits

affecting pollinator interaction including petal color [66-69], flower symmetry and size [70-72],

and production of volatile organic compounds [73-75] and nectar [76-78]. While these studies

reveal candidate genes for selection under co-evolution, herbicide resistance alleles may also

have pleiotropic effects on the ecology of weeds [79], altering plant-pollinator interactions [80].

Discovery of genes and genetic variants involved in weed-pollinator interaction and their

molecular and environmental control may create opportunities for better management of weeds

with insect-mediated pollination. For example, if management can disrupt pollinator

attraction/interaction with these weeds, the efficiency of reproduction may be reduced.

A more complete understanding of weed ecological genomics will undoubtedly elucidate

many unresolved questions regarding the genetic basis of various aspects of weediness. For

instance, when comparing populations of a species from agricultural and non-agricultural

environments, is there evidence for contemporary evolution of weedy traits selected by

agricultural management or were 'natural' populations pre-adapted to agroecosystems? Where

there is differentiation between weedy and natural populations, which traits are under selection

and what is the genetic basis of variation in those traits? When comparing between weedy

populations, is there evidence for parallel versus non-parallel evolution of weediness at the

phenotypic and genotypic levels? Such studies may uncover fundamental truths about weediness.

For example, is there a common phenotypic and/or genotypic basis for aspects of weediness

amongst diverse weed species? As genomic tools developed by the IWGC enable researchers to

address these questions, knowledge gained will help predict the potential development of newly

important weed species in new environments and cropping systems.

Population Genomics

Weed species are certainly fierce competitors, able to outcompete crops and endemic

species in their native environment, but they are also remarkable colonizers of perturbed habitats.

Weeds achieve this through high fecundity, often producing tens of thousands of seeds per

individual plant [81-83]. These large numbers in terms of demographic population size often

combine with outcrossing reproduction to generate high levels of diversity with local effective

population sizes in the hundreds of thousands [84, 85]. This has two important consequences:

weed populations retain standing genetic variation and generate many new mutations, supporting

weed success in the face of harsh control. The generation of genomic tools to monitor weed

populations at the molecular level is a game-changer to understanding weed dynamics and

precisely testing the effect of management on the genetic make-up of populations. We are now

able to assess the effect of control on genetic diversity and monitor in real time whether these

measures are effective at eroding weed populations.

Population genomic data, without any environmental or phenotypic information, can be used

to scan the genomes of weed and non-weed relatives to identify selective sweeps, pointing at loci

supporting weed adaptation on micro- or macro-evolutionary scales. Two recent within-species

examples include weedy rice, where population differentiation between weedy and domesticated

populations was used to identify the genetic basis of weedy de-domestication [86], and common

waterhemp, where consistent allelic differences among natural and agricultural collections

resolved a complex set of agriculturally adaptive alleles [87, 88]. A recent comparative

population genomic study of weedy barnyardgrass and crop millet species has demonstrated how

inter-specific investigations can resolve the signatures of crop and weed evolution [89] (also see

[90] for a non-weed climate adaptation example). Multiple sequence alignments across numerous

species provide complementary insight into adaptive convergence over deeper timescales, even

with just one genomic sample per species (e.g., [91, 92]). Thus, the new IWGC weed genomes

combined with genomes available for closely related crops (outlined by [14, 93]) and an effort to

identify other non-weed wild relatives will be invaluable in characterizing the genetic

architecture of weed adaptation and evolution across diverse species.

Weeds experience high levels of genetic selection, both artificial in response to

agricultural practices and particularly herbicides, and natural in response to the environmental

conditions they encounter [94, 95]. Using genomic analysis to identify loci that are the targets of

selection and support adaptation would point at vulnerabilities that could be leveraged against

weeds to develop new and more sustainable management strategies [96]. This is a key

motivation to develop genotype-by-environment association (GEA) and selective sweep scan

approaches, which allow researchers to resolve the molecular basis of multi-dimensional

adaptation [97, 98]. GEA approaches, in particular, have been widely used on landscape-wide

resequencing collections to determine the genetic basis of climate adaptation (e.g., [40, 99, 100]),

but have yet to be fully exploited to diagnose the genetic basis of the various aspects of

weediness [101]. Armed with data on environmental dimensions of agricultural settings, such as

focal crop, soil quality, herbicide use, and climate, GEA approaches can help disentangle how

discrete farming practices have influenced the evolution of weediness and resolve broader

patterns of local adaptation across a weed's range. Although non-weedy relatives are not

technically required for GEA analyses, inclusion of environmental and genomic data from weed

progenitors can further distinguish genetic variants underpinning weed origins from those

involved in local adaptation.

New weeds emerge frequently [102], either through hybridization between species as

documented for sea beet (*Beta vulgaris* ssp. *maritima)* hybridizing with crop beet to produce

progeny that are well adapted to agricultural conditions [103-105], or through the invasion of

alien species that find a new range to colonize. Biosecurity measures are often in place to stop

the introduction of new weeds; however, the vast scale of global agricultural commodity trade

precludes the possibility of total control. Population genomic analysis is now able to measure

gene flow between populations [87, 106-108] and identify populations of origin for invasive

species including weeds [109-111]. In particular, the invasion route of the pest fruitfly

Drosophila suzukii from Eastern Asia to North America and Europe through Hawaii was

deciphered using Approximate Bayesian Computation on high-throughput sequencing data from

a global sample of multiple populations [112]. Genomics can also be leveraged to predict

invasion rather than explain it. The resequencing of a global sample of common ragweed

(*Ambrosia artemisiifolia* L.) elucidated a complex invasion route whereby Europe was invaded

by multiple introductions of American ragweed that hybridized in Europe prior to a subsequent

introduction to Australia [113, 114]. In this context, the use of genomically-informed species

distribution models helps assess the risk associated with different source populations, which in

the case of common ragweed, suggests that a source population from Florida would allow

ragweed to invade most of northern Australia [115]. The IWGC, by federating research efforts

globally, provides a platform that can support the transformation of biosecurity from perspective

analysis towards predictive risk assessment.

Genetic Basis of Herbicide Resistance

Herbicide resistance is among the numerous weedy traits that can evolve in plant populations

exposed to agricultural selection pressures. Over-reliance on herbicides to control weeds, along

with low diversity and lack of redundancy in weed management strategies, has resulted in

globally widespread herbicide resistance [116]. To date, 268 herbicide-resistant weed species

have been reported worldwide, and at least one resistance case exists for 21 of the 31 existing

herbicide sites of action [24] – significantly limiting chemical weed control options available to

agriculturalists. This limitation of control options is exacerbated by the recent lack of discovery

of herbicides with new sites of action [117].

Herbicide resistance may result from several different physiological mechanisms. Such

mechanisms have been classified into two main groups, target-site resistance (TSR) [4, 118] and

non-target-site resistance (NTSR) [4, 119]. The first group encompasses changes that reduce

binding affinity between a herbicide and its target [120]. These changes may provide resistance

to multiple herbicides that have a common biochemical target [121] and can be effectively

managed through mixture and/or rotation of herbicides targeting different sites of action [122].

The second group (NTSR), includes alterations in herbicide absorption, translocation,

sequestration, and/or metabolism that may lead to unpredictable pleotropic cross-resistance

profiles where structurally and functionally diverse herbicides are rendered ineffective by one or

more genetic variant(s) [60]. This mechanism of resistance threatens not only the efficacy of

existing herbicidal chemistries, but also ones yet to be discovered. While TSR is well understood

because of the ease of identification and molecular characterization of target site variants, NTSR

mechanisms are significantly more challenging to research because they are often polygenic, and

the resistance causing element(s) are not well understood [123].

Improving the current understanding of metabolic NTSR mechanisms is not an easy task,

since genes of diverse biochemical functions are involved, many of which exist as extensive

gene families [121, 124]. Expression changes of NTSR genes have been implicated in several

resistance cases where the protein products of the genes are functionally equivalent across

sensitive and resistant plants, but their relative abundance leads to resistance. Thus, regulatory

elements of NTSR genes have been scrutinized to understand their role in NTSR mechanisms

[125]. Similarly, epigenetic modifications have been hypothesized to play a role in NTSR, with

much remaining to be explored [126-128]. Untargeted approaches such as genome-wide

association, selective sweep scans, linkage mapping, RNA-sequencing, and metabolomic

profiling have proven helpful to complement more specific biochemical- and chemo-

characterization studies towards the elucidation of NTSR mechanisms as well as their regulation

and evolution [60, 129-136]. Due to their complexity and importance, the IWGC has begun

addressing this subject by manually curating the annotation of NTSR genes and developing a

standard nomenclature for the gene families often involved in NTSR. This standardization will

allow researchers to quickly identify true orthologous genes between weedy species, which is a

hurdle for current research of these complex and often vast gene families.

High-quality weed genome assemblies and gene model annotations have helped and will be

crucial for investigating the landscape of NTSR genes in weeds. They can also be used to predict

the protein structure for herbicide target site and metabolism genes to predict the efficacy and

selectivity of new candidate herbicides *in silico* to increase herbicide discovery throughput.

Knowledge of the genetic basis of NTSR will aid the rational design of herbicides by 1)

screening new compounds in the presence of newly discovered NTSR proteins during early

research phases; 2) identifying conserved chemical structures that interact with these proteins;

and 3) optimizing herbicide molecular design to lower potential for resistance evolution and

increase potency/spectrum of control.

Moving forward, genomic resources will be increasingly needed and used not only for the

design of conventional small molecule herbicides, but also for next generation technologies for

sustainable weed management. Proteolysis targeting chimeras (PROTACs) have the potential to

bind desired targets with great selectivity and degrade proteins by utilizing natural protein

ubiquitination and degradation pathways within plants [137]. The combination of nanoparticles

with oligonucleotides has recently shown potential to be used in spray applications towards gene

silencing in weeds, which paves the way for a new, innovative, and sustainable method for weed

management [138, 139]. Additionally, success in the field of pharmaceutical drug discovery in

the development of molecules modulating protein-protein interactions offers another potential

avenue towards the development of herbicides with novel targets [140, 141]. High-quality

genomic references allow for the design of new weed management technologies like the ones

listed here that are specific to – and effective across – weed species but have a null effect on non-

target organisms. The tools being developed by the IWGC will have a crucial role in enabling the

development of next generation weed management strategies that will reduce our reliance on the

few chemical control options currently available to agriculturalists.

Comparative Genomics and Genome Biology

The genomes of weed species are as diverse as weed species themselves. Many weed species

belong to unique plant families with no phylogenetically close model or crop species relatives for

comparison. On all measurable metrics, weed genomes run the gamut. Some have smaller

genomes like *Cyperus* spp. (~0.26 Gb) while others are larger, such as *Avena fatua* (~11.1 Gb)

(Table 1). Some have high heterozygosity in terms of single nucleotide polymorphisms,

repetitive DNA, and structural variants, such as the *Amaranthus* spp., while others are primarily

self-pollinated and quite homozygous, such as *Poa annua* [142, 143]. Some are diploid such as

Conyza canadensis and *Echinochloa haploclada* while others are polyploid such as *C.*

sumetrensis, *E. crus-galli,* and *E. colona* [89]. The availability of genomic resources in these

diverse, unexplored branches of the tree of life allows us to identify consistencies and anomalies

in the field of genome biology.

The weed genomes published so far have focused mainly on weeds of agronomic crops, and

studies have revolved around their ability to resist key herbicides. For example, genomic

resources were vital in the elucidation of herbicide resistance cases involving target site gene

copy number variants (CNVs). Gene CNVs of 5-enolpyruvylshikimate-3-phosphate synthase

(*EPSPS*) have been found to confer resistance to the herbicide glyphosate in diverse weed

species. To date, nine species have independently evolved *EPSPS* CNVs, and species achieve

increased *EPSPS* copy number via different mechanisms [144]. For instance, the *EPSPS* CNV in

Bassia scoparia is caused by tandem duplication, which is accredited to transposable element

insertions flanking *EPSPS* and subsequent unequal crossing over events [145, 146]. In *Eleusine*

indica, a *EPSPS* CNV was caused by translocation of the *EPSPS* locus into the subtelomere

followed by telomeric sequence exchange [147]. One of the most fascinating genome biology

discoveries in weed science has been that of extra-chromosomal circular DNAs (eccDNAs) that

harbor the *EPSPS* gene in the weed species *Amaranthus palmeri* [148, 149]. In this case, the

eccDNAs autonomously replicate separately from the nuclear genome and do not reintegrate into

chromosomes, which has implications for inheritance, fitness, and genome structure [150]. These

discoveries would not have been possible without reference assemblies of weed genomes, next-

generation sequencing, and collaboration with experts in plant genomics and bioinformatics.

Another question that is often explored with weedy genomes is the nature and composition of

gene families that are associated with NTSR. Gene families under consideration often include

cytochrome P450s (CYPs), glutathione-*S*-transferases (GSTs), ABC transporters, etc. Some

questions commonly considered with new weed genomes include: how many genes are in each

of these gene families, where are they located, and which weed accessions and species have an

over-abundance of them that might explain their ability to evolve resistance so rapidly [19, 89,

151, 152]? Weed genome resources are necessary to answer questions about gene family

expansion or contraction during the evolution of weediness, including the role of polyploidy in

NTSR gene family expansion as explored by [153].

Translational Research and Communication with Weed Management Stakeholders

Whereas genomics of model plants is typically aimed at addressing fundamental questions in

plant biology, and genomics of crop species has the obvious goal of crop improvement, goals of

genomics of weedy plants also include the development of more effective and sustainable

strategies for their management. Weed genomics assists with these objectives by providing novel

molecular ecological and evolutionary insights from the context of intensive anthropogenic

management (which is lacking in model plants), and offers knowledge and resources for trait

discovery for crop improvement, especially given that many wild crop relatives are also

important agronomic weeds (e.g. [154]). For instance, crop-wild relatives are valuable for

improving crop breeding for marginal environments [155]. Thus, weed genomics presents unique

opportunities and challenges relative to plant genomics more broadly. It should also be noted that

although weed science at its core is a very applied discipline, it draws broadly from many

scientific disciplines such as, plant physiology, chemistry, ecology, and evolutionary biology, to

name a few. The successful integration of weed-management strategies, therefore, requires

extensive collaboration among individuals collectively possessing the necessary expertise [156].

Consequently, a major objective of the IWGC is to ensure that basic findings arising from weed

genomics are translated to advances in weed management and crop breeding by collaborating

broadly with breeders, applied weed scientists, outreach specialists, and practitioners.

To accomplish this objective, the IWGC must facilitate communication of weed genomics

findings to relevant stakeholders (Figure 1). With the growing complexity of herbicide resistance

management, practitioners are beginning to recognize the importance of understanding resistance

mechanisms to inform appropriate management tactics [14]. Although weed science practitioners

do not need to understand the technical details of weed genomics, their appreciation of the power

of weed genomics - together with their unique insights from field observations - will yield novel

opportunities for applications of weed genomics to weed management. In particular, combining

field management history with information on weed resistance mechanisms is expected to

provide novel insights into evolutionary trajectories [e.g., 6, 157], which can be utilized for

disrupting evolutionary adaptation. It can be difficult to obtain field history information from

practitioners, but developing an understanding among them of the importance of such

information can be invaluable. To address these aspects, the IWGC can provide funding, or at

least coordinate teams, to build extension/education programs focused on weed genomics.

Factsheets and easy-to-understand infographics can be developed and communicated to various

stakeholders through traditional and electronic media.

Conclusions

Weeds are unique and fascinating plants, having significant impacts on agriculture and

ecosystems; and yet, aspects of their biology, ecology, and genetics remain poorly understood.

Weeds represent a unique area within plant biology, given their repeated rapid adaptation to

sudden and severe shifts in the selective landscape of anthropogenic management practices. The

production of a public genomics database with reference genomes for over 50 weed species

represents a substantial step forward towards research goals that improve our understanding of

the biology and evolution of weeds. Future work is needed to improve annotations, particularly

for complex gene families involved in herbicide detoxification, structural variants, and mobile

genetic elements, given the evidence to date of the generation of adaptive genetic variation in

weeds through structural variation. As reference genome assemblies become available; standard,

affordable methods for gathering genotype information will allow for the identification of

genetic variants underlying traits of interest. Further, development of methods for functional

gene validation and hypothesis testing is needed in weeds to validate the effect of genetic

variants detected through such experiments, including systems for transformation, gene editing,

and transient gene silencing and expression. Future research should focus on utilizing weed

genomes to investigate questions about the evolutionary biology, ecology, and genetics of weedy

traits and weed population dynamics. The IWGC plans to continue the public-private partnership

model to continue to host the WeedPedia database, integrate new datasets such as genome

resequencing and transcriptomes, conduct trainings, and serve as a research coordination

network to ensure that advances in weed science from around the world are shared across the

research community (Figure 1). Bridging basic plant genomics with translational applications in

weeds is needed to deliver on the potential of weed genomics to improve weed management and

crop breeding.

Availability of data and materials

The datasets supporting the conclusions of this article is included within the article and its

additional files. All genome assemblies and related sequencing data produced by the IWGC will

be available through NCBI as part of publications reporting the first genome-wide analysis for

each species.

Competing interests

527 The authors declare that they have no competing interests.

Acknowledgements

530 The International Weed Genomics Consortium is supported by BASF SE, Bayer AG, Syngenta

531 Ltd, Corteva Agriscience, CropLife International, the Foundation for Food and Agriculture

532 Research (Award DSnew-024), and a conference grant from USDA-NIFA (Award number 2021-

533 67013-33570).

References

1.   Gianessi LP, Nathan PR. The Value of Herbicides in U.S. Crop Production. Weed

Technol. 2007;21(2):559-66.

2.   Pimentel D, Lach L, Zuniga R, Morrison D. Environmental and Economic Costs of

Nonindigenous Species in the United States. BioScience. 2000;50(1):53-65.

3.   Barrett SH. Crop mimicry in weeds. Econ Bot. 1983;37(3):255-82.

4.   Powles SB, Yu Q. Evolution in action: plants resistant to herbicides. Annu Rev Plant

Biol. 2010;61):317-47.

5.   Thurber CS, Reagon M, Gross BL, Olsen KM, Jia Y, Caicedo AL. Molecular evolution

of shattering loci in U.S. weedy rice. Mol Ecol. 2010;19(16):3271-84.

6.   Comont D, Lowe C, Hull R, Crook L, Hicks HL, Onkokesung N, et al. Evolution of

generalist resistance to herbicide mixtures reveals a trade-off in resistance management.

Nat Commun. 2020;11(1):3086.

7.   Ashworth MB, Walsh MJ, Flower KC, Vila-Aiub MM, Powles SB. Directional selection

549   for flowering time leads to adaptive evolution in *Raphanus raphanistrum* (Wild radish).

550   Evol Appl. 2016;9(4):619-29.

551   8.   Chan EK, Rowe HC, Kliebenstein DJ. Understanding the evolution of defense

552   metabolites in *Arabidopsis thaliana* using genome-wide association mapping. Genetics.

553   2010;185(3):991-1007.

554   9.   Frayling TM, Timpson NJ, Weedon MN, Zeggini E, Freathy RM, Lindgren CM, et al. A

555   common variant in the *FTO* gene is associated with body mass index and predisposes to

556   childhood and adult obesity. Science. 2007;316(5826):889-94.

557   10.   Harkess A, Zhou J, Xu C, Bowers JE, Van der Hulst R, Ayyampalayam S, et al. The

558   asparagus genome sheds light on the origin and evolution of a young Y chromosome. Nat

559   Commun. 2017;8(1):1279.

560   11.   Periyannan S, Moore J, Ayliffe M, Bansal U, Wang X, Huang L, et al. The gene *Sr33*, an

561   ortholog of barley *Mla* genes, encodes resistance to wheat stem rust race Ug99. Science.

562   2013;341(6147):786-8.

563   12.   Ågren J, Oakley CG, McKay JK, Lovell JT, Schemske DW. Genetic mapping of

564   adaptation reveals fitness tradeoffs in *Arabidopsis thaliana*. Proc Natl Acad Sci U S A.

565   2013;110(52):21077-82.

566   13.   Schartl M, Walter RB, Shen Y, Garcia T, Catchen J, Amores A, et al. The genome of the

567   platyfish, *Xiphophorus maculatus*, provides insights into evolutionary adaptation and

568   several complex traits. Nat Genet. 2013;45(5):567-72.

569   14.   Ravet K, Patterson EL, Krähmer H, Hamouzová K, Fan L, Jasieniuk M, et al. The power

570   and potential of genomics in weed biology and management. Pest Manag Sci.

571   2018;74(10):2216-25.

572   15.   Hufford MB, Seetharam AS, Woodhouse MR, Chougule KM, Ou S, Liu J, et al. De novo

573   assembly, annotation, and comparative analysis of 26 diverse maize genomes. Science.

574   2021;373(6555):655-62.

575   16.   Liao W-W, Asri M, Ebler J, Doerr D, Haukness M, Hickey G, et al. A draft human

576   pangenome reference. Nature. 2023;617(7960):312-24.

577   17.   Huang Y, Wu D, Huang Z, Li X, Merotto A, Bai L, et al. Weed genomics: yielding

578   insights into the genetics of weedy traits for crop improvement. aBIOTECH. 2023.

579   18.   Bennetzen JL, Schmutz J, Wang H, Percifield R, Hawkins J, Pontaroli AC, et al.

580   Reference genome sequence of the model plant *Setaria*. Nat Biotechnol. 2012;30(6):555-

581   61.

582   19.   Kreiner JM, Giacomini DA, Bemm F, Waithaka B, Regalado J, Lanz C, et al. Multiple

583   modes of convergent adaptation in the spread of glyphosate-resistant *Amaranthus*

584   *tuberculatus*. Proc Natl Acad Sci U S A. 2019;116(42):21076-84.

585   20.   Yu Y, Hu H, Doust AN, Kellogg EA. Divergent gene expression networks underlie

586   morphological diversity of abscission zones in grasses. New Phytol. 2020;225(4):1799-

587   815.

588   21.   International Weed Genomics Consortium Website. weedgenomics.org. Accessed 20

589   June 2023.

590   22.   USDA-ARS Germplasm Resources Information Network (GRIN). www.ars-grin.gov.

591   Accessed 20 June 2023.

23.   Buck M, Hamilton C. The Nagoya Protocol on Access to Genetic Resources and the Fair

and Equitable Sharing of Benefits Arising from their Utilization to the Convention on

Biological Diversity. RECIEL. 2011;20(1):47-61.

24.   The International Herbicide-Resistant Weed Database. www.weedscience.org. Accessed

20 June 2023.

25.   Chauhan BS, Matloob A, Mahajan G, Aslam F, Florentine SK, Jha P. Emerging

challenges and opportunities for education and research in weed science. Front Plant Sci.

2017;(8):1537.

26.   Shah S, Lonhienne T, Murray CE, Chen Y, Dougan KE, Low YS, et al. Genome-Guided

Analysis of Seven Weed Species Reveals Conserved Sequence and Structural Features of

Key Gene Targets for Herbicide Development. Front Plant Sci. 2022;13):909073.

27.   International Weed Genomics Consortium Training Resources.

https://www.weedgenomics.org/training-resources/. Accessed 20 June 2023.

28.   Blackford S. Harnessing the power of communities: career networking strategies for

bioscience PhD students and postdoctoral researchers. FEMS Microbiol Lett. 2018;365(8.

29.   Pender M, Marcotte DE, Sto Domingo MR, Maton KI. The STEM Pipeline: The Role of

Summer Research Experience in Minority Students' Ph.D. Aspirations. Educ Policy Anal

Arch. 2010;18(30):1-36.

30.   Burke A, Okrent A, Hale K. The State of U.S. Science and Engineering 2022. National

Science Foundation. https://ncses.nsf.gov/pubs/nsb20221. 2022.

31.   Wu J-Y, Liao C-H, Cheng T, Nian M-W. Using Data Analytics to Investigate Attendees'

Behaviors and Psychological States in a Virtual Academic Conference. Educ Technol

Soc. 2021;24(1):75-91.

32.   De Wet JMJ, Harlan JR. Weeds and domesticates: Evolution in the man-made habitat.

616   Econ Bot. 1975;29(2):99-108.

617   33.   Mahaut L, Cheptou PO, Fried G, Munoz F, Storkey J, Vasseur F, et al. Weeds: Against

618   the Rules? Trends Plant Sci. 2020;25(11):1107-16.

619   34.   Neve P, Vila-Aiub M, Roux F. Evolutionary-thinking in agricultural weed management.

620   New Phytol. 2009;184(4):783-93.

621   35.   Sharma G, Barney JN, Westwood JH, Haak DC. Into the weeds: new insights in plant

622   stress. Trends Plant Sci. 2021;26(10):1050-60.

623   36.   Vigueira CC, Olsen KM, Caicedo AL. The red queen in the corn: agricultural weeds as

624   models of rapid adaptive evolution. Heredity (Edinb). 2013;110(4):303-11.

625   37.   Donohue K, Dorn L, Griffith C, Kim E, Aguilera A, Polisetty CR, et al. Niche

626   construction through germination cueing: life-history responses to timing of germination

627   in *Arabidopsis thaliana*. Evolution. 2005;59(4):771-85.

628   38.   Exposito-Alonso M. Seasonal timing adaptation across the geographic range of

629   *Arabidopsis thaliana*. Proc Natl Acad Sci U S A. 2020;117(18):9665-67.

630   39.   Fournier-Level A, Korte A, Cooper MD, Nordborg M, Schmitt J, Wilczek AM. A map of

631   local adaptation in *Arabidopsis thaliana*. Science. 2011;334(6052):86-9.

632   40.   Hancock AM, Brachi B, Faure N, Horton MW, Jarymowycz LB, Sperone FG, et al.

633   Adaptation to climate across the *Arabidopsis thaliana* genome. Science.

634   2011;334(6052):83-6.

635   41.   TAG Initiative. Analysis of the genome sequence of the flowering plant *Arabidopsis

636   thaliana*. Nature. 2000;408(6814):796-815.

42. Alonso-Blanco C, Andrade J, Becker C, Bemm F, Bergelson J, Borgwardt KM, et al.

1,135 Genomes Reveal the Global Pattern of Polymorphism in *Arabidopsis thaliana*.

Cell. 2016;166(2):481-91.

43. Durvasula A, Fulgione A, Gutaker RM, Alacakaptan SI, Flood PJ, Neto C, et al. African

genomes illuminate the early history and transition to selfing in *Arabidopsis thaliana*.

Proc Natl Acad Sci U S A. 2017;114(20):5213-18.

44. Frachon L, Mayjonade B, Bartoli C, Hautekèete N-C, Roux F. Adaptation to Plant

Communities across the Genome of *Arabidopsis thaliana*. Mol Biol Evol.

2019;36(7):1442-56.

45. Fulgione A, Koornneef M, Roux F, Hermisson J, Hancock AM. Madeiran *Arabidopsis*

thaliana reveals ancient long-range colonization and clarifies demography in Eurasia.

Mol Biol Evol. 2018;35(3):564-74.

46. Fulgione A, Neto C, Elfarargi AF, Tergemina E, Ansari S, Göktay M, et al. Parallel

reduction in flowering time from de novo mutations enable evolutionary rescue in

colonizing lineages. Nat Commun. 2022;13(1):1461.

47. Kasulin L, Rowan BA, León RJC, Schuenemann VJ, Weigel D, Botto JF. A single

haplotype hyposensitive to light and requiring strong vernalization dominates

Arabidopsis thaliana populations in Patagonia, Argentina. Mol Ecol. 2017;26(13):3389-

404.

48. Picó FX, Méndez-Vigo B, Martínez-Zapater JM, Alonso-Blanco C. Natural genetic

variation of *Arabidopsis thaliana* is geographically structured in the Iberian peninsula.

Genetics. 2008;180(2):1009-21.

49.   Atwell S, Huang YS, Vilhjálmsson BJ, Willems G, Horton M, Li Y, et al. Genome-wide

660   association study of 107 phenotypes in *Arabidopsis thaliana* inbred lines. Nature.

661   2010;465(7298):627-31.

662   50.   Flood PJ, Kruijer W, Schnabel SK, van der Schoor R, Jalink H, Snel JFH, et al.

663   Phenomics for photosynthesis, growth and reflectance in *Arabidopsis thaliana* reveals

664   circadian and long-term fluctuations in heritability. Plant Methods. 2016;12(1):14.

665   51.   Marchadier E, Hanemian M, Tisné S, Bach L, Bazakos C, Gilbault E, et al. The complex

666   genetic architecture of shoot growth natural variation in *Arabidopsis thaliana*. PLoS

667   Genet. 2019;15(4):e1007954.

668   52.   Tisné S, Serrand Y, Bach L, Gilbault E, Ben Ameur R, Balasse H, et al. Phenoscope: an

669   automated large-scale phenotyping platform offering high spatial homogeneity. Plant J.

670   2013;74(3):534-44.

671   53.   Tschiersch H, Junker A, Meyer RC, Altmann T. Establishment of integrated protocols for

672   automated high throughput kinetic chlorophyll fluorescence analyses. Plant Methods.

673   2017;13):54.

674   54.   Chen X, MacGregor DR, Stefanato FL, Zhang N, Barros-Galvão T, Penfield S. A VEL3

675   histone deacetylase complex establishes a maternal epigenetic state controlling progeny

676   seed dormancy. Nat Commun. 2023;14(1):2220.

677   55.   Choi M, Scholl UI, Ji W, Liu T, Tikhonova IR, Zumbo P, et al. Genetic diagnosis by

678   whole exome capture and massively parallel DNA sequencing. Proc Natl Acad Sci U S

679   A. 2009;106(45):19096-101.

680   56.   Davey JW, Blaxter ML. RADSeq: next-generation population genetics. Brief Funct

681   Genomics. 2010;9(5-6):416-23.

682   57.   Elshire RJ, Glaubitz JC, Sun Q, Poland JA, Kawamoto K, Buckler ES, et al. A robust,

683   simple genotyping-by-sequencing (GBS) approach for high diversity species. PLoS One.

684   2011;6(5):e19379.

685   58.   MacGregor DR. What Makes a Weed a Weed? How Virus-mediated Reverse Genetics

686   Can Help to Explore the Genetics of Weediness. Outlooks Pest Manag. 2020;31(5):224-

687   29.

688   59.   Mellado-Sánchez M, McDiarmid F, Cardoso V, Kanyuka K, MacGregor DR. Virus-

689   Mediated Transient Expression Techniques Enable Gene Function Studies in Blackgrass.

690   Plant Physiol. 2020;183(2):455-59.

691   60.   Dimaano NG, Yamaguchi T, Fukunishi K, Tominaga T, Iwakami S. Functional

692   characterization of Cytochrome P450 *CYP81A* subfamily to disclose the pattern of cross-

693   resistance in *Echinochloa phyllopogon*. Plant Mol Biol. 2020;102(4-5):403-16.

694   61.   de Figueiredo MRA, Küpper A, Malone JM, Petrovic T, de Figueiredo ABTB,

695   Campagnola G, et al. An in-frame deletion mutation in the degron tail of auxin coreceptor

696   IAA2 confers resistance to the herbicide 2,4-D in *Sisymbrium orientale*. Proc Natl Acad

697   Sci U S A. 2022;119(9):e2105819119.

698   62.   Patzoldt WL, Hager AG, McCormick JS, Tranel PJ. A codon deletion confers resistance

699   to herbicides inhibiting protoporphyrinogen oxidase. Proc Natl Acad Sci U S A.

700   2006;103(33):12329-34.

701   63.   Schwartz SH, Hendrix B, Hoffer P, Sanders RA, Zheng W. Carbon Dots for Efficient

702   Small Interfering RNA Delivery and Gene Silencing in Plants. Plant Physiol.

703   2020;184(2):647-57.

704   64.   Fattorini R, Glover BJ. Molecular Mechanisms of Pollination Biology. Annu Rev Plant

705   Biol. 2020;71):487-515.

706   65.   Rollin O, Benelli G, Benvenuti S, Decourtye A, Wratten SD, Canale A, et al. Weed-

707   insect pollinator networks as bio-indicators of ecological sustainability in agriculture. A

708   review. Agron Sustain Dev. 2016;36(1):8.

709   66.   Irwin RE, Strauss SY. Flower color microevolution in wild radish: evolutionary response

710   to pollinator-mediated selection. Am Nat. 2005;165(2):225-37.

711   67.   Ma B, Wu J, Shi T-L, Yang Y-Y, Wang W-B, Zheng Y, et al. Lilac (*Syringa oblata*)

712   genome provides insights into its evolution and molecular mechanism of petal color

713   change. Commun Biol. 2022;5(1):686.

714   68.   Xing A, Wang X, Nazir MF, Zhang X, Wang X, Yang R, et al. Transcriptomic and

715   metabolomic profiling of flavonoid biosynthesis provides novel insights into petals

716   coloration in Asian cotton (*Gossypium arboreum* L.). BMC Plant Biol. 2022;22(1):416.

717   69.   Zheng Y, Chen Y, Liu Z, Wu H, Jiao F, Xin H, et al. Important Roles of Key Genes and

718   Transcription Factors in Flower Color Differences of *Nicotianaalata*. Genes (Basel).

719   2021;12(12.

720   70.   Krizek BA, Anderson JT. Control of flower size. J Exp Bot. 2013;64(6):1427-37.

721   71.   Powell AE, Lenhard M. Control of organ size in plants. Curr Biol. 2012;22(9):R360-7.

722   72.   Spencer V, Kim M. Re"CYC"ling molecular regulators in the evolution and development

723   of flower symmetry. Semin Cell Dev Biol. 2018;79):16-26.

724   73.   Amrad A, Moser M, Mandel T, de Vries M, Schuurink RC, Freitas L, et al. Gain and

725   Loss of Floral Scent Production through Changes in Structural Genes during Pollinator-

726   Mediated Speciation. Curr Biol. 2016;26(24):3303-12.

74.   Delle-Vedove R, Schatz B, Dufay M. Understanding intraspecific variation of floral scent

in light of evolutionary ecology. Ann Bot. 2017;120(1):1-20.

75.   Pichersky E, Gershenzon J. The formation and function of plant volatiles: perfumes for

pollinator attraction and defense. Curr Opin Plant Biol. 2002;5(3):237-43.

76.   Ballerini ES, Kramer EM, Hodges SA. Comparative transcriptomics of early petal

development across four diverse species of *Aquilegia* reveal few genes consistently

associated with nectar spur development. BMC Genom. 2019;20(1):668.

77.   Corbet SA, Willmer PG, Beament JWL, Unwin DM, Prys-Jones OE. Post-secretory

determinants of sugar concentration in nectar. Plant Cell Environ. 1979;2(4):293-308.

78.   Galliot C, Hoballah ME, Kuhlemeier C, Stuurman J. Genetics of flower size and nectar

volume in Petunia pollination syndromes. Planta. 2006;225(1):203-12.

79.   Vila-Aiub MM, Neve P, Powles SB. Fitness costs associated with evolved herbicide

resistance alleles in plants. New Phytol. 2009;184(4):751-67.

80.   Baucom RS. Evolutionary and ecological insights from herbicide-resistant weeds: what

have we learned about plant adaptation, and what is left to uncover? New Phytol.

2019;223(1):68-82.

81.   Bajwa AA, Latif S, Borger C, Iqbal N, Asaduzzaman M, Wu H, et al. The remarkable

journey of a weed: Biology and management of annual ryegrass (*Lolium rigidum*) in

conservation cropping systems of Australia. Plants (Basel). 2021;10(8):1505.

82.   Bitarafan Z, Andreasen C. Fecundity allocation in some European weed species

competing with crops. Agronomy. 2022;12(5):1196.

83.   Costea M, Weaver SE, Tardif FJ. The biology of Canadian weeds. 130. *Amaranthus*

retroflexus L., *A. powellii, A. powellii* S. *Watson*, and *A. hybridus* L. Can J Plant Sci.

2004;84(2):631-68.

84.   Dixon A, Comont D, Slavov GT, Neve P. Population genomics of selectively neutral

genetic structure and herbicide resistance in UK populations of *Alopecurus myosuroides*.

Pest Manag Sci. 2021;77(3):1520-29.

85.   Kersten S, Chang J, Huber CD, Voichek Y, Lanz C, Hagmaier T, et al. Standing genetic

variation fuels rapid evolution of herbicide resistance in blackgrass. Proc Natl Acad Sci U

S A. 2023;120(16):e2206808120.

86.   Qiu J, Zhou Y, Mao L, Ye C, Wang W, Zhang J, et al. Genomic variation associated with

local adaptation of weedy rice during de-domestication. Nat Commun. 2017;8(1):15323.

87.   Kreiner JM, Caballero A, Wright SI, Stinchcombe JR. Selective ancestral sorting and de

novo evolution in the agricultural invasion of *Amaranthus tuberculatus*. Evolution.

2022;76(1):70-85.

88.   Kreiner JM, Latorre SM, Burbano HA, Stinchcombe JR, Otto SP, Weigel D, et al. Rapid

weed adaptation and range expansion in response to agriculture over the past two

centuries. Science. 2022;378(6624):1079-85.

89.   Wu D, Shen E, Jiang B, Feng Y, Tang W, Lao S, et al. Genomic insights into the

evolution of *Echinochloa* species as weed and orphan crop. Nat Commun.

2022;13(1):689.

90.   Yeaman S, Hodgins KA, Lotterhos KE, Suren H, Nadeau S, Degner JC, et al. Convergent

local adaptation to climate in distantly related conifers. Science. 2016;353(6306):1431-

33.

91. Haudry A, Platts AE, Vello E, Hoen DR, Leclercq M, Williamson RJ, et al. An atlas of

over 90,000 conserved noncoding sequences provides insight into crucifer regulatory

regions. Nat Genet. 2013;45(8):891-98.

92. Sackton TB, Grayson P, Cloutier A, Hu Z, Liu JS, Wheeler NE, et al. Convergent

regulatory evolution and loss of flight in paleognathous birds. Science.

2019;364(6435):74-78.

93. Ye CY, Fan L. Orphan Crops and their Wild Relatives in the Genomic Era. Mol Plant.

2021;14(1):27-39.

94. Clements DR, Jones VL. Ten ways that weed evolution defies human management

efforts amidst a changing climate. Agronomy. 2021;11(2):284.

95. Weinig C. Rapid evolutionary responses to selection in heterogeneous environments

among agricultural and nonagricultural weeds. Int J Plant Sci. 2005;166(4):641-47.

96. Cousens RD, Fournier-Level A. Herbicide resistance costs: what are we actually

measuring and why? Pest Manag Sci. 2018;74(7):1539-46.

97. Lasky JR, Josephs EB, Morris GP. Genotype–environment associations to reveal the

molecular basis of environmental adaptation. Plant Cell. 2023;35(1):125-38.

98. Lotterhos KE. The Effect of Neutral Recombination Variation on Genome Scans for

Selection. G3-Genes Genom Genet. 2019;9(6):1851-67.

99. Lovell JT, MacQueen AH, Mamidi S, Bonnette J, Jenkins J, Napier JD, et al. Genomic

mechanisms of climate adaptation in polyploid bioenergy switchgrass. Nature.

2021;590(7846):438-44.

100.   Todesco M, Owens GL, Bercovich N, Légaré J-S, Soudi S, Burge DO, et al. Massive

793   haplotypes underlie ecotypic differentiation in sunflowers. Nature. 2020;584(7822):602-

794   07.

795   101.   Revolinski SR, Maughan PJ, Coleman CE, Burke IC. Preadapted to adapt: Underpinnings

796   of adaptive plasticity revealed by the downy brome genome. Commun Biol. 2023;6):326.

797   102.   Kuester A, Conner JK, Culley T, Baucom RS. How weeds emerge: a taxonomic and trait-

798   based examination using United States data. New Phytol. 2014;202(3):1055-68.

799   103.   Arnaud JF, Fénart S, Cordellier M, Cuguen J. Populations of weedy crop-wild hybrid

800   beets show contrasting variation in mating system and population genetic structure. Evol

801   Appl. 2010;3(3):305-18.

802   104.   Ellstrand NC, Schierenbeck KA. Hybridization as a stimulus for the evolution of

803   invasiveness in plants? Proc Natl Acad Sci U S A. 2000;97(13):7043-50.

804   105.   Nakabayashi K, Leubner-Metzger G. Seed dormancy and weed emergence: from

805   simulating environmental change to understanding trait plasticity, adaptive evolution, and

806   population fitness. J Exp Bot. 2021;72(12):4181-85.

807   106.   Busi R, Yu Q, Barrett-Lennard R, Powles S. Long distance pollen-mediated flow of

808   herbicide resistance genes in *Lolium rigidum*. Theoretical and Applied Genetics.

809   2008;117(8):1281-90.

810   107.   Délye C, Clément JAJ, Pernin F, Chauvel B, Le Corre V. High gene flow promotes the

811   genetic homogeneity of arable weed populations at the landscape level. Basic and

812   Applied Ecology. 2010;11(6):504-12.

813   108.   Roumet M, Noilhan C, Latreille M, David J, Muller MH. How to escape from crop-to-

814   weed gene flow: phenological variation and isolation-by-time within weedy sunflower

815   populations. New Phytol. 2013;197(2):642-54.

816   109.   Moghadam SH, Alebrahim MT, Mohebodini M, MacGregor DR. Genetic variation of

817   *Amaranthus retroflexus* L. and *Chenopodium album* L. (Amaranthaceae) suggests

818   multiple independent introductions into Iran. Front Plant Sci. 2023;13.

819   110.   Muller M-H, Latreille M, Tollon C. The origin and evolution of a recent agricultural

820   weed: population genetic diversity of weedy populations of sunflower (*Helianthus*

821   *annuus* L.) in Spain and France. Evol Appl. 2011;4(3):499-514.

822   111.   Wesse C, Welk E, Hurka H, Neuffer B. Geographical pattern of genetic diversity in

823   *Capsella bursa-pastoris* (Brassicaceae)—A global perspective. Ecol Evol.

824   2021;11(1):199-213.

825   112.   Fraimout A, Debat V, Fellous S, Hufbauer RA, Foucaud J, Pudlo P, et al. Deciphering the

826   routes of invasion of *Drosophila suzukii* by means of ABC random forest. Mol Biol Evol.

827   2017;34(4):980-96.

828   113.   Battlay P, Wilson J, Bieker VC, Lee C, Prapas D, Petersen B, et al. Large haploblocks

829   underlie rapid adaptation in the invasive weed *Ambrosia artemisiifolia*. Nat Commun.

830   2023;14(1):1717.

831   114.   van Boheemen LA, Hodgins KA. Rapid repeatable phenotypic and genomic adaptation

832   following multiple introductions. Mol Ecol. 2020;29(21):4102-17.

833   115.   Putra A, Hodgins K, Fournier-Level A. Assessing the invasive potential of different

834   source populations of ragweed (*Ambrosia artemisiifolia* L.) through genomically-

835   informed species distribution modelling. 2023.

836   116.   Bourguet D, Delmotte F, Franck P, Guillemaud T, Reboud X, Vacher C, et al.

837   Heterogeneity of selection and the evolution of resistance. Trends Ecol Evol.

838   2013;28(2):110-8.

839   117.   Powles S. Herbicide discovery through innovation and diversity. Adv Weed Sci.

840   2022;40(spe1):e020220074.

841   118.   Murphy BP, Tranel PJ. Target-Site Mutations Conferring Herbicide Resistance. Plants

842   (Basel). 2019;8(10):382.

843   119.   Gaines TA, Duke SO, Morran S, Rigon CAG, Tranel PJ, Küpper A, et al. Mechanisms of

844   evolved herbicide resistance. J Biol Chem. 2020;295(30):10307-30.

845   120.   Lonhienne T, Cheng Y, Garcia MD, Hu SH, Low YS, Schenk G, et al. Structural basis of

846   resistance to herbicides that target acetohydroxyacid synthase. Nat Commun.

847   2022;13(1):3368.

848   121.   Comont D, MacGregor DR, Crook L, Hull R, Nguyen L, Freckleton RP, et al. Dissecting

849   weed adaptation: Fitness and trait correlations in herbicide-resistant *Alopecurus*

850   *myosuroides*. Pest Manag Sci. 2022;78(7):3039-50.

851   122.   Neve P. Simulation modelling to understand the evolution and management of glyphosate

852   resistance in weeds. Pest Manag Sci. 2008;64(4):392-401.

853   123.   Torra J, Alcántara-de la Cruz R. Molecular Mechanisms of Herbicide Resistance in

854   Weeds. Genes (Basel). 2022;13(11.

855   124.   Délye C, Gardin JAC, Boucansaud K, Chauvel B, Petit C. Non-target-site-based

856   resistance should be the centre of attention for herbicide resistance research: *Alopecurus*

857   *myosuroides* as an illustration. Weed Res. 2011;51(5):433-37.

125. Chandra S, Leon RG. Genome-wide evolutionary analysis of putative non-specific

herbicide resistance genes and compilation of core promoters between monocots and

dicots. Genes (Basel). 2022;13(7).

126. Margaritopoulou T, Tani E, Chachalis D, Travlos I. Involvement of Epigenetic

Mechanisms in Herbicide Resistance: The Case of *Conyza canadensis*. Agriculture.

2018;8(1.

127. Pan L, Guo Q, Wang J, Shi L, Yang X, Zhou Y, et al. *CYP81A68* confers metabolic

resistance to ALS and ACCase-inhibiting herbicides and its epigenetic regulation in

Echinochloa crus-galli. J Hazard Mater. 2022;428):128225.

128. Sen MK, Hamouzová K, Košnarová P, Roy A, Soukup J. Herbicide resistance in grass

weeds: Epigenetic regulation matters too. Front Plant Sci. 2022;13):1040958.

129. Han H, Yu Q, Beffa R, González S, Maiwald F, Wang J, et al. Cytochrome P450

CYP81A10v7 in *Lolium rigidum* confers metabolic resistance to herbicides across at least

five modes of action. Plant J. 2021;105(1):79-92.

130. Kubis GC, Marques RZ, Kitamura RS, Barroso AA, Juneau P, Gomes MP. Antioxidant

Enzyme and Cytochrome P450 Activities Are Involved in Horseweed (*Conyza*

Sumatrensis) Resistance to Glyphosate. Stress. 2023;3(1):47-57.

131. Qiao Y, Zhang N, Liu J, Yang H. Interpretation of ametryn biodegradation in rice based

on joint analyses of transcriptome, metabolome and chemo-characterization. J Hazard

Mater. 2023;445):130526.

132. Rouse CE, Roma-Burgos N, Barbosa Martins BA. Physiological assessment of non–

target site restistance in multiple-resistant junglerice (*Echinochloa colona*). Weed Sci.

2019;67(6):622-32.

133.   Abou-Khater L, Maalouf F, Jighly A, Alsamman AM, Rubiales D, Rispail N, et al.

882   Genomic regions associated with herbicide tolerance in a worldwide faba bean (*Vicia*

883   *faba* L.) collection. Sci Rep. 2022;12(1):158.

884   134.   Gupta S, Harkess A, Soble A, Van Etten M, Leebens-Mack J, Baucom RS.

885   Interchromosomal linkage disequilibrium and linked fitness cost loci associated with

886   selection for herbicide resistance. New Phytol. 2023;238(3):1263-77.

887   135.   Kreiner JM, Tranel PJ, Weigel D, Stinchcombe JR, Wright SI. The genetic architecture

888   and population genomic signatures of glyphosate resistance in *Amaranthus tuberculatus*.

889   Mol Ecol. 2021;30(21):5373-89.

890   136.   Parcharidou E, Dücker R, Zöllner P, Ries S, Orru R, Beffa R. Recombinant glutathione

891   transferases from flufenacet-resistant black-grass (*Alopecurus myosuroides* Huds.) form

892   different flufenacet metabolites and differ in their interaction with pre- and post-

893   emergence herbicides. Pest Manag Sci. 2023.

894   137.   Békés M, Langley DR, Crews CM. PROTAC targeted protein degraders: the past is

895   prologue. Nat Rev Drug Discov. 2022;21(3):181-200.

896   138.   Martin K, Gaines T, Aishwarya V. Pairing Nanoparticles with FANA Technology for

897   Spray-on Gene Silencing in Weeds. Weed Science Society of America Annual Meeting;

898   Arlington, VA. 2023.

899   139.   Zabala-Pardo D, Gaines T, Lamego FP, Avila LA. RNAi as a tool for weed management:

900   challenges and opportunities. Adv Weed Sci. 2022;40(spe 1): e020220096.

901   140.   Acuner Ozbabacan SE, Engin HB, Gursoy A, Keskin O. Transient protein-protein

902   interactions. Protein Eng Des Sel. 2011;24(9):635-48.

903   141.   Lu H, Zhou Q, He J, Jiang Z, Peng C, Tong R, et al. Recent advances in the development

904   of protein–protein interactions modulators: mechanisms and clinical trials. Signal

905   Transduct Target Ther. 2020;5(1):213.

906   142.   Benson CW, Sheltra MR, Maughan PJ, Jellen EN, Robbins MD, Bushman BS, et al.

907   Homoeologous evolution of the allotetraploid genome of *Poa annua* L. BMC Genom.

908   2023;24(1):350.

909   143.   Robbins MD, Bushman BS, Huff DR, Benson CW, Warnke SE, Maughan CA, et al.

910   Chromosome-scale genome assembly and annotation of allotetraploid annual bluegrass

911   (*Poa annua* L.). Genome Biol Evol. 2022;15(1):evac180.

912   144.   Patterson EL, Pettinga DJ, Ravet K, Neve P, Gaines TA. Glyphosate Resistance and

913   *EPSPS* Gene Duplication: Convergent Evolution in Multiple Plant Species. J Hered.

914   2018;109(2):117-25.

915   145.   Jugulam M, Niehues K, Godar AS, Koo DH, Danilova T, Friebe B, et al. Tandem

916   amplification of a chromosomal segment harboring 5-enolpyruvylshikimate-3-phosphate

917   synthase locus confers glyphosate resistance in *Kochia scoparia*. Plant Physiol.

918   2014;166(3):1200-7.

919   146.   Patterson EL, Saski CA, Sloan DB, Tranel PJ, Westra P, Gaines TA. The Draft Genome

920   of *Kochia scoparia* and the Mechanism of Glyphosate Resistance via Transposon-

921   Mediated *EPSPS* Tandem Gene Duplication. Genome Biol Evol. 2019;11(10):2927-40.

922   147.   Zhang C, Johnson N, Hall N, Tian X, Yu Q, Patterson E. Subtelomeric *5-*

923   *enolpyruvylshikimate-3-phosphate synthase* copy number variation confers glyphosate

924   resistance in *Eleusine indica*. Nat Comm. 2023;14: 4865.

925   148.   Koo D-H, Molin WT, Saski CA, Jiang J, Putta K, Jugulam M, et al. Extrachromosomal

926   circular DNA-based amplification and transmission of herbicide resistance in crop weed

927   *Amaranthus palmeri*. Proc Natl Acad Sci U S A. 2018;115(13):3332-37.

928   149.   Molin WT, Yaguchi A, Blenner M, Saski CA. The EccDNA Replicon: A Heritable,

929   Extranuclear Vehicle That Enables Gene Amplification and Glyphosate Resistance in

930   *Amaranthus palmeri*. Plant Cell. 2020;32(7):2132-40.

931   150.   Jugulam M. Can non-Mendelian inheritance of extrachromosomal circular DNA-

932   mediated EPSPS gene amplification provide an opportunity to reverse resistance to

933   glyphosate? Weed Res. 2021;61(2):100-05.

934   151.   Wang L, Sun X, Peng Y, Chen K, Wu S, Guo Y, et al. Genomic insights into the origin,

935   adaptive evolution, and herbicide resistance of *Leptochloa chinensis*, a devastating

936   tetraploid weedy grass in rice fields. Mol Plant. 2022;15(6):1045-58.

937   152.   Cai L, Comont D, MacGregor D, Lowe C, Beffa R, Neve P, et al. The blackgrass genome

938   reveals patterns of non-parallel evolution of polygenic herbicide resistance. New Phytol.

939   2023;237(5):1891-907.

940   153.   Chen K, Yang H, Peng Y, Liu D, Zhang J, Zhao Z, et al. Genomic analyses provide

941   insights into the polyploidization-driven herbicide adaptation in *Leptochloa* weeds. Plant

942   Biotechnol J. 2023;21(8):1642-1658.

943   154.   Ohadi S, Hodnett G, Rooney W, Bagavathiannan M. Gene Flow and its Consequences in

944   *Sorghum spp*. Crit Rev Plant Sci. 2017;36(5-6):367-85.

945   155.   Renzi JP, Coyne CJ, Berger J, von Wettberg E, Nelson M, Ureta S, et al. How Could the

946   Use of Crop Wild Relatives in Breeding Increase the Adaptation of Crops to Marginal

947   Environments? Front Plant Sci. 2022;13:886162.

156. Ward SM, Cousens RD, Bagavathiannan MV, Barney JN, Beckie HJ, Busi R, et al.

Agricultural Weed Research: A Critique and Two Proposals. Weed Sci. 2014;62(4):672-

78.

157. Evans JA, Tranel PJ, Hager AG, Schutte B, Wu C, Chatham LA, et al. Managing the

evolution of herbicide resistance. Pest Manage Sci. 2016;72(1):74-80.

158. Montgomery JS, Giacomini D, Waithaka B, Lanz C, Murphy BP, Campe R, et al. Draft

genomes of *Amaranthus tuberculatus*, *Amaranthus hybridus* and *Amaranthus palmeri*.

Genome Biol Evol. 2020;12):1988-93.

159. Jeschke MR, Tranel PJ, Rayburn AL. DNA content analysis of smooth pigweed

(*Amaranthus hybridus*) and tall waterhemp (*A. tuberculatus*): implications for hybrid

detection. Weed Sci. 2003;51(1):1-3.

160. Rayburn AL, McCloskey R, Tatum TC, Bollero GA, Jeschke MR, Tranel PJ. Genome

size analysis of weedy *Amaranthus* species. Crop Sci. 2005;45(6):2557-62.

161. Mulligan GA. Chromosome numbers of Canadian weeds. I. Canad J Bot.

1957;35(5):779-89.

162. Meyer L, Causse R, Pernin F, Scalone R, Bailly G, Chauvel B, et al. New gSSR and

EST-SSR markers reveal high genetic diversity in the invasive plant *Ambrosia*

artemisiifolia L. and can be transferred to other invasive Ambrosia species. PLOS ONE.

2017;12(5):e0176197.

163. Pustahija F, Brown SC, Bogunić F, Bašić N, Muratović E, Ollier S, et al. Small genomes

dominate in plants growing on serpentine soils in West Balkans, an exhaustive study of 8

habitats covering 308 taxa. Plant Soil. 2013;373(1):427-53.

164.   Kubešová M, Moravcova L, Suda J, Jarošík V, Pyšek P. Naturalized plants have smaller

971   genomes than their non-invading relatives: a flow cytometric analysis of the Czech alien

972   flora. Preslia. 2010;82(1):81-96.

973   165.   Thébaud C, Abbott RJ. Characterization of invasive *Conyza* species (Asteraceae) in

974   Europe: Quantitative trait and isozyme analysis. Am J Bot. 1995;82(3):360-68.

975   166.   Garcia S, Hidalgo O, Jakovljević I, Siljak-Yakovlev S, Vigo J, Garnatje T, et al. New

976   data on genome size in 128 Asteraceae species and subspecies, with first assessments for

977   40 genera, 3 tribes and 2 subfamilies. Plant Biosyst. 2013;147(4):1219-27.

978   167.   Bennett MD, Leitch IJ, Hanson L. DNA amounts in two samples of angiosperm weeds.

979   Ann Bot. 1998;82):121-34.

980   168.   Schulz-Schaeffer J, Gerhardt S. Cytotaxonomic analysis of the *Euphorbia* spp.(leafy

981   spurge) complex. II: Comparative study of the chromosome morphology. Biol Zentralbl.

982   1989;108(1):69-76.

983   169.   Schaeffer JR, Gerhardt S. The impact of introgressive hybridization on the weediness of

984   leafy spurge. 1989 Leafy Spurge Symposium. 1989: 97-105.

985   170.   Bai C, Alverson WS, Follansbee A, Waller DM. New reports of nuclear DNA content for

986   407 vascular plant taxa from the United States. Ann Bot. 2012;110(8):1623-29.

987   171.   Aarestrup JR, Karam D, Fernandes GW. Chromosome number and cytogenetics of

988   *Euphorbia heterophylla* L. Genet Mol Res. 2008;7(1):217-22.

989   172.   Wang L, Sun X, Peng Y, Chen K, Wu S, Guo Y, et al. Genomic insights into the origin,

990   adaptive evolution, and herbicide resistance of *Leptochloa chinensis*, a devastating

991   tetraploid weedy grass in rice fields. Mol Plant. 2022;15(6):1045-58.

173.   Paril J, Pandey G, Barnett EM, Rane RV, Court L, Walsh T, et al. Rounding up the

annual ryegrass genome: High-quality reference genome of *Lolium rigidum*. Front Genet.

2022;13.

174.   Weiss-Schneeweiss H, Greilhuber J, Schneeweiss GM. Genome size evolution in

holoparasitic *Orobanche* (Orobanchaceae) and related genera. Am J Bot. 2006;93(1):148-

56.

175.   Towers G, Mitchell J, Rodriguez E, Bennett F, Subba Rao P. Biology & chemistry of

Parthenium hysterophorus L., a problem weed in India. Biol Rev. 1977;48):65-74.

176.   Moghe GD, Hufnagel DE, Tang H, Xiao Y, Dworkin I, Town CD, et al. Consequences of

whole-genome triplication as revealed by comparative genomic analyses of the wild

radish (*Raphanus raphanistrum*) and three other Brassicaceae species. Plant Cell.

2014;26(5):1925-37.

177.   Zhang X, Liu T, Wang J, Wang P, Qiu Y, Zhao W, et al. Pan-genome of *Raphanus*

highlights genetic variation and introgression among domesticated, wild, and weedy

radishes. Mol Plant. 2021;14(12):2032-55.

178.   Chytrý M, Danihelka J, Kaplan Z, Wild J, Holubová D, Novotný P, et al. Pladias

database of the Czech flora and vegetation. Preslia. 2021;93(1):1-87.

179.   Kersten S, Chang J, Huber CD, Voichek Y, Lanz C, Hagmaier T, et al. Standing genetic

variation fuels rapid evolution of herbicide resistance in blackgrass. Proc Natl Acad Sci U

S A. 2023;120(16):e2206808120.

180.   Chen J, Hall N, Saski C, Westra P, Gaines T, Patterson EL. *FHY3/FAR1* transposable

elements generate adaptive genetic variation in the *Bassia scoparia* genome. bioRxiv.

2023; DOI):10.1101/2023.05.26.542497.

181. Jarvis DE, Sproul JS, Navarro-Domínguez B, Krak K, Jaggi K, Huang Y-F, et al.

Chromosome-scale genome assembly of the hexaploid Taiwanese goosefoot "Djulis"

(*Chenopodium formosanum*). Genome Biol Evol. 2022;14(8.

182. Ferreira LAI, de Oliveira RS, Jr., Constantin J, Brunharo C. Evolution of ACCase-

inhibitor resistance in *Chloris virgata* is conferred by a Trp2027Cys mutation in the

herbicide target site. Pest Manag Sci. 2023;in press.

183. Laforest M, Martin SL, Bisaillon K, Soufiane B, Meloche S, Page E. A chromosome-

scale draft sequence of the Canada fleabane genome. Pest Manage Sci. 2020;76):2158-

69.

184. Guo L, Qiu J, Ye C, Jin G, Mao L, Zhang H, et al. *Echinochloa crus-galli* genome

analysis provides insight into its adaptation and invasiveness as a weed. Nat Commun.

2017;8(1):1031.

185. Sato MP, Iwakami S, Fukunishi K, Sugiura K, Yasuda K, Isobe S, Shirasawa K. (2023)

Telomere-to-telomere genome assembly of an allotetraploid pernicious weed,

Echinochloa phyllopogon. bioRxiv. 2023;2023.05.23.541891.

186. Wu D, Xie L, Sun Y, Huang Y, Jia L, Dong C, et al. A syntelog-based pan-genome

provides insights into rice domestication and de-domestication. Genome Biol.

2023;24(1):179.

187. Wang Z, Huang S, Yang Z, Lai J, Gao X, Shi J. A high-quality, phased genome assembly

of broomcorn millet reveals the features of its subgenome evolution and 3D chromatin

organization. Plant Commun. 2023;4(3):100557.

188. Mao Q, Huff DR. The Evolutionary Origin of *Poa annua* L. Crop Sci. 2012;52(4):1910-

22.

189. Benson CW, Sheltra MR, Maughan JP, Jellen EN, Robbins MD, Bushman BS, et al.

Homoeologous evolution of the allotetraploid genome of *Poa annua* L. Res Sq.

2023;DOI): 10.21203/rs.3.rs-2729084/v1.

190. Mo C, Wu Z, Shang X, Shi P, Wei M, Wang H, et al. Chromosome-level and graphic

genomes provide insights into metabolism of bioactive metabolites and cold-adaption of

Pueraria lobata var. *montana*. DNA Research. 2022;29(5.

191. Thielen PM, Pendleton AL, Player RA, Bowden KV, Lawton TJ, Wisecaver JH.

Reference genome for the highly transformable *Setaria viridis* ME034V. G3 (Bethesda,

Md). 2020;10(10):3467-78.

192. Yoshida S, Kim S, Wafula EK, Tanskanen J, Kim Y-M, Honaas L, et al. Genome

sequence of *Striga asiatica* provides insight into the evolution of plant parasitism. Curr

Biol. 2019;29(18):3041-52.e4.

193. Qiu S, Bradley JM, Zhang P, Chaudhuri R, Blaxter M, Butlin RK, et al. Genome-enabled

discovery of candidate virulence loci in *Striga hermonthica*, a devastating parasite of

African cereal crops. New Phytol. 2022;236(2):622-38.

194. Nunn A, Rodríguez-Arévalo I, Tandukar Z, Frels K, Contreras-Garrido A, Carbonell-

Bejerano P, et al. Chromosome-level *Thlaspi arvense* genome provides new tools for

translational research and for a newly domesticated cash cover crop of the cooler

climates. Plant Biotechnol J. 2022;20(5):944-63.

1057

 Captions for Figures

1059   Figure 1. The International Weed Genomics Consortium (IWGC) collected input from the weed

1060   genomics community to develop plans for weed genome sequencing, annotation, user-friendly

1061   genome analysis tools, and community engagement.

1062

1063   Table 1. Genome assemblies of 31 weed species completed or ongoing by the International

1064   Weed Genomics Consortium. All completed genomes are platinum assembly quality, defined as

1065   having chromosome-length scaffolds (i.e., 1-3 scaffolds per chromosome) for the assembly,

1066   unless indicated by *. Genome size estimated from flow cytometry or published references as

1067   indicated. + indicates that verification is currently in progress for cytogenetic information.

Scientific name	Common name	Haplotypes in Assembly	Anticipated Availability Date	Ploidy	x	n	Genome Size Estimate (Gbp)
Amaranthus hybridus	smooth pigweed	1; Previous version [158]	November 2023	diploid	16	16	0.509 [159]
Amaranthus palmeri	Palmer amaranth	Previous version [158]	June 2024	diploid	17	17	0.445 [160]
Amaranthus retroflexus	redroot pigweed		In progress	diploid	16	16	0.592 [160]
Amaranthus tuberculatus	common waterhemp	2; Previous version [158]	November 2023	diploid	16	16	0.694 [160]
Ambrosia artemisiifolia	common ragweed	Previous version [113]	In progress	diploid [161, 162]	18	18	1.152 [163]
Ambrosia trifida	giant ragweed	2	December 2023	diploid [161]	12	12	1.872 [164]

Apera spica-venti	loose silkybent	2	November 2023	diploid	7	7	4.622
Avena fatua	wild oat	1	November 2023	hexaploid (Additional file 1)	7	21	11.248
Chenopodium album	common lambsquarters	1	November 2023	hexaploid	9	27	1.59
Cirsium arvense	Canada thistle		In progress	diploid	17	17	1.415
Convolvulus arvensis	field bindweed		In progress	diploid[+]	12[+]	12[+]	0.652 [163]
Conyza bonariensis (Erigeron bonariensis)	hairy fleabane		In progress	hexaploid [165]	9	27	2.043 [166]
Conyza sumatrensis (Erigeron sumatrensis)	Sumatran fleabane	1	November 2023	hexaploid	9	27	1.874
Cyperus esculentus	yellow nutsedge	2	December 2023	diploid	54	54	0.588 [167]
Cyperus rotundus	purple nutsedge	2	December 2023	diploid	54	54	0.49 [167]
Digitaria insularis	sourgrass	1	November 2023	tetraploid	9	18	1.529
Digitaria ischaemum	hairy crabgrass		November 2024	tetraploid	9	18	0.625
Echinochloa colona	junglerice (weedy genotype)	See crop genotype assembly by [89]	October 2024	hexaploid	9	27	1.372 [167]
Euphorbia esula	leafy spurge		In progress	hexaploid[+] [based on 168, 169]	10[+]	60[+]	2.3 [170]

Euphorbia heterophylla	wild poinsettia		July 2024	diploid [171]	14	14	Unknown, in progress
Leptochloa chinensis	Chinese sprangletop	2; See also [172]	November 2023	tetraploid	10	20	0.454
Lolium rigidum	annual ryegrass	2; See also [173]	November 2023	diploid (Additional file 1)	7	7	2.41
Orobanche cernua	nodding broomrape		In progress	diploid	19	19	1.421 [174]
Orobanche crenata	crenate broomrape		In progress	diploid	19	19	2.787 [174]
Orobanche minor	small broomrape		In progress	diploid	19	19	1.792 [174]
Parthenium hysterophorus	ragweed parthenium		In progress	diploid [175]	17	17	Unknown, in progress
Phalaris minor	little seed canary grass	1	November 2023	tetraploid (Additional file 1)	7	14	5.851
Raphanus raphanistrum	wild radish	Previous versions [176, 177]	In progress	diploid	9	9	0.515 [176]
Salsola tragus	Russian thistle	2	November 2023	tetraploid (Additional file 1)	9	18	1.319
Sorghum halepense	johnsongrass	2	January 2024	tetraploid	10	20	1.752
Verbascum blattaria	moth mullein	1	December 2023	diploid	15	15	0.344 [178]

1068

1069

1070

1071   Table 2. Genome assemblies and genomic information for 22 weed species produced by other

1072   groups independently of the International Weed Genomics Consortium. Haploid (1n) genome

1073   size estimations are either calculated through flow cytometry or k-mer estimation.

Scientific name	Common name	x	n	1n Genome Size Estimate (Gbp)	Genome Assembly Size (Gbp)
Alopecurus myosuroides	blackgrass	7	7	3.56 [179]	3.4-3.56 [152, 179]
Bassia scoparia	kochia	9	9	0.969 [180]	0.970 [180]
Bromus tectorum	cheatgrass	7	7		2.48 [101]
Chenopodium formosanum (domesticated genotype of *C. album*)	Djulis	9	27	1.69 [181]	1.59 [181]
Chloris virgata	feather finger grass	20	20		0.336 [182]
Conyza canadensis (*Erigeron canadensis*)	horseweed	9	9	0.425 [183]	0.426 [183]
Echinochloa colona (crop genotype)	junglerice	9	27	1.18 [89]	1.13 [89]

Echinochloa crus-galli	barnyardgrass	9	27	1.4 [184]	1.34 [89]
Echinochloa oryzicola (syn. *E. phyllopogon*)	late watergrass	9	18	1.0 [184, 185]	0.95 [89] 1.0 [185]
Eleusine indica	goosegrass	9	9		0.510 [147]
Ipomoea purpurea	common morning-glory	15	15	0.81 [134]	0.60 [134]
Lolium perenne	perennial ryegrass	7	7		2.63 (Bushman and Robbins, pers. comm.)
Oryza sativa f. spontanea	weedy rice	12	12		0.37 [186]
Panicum miliaceum	wild proso millet	9	18		0.85 [187]
Poa annua	annual bluegrass	7	14	1.78 [143]	1.89 [188]
Poa infirma	early meadow-grass	7	7	1.17 [189]	1.13 [189]
Poa supina	supine bluegrass	7	7	0.66 [189]	0.64 [189]
Pueraria montana var. *lobata*	kudzu	11	11		0.98 [190]
Setaria viridis	green foxtail	9	9	0.40 [191]	0.40 [191]
Striga asiatica	red witchweed	12	12	0.6 [192]	0.47 [192]

Striga hermonthica	purple witchweed	10	20	1.48 [193]	0.96 [193]
Thlaspi arvensis	field pennycress	7	7	0.5 [194]	0.53 [194]

1074

1075

1076

1077 **Additional Files**

1078 Additional File 1 (.docx). Methods and results for visualizing and counting the metaphase

1079 chromosomes of (1A): diploid *Lolium rigidum*; (1B): hexaploid *Avena fatua*; (1C): diploid

1080 *Phalaris minor*; and (1D): tetraploid *Salsola tragus*.

1081

1082 Additional File 2 (.docx). List of completed and in-progress genome assemblies of weed species

1083 pollinated by insects.