Effects of psychedelics on neurogenesis: A systematic review of pre-clinical studies

Rafael Vitor Lima da Cruz[1]*, Richardson N. Leão[1] , Thiago C. Moulin[2,3]

1. Neurodynamics Lab, Brain Institute (ICe) Universidade Federal do Rio Grande do Norte, Natal, Brazil.
2. Department of Experimental Medical Science, Lund University, Lund, Sweden.
3. Department of Surgical Sciences, Division of Functional Pharmacology and Neuroscience, Uppsala University, Uppsala, Sweden.
*Corresponding author: rafael.lima@neuro.ufrn.br

Abstract

In the mammalian brain, new neurons continue to be generated throughout life in a process known as adult neurogenesis. The role of adult-generated neurons has been broadly studied across laboratories, and mounting evidence suggests a strong link to the HPA axis and concomitant malfunction in patients diagnosed with mood disorders. Psychedelic compounds, such as phenethylamines, tryptamines, cannabinoids, and a variety of ever-growing chemical categories, emerge as therapeutic options for neuropsychiatric disorders, while numerous reports link their effects to increased adult neurogenesis. In this systematic review, we examine studies assessing neurogenesis after psychedelics interventions and aim to provide a comprehensive picture of how this vast category of compounds regulates the generation of new neurons. Following PRISMA guidelines, we conducted a literature search on PubMed and Science Direct databases considering all articles published until January 31, 2022. We selected studies containing both the words "neurogenesis" and "psychedelics", using either *in vivo* or *in vitro* models, and employing classical or atypical psychedelics at all ontogenetic windows. Of a total of 205 articles, 68 met all the necessary conditions for further review. Our findings were divided into five main categories of psychedelics: CB1 agonists, NMDA antagonists, harmala alkaloids, tryptamines, and entactogens. We then described the outcomes of neurogenesis assessments and investigated related results on the effects of psychedelics on brain plasticity and behavior within our sample. In summary, this review presents an extensive study into how different psychedelics may affect the birth of new neurons and other brain-related processes. Such knowledge may be valuable for future research on novel therapeutic strategies for neuropsychiatric disorders.

Keywords: Psychedelics; Hallucinogens; Neurogenesis; Dentate gyrus; Plasticity; Major Depression.

Introduction

According to the Global Burden Diseases study, used by the World Health Organization (WHO) for strategic planning, 264 million people, or about 4.5% of the world population, suffer from Major Depressive disorder (MDD). It is recognized as one of the most debilitating illnesses on a global scale, substantially significantly affecting daily activities, quality of life, cognitive abilities, and work productivity (James et al., 2018). MDD can be defined as a sustained anhedonic state that can be continuous or episodical, strongly influences self-esteem and significantly impacts social, family and professional life (Lépine & Briley, 2011). Together with other mood and anxiety disorders, those diseases are the most prevalent mental illnesses and the third most prevalent cause of disability, contributing to the global burden of disease (WHO, 2012). The majority of pharmacological interventions aiming to treat mood disorders such as MDD are benzodiazepines or Selective Serotonin Reuptake Inhibitors (SSRIs). However, these classes of drugs do not elicit positive outcomes for about 50 to 60% of patients, which is then characterized as treatment-resistant depression (TRD) (Nestler et al., 2002). SSRIs, the most modern class of antidepressants, are taken daily with an onset of the desired effects close to one month after the beginning of treatment. Yet, these medications can trigger adverse effects that appear early on and last for the duration of the therapy. These drugs also have a high risk of being misused, as individuals undergoing treatment tend to become physically dependent or addicted, even with the accompanying lethargy induced by them (Wong and Licinio, 2001).

The pathophysiology of depression is not yet fully understood; however, empirical data from classical antidepressants led to the widely accepted monoamine hypothesis, which predicts that this disorder arises from a deficiency or imbalance of monoamine neurotransmitters. It is worth noting that a number of studies provide support for this theory. First, standard antidepressants primarily operate on the monoamine neurochemical route, with the goal of re-establishing dopamine (DA), noradrenaline (NA), and serotonin (5-HT) levels to homeostatic concentrations

(Heninger et al., 1996); Second, monoamine antagonists like reserpine, typically used for arterial hypertension, can induce depressive symptoms when taken over extended periods (Baumeister et al., 2003; Freis, 1954; De Freitas et al., 2016); Third, treatments for MDD and anxiety disorders usually require chronic, daily dosages for at least a month to produce meaningful effects (Kempermann, 2002). The latter observation has also led to a reinterpretation of the long-standing monoamine hypothesis of depression to what is now termed the neurogenic hypothesis of depression. This revised theory suggests that depression correlates with a decrease in the formation of new neurons in the adult brain, a process that seems to be revived by prolonged antidepressant treatment (Jacobs et al., 2000).

Adult neurogenesis is the process by which new neurons are continuously added to the brain throughout the life of an organism. Neurogenesis seems to be ubiquitous to all species with a central nervous system (Barker et al., 2011), and for many of them, the process is confined to specific regions (Barnea and Pravosudov, 2011; Drew et al., 2013). In rodents, it is restricted to two zones, the olfactory bulb, driven by the neural stem cells (NSCs) located in the subventricular zone (SVZ), and the dentate gyrus sub-region from the hippocampus, driven by the radial glial-like cells (RGL) (Laplagne et al., 2006). The foundations of the neurogenic theory of depression are supported by empirical data from clinical and preclinical studies drawn to understand how the neurogenesis process is reverted to homeostatic levels when SSRI chronic treatment is applied (Miller and Hen, 2015). However promising, alternative pathways to the proposed hypothesis are under discussion (Data-Franco et al., 2017; N. X. Li et al., 2022; Raphael Mechoulam & Parker, 2013; Sanches et al., 2021; Yuan et al., 2015) and new biochemical routes to treat depression are emerging, including the induction of neurogenesis independent of direct 5-HT modulation (Idell et al., 2017; Reiche et al., 2018). Among the chemical candidates for novel antidepressants, encouraging results have been found with the use of psychedelics (Aleksandrova & Phillips, 2021; DeVos & Miller, 2013; Muttoni et al., 2019).

Research with hallucinogens began in the 1960s when leading psychiatrists observed therapeutic potential in the compounds today referred to as psychedelics (Osmond, 1957; Vollenweider and Kometer, 2010). Those psychotomimetic drugs are often, but not exclusively, serotoninergic agents (Belouin and Henningfield, 2018; Sartori and Singewald, 2019a) and were central to the anti-war mentality in the "hippie movement". This social movement brought much attention to the topic, leading to the 1971 UN convention of psychotropic substances that classified psychedelics as class A drugs, enforcing maximum penalties for possession and use, including for research purposes (Ninnemann et al., 2012), essentially criminalizing psychiatric studies at that time. Although those pioneer studies are now considered as lacking scientific or statistical rigor (Vollenweider and Kometer, 2010), they were the first to suggest the clinical use of these substances, which has been supported by recent data from both animal and human studies (Danforth et al., 2016; Nichols, 2004; Sartori and Singewald, 2019b). Moreover, some psychedelics are currently used as treatment options for psychiatric disorders. For instance, ketamine is prescriptible to treat TRD in USA and Israel, with many other countries implementing this treatment (Mathai et al., 2020); other drugs, such as the 3,4-Methylenedioxy methamphetamine (MDMA), are in the last stages of clinical research and might be employed for the treatment of post-traumatic stress disorder (PTSD) with assisted psychotherapy (Emerson et al., 2014; Feduccia and Mithoefer, 2018; Sessa, 2017) while Australia is the first nation to legalize the compassionate use of psilocybin for mental health issues such as mood disorders .

Psychedelics act mainly through brain plasticity, initially changing neuronal functionality at the molecular level by producing electrophysiological changes that stimulate neurotrophic signaling (Browne & Lucki, 2013; Castrén et al., 2007; Magaraggia et al., 2021; Muscat et al., 2021). Ultimately, they induce neurite growth (Numakawa et al., 2010; Saengsawang & Rasenick, 2016; V. B. B. Thompson et al., 2012), synaptic remodeling (R.-J. Liu et al., 2013; Zhou & Song, 2001), neurogenesis (García-Cabrerizo & García-Fuster, 2016; Lima da Cruz et al., 2018; F. Liu et al., 2017), and oxidative stress reduction (Frecska et al., 2013, 2016; Szabo, 2015). Thus, it is believed that psychedelics can create a window of opportunity for therapists to introduce cognitive-behavioral treatment strategies and produce long-lasting effects, which are independent of the classical pharmacological approaches to treat the hypothesized neurotransmitter imbalance (Keeler et al., 2021; Nichols, 2016; Worrell and Gould, 2021). Such a holistic and personalized approach can better integrate patients into the treatment process, reducing the current disconnection between popular beliefs on mental illnesses and scientific-guided psychiatric interventions (Healy, 2004; Lacasse & Leo, 2005).

Despite the encouraging perspectives on the applications of psychedelics, their safe employment requires a deeper understanding of their mechanisms, as the currently available compounds generally target multiple neurotransmitter systems and may lead to undesired effects (Belouin & Henningfield, 2018; Brunton et al., 2011; Geyer et al., 2009). Moreover, the effects on brain physiology are shown to depend on ontogeny (J. Liu et al., 2006; Riga et al., 2016; Skaper & Di Marzo, 2012), gender (Lee et al., 2014; Realini et al., 2011; Rubino et al., 2008), dose (Fortunato et al., 2009; Maeda et al., 2007; Marinova et al., 2017) and chemical interactions (Canales & Ferrer-Donato, 2014; Zuo et al., 2018). For this reason, we sought to cover the effects of such compounds on the plasticity process of neurogenesis (Christie & Cameron, 2006; Kempermann, 2012). To categorize those compounds, we adapted a classification done elsewhere (Calvey & Howells, 2018). Finally, we discuss findings encompassing any effect in the molecular, cellular, physiological and behavioral levels reported for *in vivo* or *in vitro* models related to neurogenesis.

Methods

Search and inclusion criteria

We followed the Preferred Reported Items for Systematic Reviews and Meta-analysis (PRISMA) guidelines as a way to standardize the workflow (Liberati et al., 2009). All relevant articles were selected during a search using the PubMed and Science Direct databases in January 2023, where we searched for the combination of the words "psychedelics" and "neurogenesis", including their MeSH terms, without limitations regarding publication date. Titles and abstracts were first scanned for articles presenting original results involving the assessment of neurogenesis with the intervention of psychedelic compounds. Psychedelics were defined and classified as described in (Calvey & Howells, 2018). Articles not written in English were also considered, but none passed our inclusion criteria, leaving only articles written in English to our final sample. Posterior to the systematic search, a reference screening on all reviews gathered was performed. Experimental articles with titles containing any psychedelic drug were reached for abstract analysis under the same inclusion criteria. Selected articles underwent an elimination process, excluding duplicates, absent abstracts, conference notes, editorials, book chapters, and case reports. Articles that did not intend to test or discuss psychedelics in any organic model and did not report any effect of psychedelics on neurogenesis or neuronal plasticity were removed. Then, we sorted reports by the drug chemical group for further scrutiny. Reviews were not included in the analysis but considered in the discussion of results. The PRISMA flowchart for article selection is depicted in **Figure 1.**

Data extraction and analysis

Articles were organized in an Excel file by (1) article information, containing first and last author, author affiliation, journal, title, year of publication, and psychedelic chemical group; (2) methods information, concerning dose and/or concentration, treatment regimen, experimental model of choice, techniques used; and (3) summary of results, highlighting outcome variables and reported effect on neurogenesis. For analysis, a study location map (**Figure 2**) was generated using a web tool (www.mapinseconds.com), analysis of publications over the years (**Figure 3**) were performed using Microsoft Office Excel and GraphPad Prism.

Methodological quality assessment

We performed a qualitative analysis of the *in vivo,* animal and human studies, applying the criteria from SYRCLE (Hooijmans et al., 2014) and OHAT (OHAT, 2015) risk of bias (RoB) assessment tools, respectively. Both tools are focused on methods application, randomization of subjects, experimenter blinding, outcome measurements and report methods. The results are reported on **Figure 4** for *in vivo* studies, and **Figure 5** for Human studies. Excel tables were generated for all categories, summarizing the method and results. *In vitro* studies were not analyzed for risk of bias. Considering the purpose of this review, electrophysiological assessment of acute slices was considered *ex vivo* and thus analyzed with the animal studies via SYRCLE RoB. However, long-term cultured neurons, even when removed from embryos, were not considered *ex vivo* but *in vitro*. When articles contain *in vitro* and *in vivo* experiments, their results are reported independently.

For the first criteria of SYRCLE RoB: *1- Was the allocation sequence adequately generated and applied?* concerning randomization of treatment, due to the poor reporting of our sample in this category, authors that at least described some randomization no matter which one, were graded positively.

Also, for the second criteria of the SYRCLE RoB: *2- Were the groups similar at baseline, or were they adjusted for confounders in the analysis?* We considered as essential for quality of report in these criteria age and gender, and when necessary, disease model onset as the minimum requirement for a study to be classified positively.

Figure 1. Flowchart depicting the selection of articles in accordance with PRISMA guidelines.

Results

Article search and inclusion

The present systematic review followed the recommendations of the PRISMA Statement (Liberati et al., 2009) and included articles from two databases (PubMed and Science Direct) to identify articles related to the effects of psychedelics on neurogenesis and neuronal plasticity. A total of 205 articles were initially identified, of which 75 were from PubMed and 130 were from Science Direct. First, we assessed the abstract of selected articles and removed conference notes, editorials, book chapters, and case reports (n=69) and excluded 32 publications that did not test or discuss the effects of psychedelics on neurogenesis or neuronal plasticity. Four additional articles were excluded after a full-text analysis for the same reasons. Lastly, narrative reviews were screened for additional article selection, following the same criteria mentioned above,

resulting in 21 additional references. Our selection process resulted in 68 original research articles for further analysis. **Figure 1** summarizes the searching and selection method.

Following the article selection, we extracted metadata such as location, year of publication, publisher, and author information. We plotted the locations of the laboratories that published articles on the topic on a map **(Figure 2)**. It is worth noting that research on psychedelics was largely prohibited by the UN Act of 1971 (Ninnemann et al., 2012), making their research illegal in most countries (Rucker et al., 2018). As a result, the oldest articles on these databases related to the topic date back to 2001. This is likely due in part to the fact that neurogenesis in adult mammals was only described in 1977 (Kaplan & Hinds, 1977) and the process was not widely accepted as occurring in humans until 1998

(Eriksson et al., 1998), which is still controversial for part of the neuroscience community (Sorrells et al., 2018). Interest in psychedelics has increased over the years, with peak publication years in 2013 (8 experimental articles and 2 reviews), 2018 (9 experimental articles and 6 reviews), and 2020 (8 experimental articles and 14 reviews), as shown in **Figure 3**.

Figure 2. Localization of laboratories publishing articles using psychedelics to study neurogenesis.

Figure 3. Distribution of included articles throughout the years

Descriptive statistics

In this study, we identified 68 experimental articles examining the effects of psychedelics on various outcomes. Of these, 6 were conducted in humans, 44 were conducted *in vivo*, and 11 were conducted *in vitro*. In addition, 7 of the studies used both in vivo and in vitro approaches. We report the results of each chemical category independently in the relevant sections of this article. Among the studies, the largest number were conducted in the USA (n=18), with a focus on ketamine (n=7) and cannabinoids (n=4). These substances have gained legal status in some states in the USA in recent years (Andrade, 2017; Carliner et al., 2017). Many of the articles from this country focus on the potential teratogenic effects or collateral damage associated with the use or abuse of these substances. Brazil was the second most represented country, with 9 original articles focusing on β-carbolines and tryptamines (n=8). In Brazil, the *Ayahuasca* tea is legal for religious purposes (Svobodny, 2014). The majority of the articles in this study utilize the *Ayahuasca* concoction or its main components, as it is a traditional drink consumed ritualistically by indigenous people in the Amazonian basin (Labate & Cavnar, 2013). It is typically made from two plant species, *B. caapi* and *P. viridis*, though there may be significant variations depending on the region (de Rios & Grob, 2005). In recent years, *Ayahuasca* has gained a reputation as a miraculous medicine and has become more widely available globally, leading to concerns about its environmental impact (Holman, 2010).

Human studies

Of the 68 experimental articles identified in this study, 6 involved human participants. One of these was a case-control study examining the balance of CBD and THC phytocannabinoids in hair samples and its relationship to hippocampal mass (Demirakca et al., 2011). The other five studies explored the effects of ketamine in humans, including one cross-sectional study conducted with ketamine-addicted patients in Guangzhou, China (Fan et al., 2015), and four controlled trials (Dakwar et al., 2018; Duncan et al., 2013; Haile et al., 2014; Rybakowski et al., 2013). A summary of the findings from these studies can be found in **Supp. Tables 1-6**, and a quality assessment using the OHAT risk of bias tool is presented in **Figure 5**. The results of these studies are discussed in the following sections.

In vitro studies

Of the 18 original articles that included *in vitro* experiments, the majority utilized cannabinoid agonists (52.6%) and tryptamines (21%). The in vitro model most commonly used was cultured cortical neurons extracted from rats at different stages of embryonic development (26%). The main findings and discussion points of each of these studies are outlined in **Tables 1-6**.

In vivo studies

Of the 51 original articles that included in vivo experiments in non-human animals, the most commonly used psychedelics were cannabinoid agonists (29.4%), NMDA antagonists (27.4%), and tryptamines (17.6%); however, a great range of drugs and dosages were used in these studies. The animal models most frequently used were rats (66.6%), followed by mice (31.4%). Some studies used more than one animal model, including chick embryos, Xenopus frogs, and drosophila larvae. We performed a quality of report analysis using the SYRCLE Risk of Bias tool (Hooijmans et al., 2014), and results are summarized in **Figure 4**. Nearly all of the in vivo studies did not report a valid method of randomization, so we reduced the quality level of the report as recommended by the SYRCLE initiative. We graded these studies as "Yes" if the authors at least mentioned some method of randomization, regardless of its robustness. The second major source of bias in these studies was the lack of control over housing conditions, as only one article described the housing of experimental animals during the experiment in a satisfactory manner., In addition, for criteria 7 (40%), 3 (24%), 5 (14%), and 6 (12%), the number of studies graded as "Yes" did not reach 50% of the total pool, indicating a lack of rigor in in vivo studies, which is a common issue not only in psychedelic science but across the life sciences. This highlights the importance of increasing reproducibility in biological research (Begley & Ioannidis, 2015; Ioannidis, 2014). In the following sections, we will delve into each chemical category in more depth, with percentages in parentheses indicating the proportion of the total number of articles (n=68) unless otherwise stated.

Figure 4. Results of SYRCLE Risk of Bias Quality assessment of included studies using experimental animals.

Figure 5. Results of OHAT Risk of Bias quality assessment of included studies using human subjects.

CB1 agonists

Cannabinoids are unique in the central nervous system (CNS). The CB1 receptor was first cloned in Tom Bonner's lab (Matsuda et al., 1990) and is predominantly expressed in neurons, where it is responsible for the psychoactive effects of cannabinoids (Raphael Mechoulam & Parker, 2013). The CB1 receptor remained an "endogenous orphan" until 1999, when Raphael Mechoulam's team discovered the arachidonoylethanolamide (AEA) and the 2-arachidonoylglycerol (2-AG)(Martin et al., 1999). In contrast, the CB2 receptor is less prevalent in the brain but is widely expressed in the immune system (Onaivi, 2006; Pertwee, 2006). While we have a good understanding of the effects of cannabinoids on humans based on observational studies of subjects who smoke plants in the Cannabis genus,

which contain the phytocannabinoids THC and CBD (Bonini et al., 2018; Chang & Chronicle, 2007), the mechanisms underlying their homeostatic function are only now beginning to be understood (de Fonseca et al., 2005).

The endocannabinoids system (eCB system) is a complex and widespread system that acts retrogradely as a feedback channel, influencing the release of all classical neurotransmitters (Skaper & Di Marzo, 2012) and enabling short- and long-term adaptive responses that lead to plasticity changes through mechanisms that are not yet fully understood (Araque et al., 2017). While not all cannabinoids are psychoactive and only some are considered psychedelics (Bonini et al., 2018), we have chosen to include CB1 agonists because their availability as medicine is increasing worldwide, and because their synthetic analogs have varying potencies and behavioral effects that have not yet been fully described.

In our review, we identified 21 *in vitro* and *in vivo* studies examining the effects of five different cannabinoids: synthetic compounds HU210 (23.8%), Win55212-2 (9.5%), and O-2545 (4.7%), the endocannabinoid AEA (14.2%), and the phytocannabinoid THC (61.9%). Only one report included humans, a case-control study (Demirakca et al., 2011). Most of the experiments were conducted *in vivo* (71.42%), with a focus on adolescent murine models (47%) in an effort to better understand the widespread misuse of Cannabis plants, which is the most commonly used illicit drug globally, particularly among youth (Peacock et al., 2018). All of the studies used molecular or cellular methods of measurement, with a lesser focus on behavioral (38%) and electrophysiological (9.5%) measures. Descriptive results are summarized in **Supp Table 1**.

The relationship between neurogenesis and THC, as detailed in Table 2, appears conflicting due to differing study designs. For instance, the THC dosage varies greatly among studies, with some administering significantly high doses up to 30 mg/kg (Downer et al., 2007; Kochman et al., 2006) — levels not generally therapeutic and potentially excessive even for high tolerance individuals (Karila et al., 2014). Conversely, other studies have examined a wide range of THC doses, from 0.75 to 10 mg/kg (Beiersdorf et al., 2020; Cuccurazzu et al., 2018; Downer et al., 2007; Kochman et al., 2006; Leishman et al., 2018; Poulia et al., 2021; Realini et al., 2011; Rubino et al., 2008; Steel et al., 2014; Suliman et al., 2018). The time point of intervention and analysis also significantly diverges across the studies. Three distinct intervention periods are identifiable: embryonic/early post-natal phase exposure (Beiersdorf et

al., 2020; Berghuis et al., 2007; Downer et al., 2007; Psychoyos et al., 2008); exposure during adolescence, specifically in rats between P28-P45 (Cuccurazzu et al., 2018; Lee et al., 2014; Poulia et al., 2021; Rubino et al., 2008; Steel et al., 2014; Suliman et al., 2018); and exposure during adulthood (Jiang et al., 2005; Kochman et al., 2006; Leishman et al., 2018; Rueda et al., 2002), as well as in-vitro studies with various models (Shum et al., 2020; Stanslowsky et al., 2017; F. Zhang et al., 2009; Zhou & Song, 2001).

As shown in **Table 1**, several studies reported negligible impact of acute or chronic CB1 agonists on neurogenesis (Berghuis et al., 2007; Downer et al., 2007; Kochman et al., 2006; Stanslowsky et al., 2017; Steel et al., 2014). Importantly, it should be noted that some studies focused not directly on neurogenesis, but on other indirect plasticity markers (Berghuis et al., 2007; Poulia et al., 2021; Stanslowsky et al., 2017) or teratogenic morphological effects (Beiersdorf et al., 2020; Downer et al., 2007; Psychoyos et al., 2008). Yet, three studies identified positive effects on adult plasticity, specifically with regards to the survival of newborn neurons and neurite growth, both in vivo and in culture (Jiang et al., 2005; Suliman et al., 2018; F. Zhang et al., 2009).

Further into our analysis, we identified notable findings that warrant a more detailed discussion, each contributing unique insights into the complex effects of cannabinoids on neural functioning and development. Parolaro's group, for instance, employed a chronic, escalating THC regimen (11 days, twice a day) spanning three distinct age windows (2.5mg/kg in P35-P37; 5mg/kg in P38-P41; 10mg/kg in P42-P45). Their findings highlighted three significant outcomes: Firstly, they found that chronic THC exposure during adolescence indirectly reduced Brain-Derived Neurotrophic Factor (BDNF) via the CREB pathway in the hippocampus and prefrontal cortex (PFC) of males, while paradoxically increasing BDNF in the nucleus accumbens (NAc) of females (Rubino et al., 2008). Secondly, they observed a reduction in cell proliferation in females, an effect that was prevented by the fatty acid amide hydrolase (FAHH) inhibitor URB597 (Poulia et al., 2021; Realini et al., 2011) Finally, extending their earlier work, they demonstrated that such a THC exposure regimen could decrease not only the proliferation but also the survival rates of adult-born granule cells (abGC) in female rats (Cuccurazzu et al., 2018).

Broadening the perspective, Antoniou's group in Greece demonstrated that chronic exposure to THC during

adolescence disrupts the balance of dopamine transmission in several brain regions, including the prefrontal cortex, hippocampus, nucleus accumbens, and striatum. These findings provide a broader understanding of the molecular consequences of cannabis abuse during adolescence (Poulia et al., 2021). Lee et al. used a similar chronic exposure design, escalating through adolescence with the synthetic cannabinoid HU210, with dose ranges in the same age windows mentioned earlier (0.025 – 0.050 – 0.100 mg/kg). Their work resulted in the discovery of reduced proliferation and survival rates, but only in male mice evaluated at P70 (Lee et al., 2014).

In a study involving a transgenic model of Alzheimer's Disease (AD) APP23/PS45, Chen et al. administered a daily dose of HU210 (0.010mg/kg) chronically for 10 and 20 days to both old and adolescent mice. Their results showed a reduction in BrdU+ cell numbers in young mice, and a notable observation was that most of the animals (16/20) did not survive until the end of the planned treatment (20 days), which led the authors to halve the treatment duration, indicating the potentially harmful effects of HU210, especially given its 100-800 times higher potency compared to THC (Chen et al., 2010; R. Mechoulam et al., 1988). However, the authors' focus was primarily on AD-related features, which limits extrapolation of these neurogenesis reduction findings to a scenario involving lower doses and healthy conditions.

Leishman and colleagues took a different approach, administering a single 3 mg/kg THC injection to female CD1 rats at three different age windows (P35 – P60 – P90), followed by lipidome and transcriptome analyses 2 hours later. They found eCB levels were reduced across the brain, with the greatest impact observed in the adult hippocampus (Leishman et al., 2018). Additionally, Beiersdorf et al.'s study involved proteomics analyses of a pre-natal intervention (THC 1, 5 mg/kg from P5-P16 or P5-P35 days s.c.) and reported severe deficits in mitochondrial function at P48 and P120, indicating long-term consequences from early THC exposure (Beiersdorf et al., 2020).

Moreover, Rueda et al. reported that 24-hour *in vitro* exposure to the eCB AEA (5 µm) reduced neurite growth, with this reduction mediated by ERK signaling. Their findings were based on a variety of biochemical tools, and they also noted a decrease in the survival rates for abGC in adult rats (exact age not reported) following a 4-day *in vivo* exposure to M-AEA (daily 5mg/kg IP). (Rueda et al., 2002). Similarly, Zhou and Song's study, which involved acute exposure of neuroblastoma N1E-115 cells to HU210 (0.01-

100nM), showed that neurite growth was greatly dampened in a dose-dependent manner. This effect was mediated by intracellular cAMP (Zhou & Song, 2001). Lastly, Shum et al. proposed another mechanism for CB1R mediated dampening of neurite growth in vitro: the involvement of ERK and Akt signaling pathways. They showed that neurite growth ratio could be restored to control levels by the CB1R antagonist SR141716, at least in the case of human induced Pluripotent Stem Cells (hiNSC) differentiated into neurons (Shum et al., 2020).

We believe that extrapolating these results to a wider population warrants caution, particularly considering the gender discrepancies observed in murine models and their possible lack of translationability to humans. Such differences challenge the direct applicability of these findings to patients (Naqvi et al., 2019). In regions where Cannabis sativa is legal, an opportunity exists for researchers to conduct human studies investigating the gender-specific effects and side effects of cannabis use, as exemplified by the research from two Canadian studies (Matheson et al., 2020; Matteau-Pelletier et al., 2020).

From a neurobiological standpoint, it is postulated that CB1 agonists could potentially influence neurogenesis by modulating 5-HT and ACh neurons in the DRN and MS, respectively. These neurotransmitters are known to positively regulate different steps in neurogenesis (Raphael Mechoulam & Parker, 2013). However, the intricate relationship between the eCB system and other neurotransmitters, further complicated by species-specific variations, adds a layer of complexity to this field of study. Amidst these persisting ambiguities surrounding the mechanistic action of cannabinoids on the brain, it is pertinent to note the reported benefits of both phytocannabinoids and their synthetic counterparts. These compounds have been observed to confer antidepressant and anxiolytic effects in patients experiencing depression and those undergoing chemotherapy (Rodriguez Bambico et al., 2009). This is in the backdrop of a longstanding relationship between our species and these compounds, a relationship that can be traced back to as early as 12,000 years A.C (Ren et al., 2021), In light of their proven safety, it is prudent to continue the exploration of the potential benefits these compounds may offer. Importantly, considerations of the implications of their use and misuse should be informed by a global perspective, taking into account data collected from human populations across the world (Ferber et al., 2019; Garani et al., 2021).

Table 1 Results reported by the reviewed CB1 agonists studies

Outcome variable	Reported Results	Effect on Neurogenesis	Reference
***In vivo* studies**			
			(Berghuis et al., 2007)
Behavioral assessment (MWM; FC) **Physiological assessment** (Aβ deposition and β-Amyloid Precursor Protein; BRDU + cells)	No effect on all tests except for HU210 ↓ BRDU+ cells in young mice	HU210 ↓ BRDU+ cells only in young mice	(Chen et al., 2010)
Behavioral assessment (FST, NSFT) **Physiological assessment** (BrdU+, NeuN levels, TUNEL labeling)	Acute HU210 treatment ↑ BrdU+ cells; Chronic HU210 does not change differentiation; Chronic HU210 ↓ score in FST and NSFT; X-ray ablation of adult Hipp neurogenesis prevents Hu210 effect on FST, NFST.	Acute HU210 treatment ↑ BrdU+ cells; Chronic HU210 does not change differentiation; X-ray ablation of adult Hipp neurogenesis prevents Hu210 effect on FST, NFST	(Jiang et al., 2005)
Behavioral assessment (Stereotypy) **Physiological assessment** (BrdU+ cells, NeuN, Cort)	HU210 in adolescence ↓ BrdU+, NeuN+ cells in male but not female; HU210 in adolescence ↑ Cort levels in male but no female; HU210 in adolescence ↑ Stereotypy in female but not male.	HU210 in adolescence ↓ proliferation and survival in adulthood of male mice	(Lee et al., 2014)
Physiological assessment (BrdU+, NeuN+ cells)	M-AEA ↓ BrdU+ cells and NeuN+ cells.	M-AEA ↓ newborn cell survival.	(Rueda et al., 2002)
Physiological assessment (Lipidome and Transcriptome fold changes)	THC changes Lipidome and Transcriptome in an age dependent manner differentially for each brain region.	Negatively impact genes and lipids implicated in neurogenesis, effect is higher on the adult brain.	(Leishman et al., 2018)
Physiological assessment (JNK; Casp-3)	THC ↑JNK and Casp-3 activity.	THC ↑ neonatal cell apoptosis.	(Downer et al., 2007)
Behavioral assessment (EPM; OF; FST; SPT) **Physiological assessment** (CB1r levels; pCREB)	THC ↑ score in FST only in females; THC ↑ score in SPT for both genders; THC ↓ CB1r levels in Amg, VTA and Nac in females; THC ↓ CREB activity in Hipp and PFC in males but ↑ in Nac in females; THC ↓ BDNF indirectly by CREB in Hipp and PFC in males; THC ↑BDNF in NAc of females.	THC ↓ BDNF indirectly by CREB in Hipp and PFC in males but ↑ BDNF in NAc of females.	(Rubino et al., 2008)
Behavioral assessment (FST, PFP, NOR, SIT)	THC ↑ score in FST, PFP; THC ↓ Score in NOR; THC ↓ BrdU+ cells;	THC in adolescence ↓ BRDU+ cells in adulthood, Reversed by URB597	(Realini et al., 2011)

Physiological assessment (BRDU+ cells)	THC has no effect on pCREB or CB1r density in VTA hip or Nac In adult female.		
Physiological assessment (CB1r, PSD95, Syn1, Syn3, Ki67, DCX, PSA-NCAM, BrdU+ cells)	THC ↓ CB1r, PSD95,Syn1, Syn3 increase by cognitive training; THC has no effect on PSA-NCAM and DCX + cells increase by cognitive training; THC has no effect on Hipp BRDU + cells	Paradoxal results: THC ↓ plasticity markers (CB1r,Syn1,syn3,PSD95) increase by cognitive training however no change is observed in the overall cell proliferation (BrdU+ cells) and cell death (DCX+ and PSA-NCAM + cells) ; no change is found in the absence of cognitive training.	(Steel et al., 2014)
Behavioral assessment (NOD) **Physiological assessment** (BrdU + cells, GFAP, DCX, Nestin, Tuj-1, BDNF)	THC ↑ BrdU + cells, GFAP, Nestin, DCX, Tuj-1 in all doses (peak at 1.5mg/kg); THC ↑ score in NOD.	THC ↑ proliferation and survival; THC ↑ behavior related to pattern separation.	(Suliman et al., 2018)
Behavioral assessment (FST) **Physiological assessment** (CB1r, DCX, Neurite Growth) **Electrophysiological assessment** (fPSP in PFC L2→L5; fPSP in DG MPP→DG)	THC ↑ score in FST; THC ↓ CB1r density, DCX+ cells, neurite growth; THC ↓ fPSP amp. in PFC L2→L5; THC ↓ fPSP slope in DG MPP→DG.	THC in adolescence ↓ survival in adulthood; THC in adolescence ↓ plasticity in adulthood.	(Cuccurazzu et al., 2018)
Physiological assessment (BrdU + cells)	No effect	Proliferation (=)	(Kochman et al., 2006)
Physiological assessment (CA1 histology, proteomics)	Neonatal THC 5mg/kg exposure for 12 days reduce number and diversity of neuronal types in CA1 of adult mice; THC exposure from P5-35 downregulate respiratory chain proteins in the mitochondrial complex in adulthood (P48 and P120);	Neonatal THC 5mg/kg exposure for 12 days reduce number and diversity of neuronal types in CA1 of adult mice; THC exposure from P5-35 downregulate respiratory chain proteins in the mitochondrial complex in adulthood (P48 and P120);	
Physiological assessment (BW, DA, BDNF, TrkB levels, SOX2, DCX) **Behavioral assessment** (OF,PPI,ASR,DT, OLT, MWM)	THC treatment ↑ locomotor activity; THC tratment did not change BW; THC treatment did not change PPI response THC treatment did not change DT tests performance; THC treatment reduce spatial memory efficiency on MWM but not learning of cognitive flexibility; THC treatment ↑ CB1R expression in PFC on the P90 day; THC treatment ↑ TrkB expression in Hipp on the P90 day; THC treatment ↓ BDNF in PFC and Hipp on the P90 day; THC treatment ↑ DOPAC/DA ratio in Hipp on the P90 day; THC treatment ↓ DOPAC/DA ratio on PFC on the P90 day;	THC treatment reduce spatial memory efficiency on MWM but not learning of cognitive flexibility; THC treatment ↓ BDNF in PFC and Hipp on the P90 day; THC treatment ↓ DCX+ and SOX2+ cells in DG on the P90 day.	

	THC treatment ↓ DA and HVA levels on Dorsal striatum on the P90 day; THC treatment ↑ DOPAC levels in the Nacc on the P90 day; THC treatment ↓ DCX+ and SOX2+ cells in DG on the P90 day.		
In vitro studies			
Physiological assessment (morphology, neurite growth; Rap1 and B-Raf activation)	AEA ↓ neurite growth via CB1 signaling, blocking even NGF neurite growth increase in all models used; AEA ↓ Rap1 and B-Raf activation of ERK signaling cascade through CB1R mediated effect.	AEA ↓ neurite growth via CB1 signaling, blocking even NGF neurite growth increase in all models used; AEA ↓ Rap1 and B-Raf activation of ERK signaling cascade through CB1R mediated effect.	(Rueda et al., 2002)
Physiological assessment (CBR expression levels, Map2, Tuj1, Sox1, Nestin, TH, DAT mRNA levels, Methylation rate for DAT, MAP2 and CB1R genes) **Electrophysiological assessment** (Passive and active membrane properties, mPSC)	smNPCs cells express a higher rate of CB1R than CB2R smNPCS; Chronic THC and AEA treatment all doses have no effect on Tuj1, Map2, TH, DAT and CB1R mRNA levels compared to control medium; Chronic 10 μM THC and AEA treatment ↓ INa Inward and IK outward currents; Chronic CB treatment ↓ the number of mEPSC; Chronic 1 μM AEA ↑ frequency of sEPSC.	smNPCs cells express a higher rate of CB1R than CB2R smNPCS; Chronic THC and AEA treatment all doses have no effect on Tuj1, Map2, TH, DAT and CB1R mRNA levels compared to control medium; Chronic 10 μM THC and AEA treatment ↓ INa Inward and IK outward currents; Chronic CB treatment ↓ the number of sEPSC; Chronic 1 μM AEA ↑ frequency of sEPSC.	(Stanslowsky et al., 2017)
Physiological assessment (CB1R expression and localization, Neurite outgrowth, Rho-A-B-C activity, ERK1 phosphorylation levels)	Win55212-2 at 20 μM induced cone growth retraction in mice cultured E13.5 GABAergic neurons, whereas CB1 antagonist induced cone growth attraction CB1 agonist treatment change the axonal cone growth angle away from cathode in galvanotropic embryonic spinal neurons of X. Laevis. CB1 and BDNF ↑ ERK phosphorylation AEA induce CB1R removal from filopodia ROCK blocker converted the Chemorepellent effect of CB1R agonist into Chemo attractive effect	Win55212-2 at 20 μM induced cone growth retraction in mice cultured E13.5 GABAergic neurons, whereas CB1 antagonist induced cone growth attraction CB1 agonist treatment change the axonal cone growth angle away from cathode in galvanotropic embryonic spinal neurons of X. Laevis. CB1 and BDNF ↑ ERK phosphorylation AEA induce CB1R removal from filopodia ROCK blocker converted the Chemorepellent effect of CB1R agonist into Chemo attractive effect	(Berghuis et al., 2007)
Physiological assessment (proliferation, cAMP levels, PI3K/Akt and ERK activation, TuJ1 expression)	48 hours exposure to HU210 to cultured hippocampal NS/PC cells ↑ proliferation via cAMP increase and ERK signaling; 8 days exposure to HU210 to cultured hippocampal NS/PC cells do not change TuJ1 expression patterns.	48 hours exposure to HU210 to cultured hippocampal NS/PC cells ↑ proliferation via cAMP increase and ERK signaling; 8 days exposure to HU210 to cultured hippocampal NS/PC cells do not change TuJ1 expression patterns.	(Jiang et al., 2005)
Physiological assessment (CBR expression, Forskolin stimulated cAMP accumulation, morphology)	N1E-115 cells express CB1 but not CB2 receptor; HU210 ↑ forskolin stimulated cAMP accumulation; HU210 ↓ neurite growth in N1E-115 in a dose-dependent manner; HU210 ↓ of neurite growth is mediated by cAMP increase.	HU210 ↓ neurite growth in N1E-115 in a dose-dependent manner; HU210 ↓ of neurite growth is mediated by cAMP increase.	(Zhou & Song, 2001)
Physiological assessment (Neurite growth, TRPV1 mediated Ca2+ influx)	HU210 ↑ neurite growth compared to control only when PC12 cells are raised in hyperglycemic medium; HU210 ↓ capsaicin Ca2+ influx.	HU210 ↑ Neurite growth in PC12 cells raised in hyperglycemic medium.	(F. Zhang et al., 2009)

Morphological assessment (central and peripheral Nervous System formation)	O-2545 gastrulation has strong teratogenic effect that escalates with dose.	O-2545 gastrulation has strong teratogenic effect that escalates with dose.	(Psychoyos et al., 2008)
Physiological assessment (JNK, Casp-3 activation)	THC ↑ JNK and casp-3 activation via CB1R in a dose-dependent manner	THC ↑ JNK and casp-3 activation via CB1R in a dose-dependent manner	(Downer et al., 2007)
Physiological assesment (CB1R expression and localization, Neurite outgrowth, ERK and akt phosphorylation)	hiPSC differentiated into neurons express CB1R, DAGLA, FAAH and MGLL; 1 μM 2-AG or 3 μM THC for 24 hrs ↓ neurite outgrowth in neurons derived from hiPSC without reducing total number of branches; 1 μM 2-AG or 3μM THC for 30 mins ↓ pERK in neurons derived from hiPSC; 3μM THC for 15 and 30 mins ↓ pAkt in neurons derived from hiPSC; control levels of pERK, pAkt and neurite growth are restored if CB1R antagonist is applied together;	1 μM 2-AG or 3 μM THC for 24 hours ↓ neurite outgrowth in neurons; control levels of pERK, pAkt and neurite growth are restored if CB1R antagonist is applied together	(Shum et al., 2020)
Physiological assessment (Neurite outgrowth, mitochondrial function)	THC ↓ cell growth of cortical neurons from mice cultured for 24h in a dose-dependent manner; THC at 7.5 μM disrupt mitochondrial membrane via CB1R activation and negatively impact mitochondrial function;	THC ↓ cell growth of cortical neurons from mice cultured for 24h in a dose-dependent manner;	(Beiersdorf et al., 2020)
Human studies			
Physiological assessment (CT scan of Hipp volume)	High THC/CBD ratio correlates with ↓ of Hipp volume (causality not assured)	High THC/CBD ratio ↓ Hipp volume	(Demirakca et al., 2011)

MWM (Morris Water Maze); FC (Fear Conditioning); BrdU (Bromodeoxyuridine); FST (Forced Swim Test); NFST (Novelty suppressed Feeding Test); NOD (Novel Object Discrimination); cort (corticosterone); NeuN (Hexaribonucleotide Binding Protein-3); DCX (doublecortin); Tuj-1 (Neuron-specific class III beta-tubulin); M-AEA (Methanamide); JNK (c-Jun N-terminal kinases); Casp-3 (caspase 3); fPSP (Field Post-Synaptic Potential); mEPSC (miniature spontaneous Excitatory Post Synaptic Currents); DG (Dentate Gyrus); VTA (Ventral Tegmental Area); Hipp (Hippocampus); MPP (Medial Perforant Path); Rap1 (Ras-proximate-1 or Ras-related protein 1); B-raf (serine/threonine-protein kinase B-Raf); ERK (extracellular signal-regulated kinase); PPI (Pre-pulse Inhibition); ASR(Acoustic Startle reflex); DT (Discrimination tests); OLT(Object location test); smNPCs (small molecule Neuronal Precursor Cells); Map2 (Microtubule-associated protein 2); sox1 (SRY-box transcription factor 1); Nestin (neuroepithelial stem cell protein); TH (Tyrosine Hydroxylase); DAT (Dopamine Transporter); Rho -A-B-C (Rho family of GTPAses Ras); PSA-NCAM (polysialiated neural cell adhesion molecule); ERK1 (extracellular signal-regulated kinases); ROCK (serine-threonine kinase Rho kinase); pCREB (phosphorylated cAMP response element-binding protein); syn1 (synapsin-1); syn3 (synapsin-3); PSD95 (Post-synaptic Density protein 95); NS/PC (Neural Stem Progenitor Cells); cAMP (cyclic Adenosine monophosphate); PI3K/Akt (phosphatidylinositol 3-kinases/Protein Kinase B); N1E-115 (Mouse Neuroblastoma cell line); TRPV1 (transient receptor potential cation channel subfamily V member 1); PC12 (Rat adrenal medulla pheochromocytoma derived cell line);hiPSC (human induced Pluripotent Steam Cell; DAGLA (Diacylglycerol lipase); FAAH (Fatty acid amide hydrolase); MGLL (Monoacylglycerol lipase); CT (Computerized tomography).

NMDA antagonists

Ketamine and phencyclidine (PCP), synthetic drugs originally discovered at the Parke-Davis Research Center, are arguably the most studied NMDA antagonists (Hoefle & Warner-Lambert, 2001; Pochwat et al., 2014). Initially employed as anesthetics in the late 20th century (Domino & Warner, 2010; Journey & Bentley, 2021), these drugs induce a dissociative state (Domino & Warner, 2010; Journey & Bentley, 2021) and can lead to side effects ranging from hallucinations to mania (Javitt & Zukin, 1991; Pochwat et al., 2014; Ruiz & Strain, 2011). The discovery of ketamine's rapid antidepressant effect (Berman et al., 2000) was considered the greatest breakthrough in depression research over the past half-century (Data-Franco et al., 2017; Hashimoto, 2019; Sial et al., 2020). Within our review, 20 of the included studies utilized NMDA antagonists, accounting for the second most studied chemical group of psychedelics. The time point of analysis differs between laboratories, but the single-dose paradigm, employed by 80% of ketamine studies, remains the most remarkable aspect of ketamine treatment (see descriptive results in **Supp. Table 2**). Ketamine's facilitatory effect on plasticity has been thoroughly examined through physiological and behavioral measures, and to a lesser degree, electrophysiological methods (**Table 2**). Across diverse protocols, the focus was primarily on elucidating the cellular pathway through which ketamine boosts Brain-Derived Neurotrophic Factor (BDNF), subsequently promoting synaptogenesis, neurite growth, and neurogenesis. Among the in vivo ketamine studies (n = 10), the most frequently used behavioral assessment was the forced swim test (FST; 40%), followed by the open field test (OF; 20%). Other tests, such as the novel object recognition test (NOR), novelty suppressed feeding test (NSFT), splashing test (ST), and Morris water maze (MWM), were each utilized in 10% of the studies. Broadly, these behavioral tests were deployed to evaluate cognitive abilities known to be impaired in major depressive disorders.

Ketamine has been shown to enhance synaptogenesis and neurite growth in young adult mice when administered at a subanesthetic dose ranging from 0.25-10 mg/kg/day (Akinfiresoye & Tizabi, 2013; N. Li et al., 2010; R.-J. Liu et al., 2013; Ly et al., 2018). However, three studies using doses beyond this range reported potentially harmful effects on neurogenesis and associated behaviors. First, Zuo and colleagues administered 30mg/kg/day of ketamine to male rats between postnatal day 35-42 (P35-P42), along with 2-4g/kg/day of alcohol over 14 days. They observed a reduction in synapsin expression in the hippocampus, a phenomenon that occurred even in the absence of alcohol. The study also reported that ketamine treatment in adolescent mice induced hyperlocomotion, anxiety (assessed using OF), ataxia, stereotyped behavior, mitochondrial damage, and morphological abnormalities in hippocampal and cortical cells. Additionally, dopamine (DA) and glutamate levels increased in the cortex and hippocampus, suggesting a possible disruption of the CREB pathway causing the aforementioned adverse effects (Zuo et al., 2018). Next, Huang et al. (2016) delivered four sub-anesthetic intraperitoneal (IP) injections of 10mg/kg ketamine over 4 hours to P7 rats. This regimen led to a reduction in neuronal and glial proliferation without impacting neuronal differentiation (as assessed by B-Tubulin III/BrdU+ co-labeling) up to 14 days post-injection. However, the levels of labeled cells in all assays returned to control levels 21 days post-injection (Huang et al., 2016). Lastly, Schiavone et al. (2020) elucidated the mechanisms underlying the neurochemical imbalance and oxidative stress observed in adulthood following intermittent administration of 30 mg/kg ketamine to neonatal rats (P7, P9, P11). A two-fold increase in NOX2 immediately preceding the chemical insult persisted into adulthood, resulting in other observed imbalances. Although the authors did not measure the number of neural progenitors in the SGZ, it's been found that NOX2 knockout mice exhibit a reduced number of radial glial-like (RGL) cells compared to wild-type mice in adulthood (Dickinson et al., 2011), indicating a possible bimodal, dose-dependent effect of ketamine on RGL proliferation during this specific ontogenetic window of exposure via NOX2 overexpression (Schiavone et al., 2020).

The prevailing consensus in the scientific literature suggests that ketamine positively influences the subprocesses of adult neurogenesis via the upregulation of BDNF (Browne & Lucki, 2013; Duncan et al., 2013; Haile et al., 2014; Keilhoff et al., 2004; Lepack et al., 2015; Rybakowski et al., 2013). However, a study by Goulart et al. (2010) presents an intriguing contrast. They observed a dose-dependent reduction in the BDNF increase stimulated by the Novel Object Recognition (NOR) task, assessed approximately 5 hours post-training/injections (Goulart et al., 2010). This seemingly contradictory finding may be partially explicated by the research conducted by Huang and colleagues. Although they did not measure neurotrophin levels, their

observations suggested a transient reduction in proliferation and differentiation, possibly mediated by BDNF (Huang et al., 2016). This could provide a plausible mechanism for the ketamine-induced disruption of the consolidation phase of long-term recognition memory observed by Goulart and colleagues. Interestingly, despite the disruption in the migration of adult-born Granule Cells (abGC), leading to an increased number of GCs in the SGZ until at least P44, the cognitive deficits persisted until at least P60 as assessed by Morris Water Maze (MWM) performance (Huang et al., 2016). Nevertheless, it is important to note that Huang et al.'s study involved a potent intoxication early in life (P7), and their results do not mirror those obtained in adult animals (Browne & Lucki, 2013; Data-Franco et al., 2017; Duman & Li, 2012; Duncan et al., 2013; Ly et al., 2018; Olson, 2018). This discrepancy reinforces the need for further research to elucidate the dynamic interactions between ketamine and neurotrophins under conditions that stimulate neurogenesis.

The majority of studies on phencyclidine (PCP) have sought to elucidate the neurocognitive damage caused by chronic exposure in utero or neonatally. All of these studies employed physiological assessments, and nearly half of them incorporated tests for behavioral evaluation (Tanimura et al., 2009; Toriumi et al., 2012). The Bromodeoxyuridine (BrdU) label for cells undergoing the S-phase was the most frequently used method (80%). The consensus from these studies is that early exposure to PCP has a significantly detrimental impact on neurogenesis (Maeda et al., 2007; Tanimura et al., 2009; Toriumi et al., 2012) and overall plasticity (Toriumi et al., 2012; Q. Zhang et al., 2016). In contrast, a study by Liu et al. (2006) in Japan reported different results. They administered 7.5 mg/kg PCP acutely and chronically (for either 5 or 14 days) for comparison within groups. They found that acute and 5-day chronic exposure resulted in minor or no effects on cell proliferation, differentiation, and survival, in the dentate gyrus of adult rats when compared to the saline control. Interestingly, the decrease induced by these treatments was restored to control levels after one week of abstinence (J. Liu et al., 2006). However, their study lack detailed information on the exact age of the animals, a factor that

strongly influences adult neurogenesis (Kuhn et al., 1996), only stating that was conducted on young adult animals, not neonatal or in utero subjects. In summary, it is reasonable to assert that exposure to any psychoactive substance during the early stages of neural development could induce detrimental effects. Moreover, the safety of PCP as a therapeutic agent needs further evaluation in adult models, and preferably in disease-relevant models, rather than focusing solely on its teratogenic effects.

Human trials with ketamine shed light on various processes contributing to plasticity. For instance, a cross-sectional study conducted by Fan and colleagues revealed that ketamine addiction can lead to low-grade inflammation (Fan et al., 2015). Further, three other studies specifically focused on patients with major depressive disorder (MDD), particularly treatment-resistant depression (TRD) (Duncan et al., 2013; Haile et al., 2014; Rybakowski et al., 2013). These studies found a correlation between non-responsiveness to ketamine treatment in TRD patients and a decrease in BDNF levels 7 and 14 days after a single 0.5mg/kg ketamine infusion (Rybakowski et al., 2013). nterestingly, this same dose, when applied in a similar protocol, led to an increase in serum BDNF levels as assessed 230 minutes after the infusion. This increase was accompanied by an increase in Slow Wave Sleep (SWS) duration in responders (individuals characterized by a decrease of more than 50% in their baseline Montgomery-Asberg Depression Rating Scale (MADRS) score (Duncan et al., 2013). A similar study conducted in Texas, USA, yielded a very similar results (Haile et al., 2014).

Moreover, in a separate human-controlled trial at the New York State Psychiatric Institute, ketamine administered to cocaine-addicted patients, who weren't seeking treatment or abstinence, reduced their cocaine self-consumption and had high scores on the Hood Mysticism Scale (HMS) when compared to other scales like the Clinician Administered Dissociative States Scale (CADSS) and the Near-Death Experience Scale (NDES). This provides some insight into which subjective effects of the ketamine psychedelic experience might be beneficial for addiction treatment, suggesting these aspects should be monitored in assisted psychotherapy strategies (Dakwar et al., 2018).

Table 2 Results reported by the reviewed NMDA antagonists' studies.

Outcome variable	Reported Results	Effect on Neurogenesis	Reference
In vivo studies			
Physiological assessment (Glu, Glutamine, GABA, and PV levels; NAc DA levels and DA turnover; 8-OHdG, NOX1, NOX2)	Ketamine ↑ PFC Glu and Glutamine in P11 mice; Ketamine ↑ PFC Glu and GABA levels in P70 mice; Ketamine ↓ PFC PV, NOX1 in Both ages; Ketamine ↓ Nac DA levels in Both ages; Ketamine ↑ NAc DA turnover at both ages; Ketamine ↑ Nac and PFC, 8-OHdG and NOX2 levels in both ages; Ketamine ↓ NAc NOX1 at P11; Ketamine ↑ NAc NOX1 at P70.	Ketamine ↑ Nac and PFC, 8-OHdG and NOX2 levels in both ages.	(Schiavone et al., 2020)
Behavioral assessment (OF) **Physiological assessment** (Cell morphology, Syn, Glu, DA, Cit-C, Casp3 and Creb pathway proteins levels)	Ketamine ↑ Locomotor activity, ataxia and stereotypy; Ketamine ↑ cell morphology abnormalities in CA1, CA2 and CA3; Ketamine ↓ Syn levels in Hipp; Ketamine ↑ DA, Glu, Cit-C and Casp3 activity and levels on Cortex and Hipp; Ketamine ↓ CREB pathway protein levels.	Ketamine ↓ Syn levels in Hipp; Ketamine ↓ CREB pathway protein levels	(Zuo et al., 2018)
Behavioral assessment (FST) **Physiological assessment** (mTOR and GSK-3β activity, synaptogenesis) **Electrophysiological assessment** (sEPSC)	Ket1/li10 ↑ mTOR signaling; ketamine 10mg/kg and ket/Li10 ↑ sEPSC frequency and synaptogenesis; Ketamine 10mg/kg and Ket/li1 ↓ score in FST.	Ketamine ↑ Neurite outgrowth	(R.-J. Liu et al., 2013)
Physiological assessment (BrdU+ cells)	Ketamine ↑ BrdU+ cells after 2 and 3 weeks.	Ketamine ↑ Proliferation; Ketamine ↑ survival.	(Keilhoff et al., 2004)
Behavioral assessment (NOR) **Physiological assessment** (BrdU+ cells, BDNF levels)	Ketamine 8, 20 mg/kg after training ↓ score in NOR; Ketamine ↓ BDNF levels increase after cognitive training in NOR.	Ketamine ↓ BDNF levels increase after cognitive training in NOR	(Goulart et al., 2010)
Behavioral assessment (FST, NSFT) **Physiological assessment** (mTOR signaling, Neurite outgrowth) **Electrophysiological assessment** (sEPSC)	Ketamine 5, 10 mg/kg ↑ mTOR signaling in PFC until 1hr after injection; Ketamine 10 mg/kg ↑ Synaptogenesis and sEPSC in PFC; Ketamine increases in mTOR signaling and Synaptogenesis is Abolished by NBQX or ERK blocker pre-treatment; Ketamine ↓ score in FST and NFST; Ketamine reduction in FST and NFST are abolished by Rapamycin, and ERK blocker pre-treatment.	Ketamine ↑ synaptogenesis and sEPSC in PFC; Ketamine increase in mTOR signaling and Synaptogenesis is Abolished by NBQX or ERK blocker pre-treatment; Ketamine ↓ score in FST/ NFST.	(N. Li et al., 2010)
Behavioral assessment (OF, FST, SPT) **Physiological assessment** (BDNF, Syn and d-mTOR levels)	Ket 0.5mg/kg and Ket/AMPA 0.25, 0.5mg/kg ↓ FST score; ket/AMPA ↑ SPT score; ket 0.5 and ket/AMPA 0.25,0.5mg/kg ↑ BDNF, Syn and d-mTOR levels.	ket 0.5 and ket/AMPA 0.25,0.5mg/kg ↑ BDNF, Syn and d-mTOR levels	(Akinfiresoye & Tizabi, 2013)
Behavioral assessment (FST) **Physiological assessment** (BDNF Levels)	mPFC BDNF block prevent FST score reduction by ketamine; L-Type channel block prevents FST score reduction by ketamine.	mPFC BDNF block prevent FST score reduction by ketamine	(Lepack et al., 2015)

Behavioral assessment (MWM) **Physiological assessment** (BrdU+ cells)	Ketamine ↓ BrdU+ cells in proliferation essay; Ketamine ↑ BrdU+ neurons in differentiation essay; Ketamine ↓ BrdU+ astrocytes in differentiation essay; Ketamine did not change the apoptosis rate in the DG; Ketamine dampens the migration of Newborn Granule cells to outer GCLs; Ketamine dampens RGL growth; P7 exposure to Ketamine ↓ score in MWM.	Ketamine ↓ BrdU+ cells in proliferation essay; Ketamine ↑ BrdU+ neurons in differentiation essay; Ketamine ↓ BrdU+ astrocytes in differentiation essay; Ketamine did not change the apoptosis rate in the DG; Ketamine dampens the migration of Newborn Granule cells to outer GCLs; Ketamine dampens RGL growth; P7 exposure to Ketamine ↓ score in MWM.	(Huang et al., 2016)
Physiological assessment (Neurite outgrowth, Spine density)	Ketamine ↑ spinogenesis in rats Pyr PFC neurons.	Ketamine ↑ spinogenesis in rats Pyr PFC neurons.	(Ly et al., 2018)
Morphological assessment (BW) **Physiological assessment** (PFC cell density and number, PFC cell migration, PFC, PCNA+, PAG+, Pax6+, Tbr2+, BrdU+ cells, expression levels of Notch2 and Ntn1) **Behavioral assessment** (NOR, FST, PPI)**.**	PCP ↓ BW on P1; PCP ↓ NOR and PPI score; PCP ↑ FST score; PCP ↓ PFC cell density and PAG+ cells on P56 mice at PFC; PCP ↓ BrdU+ cells between E12-E13 and on P7 mice at PFC; PCP ↓ PCNA+, Pax6+, Tbr2+ cells on E13.5 mice; PCP ↓ expression levels of Notch2 and ntn1.	PCP ↑ FST score; PCP ↓ NOR and PPI score; PCP ↓ PFC cell density and PAG+ cells on P56 mice at PFC; PCP ↓ BrdU+ cells between E12-E13 and on P7 mice at PFC; PCP ↓ PCNA+, Pax6+, Tbr2+ cells on E13.5 mice; PCP ↓ expression levels of Notch2 and ntn1, disrupting proliferation	(Toriumi et al., 2012)
Physiological assessment (BrdU+ cells)	Chronic PCP ↓ BrdU + cells; Other treatments has no effects over the number of BrdU+ cells in any other paradigm.	Chronic PCP ↓ BrdU + cells; Other treatments has no effects over the number of BrdU+ cells; Chronic PCP did not alter survival or differentiation of abGC.	(J. Liu et al., 2006)
Behavioral assessment (locomotion) **Physiological assessment** (BrdU+ cells)	PCP ↑ BrdU+ cells at P21 and P56; PCP ↓ Locomotor activity.	PCP ↑ Survival; PCP ↑ Proliferation.	(Tanimura et al., 2009)
Physiological assessment (BrdU+ cells)	PCP 3, 10 mg/kg ↓ BrdU+ cells in proliferation and Survival essays	PCP 3, 10 mg/kg ↓ BrdU+ cells in proliferation and Survival essays	(Maeda et al., 2007)
In vitro **studies**			

Physiological assessment (Neurite growth; Synaptophysin, PSD95 and BDNF levels; Akt and GSK3β phosphorylation levels; NRG1 activation levels)	PCP ↓ neurite growth PCP ↓ Synaptophysin and PSD95 levels PCP ↓ phosphorylation of Akt and GSK3β without affecting the expression levels PCP reduction of neurite growth and Akt GSK3β phosphorylation is dependent of NRG1 activation PCP ↓ BDNF levels, and this reduction is NRG1 activation dependent PCP ↓ NRG1 expression levels	PCP ↓ neurite Growth; PCP reduction of neurite growth and Akt GSK3β phosphorylation is dependent of NRG1 activation; PCP ↓ BDNF levels, and this reduction is NRG1 activation dependent.	(Q. Zhang et al., 2016)
Human studies			
Survey assessment (PANSS) **Physiological assessment** (TNF-α, IL-6 and IL-18)	Ketamine addicted has increased IL-6, IL-8 and PANSS score; Ketamine addicted has reduced TNF-α levels	Ketamine addiction produce chronic low-grade inflammation	(Fan et al., 2015)
Survey assessment (Subjective mystic effects of ketamine, cocaine craving and self-administration self-report)	HMS correlates with Overall improvement Ketamine treatment reduced cocaine self-administration	Unclear	(Dakwar et al., 2018)
Survey assessment (HDRS) **Physiological assessment** (serum BDNF, NGF, NTF3, NTF4 and GDNF)	BDNF levels in non-responders was reduced 7 and 14 days after treatment A single dose of ketamine reduced score in HDRS in all patients 14 days after treatment 48% of patients had a HDRS score ≤ 7 points 14 days after treatment	Non-responsiveness to ketamine treatment positively correlates with low BDNF levels 7 and 14 days after treatment	(Rybakowski et al., 2013)
Survey assessment (MADRS) **Physiological assessment** (Serum BDNF) **Electrophysiological assessment** (SWA)	Single ketamine dose ↓ MADRS score; Single ketamine dose ↑ Serum BDNF levels 230 minutes after; Single ketamine dose ↑ total sleep time, SWS and REM; SWA increase and BDNF increase positively correlates in responders.	Single dose of ketamine ↑ BDNF serum levels in responders	(Duncan et al., 2013)
Survey assessment (MADRS) **Physiological assessment** (Serum BDNF)	Single ketamine dose ↓ MADRS score in ~50% of patients; BDNF increase negatively correlates with MADRS score; Single dose of ketamine ↑ BDNF serum levels in responders.	Single dose of ketamine ↑ BDNF serum levels in responders	(Haile et al., 2014)

Nac (Nucleus accumbens); PFC (Prefrontal Cortex); Glu (Glutamate); DA (Dopamine); PV (Parvalbumin); 8-OHdG (8-Oxo-2'-deoxyguanosine); NOX1 (NADPH oxidase 1); NOX2 (NADPH oxidase 2); syn (synapsin); OF (Open Field Test); FST (Forced Swim Test); cit-c (cytochrome complex); Casp3 (Caspase 3); Creb (cAMP response element-binding protein); mTor (mechanistic target of rapamycin); GSK-3β (Glycogen synthase kinase 3 beta); sEPSC (spontaneous Excitatory Post-synaptic Current); pyr (pyramidal neuron); abGC (adult born Granule Cells); GCL (Granule Cell Layer); RGL (Radial-Glia Like Cells); NBQX (2,3-dioxo-6-nitro-7-sulfamoyl-benzo[f]quinoxaline); ERK (extracellular signal-regulated kinase); BrdU (Bromodeoxyuridine); NOR (Novel Object Recognition Test); BDNF (Brain Derived Neurotrophic Factor); MWM (Morris Water Maze Test); BW (Body Weight); PCNA (Proliferating cell nuclear antigen); PAG (Phosphate activated Glutaminase); Pax6 (Paired box protein Pax-6); Tbr2 (T-box brain protein 2); Notch2 (Neurogenic locus notch homolog protein 2); Ntn1 (Netrin-1); PPI (Pre-Pulse inhibition Test); PSD95 (Post-synaptic Density Protein 95); Akt (Protein Kinase B); NRG1 (Neuregulin 1); PANSS (Positive and Negative Syndrome Scale); TNF-α (Tumor necrosis factor-α); IL-6 (Interleukin 6); IL -18 (interferon-gamma inducing factor); HMS (Hood Mysticism Scale); CADSS (Clinician Administered Dissociative States Scale); NDES (Near-Death Experience Scale); HDRS (Hamilton Depression Rating Scale); NGF (Nerve Growth Factor); NTF3 (neurotrophin-3); NTF4 (neurotrophin-4); GDNF (glial- derived neurotrophic factor); SWA (Slow-Wave activity); MADRS (Montgomery–Åsberg Depression Rating Scale).

Harmala alkaloids

Harmala alkaloids are indole nitrogenated compounds featuring a β-carboline heterocyclic structure, first identified in the seeds of *Peganum harmala*, also known as Syrian Rue (Herraiz et al., 2010). Recently, these compounds have attracted increased attention due to their presence in *ayahuasca* brews. The primary compounds in this category include harman, harmine, harmaline, and tetrahydroharmine (de Oliveira Silveira et al., 2020). While these β-carbolines are not inherently psychedelic, as they don't bind to 5-HT receptors, they can influence serotonin signaling by inhibiting the Monoamine oxidase enzyme (MAO), thus producing behavioral effects somewhat similar to Selective Serotonin Reuptake Inhibitors (SSRIs) (Domínguez-Clavé et al., 2016). Often referred to as MAO inhibitors (MAOis), these compounds are ubiquitous in the various forms of *ayahuasca* prepared by indigenous populations (McKenna et al., 1984). Interestingly, in some versions of *ayahuasca*, DMT is even absent (Callaway, 2005; Rivier & Lindgren, 1972). Their antidepressant effect has been explored in psychiatry, with some MAOis even being marketed under names like Neuralex and Marplan, among other synthetic functional analogs (Fagervall & Ross, 1986; Robinson, 1973). However, most MAOis were withdrawn from the market due to hepatotoxicity (Lopez-Munoz & Alamo, 2009), although it may not be the case of the harmala alkaloids (Brito-da-Costa et al., 2020). The cellular mechanisms of such compounds are still being explored, more recently under the light of the neurogenic hypothesis for MDD, as their effects on neurogenesis and plasticity processes are uncovered (Ferraz et al., 2019).

All studies involving harmala alkaloids (6 included, **Supp. Table 3**) focus on their potential antidepressant effects. *In vivo* studies also assessed hippocampal BDNF in conjunction with behavioral methods. Half of articles from this category originated from a Brazilian research team led by João Quevedo, with Jucélia Fortunato as a key contributor, which conducted a series of rat experiments revealing the potential antidepressant effects of the harmine alkaloid (Fortunato et al., 2009, 2010a, 2010b). The team initially found that male rats at post-natal day 60 (P60) showed improved performance in the forced swim test (a measure of depressive-like behavior) after being treated with 10 or 15mg/kg of harmine one hour prior to the test. Their locomotor activity (measured by open field test) remained normal. The improvement was comparable to that produced by the tricyclic antidepressant imipramine.

Intriguingly, rats treated with 15 mg/kg of harmine demonstrated a two-fold increase in BDNF levels immediately after the test, while those treated with imipramine did not (Fortunato et al., 2009). The following year, the team reported similar results after chronic administration (14 days) of harmine (Fortunato et al., 2010a).

A critique of their initial studies pertained to the relevance of the results to depression pathophysiology. To address this, they later generated a study incorporating a Chronic Mild Stress (CMS) model of depression in rats. There, they administered 15 mg/kg of harmine over a 7-day treatment period, using a similar protocol but without the imipramine control. Similar results were found. Moreover, the forced swim test (FST) assessment was replaced with the sucrose preference test (SPT) to measure anhedonia-like phenotypes, a core symptom of depression. The study demonstrated that harmine-treated mice exhibited less anhedonia only after CMS. The study also found that harmine treatment reduced blood levels of adrenocorticotropin (ACTH), a stress hormone, while adrenal weight was similar to that of non-CMS rats. Contrary to their previous findings, BDNF levels after treatment were not affected. Such discrepancy could be due to the different time-points of sample collection employed in the study (first after a stressful situation, second after a mildly pleasant situation), suggesting that harmine may dynamically influence BDNF levels (Fortunato et al., 2010b).

Harmine was also studied in Jiangsu, China, by a team led by Professor Chao Huang and Liu and colleagues, who used Chronic Unpredictable Stress (CUS) to model depressive symptoms in mice. They examined the effects of chronic administration of 10 and 20 mg/kg of harmine using the forced swim test (FST), sucrose preference test (SPT), and tail suspension test (TST) — the three most commonly used tests for measuring depressive symptoms in mouse models. It was reported a 40% effect favoring the antidepressant action of harmine in all behavioral tests, which was comparable to the positive control, fluoxetine (F. Liu et al., 2017). In addition to behavioral tests, biochemical markers were investigated. The team described that 20 mg/kg of harmine increased BDNF levels in the hippocampus and prefrontal cortex, along with glutamate transporter 1 (GLT-1) expression levels. They also found that harmine

promoted the survival of newborn cells, as evidenced by doublecortin (DCX) immunofluorescence, and prevented the reduction in glial fibrillary acidic protein (GFAP) induced by the CUS protocol (F. Liu et al., 2017). Lastly, the authors tested the hypothesis that these regulatory effects might be due to the restoration of astrocytic function by administering L-Alpha-Aminoadipic Acid (L-AAA), an astrocyte-specific gliotoxin, intracerebroventricularly (i.c.v). Indeed, this treatment shifted the results towards those of the saline-CUS group (F. Liu et al., 2017).

Two other studies conducted entirely in vitro reported that harmine increased cell proliferation. One study from Rio de Janeiro, Brazil, showed that the presence of 7.5 µM harmine in the culture medium of human neural progenitor cells (hNPCs) increased proliferation by 71.5% (Dakic, de Moraes Maciel, et al., 2016). A Spanish team went further and tested four different harmala alkaloids (harman, harmine, tetrahydroharmine, and harmol) at a 1 µM concentration, acutely (3 days) and chronically (7 days), on neurospheres derived from the subgranular zone (SGZ) and subventricular zone (SVZ) of adult mice. The treatment enhanced all stages related to neurogenesis in vitro: proliferation, migration, and differentiation (Morales-García et al., 2017). Altogether, the current body of research suggests that harmala alkaloids have a positive impact on neurogenesis and neuroplasticity, particularly within the framework of depression models (**Table 3**).

The antidepressant potential of MAO inhibitors (MAOis) is well known and has been explored in the past. However, more research is needed regarding the side effects of harmala alkaloids (Wimbiscus et al., 2010). Future studies should focus on expanding our understanding of the long-term safety of these compounds, comparing them with synthetic MAOis, preferably those still on the market. This research will help determine whether their positive effects can counterbalance potential hepatotoxicity and severe interactions with tyramine-rich foods. By identifying which part of the harmala alkaloid structure is responsible for differences in action and consequences, we can pave the way for the development of safer analogs.

Table 3 Results reported by the reviewed Harmala alkaloids studies

Outcome variable	Reported Results	Effect on Neurogenesis	Reference
In vivo			
Behavioral assessment (FST; OF) **Physiological** (Hipp BDNF)	10 and 15 mg/kg Harmine ↓ score in FST; 15mg/kg Harmine ↑ Hipp BDNF.	Harmine ↑ hipp BDNF.	(Fortunato et al., 2009)
Behavioral assessment (FST; OF) **Physiological** (Hipp BDNF)**.**	Harmine ↓ score in FST; Harmine ↑ Hipp BDNF.	Harmine ↑ hipp BDNF.	(Fortunato et al., 2010a)
Behavioral assessment (SPT, crossing and rearings on OFT) **Physiological assessment** (Adrenal Weight, blood ACTH and Hipp BDNF)	Harmine ↑ SPT score; Harmine ↓ Adrenal Weight; Harmine ↓ blood ACTH; Harmine ↓ hipp BDNF increase through CMS model.	Harmine ↓ hipp BDNF increase through CMS model.	(Fortunato et al., 2010b)
Behavioral assessment (TST; FST; SPT) **Physiological assessment** (Hipp and PFC BDNF; GLT-1; GFAP)	10 and 20 mg/kg Harmine ↓score in TST, FST; Harmine ↑ SPT; 20mg/kg harmine ↑ hipp and PFC BDNF, DCX+ cells, GLT-1, GFAP.	Harmine ↑Hipp and PFC BDNF; Harmine ↑DCX+; Harmine ↑GLT-1.	(F. Liu et al., 2017)
In vitro **studies**			
Physiological assessment (EdU+, Nestin, GFAP cells)	Harmine ↑ proliferation and neuronal differentiation.	Harmine ↑ Proliferation and differentiation of hNPCs.	(Dakic, Maciel, et al., 2016)
Physiological assessment (Musashi1, Nestin, Sox-2, Tuj-1, MAP-2, CNPase, GFAP levels, Ki67 +, DCX+ Cells)	Chronic B-Carbolines ↓ Musashi-1, Sox-2, Nestin levels in both NPC populations; Chronic B-Carbolines ↑ # and size of adult neurospheres in both NPC populations; Chronic B-Carbolines ↑ ki67+ cells and PCNA levels; Acute B-Carbolines ↑ Tuj1 and MAP-2 levels in both NPC; populations, with exception to Harmine at SGZ for Tuj1 ; Chronic Harmol and Harmaline ↑CNPase levels at SVZ population; Chronic B-Carboline ↑ GFAP in both NPC populations, with exception to Harmine at SGZ.	B-Carbolines ↑ Proliferation, migration and differentiation into adult neurons.	(Morales-García et al., 2017)

FST (Forced Swim Test); OF (Open Field Test); BDNF (Brain Derived Neurotrophic Factor); SPT (Sucrose Preference Test); TST (Tail Suspension Test); PFC (Pre Frontal Cortex); Hipp (Hippocampus); EdU (5-Ethynyl-2'-deoxyuridine); GFAP (Glial Fibrillary Acidic Protein) ; Nestin (Neuroepithelial Stem Cell protein); Sox-2 (sex determining region Y-box 2); Tuj-1 (Neuron-specific class III beta-tubulin); Map-2 (Microtubule-associated protein 2); CNPase (2',3'-Cyclic-nucleotide 3'-phosphodiesterase) ; Ki67 (Marker Of Proliferation Ki-67); SVZ (Subventricular Zone); NPC (Neural Progenitor Cell); PCNA (Proliferating cell nuclear antigen); CMS (Chronic Mild Stress); DCX (Doublecortin); SGZ (Subgranular Zone); GLT-1 (glutamate transporter 1);   hNPCs (human Neural Progenitor Cells)

Psychoactive tryptamines

Tryptamines are indolamines neurotransmitters that originate from tryptophan (Araújo et al., 2015). All psychoactive tryptamines are potent 5-HT2A agonists, and many also demonstrate affinity for 5-HT1 and other 5-HT2 receptors. Notably, they may also interact with ionotropic and metabotropic glutamate receptors, dopamine, acetylcholine, and trace amine-associated receptor (TAAR) (Carbonaro & Gatch, 2016). Certain DMT analogs have shown activity via sigma-1 receptors as well (Koornneef et al., 2009). This broad receptor affinity equips psychoactive tryptamines with the capability to induce altered states of consciousness, characterized by heightened introspection (de Araujo et al., 2012) and changes in sensory perception, mood, and thought (Calvey & Howells, 2018).

While psychoactive tryptamines, used by ancient civilizations globally for millennia (Desmarchelier et al., 1996), were first clinically investigated in the mid-20th century for the treatment of mood and drug abuse disorders, and some advocates proposed their systematic use by healthy individuals to enhance cognition (Nichols, 2016; Williams et al., 2002). Despite their shared primary pharmacologic pathway—5-HT2A receptor agonism—these tryptamines can generate different experiences due to their variable affinity for other receptors. The current review includes eleven studies that meet the inclusion criteria, investigating the effects of psilocybin (18%), ibogaine (9%), and DMT analogs, including the ayahuasca concoction (73%). These tryptamines were predominantly assessed using physiological and behavioral methodologies, with electrophysiological methods used less frequently (**Supp. Table 4**).

As shown in **Table 4**, the majority of the included studies provide evidence that, irrespective of their specific type, tryptamines have the potential to enhance aspects of neuroplasticity (Catlow et al., 2013; Dunlap et al., 2020; Katchborian-Neto et al., 2020; Lima da Cruz et al., 2018; Ly et al., 2018; Marton et al., 2019; Morales-Garcia et al., 2020). Remaining studies, although not directly assessing neurogenesis, reported no adverse effects on behavior or cellular physiology (Castro-Neto et al., 2013; Wankhar et al., 2020; Winne et al., 2020). For instance, Catlow et al. (2013) discovered that a single low dose (0.1-0.5 mg/kg) of psilocybin, administered 24 hours prior to a Trace Fear Conditioning test, initially heightened fear response but facilitated fear extinction by the third trial, an effect not observed until the tenth trial in control animals. Conversely, high doses (1-1.5 mg/kg) showed no behavioral impact and reduced BrdU-labeled cells 18 days post-treatment. (Catlow et al., 2013).

Another team, based in Shillong, India, investigated the acute and sub-acute (7 days) effects of 0.7 mg/kg 1-Methylpsilocin, a psilocin tryptamine derivative. Their research aimed to assess the potential of 5-HT2CR exploitation in mitigating symptoms common to mood disorders. Following a Chronic Unpredictable Stress (CUS) protocol in rats, they employed Open Field (OF), Forced Swim Test (FST), and Elevated Plus Maze (EPM) tests in conjunction with tissue sample analysis. Their findings indicated that 1-Methylpsilocin alleviated behavioral signs of anxiety and anhedonic phenotype by the seventh day post-CUS. Furthermore, they demonstrated that this treatment could rescue subjects from CUS-induced mitochondrial damage and imbalanced 5-HT/5-HT2CR levels in the PFC and hippocampus. Remarkably, the treatment also mitigated hippocampal cell morphological abnormalities and the CUS-induced rise in corticosterone levels (Wankhar et al., 2020).

Ibogaine, a naturally occurring psychoactive tryptamine found in certain plants of the Apocynaceae family native to Central Africa, has been investigated for its potential to mitigate drug-seeking behavior (Alper et al., 2008). Researchers in Uruguay examined the compound's influence on neuroplasticity in mesocorticolimbic areas of the rat brain, uncovering intricate interactions between NGF, GDNF, and BDNF, both in terms of mRNA and protein products. Briefly, 20 or 40 mg/kg doses of ibogaine (I_{20} and I_{40}) dose-dependently upregulated BDNF and NGF mRNA in PFC, Ventral Tegmental Area (VTA), Substantia Nigra (SN), and Nucleus Accumbens (Nacc) 24 hours post-injection. However, 3 hours post-injection, a decline of 1.7 and 2-fold in PFC-BDNF mRNA was observed for the I_{20} and I_{40} groups, respectively. Notably, I_{40} also elevated GDNF levels in VTA and SN 24 hours later. In terms of protein levels, upregulation was confined to proBDNF-NAcc (for both I_{20} and I_{40}) and VTA-GDNF exclusively for I40, with a possible tendency for increased BDNF in NAcc and VTA that warrants further investigation (Marton et al., 2019).

The indole alkaloid 5-Methoxy-N, N-dimethyltryptamine (5-MeO-DMT), a structural analog of DMT found in the venom of the *Incilius alvarius* toad, has been the subject of several studies (Araújo et al., 2015). In a series of three experiments conducted by our team, we investigated the

effects of 5-MeO-DMT on adult neurogenesis. Our first two experiments showed an increase in the proliferation and survival of newborn granule cells in the ventral dentate gyrus following a single intracerebroventricular injection of 100µg of 5-MeO-DMT. In the third experiment, we employed the whole cell patch clamp technique to measure the passive and active membrane properties of the identified doublecortin-positive cells. The results revealed that the adult-born granule cells (abGCs) from treated animals exhibited electrophysiological properties more akin to mature granule cells. In a separate study, Winne et al. examined salicylate-induced anxiety-like behavior. Their findings suggested that salicylate can induce theta-2 and slow gamma oscillations in young mice, correlating with anxiogenic behavior. This effect was mitigated by a single 20 mg/kg intraperitoneal injection of 5-MeO-DMT administered a week prior to the test (Winne et al., 2020).

Moreover, researchers from the University of São Paulo, Brazil, conducted a study on adult rats to assess changes in neurotransmitter levels 40 minutes post-ingestion of *ayahuasca*. Their findings indicated that all tested ayahuasca concentrations (250, 500, 800 mg/kg oral) elevated GABA and 5-HT levels in the hippocampus and reduced glycine and GABA levels in the amygdala, following a dose-dependent pattern. Moreover, levels of all monoamines in the amygdala increased, yet the rate of serotonin consumption was reduced in the amygdala. Only the highest dose of 800 mg/kg could reproduce this effect in the hippocampus (Castro-Neto et al., 2013). Regarding DMT analogs, noteworthy efforts are being made by David E. Olson's lab at UC Davis to develop psychedelics without hallucinogenic properties while maintaining their neuroplastic effects, thus potentially reducing undesirable effects for some patients. Despite certain criticisms, Dunlap and colleagues reported successful results, demonstrating that isoDMT and 5-MeO-isoDMT enhanced neurite growth in cultured cortical neurons, much like their natural counterparts. Interestingly, in further experiments, these compounds did not elicit the head-twitch response (HTR) - a behavior commonly observed in mice when a hallucinogenic 5-HT2A agonist is administered (Dunlap et al., 2020).

Future research exploring psychoactive tryptamines as potential therapeutic tools should focus on the modification of these molecules' structure (Structure-Activity Relationship - SAR), administration methods, and interactions with other drugs. This can be achieved through two main strategies: (1) eliminating the hallucinogenic properties, as exemplified by the work of Olson and his team at UC Davis, who are developing psychotropic drugs that retain their overall benefits for mental health while minimizing subjective effects; and (2) reducing the duration of the psychedelic experience to improve treatment readiness, reduce costs, and broaden patient accessibility. The outcome of these strategies would enable the administration of tryptamines without the requirement for patients to be monitored by healthcare professionals during the active hours of the drug experience.

An alternative route of treatment employing these compounds is currently being explored by syncretic practices like *União do Vegetal* (UDV) and *Santo Daime* in Brazil, among others globally (Labate and Cavnar, 2014; Svobodny, 2014). These groups administer the drugs in a traditional setting that merges Amerindian practices, Christianity, and (pseudo)scientific principles. Despite their obvious limitations, through these settings we are gaining insights into the drug's effects on individuals with diverse backgrounds, serving as a prototype for psychedelic-assisted psychotherapy. In this context, it is believed that the hallucinogenic properties of the drugs are not only desirable but also necessary to aid individuals in dealing with their traumas and behaviors by reshaping their consciousness with the assistance of experienced staff. However, this approach has been the subject of criticism globally as the number of fatal accidents begins to rise (Hearn, 2013; Holman, 2010), as practitioners are increasingly ill-equipped to handle the growing complexity of mental health issues presented by individuals seeking their services.

Table 4 Results reported by the reviewed psychoactive tryptamines studies

Outcome variable	Reported Results	Effect on Neurogenesis	Reference
In vivo studies			
Physiological assessment (Neurite outgrowth, Spine density) **Electrophysiological assessment** (sEPSC)	DOI, LSD ↑ Neurite outgrowth in Drosophila and Rats; DMT and ketamine ↑ Spine density in Pyr PFC neurons; DMT ↑ sEPSC frequency and amplitude in Pyr PFC neurons.	DOI, LSD ↑ Neurite outgrowth in Drosophila and Rats; DMT and ketamine ↑ Spine density in Pyr PFC neurons; DMT ↑ sEPSC frequency and amplitude in Pyr PFC neurons.	(Ly et al., 2018)
Physiological assessment (BrdU+ cells, DCX+ cells, Neurite outgrowth) **Electrophysiological assessment** (Passive and active membrane properties)	5-MeO-DMT ↑ BrdU+ cells in vDG; 5-MeO-DMT ↑ DCX+ neurons in vDG; 5-MeO-DMT ↓ AHP and AP threshold in young GC on vDG; 5-MeO-DMT ↑ maturation rate of young GC on vDG; 5-MeO-DMT ↑ Nº of branches on young GC at vDG.	5-MeO-DMT ↑ BrdU+ cells in vDG; 5-MeO-DMT ↑ DCX+ neurons in vDG; 5-MeO-DMT ↓ AHP and AP threshold in young GC on vDG; 5-MeO-DMT ↑ maturation rate of young GC on vDG; 5-MeO-DMT ↑ Nº of branches on young GC at vDG.	(Lima da Cruz et al., 2018)
Behavioral assessment (OF) **Electrophysiological assessment** (LFP)	5-MeO-DMT ↓ anxiety-like behavior and Electrophysiological signature of anxiety.	5-MeO-DMT reduces anxiety similar to anti-depressant that increase neurogenesis	(Winne et al., 2020)
Behavioral assessment (OF) **Physiological assessment** (Nac, PFC, VTA and SN mRNA and protein levels for GDNF, BDNF and NGF)	Ibogaine 40mg/kg ↑ GDNF mRNA in VTA and SN 24h after injection; Ibogaine 20, 40 mg/kg ↑ BDNF mRNA in all areas 24h after injection; Ibogaine 20,40 mg/kg ↓ BDNF mRNA in PFC 3h after injection; Ibogaine 40mg/kg ↑ NGF mRNA in all areas 24h after injection; Ibogaine 20mg/kg ↑ NGF mRNA in PFC and VTA 24h after injection; Ibogaine 40 mg/kg ↑ GDNF protein levels in VTA 24h after injection; Ibogaine 20, 40 mg/kg ↑ proBDNF protein levels in Nac 24h after injection.	Ibogaine 20, 40 mg/kg ↑ BDNF mRNA in all areas 24h after injection; Ibogaine 20,40 mg/kg ↓ BDNF mRNA in PFC 3h after injection; Ibogaine 40mg/kg ↑ NGF mRNA in all areas 24h after injection; Ibogaine 20mg/kg ↑ NGF mRNA in PFC and VTA 24h after injection; Ibogaine 20, 40 mg/kg ↑ proBDNF protein levels in Nac 24h; after injections do not increase BDNF protein levels in any area 24h after injection.	(Marton et al., 2019)
Behavioral assessment (OF, FST, EPM) **Physiological assessment** (5-HT levels, TPH-2, $Ca^2ATPase$ and ETC-1 activity, 5-HT2C-R expression, Cell morphology)	MP ↑ 5-HT levels in Hipp and PFC in the CUS model but not in healthy animals; MP ↑ TPH-2 activity in Hipp in the CUS model but not in healthy animals; MP ↑ 5-HT2c-R expression in PFC on CUS and healthy animals; MP ↑ 5-HT2c-R expression in Hipp on CUS but not on healthy animals; MP ↑ $Ca^2ATPase$ activity in Hipp and PFC in the CUS model;	MP ↑ 5-HT levels in Hipp and PFC in the CUS model but not in healthy animals; MP ↑ TPH-2 activity in Hipp in the CUS model but not in healthy animals; MP ↑ 5-HT2c-R expression in PFC on CUS and healthy animals; MP ↑ 5-HT2c-R expression in Hipp on CUS but not on healthy animals; MP ↑ Ca2ATPase activity in Hipp and PFC in the CUS model;	(Wankhar et al., 2020)

	MP rescue ETC-1 downregulation induced by CUS back to control levels; MP reduce anxiety levels and anhedonical phenotype in CUS animals; MP reduce cell morphological abnormalities induced by CUS in Hipp and PFC.	MP rescue ETC-1 downregulation induced by CUS back to control levels; MP reduce anxiety levels and anhedonical phenotype in CUS animals; MP reduce cell morphological abnormalities induced by CUS in Hipp and PFC.	
Behavioral assessment (OF) **Physiological assessment** (BrdU+ cells) **Electrophysiological assessment** (Active membrane properties)	BW723C86 ↑ firing rate on pet1+ DRN ex vivo; BW723C86 prevent 5-HT1A agonist induced inhibition of pet1+DRN in vivo; MDMA fail to ↑ locomotor activity on 5-HT2B-KO; DOI 5mg/kg fail to ↑ HTR in 5-HT2B-KO; Fluoxetine fails to ↑ BrdU+ cells in 5-HT2B-KO.	BW723C86 ↑ firing rate on pet1+ DRN ex vivo; BW723C86 prevent 5-HT1A agonist induced inhibition of pet1+DRN in vivo; DOI fail to ↑ HTR in 5-HT2B-KO; Fluoxetine fails to ↑ BrdU+ cells in 5-HT2B-KO.	(Belmer et al., 2018)
Behavioral assessment (TFC) **Physiological assessment** (BrdU+ cells)	Psilocybin do not change TFC score for acquisition; Psilocybin 0.1,0.5 mg/kg ↑ fear response on trial 1 of TFC for extinction; Psylocibin ↓ fear response on trial 10 of TFC for extinction; Psylocibin 1mg/kg ↓ BrdU+ neurons in DG; NBOMe ↓ BrdU+ neurons in DG.	Psilocybin do not change TFC score for acquisition; Psilocybin 0.1,0.5 mg/kg ↑ fear response on trial 1 of TFC for extinction; Psylocibin ↓ fear response on trial 10 of TFC for extinction; Psylocibin 1mg/kg ↓ BrdU+ neurons in DG; NBOMe ↓ BrdU+ neurons in DG.	(Catlow et al., 2013)
Physiological assessment (AA and monoamines levels, utilization rate of monoamines)	Ayahuasca ↑ GABA levels in Hipp; Ayahuasca ↓ Glycine and GABA levels on Amygdala (dose dependent); Ayahuasca ↑ 5-HT on Hipp; Ayahuasca ↑ all monoamines on Amygdala; Ayahuasca 800mg/kg ↓ 5-HT use rate on Hipp; Ayahuasca ↓ monoamine consumption on amygdala.	Ayahuasca modifies 5-HT levels and consumption in a complex manner throughout the brain.	(Castro-Neto et al., 2013)
Behavioral assessment (Task oriented behavior; HTR)	DMT synthetic isoforms has comparable effect to its natural counterparts in a Zebrafish cue-oriented locomotion protocol; 5-MeO-IsoDMT do not produce any HTR still have in vitro psychoplastogens properties	5-MeO-IsoDMT do not produce any HTR still have in vitro psychoplastogens properties	(Dunlap et al., 2020)
Physiological assesment (BrdU+ and DCX+ cells) **Behavioral assesment** (MWM; NORT)	DMT ↑ survival and proliferation of newborn granule cells BrdU+ and DCX+ which is blocked by sigma-1R antagonist; DMT ↓ escape latency in MWM is blocked by 5-HT2AR antagonist; DMT ↑ exploration time in NORT.	DMT ↑ survival and proliferation of newborn granule cells BrdU+ and DCX+ which is blocked by sigma-1R antagonist; DMT ↓ escape latency in MWM is blocked by 5-HT2AR antagonist; DMT ↑ exploration time in NORT.	(Morales-Garcia et al., 2020)
	In vitro studies		
Physiological assessment (cell viability by MTT and Calcein assay; ki67+ and PI+ cells)	AYA1, CEB, CEP all doses ↑ cell viability with 48h incubation; AYA2 1, 1.5 µg/mL ↑ cell viability with 72h incubation.	all samples exhibited a neuroprotective effect in at least one time point and/or dose against the 6-OHDA induced neurodegeneration*.	(Katchborian-Neto et al., 2020)

Physiological assessments (Neurite growth)	DMT, isoDMT and 1-Me-DMT ↑ neurite growth in a dose comparable manner; 5-ht2A binding is necessary to neurite growth effect of all psychoplastogens tested.	DMT, isoDMT and 1-Me-DMT ↑ neurite growth in a dose comparable manner.	(Dunlap et al., 2020)
Physiological assessments (Neurite growth, Spinogenesis)	DOI, DMT, LSD ↑ Neurite growth and spinogenesis; The effects of psychoplastogens are similar to increased BDNF in culture medium, and both act through TrkB and mTOR cascade as well 5-ht2A signaling	DOI, DMT, LSD ↑ Neurite growth and spinogenesis The effects of psychoplastogens are similar to increased BDNF in culture medium, and both act through TrkB and mTOR cascade as well 5-ht2A signaling	(Ly et al., 2018)
Physiological assessment (Musashi1, Nestin, Sox-2, PCNA, Tuj, Map-2, GADPH, GFAP, CNPase levels)	DMT ↓ of Musashi1, Nestin and sox2 expression levels are blocked by sigma-1R antagonist after 7 days treatment; DMT ↑ number of neurospheres and PCNA expression levels are blocked by sigma-1R antagonist after 7 days treatment; DMT ↑ tuj, Map-2 GFAP and CNPase are blocked by sigma-1R antagonist after 7 days treatment;	DMT ↓ of Musashi1, Nestin and sox2 expression levels are blocked by sigma-1R antagonist after 7 days treatment; DMT ↑ number of neurospheres and PCNA expression levels are blocked by sigma-1R antagonist after 7 days treatment; DMT ↑ tuj, Map-2 GFAP and CNPase are blocked by sigma-1R antagonist after 7 days treatment;	(Morales-Garcia et al., 2020)

vDG (ventral Dentate Gyrus); sEPSC (spontaneous Excitatory Post Synaptic Current); Pyr (Pyramidal neurons); Nac (Nucleus accumbens); PFC (Prefrontal Cortex); SN (Substantia Nigra); Hipp (Hippocampus); VTA (Ventral Tegmental Area); CUS (Chronic Unpredictable Stress model); OF (Open Field); TST (Tail Suspension Test); FST (Forced Swim Test); EPM (Elevated Plus Maze); 5-HT (Serotonin); TPH-2 (Tryptophan Hydroxylase 2); ETC-1 (Mitochondrial electron transporter complex-I); DCX (doublecortin); GC (Granule Cells); BrdU (Bromodeoxyuridine); AHP (After Hyperpolarization); AP (Action Potential); LFP (Local Field Potential); TFC (Trace Fear Conditioning); AA (Amino acids) HTR (Head-Twitch Response); GDNF (Glial Cell Derived Neurotrophic Factor); BDNF (Brain Derived Neurotrophic Factor); NGF (Nerve Growth Factor); pet1 [pheochromocytoma 12 ETS (E26 transformation-specific)]; DRN (Dorsal Raphe Nucleus); MTT (3-[4,5-Dimethylthiazol-2-yl]- 2,5-diphenyl tetrazolium bromide); Ki67 (Marker Of Proliferation Ki-67); AYA1 (santo Daime's Ayahuasca decoction); CEB (*Banisteriopsis caapi* crude extract); CEP (*Psychotria viridis* crude extract); AYA2 (Ayahuasca prepared by authors); TrkB (Tropomyosin receptor kinase B); mTOR (mechanistic target of rapamycin).

Entactogens/ empathogens

As atypical psychedelics (Calvey & Howells, 2018), those compounds typically induce an emotional sense of "oneness," notably enhancing empathy on top of the usual psychedelic experience (Nichols, 1986). These unique properties distinguish them from other classes of drugs. All entactogens bind to 5-HT$_{1A}$ and 5-HT$_2$ receptors with varying affinities. Some phenethylamines, such as MDMA (3,4-methylenedioxymethamphetamine) and its analogs, affect other monoamines by blocking their uptake and forcing their release into the synaptic cleft (Papaseit et al., 2020).

Although some tryptamines are included as entactogens (Nagai et al., 2007) our review found only articles exploring the effects of MDMA within our scope. MDMA was initially developed in 1912 by Merck chemist Anton Köllisch to be an appetite suppressant, but it was never sold for this purpose (Kalant, 2001). In the 1970s, psychopharmacologist Alexander Shulgin synthesized it and, after self-experimentation, suggested its use as a medicine to aid patients undergoing therapy for PTSD and addiction disorders (Shulgin & Shulgin, 1991). In the 1980s, MDMA became a street drug, often sold illicitly as "molly" or "ecstasy" (NIDA, 2020). Today, MDMA is in the final stages of clinical research for approval as a psychotherapeutic tool to treat PTSD (Emerson et al., 2014). The included articles explore the effects of MDMA during embryonic and/or neonatal stages (57.14%), adolescence (28.57%), and adulthood (28.57%) in murine models, with some articles examining more than one ontogenetic window (**Supp. Table 5**).

The findings of these studies, as summarized in Table 5, illustrate that the effects of MDMA on neurogenesis can vary depending on the dosage and duration of administration. In a study by Hernández-Rabaza et al., researchers attempted to mimic the binge consumption of MDMA often seen in social settings such as raves. They administered a dose of 40mg/kg over two days to male rats (the age of the rats was not reported). This treatment did not affect the proliferation of new cells. However, when assessed 14 days after the treatment, there was a noticeable 40% decrease in new cells (BrdU+ cells) in the dentate gyrus compared to the control group. The researchers also examined potential abnormalities in the growth of new neurons but found no difference between the MDMA-treated and control groups. (Hernández-Rabaza et al., 2006). Moreover, a study by Catlow et al. investigated the effects of a longer, chronic administration of MDMA. Rats were

given a lower dose of 5 mg/kg per day for 10 days. The researchers observed an initial increase in new cells (BrdU+ cells) immediately after the MDMA treatment, but a significant decrease (around 40-50%) was noted 14 days later. The survival of these new neurons, measured by co-labeling BrdU and NeuN, was also reduced by approximately 40-50%. (Catlow et al., 2010).

The results presented by Catlow (2010) and Hernández-Rabaza et al. (2006) prompt further investigation into potential differential effects of MDMA across developmental stages. García & García (2016) report that a 60mg/kg dose of MDMA (3x5mg/kg over 4 days) more significantly impacted proliferation in adult rats (P48-P58) compared to adolescent rats (P27-P37), as demonstrated by ki67 and BrdU labeling (García-Cabrerizo & García-Fuster, 2016). Given the unreported age of the rats in Hernández-Rabaza et al.'s (2006) study, a direct comparison with García & García's findings is not recommended. However, the reported age range of rats in Catlow et al.'s study (P28-39) corresponds to adolescence/young adulthood, introducing an element of ambiguity.

Further, García & García's study also revealed a positive correlation between mature BDNF (mBDNF) levels and numbers of BrdU+ cells (r=0.492), with a stronger correlation observed between 5-HT2CR expression and mBDNF (r = 0.713). This led the authors to posit that chronic MDMA exposure (60 mg/kg) negatively affects overall plasticity, with more pronounced effects in adult animals. Interestingly, no significant impact was found when a lower dosage of 15mg/kg (3x5 mg within 1 day) was administered (García-Cabrerizo & García-Fuster, 2016).

With the growing popularity and widespread use of MDMA among young people and adults (Armenian & Rodda, 2020), three other studies focused on examining the effects of MDMA during early stages of life, given concerns about the drug's potential teratogenic properties. In particular, a team of researchers from Seoul, Korea, investigated the impact on adult neurogenesis in embryos from dams exposed to 1.25 or 20 mg/kg MDMA for 36 days (E6-P21) via oral gavage. At 11 weeks post-exposure, animals were treated with BrdU. The study found that both doses led to reduced cell proliferation compared to saline, but only the higher dose of 20 mg/kg resulted in decreased survival of abGC as indicated by BrdU/NeuN labeling. Notably, only comparisons with female offspring in the experimental groups were reported in survival assays (Cho et al., 2008).

In a teratology study conducted in Ohio, USA, Thompson et al. (2012) found that maternal ingestion of MDMA at a similar dose during E14-E20 increased fiber density in the prelimbic cortex (PLC) and CA1. Meanwhile, the expression of the norepinephrine transporter (NET) was elevated in the cornus ammonis regions, but not in the dentate gyrus (DG), Locus Coeruleus (LC), or PLC (V. B. B. Thompson et al., 2012). The authors suggested further exploration of NET upregulation to strengthen the association between downstream morphological (increased plasticity) and neurochemical abnormalities (upregulation of NE in the PFC and Nac) with previously reported cognitive and behavioral changes related to novelty (Koprich et al., 2003; V. B. Thompson et al., 2009; V. B. B. Thompson et al., 2012) However, the authors emphasized that the existing data does not suffice to determine causality. Subsequently, Canales and Ferrer-Donato (2014) demonstrated that a dose of 10mg/kg MDMA administered during a shorter pregnancy window (3 days, E13-E15) combined with alcohol significantly reduced neurogenesis (both proliferation and survival) by approximately 40% in adult female offspring compared to saline. Behavioral assessment via the Radial Arm Maze (RM) revealed about a 30% reduction in locomotion and a roughly 50% increase in working memory errors. Notably, the within-group variance was quite narrow, and only the group exposed to both drugs exhibited these impairments (Canales & Ferrer-Donato, 2014).

We conclude that intense abuse of MDMA with high acute doses, as much as 40 mg/kg (equivalent to a 3.6 grams dose for an average 90kg human), results in enduring imbalances in the monoamine system and subsequent downregulation of neurogenesis. Conversely, low doses (equivalent to 0.112 grams for an average 90kg human) have no detectable effects on morphological abnormalities in adult brain tissue. It is noteworthy that the phase 3 studies currently being conducted by the Multidisciplinary Association for Psychedelic Studies (MAPS) are utilizing doses ranging from 0.08-0.18 grams of MDMA per patient (Mithoefer et al., 2019). Therefore, neurobiological studies should examine very low doses of MDMA to answer the question of how MDMA impacts brain physiology. It is crucial to emphasize that the use of psychoactive substances is never recommended for pregnant women.

Table 5 Results reported by the reviewed Entactogens studies

Outcome variable	Reported Results	Effect on Neurogenesis	Reference
In vivo studies			
Physiological assessment (BrdU+ cells in offspring)	Embryonic MDMA exposure ↓BrdU+ cells in offspring.	Embryonic MDMA ↓ BrdU+ cells in adulthood.	(Cho et al., 2008)
Behavioral assessment (SC, CWM, MWM) **Physiological assessment** (BW)	MDMA ↑ score in CWM; MDMA ↑ Path length, Heading error in MWM; MDMA ↓ quad accuracy and crossovers in MWM; MDMA ↓ BW.	MDMA ↓ behavior related do pattern separation.	(Schaefer et al., 2013)
Physiological assessment (BRDU+, DCX+, KI67+ cells, neurite growth)	MDMA has no effect on BrdU+ , Ki67+ cells and Neurite growth; MDMA ↓ DCX+ cells.	MDMA has no effect on proliferation; MDMA ↓ survival.	(Hernández-Rabaza et al., 2006)
Behavioral assessment (CPP) **Physiological assessment** (BrdU+, NeuN+ cells)	MDMA ↑ CPP only at 2.5mg/kg dose; 5mg/kg MDMA ↑ BrdU+ cells 24hrs after BRDU injection; MDMA ↓ BrdU + cells 14 days after BrdU injection; 5mg/kg MDMA ↓ colableled BrdU+ and NeuN+ cells 14 days after BrdU injection.	5mg/kg MDMA ↑ BrdU+ cells 24hrs after BRDU injection; MDMA ↓ BrdU + cells 14 days after BrdU injection (dose dependent effect); 5mg/kg MDMA ↓ colableled BrdU+ and NeuN+ cells.	(Catlow et al., 2010)
Physiological assessment (Fiber density on PLC and Hipp, NET density, NE levels)	MDMA ↑ fiber density in PLC and CA1; MDMA ↑ NET density in CA1. CA2 and CA3 but not on DG; MDMA ↑ NE levels on PFC and Nac.	MDMA promote neurite outgrowth in a complex manner.	(V. B. B. Thompson et al., 2012)
Behavioral assessment (RM) **Physiological assessment** (BrdU+, DCX+ cells)	ETOH/MDMA ↑ WM errors in RM; ETOH/MDMA ↓ Locomotor activity in RM; ETOH/MDMA ↓ BrdU+ and DCX+ cells.	ETOH/MDMA ↓ BrdU+ and DCX+ cells.	(Canales & Ferrer-Donato, 2014)
Physiological assessment (BW, BrdU+, Ki-67, proBDNF and mBDNF and 5-HT2cR)	MDMA has no effect on BW; Chronic MDMA ↓ Ki-67 + cells; Chronic MDMA ↓ BrdU+ cells only in P42 mice; Chronic MDMA ↓ proBDNF; chronic MDMA ↓ mBDNF only in P21 mice; 5-HT2cR levels increase as mBDNF in P42 chronic MDMA treated mice.	Chronic MDMA ↓ Ki-67 + cells; Chronic MDMA ↓ BrdU+ cells only in P42 mice; Chronic MDMA ↓ proBDNF; chronic MDMA ↓ mBDNF only in P21 mice.	(García-Cabrerizo & García-Fuster, 2016)

SC (Straight Channel test); CWM (Cincinnati Water Maze); MWM (Morris Water Maze); BW (Body Weight); BrdU (Bromodeoxyuridine); DCX (Doublecortin); KI67 (Marker of Proliferation Ki-67); CPP (Conditioned Place Preference); PLC (Pre-limbic Cortex); Hipp (Hippocampus); NET (Noradrenergic Transporter); NE (Noradrenaline); DG (Dentate Gyrus); PFC (Prefrontal Cortex); Nac (Nucleus Accumbens); ETOH (Ethanol); RM (Radial Maze); proBDNF (precursor protein of BDNF); mBDNF (mature BDNF).

Other Psychedelics

In this section, we go through additional articles that employed psychedelic drugs from chemical categories not previously discussed. These include the substituted amphetamine with psychedelic effects 2,5-Dimethoxy-4-iodoamphetamine (DOI) (**Supp. Table 6**) (Belmer et al., 2018; Berthoux et al., 2019; Jha et al., 2008; Jones et al., 2009; Ly et al., 2018; Marinova et al., 2017) and the substituted phenethylamine 25I-NBOMe (**Supp. Table 5**) (Catlow et al., 2013).

A group from the French National Institute of Health and Medical Research (INSERM) assessed the capability of the 5-HT$_{2B}$R to regulate Dorsal Raphe Nucleus (DRN) serotonin release. Activation of pet1+ DRN cells by the tryptamine BW723C86 (1µM), a 5-HT$_{2B}$ receptor agonist, administered *ex vivo* increased the spontaneous firing rate. Intriguingly, in subsequent experiments, 5mg/kg DOI in vivo failed to boost the HTR in animals with a knockout of the 5-HT$_{2B}$R, and 20 mg/kg MDMA in the same genotype also failed to replicate the increase in locomotor response observed in control mice. These behaviors were evaluated using the Open Field (OF) test (Belmer et al., 2018), indicating that 5-HT$_{2B}$R may play a larger role in the behavioral effects of psychedelics on mice than previously believed.

A research team at the Tata Institute of Fundamental Research in Mumbai, India, using the BrdU labeling technique, reported that neither LSD (0.5 mg/kg) nor DOI (8 mg/kg) could increase proliferation in rats (age not reported) under either chronic (7 days) or acute treatment regimens (Jha et al., 2008). Another group from the Northwestern University Feinberg School of Medicine in Chicago, Illinois, reported that 1µM DOI could transiently promote dendritic remodeling in cultured cortical pyramidal neurons from rats, and this plasticity was mediated by Kalirin-7. The change involved the enlargement of spines, which began 30 minutes after exposure and returned to control size 60 minutes later (Jones et al., 2009). Ly and colleagues from UC Davis, in an article discussed previously (**Table 2**), extended the scope of their findings to other species, demonstrating that LSD and DOI (dose not reported) could promote neuritogenesis in cultured neurons from mice and Drosophila larvae (Ly et al., 2018). Additionally, Marinova and colleagues employed human neuroblastoma SK-N-SH cells, TrkA is necessary for neurite extension (Marinova et al., 2017).

The only report found to investigate DOI using electrophysiological methods was conducted by Berthoux and colleagues. They used 1µM DOI on ex vivo slices from P14-P21 mice and reported that DOI induced Long-Term Depression (LTD) in mPFC layer 5 pyramidal neurons, as evidenced by whole-cell patch-clamp. The amplitude of the AMPA ESPC (excitatory postsynaptic current) coming from layer 1 Pyramidal stimuli was almost halved, and this decrease was mediated by 5-HT2AR since mice knocked out for 5-HT2AR did not show the same response. The internalization of AMPA receptors containing the GluA2 subunit through Protein kinase C (PKC) was necessary for DOI-induced LTD but not when LTD was replicated by a fast electrical pairing protocol (Berthoux et al., 2019). Moreover, in an article already discussed in the tryptamine section (**Table 2**), Catlow and colleagues used the substituted phenethylamine 25I-NBOMe in a single experiment to explore if 5-HT2AR activation affects proliferation of adult-born granule cells (abGC) in mice (age not reported). All tested doses (0.1, 0.3, and 1 mg/kg) reduced proliferation, but only the 1 mg/kg dose had a significant effect on the number of BrdU+/NeuN+ cells compared to saline-treated mice (Catlow et al., 2013).

In summary, these studies suggest that the substituted amphetamine DOI and the substituted phenethylamine 25I-NBOMe may have a range of effects on neural plasticity and behavior. These effects appear to be modulated by different serotonin receptors, including 5-HT2BR and 5-HT2AR. More research is needed to further understand the molecular mechanisms and the functional implications of these changes. Also, the specificity of the effects of these substances on different stages of neurogenesis needs to be further investigated.

Table 6 Results reported by the reviewed studies using other psychedelics.

Outcome variable	Reported Results	Effect on Neurogenesis	Reference
	In vivo studies		
Electrophysiological assessment (Layer1 -↑ Layer 5 EPSC)	Doi ↓ Layer 1 AMPA ESPC Amplitude in Layer 5 Pyr mPFC; mPFC layer 5 Pyr of 5-HT$_{2A}$-KO mice do not react to DOI treatment; GluA2 subunit internalizations necessary for DOI-induced LTD in layer 5 Pyr mPFC; PKC inhibitor block DOI-induced LTD in layer 5 Pyr mPFC.	DOI promotes activity dependent plasticity mediated by 5-HT$_{2A}$R and PKC.	(Berthoux et al., 2019)
Physiological assessment (BrdU + cells)	Nor DOI or LSD ↑ BrdU+ Cell on DG; Ketanserin ↓ on acute but ↑ on chronic treatment.	Proliferation (=).	(Jha et al., 2008)
Behavioral assessment (OF) Physiological assessment (BRdU+ cells) **Electrophysiological assessment (Active membrane properties)**	BW723C86 ↑ firing rate on pet1+ DRN ex vivo; BW723C86 prevent 5-HT1A agonist induced inhibition of pet1+DRN in vivo; MDMA fail to ↑ locomotor activity on 5-HT2B-KO; DOI 5mg/kg fail to ↑ HTR in 5-HT2B-KO; Fluoxetine fails to ↑ BRdU+ cells in 5-HT2B-KO.	5-HT2B-KO mice shows insensibility to treatment by SSRI, MDMA and DOI as evaluated by authors.	(Belmer et al., 2018)
	In vitro studies		
Physiological assessment (5-HT2AR, PSD95, Kalirin-7 and bassoon localization; MUPP1 activity; Dendritic spine remodeling; PAK phosphorylation levels)	5-HT2AR colocalizes with PSD95, kalirin-7 and bassoon proteins mainly on small and medium size spines on both shaft and heads; MUPP1 activity promotes 5-HT2AR migration to cell periphery on Cos-7 mature cells; DOI promotes spine shape remodeling without changing global spine density through PAK phosphorylation and subsequent Kalirin-7 activity in cultured cortical neurons of rats.	DOI transiently promote dendritic spine remodeling towards larger spines.	(Jones et al., 2009)
Physiological assessment (TrkA tyr Phosphorylation levels at AA residue Tyr490 and Tyr785, Neurite Extension)	DOI all doses for 30 or 120 minutes ↑ TrkA phosphorylation in SK-N-SH cells; DOI all doses for 30 or 120 minutes ↑ TrkA phosphorylation at AA residue Tyr490 but not Tyr785; 5-HT2A and 5-HT2B block do not affect TrkA phosphorylation induced by DOI; Chronic DOI 1,5 μM ↑ neurite extension, and is blocked by TrkA inhibitor.	Chronic DOI 1,5 μM ↑ neurite extension, and is blocked by TrkA inhibitor.	(Marinova et al., 2017)

EPSC (Excitatory Post Synaptic Current); Pyr (Pyramidal); mPFC (medial Prefrontal Cortex); GluA2 (AMPA receptor subunit GluA2); LTD (Long term Depression); PKC (Protein Kinase C); BrdU (Bromodeoxyuridine); DG (Dentate Gyrus); PSD95 (Post-synaptic Density Protein 95); bassoon (Pre synaptic cytomatrix protein bassoon); MUPP1 (Multiple PDZ protein-1); Cos-7 (Fibroblast-like cell line from monkey kidney); PAK (P21-activated kinase); TrkA (Tropomyosin receptor kinase A); AA (Amino Acid); Tyr (Tyrosine).

Conclusion

This systematic review sought to reconcile the diverse outcomes observed in studies investigating the impact of psychedelics on neurogenesis. It has become evident that a multitude of factors, including dosage, treatment frequency, and the choice of model, significantly influence study results. Additionally, this review has integrated studies examining related aspects of neuroplasticity, such as neurotrophic factor regulation and synaptic remodeling, regardless of the specific brain regions investigated, in recognition of the potential transferability of these findings.

One of the key takeaways from this review is the critical role of dosage in determining the effects of psychedelics. Therapeutic dosages, whether administered acutely or chronically, have been shown to stimulate neurotrophic factor production, proliferation and survival of adult-born granule cells (abGC), and neuritogenesis. While the precise mechanisms underlying these effects remain to be fully elucidated, the capacity of some psychedelics to induce neuroplastic changes is incontrovertible. Therefore, it is essential to conduct rigorous preclinical and clinical trials to fully understand the mechanisms of action, the optimal dosages and treatment regimens, and the potential risks and side effects. Importantly, the effects of these substances on different stages of life, from embryonic development to adulthood, need to be further investigated. Moreover, it is also crucial for future studies in relevant disease models, such as depression, anxiety, and Alzheimer's disease, to carefully calibrate experimental parameters, including the age of the animals, treatment regimen, and time point of analysis.

As psychedelics edge closer to mainstream therapeutic use, it is of utmost importance for mental health professionals to appreciate the role of set and setting (Hartogsohn, 2017) in shaping the psychedelic experience. Drug developers, too, should carefully evaluate contraindications and potential interactions, given the unique pharmacological profiles of these compounds and the relative lack of familiarity with them within the psychiatric community. It is recommended that practitioners intending to work with psychedelics undergo supervised clinical training and achieve professional certification, akin to the practices upheld by Amerindian traditions with a long history of using these substances (Desmarchelier et al., 1996; Labate & Cavnar, 2014; Naranjo, 1979; Svobodny, 2014). This will ensure the safe and efficacious deployment of these powerful tools for neuroplasticity in the therapeutic setting.

The rapidly evolving field of psychedelics in neuroscience is providing exciting opportunities for therapeutic intervention. Nevertheless, it is crucial to approach this potential with due diligence, taking into consideration the intricate balance of variables that contribute to the outcomes observed in pre-clinical models. The effects of psychedelics on neuroplasticity underline their potential utility in various neuropsychiatric conditions, but also stress the need for thorough understanding and careful handling.

Acknowledgments

RVLC thanks the colleagues from the Brain Institute and mentors from the PPG in Psychobiology (UFRN) for support and guidance, as well as CAPES (*Coordenação de Aperfeiçoamento de Pessoal de Nível Superior*, Brazil) for the scholarship funding. TCM is grateful for the funding from the Royal Physiographic Society of Lund.

Declaration of competing interests

None.

References

Akinfiresoye, L., & Tizabi, Y. (2013). Antidepressant effects of AMPA and ketamine combination: role of hippocampal BDNF, synapsin, and mTOR. *Psychopharmacology*, *230*(2), 291–298. https://doi.org/10.1007/s00213-013-3153-2

Aleksandrova, L. R., & Phillips, A. G. (2021). Neuroplasticity as a convergent mechanism of ketamine and classical psychedelics. *Trends in Pharmacological Sciences*, *42*(11), 929–942. https://doi.org/10.1016/j.tips.2021.08.003

Alper, K. R., Lotsof, H. S., & Kaplan, C. D. (2008). The ibogaine medical subculture. *Journal of Ethnopharmacology*. https://doi.org/10.1016/j.jep.2007.08.034

Andrade, C. (2017). Ketamine for Depression, 1: Clinical Summary of Issues Related to Efficacy, Adverse Effects, and Mechanism of Action. *The Journal of Clinical Psychiatry*, *78*(04), e415–e419. https://doi.org/10.4088/JCP.17f11567

Araque, A., Castillo, P. E., Manzoni, O. J., & Tonini, R. (2017). Synaptic functions of endocannabinoid signaling in health and disease. *Neuropharmacology*, *124*, 13–24. https://doi.org/10.1016/j.neuropharm.2017.06.017

Araújo, A. M., Carvalho, F., Bastos, M. D. L., Guedes de Pinho, P., & Carvalho, M. (2015). The hallucinogenic world of tryptamines: an updated review. *Archives of Toxicology*. https://doi.org/10.1007/s00204-015-1513-x

Armenian, P., & Rodda, L. N. (2020). Two Decades of Ecstasy: Shifting Demographic Trends in Decedents Using MDMA. *Journal of Analytical Toxicology*. https://doi.org/10.1093/jat/bkaa193

Barker, J. M., Boonstra, R., & Wojtowicz, J. M. (2011). From pattern to purpose: how comparative studies contribute to understanding the function of adult neurogenesis. *European Journal of Neuroscience*, *34*(6), 963–977. https://doi.org/10.1111/j.1460-9568.2011.07823.x

Barnea, A., & Pravosudov, V. (2011). Birds as a model to study adult neurogenesis: Bridging evolutionary, comparative and neuroethological approaches. *European Journal of Neuroscience*. https://doi.org/10.1111/j.1460-9568.2011.07851.x

Baumeister, A. A., Hawkins, M. F., & Uzelac, S. M. (2003). The Myth of Reserpine-Induced Depression: Role in the Historical Development of the Monoamine Hypothesis. *Journal of the History of the Neurosciences*, *12*(2), 207–220. https://doi.org/10.1076/jhin.12.2.207.15535

Begley, C. G., & Ioannidis, J. P. A. (2015). Reproducibility in Science: improving the standard for basic and preclinical research. *Circulation Research*, *116*(1), 116–126. https://doi.org/10.1161/CIRCRESAHA.114.303819

Beiersdorf, J., Hevesi, Z., Calvigioni, D., Pyszkowski, J., Romanov, R., Szodorai, E., Lubec, G., Shirran, S., Botting, C. H., Kasper, S., Guy, G. W., Gray, R., Marzo, V. Di, Harkany, T., & Keimpema, E. (2020). Adverse effects of Δ9-tetrahydrocannabinol on neuronal bioenergetics during postnatal development. *JCI Insight*, *5*(23). https://doi.org/10.1172/jci.insight.135418

Belmer, A., Quentin, E., Diaz, S. L., Guiard, B. P., Fernandez, S. P., Doly, S., Banas, S. M., Pitychoutis, P. M., Moutkine, I., Muzerelle, A., Tchenio, A., Roumier, A., Mameli, M., & Maroteaux, L. (2018). Positive regulation of raphe serotonin neurons by serotonin 2B receptors. *Neuropsychopharmacology*, *43*(7), 1623–1632. https://doi.org/10.1038/s41386-018-0013-0

Belouin, S. J., & Henningfield, J. E. (2018). Psychedelics: Where we are now, why we got here, what we must do. *Neuropharmacology*, *142*, 7–19. https://doi.org/10.1016/j.neuropharm.2018.02.018

Berghuis, P., Rajnicek, A. M., Morozov, Y. M., Ross, R. A., Mulder, J., Urban, G. M., Monory, K., Marsicano, G., Matteoli, M., Canty, A., Irving, A. J., Katona, I., Yanagawa, Y., Rakic, P., Lutz, B., Mackie, K., & Harkany, T. (2007). Hardwiring the Brain: Endocannabinoids Shape Neuronal Connectivity. *Science*, *316*(5828), 1212–1216. https://doi.org/10.1126/science.1137406

Berman, R. M., Cappiello, A., Anand, A., Oren, D. A., Heninger, G. R., Charney, D. S., & Krystal, J. H. (2000). Antidepressant effects of ketamine in depressed patients. *Biological Psychiatry*, *47*(4), 351–354. https://doi.org/10.1016/S0006-3223(99)00230-9

Berthoux, C., Barre, A., Bockaert, J., Marin, P., & Bécamel, C. (2019). Sustained Activation of Postsynaptic 5-HT 2A Receptors Gates Plasticity at Prefrontal Cortex Synapses. *Cerebral Cortex*, *29*(4), 1659–1669. https://doi.org/10.1093/cercor/bhy064

Bonini, S. A., Premoli, M., Tambaro, S., Kumar, A., Maccarinelli, G., Memo, M., & Mastinu, A. (2018). Cannabis sativa: A comprehensive ethnopharmacological review of a medicinal plant with a long history. In *Journal of Ethnopharmacology* (Vol. 227, pp. 300–315). https://doi.org/10.1016/j.jep.2018.09.004

Brito-da-Costa, A. M., Dias-da-Silva, D., Gomes, N. G. M., Dinis-Oliveira, R. J., & Madureira-Carvalho, Á. (2020). Toxicokinetics and Toxicodynamics of Ayahuasca Alkaloids N,N-Dimethyltryptamine (DMT), Harmine, Harmaline and Tetrahydroharmine: Clinical and Forensic Impact. *Pharmaceuticals*, *13*(11), 334. https://doi.org/10.3390/ph13110334

Browne, C. A., & Lucki, I. (2013). Antidepressant effects of ketamine: Mechanisms underlying fast-acting novel antidepressants. In *Frontiers in Pharmacology*. https://doi.org/10.3389/fphar.2013.00161

Brunton, L. L., Chabner, B. A., & Knollmann, B. C. (2011). *Goodman & Gilman's: The Pharmacological Basis of Therapeutics* (J. F. Shanahan & C. Naglieri (eds.); 12ª Editio). McGraw - Hill Book Company.

Callaway, J. C. (2005). Various Alkaloid Profiles in Decoctions of Banisteriopsis Caapi. *Journal of Psychoactive Drugs*, *37*(2), 151–155. https://doi.org/10.1080/02791072.2005.10399796

Calvey, T., & Howells, F. M. (2018). An introduction to psychedelic neuroscience. In *Progress in Brain Research* (1st ed., Vol. 242, pp. 1–23). Elsevier B.V. https://doi.org/10.1016/bs.pbr.2018.09.013

Canales, J. J., & Ferrer-Donato, A. (2014). Prenatal Exposure to Alcohol and 3,4-Methylenedioxymethamphetamine (Ecstasy) Alters Adult Hippocampal Neurogenesis and Causes Enduring Memory Deficits. *Developmental Neuroscience*, *36*(1), 10–17. https://doi.org/10.1159/000356820

Carbonaro, T. M., & Gatch, M. B. (2016). Neuropharmacology of N,N-Dimethyltryptamine. *Brain Research Bulletin*. https://doi.org/10.1016/j.brainresbull.2016.04.016

Carliner, H., Brown, Q. L., Sarvet, A. L., & Hasin, D. S. (2017). Cannabis use, attitudes, and legal status in the U.S.: A review. *Preventive Medicine*, *104*, 13–23. https://doi.org/10.1016/j.ypmed.2017.07.008

Castrén, E., Voikar, V., & Rantamaki, T. (2007). Role of neurotrophic factors in depression. *Current Opinion in Pharmacology*, *7*(1), 18–21. https://doi.org/10.1016/j.coph.2006.08.009

Castro-Neto, E. F. de, Cunha, R. H. da, Silveira, D. X. da, Yonamine, M., Gouveia, T. L. F., Cavalheiro, E. A., Amado, D., & Naffah-Mazzacoratti, M. da G. (2013). Changes in aminoacidergic and monoaminergic neurotransmission in the hippocampus and amygdala of rats after ayahuasca ingestion. *World Journal of Biological Chemistry*, *4*(4), 141. https://doi.org/10.4331/wjbc.v4.i4.141

Catlow, B. J., Badanich, K. A., Sponaugle, A. E., Rowe, A. R., Song, S., Rafalovich, I., Sava, V., Kirstein, C. L., & Sanchez-Ramos, J. (2010). Effects of MDMA ("ecstasy") during adolescence on place conditioning and hippocampal neurogenesis. *European Journal of Pharmacology*, *628*(1–3), 96–103. https://doi.org/10.1016/j.ejphar.2009.11.017

Catlow, B. J., Song, S., Paredes, D. A., Kirstein, C. L., & Sanchez-Ramos, J. (2013). Effects of psilocybin on hippocampal neurogenesis and extinction of trace fear conditioning. *Experimental Brain Research*, *228*(4), 481–491. https://doi.org/10.1007/s00221-013-3579-0

Chang, L., & Chronicle, E. P. (2007). Functional Imaging Studies in Cannabis Users. *The Neuroscientist*, *13*(5), 422–432. https://doi.org/10.1177/1073858406296601

Chen, B., Bromley-Brits, K., He, G., Cai, F., Zhang, X., & Song, W. (2010). Effect of Synthetic Cannabinoid HU210 on Memory Deficits and Neuropathology in Alzheimers Disease Mouse Model. *Current Alzheimer Research*, *7*(3), 255–261. https://doi.org/10.2174/156720510791050948

Cho, K. O., Rhee, G. S., Kwack, S. J., Chung, S. Y., & Kim, S. Y. (2008). Developmental exposure to 3,4-methylenedioxymethamphetamine results in downregulation of neurogenesis in the adult mouse hippocampus. *Neuroscience*, *154*(3), 1034–1041. https://doi.org/10.1016/j.neuroscience.2008.04.040

Christie, B. R., & Cameron, H. A. (2006). Neurogenesis in the Adult Hippocampus. *European Journal of Neuroscience*, *207*, 199–207. https://doi.org/10.1002/hipo.20151

Cuccurazzu, B., Zamberletti, E., Nazzaro, C., Prini, P., Trusel, M., Grilli, M., Parolaro, D., Tonini, R., & Rubino, T. (2018). Adult Cellular Neuroadaptations Induced by Adolescent THC Exposure in Female Rats Are Rescued by Enhancing Anandamide Signaling. *International Journal of Neuropsychopharmacology*, *21*(11), 1014–1024. https://doi.org/10.1093/ijnp/pyy057

Dakic, V., de Moraes Maciel, R., Drummond, H., Nascimento, J. M., Trindade, P., & Rehen, S. K. (2016). Harmine stimulates proliferation of human neural progenitors. *PeerJ*, *2016*(12), 1–13. https://doi.org/10.7717/peerj.2727

Dakic, V., Maciel, R. de M., Drummond, H., Nascimento, J. M., Trindade, P., & Rehen, S. K. (2016). Harmine stimulates proliferation of human neural progenitors. *PeerJ*, *4*(12), e2727. https://doi.org/10.7717/peerj.2727

Dakwar, E., Nunes, E. V., Hart, C. L., Hu, M. C., Foltin, R. W., & Levin, F. R. (2018). A sub-set of psychoactive effects may be critical to the behavioral impact of ketamine on cocaine use disorder: Results from a randomized, controlled laboratory study. *Neuropharmacology*, *142*, 270–276. https://doi.org/10.1016/j.neuropharm.2018.01.005

Danforth, A. L., Struble, C. M., Yazar-Klosinski, B., & Grob, C. S. (2016). MDMA-assisted therapy: A new treatment model for social anxiety in autistic adults. *Progress in Neuro-Psychopharmacology and Biological Psychiatry*. https://doi.org/10.1016/j.pnpbp.2015.03.011

Data-Franco, J., Singh, A., Popovic, D., Ashton, M., Berk, M., Vieta, E., Figueira, M. L., & Dean, O. M. (2017). Beyond

the therapeutic shackles of the monoamines: New mechanisms in bipolar disorder biology. *Progress in Neuro-Psychopharmacology and Biological Psychiatry*, *72*, 73–86. https://doi.org/10.1016/j.pnpbp.2016.09.004

de Araujo, D. B., Ribeiro, S., Cecchi, G. a, Carvalho, F. M., Sanchez, T. a, Pinto, J. P., de Martinis, B. S., Crippa, J. a, Hallak, J. E. C., & Santos, A. C. (2012). Seeing with the eyes shut: neural basis of enhanced imagery following Ayahuasca ingestion. *Human Brain Mapping*, *33*, 2550–2560. https://doi.org/10.1002/hbm.21381

de Fonseca, F. R., del Arco, I., Bermudez-Silva, F. J., Bilbao, A., Cippitelli, A., & Navarro, M. (2005). The endocannabinoid system: Physiology and pharmacology. *Alcohol and Alcoholism*. https://doi.org/10.1093/alcalc/agh110

De Freitas, C. M., Busanello, A., Schaffer, L. F., Peroza, L. R., Krum, B. N., Leal, C. Q., Ceretta, A. P. C., Da Rocha, J. B. T., & Fachinetto, R. (2016). Behavioral and neurochemical effects induced by reserpine in mice. *Psychopharmacology*. https://doi.org/10.1007/s00213-015-4118-4

de Oliveira Silveira, G., Guimarães dos Santos, R., Rebello Lourenço, F., Novak Rossi, G., Hallak, J., & Yonamine, M. (2020). Stability Evaluation of DMT and Harmala Alkaloids in Ayahuasca Tea Samples. *Molecules*, *25*(9), 2072. https://doi.org/10.3390/molecules25092072

de Rios, M. D., & Grob, C. S. (2005). Ayahuasca Use in Cross-Cultural Perspective. *Journal of Psychoactive Drugs*, *37*(2), 119–121. https://doi.org/10.1080/02791072.2005.10399790

Demirakca, T., Sartorius, A., Ende, G., Meyer, N., Welzel, H., Skopp, G., Mann, K., & Hermann, D. (2011). Diminished gray matter in the hippocampus of cannabis users: Possible protective effects of cannabidiol. *Drug and Alcohol Dependence*, *114*(2–3), 242–245. https://doi.org/10.1016/j.drugalcdep.2010.09.020

Desmarchelier, C., Gurni, A., Ciccia, G., & Giulietti, A. M. (1996). Ritual and medicinal plants of the Ese'ejas of the Amazonian rainforest (Madre de Dios, Perú). *Journal of Ethnopharmacology*. https://doi.org/10.1016/0378-8741(96)01390-6

DeVos, S. L., & Miller, T. M. (2013). Direct intraventricular delivery of drugs to the rodent central nervous system. *Journal of Visualized Experiments : JoVE*, *75*, e50326. https://doi.org/10.3791/50326

Dickinson, B. C., Peltier, J., Stone, D., Schaffer, D. V, & Chang, C. J. (2011). Nox2 redox signaling maintains essential cell populations in the brain. *Nature Chemical Biology*, *7*(2), 106–112. https://doi.org/10.1038/nchembio.497

Domínguez-Clavé, E., Soler, J., Elices, M., Pascual, J. C., Álvarez, E., de la Fuente Revenga, M., Friedlander, P., Feilding, A., & Riba, J. (2016). Ayahuasca: Pharmacology, neuroscience and therapeutic potential. *Brain Research Bulletin*, *126*(Pt 1), 89–101. https://doi.org/10.1016/j.brainresbull.2016.03.002

Domino, E. F., & Warner, D. S. (2010). Taming the Ketamine Tiger. *Anesthesiology*, *113*(3), 678–684. https://doi.org/10.1097/ALN.0b013e3181ed09a2

Downer, E. J., Gowran, A., & Campbell, V. A. (2007). A comparison of the apoptotic effect of Δ9-tetrahydrocannabinol in the neonatal and adult rat cerebral cortex. *Brain Research*, *1175*(1), 39–47. https://doi.org/10.1016/j.brainres.2007.07.076

Drew, L. J., Fusi, S., & Hen, R. (2013). Adult neurogenesis in the mammalian hippocampus: why the dentate gyrus? *Learning & Memory (Cold Spring Harbor, N.Y.)*, *20*(12), 710–729. https://doi.org/10.1101/lm.026542.112

Duman, R. S., & Li, N. (2012). A neurotrophic hypothesis of depression: Role of synaptogenesis in the actions of NMDA receptor antagonists. In *Philosophical Transactions of the Royal Society B: Biological Sciences* (Vol. 367, Issue 1601, pp. 2475–2484). https://doi.org/10.1098/rstb.2011.0357

Duncan, W. C., Sarasso, S., Ferrarelli, F., Selter, J., Riedner, B. A., Hejazi, N. S., Yuan, P., Brutsche, N., Manji, H. K., Tononi, G., & Zarate, C. A. (2013). Concomitant BDNF and sleep slow wave changes indicate ketamine-induced plasticity in major depressive disorder. *International Journal of Neuropsychopharmacology*, *16*(2), 301–311. https://doi.org/10.1017/S1461145712000545

Dunlap, L. E., Azinfar, A., Ly, C., Cameron, L. P., Viswanathan, J., Tombari, R. J., Myers-Turnbull, D., Taylor, J. C., Grodzki, A. C., Lein, P. J., Kokel, D., & Olson, D. E. (2020). Identification of Psychoplastogenic N , N -Dimethylaminoisotryptamine (isoDMT) Analogues through Structure–Activity Relationship Studies. *Journal of Medicinal Chemistry*, *63*(3), 1142–1155. https://doi.org/10.1021/acs.jmedchem.9b01404

Emerson, A., Ponté, L., Jerome, L., & Doblin, R. (2014). History and Future of the Multidisciplinary Association for Psychedelic Studies (MAPS). *Journal of Psychoactive Drugs*. https://doi.org/10.1080/02791072.2014.877321

Eriksson, P. S., Perfilieva, E., Björk-Eriksson, T., Alborn, A. M., Nordborg, C., Peterson, D. A., & Gage, F. H. (1998). Neurogenesis in the adult human hippocampus. *Nature Medicine*. https://doi.org/10.1038/3305

Fagervall, I., & Ross, S. B. (1986). Inhibition of mono amine oxidase in monoaminergic neurones in the rat brain by irreversible inhibitors. *Biochemical Pharmacology*, *35*(8), 1381–1387. https://doi.org/10.1016/0006-2952(86)90285-

Fan, N., Luo, Y., Xu, K., Zhang, M., Ke, X., Huang, X., Ding, Y., Wang, D., Ning, Y., Deng, X., & He, H. (2015). Relationship of serum levels of TNF-α, IL-6 and IL-18 and schizophrenia-like symptoms in chronic ketamine abusers. *Schizophrenia Research*, *169*(1–3), 10–15. https://doi.org/10.1016/j.schres.2015.11.006

Feduccia, A. A., & Mithoefer, M. C. (2018). MDMA-assisted psychotherapy for PTSD: Are memory reconsolidation and fear extinction underlying mechanisms? In *Progress in Neuro-Psychopharmacology and Biological Psychiatry*. https://doi.org/10.1016/j.pnpbp.2018.03.003

Ferber, S. G., Namdar, D., Hen-Shoval, D., Eger, G., Koltai, H., Shoval, G., Shbiro, L., & Weller, A. (2019). The "Entourage Effect": Terpenes Coupled with Cannabinoids for the Treatment of Mood Disorders and Anxiety Disorders. *Current Neuropharmacology*, *18*(2), 87–96. https://doi.org/10.2174/1570159x17666190903103923

Ferraz, C. A. A., de Oliveira Júnior, R. G., Picot, L., da Silva Almeida, J. R. G., & Nunes, X. P. (2019). Pre-clinical investigations of β-carboline alkaloids as antidepressant agents: A systematic review. *Fitoterapia*, *137*(March). https://doi.org/10.1016/j.fitote.2019.104196

Fortunato, J. J., Réus, G. Z., Kirsch, T. R., Stringari, R. B., Fries, G. R., Kapczinski, F., Hallak, J. E., Zuardi, A. W., Crippa, J. A., & Quevedo, J. (2010a). Chronic administration of harmine elicits antidepressant-like effects and increases BDNF levels in rat hippocampus. *Journal of Neural Transmission*, *117*(10), 1131–1137. https://doi.org/10.1007/s00702-010-0451-2

Fortunato, J. J., Réus, G. Z., Kirsch, T. R., Stringari, R. B., Fries, G. R., Kapczinski, F., Hallak, J. E., Zuardi, A. W., Crippa, J. A., & Quevedo, J. (2010b). Effects of β-carboline harmine on behavioral and physiological parameters observed in the chronic mild stress model: Further evidence of antidepressant properties. *Brain Research Bulletin*, *81*(4–5), 491–496. https://doi.org/10.1016/j.brainresbull.2009.09.008

Fortunato, J. J., Réus, G. Z., Kirsch, T. R., Stringari, R. B., Stertz, L., Kapczinski, F., Pinto, J. P., Hallak, J. E., Zuardi, A. W., Crippa, J. A., & Quevedo, J. (2009). Acute harmine administration induces antidepressive-like effects and increases BDNF levels in the rat hippocampus. *Progress in Neuro-Psychopharmacology and Biological Psychiatry*, *33*(8), 1425–1430. https://doi.org/10.1016/j.pnpbp.2009.07.021

Frecska, E., Bokor, P., & Winkelman, M. (2016). The Therapeutic Potentials of Ayahuasca: Possible Effects against Various Diseases of Civilization. *Frontiers in Pharmacology*, *7*(March), 1–17. https://doi.org/10.3389/fphar.2016.00035

Frecska, E., Szabo, A., Winkelman, M. J., Luna, L. E., & McKenna, D. J. (2013). A possibly sigma-1 receptor mediated role of dimethyltryptamine in tissue protection, regeneration, and immunity. *Journal of Neural Transmission (Vienna, Austria : 1996)*, *120*(9), 1295–1303. https://doi.org/10.1007/s00702-013-1024-y

Freis, E. D. (1954). Mental Depression in Hypertensive Patients Treated for Long Periods with Large Doses of Reserpine. *New England Journal of Medicine*. https://doi.org/10.1056/nejm195412162512504

Garani, R., Watts, J. J., & Mizrahi, R. (2021). Endocannabinoid system in psychotic and mood disorders, a review of human studies. *Progress in Neuro-Psychopharmacology and Biological Psychiatry*, *106*(August 2020), 110096. https://doi.org/10.1016/j.pnpbp.2020.110096

García-Cabrerizo, R., & García-Fuster, M. J. (2016). Comparative effects of amphetamine-like psychostimulants on rat hippocampal cell genesis at different developmental ages. *NeuroToxicology*, *56*, 29–39. https://doi.org/10.1016/j.neuro.2016.06.014

Geyer, M. A., Nichols, D. E., & Vollenweider, F. X. (2009). Serotonin-Related Psychedelic Drugs. In *Encyclopedia of Neuroscience* (pp. 731–738). Elsevier. https://doi.org/10.1016/B978-008045046-9.01160-8

Goulart, B. K., de Lima, M. N. M., de Farias, C. B., Reolon, G. K., Almeida, V. R., Quevedo, J., Kapczinski, F., Schröder, N., & Roesler, R. (2010). Ketamine impairs recognition memory consolidation and prevents learning-induced increase in hippocampal brain-derived neurotrophic factor levels. *Neuroscience*, *167*(4), 969–973. https://doi.org/10.1016/j.neuroscience.2010.03.032

Haile, C. N., Murrough, J. W., Iosifescu, D. V., Chang, L. C., Al Jurdi, R. K., Foulkes, A., Iqbal, S., Mahoney, J. J., De La Garza, R., Charney, D. S., Newton, T. F., & Mathew, S. J. (2014). Plasma brain derived neurotrophic factor (BDNF) and response to ketamine in treatment-resistant depression. *International Journal of Neuropsychopharmacology*, *17*(2), 331–336. https://doi.org/10.1017/S1461145713001119

Hartogsohn, I. (2017). Constructing drug effects: A history of set and setting. *Drug Science, Policy and Law*. https://doi.org/10.1177/2050324516683325

Hashimoto, K. (2019). Rapid-acting antidepressant ketamine, its metabolites and other candidates: A historical overview and future perspective. *Psychiatry and Clinical Neurosciences*, *73*(10), 613–627. https://doi.org/10.1111/pcn.12902

Healy, D. (2004). *Let them eat PROZAC The Unhealthy Relationship Between the Pharmaceutical Industry and*

Depression. NYU Press.

Hearn, K. (2013). The Dark Side of Ayahuasca. In *Men's Journal*.

Heninger, G., Delgado, P., & Charney, D. (1996). The Revised Monoamine Theory of Depression: A Modulatory Role for Monoamines, Based on New Findings From Monoamine Depletion Experiments in Humans. *Pharmacopsychiatry*, *29*(01), 2–11. https://doi.org/10.1055/s-2007-979535

Hernández-Rabaza, V., Domínguez-Escribà, L., Barcia, J. A., Rosel, J. F., Romero, F. J., García-Verdugo, J. M., & Canales, J. J. (2006). Binge administration of 3,4-methylenedioxymethamphetamine ("ecstasy") impairs the survival of neural precursors in adult rat dentate gyrus. *Neuropharmacology*, *51*(5), 967–973. https://doi.org/10.1016/j.neuropharm.2006.06.019

Herraiz, T., González, D., Ancín-Azpilicueta, C., Arán, V. J., & Guillén, H. (2010). β-Carboline alkaloids in Peganum harmala and inhibition of human monoamine oxidase (MAO). *Food and Chemical Toxicology*, *48*(3), 839–845. https://doi.org/10.1016/j.fct.2009.12.019

Hoefle, M. L., & Warner-Lambert, P. D. (2001). The early history of Parke-Davis and Company. *Bulletin for the History of Chemestry*, *25*(1), 28.

Holman, C. L. (2010). Spirituality for sale an analysis of ayahuasca tourism. In *Dissertation Abstracts International Section A: Humanities and Social Sciences*.

Hooijmans, C. R., Rovers, M. M., de Vries, R. B., Leenaars, M., Ritskes-Hoitinga, M., & Langendam, M. W. (2014). SYRCLE's risk of bias tool for animal studies. *BMC Medical Research Methodology*, *14*(1), 43. https://doi.org/10.1186/1471-2288-14-43

Huang, H., Liu, C.-M., Sun, J., Hao, T., Xu, C.-M., Wang, D., & Wu, Y.-Q. (2016). Ketamine Affects the Neurogenesis of the Hippocampal Dentate Gyrus in 7-Day-Old Rats. *Neurotoxicity Research*, *30*(2), 185–198. https://doi.org/10.1007/s12640-016-9615-7

Idell, R. D., Florova, G., Komissarov, A. A., Shetty, S., Girard, R. B. S., & Idell, S. (2017). The fibrinolytic system: A new target for treatment of depression with psychedelics. *Medical Hypotheses*, *100*, 46–53. https://doi.org/10.1016/j.mehy.2017.01.013

Ioannidis, J. P. A. (2014). How to Make More Published Research True. *PLoS Medicine*. https://doi.org/10.1371/journal.pmed.1001747

Jacobs, B. L., van Praag, H., & Gage, F. H. (2000). Adult brain neurogenesis and psychiatry: a novel theory of depression. *Molecular Psychiatry*, *5*(3), 262–269. https://doi.org/10.1038/sj.mp.4000712

James, S. L., Abate, D., Abate, K. H., Abay, S. M., Abbafati, C., Abbasi, N., Abbastabar, H., Abd-Allah, F., Abdela, J., Abdelalim, A., Abdollahpour, I., Abdulkader, R. S., Abebe, Z., Abera, S. F., Abil, O. Z., Abraha, H. N., Abu-Raddad, L. J., Abu-Rmeileh, N. M. E., Accrombessi, M. M. K., … Murray, C. J. L. (2018). Global, regional, and national incidence, prevalence, and years lived with disability for 354 diseases and injuries for 195 countries and territories, 1990–2017: a systematic analysis for the Global Burden of Disease Study 2017. *The Lancet*, *392*(10159), 1789–1858. https://doi.org/10.1016/S0140-6736(18)32279-7

Javitt, D. C., & Zukin, S. R. (1991). Recent advances in the phencyclidine model of schizophrenia. *The American Journal of Psychiatry*, *148*(10), 1301–1308. https://doi.org/10.1176/ajp.148.10.1301

Jha, S., Rajendran, R., Fernandes, K. A., & Vaidya, V. A. (2008). 5-HT2A/2C receptor blockade regulates progenitor cell proliferation in the adult rat hippocampus. *Neuroscience Letters*, *441*(2), 210–214. https://doi.org/10.1016/j.neulet.2008.06.028

Jiang, W., Zhang, Y., Xiao, L., Van Cleemput, J., Ji, S. P., Bai, G., & Zhang, X. (2005). Cannabinoids promote embryonic and adult hippocampus neurogenesis and produce anxiolytic- and antidepressant-like effects. *Journal of Clinical Investigation*, *115*(11), 3104–3116. https://doi.org/10.1172/JCI25509

Jones, K. A., Srivastava, D. P., Allen, J. A., Strachan, R. T., Roth, B. L., & Penzes, P. (2009). Rapid modulation of spine morphology by the 5-HT2A serotonin receptor through kalirin-7 signaling. *Proceedings of the National Academy of Sciences of the United States of America*, *106*(46), 19575–19580. https://doi.org/10.1073/pnas.0905884106

Journey, J. D., & Bentley, T. P. (2021). *Phencyclidine Toxicity.* StatPearls.

Kalant, H. (2001). The pharmacology and toxicology of "ecstasy" (MDMA) and related drugs. *CMAJ : Canadian Medical Association Journal = Journal de l'Association Medicale Canadienne*, *165*(7), 917–928.

Kaplan, M., & Hinds, J. (1977). Neurogenesis in the adult rat: electron microscopic analysis of light radioautographs. *Science*, *197*(4308), 1092–1094. https://doi.org/10.1126/science.887941

Karila, L., Roux, P., Rolland, B., Benyamina, A., Reynaud, M., Aubin, H.-J., & Lancon, C. (2014). Acute and Long-Term Effects of Cannabis Use: A Review. *Current Pharmaceutical Design*, *20*(25), 4112–4118.

https://doi.org/10.2174/13816128113199990620

Katchborian-Neto, A., Santos, W. T., Nicácio, K. J., Corrêa, J. O. A., Murgu, M., Martins, T. M. M., Gomes, D. A., Goes, A. M., Soares, M. G., Dias, D. F., Chagas-Paula, D. A., & Paula, A. C. C. (2020). Neuroprotective potential of Ayahuasca and untargeted metabolomics analyses: applicability to Parkinson's disease. *Journal of Ethnopharmacology*, *255*, 112743. https://doi.org/10.1016/j.jep.2020.112743

Keilhoff, G., Bernstein, H. G., Becker, A., Grecksch, G., & Wolf, G. (2004). Increased neurogenesis in a rat ketamine model of schizophrenia. *Biological Psychiatry*, *56*(5), 317–322. https://doi.org/10.1016/j.biopsych.2004.06.010

Kempermann, G. (2002). Regulation of adult hippocampal neurogenesis - implications for novel theories of major depression 1. *Bipolar Disorders*, *4*(1), 17–33. https://doi.org/10.1034/j.1399-5618.2002.40101.x

Kempermann, G. (2012). New neurons for "survival of the fittest." *Nature Reviews Neuroscience*, *13*(10), 727–736. https://doi.org/10.1038/nrn3319

Kochman, L. J., dos Santos, A. A., Fornal, C. A., & Jacobs, B. L. (2006). Despite strong behavioral disruption, Δ9-tetrahydrocannabinol does not affect cell proliferation in the adult mouse dentate gyrus. *Brain Research*, *1113*(1), 86–93. https://doi.org/10.1016/j.brainres.2006.07.080

Koornneef, M., Hanhart, C. J., Veen, J. H. Van Der, Josephs, E., Lockwood, A., Myllylä, S., Oakley, C., Palmer, R., Fontanilla, D., Johannessen, M., Hajipour, A. R., Cozzi, N. V, Jackson, M. B., & Ruoho, A. E. (2009). *The Hallucinogen N,N-Dimethyltryptamine (DMT) Is an Endogenous Sigma-1 Receptor Regulator. February*, 934–937.

Koprich, J. B., Chen, E.-Y., Kanaan, N. M., Campbell, N. G., Kordower, J. H., & Lipton, J. W. (2003). Prenatal 3,4-methylenedioxymethamphetamine (ecstasy) alters exploratory behavior, reduces monoamine metabolism, and increases forebrain tyrosine hydroxylase fiber density of juvenile rats. *Neurotoxicology and Teratology*, *25*(5), 509–517. https://doi.org/10.1016/S0892-0362(03)00091-6

Kuhn, H. G., Dickinson-Anson, H., & Gage, F. H. (1996). Neurogenesis in the dentate gyrus of the adult rat: age-related decrease of neuronal progenitor proliferation. *The Journal of Neuroscience : The Official Journal of the Society for Neuroscience*.

Labate, B. C., & Cavnar, C. (2013). The therapeutic use of Ayahuasca. In *The Therapeutic Use of Ayahuasca*. https://doi.org/10.1007/978-3-642-40426-9

Labate, B. C., & Cavnar, C. (2014). Ayahuasca Shamanism in the Amazon and Beyond. In B. C. Labate & C. Cavnar (Eds.), *Ayahuasca Shamanism in the Amazon and Beyond*. Oxford University Press. https://doi.org/10.1093/acprof:oso/9780199341191.001.0001

Lacasse, J. R., & Leo, J. (2005). Serotonin and Depression: A Disconnect between the Advertisements and the Scientific Literature. *PLoS Medicine*, *2*(12), e392. https://doi.org/10.1371/journal.pmed.0020392

Laplagne, D. A., Espósito, M. S., Piatti, V. C., Morgenstern, N. A., Zhao, C., Van Praag, H., Gage, F. H., & Schinder, A. F. (2006). Functional convergence of neurons generated in the developing and adult hippocampus. *PLoS Biology*, *4*(12), 2349–2360. https://doi.org/10.1371/journal.pbio.0040409

Lee, T. T.-Y., Wainwright, S. R., Hill, M. N., Galea, L. A. M., & Gorzalka, B. B. (2014). Sex, drugs, and adult neurogenesis: Sex-dependent effects of escalating adolescent cannabinoid exposure on adult hippocampal neurogenesis, stress reactivity, and amphetamine sensitization. *Hippocampus*, *24*(3), 280–292. https://doi.org/10.1002/hipo.22221

Leishman, E., Murphy, M., Mackie, K., & Bradshaw, H. B. (2018). Δ 9 -Tetrahydrocannabinol changes the brain lipidome and transcriptome differentially in the adolescent and the adult. *Biochimica et Biophysica Acta (BBA) - Molecular and Cell Biology of Lipids*, *1863*(5), 479–492. https://doi.org/10.1016/j.bbalip.2018.02.001

Lepack, A. E., Fuchikami, M., Dwyer, J. M., Banasr, M., & Duman, R. S. (2015). BDNF Release Is Required for the Behavioral Actions of Ketamine. *International Journal of Neuropsychopharmacology*, *18*(1), pyu033–pyu033. https://doi.org/10.1093/ijnp/pyu033

Lépine, J. P., & Briley, M. (2011). The increasing burden of depression. *Neuropsychiatric Disease and Treatment*. https://doi.org/10.2147/NDT.S19617

Li, N., Lee, B., Liu, R.-J., Banasr, M., Dwyer, J. M., Iwata, M., Li, X.-Y., Aghajanian, G., & Duman, R. S. (2010). mTOR-Dependent Synapse Formation Underlies the Rapid Antidepressant Effects of NMDA Antagonists. *Science*, *329*(5994), 959–964. https://doi.org/10.1126/science.1190287

Li, N. X., Hu, Y. R., Chen, W. N., & Zhang, B. (2022). Dose effect of psilocybin on primary and secondary depression: a preliminary systematic review and meta-analysis. *Journal of Affective Disorders*, *296*(June 2021), 26–34. https://doi.org/10.1016/j.jad.2021.09.041

Liberati, A., Tetzlaff, J., & Altman, D. G. (2009). Preferred Reporting Items for Systematic Reviews and Meta-Analyses:

The PRISMA Statement. *PLoS Medicine*, *6*(7), e1000097. https://doi.org/10.1371/journal.pmed.1000097

Lima da Cruz, R. V., Moulin, T. C., Petiz, L. L., & Leão, R. N. (2018). A Single Dose of 5-MeO-DMT Stimulates Cell Proliferation, Neuronal Survivability, Morphological and Functional Changes in Adult Mice Ventral Dentate Gyrus. *Frontiers in Molecular Neuroscience*, *11*. https://doi.org/10.3389/fnmol.2018.00312

Liu, F., Wu, J., Gong, Y., Wang, P., Zhu, L., Tong, L., Chen, X., Ling, Y., & Huang, C. (2017). Harmine produces antidepressant-like effects via restoration of astrocytic functions. *Progress in Neuro-Psychopharmacology and Biological Psychiatry*, *79*(June), 258–267. https://doi.org/10.1016/j.pnpbp.2017.06.012

Liu, J., Suzuki, T., Seki, T., Namba, T., Tanimura, A., & Arai, H. (2006). Effects of repeated phencyclidine administration on adult hippocampal neurogenesis in the rat. *Synapse*, *60*(1), 56–68. https://doi.org/10.1002/syn.20275

Liu, R.-J., Fuchikami, M., Dwyer, J. M., Lepack, A. E., Duman, R. S., & Aghajanian, G. K. (2013). GSK-3 Inhibition Potentiates the Synaptogenic and Antidepressant-Like Effects of Subthreshold Doses of Ketamine. *Neuropsychopharmacology*, *38*(11), 2268–2277. https://doi.org/10.1038/npp.2013.128

Lopez-Munoz, F., & Alamo, C. (2009). Monoaminergic Neurotransmission: The History of the Discovery of Antidepressants from 1950s Until Today. *Current Pharmaceutical Design*, *15*(14), 1563–1586. https://doi.org/10.2174/138161209788168001

Ly, C., Greb, A. C., Cameron, L. P., Wong, J. M., Barragan, E. V., Wilson, P. C., Burbach, K. F., Soltanzadeh Zarandi, S., Sood, A., Paddy, M. R., Duim, W. C., Dennis, M. Y., McAllister, A. K., Ori-McKenney, K. M., Gray, J. A., & Olson, D. E. (2018). Psychedelics Promote Structural and Functional Neural Plasticity. *Cell Reports*, *23*(11), 3170–3182. https://doi.org/10.1016/j.celrep.2018.05.022

Maeda, K., Sugino, H., Hirose, T., Kitagawa, H., Nagai, T., Mizoguchi, H., Takuma, K., & Yamada, K. (2007). Clozapine prevents a decrease in neurogenesis in mice repeatedly treated with phencyclidine. *Journal of Pharmacological Sciences*, *103*(3), 299–308. https://doi.org/10.1254/jphs.FP0061424

Magaraggia, I., Kuiperes, Z., & Schreiber, R. (2021). Improving cognitive functioning in major depressive disorder with psychedelics: A dimensional approach. *Neurobiology of Learning and Memory*, *183*(May), 107467. https://doi.org/10.1016/j.nlm.2021.107467

Marinova, Z., Walitza, S., & Grünblatt, E. (2017). The hallucinogen 2,5-dimethoxy-4-iodoamphetamine hydrochloride activates neurotrophin receptors in a neuronal cell line and promotes neurites extension. *Journal of Neural Transmission*, *124*(6), 749–759. https://doi.org/10.1007/s00702-017-1706-y

Martin, B. R., Mechoulam, R., & Razdan, R. K. (1999). Discovery and characterization of endogenous cannabinoids. *Life Sciences*, *65*(6–7), 573–595. https://doi.org/10.1016/S0024-3205(99)00281-7

Marton, S., González, B., Rodríguez-Bottero, S., Miquel, E., Martínez-Palma, L., Pazos, M., Pedro Prieto, J., Rodríguez, P., Sames, D., Seoane, G., Scorza, C., Cassina, P., & Carrera, I. (2019). Ibogaine administration modifies GDNF and BDNF expression in brain regions involved in mesocorticolimbic and nigral dopaminergic circuits. *Frontiers in Pharmacology*, *10*(MAR), 193. https://doi.org/10.3389/fphar.2019.00193

Mathai, D. S., Meyer, M. J., Storch, E. A., & Kosten, T. R. (2020). The relationship between subjective effects induced by a single dose of ketamine and treatment response in patients with major depressive disorder: A systematic review. *Journal of Affective Disorders*. https://doi.org/10.1016/j.jad.2019.12.023

Matheson, J., Sproule, B., Di Ciano, P., Fares, A., Le Foll, B., Mann, R. E., & Brands, B. (2020). Sex differences in the acute effects of smoked cannabis: evidence from a human laboratory study of young adults. *Psychopharmacology*, *237*(2), 305–316. https://doi.org/10.1007/s00213-019-05369-y

Matteau-Pelletier, L., Bélanger, R. E., Leatherdale, S., Desbiens, F., & Haddad, S. (2020). Sex-Related Differences in Adolescent Cannabis Use: Influences of School Context and School Connectedness. *The Journal of School Health*, *90*(11), 878–886. https://doi.org/10.1111/josh.12950

McKenna, D. J., Towers, G. H. N., & Abbott, F. (1984). Monoamine oxidase inhibitors in South American hallucinogenic plants: Tryptamine and β-carboline constituents of Ayahuasca. *Journal of Ethnopharmacology*, *10*(2), 195–223. https://doi.org/10.1016/0378-8741(84)90003-5

Mechoulam, R., Feigenbaum, J. J., Lander, N., Segal, M., Järbe, T. U. C., Hiltunen, A. J., & Consroe, P. (1988). Enantiomeric cannabinoids: stereospecificity of psychotropic activity. *Experientia*, *44*(9), 762–764. https://doi.org/10.1007/BF01959156

Mechoulam, Raphael, & Parker, L. A. (2013). The endocannabinoid system and the brain. *Annual Review of Psychology*, *64*(1), 21–47. https://doi.org/10.1146/annurev-psych-113011-143739

Miller, B. R., & Hen, R. (2015). The Current State of the Neurogenic Theory of Depression and Anxiety. *Current Opinions in Neurobiology*, *0*(February), 51–58. https://doi.org/10.1016/j.conb.2014.08.012.THE

Mithoefer, M. C., Feduccia, A. A., Jerome, L., Mithoefer, A., Wagner, M., Walsh, Z., Hamilton, S., Yazar-Klosinski, B., Emerson, A., & Doblin, R. (2019). MDMA-assisted psychotherapy for treatment of PTSD: study design and rationale for phase 3 trials based on pooled analysis of six phase 2 randomized controlled trials. *Psychopharmacology*, *236*(9), 2735–2745. https://doi.org/10.1007/s00213-019-05249-5

Morales-Garcia, J. A., Calleja-Conde, J., Lopez-Moreno, J. A., Alonso-Gil, S., Sanz-SanCristobal, M., Riba, J., & Perez-Castillo, A. (2020). N,N-dimethyltryptamine compound found in the hallucinogenic tea ayahuasca, regulates adult neurogenesis in vitro and in vivo. *Translational Psychiatry*, *10*(1). https://doi.org/10.1038/s41398-020-01011-0

Morales-García, J. A., De La Fuente Revenga, M., Alonso-Gil, S., Rodríguez-Franco, M. I., Feilding, A., Perez-Castillo, A., & Riba, J. (2017). The alkaloids of Banisteriopsis caapi, the plant source of the Amazonian hallucinogen Ayahuasca, stimulate adult neurogenesis in vitro. *Scientific Reports*, *7*(1), 1–13. https://doi.org/10.1038/s41598-017-05407-9

Muscat, S.-A., Hartelius, G., Crouch, C. R., & Morin, K. W. (2021). An Integrative Approach to Ketamine Therapy May Enhance Multiple Dimensions of Efficacy: Improving Therapeutic Outcomes With Treatment Resistant Depression. *Frontiers in Psychiatry*, *12*(November), 1–20. https://doi.org/10.3389/fpsyt.2021.710338

Muttoni, S., Ardissino, M., & John, C. (2019). Classical psychedelics for the treatment of depression and anxiety: A systematic review. *Journal of Affective Disorders*, *258*(April), 11–24. https://doi.org/10.1016/j.jad.2019.07.076

Nagai, F., Nonaka, R., & Satoh Hisashi Kamimura, K. (2007). The effects of non-medically used psychoactive drugs on monoamine neurotransmission in rat brain. *European Journal of Pharmacology*, *559*(2–3), 132–137. https://doi.org/10.1016/j.ejphar.2006.11.075

Naqvi, S., Godfrey, A. K., Hughes, J. F., Goodheart, M. L., Mitchell, R. N., & Page, D. C. (2019). Conservation, acquisition, and functional impact of sex-biased gene expression in mammals. *Science*, *365*(6450), eaaw7317. https://doi.org/10.1126/science.aaw7317

Naranjo, P. (1979). Hallucinogenic plant use and related indigenous belief systems in the ecuadorian amazon. *Journal of Ethnopharmacology*, *1*(2), 121–145. https://doi.org/10.1016/0378-8741(79)90003-5

Nestler, E. J., Barrot, M., DiLeone, R. J., Eisch, A. J., Gold, S. J., & Monteggia, L. M. (2002). Neurobiology of depression. In *Neuron* (Vol. 34, Issue 1, pp. 13–25). https://doi.org/10.1016/S0896-6273(02)00653-0

Nichols, D. E. (1986). Differences Between the Mechanism of Action of MDMA, MBDB, and the Classic Hallucinogens. Identification of a New Therapeutic Class: Entactogens. *Journal of Psychoactive Drugs*, *18*(4), 305–313. https://doi.org/10.1080/02791072.1986.10472362

Nichols, D. E. (2004). Hallucinogens. *Pharmacology & Therapeutics*, *101*(2), 131–181. https://doi.org/10.1016/j.pharmthera.2003.11.002

Nichols, D. E. (2016). Psychedelics. *Pharmacological Reviews*, *68*(2), 264–355. https://doi.org/10.1124/pr.115.011478

Ninnemann, A., Stuart, G. L., & Andersen, S. M. (2012). Convention on Psychotropic Substances, 1971. In *Encyclopedia of Drug Policy*. SAGE Publications, Inc. https://doi.org/10.4135/9781412976961.n78

Numakawa, T., Suzuki, S., Kumamaru, E., Adachi, N., Richards, M., & Kunugi, H. (2010). BDNF function and intracellular signaling in neurons. *Histology and Histopathology*, *25*(2), 237–258. https://doi.org/10.14670/HH-25.237

OHAT. (2015). *OHAT Risk of Bias Rating Tool for Human and Animal Studies Organization of This Document Indirectness , Timing , and Other Factors Related to Risk of Bias*. January, 1–37.

Olson, D. E. (2018). Psychoplastogens: A Promising Class of Plasticity-Promoting Neurotherapeutics. *Journal of Experimental Neuroscience*, *12*, 117906951880050. https://doi.org/10.1177/1179069518800508

Onaivi, E. S. (2006). Neuropsychobiological Evidence for the Functional Presence and Expression of Cannabinoid CB2 Receptors in the Brain. *Neuropsychobiology*, *54*(4), 231–246. https://doi.org/10.1159/000100778

Osmond, H. (1957). A REVIEW OF THE CLINICAL EFFECTS OF PSYCHOTOMIMETIC AGENTS. *Annals of the New York Academy of Sciences*, *66*(3), 418–434. https://doi.org/10.1111/j.1749-6632.1957.tb40738.x

Papaseit, E., Pérez-Mañá, C., Torrens, M., Farré, A., Poyatos, L., Hladun, O., Sanvisens, A., Muga, R., & Farré, M. (2020). MDMA interactions with pharmaceuticals and drugs of abuse. *Expert Opinion on Drug Metabolism & Toxicology*, *16*(5), 357–369. https://doi.org/10.1080/17425255.2020.1749262

Peacock, A., Leung, J., Larney, S., Colledge, S., Hickman, M., Rehm, J., Giovino, G. A., West, R., Hall, W., Griffiths, P., Ali, R., Gowing, L., Marsden, J., Ferrari, A. J., Grebely, J., Farrell, M., & Degenhardt, L. (2018). Global statistics on alcohol, tobacco and illicit drug use: 2017 status report. *Addiction*, *113*(10), 1905–1926. https://doi.org/10.1111/add.14234

Pertwee, R. G. (2006). The pharmacology of cannabinoid receptors and their ligands: An overview. *International Journal*

of Obesity, *30*, S13–S18. https://doi.org/10.1038/sj.ijo.0803272

Pochwat, B., Pałucha-Poniewiera, A., Szewczyk, B., Pilc, A., & Nowak, G. (2014). NMDA antagonists under investigation for the treatment of major depressive disorder. *Expert Opinion on Investigational Drugs*, *23*(9), 1181–1192. https://doi.org/10.1517/13543784.2014.918951

Poulia, N., Delis, F., Brakatselos, C., Polissidis, A., Koutmani, Y., Kokras, N., Dalla, C., Politis, P. K., & Antoniou, K. (2021). Detrimental effects of adolescent escalating low-dose Δ9-tetrahydrocannabinol leads to a specific bio-behavioural profile in adult male rats. *British Journal of Pharmacology*, *178*(7), 1722–1736. https://doi.org/10.1111/bph.15394

Psychoyos, D., Hungund, B., Cooper, T., & Finnell, R. H. (2008). A cannabinoid analogue of Δ9-tetrahydrocannabinol disrupts neural development in chick. *Birth Defects Research Part B - Developmental and Reproductive Toxicology*, *83*(5), 477–488. https://doi.org/10.1002/bdrb.20166

Realini, N., Vigano', D., Guidali, C., Zamberletti, E., Rubino, T., & Parolaro, D. (2011). Chronic URB597 treatment at adulthood reverted most depressive-like symptoms induced by adolescent exposure to THC in female rats. *Neuropharmacology*, *60*(2–3), 235–243. https://doi.org/10.1016/j.neuropharm.2010.09.003

Reiche, S., Hermle, L., Gutwinski, S., Jungaberle, H., Gasser, P., & Majić, T. (2018). Serotonergic hallucinogens in the treatment of anxiety and depression in patients suffering from a life-threatening disease: A systematic review. *Progress in Neuro-Psychopharmacology & Biological Psychiatry*, *81*, 1–10. https://doi.org/10.1016/j.pnpbp.2017.09.012

Ren, G., Zhang, X., Li, Y., Ridout, K., Serrano-Serrano, M. L., Yang, Y., Liu, A., Ravikanth, G., Nawaz, M. A., Mumtaz, A. S., Salamin, N., & Fumagalli, L. (2021). Large-scale whole-genome resequencing unravels the domestication history of Cannabis sativa. *Science Advances*, *7*(29), eabg2286. https://doi.org/10.1126/sciadv.abg2286

Riga, M. S., Bortolozzi, A., Campa, L., Artigas, F., & Celada, P. (2016). The serotonergic hallucinogen 5-methoxy-N,N-dimethyltryptamine disrupts cortical activity in a regionally-selective manner via 5-HT 1A and 5-HT 2A receptors. *Neuropharmacology*, *101*, 370–378. https://doi.org/10.1016/j.neuropharm.2015.10.016

Rivier, L., & Lindgren, J.-E. (1972). "Ayahuasca," the South American Hallucinogenic Drink: An Ethnobotanical and Chemical Investigation. *Economic Botany*, *26*(2), 101–129.

Robinson, D. S. (1973). The Monoamine Oxidase Inhibitor, Phenelzine, in the Treatment of Depressive-Anxiety States. *Archives of General Psychiatry*, *29*(3), 407. https://doi.org/10.1001/archpsyc.1973.04200030093015

Rodriguez Bambico, F., Duranti, A., Tontini, A., Tarzia, G., & Gobbi, G. (2009). Endocannabinoids in the Treatment of Mood Disorders: Evidence from Animal Models. *Current Pharmaceutical Design*, *15*(14), 1623–1646. https://doi.org/10.2174/138161209788168029

Rubino, T., Vigano', D., Realini, N., Guidali, C., Braida, D., Capurro, V., Castiglioni, C., Cherubino, F., Romualdi, P., Candeletti, S., Sala, M., & Parolaro, D. (2008). Chronic Δ9-Tetrahydrocannabinol During Adolescence Provokes Sex-Dependent Changes in the Emotional Profile in Adult Rats: Behavioral and Biochemical Correlates. *Neuropsychopharmacology*, *33*(11), 2760–2771. https://doi.org/10.1038/sj.npp.1301664

Rucker, J. J. H., Iliff, J., & Nutt, D. J. (2018). Psychiatry & the psychedelic drugs. Past, present & future. *Neuropharmacology*, *142*, 200–218. https://doi.org/10.1016/j.neuropharm.2017.12.040

Rueda, D., Navarro, B., Martínez-Serrano, A., Guzmán, M., & Galve-Roperh, I. (2002). The endocannabinoid anandamide inhibits neuronal progenitor cell differentiation through attenuation of the Rap1/B-Raf/ERK pathway. *Journal of Biological Chemistry*, *277*(48), 46645–46650. https://doi.org/10.1074/jbc.M206590200

Ruiz, P., & Strain, E. (2011). Substance Abuse: A Comprehensive Textbook. In Lowinson & P. Ruiz (Eds.), *Wolters Kluwer*. Wolters Kluwer.

Rybakowski, J. K., Permoda-Osip, A., Skibinska, M., Adamski, R., & Bartkowska-Sniatkowska, A. (2013). Single ketamine infusion in bipolar depression resistant to antidepressants: Are neurotrophins involved? *Human Psychopharmacology*, *28*(1), 87–90. https://doi.org/10.1002/hup.2271

Saengsawang, W., & Rasenick, M. M. (2016). G Protein-Coupled Receptors. In *Encyclopedia of Cell Biology* (pp. 51–55). Elsevier. https://doi.org/10.1016/B978-0-12-394447-4.30004-9

Sanches, M., Quevedo, J., & Soares, J. C. (2021). New agents and perspectives in the pharmacological treatment of major depressive disorder. In *Progress in Neuro-Psychopharmacology and Biological Psychiatry* (Vol. 106, p. 9). https://doi.org/10.1016/j.pnpbp.2020.110157

Sartori, S. B., & Singewald, N. (2019). Novel pharmacological targets in drug development for the treatment of anxiety and anxiety-related disorders. *Pharmacology and Therapeutics*, *204*. https://doi.org/10.1016/j.pharmthera.2019.107402

Schaefer, T. L., Grace, C. E., Braun, A. A., Amos-Kroohs, R. M., Graham, D. L., Skelton, M. R., Williams, M. T., & Vorhees, C. V. (2013). Cognitive impairments from developmental exposure to serotonergic drugs: Citalopram and MDMA. *International Journal of Neuropsychopharmacology*, *16*(6), 1383–1394. https://doi.org/10.1017/S1461145712001447

Schiavone, S., Morgese, M. G., Bove, M., Colia, A. L., Maffione, A. B., Tucci, P., Trabace, L., & Cuomo, V. (2020). Ketamine administration induces early and persistent neurochemical imbalance and altered NADPH oxidase in mice. *Progress in Neuro-Psychopharmacology and Biological Psychiatry*, *96*(January 2019). https://doi.org/10.1016/j.pnpbp.2019.109750

Sessa, B. (2017). MDMA and PTSD treatment: "PTSD: From novel pathophysiology to innovative therapeutics." In *Neuroscience Letters*. https://doi.org/10.1016/j.neulet.2016.07.004

Shum, C., Dutan, L., Annuario, E., Warre-Cornish, K., Taylor, S. E., Taylor, R. D., Andreae, L. C., Buckley, N. J., Price, J., Bhattacharyya, S., & Srivastava, D. P. (2020). Δ9-tetrahydrocannabinol and 2-AG decreases neurite outgrowth and differentially affects ERK1/2 and Akt signaling in hiPSC-derived cortical neurons. *Molecular and Cellular Neuroscience*, *103*, 103463. https://doi.org/10.1016/j.mcn.2019.103463

Sial, O. K., Parise, E. M., Parise, L. F., Gnecco, T., & Bolaños-Guzmán, C. A. (2020). Ketamine: The final frontier or another depressing end? In *Behavioural Brain Research* (Vol. 383, p. 18). https://doi.org/10.1016/j.bbr.2020.112508

Skaper, S. D., & Di Marzo, V. (2012). Endocannabinoids in nervous system health and disease: The big picture in a nutshell. *Philosophical Transactions of the Royal Society B: Biological Sciences*, *367*(1607), 3193–3200. https://doi.org/10.1098/rstb.2012.0313

Sorrells, S. F., Paredes, M. F., Cebrian-Silla, A., Sandoval, K., Qi, D., Kelley, K. W., James, D., Mayer, S., Chang, J., Auguste, K. I., Chang, E. F., Gutierrez, A. J., Kriegstein, A. R., Mathern, G. W., Oldham, M. C., Huang, E. J., Garcia-Verdugo, J. M., Yang, Z., & Alvarez-Buylla, A. (2018). Human hippocampal neurogenesis drops sharply in children to undetectable levels in adults. *Nature*. https://doi.org/10.1038/nature25975

Stanslowsky, N., Jahn, K., Venneri, A., Naujock, M., Haase, A., Martin, U., Frieling, H., & Wegner, F. (2017). Functional effects of cannabinoids during dopaminergic specification of human neural precursors derived from induced pluripotent stem cells. *Addiction Biology*, *22*(5), 1329–1342. https://doi.org/10.1111/adb.12394

Steel, R. W. J., Miller, J. H., Sim, D. A., & Day, D. J. (2014). Delta-9-tetrahydrocannabinol disrupts hippocampal neuroplasticity and neurogenesis in trained, but not untrained adolescent Sprague-Dawley rats. *Brain Research*, *1548*, 12–19. https://doi.org/10.1016/j.brainres.2013.12.034

Suliman, N. A., Taib, C. N. M., Moklas, M. A. M., & Basir, R. (2018). Delta-9-Tetrahydrocannabinol (Δ9-THC) Induce Neurogenesis and Improve Cognitive Performances of Male Sprague Dawley Rats. *Neurotoxicity Research*, *33*(2), 402–411. https://doi.org/10.1007/s12640-017-9806-x

Svobodny, G. (2014). The internationalization of Ayahuasca. *Journal of Ethnopharmacology*, *153*(3), 933. https://doi.org/10.1016/j.jep.2014.02.018

Szabo, A. (2015). Psychedelics and Immunomodulation: Novel Approaches and Therapeutic Opportunities. *Frontiers in Immunology*, *6*(July), 1–11. https://doi.org/10.3389/fimmu.2015.00358

Tanimura, A., Liu, J., Namba, T., Seki, T., Matsubara, Y., Itoh, M., Suzuki, T., & Arai, H. (2009). Prenatal phencyclidine exposure alters hippocampal cell proliferation in offspring rats. *Synapse*, *63*(9), 729–736. https://doi.org/10.1002/syn.20660

Thompson, V. B. B., Koprich, J. B. B., Chen, E. Y. Y., Kordower, J. H. H., Terpstra, B. T. T., & Lipton, J. W. W. (2012). Prenatal exposure to MDMA alters noradrenergic neurodevelopment in the rat. *Neurotoxicology and Teratology*, *34*(1), 206–213. https://doi.org/10.1016/j.ntt.2011.09.005

Thompson, V. B., Heiman, J., Chambers, J. B., Benoit, S. C., Buesing, W. R., Norman, M. K., Norman, A. B., & Lipton, J. W. (2009). Long-term behavioral consequences of prenatal MDMA exposure. *Physiology & Behavior*, *96*(4–5), 593–601. https://doi.org/10.1016/j.physbeh.2008.12.013

Toriumi, K., Mouri, A., Narusawa, S., Aoyama, Y., Ikawa, N., Lu, L., Nagai, T., Mamiya, T., Kim, H.-C., & Nabeshima, T. (2012). Prenatal NMDA Receptor Antagonism Impaired Proliferation of Neuronal Progenitor, Leading to Fewer Glutamatergic Neurons in the Prefrontal Cortex. *Neuropsychopharmacology*, *37*(6), 1387–1396. https://doi.org/10.1038/npp.2011.324

Vollenweider, F. X., & Kometer, M. (2010). The neurobiology of psychedelic drugs: implications for the treatment of mood disorders. *Nature Reviews Neuroscience*, *11*(9), 642–651. https://doi.org/10.1038/nrn2884

Wankhar, W., Syiem, D., Pakyntein, C. L., Thabah, D., & Sunn, S. E. (2020). Effect of 5-HT2C receptor agonist and antagonist on chronic unpredictable stress (CUS) - Mediated anxiety and depression in adolescent Wistar albino rat: Implicating serotonin and mitochondrial ETC-I function in serotonergic neurotransmission. *Behavioural Brain*

Research, *393*, 112780. https://doi.org/10.1016/j.bbr.2020.112780

WHO. (2012). Depression, a global public health concern. *WHO Department of Mental Health and Substance Abuse*, 1–8. https://doi.org/10.1007/978-3-642-11688-9_20

Williams, G. V, Rao, S. G., & Goldman-Rakic, P. S. (2002). *The Physiological Role of 5-HT 2A Receptors in Working Memory*.

Wimbiscus, M., Kostenko, O., & Malone, D. (2010). MAO inhibitors: Risks, benefits, and lore. *Cleveland Clinic Journal of Medicine*, *77*(12), 859–882. https://doi.org/10.3949/ccjm.77a.09103

Winne, J., Boerner, B. C., Malfatti, T., Brisa, E., Doerl, J., Nogueira, I., Leão, K. E., & Leão, R. N. (2020). Anxiety-like behavior induced by salicylate depends on age and can be prevented by a single dose of 5-MeO-DMT. *Experimental Neurology*, *326*(October 2019), 113175. https://doi.org/10.1016/j.expneurol.2020.113175

Wong, M. L., & Licinio, J. (2001). Research and treatment approaches to depression. *Nature Reviews Neuroscience*, *2*(5), 343–351. https://doi.org/10.1038/35072566

Yuan, T.-F., Paes, F., Arias-Carrión, O., Ferreira Rocha, N., de Sá Filho, A., & Machado, S. (2015). Neural Mechanisms of Exercise: Anti-Depression, Neurogenesis, and Serotonin Signaling. *CNS & Neurological Disorders - Drug Targets*. https://doi.org/10.2174/1871527315666151111124402

Zhang, F., Challapalli, S. C., & Smith, P. J. W. (2009). Cannabinoid CB1 receptor activation stimulates neurite outgrowth and inhibits capsaicin-induced Ca2+ influx in an in vitro model of diabetic neuropathy. *Neuropharmacology*, *57*(2), 88–96. https://doi.org/10.1016/j.neuropharm.2009.04.017

Zhang, Q., Yu, Y., & Huang, X. F. (2016). Olanzapine Prevents the PCP-induced Reduction in the Neurite Outgrowth of Prefrontal Cortical Neurons via NRG1. *Scientific Reports*, *6*(1), 19581. https://doi.org/10.1038/srep19581

Zhou, D., & Song, Z. H. (2001). CB1 cannabinoid receptor-mediated neurite remodeling in mouse neuroblastoma N1E-115 cells. *Journal of Neuroscience Research*, *65*(4), 346–353. https://doi.org/10.1002/jnr.1160

Zuo, D., Liu, Y., Liu, Z., Cui, J., Zhou, X., Liu, Y., Li, Z., & Wu, Y. (2018). Alcohol aggravates ketamine-induced behavioral, morphological and neurochemical alterations in adolescent rats: The involvement of CREB-related pathways. *Behavioural Brain Research*, *349*, 80–90. https://doi.org/10.1016/j.bbr.2018.05.003