Allosteric regulation of kinase activity in living cells

Shivani Godbole[1] and Nikolay V. Dokholyan[1,2,3,4,5,*]

[1]Department of Pharmacology, Penn State College of Medicine, Hershey, PA 17033-0850, USA.
[2]Department of Biomedical Engineering, Penn State University, University Park, PA 16802, USA.
[3]Department of Engineering Science and Mechanics, Penn State University, University Park, PA 16802, USA
[4]Department of Biochemistry & Molecular Biology, Penn State College of Medicine, Hershey, PA 17033-0850, USA.
[5]Department of Chemistry, Penn State University, University Park, PA 16802, USA.

[*]Correspondence
Nikolay V. Dokholyan, dokh@psu.edu

Abstract

The dysregulation of protein kinases is associated with multiple diseases due to the kinases' involvement in a variety of cell signaling pathways. Manipulating protein kinase function, by controlling the active site, is a promising therapeutic and investigative strategy to mitigate and study diseases. Kinase active sites share structural similarities making it difficult to specifically target one kinase, allosteric control allows specific regulation and study of kinase function without directly targeting the active site. Allosteric sites are distal to the active site but coupled via a dynamic network of inter-atomic interactions between residues in the protein. Establishing an allosteric control over a kinase requires understanding the allosteric wiring of the protein. Computational techniques offer effective and inexpensive mapping of the allosteric sites on a protein. Here, we discuss methods to map and regulate allosteric communications in proteins, and strategies to establish control over kinase functions in live cells and organisms. Protein molecules, or "sensors" are engineered to function as tools to control allosteric activity of the protein as these sensors have high spatiotemporal resolution and help in understanding cell phenotypes after immediate activation or inactivation of a kinase. Traditional methods used to study protein functions, such as knockout, knockdown, or mutation, cannot offer a sufficiently high spatiotemporal resolution. We discuss the modern repertoire of tools to regulate protein kinases as we enter a new era in deciphering cellular signaling and developing novel approaches to treat diseases associated with signal dysregulation.

Introduction

Protein kinases expedite the cell signaling processes by facilitating the transfer of a phosphate from the nucleotide triphosphate to the target protein. Kinase-mediated phosphorylation is critical for a wide spectrum of cell signaling processes such as transcription, translation, cell proliferation, receptor downregulation, metabolism, and apoptosis[1]. The dysregulation of kinases alters cell function and may lead to diseases such as schizophrenia[2], Alzheimer's disease[3], cancer[4], and cardiovascular diseases[5]. Kinases undergo dynamic conformational changes to elicit their function during cell signaling. All kinases have a conserved residue sequence that forms the nucleotide-binding domain to bind to adenosine triphosphate (ATP) or guanosine triphosphate (GTP)[1,6]. The binding of the nucleotides is facilitated by phosphorylation of the activation loop[7]. Therapies targeting kinases focus on the nucleotide-binding sites (active/ orthosteric sites) that regulate the kinase function. Therapies targeting orthosteric sites to regulate kinase activity compete with the native ligand (ATP or GTP) to bind to the active site and need to be highly selective for the active site. Due to the conservation of the nucleotide-binding domain across all protein kinases, specifically targeting the active/ orthosteric sites of one type of kinase is not possible.

Kinase activity is also regulated by autoinhibitory domains[8,9]. Autoinhibitory domains suppress the interaction between the kinases and the effector molecule (nucleotides). The autoinhibitory domains are the sites distal to the active sites that affect the activity of the kinase at the active site through the induction of conformational and dynamic changes in the protein structure[10–12]. Allosteric regulation does not involve direct competition with the native ligand. Instead, it perturbs active site structure or dynamics. The kinases, especially in diseased conditions, may develop resistance to orthosterically regulated therapies by incorporating mutations at the active site, thus enabling kinases to elicit catalytic activity while disallowing the binding of the orthosteric regulators. Allosteric control of kinase function allows the bypass of this resistance.[11]

Finding allosteric sites that effectively regulate active sites is critical for establishing allosteric control over a particular kinase. X-ray crystallography[13] and nuclear magnetic resonance (NMR) spectroscopy[14] techniques are widely used to observe the structures of a protein. The changes occurring in the structure of protein molecules during the allosteric binding of effector molecules can be mapped by comparing structures at different points in time. However, these techniques are expensive and time-consuming. Modern computational approaches offer an inexpensive and fast assessment of protein allosteric "wiring". However, the computational approaches only offer hypothetical predictions and still need to be tested experimentally. Experimental validation does not require the structural characterization of the protein, instead, a direct test of the predictions in activity assays demonstrates the functioning of the established allosteric control[15]. Here, we discuss computational and experimental approaches to map allosteric communications, and tools used to control protein function through allosteric regulation.

Methods to map allosteric communications

Computer simulations developed to predict allosteric communications offer faster and more cost-effective methods of studying allosteric communication in proteins compared to traditional experimental structural characterization techniques[16]. Experimentally, NMR spectroscopy and X-ray crystallography are used in conjunction to map allostery, Boulton and Melacini[17] hypothesized that the crystal structures of proteins solved by these techniques would provide information about the changes in the conformation of proteins upon binding of an allosteric regulator. Sites that are susceptible to a conformational change upon binding of the modulator/regulator were termed "hot spots". The high-resolution images from NMR

spectroscopy provided the opportunity to study the conformational dynamics as a result of
allostery. By combining the high-resolution images of the protein structure, and the molecular
dynamics simulations, predictions can be made about the allosteric communications within
proteins. Lisi and Loria[18] described allostery using NMR studies paired with the computational
predictions in enzyme catalysts; PTP1B (protein tyrosine phosphatase 1B)[19], which is known to
be associated with obesity and diabetes, and PKA-C (protein kinase C)[20], which is responsible
for cell signaling and proliferation. Jain et al.[21] used NMR spectroscopy to configure the active
and inactive structures of Cdc42-LOV2 construct. Light oxygen voltage (LOV2) protein is used
as an optogenetic tool since it is active under blue light and is used to control protein function
allosterically. The authors elucidated the structural changes of the Cdc42-LOV2 occurring upon
irradiation by blue light converting the construct from an inactive to an active state. This
provides an insight on the allosteric communication within proteins through structural
elucidation. NMR spectroscopy is able to map both structural and dynamic changes that occur
within an allosteric system, but may not be sufficient to determine the structural and dynamic
changes occurring in complex allosteric systems.[17]

Computational methods provide a faster means of deciphering protein interactions.
Gadiyaram et al.[22] developed a graph spectral analysis of protein structures to predict allostery
using non-covalent interactions between amino acid residues to create a protein structure
network. The authors put their theories to the test by predicting the side-chain interactions
between residues that form the G-protein coupled receptors (GPCRs) (specifically the β2-
adrenergic receptor) and the HIV proteases. Another computational strategy, multiple sequence
alignments or MSA[23] is the process of aligning two or more protein sequences to discover the
conserved regions of the proteins. Lockless and Ranganathan[24] made use of MSA to map
allosteric communications in a postsynaptic density protein, Drosophila disc large tumor
suppressor, and zonula occludens-1 (PDZ) protein domain. They hypothesized that
perturbations in one residue of the protein sequence would cause thermodynamic changes in
the residue adjacent to it. If the protein sequences are conserved, similar changes in the
vibrational frequencies should be seen in all proteins possessing the conserved residue
sequence. The theory had two developmental parameters – conservation of the residues at a
given site of the protein as observed in the MSA, and statistical coupling of two residues (nodes)
of the given amino acid sequence. This method was validated by analyzing the conserved
sequences of the PDZ and the Poxvirus and zinc finger (POZ) protein domains using statistical
coupling analysis[25] and thermodynamic mutation cycles[26], which measures the strength of the
interaction of the residues. Ambroggio and Kulhman[27] developed an algorithm to engineer
proteins that can undergo structural protein-folding changes using MSA. The algorithm
Ambroggio and Kulhman developed could predict stable structural conformations of both the
changed and native protein instead of one protein conformation favoring the other. To test the
algorithm, the authors made use of the Sw2 sequence, which could switch the conformations
between the coiled coil and the zinc finger states. The information provided by MSA is
susceptible to errors[28] as the "true alignment" of proteins is not possible limiting the accuracy of
allosteric communication predictions using MSA[29].

The calculations necessary to describe protein structure flexibility and allostery of large
molecular systems are computationally intensive. Molecular dynamic (MD) simulations help to
overcome this limitation in predicting the interactions between amino acid residues. MD
simulations use atomic interaction approximations to simulate atomic motions on the basis of
Newtonian physics and reduce the computational complexity of this systems[30]. MD simulations
are utilized to develop algorithms that predict allosteric communications. The algorithms not only
help in predicting allosteric communication pathways within proteins but also help in determining
the potential allosteric sites in the protein. Proctor et al.[31] analyzed the allosteric
communications in mutations of cystic fibrosis transmembrane conductance regulator protein
(CFTR) and identified the allosteric hot spots using molecular dynamic simulations. These hot

spots could be targeted to restore the transcription and translation of CFTR and provide an
allosteric therapeutic approach. MD simulations can predict residue interactions within proteins
at a high spatiotemporal resolution and provide a faster method of analysis of allosteric
communication within a protein compared to experimental methods used for structure
elucidation. The timescales play a role in predicting residue interactions since allosteric
communications are dynamic in nature. Short time scales (nanosecond) are not always
accurately predicted by MD simulations,[32] this can be overcome using coarse-grained
simulations (a type of multiscale modeling method that simulates complex behavior of system in
a more realistic timescale) or discrete molecular dynamic (DMD) simulations that reach time
scales in the range of microseconds.[33,34] DMD simulations have been successfully used to
predict allosteric communications in proteins.[35,36]

Berezovsky et al.[37] developed a structure-based statistical mechanical model of allostery
(SBSMMA). This model predicts allostery by quantifying the thermodynamic perturbations of
proteins (upon ligand binding or mutations in the protein) by estimating the free energy of every
residue present in the ligand-bound and ligand-free proteins using an allosteric signaling map
(ASM). SBSMMA identifies potential exosites, protein regions that are likely to govern the
allosteric regulation through reverse perturbations. AlloSigMA[38] is a server that provides a
quantitative assessment of protein-energy dynamics by estimating the changes in the relative
energy of the residues at the allosteric site. AlloMAPS[39] is a database that provides data on the
predicted energy changes and the allosteric pathways of the molecule predicted using
SBSMMA[37]. SBSMMA provides a way to discover allosteric sites on the protein and elucidate
allosteric mechanisms within proteins.

Treating the allosteric phenomenon as signal propagation through heterogeneous
media, Wang et al.[40] designed a physics-based computational approach for mapping allosteric
communications called Ohm[40]. The multifaceted platform can be used to predict the allosteric
sites along with the possible pathways, identify the critical residues, and predict the possible
allosteric correlation between the residues. The platform is built on a perturbation propagation
algorithm that measures the perturbation of residues by calculating the probability of interactions
between the two residues (obtained by a probability matrix). The predictions were validated by
predicting the allosteric communication during phosphorylation of CheY[41] (a protein found in the
flagellar motor of bacteria) and the Caspase-I protein[42] (involved in inflammatory and host-
defense response). Allosteric communication within proteins can be predicted by a variety of
computational methods that provide closely realistic predictions in a short span of time. These
predictions can be validated using experimental techniques.

Tools to control allostery

Conceptually, a "sensor" domain is engineered into an allosteric site to establish allosteric
control over a protein. This sensor domain responds to stimuli, external or internal, and
undergoes conformational changes that propagate through allosteric paths to the active site.
The active site then is altered upon the application of the stimuli. Some of the examples are
where "sensor" proteins interact with small molecules or light and produce a conformational and
dynamic change in the target protein leading to either activation or deactivation of the protein.

Optogenetic control. Photoreceptors are molecules that can change their conformation upon
irradiation by light. The light-sensitive domains of the photoreceptors can be used as sensor
domains to control protein function[43]. The phytochrome B (PhyB)[44] is used extensively as a red
light photoreceptor, since many optical systems do not have far-infrared light range lasers, red
light photoreceptors have limitations to their usage.[44] Light Oxygen Voltage (LOV2) and
Cryptochromes are the two main types of blue light photoreceptors used to control protein
function. The (LOV2) domain of phototropin 1[45] obtained from *Avena sativa* has been shown to

produce dramatic conformational changes upon exposure to blue light and is used to control
protein function. Zimmerman, et al.[46] developed LOV2 photoswitches and validated them by
inserting them in various protein domains and determining the activity of the protein upon
exposure to blue light. Dagliyan et al.[47] developed a photo-inhibitable Src kinase domain by
inserting LOV2 in the L1 loop connecting β-strands of Src kinase causing inhibition of the kinase
upon activation of LOV2. However, insertion of the LOV2 domain into the protein without
causing structural perturbations in the native protein structure is difficult due to limitations arising
from the lack of availability of caging surfaces leading to non-specific activation of the
photoreceptors. Caging surfaces or photocages are photosensitive groups that prevent the
protein from reacting to small molecules or other proteins that might produce a biological
function of the protein. The caging groups ensure activation of proteins only upon irradiation of
the proteins by light of suitable wavelength[48]. To overcome this limitation, He et al.[49] developed
a circular permutant of LOV2 (cpLOV2). The cpLOV2 is constructed by introduction of a glycine
and serine-rich linker in the surface-exposed loop. The construct remains compatible with the
existing LOV2-based engineered proteins with the advantage of having more caging surfaces
thus improving its applicability (Figure 2).
Owing to the difficulty of optogenetic tools to control allosteric activity, arising due to the
limitations in achieving subcellular control of activity[50], Shaaya et al.[51] developed an optogenetic
tool, LightR, that enables subcellular control of protein and provides a tight temporal regulation
of protein activity. LightR comprises two tandemly connected Vivid photoreceptor[52] domains that
form an antiparallel homodimer upon irradiation by blue light causing its activation. The use of
LightR in controlling allosteric activity was validated in Src kinase, where irradiation with blue
light caused activation of Src kinase. LightR provides better spatiotemporal resolution than
LOV2 due to the subcellular control of proteins (Figure 3).
219
Chemogenetic control. Although optogenetics provides a high resolution of control of protein
activity, optogenetic tools are often challenging to translate from cellular to animal models due
to the limitation of the extent of absorption of light by animals that are non-transparent. To
overcome this limitation, Dagliyan et al.[53] developed a protein switch (uniRapR) whose activity
can be controlled by a small molecule (rapamycin). uniRapR is a complex of FK506-binding
protein (FKBP12) and FKBP12-rapamycin binding protein[54] built into a single-chain protein.
uniRapR binds selectively to rapamycin and upon binding induces a conformational change in
the structure. uniRapR can be used to control protein function allosterically. To test the
functionality of uniRapR, the authors inserted this construct into Src kinase and focal adhesion
kinase (FAK) and compared the changes in phosphorylation of the two enzymes before and
after the addition of rapamycin. The activity of uniRapR was further validated using a mutant
strain of FAK with a point mutation that caused inhibition of FAK activity, the FAK activity was
recovered by uniRapR which validated its use as an allosteric protein function regulator.
Guntas et al.[55] developed a maltose-binding allosteric regulator using *in vitro*
recombination techniques. The wild-type beta-lactamase enzyme (BLA) activity is rendered
unaffected by *E. coli* maltose binding protein. Upon recombination of the two genes, TEM-1-
beta-lactamase and *E. coli.* maltose binding gene[55], the enzyme activity was seen to be
regulated upon the addition of maltose. The change in activity was analyzed by measuring the
rate of β-lactam hydrolysis upon an increase in the concentration of maltose. Chemogenetic
control of proteins has better *in vivo* resolution since light can only be absorbed up to a certain
depth and has a shorter timescale of activation (Figure 2).

Allosteric regulation of protein kinases

Kinases are considered to be biological switches due to their ability to change structural
conformation upon activation.[56] Since kinases make up 2% of the protein-coding genome[57] in

humans and are implicated in multiple diseases this family of proteins is an important
therapeutic target. Protein kinases phosphorylate serines, threonines, or tyrosines by
transferring the γ-phosphate of ATP to the hydroxyl group causing structural changes in the
protein, eliciting its function as an enzyme.[58] Due to the widespread role of kinases in nature,
they are considered to be desirable targets for studying cellular signaling. In eukaryotic cells,
across all different types of protein kinases there are close to 250 residues worth of conserved
sequences.[56] One conserved region is attributed to be the nucleotide-binding domain (NBD)
which is a cleft that forms the binding pocket for ATP and is located between the N-terminal lobe
and the C-terminal lobe of the protein. The flexibility of the protein is maintained by two
ensembles of hydrophobic residues or "spines" that connect the α-helix of the C lobe to the β-
sheet of the N lobe.[59] These spines are called the "regulatory spine (R spine)", due to their
ability to regulate the assembly and disassembly of the active kinase, and the "catalytic spine (C
spine)", since these residues balance the catalytic activity of the kinases upon ATP binding. The
changes in structure of the kinases due to their complex structure-activity relationship are likely
to be resolved due to the presence of the spines.[60] The dynamic structure of kinases has proved
to be of great importance for the functionality of the protein and the NMR structures confirming
these dynamic structural changes have underscored the presence and importance of allostery
of protein kinases[61] (Figure 1).

262
263 **Figure 1** Domains of kinase protein responsible for the regulation of function
264

Several approaches have been developed using the pre-existing tools to control protein
function for the allosteric regulation of protein kinases. Kinase activity can be controlled by
ligands, Zorba et al.[62] made use of monobodies[63] (a type of synthetic binding protein) to control
allostery in Aurora A (AurA)[64]. AurA is a protein kinase responsible for the maturation of the
centrosome and duplication through its effect on the microtubule spindle formation in mitosis[65].
AurA serves to be a major oncogene since its overexpression has been associated with a
variety of cancers. The authors used monobodies[66] to bind to the allosteric sites of AurA and
produce complete inhibition or complete activation without having to actively compete with ATP
to bind to the NBD. The effect of the binding of monobodies to AurA, and the activity of the
protein, were mapped using X-ray crystallography. The binding of the monobody showed a 20-
fold inhibition of the kinases proving its effectiveness as an allosteric inhibitor and demonstrating
promising therapeutic potential. Wojcik et al.[67] designed high-affinity monobodies to bind to the
kinase Bcr-Abl formed by the fusion of breakpoint cluster region (Bcr) and Abelson tyrosine
kinase 1 (Abl1). Bcr-Abl activity leads to the development of chronic myelogenous leukemia.[68]
Studies have shown an interaction between Src homology domain (SH2) and tyrosine kinase,
critical to kinase activity and mutations arising at these interfaces (I164E), leads to inhibition of
the kinase activity and oncogenesis induced by Bcr-Abl.[69] The authors developed monobodies
for these critical residues to control the kinase activity of Bcr-Abl allosterically. The monobodies
were developed using combinatorial synthesis and the binding affinity was tested using chemical
shift perturbation assays that identify binding sites and predict the binding efficiency of ligands.
Inhibition of interaction of the SH2 domain and the kinase domain due to monobodies
decreased the activity of Bcr-Abl (validated by western blot and immunohistochemistry
experiments). Monobodies can be used as a tool for controlling kinase activity, providing an
alternative approach to chemogenetic and optogenetic control of protein function as
monobodies do not require engineering of the target kinase. (Figure 2).
Dagliyan et al.[70] developed a rapamycin-binding domain that destabilized the NBD,
thereby disrupting the kinase activity in the P21-activated kinase (PAK1). PAK1 is known to play
a role in angiogenesis, neurogenesis, and cell motility.[71] The dysregulation of this protein leads
to the development of speech impairment, seizures, and macrocephaly. The authors developed
a protein domain that is activated upon dopamine binding to the PAK1 domain. PAK1 is
overexpressed in breast cancer and hence is identified as an oncogene.[72] The authors tested
this rapamycin-PAK construct (RapR-PAK1) in breast cancer cells to observe changes in the
kinase activity due to the effect of rapamycin. The construct was successfully able to control the
kinase activity of PAK, demonstrating the applicability of ligands to control protein activity
allosterically by either inhibiting activity or inducing it.
300

301
302 **Figure 2 A** Allosteric activation of kinase (ERK2 kinase, PDB ID: 4GT3) by uniRapR (PDB ID:
303 7F2J) domain that binds to small molecule (rapamycin) causing activation of the protein. **B**
304 Allosteric inhibition of kinase (ERK2 kinase, PDB ID: 4GT3) by insertion of an optogenetic
305 control protein, LOV2 (PDB ID: 2V0W), causing conformational change upon irradiation by blue
306 light. **C** Activation of kinase using monobodies (PDB ID: 3RZW).
307
Karginov et al.[73] used a similar strategy for regulating kinase activity, but instead of using
a ligand-based approach, the authors used light as the input signal for controlling activity. The
authors built a photoactivatable analog of rapamycin to regulate the dimerization of proteins.
The photo-inducible small molecules are generated by the installation of "caging groups"[74,75]
that keep the molecule inactive until removed. Caging groups sequester the site on the ligand
necessary for activation. Dissociation of the caging groups from the activation site through
irradiation by light leads to activation of the small molecules. The authors used "caged-
rapamycin (pRap)" to regulate both the kinase activity and protein dimerization.
Zhou et al.[76] developed an optically-inducible, single-chain protein that could undergo
conformational changes upon exposure to light. The authors induced modifications to
DronpaN145, a photo-inducible green fluorescent protein to form pdDronpa1[77] that could
dimerize at the kinase active site in the presence of 500 nm cyan and stymie kinase activity.
Upon removal of light, the kinase activity was restored. The authors used dual specificity
mitogen-activated protein kinase 1 (MEK1) to test the pdDronpa1 construct. The authors were
able to successfully control the kinase activity of MEK1 providing a unique and versatile
approach toward optogenetics.
Vishweshwaraiah et al.[78] developed a protein-based nanocomputing agent[79] containing
two inputs that had opposing effects on the kinase activity, forming a logic OR gate to regulate
allosteric communications. The authors inserted a chemogenetic (uniRapR[53]) and an
optogenetic domain (LOV2[80]) within FAK. The uniRapR construct was inserted in the allosteric
site of the kinase domain, predicted using Ohm[40], such that it could activate the protein upon
interaction with rapamycin. This activation pathway of the protein upon interaction with
rapamycin was predicted by DMD simulations. The LOV2 construct was inserted in loop 2 of the
FERM domain (allosteric site on FERM domain was predicted using Ohm) leading to an
inhibition of kinase activity by inactivation of the protein[81]. Inactivation of the protein was
predicted using DMD simulations in blue light on/off conditions. The authors observed a
decrease in kinase activity upon exposure to blue light and an increase in kinase activity upon
exposure to rapamycin upon live cell imaging (changes in phenotype were used to determine

kinase activity). This successful study provides a basis for the development of rationally designed nano-computing agents and can be used in diagnostics, drug delivery, regulation of cell signaling, and metabolic flux. In order to demonstrate that the nanocomputing agents are effective in other proteins as well, Chen et al. developed a Src-uniRapR-LOV2 construct that provided a dual control on the function of Src kinase. The uniRapR construct was able to activate the Src kinase while the LOV2 construct rendered the protein inactive upon irradiation by blue light. The on/off protein switch was tested by visualization of the focal adhesion. With rapamycin as the first signal, followed by blue light as the second.[82] Along with computational methods such as MSA, MD, and DMD simulations; molecular docking and *in silico* modeling[83] paired with mass spectroscopy have also been used to define protein structures and map allosteric sites in proteins. Through microscopic techniques, such as cryo-EM, two-dimensional structures of proteins can be acquired and morphed into three-dimensional models using artificial intelligence[84]. In addition to small molecules, biologics such as antibodies or various input stimuli (such as pH, temperature, or pressure) can be made use of to regulate protein function by exploitation of allosteric communication. Protein allostery proves to be an essential tool in controlling protein function in translational research.

Figure 3 Sensor proteins used to control function of kinase protein allosterically. Chemogenetic tools (LightR, LOV2), optogenetic tool (UniRapR, Maltose binding protein), synthetic protein tool (Monobody).

Conclusion

The human cell kinome consists of 538 kinases that are involved in a variety of cellular processes. Kinases have a conserved active site residue sequence making it challenging to target a specific kinase with ligands acting on the active site, hence, allosteric regulation kinases provides the ability to specifically target different types of kinases. Finding the allosteric sites on the proteins and visualizing the changes occurring in the protein conformation is experimentally time-consuming and an expensive endeavor. To overcome this limitation, computational methods are developed that can predict allosteric communication within proteins. Advancements in technology have paved the way to engineer proteins and regulate their activity. Various methods such as chemogenetics, optogenetics, and synthetic proteins are used as allosteric tools to control protein function. Kinases are an important family of proteins as they are implicated in a variety of diseases, hence allosteric regulation of kinases could be used as a potential therapeutic approach. Combining the various allosteric control tools help develop nanocomputing agents with more than one input function leading to a different output function of the protein, providing means of controlling different aspects of protein function. New tools can be developed that react to different inputs such as temperature, pH, and pressure along with the preexisting tools interacting with light or small molecules. These allosteric control sensor proteins provide a better spatiotemporal resolution in controlling protein function and prove to be a potential therapeutic approach targeting diseases that arise from the dysregulation of proteins.

Acknowledgments
We thank Dr. Richard Mailman, Brianna Hnath and Jiaxing Chen for valuable discussions. We acknowledge support from the National Institutes for Health (R35 GM134864 to N.V.D.) and the Passan Foundation.

References:

1. Cheng, H.-C., Qi, R. Z., Paudel, H. & Zhu, H.-J. Regulation and Function of Protein Kinases and

    Phosphatases. *Enzyme Res.* **2011**, 1–3 (2011).

2. Bentea, E. *et al.* Kinase network dysregulation in a human induced pluripotent stem cell model of

    DISC1 schizophrenia. *Mol. Omics* **15**, 173–188 (2019).

3. Ohno, M. Roles of eIF2α kinases in the pathogenesis of Alzheimer’s disease. *Front. Mol.*

    Neurosci. **7**, (2014).

4. Sarkar, N., Singh, A., Kumar, P. & Kaushik, M. Protein kinases: Role of their dysregulation in

    carcinogenesis, identification and inhibition. *Drug Res.* **73**, 189–199 (2023).

5. Chen, J. *et al.* Protein kinases in cardiovascular diseases. *Chin. Med. J. (Engl.)* **135**, 557–570 (2022).

6.   Taylor, S. S. & Kornev, A. P. Protein kinases: evolution of dynamic regulatory proteins. *Trends*

Biochem. Sci. **36**, 65–77 (2011).

7.   Adams, J. A. Activation Loop Phosphorylation and Catalysis in Protein Kinases: Is There Functional

Evidence for the Autoinhibitor Model? *Biochemistry* **42**, 601–607 (2003).

8.   Soderling, T. R. Protein kinases. Regulation by autoinhibitory domains. *J. Biol. Chem.* **265**, 1823–1826

(1990).

9.   Li, M., Rehman, A. U., Liu, Y., Chen, K. & Lu, S. Dual roles of ATP-binding site in protein kinases:

Orthosteric inhibition and allosteric regulation. in *Advances in Protein Chemistry and Structural*

Biology vol. 124 87–119 (Elsevier, 2021).

10. Nussinov, R. & Tsai, C.-J. Allostery in Disease and in Drug Discovery. *Cell* **153**, 293–305 (2013).

11. Cui, Q. & Karplus, M. Allostery and cooperativity revisited. *Protein Sci.* **17**, 1295–1307 (2008).

12. Guo, J. & Zhou, H.-X. Protein Allostery and Conformational Dynamics. *Chem. Rev.* **116**, 6503–6515

(2016).

13. Keedy, D. A. Journey to the center of the protein: allostery from multitemperature multiconformer

X-ray crystallography. *Acta Crystallogr. Sect. Struct. Biol.* **75**, 123–137 (2019).

14. Grutsch, S., Brüschweiler, S. & Tollinger, M. NMR Methods to Study Dynamic Allostery. *PLOS Comput.*

Biol. **12**, e1004620 (2016).

15. Vishweshwaraiah, Y. L., Chen, J. & Dokholyan, N. V. Engineering an Allosteric Control of Protein

Function. *J. Phys. Chem. B* **125**, 1806–1814 (2021).

16. Dokholyan, N. V. Controlling Allosteric Networks in Proteins. *Chem. Rev.* **116**, 6463–6487 (2016).

17. Boulton, S. & Melacini, G. Advances in NMR Methods To Map Allosteric Sites: From Models to

Translation. *Chem. Rev.* **116**, 6267–6304 (2016).

18. Lisi, G. P. & Loria, J. P. Allostery in enzyme catalysis. *Curr. Opin. Struct. Biol.* **47**, 123–130 (2017).

19. Zhang, Z.-Y. Drugging the Undruggable: Therapeutic Potential of Targeting Protein Tyrosine

Phosphatases. *Acc. Chem. Res.* **50**, 122–129 (2017).

20. Xiao, Y., Liddle, J. C., Pardi, A. & Ahn, N. G. Dynamics of Protein Kinases: Insights from Nuclear

Magnetic Resonance. *Acc. Chem. Res.* **48**, 1106–1114 (2015).

21. Jain, A., Dokholyan, N. V. & Lee, A. L. Allosteric inactivation of an engineered optogenetic GTPase.

Proc. Natl. Acad. Sci. **120**, e2219254120 (2023).

22. Gadiyaram, V., Vishveshwara, S. & Vishveshwara, S. From Quantum Chemistry to Networks in

Biology: A Graph Spectral Approach to Protein Structure Analyses. *J. Chem. Inf. Model.* **59**, 1715–

1727 (2019).

23. Sofi, M. Y., Shafi, A. & Masoodi, K. Z. Multiple sequence alignment. in *Bioinformatics for Everyone*

47–53 (Elsevier, 2022). doi:10.1016/B978-0-323-91128-3.00011-2.

24. Lockless, S. W. & Ranganathan, R. Evolutionarily Conserved Pathways of Energetic Connectivity in

Protein Families. *Science* **286**, 295–299 (1999).

25. Teşileanu, T., Colwell, L. J. & Leibler, S. Protein Sectors: Statistical Coupling Analysis versus

Conservation. *PLOS Comput. Biol.* **11**, e1004091 (2015).

26. Pagano, L. *et al.* Double Mutant Cycles as a Tool to Address Folding, Binding, and Allostery. *Int. J.*

Mol. Sci. **22**, 828 (2021).

27. Ambroggio, X. I. & Kuhlman, B. Computational Design of a Single Amino Acid Sequence that Can

Switch between Two Distinct Protein Folds. *J. Am. Chem. Soc.* **128**, 1154–1161 (2006).

28. Kemena, C. & Notredame, C. Upcoming challenges for multiple sequence alignment methods in the

high-throughput era. *Bioinforma. Oxf. Engl.* **25**, 2455–2465 (2009).

29. Nuin, P. A., Wang, Z. & Tillier, E. R. The accuracy of several multiple sequence alignment programs for

proteins. *BMC Bioinformatics* **7**, 471 (2006).

30. Durrant, J. D. & McCammon, J. A. Molecular dynamics simulations and drug discovery. *BMC Biol.* **9**,

71 (2011).

31. Proctor, E. A. *et al.* Rational coupled dynamics network manipulation rescues disease-relevant

mutant cystic fibrosis transmembrane conductance regulator. *Chem. Sci.* **6**, 1237–1246 (2015).

32. Hertig, S., Latorraca, N. R. & Dror, R. O. Revealing Atomic-Level Mechanisms of Protein Allostery with

Molecular Dynamics Simulations. *PLOS Comput. Biol.* **12**, e1004746 (2016).

33. Shirvanyants, D., Ding, F., Tsao, D., Ramachandran, S. & Dokholyan, N. V. Discrete Molecular

Dynamics: An Efficient And Versatile Simulation Method For Fine Protein Characterization. *J. Phys.*

Chem. B **116**, 8375–8382 (2012).

34. Proctor, E. A., Ding, F. & Dokholyan, N. V. Discrete molecular dynamics. *WIREs Comput. Mol. Sci.* **1**,

80–92 (2011).

35. Bowerman, S. & Wereszczynski, J. Detecting Allosteric Networks Using Molecular Dynamics

Simulation. *Methods Enzymol.* **578**, 429–447 (2016).

36. Dokholyan, N. V., Buldyrev, S. V., Stanley, H. E. & Shakhnovich, E. I. Discrete molecular dynamics

studies of the folding of a protein-like model. *Fold. Des.* **3**, 577–587 (1998).

37. Tee, W.-V., Tan, Z. W., Lee, K., Guarnera, E. & Berezovsky, I. N. Exploring the Allosteric Territory of

Protein Function. *J. Phys. Chem. B* **125**, 3763–3780 (2021).

38. Guarnera, E., Tan, Z. W., Zheng, Z. & Berezovsky, I. N. AlloSigMA: allosteric signaling and mutation

analysis server. *Bioinformatics* **33**, 3996–3998 (2017).

39. Tan, Z. W., Tee, W.-V., Guarnera, E., Booth, L. & Berezovsky, I. N. AlloMAPS: allosteric mutation

analysis and polymorphism of signaling database. *Nucleic Acids Res.* **47**, D265–D270 (2019).

40. Wang, J. *et al.* Mapping allosteric communications within individual proteins. *Nat. Commun.* **11**,

3862 (2020).

41. Lee, S.-Y. *et al.* Crystal Structure of Activated CheY. *J. Biol. Chem.* **276**, 16425–16431 (2001).

42. Clark, A. C. Caspase Allostery and Conformational Selection. *Chem. Rev.* **116**, 6666–6706 (2016).

43. Padmanabhan, S., Pérez-Castaño, R., Osete-Alcaraz, L., Polanco, M. C. & Elías-Arnanz, M. Vitamin

B12 photoreceptors. in *Vitamins and Hormones* vol. 119 149–184 (Elsevier, 2022).

44. Ni, M., Tepperman, J. M. & Quail, P. H. Binding of phytochrome B to its nuclear signalling partner

PIF3 is reversibly induced by light. *Nature* **400**, 781–784 (1999).

45. Halavaty, A. S. & Moffat, K. N- and C-Terminal Flanking Regions Modulate Light-Induced Signal

Transduction in the LOV2 Domain of the Blue Light Sensor Phototropin 1 from *Avena sativa* ′.

Biochemistry **46**, 14001–14009 (2007).

46. Zimmerman, S. P., Kuhlman, B. & Yumerefendi, H. Engineering and Application of LOV2-Based

Photoswitches. in *Methods in Enzymology* vol. 580 169–190 (Elsevier, 2016).

47. Dagliyan, O. *et al.* Engineering extrinsic disorder to control protein activity in living cells. *Science* **354**,

1441–1444 (2016).

48. Bojtár, M. *et al.* Conditionally Activatable Visible-Light Photocages. *J. Am. Chem. Soc.* **142**, 15164–

15171 (2020).

49. He, L. *et al.* Circularly permuted LOV2 as a modular photoswitch for optogenetic engineering. *Nat.*

Chem. Biol. **17**, 915–923 (2021).

50. Wang, J. *et al.* Time-resolved protein activation by proximal decaging in living systems. *Nature* **569**,

509–513 (2019).

51. Shaaya, M. *et al.* Light-regulated allosteric switch enables temporal and subcellular control of

enzyme activity. *eLife* **9**, e60647 (2020).

52. Nihongaki, Y., Suzuki, H., Kawano, F. & Sato, M. Genetically Engineered Photoinducible

Homodimerization System with Improved Dimer-Forming Efficiency. *ACS Chem. Biol.* **9**, 617–621

(2014).

53. Dagliyan, O. *et al.* Rational design of a ligand-controlled protein conformational switch. *Proc. Natl.*

Acad. Sci. **110**, 6800–6804 (2013).

54. Choi, J., Chen, J., Schreiber, S. L. & Clardy, J. Structure of the FKBP12-Rapamycin Complex Interacting

with Binding Domain of Human FRAP. *Science* **273**, 239–242 (1996).

55. Guntas, G. & Ostermeier, M. Creation of an Allosteric Enzyme by Domain Insertion. *J. Mol. Biol.* **336**,

263–273 (2004).

56. Taylor, S. S., Keshwani, M. M., Steichen, J. M. & Kornev, A. P. Evolution of the eukaryotic protein

kinases as dynamic molecular switches. *Philos. Trans. R. Soc. B Biol. Sci.* **367**, 2517–2528 (2012).

57. Manning, G., Whyte, D. B., Martinez, R., Hunter, T. & Sudarsanam, S. The Protein Kinase Complement

of the Human Genome. *Science* **298**, 1912–1934 (2002).

58. Cohen, P. The origins of protein phosphorylation. *Nat. Cell Biol.* **4**, E127–E130 (2002).

59. Johnson, L. N. & Lewis, R. J. Structural Basis for Control by Phosphorylation. *Chem. Rev.* **101**, 2209–

2242 (2001).

60. Taylor, S. S., Ilouz, R., Zhang, P. & Kornev, A. P. Assembly of allosteric macromolecular switches:

lessons from PKA. *Nat. Rev. Mol. Cell Biol.* **13**, 646–658 (2012).

61. Masterson, L. R., Cembran, A., Shi, L. & Veglia, G. Allostery and Binding Cooperativity of the Catalytic

Subunit of Protein Kinase A by NMR Spectroscopy and Molecular Dynamics Simulations. in *Advances*

in Protein Chemistry and Structural Biology vol. 87 363–389 (Elsevier, 2012).

62. Zorba, A. *et al.* Allosteric modulation of a human protein kinase with monobodies. *Proc. Natl. Acad.*

Sci. **116**, 13937–13942 (2019).

63. Sha, F., Salzman, G., Gupta, A. & Koide, S. Monobodies and other synthetic binding proteins for

expanding protein science: Monobodies and Other Synthetic Binding Proteins. *Protein Sci.* **26**, 910–

924 (2017).

64. Zhou, H. *et al.* Tumour amplified kinase STK15/BTAK induces centrosome amplification, aneuploidy

and transformation. *Nat. Genet.* **20**, 189–193 (1998).

65. Glover, D. M., Leibowitz, M. H., McLean, D. A. & Parry, H. Mutations in aurora prevent centrosome

separation leading to the formation of monopolar spindles. *Cell* **81**, 95–105 (1995).

66. Koide, A., Wojcik, J., Gilbreth, R. N., Hoey, R. J. & Koide, S. Teaching an Old Scaffold New Tricks:

Monobodies Constructed Using Alternative Surfaces of the FN3 Scaffold. *J. Mol. Biol.* **415**, 393–405

(2012).

67. Wojcik, J. *et al.* Allosteric Inhibition of Bcr-Abl Kinase by High Affinity Monobody Inhibitors Directed

to the Src Homology 2 (SH2)-Kinase Interface. *J. Biol. Chem.* **291**, 8836–8847 (2016).

68. Deininger, M. W. N., Goldman, J. M. & Melo, J. V. The molecular biology of chronic myeloid leukemia.

Blood **96**, 3343–3356 (2000).

69. Grebien, F. *et al.* Targeting the SH2-Kinase Interface in Bcr-Abl Inhibits Leukemogenesis. *Cell* **147**,

306–319 (2011).

70. Dagliyan, O. *et al.* Engineering Pak1 Allosteric Switches. *ACS Synth. Biol.* **6**, 1257–1262 (2017).

71. Hofmann, C., Shepelev, M. & Chernoff, J. The genetics of Pak. *J. Cell Sci.* **117**, 4343–4354 (2004).

72. Radu, M., Semenova, G., Kosoff, R. & Chernoff, J. PAK signalling during the development and

progression of cancer. *Nat. Rev. Cancer* **14**, 13–25 (2014).

73. Karginov, A. V. *et al.* Light Regulation of Protein Dimerization and Kinase Activity in Living Cells Using

Photocaged Rapamycin and Engineered FKBP. *J. Am. Chem. Soc.* **133**, 420–423 (2011).

74. Riggsbee, C. W. & Deiters, A. Recent advances in the photochemical control of protein function.

Trends Biotechnol. **28**, 468–475 (2010).

75. Mayer, G. & Heckel, A. Biologically Active Molecules with a "Light Switch". *Angew. Chem. Int. Ed.* **45**,

4900–4921 (2006).

76. Zhou, X. X., Fan, L. Z., Li, P., Shen, K. & Lin, M. Z. Optical control of cell signaling by single-chain

photoswitchable kinases. *Science* **355**, 836–842 (2017).

77. Zhou, X. X., Chung, H. K., Lam, A. J. & Lin, M. Z. Optical Control of Protein Activity by Fluorescent

Protein Domains. *Science* **338**, 810–814 (2012).

78. Vishweshwaraiah, Y. L., Chen, J., Chirasani, V. R., Tabdanov, E. D. & Dokholyan, N. V. Two-input

protein logic gate for computation in living cells. *Nat. Commun.* **12**, 6615 (2021).

79. Dokholyan, N. V. Nanoscale programming of cellular and physiological phenotypes: inorganic meets

organic programming. *Npj Syst. Biol. Appl.* **7**, 15 (2021).

80. Lee, J. *et al.* Surface Sites for Engineering Allosteric Control in Proteins. *Science* **322**, 438–442 (2008).

81. Lietha, D. *et al.* Structural Basis for the Autoinhibition of Focal Adhesion Kinase. *Cell* **129**, 1177–1187

(2007).

82. Chen, J., Vishweshwaraiah, Y. L., Mailman, R. B., Tabdanov, E. D. & Dokholyan, N. V. A

noncommutative combinatorial protein logic circuit controls cell orientation in nanoenvironments.

Sci. Adv. **9**, eadg1062 (2023).

83. Yuan, J. *et al.* In Silico Prediction and Validation of CB2 Allosteric Binding Sites to Aid the Design of

Allosteric Modulators. *Molecules* **27**, 453 (2022).

84. Chirigati, F. Predicting protein structure from cryo-EM data. *Nat. Comput. Sci.* **1**, 96–96 (2021).

549