Homeostatic macrophages prevent preterm birth and improve neonatal outcomes by mitigating *in utero* sterile inflammation

Running title: Homeostatic Macrophages Mitigate *In Utero* Sterile Inflammation

Valeria Garcia-Flores[1,2,3], Zhenjie Liu[1,2], Roberto Romero[1,4,5], Roger Pique-Regi[2,6], Yi Xu[1,2],

Derek Miller[1,2,3], Dustyn Levenson[3,7], Jose Galaz[1,2,8], Andrew D. Winters[1,9],

Marcelo Farias-Jofre[1,2,8], Jonathan J. Panzer[9], Kevin R. Theis[1,2,9],

Nardhy Gomez-Lopez[1,2,3,9,10,11]

[1]Pregnancy Research Branch, Division of Obstetrics and Maternal-Fetal Medicine, Division of Intramural Research, *Eunice Kennedy Shriver* National Institute of Child Health and Human Development, National Institutes of Health, U.S. Department of Health and Human Services (NICHD/NIH/DHHS); Detroit, Michigan, and Bethesda, Maryland, USA

[2]Department of Obstetrics and Gynecology, Wayne State University School of Medicine; Detroit, Michigan, USA

[3]Center for Reproductive Health Sciences, Department of Obstetrics and Gynecology, Washington University School of Medicine; St. Louis, Missouri, USA

[4]Department of Obstetrics and Gynecology, University of Michigan; Ann Arbor, Michigan, USA

[5]Department of Epidemiology and Biostatistics, Michigan State University; East Lansing, Michigan, USA

[6]Center for Molecular Medicine and Genetics, Wayne State University; Detroit, MI, USA

[7]Department of Physiology, Wayne State University School of Medicine; Detroit, MI, USA

[8]Division of Obstetrics and Gynecology, School of Medicine, Faculty of Medicine, Pontificia Universidad Catolica de Chile; Santiago, Chile

[9]Department of Biochemistry, Microbiology, and Immunology, Wayne State University School

of Medicine; Detroit, Michigan, USA

[10]Department of Pathology and Immunology, Washington University School of Medicine; St.

Louis, Missouri, 63110, USA

[11]Lead contact

The study was conducted at the Perinatology Research Branch, NICHD/NIH/DHHS, in Detroit,

Michigan; the Branch has since been renamed as the Pregnancy Research Branch,

NICHD/NIH/DHHS

Correspondence: nardhy@wustl.edu

SUMMARY

Preterm birth (PTB), often preceded by preterm labor, is a major cause of neonatal morbidity and mortality worldwide. Most PTB cases involve intra-amniotic inflammation without detectable microorganisms, termed *in utero* sterile inflammation, for which there is no established treatment. Here, we propose homeostatic macrophages to prevent PTB and adverse neonatal outcomes caused by *in utero* sterile inflammation. Single-cell atlases of the maternal-fetal interface revealed that homeostatic maternal macrophages are reduced with human labor. M2 macrophage treatment prevented PTB and reduced adverse neonatal outcomes in mice with *in utero* sterile inflammation. Specifically, M2 macrophages halted premature labor by suppressing inflammatory responses in the amniotic cavity, including inflammasome activation, and mitigated placental and offspring lung inflammation. Moreover, M2 macrophages restored neonatal gut homeostasis and enhanced resistance to systemic bacterial infection. Our findings show that M2 macrophages are a promising strategy to mitigate PTB and improve neonatal outcomes from *in utero* sterile inflammation.

INTRODUCTION

Preterm birth, the leading cause of neonatal morbidity and mortality worldwide[1,2], is often preceded by spontaneous preterm labor, a syndrome of multiple etiologies[3]. Among the known and proposed causes of preterm labor, intra-amniotic inflammation is the best characterized and accounts for a large proportion of cases[3-5]. The recent incorporation of next generation sequencing in obstetrics has revealed that most cases of intra-amniotic inflammation occur in the absence of invading microbes in the amniotic cavity[6-9], resulting in the discovery of sterile intra-amniotic inflammation (hereafter referred to as *in utero* sterile inflammation). Hence, this new condition is diagnosed by elevated concentrations of inflammatory mediators such as interleukin (IL)-6 in amniotic fluid, in the absence of detectable microorganisms using culture and molecular microbiological techniques[6-10]. Importantly, *in utero* sterile inflammation has been linked to adverse short- and long-term outcomes for the offspring of women with this clinical condition[7,11]. Specifically, women with *in utero* sterile inflammation are at greater risk of having placentas affected by acute histologic chorioamnionitis[12], which is linked to the development of deleterious neonatal conditions such as bronchopulmonary dysplasia[13-15] and necrotizing enterocolitis[16,17], likely due to exposure of the fetal lungs and intestine to intra-amniotic inflammation[18-20]. However, despite the strong associations between *in utero* sterile inflammation and adverse fetal and neonatal outcomes, no approved treatments currently exist.

Sterile inflammation can be triggered by danger signals or alarmins released during cellular stress or injury[21-23]. Consistently, clinical studies have also shown that women with spontaneous preterm labor and *in utero* sterile inflammation have increased amniotic fluid concentrations of alarmins[7,24-26]. Indeed, women with preterm labor and elevated amniotic concentrations of the prototypical alarmin high-mobility group box-1 (HMGB1) ($\geq$8.55 ng/mL)

delivered earlier than those with lower concentrations of this alarmin[7]. Furthermore, we have

provided mechanistic evidence showing that the *in utero* delivery of HMGB1 induces preterm

labor and birth in mice[27-29]. Moreover, our *in vitro* studies demonstrated that incubation with

HMGB1 induces the activation of the NLRP3 inflammasome[30], one of the central pathway in

triggering preterm labor and birth in women[31] and mice[32-34] experiencing *in utero* sterile

inflammation[35]. Hence, a therapeutic approach targeting the NLRP3 inflammasome, elevated

inflammatory cytokines in amniotic fluid such as IL-6, and the process of preterm labor could

represent a promising strategy for treating *in utero* inflammation and its devastating perinatal

consequences.

The maternal-fetal interface hosts a notable and heterogeneous population of

macrophages[36-43]. Specifically, we reported that macrophages expressing a homeostatic or M2-

like phenotype are more abundant in both term and preterm gestation than those expressing

inflammatory phenotypes[42], pointing to an important role for these cells in maintaining

pregnancy homeostasis. Next, we demonstrated that the depletion of maternal macrophages

results in preterm birth as well as neonatal growth restriction and increased mortality[43].

Furthermore, we also showed that the adoptive transfer of M2-polarized macrophages prevents

preterm birth induced by intra-amniotic LPS[43], providing proof-of-concept that such cells can

serve as a therapeutic approach for *in utero* sterile inflammation. Thus, here we propose the use

of M2-polarized macrophages as a cellular therapy to prevent preterm labor associated with *in*

utero sterile inflammation as well as its consequences for the offspring.

In this study, we employ a translational mechanistic approach by first leveraging our

single-cell atlases of the human maternal-fetal interface to demonstrate a labor-associated

reduction of homeostatic macrophages. Next, by using a clinically relevant animal model of *in*

utero sterile inflammation induced by the intra-amniotic injection of the alarmin HMGB1, we

investigate the potential restoration of M2-polarized macrophages (hereafter also referred to as

M2 macrophages) via adoptive transfer to prevent preterm birth and reduce the adverse neonatal

outcomes. In addition, we utilize molecular approaches to investigate the inflammatory

responses driven by HMGB1-induced *in utero* sterile inflammation and the homeostatic effects

of M2 macrophage treatment in maternal and fetal tissues targeted by transferred macrophages,

including those involved in the common pathway of parturition. Moreover, we evaluate the

damage to key fetal and neonatal organs, namely the lung and intestine, driven by exposure to *in*

utero sterile inflammation, including alterations of the gut microbiome, and whether this was

reverted by M2 macrophage treatment. Last, we challenge neonates with Group B *Streptococcus*

to determine whether M2 macrophage treatment restores neonatal immunocompetence.

Collectively, our data indicate that treatment with M2 macrophages represents a novel cellular

approach that can prevent preterm birth and ameliorate the adverse neonatal outcomes induced

by *in utero* sterile inflammation.

RESULTS

The maternal-fetal interface hosts a homeostatic macrophage population that is diminished

with labor

We first hypothesized that labor is accompanied by a reduction of homeostatic

macrophages in maternal compartments. Our previous flow cytometry studies targeting specific

macrophage subsets suggested that such a reduction occurs at the maternal-fetal interface (i.e.,

the decidua)[42,43]. However, we sought to test our hypothesis using an unbiased discovery

approach. To do this, we leveraged our previously generated single-cell atlases of the

myometrium and maternal-fetal interface[44-46]. The maternal-fetal interface includes key sites of

contact between maternal and fetal tissues: the fetal placenta embedded in the maternal decidua

basalis adjacent to the myometrium, and the fetal extraplacental membranes enclosing the

amniotic cavity and attached to the maternal decidua parietalis, next to the myometrium (Figure

1A). Our myometrial single-cell atlas includes samples collected from women with term labor as

well as term non-labor controls[45], whereas our single-cell atlases of the placenta and

extraplacental membranes also include samples from women with preterm labor in addition to

the term groups[44,46]. Given that preterm non-labor deliveries are only performed due to

pregnancy complications, such cases are not suited for use as gestational age controls for preterm

labor and thus historically have not been utilized in our studies. After normalizing data from our

three single-cell atlases, we identified seven distinct macrophage clusters, termed M1 – M7,

across all tissues (Figure 1B, Supplementary Table 1). Of note, these cluster numbers were not

chosen to correlate with the conventional M1-M2 paradigm, but rather reflect cluster number

assignments. We focused on the M1 and M2 clusters because they constituted a significant

proportion of macrophages in the myometrium, placenta, and extraplacental membranes (Figure

1B).

133   Our single-cell atlases include maternal and fetal genotyping data, which allow us to

assign an origin to each individual cell. Therefore, we next determined the origins of

macrophages in each compartment (Figure 1C). As expected, macrophages in the myometrium

were entirely of maternal origin (Figure 1C). In the placenta, both maternal and fetal

macrophages were identified, with the majority of maternal macrophages corresponding to the

M1 cluster and fetal macrophages to the M2 cluster (Figure 1C). By contrast, the extraplacental

membranes include some fetal macrophages, but the M2 cluster was predominantly of maternal

origin (Figure 1C). Next, we evaluated whether the proportions of M1 and M2 clusters, as well

as other macrophage clusters, differed between labor and non-labor samples. The only subset

that decreased during labor was the maternal M2 cluster in the myometrium (Figure 1D). We

also observed a non-significant 1.4-fold decrease in the proportion of the maternal M2 cluster in

the basal plate (maternal tissue attached to the placenta) in women with preterm labor and birth

compared to those with term labor and birth (Figure 1E). This may be due to the small sample

size and the challenges associated with collecting human preterm samples. Last, using the top 20

marker genes for the M2 cluster (Figure 1F), we performed over-representation analysis (ORA)

based on the Gene Ontology (GO). The biological processes enriched in the M2 cluster included

"homeostatic process," "regulation of immune system process," and "regulation of

developmental process," supporting the homeostatic functions of this macrophage cluster (Figure

1G and Supplementary Figure 1A). These processes were distinct from those enriched in other

macrophage clusters (Supplementary Figure 1B).

Taken together, our single-cell atlases confirmed that the myometrium and other maternal

compartments attached to the placenta host a significant population of homeostatic macrophages,

and that these cells are reduced during labor, confirming our hypothesis (Figure 1A).

M2 macrophage treatment prevents preterm birth and improves neonatal survival in a

mouse model of *in utero* sterile inflammation

Next, we determined whether homeostatic macrophages could serve as a viable

therapeutic approach to prevent preterm birth. We therefore performed adoptive transfer of M2

macrophages in a murine model of *in utero* sterile inflammation induced by the ultrasound-

guided intra-amniotic injection of the alarmin HMGB1 (Figure 2A). To ensure the clinical

relevance of this model, we injected HMGB1 at concentrations found in women with preterm

labor and *in utero* sterile inflammation[7]. The intra-amniotic delivery of HMGB1 shortened

gestational length (Figure 2B), resulting in high rates of preterm birth (Figure 2C). Notably,

treatment with M2 macrophages extended gestational length, preventing preterm birth (Figure

2B&C). Neonatal mortality was elevated in mice exposed to *in utero* sterile inflammation, given

that most preterm neonates die; yet, such mortality was mitigated by treatment with M2

macrophages (Figure 2D). Furthermore, neonates born to dams intra-amniotically injected with

HMGB1 failed to thrive, but once again this effect was ameliorated by treatment with M2

macrophages (Figure 2E). Moreover, surviving neonates that had been exposed to HMGB1-

induced *in utero* sterile inflammation displayed growth restriction; however, such impairment

was rescued by prenatal treatment with M2 macrophages (Figure 2F). Thus, these data indicate

that M2 macrophage treatment can serve as a viable therapy to not only prevent preterm birth but

also ameliorate the adverse neonatal outcomes driven by exposure to *in utero* sterile

inflammation.

M2 macrophages dampen HMGB1-induced *in utero* sterile inflammation, including

inflammasome activation, in the amniotic cavity

We next sought to uncover the mechanisms whereby adoptively transferred M2

macrophages were preventing adverse outcomes driven by HMGB1-induced *in utero* sterile

inflammation. First, to establish the kinetics of the *in utero* sterile inflammatory response, we

collected the amniotic fluid of HMGB1-injected dams at 24, 48, 72, or 96 h post-injection to

evaluate the concentrations of IL-6, total IL-1β, and TNF - classic cytokines associated with *in*

utero sterile inflammation[25,26] (Figure 3A). The intra-amniotic delivery of HMGB1 increased the

amniotic fluid concentrations of IL-6, the gold standard cytokine used to clinically diagnose *in*

utero sterile inflammation[6,7], at 72 h post-injection (Figure 3B), consistent with the observed

timing of preterm labor post-HMGB1 injection. Notably, M2 macrophage treatment not only

reduced amniotic fluid concentrations of IL-6 at 72 h post-injection, but also those of TNF and

total IL-1β (Figure 3C&D), even though these were not significantly affected by HMGB1

(Supplementary Figure 2). We then focused on the inflammasome, a key signaling pathway

implicated in the *in utero* inflammatory response triggered by alarmins, leading to the processing

of active caspase (CASP)-1 and mature IL-1β (Figure 3E)[31-33,35]. The intra-amniotic delivery of

HMGB1 caused an increase in active CASP-1 and mature IL-1β in amniotic fluid (Figure

3F&G). Notably, M2 macrophage treatment dampened such inflammasome activation by

reducing both active CASP-1 and mature IL-1β (Figure 3H&I). These data indicate that M2

macrophages exert their homeostatic effects in the amniotic cavity by dampening the HMGB1-

induced inflammatory cytokine response, including inflammasome activation.

M2 macrophages interfere with the pathway of preterm labor

The processes of term and preterm labor share a common underlying pathway including

fetal membrane activation, uterine contractility, and cervical remodeling[47-50]. Such processes are

characterized by the activation of pro-inflammatory signaling networks, including inflammasome

activation[51]. Indeed, we recently reported that inflammasome activation in the fetal membranes

and uterine tissues is essential for the onset of preterm labor in mice undergoing *in utero* sterile

inflammation[33]. However, this pathway is not involved in the cervical processes associated with

preterm labor[33,52]. Hence, we next evaluated whether M2 macrophage treatment abrogates

inflammatory processes, including inflammasome activation in the fetal membranes (Figure 4A)

and uterine tissues. Consistent with our findings in amniotic fluid, intra-amniotic injection of

HMGB1 increased levels of active CASP-1 and mature IL-1β in the fetal membranes, indicating

inflammasome activation (Figure 4B&C). Yet, M2 macrophage treatment abrogated

inflammasome activation in the fetal membranes, as indicated by reduced quantities of active

CASP-1 and mature IL-1β (Figure 4D&E). Moreover, gene expression profiling revealed that

intra-amniotic injection of HMGB1 induced the overexpression of several inflammatory genes in

the fetal membranes, including a significant increase in *Ccl17* (Figure 4F&G). Conversely, M2

macrophage treatment exhibited a broad anti-inflammatory effect by downregulating the

expression of *Tlr4*, *Cxcl1, Tnf*, and *Il1b* in the fetal membranes (Figure 4H&I). These findings

indicate that M2 macrophages disrupt the inflammatory process of preterm labor in the fetal

membranes.

220   In the uterine tissues, intra-amniotic injection of HMGB1 induced inflammasome

221 activation (Supplementary Figure 3A-H), but this was not significantly reduced by treatment

222 with M2 macrophages. However, HMGB1 failed to induce a significant inflammatory response

223 in uterine tissues, and treatment with M2 macrophages had no significant effect (Supplementary

224 Figure 4A&B). These findings suggest that HMGB1-induced preterm labor in mice did not

225 trigger a strong inflammatory response in uterine tissues, rendering M2 macrophage treatment

226 unnecessary for preventing preterm birth.

227   In addition to the fetal membranes and uterus, the complex pathway of parturition also

228 involves the initiation of an immune response in the decidual tissues[3,49,53]. However, we have

229 demonstrated that such an inflammatory response is not associated with inflammasome

230 activation[32,33,54]. Through inflammatory gene profiling, we found that HMGB1 did not strongly

231 induce an inflammatory response in the decidua; yet, M2 macrophage treatment downregulated

232 several inflammatory genes (Supplementary Figure 5A&B). This finding show that, even in the

233 absence of abnormal decidual inflammation, M2 macrophages restrict the labor-associated

234 inflammatory response at the maternal-fetal interface.

235   Together, these findings demonstrate that M2 macrophage treatment prevents preterm

236 birth induced by *in utero* sterile inflammation by interfering with the inflammatory processes

237 required to activate the pathway of preterm labor in the fetal membranes and maternal-fetal

238 interface.

240 **M2 macrophage treatment mitigates inflammation in fetal and neonatal tissues**

241   Given the robust homeostatic effects of M2 macrophages in the amniotic cavity and fetal

242 membranes, we next investigated their ability to penetrate other fetal organs. We utilized a model

wherein M2 macrophages derived from donor CD45.1$^+$ mice were transferred to recipient

CD45.2$^+$ dams injected with HMGB1. Afterwards, maternal blood, maternal and fetal tissues,

and amniotic fluid were collected to track the migration of transferred cells (Supplementary

Figure 6A). Transferred M2 macrophages retained their M2 phenotype (data not shown), as

previously reported[43]. Representative flow cytometry dot plots display the proportions of

CD45.1$^+$ M2 macrophages detected in maternal and fetal compartments (Supplementary Figure

6B). Quantification of transferred CD45.1$^+$ M2 macrophages revealed distinct kinetic patterns

across the maternal and fetal compartments (Supplementary Figure 6C&D). CD45.1$^+$ M2

macrophages were highest in the maternal blood and lung at 2 hours post-injection, decreasing to

negligible levels by 12 hours. Conversely, a gradual accumulation of CD45.1$^+$ M2 macrophages

was observed in the uterus, while modest numbers were consistently detected in the decidua at

each time point (Supplementary Figure 6C). In the fetal compartments, CD45.1$^+$ M2

macrophages were abundant in the placenta at 2 hours and declined over time, similar to the

pattern in maternal blood, suggesting localization in the intervillous space rather than the

parenchyma (Supplementary Figure 6D). A small number of CD45.1$^+$ M2 macrophages

accumulated in the fetal membranes over time, with a few reaching the amniotic cavity. In

contrast, these cells were scarcely detected in the fetal intestine and lung at 2 hours, becoming

negligible by 6 and 12 hours. The transient presence of M2 macrophages in the fetal lung and

intestine is likely due to exposure to amniotic fluid, as evidenced by their absence in the

unexposed fetal liver at any time point (Supplementary Figure 6D). These findings demonstrate

that adoptively transferred M2 macrophages primarily migrate to intrauterine compartments,

with limited penetration into fetal compartments.

Therefore, it is likely that M2 macrophages exert their homeostatic effects on the fetus

through indirect mechanisms by downregulating the inflammatory response in the amniotic

cavity and surrounding fetal membranes (Figures 3 and 4) and reducing inflammation in the

placenta. We then evaluated the effect of M2 macrophage treatment on the placenta. The intra-

amniotic injection of HMGB1 triggered the upregulation of *Tlr4*, *Gja1*, and *Ptgs2*, along with a

non-significant increase in the expression of other genes, including *Ccl17* and *Nfkb2*, in the

placenta (Supplementary Figure 7A). Treatment with M2 macrophages downregulated the

expression of *Ccl17* and *Nfkb2* (Supplementary Figure 7B). Hence, M2 macrophages effectively

dampen placental inflammation as part of their beneficial effects *in utero*.

Next, we focused on the fetal lung since preterm neonates born to women with intra-

amniotic inflammation are at high risk of developing bronchopulmonary dysplasia[13-15]. In the

fetal lung (Figure 5A), intra-amniotic injection of HMGB1 resulted in the upregulation of *Il6*,

Tnfrsf1a, *Il33*, *Nlrp6*, and *Tlr9* (Figure 5B&C). M2 macrophage treatment downregulated the

expression of *Tnfrsf1a* and *Nod1* (Figure 5D&E). Next, we explored whether the homeostatic

effects of M2 macrophages were evident in the neonatal lung by evaluating inflammatory gene

expression in this tissue. Neonates born to dams treated with M2 macrophages exhibited a

diminished inflammatory profile in the neonatal lung, characterized by reduced expression of *Il6*,

Ccl2, *Socs3*, and *Cxcl1* in comparison to offspring from untreated dams (Figure 5F&G). These

findings indicate that M2 macrophages dampen inflammatory responses in the fetal lung, with

sustained effects in the neonatal lung. This underscores the protective role of these cells in

mitigating *in utero* sterile inflammation.

In addition to the lungs, the fetal intestine is also exposed to amniotic fluid, putting

neonates born to women with intra-amniotic inflammation at high risk of developing necrotizing

enterocolitis[16,17]. Therefore, we next explored inflammatory gene expression in the fetal intestine

following an intra-amniotic injection of HMGB1. This revealed alterations in inflammatory gene

expression (Supplementary Figure 8), consistent with our previous reports on *in utero*

inflammation[55,56]. Unlike in the fetal lung, the inflammatory milieu in the fetal intestine was

downregulated, suggesting that *in utero* sterile inflammation exerts different effects across fetal

organs. Since we previously showed that treatment with an anti-inflammatory drug did not exert

a significant impact on the fetal intestine exposed to *in utero* sterile inflammaiton[55], we shifted

our focus to investigate the effects of M2 macrophage treatment on the neonatal intestine (Figure

6A). Treatment with M2 macrophages boosted the neonatal gut inflammatory response

dampened by *in utero* exposure to HMGB1 (Figure 6B, D, F). Specifically, M2 macrophage

treatment upregulated the expression of *Aim2, Cd68, Tnf, Rgs4, Ccl5,* and *Ccl17* in the small

intestine; *Casp1* and *Ccl17* in the cecum; and *Il1a, Cd68, Casp1, Jun, Kdm6b,* and *Ccl17* in the

colon (Figure 6C, E, G) of neonates. These data show that treatment with M2 macrophages

enhances the neonatal gut inflammatory profile, which is otherwise suppressed by *in utero*

exposure to sterile inflammation.

We recently demonstrated that *in utero* sterile inflammation leads to neonatal gut

dysbiosis[55]. Therefore, we proceeded to evaluate whether treatment with M2 macrophages could

mitigate this dysbiosis by performing 16S rRNA gene sequencing of the small intestine, cecum,

and colon from neonates born to dams intra-amniotically injected with HMGB1 with or without

M2 macrophage treatment (Supplementary Figure 9A). Alpha diversity metrics of the

microbiome indicated no differences in the community evenness of the small intestine, cecum, or

colon between groups (data not shown). When examining beta diversity, principal coordinate

analysis (PCoA) revealed that the neonatal microbiomes were strongly clustered by treatment

group (Supplementary Figure 9B) and underwent consistent changes in structure and

Firmicutes/Bacteroidetes ratio (Supplementary Figure 9C-E). In the small intestine, M2

macrophage treatment was associated with a shift in the microbiome structure (Supplementary

Figures 9D and 10A&B) characterized by decreased abundance of *Rodentibacter* (ASV-5) and

Ruminiclostridium (ASV-17) (Supplementary Figure 10C). Similar patterns were also observed

in the cecum (Supplementary Figures 9D and 10D&E), in which a decreased abundance of

Ruminiclostridium (ASV-17), Lachnospiraceae (ASV-24)*,* and *Lachnoclostrium* (ASV-15) was

observed alongside an increased abundance of Lachnospiraceae (ASV-39) (Supplementary

Figure 10F). Consistently, in the colon, the microbiome structure was modified by M2

macrophage treatment (Supplementary Figures 9D and 10G&H), resulting in decreased

abundance of *Ruminiclostridium* (ASV-17)*,* Lachnospiraceae (ASV-24)*,* and *Lachnoclostrium*

(ASV-15) (Supplementary Figure 10I). Treatment with M2 macrophages alone did not alter the

alpha or beta diversity of the neonatal gut microbiome (Supplementary Figure 9F-H). Thus,

prenatal treatment with M2 macrophages can change neonatal gut dysbiosis induced by *in utero*

exposure to sterile inflammation.

Collectively, the data demonstrate that prenatal treatment with M2 macrophages dampens

fetal tissue inflammation, thereby preventing subsequent neonatal organ inflammatory injury and

gut microbiome dysbiosis. These findings suggest potential mechanisms through which this

cellular therapy improves neonatal outcomes following exposure to *in utero* sterile inflammation.

M2 macrophage treatment improves neonatal ability to fight Group B *Streptococcus*

infection following *in utero* sterile inflammation

Thus far, we have demonstrated that M2 macrophage treatment enhances survival and

alleviates inflammation in neonates exposed to *in utero* sterile inflammation. Therefore, we last

assessed whether these changes would result in improved neonatal ability to combat systemic

infection with Group B *Streptococcus* (GBS), a microbe commonly associated with neonatal

infection and sepsis[57]. At two weeks of age, control neonates (i.e., pups born to dams without

any treatment) and neonates exposed *in utero* to HMGB1 alone, HMGB1 with vehicle treatment,

or HMGB1 with M2 macrophage treatment received intraperitoneal injection of GBS (Figure

7A). *In utero* exposure to HMGB1 caused low neonatal survival rates within the first five days

post-GBS infection (Figure 7B). Notably, prenatal treatment with M2 macrophages bolstered

neonatal survival to be comparable to controls (Figure 7B), indicating improved capacity to fight

GBS infection. *In utero* exposure to HMGB1 also negatively impacted neonatal weight, which

was significantly reduced compared to controls at five days post-infection (Figure 8cC). Prenatal

treatment with M2 macrophages partially rescued neonatal weight but did not fully restore it to

the control trajectory (Figure 7C), a phenomenon that requires further investigation. The

improved neonatal survival and growth driven by M2 macrophage treatment were accompanied

by reduced incidence of hypothermia (considered the murine equivalent of fever[58,59]) upon GBS

infection (Supplementary Figure 11). Thus, prenatal treatment with M2 macrophages bolsters the

capacity to clear pathogenic bacteria in neonates compromised by exposure to *in utero* sterile

inflammation.

DISCUSSION

The lack of knowledge regarding specific molecular targets in women with *in utero*

sterile inflammation has made the investigation of useful treatment options challenging. In the

current study, we provide mechanistic evidence for M2 macrophage treatment as a cellular

therapy that can prevent preterm birth resulting from *in utero* sterile inflammation. The reasoning

behind choosing M2 macrophages as an approach to prevent preterm birth and the mechanisms

through which they exert these effects are explained below. *First*, our prior studies have

indicated that both physiological and pathological labor are accompanied by a shift in

macrophage polarization, primarily at the maternal-fetal interface, with a subset of these cells

acquiring a pro-inflammatory M1-like phenotype[42,43]. Here, we build on this concept by showing

that M2-like homeostatic macrophages of maternal origin are reduced in the myometrium and

basal plate of women with term and preterm labor, respectively. This finding complements our

previous targeted approaches, which showed a reduction of decidual M2 macrophages in women

experiencing preterm labor and birth[43]. Hence, it is possible that homeostatic macrophages in the

uterine and decidua tissues help maintain a favorable environment at the maternal-fetal interface

and surrounding tissues, thereby dampening inflammatory signaling associated with the

premature onset of labor. The latter concept is supported by our prior observations that the

depletion of maternal macrophages can result in preterm labor and birth[43]. *Second*, labor

involves coordinated activation across multiple maternal and fetal tissues, conventionally

referred to as the common pathway of parturition[47-50]. Herein, we demonstrate that M2

macrophages infiltrate the maternal-fetal interface, interfere with the common pathway of labor

in the fetal membranes and decidua, and dampen fetal inflammation mediated by the placenta

and amniotic cavity. In line with this, M2 macrophages express immunomodulatory factors, such

as arginase-1 and IL-10, that can modify the tissue microenvironment and prevent local immune

activation[60,61]. Indeed, in a previous study using an animal model of LPS-induced intra-amniotic

inflammation, we found that treatment with M2 macrophages increased concentrations of IL-10

in both maternal circulation and amniotic fluid, along with a reduction in inflammatory

mediators at the maternal-fetal interface[43], providing further evidence that homeostatic

macrophages target both maternal and fetal components in the process of preterm labor. *Third*,

we previously demonstrated that the inflammasome is a key pathway involved in premature

labor induced by microbes[54,62] or alarmins[32-34,63]. As a proof of concept, we administered a

specific inhibitor of the NLRP3 inflammasome (i.e., MCC950)[64] to dams injected with the

alarmin S100B and observed reduced rates of preterm birth and adverse neonatal outcomes[32].

While NLRP3 inhibitors hold promise, further research is required to comprehensively assess

their effects on offspring before considering their use during pregnancy. The current study

demonstrates that M2 macrophages suppress inflammasome activation in the amniotic cavity and

fetal membranes, offering an alternative strategy for modulating this signaling pathway.

In utero sterile inflammation is not only detrimental due to its short-term outcomes, such

as preterm birth, but is importantly linked to long-term negative consequences for the offspring

as indicated by human data[7,11] and animal studies[27-29,32,34,55,63,65]. Such consequences are not

entirely the result of prematurity, as neonates born at term to HMGB1-injected dams still had

reduced survival compared to those that received treatment with M2 macrophages. Thus, even a

relatively brief exposure to *in utero* sterile inflammation imprints fetal tissue alterations that can

be carried over to neonatal life. This concept is supported by previous research demonstrating

that prenatal treatments or exposures during gestation can induce permanent changes in neonatal

life[55,66-68]. Of interest, recent research employing a transient gestational infection model revealed

the role of the IL-6/Th17 axis in augmenting offspring gut immunity, indicating that prenatal

immunological events can induce long-lasting effects[67]. Transient exposure to a maternally-

derived inflammatory cytokine, IL-6, was similarly explored in pregnant mice and found to

result in permanent alteration of neuronal gene programs, providing mechanistic evidence for

how prenatal inflammation drives long-term changes in neonatal neurodevelopment[68]. Strikingly,

in the current study, we report that prenatal treatment with M2 macrophages mitigates the

negative consequences of *in utero* sterile inflammation, enhancing survival, reversing growth

restriction, and overall improving neonatal infection resistance. Therefore, the current study

supports two key concepts: firstly, acute insults *in utero* can disrupt developmental

programming, resulting in impaired immunocompetence in the offspring, the full extent and

significance of which remain poorly understood; and second, the prenatal cellular therapy using

M2 macrophages demonstrated herein exerts substantial homeostatic effects, effectively

countering such alterations.

Proper formation of the gut microbiota during early life is essential for the development

of a healthy adult gut[69]. Indeed, multiple inflammatory, metabolic, neurologic, cardiovascular,

and gastrointestinal diseases have been linked to disruption of the microbiota during early life[70].

Importantly, the risk of gut dysbiosis is increased in preterm neonates compared to those

delivered at term[69], and thus the prevention or treatment of premature labor leading to preterm

birth, particularly in the context of *in utero* sterile inflammation, is essential to ensure the

development of a healthy gut microbiota and avoid long-term consequences. Several prenatal

events have been studied and found to contribute to microbiota dysbiosis in human neonates,

including maternal diet and antibiotic use, among others[69,71]. Specifically, the combined presence

of acute histologic chorioamnionitis/funisitis was linked to altered microbial composition in

neonatal fecal samples, including modified abundance of *Bacteroidetes* and *Fusobacteria*[72].

Importantly, neonates affected by acute histologic chorioamnionitis alone or acute histologic

chorioamnionitis/funisitis had higher incidence of late-onset sepsis/death, thereby providing an

association between microbiome alterations and short-term adverse events[72]. Consistent with

these studies, we demonstrated herein that adverse neonatal outcomes resulting from *in utero*

sterile inflammation are similarly linked to alterations in gut microbiome composition, and that

prenatal treatment with M2 macrophages improves these aberrations. Although the current study

did not continuously examine the neonatal gut microbiome from birth to the weaning period, our

results indicate that treatment with M2 macrophages promotes the proper maturation of a

"healthy" gut microbiota, which can contribute to reducing disease risk in neonatal life.

It is worth acknowledging that there are inherent differences between humans and murine

models. Key physiological factors such as gestational length, placental structure, immune system

regulation, and hormonal regulation vary between humans and mice, as has been extensively

described[73-75]. Therefore, we consider that any interventions that prove effective in murine

models, such as the M2 macrophages demonstrated in the current study, require additional

validation in models that more closely resemble human, such as non-human primates, to more

firmly establish their clinical relevance during pregnancy.

Collectively, the data presented herein link a reduction in maternal homeostatic

macrophages to human labor and mechanistic evidence in mice demonstrating that M2

macrophages can treat *in utero* sterile inflammation, thereby preventing preterm birth and

adverse neonatal outcomes. Specifically, treatment with M2 macrophages mitigates

inflammatory responses in the amniotic cavity and surrounding fetal membranes by inhibiting

inflammasome activation and halting inflammatory processes in the common pathway of labor,

thereby extending gestational length and preventing preterm birth. Importantly, M2 macrophage

treatment not only diminishes inflammation in the placenta as well as the fetal and neonatal lung

but also enhances neonatal intestinal responses. Neonates born to dams prenatally treated with

M2 macrophages demonstrate a distinct gut microbiome compared to those born from dams with

in utero sterile inflammation, alongside an improved ability to clear bacterial infections. In

conclusion, these findings provide mechanistic evidence that M2 macrophages can serve as a

cellular approach not only to prevent premature labor leading to preterm birth but also to

enhance fetal and neonatal immunity following *in utero* exposure to sterile inflammation.

Acknowledgements

This research was supported by the Perinatology Research Branch, Division of Obstetrics and Maternal-Fetal Medicine, Division of Intramural Research, *Eunice Kennedy Shriver* National Institute of Child Health and Human Development, National Institutes of Health, U.S. Department of Health and Human Services (NICHD/NIH/DHHS) under Contract No. HHSN275201300006C. This research was also supported by the Wayne State University Perinatal Initiative in Maternal, Perinatal and Child Health. R.R. has contributed to this work as part of his official duties as an employee of the United States Federal Government. The funders had no role in study design, data collection and interpretation, or the decision to submit the work for publication. Figures include art created with BioRender.com

Author contributions

Conceptualization: N.G.-L.

Methodology: V.G.-F., Z.L., Y.X., N.G.-L.

Validation: V.G.-F., Z.L., Y.X., N.G.-L.

Formal analysis: V.G.-F., Z.L., R.R., R.P.-R., Y.X., D.L., J.G., A.D.W., M.F.-J., J.J.P., N.G.-L.

Investigation: V.G.-F., Z.L., R.P.-R., Y.X., D.M., D.L., J.G., A.D.W., M.F.-J., J.J.P., N.G.-L.

Resources: R.R., K.R.T., N.G.-L.

Data curation: V.G.-F., Z.L., R.P.-R., Y.X., A.D.W.

Writing – original draft: V.G.-F., D.M., J.G., N.G.-L.

Writing – review & editing: V.G.-F., Z.L., R.R., R.P.-R., Y.X., D.M., D.L., J.G., A.D.W., M.F.-J., J.J.P., K.R.T., N.G.-L.

Visualization: V.G.-F., Z.L., J.G., A.D.W., N.G.-L.

Supervision: R.R., R.P.-R., K.R.T., N.G.-L.

Project administration: N.G.-L.

Funding acquisition: R.R., N.G.-L.

Conflict of interest

The authors declare no competing interests.

Figure Legends

Figure 1. Single-cell atlases of the human maternal-fetal interface reveal that maternal

homeostatic macrophages are reduced with labor. (**A**) Representative diagram showing the

placenta, extraplacental membranes, and myometrium, along with our working hypothesis that a

reduction in homeostatic macrophages may accompany the processes of labor. (**B**) Uniform

Manifold Approximation and Projection (UMAP) plots showing seven distinct macrophage

clusters (M1 – M7) in (*Top row*) the myometrium from women who delivered at term with

(n=11) or without (n=13) labor, (*Middle row*) the placenta (placental villi and basal plate) from

women who delivered at term with (n=27) or without (n=21) labor or preterm after preterm labor

(n=3), and (*Bottom row*) extraplacental membranes from women who delivered at term with

(n=27) or without (n=21) labor or preterm after preterm labor (n=3). (**C**) UMAP plots showing

maternal (blue) or fetal (red) origin of all macrophage clusters in the (*top to bottom*)

myometrium, placenta, and extraplacental membranes. (**D**) Plot showing the proportions of

maternal M2 macrophages in the myometrium from women with term non-labor (TNL) or those

with term labor (TIL). (**E**) Plot showing the proportions of maternal M2 macrophages in the

basal plate from women with term in labor (TIL) or preterm labor (PTL). P-values were

determined using Mann-Whitney U-tests. **p < 0.01. (**F**) STRING analysis showing the top 20

marker genes from the M2 cluster. (**G**) Over-representation analysis showing enriched Gene

Ontology processes in the M2 cluster. Dot size corresponds to gene count and color scaling

represents false discovery rate-adjusted p-values (q<0.05) as determined by Wilcoxon Rank Sum

test. See also Figure S1 and Table S1.

Figure 2. M2 macrophages prevent preterm birth and neonatal mortality induced by *in*

***utero* sterile inflammation. (A)** To induce *in utero* sterile inflammation, the alarmin HMGB1

was intra-amniotically administered to C57BL/6 dams under ultrasound guidance on 14.5 days

post coitum (dpc). Bone marrow-derived cells were collected from C57BL/6 mice, differentiated,

and polarized to an M2 phenotype (M2 MΦ) *in vitro*. M2 MΦ were administered intravenously

to C57BL/6 dams on 13.5 and 14.5 dpc. Dams were monitored until delivery, and neonatal

survival and weight were recorded until three weeks of age. **(B)** Gestational length shown as box

plots where midlines indicate medians and whiskers indicate minimum/maximum range. P-

values were determined using the Kruskal-Wallis test followed by two-stage linear step-up

procedure of Benjamini, Krieger, and Yekutieli post-hoc test. **(C)** Rates of preterm birth among

dams injected with PBS (n = 20), HMGB1 (n = 28), PBS + M2 MΦ (n = 14), and HMGB1 + M2

MΦ (n = 29) are shown as bar plots. P-values were determined using two-sided Fisher's exact

test. **(D)** Pie charts representing the survival at birth of preterm neonates. PBS (n = 20 litters),

HMGB1 (n = 20 litters), PBS + M2 MΦ (n = 14 litters), and HMGB1 + M2 MΦ (n = 16 litters).

(E) Kaplan-Meier survival curves from neonates at weeks 1, 2, and 3 of life. PBS (n = 20 litters),

HMGB1 (n = 20 litters), PBS + M2 MΦ (n = 14 litters), and HMGB1 + M2 MΦ (n = 16 litters).

P-values were determined using the Gehan-Breslow-Wilcoxon test. **(F)** Individual weights

(grams, g) of neonates across the first three weeks of life are shown as box plots where midlines

indicate medians and whiskers indicate minimum/maximum range. PBS (n = 11 litters), HMGB1

(n = 11 litters), PBS + M2 MΦ (n = 11 litters), and HMGB1 + M2 MΦ (n = 14 litters). P-values

were determined using the Kruskal-Wallis test followed by Dunn's post-hoc test. *$p < 0.05$; **p

< 0.01; ***$p < 0.001$.

Figure 3. M2 macrophages dampen HMGB1-induced *in utero* **sterile inflammation,**

including inflammasome activation, in the amniotic cavity. (A) Dams were intra-amniotically

injected with HMGB1 on 14.5 days *post coitum* (dpc). Amniotic fluid was collected at 24, 48,

72, or 96 h post-HMGB1 injection for cytokine determination. **(B)** Concentrations of IL-6 in the

amniotic fluid of HMGB1-injected dams at 24, 48, 72, and 96 h post-injection are shown as box

plots (n = 8 dams per time point). P-values were determined using the Kruskal-Wallis test

followed by two-stage linear step-up procedure of Benjamini, Krieger, and Yekutieli post-hoc

test. **(C)** M2 macrophages (M2 MΦ) or vehicle (PBS) were intravenously administered on 13.5

dpc and 14.5 dpc to dams followed by ultrasound-guided intra-amniotic injection of HMGB1 on

14.5 dpc. Amniotic fluid was collected at 72 h post-HMGB1 injection for cytokine

determination. **(D)** Concentrations of IL-6, TNF, and total IL-1β in the amniotic fluid of the

HMGB1+Vehicle dams (n = 16 dams) or HMGB1+M2 MΦ dams (n = 16 dams) at 72 h post-

injection are shown as box plots. P-values were determined using the two-tailed Mann-Whitney

U-test. **(E)** To determine inflammasome activation, amniotic fluid of dams that received PBS,

HMGB1, HMGB1 + Vehicle, or HMGB1 + M2 MΦ was collected at 72 h post-HMGB1

injection for immunoblotting. **(F)** Immunoblotting of active caspase (CASP)-1 expression and

mature IL-1β expression in the amniotic fluid of dams injected with PBS or HMGB1.

Representative CASP-1 immunoblot image shows 6 samples per group, and representative

mature IL-1β immunoblot image shows 3 pooled samples (pooled amniotic fluids from 3 dams)

per group. **(G)** Protein quantification of active CASP-1 in the amniotic fluid of dams injected

with PBS (n = 12) or HMGB1 (n = 13). Protein quantification of mature IL-1β in the pooled

amniotic fluid of dams injected with PBS (n = 3) or HMGB1 (n = 3). Data are shown as boxplots

where midlines indicate medians, boxes denote interquartile ranges, and whiskers indicate the

minimum/maximum range. P-values were determined using the one-tailed Mann-Whitney U-

test. **(H)** Immunoblotting of active CASP-1 expression and mature IL-1β expression in the

amniotic fluid of HMGB1+Vehicle or HMGB1+M2 MΦ dams. Representative CASP-1

immunoblot image shows 6 samples per group, and representative mature IL-1β immunoblot

image shows 3 pooled samples (pooled amniotic fluids from 3 dams) per group. **(I)** Protein

quantification of active CASP-1 in the amniotic fluid of dams injected with HMGB1+Vehicle (n

= 12) or HMGB1+M2 MΦ (n = 14). Protein quantification of mature IL-1β in the pooled

amniotic fluid of dams injected with HMGB1+Vehicle (n = 3) or HMGB1+M2 MΦ (n = 3). P-

values were determined using the one-tailed Mann-Whitney U-test. Data are shown as boxplots

where midlines indicate medians, boxes denote interquartile ranges, and whiskers indicate the

minimum/maximum range. *$p < 0.05$; **$p < 0.01$; ***$p < 0.001$. See also Figure S3.

Figure 4. M2 macrophages inhibit inflammasome activation and regulate gene expression

in the fetal membranes. (A) Dams received intra-amniotic injection of PBS (control) or

HMGB1 on 14.5 days *post coitum* (dpc). M2 macrophages (M2 MΦ) or vehicle (PBS) were

intravenously administered to dams on 13.5 dpc and 14.5 dpc followed by intra-amniotic

injection of HMGB1 on 14.5 dpc. Tissue collection was performed at 72 h post-injection to

collect the fetal membranes for immunoblotting and to determine gene expression. **(B)**

Immunoblotting of active caspase (CASP)-1, mature IL-1β, and β-actin (ACTB) in the fetal

membranes of PBS- or HMGB1-injected dams. Representative immunoblot images depict 6

samples per group in each gel. **(C)** Protein quantification of active CASP-1 and mature IL-1β

(both normalized by ACTB) in the fetal membranes of PBS- or HMGB1-injected dams (n = 6

per group). **(D)** Immunoblotting of active CASP-1, mature IL-1β, and ACTB in the fetal

membranes from HMGB1+Vehicle or HMGB1+M2 MΦ dams. Representative immunoblot

images depict 6 samples per group in each gel. **(E)** Relative quantification of active CASP-1 and

mature IL-1β (both normalized by ACTB) in the fetal membranes of HMGB1+Vehicle or

HMGB1+M2 MΦ dams (n = 6 per group). **(F)** Representative heatmaps displaying the

expression of key inflammatory genes in the fetal membranes of PBS- (n = 8) or HMGB1-

injected (n = 8) dams. **(G)** Expression of *Ccl17* in the fetal membranes of PBS- or HMGB1-

injected dams. Data are shown as boxplots where midlines indicate medians, boxes denote

interquartile ranges, and whiskers indicate the minimum/maximum range. P-values were

determined using the two-tailed Mann-Whitney U-test. **(H)** Representative heatmaps displaying

the expression of key inflammatory genes in the fetal membranes of HMGB1+Vehicle (n = 8) or

HMGB1+M2 MΦ (n = 8) dams. **(I)** Expression of *Tlr4*, *Cxcl1, Tnf,* and *Il1b* in the fetal

membranes of HMGB1+Vehicle or HMGB1+M2 MΦ dams. Data are shown as boxplots where

midlines indicate medians, boxes denote interquartile ranges, and whiskers indicate the

minimum/maximum range. P-values were determined using the two-tailed Mann-Whitney U-

test. *p < 0.05; **p < 0.01.

Figure 5. M2 macrophages ameliorate fetal and neonatal lung inflammation induced by *in*

***utero* sterile inflammation. (A)** Dams received intra-amniotic injection of PBS (control) or

HMGB1 on 14.5 days *post coitum* (dpc). M2 macrophages (M2 MΦ) or vehicle (PBS) were

intravenously administered to dams on 13.5 dpc and 14.5 dpc followed by intra-amniotic

injection of HMGB1 on 14.5 dpc. Tissue collection was performed at 72 h post-injection to

collect the fetal lung for gene expression. **(B)** Representative heatmaps displaying the expression

of inflammatory genes in the fetal lung of PBS- (n = 8) or HMGB1-injected (n = 8) dams. **(C)**

Expression of *Il6*, *Tnfrsf1a*, *Il33*, *Nlrp6*, and *Tlr9* in the fetal lung of PBS- or HMGB1-injected

dams. Data are shown as boxplots where midlines indicate medians, boxes denote interquartile

ranges, and whiskers indicate the minimum/maximum range. P-values were determined using the

two-tailed Mann-Whitney U-test. **(D)** Representative heatmaps displaying the expression of

inflammatory genes in the fetal lung of HMGB1+Vehicle (n = 8) or HMGB1+M2 MΦ (n = 8)

dams. **(E)** Expression of *Tnfrsf1a* and *Nod1* in the fetal lung of HMGB1+Vehicle or

HMGB1+M2 MΦ dams. **(F)** The lung was collected at 14-16 days of life from neonates born to

HMGB1+Vehicle or HMGB1+M2 MΦ dams to evaluate gene expression as shown in the

representative heatmap. **(G)** Expression of *Il6*, *Ccl2*, *Socs3*, and *Cxcl1* in the neonatal lung (n =

10 per group). Data are shown as boxplots where midlines indicate medians, boxes denote

interquartile ranges, and whiskers indicate the minimum/maximum range. P-values were

determined using the two-tailed Mann-Whitney U-test. *p < 0.05; **p < 0.01.

Figure 6. M2 macrophages regulate neonatal gut inflammation resulting from *in utero*

sterile inflammation. (A) M2 macrophages (M2 MΦ) or PBS (Vehicle) were intravenously

administered to dams on 13.5 dpc and 14.5 dpc followed by intra-amniotic injection of HMGB1

on 14.5 dpc. After delivery, neonates were monitored until 14-16 days of age, after which the

small intestine, cecum, and colon were collected to determine gene expression. Representative

heatmaps display gene expression in the **(B)** neonatal small intestine, **(D)** cecum, and **(F)** colon

of neonates born to HMGB1+Vehicle (n = 10 neonates) or HMGB1+M2 MΦ (n = 10 neonates)

dams. The expression of specific genes in the **(C)** neonatal small intestine (*Aim2, Cd68, Tnf,*

Rgs4, Ccl5, Ccl17), **(E)** cecum (*Casp1* and *Ccl17*), and **(G)** colon (*Il1a, Cd68, Casp1, Jun,*

Kdm6b, Ccl17) are shown as box plots. P-values were determined using the two-tailed Mann-

Whitney U-test. Data are shown as boxplots where midlines indicate medians, boxes denote

interquartile ranges, and whiskers indicate the minimum/maximum range. $*p < 0.05$; $**p < 0.01$.

See also Figures S8-S10.

Figure 7. M2 macrophages improve neonatal immunocompetence. (A) M2 macrophages (M2

MΦ) or PBS (Vehicle) were intravenously administered to dams on 13.5 dpc and 14.5 dpc

followed by intra-amniotic injection of HMGB1 on 14.5 dpc. Pregnant dams without any

treatment or injected with HMGB1 alone were also included. After delivery, surviving neonates

were monitored until 14-16 days of age, after which they received intraperitoneal injection of

Group B *Streptococcus* (GBS) and were observed for five days. **(B)** The survival rates of GBS-

infected neonates born to untreated (n = 10 neonates), HMGB1-injected (n = 10 neonates),

HMGB1+Vehicle (n = 12 neonates), or HMGB1+M2 MΦ (n = 12 neonates) dams over the five

days post-challenge are displayed as Kaplan-Meier survival curves. P-values were determined

using the Gehan-Breslow-Wilcoxon test. **(C)** Mean weights of GBS-infected neonates over the

five days post-challenge. P-values for comparing the mean weight at the end of day five (final

data point in plot) were determined using the two-tailed Mann-Whitney U-test. $*p < 0.05$; $**p <$

0.01; $***p < 0.001$. See also Figure S11.

STAR METHODS

KEY RESOURCES TABLE

(Submitted separately)

RESOURCE AVAILABILITY

Lead Contact

Further information and requests for resources and reagents should be directed to and will

be fulfilled by the lead contact, Nardhy Gomez-Lopez (nardhy.gomez-lopez@wayne.edu).

Materials Availability

This study did not generate new unique reagents.

Data and Code Availability

Single-cell RNA-sequencing data were previously reported[44-46] and are deposited in the

NIH dbGAP database (phs001886.v5.p1). 16S rRNA gene sequencing files and associated

metadata have been uploaded to the National Center for Biotechnology Information's Sequence

Read Archive (PRJNA925285). All other data are presented within the current manuscript and/or

its supplementary materials.

EXPERIMENTAL MODEL AND SUBJECT DETAILS

C57BL/6 (strain # 000664) and B6 CD45.1 (B6.SJL-*Ptprc^a Pepc^b*/BoyJ; strain # 002014)

mice were purchased from The Jackson Laboratory (Bar Harbor, ME, USA). Mice were housed

in the animal care facility at the C.S. Mott Center for Human Growth and Development at

Wayne State University (Detroit, MI, USA) under a circadian cycle (light:dark = 12:12 h). Eight-

to twelve-week-old females were mated with males of proven fertility. Females were checked

daily between 8 and 9 a.m. to investigate the appearance of a vaginal plug, which indicated 0.5

days *post coitum* (dpc). Plugged females were then housed separately from the male mice and

their weights were monitored daily. A weight gain of $\geq$ 2 grams by 12.5 dpc confirmed

pregnancy. Mice were randomly assigned to study groups prior to the experiments described

herein. Numbers of biological replicates are indicated in each figure caption. All procedures

were approved by the Institutional Animal Care and Use Committee (IACUC) at Wayne State

University under Protocol Nos. A-07-03-15, 18-03-0584, and 21-04-3506.

METHOD DETAILS

Analysis of scRNA-seq data

Data normalization and pre-processing

Data from three of our previously generated human scRNA-seq datasets[44-46] were

combined and re-analyzed to specifically investigate macrophage populations at the maternal-

fetal interface. All datasets were pre-processed and normalized with a comparable pipeline. In

brief, sequencing data were processed using Cell Ranger version 4.0.0 (10x Genomics). The

fastq files were aligned using kallisto[76], and bustools[77] was used to summarize the cell/gene

transcript counts in a matrix for each sample. In parallel, "cellranger counts" was also used to

align the scRNA-seq reads using the STAR aligner[78] to produce the bam files necessary for

demultiplexing the individual of origin based on genotype information using souporcell[79] and

demuxlet[80]. Quality metrics were calculated and each library was determined to be of excellent

quality based on 10X Genomics recommendations. Any droplet/GEM barcode assigned to a

double or ambiguous cell in demuxlet or souporcell was filtered, and only cells that could be

assigned to a pregnancy case were kept. Furthermore, any cells with less than 100 or more than

10,000 genes were filtered out, as well as those with > 25% mitochondrial reads. Cells annotated

as macrophages (details regarding the annotation process can be found in the original

publications[44-46]) were extracted and combined into a new Seurat object.

All count data matrices were normalized and combined using the "NormalizedData"

"FindVariableFeatures" and "ScaleData" methods implemented in the Seurat package in R[81,82].

Next, the Seurat "RunPCA" function was applied to obtain the first 50 principal components, and

the different libraries were integrated and harmonized using the Harmony package in R[83] while

accounting for library of origin as a potential batch effect. The top 30 harmony components were

then processed using the Seurat "runUMAP" function to embed and visualize the cells in a two-

dimensional map via the Uniform Manifold Approximation and Projection for Dimension

Reduction (UMAP) algorithm. The "FindClusters" function with a resolution of 0.1 was used to

cluster the single cells into seven distinct clusters. The two most abundant macrophage cell type

clusters (M1 and M2) were found to be equivalent to the previously annotated clusters

Macrophage-1 and Macrophage-2, respectively that we have previously reported[44-46]. The

proportions of specific macrophage clusters within a tissue were compared between groups using

Mann-Whitney U-tests.

Gene ontology

The differential expression of selected marker genes for each cell type/cluster were

identified using the Wilcoxon Rank Sum test implemented by FindAllMarkers function from

Seurat. For this analysis, we compared each cluster to all cell types. The clusterProfiler[84] was

used to perform Over-Representation Analysis (ORA) based on the Gene Ontology (GO) using

"enrichGO" function. We first compared all clusters to each other using "compareCluster" and

reported the top terms for each cluster. Since many of these terms were shared, we then focused

on the M2 cluster and compared the list of unique genes identified with a FDR of 5% for this

population against the universe of all genes expressed in macrophages. Only ORA results that

were significant after correction were reported with $q < 0.05$ being considered statistically

significant.

Intra-amniotic administration of HMGB1

Ultrasound-guided intra-amniotic injection of HMGB1 was performed as previously

reported[27-29]. Briefly, dams were anesthetized on 14.5 dpc by inhalation of 2% isoflurane

(Fluriso[TM]/Isoflurane, USP; VetOne, Boise, ID, USA) and 1–2 L/min of oxygen in an induction

chamber, and a mixture of 1.5–2% isoflurane and 1.5–2 L/min of oxygen was used to maintain

anesthesia. Mice were positioned on a heating pad and stabilized with adhesive tape, and fur was

removed from the abdomen and thorax using Nair cream (Church & Dwight Co., Inc., Ewing,

NJ, USA). Body temperature was detected with a rectal probe (VisualSonics, Toronto, ON,

Canada) throughout the procedure, and respiratory and heart rates were monitored by electrodes

embedded in the heating pad. An ultrasound probe was fixed and mobilized with a mechanical

holder, and the transducer was slowly moved toward the abdomen[85]. The ultrasound-guided

intra-amniotic injection of recombinant human HMGB1 (Biolegend, San Diego, CA, USA) at a

concentration of 9 ng dissolved in 100 μL of sterile 1X phosphate-buffered saline (PBS; Life

Technologies, Grand Island, NY, USA) was performed in each amniotic sac using a 30G needle

(BD PrecisionGlide Needle; Becton Dickinson, Franklin Lakes, NJ, USA). After ultrasound

completion, mice were placed under a heat lamp for recovery defined as when the mouse

resumed normal activities, such as walking and interacting with its environment, which typically

occurred within 10 min after removal from anesthesia. After recovery, dams were monitored via

video camera (Sony Corporation, Tokyo, Japan) to observe pregnancy and neonatal outcomes.

Isolation, differentiation, and adoptive transfer of M2 macrophages

Bone marrow cells were isolated, differentiated, and adoptively transferred following a

previously established protocol[43]. Briefly, bone marrow cells were collected from female mice

(12-16 weeks old) and treated with red blood cell lysis buffer (Ammonium Chloride Solution;

Cat# 07800, Stem Cell Technologies; Vancouver, CA). Then, cells were resuspended and

cultured in Iscove's Modified Dulbecco's Media (IMDM) medium (Thermo-Fisher Scientific;

Waltham, MA, USA) supplemented with 10% fetal bovine serum (FBS; Invitrogen, Carlsbad,

CA, USA), 1X antibiotic-antimycotic (Cat# 15240062; Thermo-Fisher Scientific), and 10 ng/ml

recombinant macrophage colony-stimulating factor (M-CSF; Cat# 576402; BioLegend, San

Diego, CA, USA) for 7 days. On day 7, the culture medium was replaced with fresh IMDM

medium supplemented with 10% FBS, 1X antibiotic-antimycotic, 10 ng/ml of recombinant IL-4

(Cat# 574302, BioLegend), and 10 ng/ml of recombinant IL-13 (Cat# 575902, BioLegend). After

14-18 h, M2 macrophages were collected by washing with ice-cold PBS (Fisher Scientific

Bioreagents, Fair Lawn, NJ, USA or Life Technologies Limited, Pailey, UK). The purity and

phenotype of macrophages after M2 polarization were routinely checked throughout the course

of the study, as we have previously reported[43]. Prior to injection, the viability of M2

macrophages was consistently confirmed to be >90%. Approximately 2-5 x 10^6 cells were

resuspended in 100 μL PBS for intravenous injection into dams on 13.5 and 14.5 dpc, prior to the

administration of HMGB1 or PBS on 14.5 dpc. This timing was chosen so that the transferred

M2 macrophages would have sufficient time to reach key maternal and fetal tissues during the

critical window of inflammation resulting in preterm birth, as previously reported[43].

Perinatal outcomes

Gestational length was calculated as the duration of time from the presence of the vaginal

plug (0.5 dpc) to the detection of the first pup in the cage bedding. Preterm birth was defined as

delivery occurring before 18.75 dpc based on the lowest gestational age at delivery observed in

the control group. The rate of preterm birth was calculated as the proportion of females

delivering preterm out of the total number of mice per group. The rate of stillbirth was defined as

the proportion of delivered pups found dead among the total number of pups. The rate of

neonatal mortality was defined as the proportion of pups found dead among the total number of

pups. Neonatal mortality rates and weights were calculated and recorded at postnatal weeks 1, 2,

and 3.

Sampling from dams intra-amniotically injected with HMGB1

Dams were euthanized at 24, 48, 72, and 96 h after intra-amniotic HMGB1 injection for

tissue collection. The amniotic fluid was collected from each amniotic sac and centrifuged at

1,300 x g for 5 min at 4°C. The resulting supernatants were stored at -20°C until analysis. The

maternal whole blood was collected from each dam, mixed with heparin (Cat# 2106-15VL;

Sigma-Aldrich, St. Louis, MO, USA), and centrifuged at 800 x g for 10 min at 4°C. Collection of

the uterus, decidua, placenta, fetal membrane, fetal lung, and fetal intestine was performed. The

placentas and fetuses were imaged and weighed during tissue dissection. The collected tissues

were snap-frozen in liquid nitrogen for immunoblotting or preserved in RNA*later* Stabilization

Solution (Cat# AM7021; Invitrogen by Thermo-Fisher Scientific, Carlsbad, CA, USA),

according to the manufacturer's instructions.

Determination of cytokine concentrations in amniotic fluid

Concentrations of IL-6, IL-1β, and TNF were determined in murine amniotic fluid using

the U-PLEX Custom Biomarker Group 1 (ms) assay (Cat# K15069M-2; Meso Scale Discovery,

Rockville, MD, USA), according to the manufacturer's instructions. Plates were read using the

MESO QuickPlex SQ 120 (Meso Scale Discovery) and analyte concentrations were calculated

using the Discovery Workbench software v4.0 (Meso Scale Discovery). The sensitivities of IL-6,

IL-1β, and TNF were 4.8 pg/ml, 3.1 pg/ml, and 1.3 pg/ml, respectively.

RNA isolation, cDNA synthesis, and qPCR analysis of tissues from pregnant mice

Total RNA was isolated from the uterus, decidua, placenta, fetal membranes, fetal lung,

and fetal intestine using QIAshredders (Cat# 79656; Qiagen, Germantown, MD, USA), RNase-

Free DNase Sets (Cat# 79254; Qiagen), and RNeasy Mini Kits (Cat# 74106; Qiagen), according

to the manufacturer's instructions. The NanoDrop 8000 spectrophotometer (Thermo Scientific,

Wilmington, DE, USA) and the Bioanalyzer 2100 (Agilent Technologies, Waldbronn, Germany)

were used to evaluate RNA concentrations, purity, and integrity. SuperScript IV VILO Master

Mix (cat# 11756050; Invitrogen by Thermo Fisher Scientific Baltics UAB, Vilnius, Lithuania)

was used to synthesize complementary (c)DNA. Gene expression profiling of the tissues was

performed on the BioMark System for high-throughput qPCR (Fluidigm, San Francisco, CA,

USA) with the TaqMan gene expression assays (Applied Biosystems, Life Technologies

Corporation, Pleasanton, CA, USA) listed in Supplementary Table 2. Negative delta cycle

threshold ($-\Delta C_T$) values were determined using multiple reference genes (*Actb*, *Gapdh*, *Gusb*,

and *Hsp90ab1*) averaged within each sample for contractility-associated and inflammatory

genes. The $-\Delta C_T$ values were normalized by calculating the Z-score of each gene with the study

groups. Heatmaps were created representing the Z-score of each $-\Delta C_T$ using GraphPad Prism

(GraphPad, San Diego, CA, USA).

Tissue lysate preparation and protein quantification

Snap-frozen fetal membrane and uterus were mechanically homogenized in PBS

containing a complete protease inhibitor cocktail (Cat# 11836170001; Roche Applied Sciences,

Mannheim, Germany) for 10 min. The resulting lysates were centrifuged at 16,100 x g

(maximum speed) for 5 min at 4°C. The total protein concentrations in amniotic fluid

(supernatant) and tissue lysate samples were tested by the Pierce BCA Protein Assay Kit (Cat#

23225; Pierce Biotechnology, Thermo-Fisher Scientific, Inc., Rockford, IL, USA) prior to

immunoblotting. Cell lysates or concentrated culture medium from murine bone marrow-derived

macrophages were utilized as positive controls for the expression of active CASP-1 and mature

IL-1β, as previously described[54].

Immunoblotting

Cell lysates or culture supernatants from murine bone marrow-derived macrophages (30

µg per well; positive control), amniotic fluid (20 µg total protein per well), fetal membrane

lysates (75 µg per well), and uterine tissue lysates (75 µg per well) were subjected to

electrophoresis in sulfate-polyacrylamide gels (Cat# NP0336BOX; Invitrogen). The proteins

were transferred onto nitrocellulose membranes (Cat# 1620145; Bio-Rad, Hercules, CA, USA),

which were then submerged in 5% blotting-graded blocking solution (Cat# 1706404; Bio-Rad)

for 30 min at room temperature. Next, the membranes were probed overnight at 4°C with anti-

mouse CASP-1 (Cat# 14-9832-82; Thermo-Fisher Scientific) or anti-mouse mature IL-1β (Cat#

63124S; Cell Signaling Technology, Danvers, MA, USA; only for fetal membranes). After

incubation with each primary antibody, membranes were incubated with HRP-conjugated anti-

rat IgG (Cat# 7077S; Cell Signaling) for CASP-1 or HRP-conjugated anti-heavy chain of rabbit

IgG (Cat# HRP-66467; Proteintech, Rosemont, IL, USA) for mature IL-1β for 1 h at room

temperature. The ChemiGlow West Chemiluminescence Substrate Kit (Cat# 60-12596-00;

ProteinSimple, San Jose, CA, USA) was used to detect chemiluminescence signals, and

corresponding images were acquired using the ChemiDoc Imaging System (Bio-Rad). The

membranes that were loaded with fetal membrane and uterine tissue lysates were re-probed for 1

h at room temperature with a mouse anti-β-actin (ACTB) monoclonal antibody (Cat# A5441,

Sigma-Aldrich). Then, the membranes were incubated with HRP-conjugated anti-mouse IgG

(Cat# 7076S; Cell Signaling) for 30 min at room temperature. The chemiluminescence signals of

the ACTB were detected as shown above. Finally, quantification was performed using ImageJ

software as previously reported [54]. In short, each individual protein band was automatically

quantified on the blot images. The internal control, β-actin, was used to normalize the target

protein expression in each fetal membrane and uterine tissue sample for relative quantification.

Immunoprecipitation of mature IL-1β

Immunoprecipitation of cleaved IL-1β from uterine tissue lysates and amniotic fluid was performed using the Pierce™ Classic IP Kit (Cat# 26146; Thermo Fisher Scientific) following the manufacturer's instructions. Amniotic fluid samples (n = 3 per study group) were pooled. Culture supernatants from murine bone marrow-derived macrophages were utilized as positive controls. Each uterine tissue lysate (1 mg protein) or pooled amniotic fluid sample (1 mg protein) was pre-cleared using the control agarose resin and incubated with rabbit anti-mouse mature IL-1β antibody overnight at 4°C to form the immune complex. Next, the immune complex was captured using Pierce Protein A/G Agarose. After several washes to remove non-bound proteins, the immune complex was eluted with sample buffer and subjected to electrophoresis in 4%−12% polyacrylamide gels followed by Western blot transfer, as described above. The blot was then incubated with rabbit anti-mouse mature IL-1β antibody, followed by incubation with an HRP-conjugated anti-heavy chain of rabbit IgG antibody. Finally, chemiluminescence signals were detected with the ChemiGlow West Substrate Kit and images were acquired using the ChemiDoc Chemiluminescence Imaging System, and quantification was performed using ImageJ software.

Tracking transferred M2 macrophages *in vivo*

Bone marrow from B6 CD45.1 mice was collected, isolated, and differentiated into M2 macrophages, as described above. Pregnant C57BL/6 recipient mice were intravenously injected with 2-5x10^6 CD45.1$^+$ donor M2 macrophages at 13.5 and 14.5 dpc followed by intra-amniotic injection of HMGB1 on 14.5 dpc, as described above. Mice were euthanized at 2, 6, or 12 h after intra-amniotic injection to collect the maternal blood, maternal lung, uterus, decidua, and

placenta as well as the fetal membrane, fetal intestine, fetal liver, fetal liver, and amniotic fluid.

Whole maternal blood (100 µL) was used directly for immunophenotyping following the

procedure described below. For maternal and fetal tissues, leukocytes were isolated by following

protocols adapted from[86]. Briefly, maternal and fetal tissues were gently minced using fine

scissors and enzymatically digested with StemPro Accutase Cell Dissociation Reagent (Cat#

A1110501, Thermo-Fisher) for 30 min at 37°C. Cells were filtered using a 100 μm cell strainer

(Cat# 22-363-549, Fisher Scientific, Fair Lawn, NY, USA) followed by washing with PBS prior

to immunophenotyping. Immediately after isolation of leukocytes, cell pellets were re-suspended

in FACS buffer and pre-incubated with rat anti-mouse CD16/CD32 Fc Block™ (Cat# 553142,

clone 2.4G2; BD Bioscience, Franklin Lakes, NJ, USA) for 10 min on ice and subsequently

incubated with specific fluorochrome-conjugated monoclonal anti-mouse antibodies shown in

Supplementary Table 3. Cells were acquired using the BD LSRFortessa flow cytometer (BD

Biosciences) with FACSDiva 9.0 software (BD Biosciences). CountBright absolute counting

beads (Thermo Fisher Scientific) were added prior to acquisition. The analysis was performed

and the figures were created by using FlowJo software v10 (FlowJo, Ashland, OR, USA).

Sampling and gene expressing profiling of neonates

Neonates born to dams that received M2 macrophages or PBS followed by HMGB1 were

sacrificed at 14 days of life, and the lung and intestine (cecum, colon, and small intestine) were

preserved in RNAlater Stabilization Solution. Gene expression analysis was performed using the

methods described above (see "RNA isolation, cDNA synthesis, and qPCR analysis").

Neonatal sampling for gut microbiome analysis

Dams underwent treatment with M2 macrophages alone, or were injected with HMGB1

and treated with vehicle or M2 macrophages, as described above. A group of non-injected dams

(control) was also included. There was no group-level separation of these dams; each was housed

alone after receiving their respective injections. After delivery, dams were housed with their

neonates. Surviving neonates were euthanized on postnatal day 14-16. Samples from the

neonatal small intestine, cecum, and colon as well as environmental controls were obtained using

sterile swabs (FLOQSwabs, Cat# 553C Copan Diagnostics, Murieta, CA, USA) under aseptic

conditions. Each experimental group included samples obtained from neonates born to three

different dams. All swab samples collected for 16S rRNA gene sequencing were stored at -80°C

until DNA extractions were performed.

DNA extraction

Genomic DNA was extracted from swabs of the small intestine, cecum, and colon (n = 35

each) and negative blank DNA extraction kit controls using the DNeasy PowerLyzer Powersoil

kit (Qiagen, Germantown, MD, USA), with minor modifications to the manufacturer's protocol

as previously described[87-89]. All swabs were randomized across extraction runs. The purified

DNA was stored at -20°C.

16s rRNA Gene Sequencing

The V4 region of the 16S rRNA gene was amplified and sequenced via the dual indexing

strategy developed by Kozich et al.[90] as previously described[87-89,91], with the exception that

library builds were performed using 32 cycles of PCR prior to the equimolar pooling of all

sample libraries for multiplex sequencing. 16S rRNA gene sequences were clustered into

amplicon sequence variants (ASVs) defined by 100% sequence similarity using DADA2 version

1.12[92] in R version 4.2.2[93] as previously described[94], with the exception that forward and reverse

reads were truncated at 240 and 215 bases, respectively. Sequences were then classified using the

silva_nr_v132_train_set database with a minimum bootstrap value of 80%, and sequences that

were derived from Archaea, chloroplast, Eukaryota, or mitochondria were removed.

Potential background DNA contaminant ASVs were identified using the R package

decontam version 1.18.0[95] and the "IsContaminant" function. One of the 82 ASVs identified as a

contaminant [Lachnospiraceae (ASV-2)] was present in > 90% of both biological and control

samples, however, it had a greater mean relative abundance in biological (12.95%) than in

control samples (1.77%) and was therefore not removed from the dataset. The two taxa with the

highest mean prevalence and abundance in control samples that were removed from the dataset

were Burkholderiaceae (ASV-42) (41.7% and 5.3%) and *Pseudomonas* (ASV-97) (50.0% and

3.8%, respectively). The mean relative abundances of these two ASVs across all biological

samples were 0.0003% and 0.006%, respectively. Other ASVs removed from the dataset

included taxa that have previously been reported to be contaminants in 16S rRNA gene

sequencing studies (e.g., *Acinetobacter*, *Cloacibacterium*, *Corynebacterium*, *Cutibacterium*,

Escherichia, *Halomonas*, *Pseudomonas*, *Ralstonia*, *Sphingomonas*, *Staphylococcus*,

Stenotrophomonas, and *Xanthomonas*)[91,96-99].

After removing contaminant ASVs, all samples were normalized to 8,715 reads using the

"rarefy_even_depth" function in phyloseq 1.42.0[100], resulting in 20 bacterial profiles with

Good's coverage[101] values ≥ 99.68% for each of the three sample types. The final dataset

contained a total of 411 ASVs. No cross-tissue comparisons were conducted.

Beta diversity

Beta diversity of bacterial profiles was characterized using the Bray-Curtis similarity

index. Principal coordinate analysis (PCoA) plots were used to visualize the Bray-Curtis

similarity of the sample profiles. Using nonparametric multivariate analysis of variance

(NPMANOVA) as implemented in vegan 2.6.4[102], differences in bacterial community structure

across treatments were evaluated for each tissue type.

Differential abundance of bacterial species

Differential abundance of the 25 most prevalent ASVs between treatment groups was

independently evaluated for the small intestine, cecum, and colon datasets using two-tailed

Mann-Whitney U-tests with Holm's correction for multiple comparisons as implemented in R

version 4.2.2. Adjusted p-values < 0.1 were considered significant.

Neonatal bacterial challenge

Group B *Streptococcus* (GBS, CNCTC 10/84, serotype V, sequence type 26) was

originally isolated from a neonate with sepsis. From an overnight culture, a sub-culture was

placed with fresh Tryptic Soy Broth (Cat# 211825; BD Bioscience, Franklin Lakes, NJ, USA)

and grown to the logarithmic phase (OD600 0.8-0.9). Additional dilution was performed using

sterile PBS to reach working concentrations. Thriving term neonates (14-16 days old) born to

dams that did not receive any treatment (control) and dams injected solely with HMGB1, PBS

(vehicle for M2 macrophages) followed by HMGB1, or M2 macrophages followed by HMGB1

were intraperitoneally injected with 2×10^{6} colony-forming units of GBS in 200 μL sterile PBS.

Survivability, body weight, and rectal temperature of the challenged neonates were checked at 4-

6 h intervals throughout the daytime for 5 days.

QUANTIFICATION AND STATISTICAL ANALYSIS

Statistical analyses were performed using GraphPad Prism (v9.5.0; GraphPad, San Diego,

CA, US) as indicated in each figure caption. Flow cytometry analysis was performed using

FlowJo software v10. Protein expression quantification was performed using ImageJ software.

To determine gene expression levels from qPCR arrays, $-\Delta C_T$ values were calculated using

averaged reference genes (*Actb, Gapdh, Gusb*, and *Hsp90ab1*) within each sample. Heatmaps

were created to represent the Z-scores. Single-cell RNA-sequencing and microbiome data

analysis were performed using R (v.4.2.2; https://www.r-project.org/), as described above. A p-

value ≤ 0.05 or adjusted p-value (q-value) ≤ 0.1 was considered statistically significant.

Supplemental Items

Document S1. Figures S1-S11 and Tables S2-S3

Table S1. Excel file containing additional data too large to fit in a PDF, related to Figure 1

Supplemental Item Legends

Figure S1. Gene Ontology of macrophage populations at the maternal-fetal interface, related to

Figure 1.

Figure S2. Intra-amniotic injection of HMGB1 does not significantly alter amniotic fluid

concentrations of total IL-1β and TNF, related to Figure 3.

Figure S3. M2 macrophages do not significantly alter HMGB1-induced inflammasome

activation in the uterine tissues.

Figure S4. Neither HMGB1 nor M2 macrophage treatment induce changes in uterine gene

expression.

Figure S5. M2 macrophages dampen inflammatory gene expression in the decidua.

Figure S6. Adoptively transferred M2 macrophages accumulate at the maternal-fetal interface

but do not reach fetal organs.

Figure S7. M2 macrophages ameliorate the HMGB1-induced inflammatory response in the

placenta.

Figure S8. HMGB1 regulates gene expression in the fetal intestine, related to Figure 6.

Figure S9. M2 macrophages change the neonatal gut microbiome after in utero exposure to

HMGB1, related to Figure 6.

Figure S10. M2 macrophages modulate microbiome dysbiosis in each compartment of the

neonatal gut, related to Figure 6.

Figure S11. M2 macrophages reduce neonatal hypothermia upon bacterial challenge, related to

Figure 7.

Table S1. Marker genes used to distinguish macrophage clusters M1-M7, related to Figure 1

Table S2. TaqMan assays utilized for qPCR.

Table S3. Antibodies utilized for flow cytometry.

REFERENCES

1. Liu, L., Oza, S., Hogan, D., Perin, J., Rudan, I., Lawn, J.E., Cousens, S., Mathers, C., and Black, R.E. (2015). Global, regional, and national causes of child mortality in 2000-13, with projections to inform post-2015 priorities: an updated systematic analysis. Lancet (London, England) *385*, 430-440. 10.1016/s0140-6736(14)61698-6.

2. Guo, X., Li, X., Qi, T., Pan, Z., Zhu, X., Wang, H., Dong, Y., Yue, H., and Sun, B. (2021). A birth population-based survey of preterm morbidity and mortality by gestational age. BMC Pregnancy Childbirth *21*, 291. 10.1186/s12884-021-03726-4.

3. Romero, R., Dey, S.K., and Fisher, S.J. (2014). Preterm labor: one syndrome, many causes. Science *345*, 760-765. 10.1126/science.1251816.

4. Goldenberg, R.L., Culhane, J.F., Iams, J.D., and Romero, R. (2008). Epidemiology and causes of preterm birth. Lancet (London, England) *371*, 75-84. 10.1016/S0140-6736(08)60074-4.

5. Bastek, J.A., Gomez, L.M., and Elovitz, M.A. (2011). The role of inflammation and infection in preterm birth. Clin Perinatol *38*, 385-406. 10.1016/j.clp.2011.06.003.

6. Romero, R., Miranda, J., Chaiworapongsa, T., Chaemsaithong, P., Gotsch, F., Dong, Z., Ahmed, A.I., Yoon, B.H., Hassan, S.S., Kim, C.J., et al. (2014). A novel molecular microbiologic technique for the rapid diagnosis of microbial invasion of the amniotic cavity and intra-amniotic infection in preterm labor with intact membranes. American journal of reproductive immunology (New York, N.Y. : 1989) *71*, 330-358. 10.1111/aji.12189.

7. Romero, R., Miranda, J., Chaiworapongsa, T., Korzeniewski, S.J., Chaemsaithong, P., Gotsch, F., Dong, Z., Ahmed, A.I., Yoon, B.H., Hassan, S.S., et al. (2014). Prevalence and clinical significance of sterile intra-amniotic inflammation in patients with preterm labor and intact membranes. American journal of reproductive immunology (New York, N.Y. : 1989) *72*, 458-474. 10.1111/aji.12296.

8. Romero, R., Miranda, J., Chaemsaithong, P., Chaiworapongsa, T., Kusanovic, J.P., Dong, Z., Ahmed, A.I., Shaman, M., Lannaman, K., Yoon, B.H., et al. (2015). Sterile and microbial-associated intra-amniotic inflammation in preterm prelabor rupture of membranes. J Matern Fetal Neonatal Med *28*, 1394-1409. 10.3109/14767058.2014.958463.

9. Romero, R., Miranda, J., Chaiworapongsa, T., Chaemsaithong, P., Gotsch, F., Dong, Z., Ahmed, A.I., Yoon, B.H., Hassan, S.S., Kim, C.J., et al. (2015). Sterile intra-amniotic inflammation in asymptomatic patients with a sonographic short cervix: prevalence and clinical significance. J Matern Fetal Neonatal Med *28*, 1343-1359. 10.3109/14767058.2014.954243.

10. Burnham, P., Gomez-Lopez, N., Heyang, M., Cheng, A.P., Lenz, J.S., Dadhania, D.M., Lee, J.R., Suthanthiran, M., Romero, R., and De Vlaminck, I. (2020). Separating the signal from the noise in metagenomic cell-free DNA sequencing. Microbiome *8*, 18. 10.1186/s40168-020-0793-4.

11. Combs, C.A., Gravett, M., Garite, T.J., Hickok, D.E., Lapidus, J., Porreco, R., Rael, J., Grove, T., Morgan, T.K., Clewell, W., et al. (2014). Amniotic fluid infection, inflammation, and colonization in preterm labor with intact membranes. Am J Obstet Gynecol *210*, 125.e121-125.e115. 10.1016/j.ajog.2013.11.032.

12. Motomura, K., Romero, R., Galaz, J., Tarca, A.L., Done, B., Xu, Y., Leng, Y., Garcia-Flores, V., Arenas-Hernandez, M., Theis, K.R., et al. (2021). RNA Sequencing Reveals Distinct Immune Responses in the Chorioamniotic Membranes of Women with Preterm Labor and Microbial or Sterile Intra-amniotic Inflammation. Infect Immun *89*. 10.1128/IAI.00819-20.

13. Lee, H.J., Kim, E.K., Kim, H.S., Choi, C.W., Kim, B.I., and Choi, J.H. (2011). Chorioamnionitis, respiratory distress syndrome and bronchopulmonary dysplasia in extremely low birth weight infants. J Perinatol *31*, 166-170. 10.1038/jp.2010.113.

14. Villamor-Martinez, E., Álvarez-Fuente, M., Ghazi, A.M.T., Degraeuwe, P., Zimmermann, L.J.I., Kramer, B.W., and Villamor, E. (2019). Association of Chorioamnionitis With Bronchopulmonary Dysplasia Among Preterm Infants: A Systematic Review, Meta-analysis, and Metaregression. JAMA Netw Open *2*, e1914611. 10.1001/jamanetworkopen.2019.14611.

15. Costa, S., Fattore, S., De Santis, M., Lanzone, A., Spanu, T., Arena, V., Tana, M., Trapani, M., Sanguinetti, M., Barnea, E.R., and Vento, G. (2023). Effect of acute histologic chorioamnionitis on bronchopulmonary dysplasia and mortality rate among extremely low gestational age neonates: A retrospective case-control study. Int J Gynaecol Obstet. 10.1002/ijgo.15290.

16. Been, J.V., Lievense, S., Zimmermann, L.J., Kramer, B.W., and Wolfs, T.G. (2013). Chorioamnionitis as a risk factor for necrotizing enterocolitis: a systematic review and meta-analysis. J Pediatr *162*, 236-242.e232. 10.1016/j.jpeds.2012.07.012.

17. Duci, M., Frigo, A.C., Visentin, S., Verlato, G., Gamba, P., and Fascetti-Leon, F. (2019). Maternal and placental risk factors associated with the development of necrotizing enterocolitis (NEC) and its severity. J Pediatr Surg *54*, 2099-2102. 10.1016/j.jpedsurg.2019.04.018.

18. Kallapur, S.G., Willet, K.E., Jobe, A.H., Ikegami, M., and Bachurski, C.J. (2001). Intra-amniotic endotoxin: chorioamnionitis precedes lung maturation in preterm lambs. Am J Physiol Lung Cell Mol Physiol *280*, L527-536. 10.1152/ajplung.2001.280.3.L527.

19. Wolfs, T.G., Kramer, B.W., Thuijls, G., Kemp, M.W., Saito, M., Willems, M.G., Senthamarai-Kannan, P., Newnham, J.P., Jobe, A.H., and Kallapur, S.G. (2014). Chorioamnionitis-induced fetal gut injury is mediated by direct gut exposure of inflammatory mediators or by lung inflammation. Am J Physiol Gastrointest Liver Physiol *306*, G382-393. 10.1152/ajpgi.00260.2013.

20. Jung, E., Romero, R., Yeo, L., Diaz-Primera, R., Marin-Concha, J., Para, R., Lopez, A.M., Pacora, P., Gomez-Lopez, N., Yoon, B.H., et al. (2020). The fetal inflammatory response syndrome: the origins of a concept, pathophysiology, diagnosis, and obstetrical implications. Semin Fetal Neonatal Med *25*, 101146. 10.1016/j.siny.2020.101146.

21. Matzinger, P. (1998). An innate sense of danger. Seminars in immunology *10*, 399-415. 10.1006/smim.1998.0143.

22. Rubartelli, A., and Lotze, M.T. (2007). Inside, outside, upside down: damage-associated molecular-pattern molecules (DAMPs) and redox. Trends in immunology *28*, 429-436. 10.1016/j.it.2007.08.004.

23. Zindel, J., and Kubes, P. (2020). DAMPs, PAMPs, and LAMPs in Immunity and Sterile Inflammation. Annu Rev Pathol *15*, 493-518. 10.1146/annurev-pathmechdis-012419-032847.

24. Romero, R., Chaiworapongsa, T., Alpay Savasan, Z., Xu, Y., Hussein, Y., Dong, Z.,
Kusanovic, J.P., Kim, C.J., and Hassan, S.S. (2011). Damage-associated molecular
patterns (DAMPs) in preterm labor with intact membranes and preterm PROM: a study
of the alarmin HMGB1. J Matern Fetal Neonatal Med *24*, 1444-1455.
10.3109/14767058.2011.591460.
25. Romero, R., Grivel, J.C., Tarca, A.L., Chaemsaithong, P., Xu, Z., Fitzgerald, W., Hassan,
S.S., Chaiworapongsa, T., and Margolis, L. (2015). Evidence of perturbations of the
cytokine network in preterm labor. Am J Obstet Gynecol *213*, 836.e831-836.e818.
10.1016/j.ajog.2015.07.037.
26. Bhatti, G., Romero, R., Rice, G.E., Fitzgerald, W., Pacora, P., Gomez-Lopez, N., Kavdia,
M., Tarca, A.L., and Margolis, L. (2020). Compartmentalized profiling of amniotic fluid
cytokines in women with preterm labor. PLoS One *15*, e0227881.
10.1371/journal.pone.0227881.
27. Gomez-Lopez, N., Romero, R., Plazyo, O., Panaitescu, B., Furcron, A.E., Miller, D.,
Roumayah, T., Flom, E., and Hassan, S.S. (2016). Intra-Amniotic Administration of
HMGB1 Induces Spontaneous Preterm Labor and Birth. American journal of
reproductive immunology (New York, N.Y. : 1989) *75*, 3-7. 10.1111/aji.12443.
28. Galaz, J., Romero, R., Arenas-Hernandez, M., Panaitescu, B., Para, R., and Gomez-
Lopez, N. (2021). Betamethasone as a potential treatment for preterm birth associated
with sterile intra-amniotic inflammation: a murine study. J Perinat Med *49*, 897-906.
10.1515/jpm-2021-0049.
29. Galaz, J., Romero, R., Arenas-Hernandez, M., Farias-Jofre, M., Motomura, K., Liu, Z.,
Kawahara, N., Demery-Poulos, C., Liu, T.N., Padron, J., et al. (2022). Clarithromycin
prevents preterm birth and neonatal mortality by dampening alarmin-induced maternal-
fetal inflammation in mice. BMC Pregnancy Childbirth *22*, 503. 10.1186/s12884-022-
04764-2.
30. Plazyo, O., Romero, R., Unkel, R., Balancio, A., Mial, T.N., Xu, Y., Dong, Z., Hassan,
S.S., and Gomez-Lopez, N. (2016). HMGB1 Induces an Inflammatory Response in the
Chorioamniotic Membranes That Is Partially Mediated by the Inflammasome. Biology of
reproduction *95*, 130. 10.1095/biolreprod.116.144139.
31. Gomez-Lopez, N., Romero, R., Panaitescu, B., Leng, Y., Xu, Y., Tarca, A.L., Faro, J.,
Pacora, P., Hassan, S.S., and Hsu, C.D. (2018). Inflammasome activation during
spontaneous preterm labor with intra-amniotic infection or sterile intra-amniotic
inflammation. American journal of reproductive immunology (New York, N.Y. : 1989)
80, e13049. 10.1111/aji.13049.
32. Gomez-Lopez, N., Romero, R., Garcia-Flores, V., Leng, Y., Miller, D., Hassan, S.S.,
Hsu, C.D., and Panaitescu, B. (2019). Inhibition of the NLRP3 inflammasome can
prevent sterile intra-amniotic inflammation, preterm labor/birth, and adverse neonatal
outcomesdagger. Biol Reprod *100*, 1306-1318. 10.1093/biolre/ioy264.
33. Galaz, J., Motomura, K., Romero, R., Liu, Z., Garcia-Flores, V., Tao, L., Xu, Y., Done,
B., Arenas-Hernandez, M., Kanninen, T., et al. (2023). A key role for NLRP3 signaling
in preterm labor and birth driven by the alarmin S100B. Transl Res *259*, 46-61.
10.1016/j.trsl.2023.04.004.
34. Motomura, K., Romero, R., Garcia-Flores, V., Leng, Y., Xu, Y., Galaz, J., Slutsky, R.,
Levenson, D., and Gomez-Lopez, N. (2020). The alarmin interleukin-1α causes preterm

birth through the NLRP3 inflammasome. Molecular human reproduction *26*, 712-726. 10.1093/molehr/gaaa054.

35. Gomez-Lopez, N., Motomura, K., Miller, D., Garcia-Flores, V., Galaz, J., and Romero, R. (2019). Inflammasomes: Their Role in Normal and Complicated Pregnancies. J Immunol *203*, 2757-2769. 10.4049/jimmunol.1900901.

36. Cohen, P.E., Nishimura, K., Zhu, L., and Pollard, J.W. (1999). Macrophages: important accessory cells for reproductive function. J Leukoc Biol *66*, 765-772. 10.1002/jlb.66.5.765.

37. Repnik, U., Tilburgs, T., Roelen, D.L., van der Mast, B.J., Kanhai, H.H., Scherjon, S., and Claas, F.H. (2008). Comparison of macrophage phenotype between decidua basalis and decidua parietalis by flow cytometry. Placenta *29*, 405-412. 10.1016/j.placenta.2008.02.004.

38. Houser, B.L., Tilburgs, T., Hill, J., Nicotra, M.L., and Strominger, J.L. (2011). Two unique human decidual macrophage populations. J Immunol *186*, 2633-2642. 10.4049/jimmunol.1003153.

39. Svensson, J., Jenmalm, M.C., Matussek, A., Geffers, R., Berg, G., and Ernerudh, J. (2011). Macrophages at the fetal-maternal interface express markers of alternative activation and are induced by M-CSF and IL-10. J Immunol *187*, 3671-3682. 10.4049/jimmunol.1100130.

40. Care, A.S., Diener, K.R., Jasper, M.J., Brown, H.M., Ingman, W.V., and Robertson, S.A. (2013). Macrophages regulate corpus luteum development during embryo implantation in mice. J Clin Invest *123*, 3472-3487. 10.1172/JCI60561.

41. Svensson-Arvelund, J., Mehta, R.B., Lindau, R., Mirrasekhian, E., Rodriguez-Martinez, H., Berg, G., Lash, G.E., Jenmalm, M.C., and Ernerudh, J. (2015). The human fetal placenta promotes tolerance against the semiallogeneic fetus by inducing regulatory T cells and homeostatic M2 macrophages. J Immunol *194*, 1534-1544. 10.4049/jimmunol.1401536.

42. Xu, Y., Romero, R., Miller, D., Kadam, L., Mial, T.N., Plazyo, O., Garcia-Flores, V., Hassan, S.S., Xu, Z., Tarca, A.L., et al. (2016). An M1-like Macrophage Polarization in Decidual Tissue during Spontaneous Preterm Labor That Is Attenuated by Rosiglitazone Treatment. J Immunol *196*, 2476-2491. 10.4049/jimmunol.1502055.

43. Gomez-Lopez, N., Garcia-Flores, V., Chin, P.Y., Groome, H.M., Bijland, M.T., Diener, K.R., Romero, R., and Robertson, S.A. (2021). Macrophages exert homeostatic actions in pregnancy to protect against preterm birth and fetal inflammatory injury. JCI insight *6*. 10.1172/jci.insight.146089.

44. Pique-Regi, R., Romero, R., Tarca, A.L., Sendler, E.D., Xu, Y., Garcia-Flores, V., Leng, Y., Luca, F., Hassan, S.S., and Gomez-Lopez, N. (2019). Single cell transcriptional signatures of the human placenta in term and preterm parturition. Elife *8*. 10.7554/eLife.52004.

45. Pique-Regi, R., Romero, R., Garcia-Flores, V., Peyvandipour, A., Tarca, A.L., Pusod, E., Galaz, J., Miller, D., Bhatti, G., Para, R., et al. (2022). A single-cell atlas of the myometrium in human parturition. JCI insight *7*. 10.1172/jci.insight.153921.

46. Garcia-Flores, V., Romero, R., Tarca, A.L., Peyvandipour, A., Xu, Y., Galaz, J., Miller, D., Chaiworapongsa, T., Chaemsaithong, P., Berry, S.M., et al. (2024). Deciphering maternal-fetal cross-talk in the human placenta during parturition using single-cell RNA sequencing. Science translational medicine *16*, eadh8335. 10.1126/scitranslmed.adh8335.

47. Romero, R., Mazor, M., Munoz, H., Gomez, R., Galasso, M., and Sherer, D.M. (1994). The preterm labor syndrome. Ann N Y Acad Sci *734*, 414-429. 10.1111/j.1749-6632.1994.tb21771.x.

48. Norwitz, E.R., Robinson, J.N., and Challis, J.R. (1999). The control of labor. N Engl J Med *341*, 660-666. 10.1056/NEJM199908263410906.

49. Romero, R., Espinoza, J., Kusanovic, J.P., Gotsch, F., Hassan, S., Erez, O., Chaiworapongsa, T., and Mazor, M. (2006). The preterm parturition syndrome. BJOG *113 Suppl 3*, 17-42. 10.1111/j.1471-0528.2006.01120.x.

50. Smith, R. (2007). Parturition. N Engl J Med *356*, 271-283. 10.1056/NEJMra061360.

51. Gomez-Lopez, N., Galaz, J., Miller, D., Farias-Jofre, M., Liu, Z., Arenas-Hernandez, M., Garcia-Flores, V., Shaffer, Z., Greenberg, J.M., Theis, K.R., and Romero, R. (2022). The immunobiology of preterm labor and birth: intra-amniotic inflammation or breakdown of maternal-fetal homeostasis. Reproduction *164*, R11-R45. 10.1530/REP-22-0046.

52. Willcockson, A.R., Nandu, T., Liu, C.L., Nallasamy, S., Kraus, W.L., and Mahendroo, M. (2018). Transcriptome signature identifies distinct cervical pathways induced in lipopolysaccharide-mediated preterm birth. Biology of reproduction *98*, 408-421. 10.1093/biolre/iox180.

53. Norwitz, E.R., Bonney, E.A., Snegovskikh, V.V., Williams, M.A., Phillippe, M., Park, J.S., and Abrahams, V.M. (2015). Molecular Regulation of Parturition: The Role of the Decidual Clock. Cold Spring Harb Perspect Med *5*. 10.1101/cshperspect.a023143.

54. Motomura, K., Romero, R., Galaz, J., Tao, L., Garcia-Flores, V., Xu, Y., Done, B., Arenas-Hernandez, M., Miller, D., Gutierrez-Contreras, P., et al. (2022). Fetal and maternal NLRP3 signaling is required for preterm labor and birth. JCI insight *7*. 10.1172/jci.insight.158238.

55. Farias-Jofre, M., Romero, R., Galaz, J., Xu, Y., Miller, D., Garcia-Flores, V., Arenas-Hernandez, M., Winters, A.D., Berkowitz, B.A., Podolsky, R.H., et al. (2023). Blockade of IL-6R prevents preterm birth and adverse neonatal outcomes. EBioMedicine *98*, 104865. 10.1016/j.ebiom.2023.104865.

56. Gershater, M., Romero, R., Arenas-Hernandez, M., Galaz, J., Motomura, K., Tao, L., Xu, Y., Miller, D., Pique-Regi, R., Martinez, G., 3rd, et al. (2022). IL-22 Plays a Dual Role in the Amniotic Cavity: Tissue Injury and Host Defense against Microbes in Preterm Labor. J Immunol *208*, 1595-1615. 10.4049/jimmunol.2100439.

57. Furuta, A., Brokaw, A., Manuel, G., Dacanay, M., Marcell, L., Seepersaud, R., Rajagopal, L., and Adams Waldorf, K. (2022). Bacterial and Host Determinants of Group B Streptococcal Infection of the Neonate and Infant. Front Microbiol *13*, 820365. 10.3389/fmicb.2022.820365.

58. Saito, H., Sherwood, E.R., Varma, T.K., and Evers, B.M. (2003). Effects of aging on mortality, hypothermia, and cytokine induction in mice with endotoxemia or sepsis. Mechanisms of ageing and development *124*, 1047-1058. 10.1016/j.mad.2003.08.002.

59. Copeland, S., Warren, H.S., Lowry, S.F., Calvano, S.E., and Remick, D. (2005). Acute inflammatory response to endotoxin in mice and humans. Clin Diagn Lab Immunol *12*, 60-67. 10.1128/cdli.12.1.60-67.2005.

60. Lavin, Y., and Merad, M. (2013). Macrophages: gatekeepers of tissue integrity. Cancer Immunol Res *1*, 201-209. 10.1158/2326-6066.CIR-13-0117.

61. Roszer, T. (2015). Understanding the Mysterious M2 Macrophage through Activation Markers and Effector Mechanisms. Mediators Inflamm *2015*, 816460. 10.1155/2015/816460.

62. Faro, J., Romero, R., Schwenkel, G., Garcia-Flores, V., Arenas-Hernandez, M., Leng, Y., Xu, Y., Miller, D., Hassan, S.S., and Gomez-Lopez, N. (2019). Intra-amniotic inflammation induces preterm birth by activating the NLRP3 inflammasome. Biology of reproduction *100*, 1290-1305. 10.1093/biolre/ioy261.

63. Motomura, K., Romero, R., Plazyo, O., Garcia-Flores, V., Gershater, M., Galaz, J., Miller, D., and Gomez-Lopez, N. (2021). The alarmin S100A12 causes sterile inflammation of the human chorioamniotic membranes as well as preterm birth and neonatal mortality in mice. Biology of reproduction *105*, 1494-1509. 10.1093/biolre/ioab188.

64. Coll, R.C., Robertson, A.A., Chae, J.J., Higgins, S.C., Munoz-Planillo, R., Inserra, M.C., Vetter, I., Dungan, L.S., Monks, B.G., Stutz, A., et al. (2015). A small-molecule inhibitor of the NLRP3 inflammasome for the treatment of inflammatory diseases. Nat Med *21*, 248-255. 10.1038/nm.3806.

65. Galaz, J., Romero, R., Greenberg, J.M., Theis, K.R., Arenas-Hernandez, M., Xu, Y., Farias-Jofre, M., Miller, D., Kanninen, T., Garcia-Flores, V., and Gomez-Lopez, N. (2023). Host-microbiome interactions in distinct subsets of preterm labor and birth. iScience *26*, 108341. 10.1016/j.isci.2023.108341.

66. Gomez de Aguero, M., Ganal-Vonarburg, S.C., Fuhrer, T., Rupp, S., Uchimura, Y., Li, H., Steinert, A., Heikenwalder, M., Hapfelmeier, S., Sauer, U., et al. (2016). The maternal microbiota drives early postnatal innate immune development. Science *351*, 1296-1302. 10.1126/science.aad2571.

67. Lim, A.I., McFadden, T., Link, V.M., Han, S.J., Karlsson, R.M., Stacy, A., Farley, T.K., Lima-Junior, D.S., Harrison, O.J., Desai, J.V., et al. (2021). Prenatal maternal infection promotes tissue-specific immunity and inflammation in offspring. Science *373*. 10.1126/science.abf3002.

68. Mirabella, F., Desiato, G., Mancinelli, S., Fossati, G., Rasile, M., Morini, R., Markicevic, M., Grimm, C., Amegandjin, C., Termanini, A., et al. (2021). Prenatal interleukin 6 elevation increases glutamatergic synapse density and disrupts hippocampal connectivity in offspring. Immunity *54*, 2611-2631 e2618. 10.1016/j.immuni.2021.10.006.

69. Cuna, A., Morowitz, M.J., Ahmed, I., Umar, S., and Sampath, V. (2021). Dynamics of the preterm gut microbiome in health and disease. Am J Physiol Gastrointest Liver Physiol *320*, G411-G419. 10.1152/ajpgi.00399.2020.

70. Tamburini, S., Shen, N., Wu, H.C., and Clemente, J.C. (2016). The microbiome in early life: implications for health outcomes. Nat Med *22*, 713-722. 10.1038/nm.4142.

71. Ficara, M., Pietrella, E., Spada, C., Della Casa Muttini, E., Lucaccioni, L., Iughetti, L., and Berardi, A. (2020). Changes of intestinal microbiota in early life. J Matern Fetal Neonatal Med *33*, 1036-1043. 10.1080/14767058.2018.1506760.

72. Puri, K., Taft, D.H., Ambalavanan, N., Schibler, K.R., Morrow, A.L., and Kallapur, S.G. (2016). Association of Chorioamnionitis with Aberrant Neonatal Gut Colonization and Adverse Clinical Outcomes. PLoS One *11*, e0162734. 10.1371/journal.pone.0162734.

73. Mitchell, B.F., and Taggart, M.J. (2009). Are animal models relevant to key aspects of human parturition? Am J Physiol Regul Integr Comp Physiol *297*, R525-545. 10.1152/ajpregu.00153.2009.

74. Soncin, F., Natale, D., and Parast, M.M. (2015). Signaling pathways in mouse and human
trophoblast differentiation: a comparative review. Cell Mol Life Sci *72*, 1291-1302.
10.1007/s00018-014-1794-x.
75. McCarthy, R., Martin-Fairey, C., Sojka, D.K., Herzog, E.D., Jungheim, E.S., Stout, M.J.,
Fay, J.C., Mahendroo, M., Reese, J., Herington, J.L., et al. (2018). Mouse models of
preterm birth: suggested assessment and reporting guidelines. Biology of reproduction
99, 922-937. 10.1093/biolre/ioy109.
76. Bray, N.L., Pimentel, H., Melsted, P., and Pachter, L. (2016). Near-optimal probabilistic
RNA-seq quantification. Nat Biotechnol *34*, 525-527. 10.1038/nbt.3519.
77. Melsted, P., Booeshaghi, A.S., Liu, L., Gao, F., Lu, L., Min, K.H.J., da Veiga Beltrame,
E., Hjorleifsson, K.E., Gehring, J., and Pachter, L. (2021). Modular, efficient and
constant-memory single-cell RNA-seq preprocessing. Nat Biotechnol *39*, 813-818.
10.1038/s41587-021-00870-2.
78. Dobin, A., Davis, C.A., Schlesinger, F., Drenkow, J., Zaleski, C., Jha, S., Batut, P.,
Chaisson, M., and Gingeras, T.R. (2013). STAR: ultrafast universal RNA-seq aligner.
Bioinformatics *29*, 15-21. 10.1093/bioinformatics/bts635.
79. Heaton, H., Talman, A.M., Knights, A., Imaz, M., Gaffney, D.J., Durbin, R., Hemberg,
M., and Lawniczak, M.K.N. (2020). Souporcell: robust clustering of single-cell RNA-seq
data by genotype without reference genotypes. Nat Methods *17*, 615-620.
10.1038/s41592-020-0820-1.
80. Kang, H.M., Subramaniam, M., Targ, S., Nguyen, M., Maliskova, L., McCarthy, E.,
Wan, E., Wong, S., Byrnes, L., Lanata, C.M., et al. (2018). Multiplexed droplet single-
cell RNA-sequencing using natural genetic variation. Nat Biotechnol *36*, 89-94.
10.1038/nbt.4042.
81. Stuart, T., Butler, A., Hoffman, P., Hafemeister, C., Papalexi, E., Mauck, W.M., 3rd,
Hao, Y., Stoeckius, M., Smibert, P., and Satija, R. (2019). Comprehensive Integration of
Single-Cell Data. Cell *177*, 1888-1902 e1821. 10.1016/j.cell.2019.05.031.
82. Hafemeister, C., and Satija, R. (2019). Normalization and variance stabilization of single-
cell RNA-seq data using regularized negative binomial regression. Genome Biol *20*, 296.
10.1186/s13059-019-1874-1.
83. Korsunsky, I., Millard, N., Fan, J., Slowikowski, K., Zhang, F., Wei, K., Baglaenko, Y.,
Brenner, M., Loh, P.R., and Raychaudhuri, S. (2019). Fast, sensitive and accurate
integration of single-cell data with Harmony. Nat Methods *16*, 1289-1296.
10.1038/s41592-019-0619-0.
84. Yu, G., Wang, L.G., Han, Y., and He, Q.Y. (2012). clusterProfiler: an R package for
comparing biological themes among gene clusters. OMICS *16*, 284-287.
10.1089/omi.2011.0118.
85. Galaz, J., Romero, R., Arenas-Hernandez, M., Panaitescu, B., Garcia-Flores, V., and
Gomez-Lopez, N. (2020). A Protocol for Evaluating Vital Signs and Maternal-Fetal
Parameters Using High-Resolution Ultrasound in Pregnant Mice. STAR Protoc *1*,
100134. 10.1016/j.xpro.2020.100134.
86. Arenas-Hernandez, M., Sanchez-Rodriguez, E.N., Mial, T.N., Robertson, S.A., and
Gomez-Lopez, N. (2015). Isolation of Leukocytes from the Murine Tissues at the
Maternal-Fetal Interface. Journal of visualized experiments : JoVE, e52866.
10.3791/52866.

87. Theis, K.R., Romero, R., Greenberg, J.M., Winters, A.D., Garcia-Flores, V., Motomura,
K., Ahmad, M.M., Galaz, J., Arenas-Hernandez, M., and Gomez-Lopez, N. (2020). No
Consistent Evidence for Microbiota in Murine Placental and Fetal Tissues. mSphere *5*.
10.1128/mSphere.00933-19.
88. Theis, K.R., Romero, R., Winters, A.D., Jobe, A.H., and Gomez-Lopez, N. (2020). Lack
of Evidence for Microbiota in the Placental and Fetal Tissues of Rhesus Macaques.
mSphere *5*. 10.1128/mSphere.00210-20.
89. Winters, A.D., Romero, R., Greenberg, J.M., Galaz, J., Shaffer, Z.D., Garcia-Flores, V.,
Kracht, D.J., Gomez-Lopez, N., and Theis, K.R. (2022). Does the Amniotic Fluid of
Mice Contain a Viable Microbiota? Frontiers in immunology *13*, 820366.
10.3389/fimmu.2022.820366.
90. Kozich, J.J., Westcott, S.L., Baxter, N.T., Highlander, S.K., and Schloss, P.D. (2013).
Development of a dual-index sequencing strategy and curation pipeline for analyzing
amplicon sequence data on the MiSeq Illumina sequencing platform. Appl Environ
Microbiol *79*, 5112-5120. 10.1128/aem.01043-13.
91. Winters, A.D., Romero, R., Gervasi, M.T., Gomez-Lopez, N., Tran, M.R., Garcia-Flores,
V., Pacora, P., Jung, E., Hassan, S.S., Hsu, C.D., and Theis, K.R. (2019). Does the
endometrial cavity have a molecular microbial signature? Sci Rep *9*, 9905.
10.1038/s41598-019-46173-0.
92. Callahan, B.J., McMurdie, P.J., Rosen, M.J., Han, A.W., Johnson, A.J., and Holmes, S.P.
(2016). DADA2: High-resolution sample inference from Illumina amplicon data. Nat
Methods *13*, 581-583. 10.1038/nmeth.3869.
93. Team, R.C. (2013). R: A language and environment for statistical computing.
94. Romero, R., Tarca, A.L., Gomez-Lopez, N., Winters, A.D., Panzer, J., Lin, H., Gudicha,
D.W., Galaz, J., Farias-Jofre, M., Kracht, D.J., et al. (2022). The vaginal microbiota in
early pregnancy identifies a subset of women at risk for early preterm prelabor rupture of
membranes and preterm birth. Research Square. https://doi.org/10.21203/rs.3.rs-
2359402/v1.
95. Davis, N.M., Proctor, D.M., Holmes, S.P., Relman, D.A., and Callahan, B.J. (2018).
Simple statistical identification and removal of contaminant sequences in marker-gene
and metagenomics data. Microbiome *6*, 226. 10.1186/s40168-018-0605-2.
96. Salter, S.J., Cox, M.J., Turek, E.M., Calus, S.T., Cookson, W.O., Moffatt, M.F., Turner,
P., Parkhill, J., Loman, N.J., and Walker, A.W. (2014). Reagent and laboratory
contamination can critically impact sequence-based microbiome analyses. BMC Biol *12*,
87. 10.1186/s12915-014-0087-z.
97. Glassing, A., Dowd, S.E., Galandiuk, S., Davis, B., and Chiodini, R.J. (2016). Inherent
bacterial DNA contamination of extraction and sequencing reagents may affect
interpretation of microbiota in low bacterial biomass samples. Gut pathogens *8*, 24.
10.1186/s13099-016-0103-7.
98. Eisenhofer, R., Minich, J.J., Marotz, C., Cooper, A., Knight, R., and Weyrich, L.S.
(2019). Contamination in Low Microbial Biomass Microbiome Studies: Issues and
Recommendations. Trends in microbiology *27*, 105-117. 10.1016/j.tim.2018.11.003.
99. Karstens, L., Asquith, M., Davin, S., Fair, D., Gregory, W.T., Wolfe, A.J., Braun, J., and
McWeeney, S. (2019). Controlling for Contaminants in Low-Biomass 16S rRNA Gene
Sequencing Experiments. mSystems *4*. 10.1128/mSystems.00290-19.

100.   McMurdie, P.J., and Holmes, S. (2013). phyloseq: an R package for reproducible
1355   interactive analysis and graphics of microbiome census data. PLoS One *8*, e61217.
1356   10.1371/journal.pone.0061217.
1357   101.   Good, I.J. (1953). The population frequencies of species and the estimation of population
1358   parameters. Biometrika *40*, 237-264. 10.1093/biomet/40.3-4.237.
1359   102.   Dixon, P. (2003). VEGAN, a package of R functions for community ecology. Journal of
1360   Vegetation Science *14*, 927-930. https://doi.org/10.1111/j.1654-1103.2003.tb02228.x.

1361

1362

1363

1364

A

Placenta
Extraplacental membranes
Myometrium

Preterm delivery
Term delivery
Homeostatic macrophages

L A B O R ?

Gestational age   37 wks.   40 wks.

B

Term delivery

No labor   In labor

Macrophages:
M1 M5
M2 M6
M3 M7
M4

Preterm delivery
In labor

Myometrium

Placenta

Placenta

Membranes

UMAP_2

-5 0 5 10   -5 0 5 10
UMAP_1

C

Maternal origin
Fetal origin

M2 maternal macrophages

D
Myometrium

Proportion (%)

**

TNL   TIL

E
Maternal side of the placenta (basal plate)

Proportion (%)

1.4-fold

TIL   PTL

G

Cell morphogenesis
Regulation of anatomical structure morphogenesis
Import into cell
Positive regulation of immune system process
Homeostatic process
Response to endogenous stimulus
Regulation of immune system process
Cell development
Anatomical structure morphogenesis
Regulation of developmental process

0.1

0.2

GeneRatio

p.adjust
0.010
0.015
0.020

Count
20
40
60

F

EGFL7   PDK4   FCGBP
FSCN1   TPM1   DAB2
    CD36   MAF
PYGL   MEF2C
    CSF1R   CD209
GPR34   F13A1   SCN9A
SLC18B1   LYVE1
    SESN1   STAB1   WWP1

A *In utero* sterile inflammation

Model of *in utero* sterile inflammation

Alarmin HMGB1

M2 Mφ treatment

Ultrasound-guided intra-amniotic injection

Needle

Fetus

Amniotic cavity

Pregnancy and neonatal outcomes

Week 1

Week 2

Week 3

B Gestational length

Days *post coitum* (dpc)

Preterm birth

PBS	+	−	+	−
HMGB1	−	+	−	+
M2 Mφ	−	−	+	+

C Preterm birth rate

Percentage (%)

0% 25% 22%↓ 3%

PBS	+	−	+	−
HMGB1	−	+	−	+
M2 Mφ	−	−	+	+

D Neonatal mortality at birth

PBS — 100%

HMGB1 — 77%, 20%, 3%

PBS+M2 Mφ — 100%

HMGB1+M2 Mφ — 93%, 3%, 4%

☐ Term ☒ Preterm alive ☐ Preterm dead

E Neonatal survival after birth

Survival (%)

— PBS
— HMGB1
--- PBS + M2 Mφ
--- HMGB1 + M2 Mφ

(weeks)

F Week 1 → Week 2 → Week 3

Neonatal weight (g) *** **

Neonatal weight (g) *** *

Neonatal weight (g) *** *

● PBS ● HMGB1 ● PBS + M2 Mφ ● HMGB1 + M2 Mφ

A Membrane activation

Cervical dilatation — Labor — Uterine contractility

Inflammasome activation — Active CASP-1 — Mature IL-1β

Inflammation

Fetal membrane

HMGB1 (or PBS)

M2 Mϕ (or Vehicle)

Treatment

B
(+) Ctrl.   PBS   HMGB1

Active CASP-1 (20 kDa)
Mature IL-1β (17 kDa)
ACTB (42 kDa)

Dam   1 2 3 4 5 6   1 2 3 4 5 6

D
(+) Ctrl.   HMGB1 + Vehicle   HMGB1 + M2 Mϕ

Active CASP-1 (20 kDa)
Mature IL-1β (17 kDa)
ACTB (42 kDa)

Dam   1 2 3 4 5 6   1 2 3 4 5 6

C
Active CASP-1 — Relative quantification — *
Mature IL-1β — Relative quantification — **
○ PBS   ● HMGB1

E
Active CASP-1 — Relative quantification — **
Mature IL-1β — Relative quantification — **
● HMGB1 + Vehicle   ● HMGB1 + M2 Mϕ

F
PBS   HMGB1

Z-score
Tlr4
Cxcl1
Tnf
Ccl17
Il1b

H
HMGB1+Veh   HMGB1+M2 Mϕ

Z-score
Tlr4
Cxcl1
Tnf
Ccl17
Il1b

G
Ccl17
Expression ($-\Delta C_T$) — **
○ PBS   ○ HMGB1

I
Tlr4 — Expression ($-\Delta C_T$) — *
Cxcl1 — Expression ($-\Delta C_T$) — *
Tnf — Expression ($-\Delta C_T$) — **
Il1b — Expression ($-\Delta C_T$) — **
○ HMGB1 + Vehicle   ○ HMGB1 + M2 Mϕ

A

Control
Bacterial challenge
HMGB1
Group B
Streptococcus

HMGB1 + Vehicle
HMGB1 + M2 Mφ
Neonate (14-16 days old)

B

Neonatal survival

Survival (%)

— Control
— HMGB1
— HMGB1 + Vehicle
— HMGB1 + M2 Mφ

days post-GBS infection

*
**
**

C

Neonatal weight

Δ weight (g)

— Control
— HMGB1
— HMGB1 + Vehicle
— HMGB1 + M2 Mφ

days post-GBS infection

***
**
*