Passaging Human Tauopathy Patient Samples in Cells Generates Heterogeneous Fibrils with a Subpopulation Adopting Disease Folds

Zhikai Zeng[1], Karen Tsay[2], Vishnu Vijayan[2], Matthew P. Frost[3], Shikhar Prakash[4], Athena Quddus[5], Alexa Albert[6], Michael Vigers[7], Madhur Srivastava[8], Amanda L. Woerman[3,9*] and Songi Han[2*]

[1]Department of Chemistry and Biochemistry, University of California Santa Barbara, Santa Barbara, 93105, CA, USA.
[2]Department of Chemistry, Northwestern University, Evanston, 60208, IL, USA.
[3]Department of Biology and Institute for Applied Life Sciences, University of Massachusetts Amherst, Amherst, 01003, MA, USA.
[4]Department of Systems Engineering, Cornell University, Ithaca, 14853, NY, USA.
[5]Department of Molecular, Cellular, and Developmental Biology, University of California Santa Barbara, Santa Barbara, 93105, CA, USA.
[6]Department of Psychological and Brain Sciences, University of California Santa Barbara, Santa Barbara, 93105, CA, USA.
[7]Department of Chemical Engineering, University of California Santa Barbara, Santa Barbara, 93105, CA, USA.
[8]ACERT (National Biomedical Center for Advanced ESR Technology), Department of Chemistry and Chemical Biology, Cornell University, Ithaca, 14853, NY, USA.
[9]Department of Microbiology, Immunology, and Pathology, Colorado State University, Fort Collins, 80521, CO, USA.

*Corresponding author(s). E-mail(s):
amanda.woerman@colostate.edu; songi.han@northwestern.edu;

Contributing authors: zeng@ucsb.edu;
karen.tsay@northwestern.edu; vishnu@umail.ucsb.edu;
mfrost@uchc.edu; sp868@cornell.edu; aquddus@ucsb.edu;
alexacalbert@ucsb.edu; mvigers@ucsb.edu;
madhur.srivastava@cornell.edu;

Abstract

The discovery by cryo-electron microscopy (cryo-EM) that the neuropathological hallmarks of different tauopathies, including Alzheimer's disease, corticobasal degeneration (CBD), and progressive supranuclear palsy (PSP), are caused by unique misfolded conformations of the protein tau is among the most profound developments in neurodegenerative disease research. To capitalize on these discoveries for therapeutic development, one must achieve *in vitro* replication of tau fibrils that adopt the representative tauopathy disease folds, which represents a grand challenge for the field. A widely used approach has been seeded propagation using pathological tau fibrils derived from post-mortem patient samples in biosensor cells that expresses a fragment of the tau protein known as K18, or Tau4RD, containing the microtubule-binding repeat domain of tau as the substrate. The new insights from cryo-EM raised the question of whether the Tau4RD fragment is capable of adopting characteristic tau folds found in CBD and PSP patient fibrils, and whether cell-passaged and amplified tau fibrils can be used as seeds to achieve cell-free assembly of recombinant 4R tau into fibrils without the addition of cofactors. Using Double Electron Electron Resonance (DEER) spectroscopy, we discovered that cell-passaged pathological seeds generate heterogeneous fibrils that are, however, distinct between the CBD and PSP lysate-seeded fibrils, and vastly different from heparin-induced tau fibril structures. Moreover, the lysate-seeded fibrils contain a characteristic sub-population that resembles the disease fold corresponding to the respective starting patient sample. These findings indicate that templated propagation using CBD and PSP patient-derived fibrils is possible with a tau fragment that does not contain the entire pathological fibril core, but also that additional mechanisms must be tuned to converge on a homogeneous fibril population.

Keywords: Tau, seeding, strains, structure, DEER

1 Introduction

Tauopathies are a varied group of neurodegenerative diseases defined by the deposition of fibrillar aggregates of the microtubule-binding protein tau (MAPT, or tau) in the brain. Notably, the prevalence of these deposits is closely related to clinically observable neurodegeneration [1, 2]. In the disease

state, tau misfolds from a soluble monomer into an insoluble fibril with a β-sheet-rich structure capable of recruiting naïve tau to self-replicate, consistent with the prion mechanism of disease. Although tau pathologies specific to each disease vary in morphology and isoform composition [3], the defining difference between the clinical phenotypes is thought to arise from the shape, or conformation, that tau adopts. This is the basis of the strain hypothesis, or the idea that protein conformation is the disease hallmark [4–6].

In 2013, the Goedert lab showed that brain homogenates prepared from a variety of deceased tauopathy patient samples, including progressive supranuclear palsy (PSP), corticobasal degeneration (CBD), and argyrophilic grain disease (AGD), induced tau pathology following intracerebral injection into a transgenic mouse model expressing human tau [7]. The induced inclusions exhibited immunostaining patterns similar to those seen in the corresponding human pathology, suggesting that each disease may be caused by a distinct tau strain [7, 8]. Subsequent work investigating tau strain biology employed cell models expressing the microtubule-binding repeat domain of tau, including the fragment spanning residues 244-378 known as K18, with a C-terminal fluorescent reporter protein [9–11]. These studies reported that tau strains are distinct with respect to seeding capacity, isoform involvement, cell-type specificity, and protease digestion or guanidine denaturation patterns [9–16]. However, it was advances in cryo-electron microscopy (cryo-EM) that facilitated a major leap in our understanding of tau strains on a molecular level, resulting in the resolution of misfolded tau folds present in several tauopathies, including Alzheimer's disease (AD) [17], Pick's disease, [18],CBD [19, 20], chronic traumatic encephalopathy (CTE) [21], PSP [22, 23], AGD [23], and globular glial tauopathy (GGT) [23] patient samples. Goedert, Scheres and coworkers also reported on the tau fibril structure derived from cells infected with human AD tissue-derived seeds to adopt the AD tau fibril structure[24].This important body of work has shown that each tauopathy is associated with tau fibrils that adopt distinctly misfolded tau conformations.

Recombinant tau fibrils induced by polyanionic cofactors, such as heparin, were commonly used to study tau aggregation [25], but cryo-EM [26] and Double Electron Electron Resonance (DEER) [27] have revealed that cofactor-induced fibrils are heterogeneous and distinct from patient-derived fibrils, and are thus not disease relevant. This illustrated a need for developing accessible and reliable methods for replicating disease folds. Cell-free methods are needed for scale-up and facilitate structural biology studies. Previous studies have shown that patient-derived fibrils can propagate in cell-free or *in vitro* systems, as evidenced by the similar morphological and biochemical properties between the tau seeds and resulting fibrils [13, 14, 28–34]. A recent study by Scheres and coworkers further strengthened the concept that templating defines the prion property of pathological tau fibrils by identifying intermediate structures along aggregation pathways[35]. That this debate is evolving and not settled yet is shown in another recent study by Hong and coworkers that replicated AD fibril structures from recombinant tau using AD patient-derived seed that,

however, did not display any seeding competency, implying that templated aggregation may need additional factors than the template alone[36]. In the study presented here, the generated tau fibrils are seeding competent. However, whether or not the atomic conformational features of the disease folds are conserved and replicated by second generation seeding is not known to date.

We set out to test the hypothesis that patient-derived fibrils induce a relevant, tauopathy-specific, protein misfolding of a recombinant tau monomer fragment after passaging through Tau4RD (residues 244-378) reporter cells. While cryo-EM and solid-state NMR have been crucial for unraveling the atomic basis of tau strains at high-resolution, these are not the best tools for readily capturing dynamically evolving and long-range intra-tau distances, nor for studying a heterogeneous fibril population. In order to study *whether* and *how* tau fibrils template protein misfolding in a strain-specific manner at sufficiently high-resolution to differentiate between key disease folds, a structural biology tool is needed that can capture the conformational ensemble of fibrils with both disordered and partially ordered fibril regions and report on the distribution of a heterogeneous fibril population. DEER, a pulsed dipolar electron paramagnetic resonance (EPR) spectroscopic method, is such a tool capable of reporting on the distribution of conformations of doubly spin-labeled proteins regardless of its order and disorder, and hence captures the complete population of tau fibrils formed upon seeded aggregation using CBD and PSP patient-derived materials.

In this study, we isolated tau fibrils from post-mortem CBD and PSP patient samples, which we used to establish monoclonal Tau4RD*LM-YFP cell lines expressing Tau4RD (residues 244-378 with the P301L and V337M mutations), also referred to as biosensor cell lines, that stably propagate either CBD or PSP patient-derived fibrils, as shown in Fig. 1a. Lysates collected from these cells were then added to recombinant 4R tau monomer, either Tau187 (residues 255-441) or the 0N4R tau isoform, to seed protein misfolding. DEER was used to measure the probability distribution, $P(r)$, of intra-molecular distances r, spanning 1.5 to several nm across a pair of spin-labels that are both simultaneously attached to the same tau molecule. We carefully selected three pairs of spin-labels to capture distances across tau segments that can differentiate between CBD and PSP folds - residues 351 & 373, 334 & 360, and 340 & 378 (illustrated in Fig. 1b). A similar approach using select spin label pairs on tau molecules to evaluate the protein fold of tau stacked into fibrils was demonstrated in earlier studies [27, 37, 38]. The theoretical distance distribution for the above-listed spin-label pairs on the reported CBD and PSP folds was computed with the RotamerConvolveMD method [39] (shown in Fig. 1c). The doubly spin-labeled tau was diluted with unlabeled cysteine-less tau to capture intra-molecular distances that are sensitive to the fold that the tau protein adopts when stacked into fibrils, and not inter-molecular distances that report on the fibril packing order. We compared the measured $P(r)$ with the simulated $P(r)$ based on the cryo-EM structures of CBD and PSP tau to

understand whether fibrils seeded with cell-passaged CBD and PSP are conformationally distinct from each other, and from heparin-induced fibrils. We validated this comparison using the novel frequency pattern analysis for DEER data, both qualitatively by visualizing the localized frequency differences, and quantitatively by calculating the structure similarity index measure (SSIM) between differently generated fibrils. Moreover, we set out to discover whether a sub-population of the heterogeneous fibril mixture represents intra-molecular tau distances contained in the CBD and PSP disease folds.

While CBD and PSP patient-derived fibrils are used to induce tau misfolding, the cellular substrate is Tau4RD (spanning residues 244-378), which is shorter than either fibril core and contains no post-translational modifications (PTMs). The cell-derived Tau4RD fibrils were then used as seeds to induce aggregation of 4R tau constructs that contain the entire core of the CBD and PSP fibrils, but again without PTMs, without the use of co-factors in a cell-free system. Hence, it is plausible to think that templating would not be effective and no remnants of the CBD and PSP structural properties would remain in the final tau fibril formed. On the other hand, our recent studies demonstrated that a much smaller 19 residue tau fragment can still form fibrils containing a representative fold found in 4R tauopathies, and it can serve as effective 4R tau isoform-selective seeds for aggregation[40, 41]. Hence, this study set out to determine whether templated misfolding of tau differs between human tauopathy strains when the seeding source is produced via the most widely used cellular biosensor cells expressing Tau4RD as substrate.

2 Results

2.1 Cell-passaged tauopathy patient samples template misfolding of recombinant tau in a cell-free system

To investigate tau misfolding in CBD and PSP, we first established Tau4RD*LM-YFP cells that stably propagate pathogenic tau isolated from brain samples received from deceased CBD and PSP patients (schematic shown in Fig. 1a). Using similar methods as previously described [11], tau prions were isolated from one control, one CBD, and one PSP brain homogenate with sodium phosphotungstate (PTA), which selectively precipitates insoluble protein aggregates [42, 43]. The resulting protein pellets were then resuspended in Dulbecco's phosphate buffered saline (DPBS) and incubated with either Tau3RD*VM-YFP cells or Tau4RD*LM-YFP cells for 4 days. Consistent with our previous findings [42], the CBD and PSP patient samples induced tau misfolding and aggregation in the Tau4RD*LM-YFP cells, but had no effect on the Tau3RD*VM-YFP cell line (Fig. 2a & b). We then generated monoclonal cell lines that stably propagate either CBD or PSP tau via single-cell plating of infected cells. Expression levels of the 4RD-YFP construct were confirmed via Western blot using lysate collected in a radioimmunoprecipitation assay (RIPA) buffer used for cell lysis (Fig. 2c).

For seeding experiments, lysates from Tau4RD*LM-YFP cells infected with CBD and PSP patient samples were collected in 1X protease inhibitor and were further diluted in 20 mM HEPES buffer (pH 7.4) to 5-15% (protein mass percentage) to seed the fibrillization of recombinant Tau187. The two naturally occurring cysteines in this construct were mutated to serines (C291S and C322S) to avoid the formation of disulfide bonds that are not present in the pathological tau fibril folds [19, 21], and instead may inhibit tau aggregation [44]. To determine if cell-passaged lysates can induce the fibrillization of tau in a cell-free aggregation assay, thioflavin T (ThT) fluorescence measurements were collected over the course of 24 hours. ThT is a fluorescent dye that binds specifically to β-sheet structures of amyloid proteins [45] and provides an *in situ* assessment of β-sheet abundance. For similar fibril types that are ThT-active, this assay has been shown to be semi-quantitative. The cell-passaged CBD and PSP lysates induced a robust increase in ThT fluorescence of Tau187 in a concentration-dependent manner (fluorescence intensity increased when 15% lysate was used compared to 5%; Fig. 3a). The negative control, lysate from uninfected Tau4RD cells, did not induce fibrillization (Fig. 3a). Not surprisingly, when polydisperse heparin (average molecular mass 15 kDa, Galen Lab Inc.) was added to the recombinant tau at a tau:heparin molar ratio of 4:1 (equivalent to 17% heparin by mass), heparin induced greater fibril quantities than the CBD or PSP lysates, as assessed by a significantly greater increase in the ThT max fluorescence emission (*** = $P < 0.001$). Furthermore, the cell-passaged lysates were also used to successfully seed the fibrillization of recombinant full-length (FL) 0N4R tau, but less than what heparin induced as determined via ThT fluorescence (Fig. S1). We focused on Tau187 in subsequent experiments as it has a more robust aggregation propensity compared to FL 0N4R tau. Tau187 contains the entire Tau4RD sequence, as well as all regions included in the CBD and PSP fibril cores, while the N terminus that slows aggregation is truncated, and hence is best suited to study the seeded aggregation by cell-passaged CBD or PSP lysates.

To ensure that the measured fluorescence increase in the ThT assay was due to the formation of tau fibrils from recombinant tau monomers rather than from soluble tau species of the cell-passaged seeds, we pelleted the tau aggregates from the lysates by ultracentrifugation at 100,000 x g for 1 hour at 4 °C, removing any soluble small molecule species in the supernatant. The remaining tau fibrils were then resuspended in 20 mM HEPES buffer (pH 7.4) and used to seed recombinant Tau187 fibrillization. ThT fluorescence showed that the pelleted samples induced fibrillization, whereas the supernatant did not (Fig. S2). Moreover, the pelleted samples induced comparable fluorescence plateaus as the cell lysates, indicating that the seeding capacity of the cell-passaged fibrils originates from the aggregated insoluble tau, not from soluble species in the lysates. We conclude that the cell-passaged fibrils induce misfolding and fibriliization of recombinant tau, and that the kinetics of this process is influenced by the quantity of the seed used. To maximize the seeded fibril quantity, 15% cell-passaged seeds were used in subsequent experiments.

The morphologies of the heparin-induced and lysate-seeded fibrils were characterized by negative stain Transmission Electron Microscopy (TEM). As shown in Fig. 3b, heparin-induced fibrils were predominantly long, straight, and well-separated from one another with a few of the fibrils exhibiting a wavy appearance. We also observed long and straight, but more associated, fibrils for both CBD and PSP lysate-seeded fibrils. The CBD lysate-seeded fibrils have a diameter of 20 nm and are devoid of any helical twists. Earlier reports suggest that CBD fibrils are heterogeneous, containing both narrow, straight fibrils and wider twisted, ribbon-like fibrils [46–48]. The morphology we observe in this study is closer to the narrow, straight filaments reported for CBD [19]. The TEM images of PSP-induced fibrils showed fibrils with a very similar appearance, featuring slightly smaller diameter fibrils (16-18 nm) with the appearance of straight tubules, in agreement with the previous reports of brain-derived PSP fibril morphology[47, 48].

To determine if the heparin-induced and lysate-seeded fibrils exhibit prion activity, we incubated both the Tau3RD*VM-YFP and Tau4RD*LM-YFP cells with all three fibril types, as well as monomeric Tau187. Indeed, all three fibrils induced tau aggregation in the Tau4RD*LM-YFP cells but had no effect on the Tau3RD*VM-YFP cells (Fig. 3c & d), consistent with these being 4R fibrils. In contrast, monomeric Tau187 had no effect in either cell line tested.

2.2 Tau fibrils seeded with cell-passaged patient samples have distinct conformations with sub-populations that recapitulate disease-specific folds

Although cryo-EM tau structures [19–23, 23, 49] have rightfully captured the field's attention, cryo-EM requires a homogeneous fibril population from brain samples that can be obtained after extensive purification from large quantities of brain material. Moreover, only large, detergent-insoluble fibrils have been successfully imaged, which may not include critical intermediates and dynamic fibril populations en route to pathological tau aggregation. In contrast, DEER spectroscopy provides an efficient and direct approach to characterize partially ordered and/or heterogeneous fibrils, or even of disordered proteins and partially aggregated proteins. It reports on the probability distribution of intra-molecular distances in the 1.5 to 8 nm ranges across a pair of spin-labels attached simultaneously to a single tau molecule. Therefore, DEER can be used for discovery work on heterogeneous samples or to capture transient conformational ensembles populated along the pathway of seeded aggregation. After conditions for structural convergence are achieved, guided by DEER studies, cryo-EM and other higher-resolution structural biology tools can be used to characterize the end product.

Consistently distinct fibril morphologies observed by negative stain TEM may have different underlying protein folds. However, there are many examples where similar protein folds give rise to distinct fibril morphologies, presumably due to differences in quaternary packing of protofibrils along the fibril axis or inter-fibril interactions, to name a few factors [17, 19, 50]. Some studies have

shown that different protein folds can also give rise to indistinguishable fibril morphologies [20], and vice versa. Hence, protein folds within a fibril cannot be derived from ultrastructural properties, but must be directly measured.

To investigate the conformational features of the lysate-seeded fibrils, we employed DEER to measure the distribution of intra-tau distances. DEER has the unique ability to measure the distribution of distances between a pair of electron spin-labels covalently tethered to tau at two sites modified by site-directed mutagenesis to cysteines. DEER is also solely sensitive to paramagnetic probes such as MTSL [S-(1-oxyl-2,2,5,5-tetramethyl-2,5-dihydro-1H-pyrrol-3-yl)methyl methanesulfonothioate] and is not affected by the disorder or size of the protein, and hence it is particularly well suited for tracking protein aggregation. In contrast, the effects of protein crowding, sample heterogeneity, and protein intrinsic disorder render the use of NMR, crystallography, or cryo-EM challenging or infeasible to investigate a heterogeneous population of tau fibrils. Previously, we presented DEER-derived distance distributions from pairwise spin-labeled tau included in fibrils induced by the addition of the co-factor heparin. The results show that these fibrils are structurally heterogeneous judging by the width of the intra-tau distance distribution, $P(r)$. These structures were found to be clearly distinct from the tau fibrils present in AD patient samples when comparing the theoretical distance distribution for spin-label pairs on the reported AD fold with the experimentally measured one [27]. This result was subsequently confirmed with cryo-EM structures of heparin-induced tau fibrils that revealed a complete lack of homology between the heparin-induced and patient-derived fibrils [26].

In this study, we followed a similar procedure as reported in Fichou *et al.* [27] by rationally selecting the spin-label pairs to be positioned along the outer layer of the fibril cores of the CBD and PSP folds to differentiate between the two cryo-EM structures [19, 21]. Critically, we selected spin-label pairs that yield significantly different $P(r)$ between the monomeric and aggregated state (Fig. S4) to ensure that the change of the mean distance and distance distribution can unambiguously report on tau misfolding. All labeled sites were chosen due to their location outside of the heparin-induced fibrils core as identified by cryo-EM [26] and hence should differentiate between heparin-induced and patient-derived structures. The pairs of sites that fulfill the criteria were residues 351 & 373, 334 & 360, and 340 & 378. After selecting our spin-label pairs, we introduced two cysteines at a time into Tau187 at these desired sites by site-directed mutagenesis and spin-labeling (SDSL) of MTSL to the cysteine residues [51, 52]. The expected pairwise distance distributions, taking the rotamer distribution into account, are shown as grey-filled features in Fig. 4. None of the distance pairs are included in the core of the heparin fibril structure solved to date, while each of them are distinct between CBD and PSP.

To ensure that DEER measurements capture intra-molecular (not intermolecular) distances, we diluted the doubly spin-labeled Tau187 with unlabeled Tau187 (i.e., without mutation to cysteines, and hence also no labels)

to get spin-diluted Tau187 (10% doubly spin-labeled Tau187 + 90% unlabeled Tau187), such that statistically the closest distances are from the pair of spin-labels attached to the same Tau187 molecule. Effects from longer inter-molecular distances between spin-labels are corrected by dividing a background function from the time-domain DEER decay that is modulated by any residual inter-molecular spin-spin interactions. Seeding assays conducted using spin-diluted Tau187 yielded similar ThT fluorescence intensity profiles as the aggregation assay using unlabeled Tau187 alone (Fig. S2). To prepare samples for DEER measurements, we added heparin or 15% lysates, respectively, to induce aggregation for 24 h. Samples were then subjected to dialysis (molecular weight cut off 25 kDa) against a D_2O buffer with 20 mM HEPES to remove excess monomer and to reduce the ^1H concentration of water to aid DEER measurements. We then derived the distribution of intra-tau spin-label distance, P(r), from the experimental time-domain DEER data, V(t), which is modulated by convolved dipolar coupling frequencies between pairs of spin-labels. The reconstruction of the distance distribution, P(r), from V(t) was determined using the DeerLab fitting algorithm and software [53]. The 95% confidence intervals were determined using the bootstrapping method with a sample size of 100, and shown in Fig. 4. Triplicates of each of the DEER time-domain and P(r) data is shown for the monomer and three types of fibrils, and for each of the three spin label pairs in Fig. S4, S5 and S6. Visually, both the reproducibility between replicates and the distinction between different fibril types and distances is remarkable.

When comparing the DEER-derived P(r) in the monomer and heparin-induced fibril state across sites 351-373, 334-360, and 340-378 (shown in Fig. S4), it is clear that the protein region spanning these distances in the monomer is not simply intrinsically disordered, given that their P(r) contains distinct features and is narrower than the P(r) of heparin-induced fibrils that exhibit a single, broad, peak. This is particularly clear for the distances spanning 334-360 and 340-378. Sites 334, 351, 360, 373, and 378 have not been reported to be part of any heparin-induced fibril fold according to cryo-EM [26], further highlighting the ability of DEER to report on less homogeneous and more dynamic protein folds in monomers and partially structured fibrils.

Before presenting a more in-depth, unbiased, analysis of similarity and difference of the DEER data, we first examine the pair-wise distances present in the experimental P(r) in the context of distances expected from the reported cryo-EM CBD and PSP folds by comparing their V(t) and P(r) (Fig. 4). The heparin-induced fibrils (green) yielded the broadest distance distribution across all distance pairs, and distinct from the CBD (pink) and PSP (blue) lysate-seeded fibrils. The P(r) of CBD and PSP contain the expected distances within their profile, however it is difficult to tell apart by eye whether they are distinct from each other.

We hence needed a method to assess the DEER data in an unbiased fashion. For this, we utilized a new data analysis methodology developed by Srivastava, Freed and coworkers, which uses a frequency pattern recognition technique that

combines discretized continuous wavelet transform (CWT) of pulsed dipolar spectroscopy data and structure similarity index measure (SSIM), a technique commonly used in image analysis [54**?**]. This method relies on the ability of CWT to provide time-frequency representation of the data, and therefore allows for the analysis of localized variation in the signal. By calculating the CWT, we can distinguish between the DEER signal, noise, and artifacts directly from the experimental data by decomposing the DEER signal into different frequency components, thereby allowing for a more detailed comparison of the raw time-domain data V(t), while avoiding the interfering effects of background correction and other pre-processing error that can occur in extracting the P(r) shape. This approach renders this analysis less bias-prone [54]. By utilizing the SSIM analysis, we can capture small differences between the spectrograms of the CWT of two data sets by providing a quantitative measure of similarity, with index < 1 implying there is a difference between the spectrograms [55]. Additionally, the similarity gradient plot, which visually represents the difference between two spectrograms, allows for a qualitative comparison between spectrograms by revealing the location of the differences in the frequency profile. This method is not yet able to correlate the difference in frequency profile with differences in P(r), but the SSIM index can evaluate whether the DEER data are differentiable or not. The result of this frequency pattern recognition is displayed as a similarity gradient plot, along with the SSIM index, to compare the spectrograms of heparin-induced and CBD and PSP lysate-seeded fibrils (Fig. 5). In the Heparin vs CBD and CBD vs PSP comparison in Fig. 5a, the SSIM analysis shows that the Heparin data is significantly different from both CBD and PSP data (SSIM value between 0.5 and 0.7 for all spin label pairs); whereas, the PSP is more similar to the CBD data (SSIM value above 0.9). However, there are still notable differences in the high frequency region between the CBD and PSP data for all three spin label pairs. This is reflected in distinct color variations in the similarity gradient plot, consistently visible in the higher frequency range (top of the plot) for all three spin label pairs. For DEER data across sites 340-378, there are also distinct differences in the medium to low frequency range (center of the plot,) as reflected in two distinct green streaks across the time domain. We conclude that the heparin-induced fibrils are structurally unique and more heterogeneous in the microtubule binding region of the protein judging by the width of P(r). The CBD and PSP lysate-seeded fibrils (in pink and blue, respectively) exhibited distinct patterns from that of heparin-induced fibrils. This is especially notable in the P(r, 351-373) data that features two dominant distance features with narrower widths compared to that of heparin-induced fibrils, suggesting that multiple distinct populations of ordered tau conformers are present.

Given the complexity of the observed P(r) shape, it is non-obvious whether the patterns of the peaks representing the CBD vs PSP lysate-seeded fibrils are reproducible until triplicates for the time-domain and P(r) data (all data shown in Fig. S5 - 7) and their CWT (all data shown in Fig. S8 - 14, Table S1) are compared to each other. The triplicate data show stunning reproducibility

between the repeats. The triplicates were derived from lysate-seeded samples from at least three different *in vitro* seeding experiments performed on different batches of recombinant Tau187 using preparations of different cell-passaged seeds. The cell-passaged seeds were derived from multiple cellular batches prepared at different times spanning multiple years. However, the cells were infected with the same biological CBD or PSP patient-derived insoluble tau material. These results reveal that the cell-passaged seeds derived from CBD or PSP human tissues, even after multiple passaging through biosensor cells that express tau fragments that do not contain the full core-forming sequence, still exert reproducible misfolding and templating effects on tau aggregation in the cell and, upon harvesting, in vitro.

An important observation is that each pattern of P(r) is still broad, suggesting that the templating induced by seeding does not generate converged fibril structures with the commonly used seed material and widely used seeding protocol. Critically, a dominant peak near the predicted spin-label distance (indicated by dotted lines) suggests that a sub-population of fibrils resembling the CBD and PSP disease folds may have formed. Specifically, a mean distance spanning sites 351-373 of 6.5 nm is expected for the CBD fold, while a dominant experimental peak is found at distances of 5-7 nm in P(r, 351-373). In contrast, a mean distance of 3 nm is expected in P(r, 351-373) for the PSP fold, and again, one of the dominant experimental peaks is found spanning distances of 2-4 nm (Fig. 4a). Notably, this intensity at 2-4 nm is missing in P(r, 351-373) of the CBD lysate-seeded tau fibril population.

The experimental P(r, 334-360) of CBD lysate-seeded fibrils (pink solid line) contains the expected distance distribution (pink dotted line) centered around 2.5 nm. The P(r, 334-360) of PSP lysate-seeded fibrils has a dominant peak centered around 5-7 nm, which is closer to the expected distance of 7.5 nm. In contrast, the P(r, 340-378) of CBD or PSP lysate-seeded fibrils contain a major peak around 7-10 nm. This is slightly greater than the expected mean distance of 7.5 nm for the CBD fold, and much greater than the expected 5.5 nm for the PSP fold, but certainly closer to the expected distances compared to that from P(r, 351-373) or P(r, 334-360). To validate the significance of the just discussed differences in P(r), we examined the similarity gradient plot between the CBD vs PSP lysated-seeded fibrils (Fig. 5). They show differences in the higher frequency region (i.e., short distance region) for all three distance pairs, and also in the medium frequency range (i.e., longer distance region) for the 340-378 pair. This is consistent with the P(r, 340-378) featuring longer distances, spanning 5.5-8 nm, for CBD or PSP lysated-seeded fibrils.

Notably, site 378 lies at the most terminal position of the structure, with respect to the expected CBD or PSP fibril core [19, 21], compared to all other spin-labeled sites tested here. The Tau4RD expressed in the Tau4RD*LM-YFP cells also ends at residue 378, and the subsequent C-terminal residues present in both the CBD and PSP folds may be key to stabilizing the structures. It is, therefore plausible that the terminal segment around site 378 is not tightly

folded into the CBD or PSP cell-passaged seeds used here, resulting in additional broadening or lengthening of the distance captured in the experimental P(r) shape for the induced fibril. The interpretation of P(r, 340-378) data for very broad features spanning distances exceeding 7 nm as found with CBD and PSP lysate-seeded fibrils must be viewed with caution according to benchmarks set for DEER [56]. However, the remarkable reproducibility of P(r) patterns between triplicates (Fig. S7, S11-S14; Table S1) demonstrates that meaningful structural features are contained in the dipolar coupling distributions between spin-labels at sites 340-378, even if their direct interpretation is tenuous.

Each P(r) of CBD or PSP lysate-seeded fibrils contain populations that feature greater distances between sites 351-373, 334-360, and 340-378, compared to heparin-induced fibrils, demonstrating that these sites are located at the outer core of a sub-population of tau fibrils and adopt more extended conformations, as expected from the PDB structure of the CBD and PSP folds. In other words, the *in vitro* CBD and PSP lysate-seeded fibrils contain sub-populations of fibrils that adopt structural features that are consistent with the tau structures found by cryo-EM, and certainly distinct from heparin-induced fibrils or amorphously tau aggregates. These findings indicate that cell-passaged CBD and PSP can transmit their structural properties to naïve tau by templated misfolding, at least in part, even though the biosensor cell lines used in this study express Tau4RD spanning residues 244-378, which is a few amino acids short of the tau segment that forms the PSP and CBD core. The results also show that seeding with cell-passaged patient-derived tau populates multiple fibril structures. It is conceivable that the particle picking and class averaging approach of cryo-EM result in resolving only the dominant fibril sub-population, failing to capture other conformations that may exist in the human brain. Future cryo-EM studies will benefit from experimental approaches to measure the complete ensemble of the fibril population that guide the discovery of mechanisms and conditions to achieve structural convergence to generate pathological tau fibrils.

3 Discussion

In this study, we successfully used cell-passaged CBD and PSP patient samples to induce recombinant tau fibrillization without the addition of any co-factors or the use of sonication. The resulting fibrils were able to reinfect the Tau4RD*LM-YFP cells showing isoform specificity (Fig. 3c & d). Using DEER-based distance measurements at sufficiently high-resolution and subnanometer levels, we found that the CBD and PSP lysate-seeded fibrils are structurally distinct from one another, as well as from heparin-induced tau fibrils (Fig. 4). Together, these results indicate that we can propagate and measure distinct fibril structures, but that additional factors must be discovered and controlled to achieve structural convergence. Our study shows that DEER is an effective adjunctive tool to cryo-EM to evaluate strain-specific tau misfolding for any experimental condition, independent of whether convergence of fibril structure is achieved or not.

Our results are consistent with the predominant hypothesis that tau fibril propagation uses the prion mechanism to self-template distinct conformations or strains. Notably, this is the first study that examined directly the structural property of the cellular lysated-seeded fibril ensemble. The outcome was non-obvious, and could have been that all structural memory of the patient sample is lost after multiple passaging through the cell and subsequent in vitro seeding, or that it prevails much more strongly to generate a dominant tau fibril population adopting CBD or PSP folds. Instead, our study demonstrated that templating occurs and is propagated even under imperfect conditions, but that a highly heterogeneous fibril population is generated. While cryo-EM has revealed a diverse array of tau fibril conformations [19–23, 23, 49], the ability to replicate these structures upon seeding has only recently been completed at high-resolution. An exciting study was published recently by Tarutani *et al.* reporting the first cryo-EM structures of cell-passaged insoluble tau fibrils [24]. Complementing this exciting work, we demonstrated with the help of DEER that can uniquely capture an ensemble of structures within a larger population that multiple fibril structures are present, not just a homogeneous or dominant sub-population.

Previously, morphological and biochemical properties were used to demonstrate that distinct tau conformations exist [13, 14, 28, 28, 29, 34], and similar features between the seeded fibrils and the starting seed, or tau prion, have been used to determine the fidelity of tau templating [13, 14, 28–34, 38]. For example, Kaufman *et al.* [10] isolated and characterized 18 tau strains that originated from either recombinant protein, tauopathy mouse brain, or deceased human tauopathy patient samples. The isolated strains were categorized using multiple approaches, including inclusion morphology, detergent solubility, seeding activity, proteolytic digestion, and toxicity. After inoculating transgenic mouse brains with the 18 strains, the unique patterns of neuropathological lesions induced by each strain were faithfully transmitted, indicating that the seeded fibrils in mice are similar to the starting material in the inocula. In a more recent study, Xu *et al.* [34] showed that AD patient samples may be used to template recombinant tau misfolding, resulting in fibrils that adopt the same conformation. While their analyses lacked confirmation via cryo-EM, the conformation-dependent antibodies DMR7 and MC1 were shown to bind to the AD-induced fibrils. In contrast, others have found that the AD seeded fibrils adopt conformations that are distinct from the starting material, as determined by unique protease digestion patterns and macroscopic morphology of induced fibrils [13, 57]. The current state of the literature shows that robust structural biology tools are needed to effectively guide this evolving debate regarding strain-specific templated misfolding of tau. Studies such as ours will contribute to the critical ongoing debate on the molecular basis of tau prions and shape-selected fibril formation.

Using DEER and frequency pattern recognition, we detected partial distance features of seeded tau fibrils that are similar to the expected distances measured from the cryo-EM structures of CBD and PSP tau. These findings

indicate that we achieved some convergence with these structures using seeding, even when using imperfect cellular seed material. One might view these results only as partial success given that the fibril structures did not converge to a single structure resembling the CBD or PSP fold. While achieving complete convergence to disease fold is an ultimate goal, the data presented here highlights an optimistic picture, namely that templated aggregation is a potent mechanism to generate tau "strains", as demonstrated when using heterogeneous patient-derived seeds, imperfect cellular tau substrates and models, and imperfect in vitro tau substrates. No other experimental technique could reveal that tauopathy seeds derived from cell lysates do, on the one hand, generate the originally targeted disease fold, while on the other hand also report that many other fibril structures coexist with the disease fold. The ability to capture the entire conformational population is powerful because this tool offers the opportunity to learn what dominant and relevant factors are needed to achieve conformation selection and convergence.

Importantly, there are a few factors that may account for this partial convergence, rather than complete replication of fibril structure. First, the cell-passaged seeds were generated in cells expressing Tau4RD which lacks the N- and C-termini. Although the Tau4RD in cells may be templated by human materials, residues outside of the microtubule binding region (such as the rest of C-terminal region after residue 378) may contribute to tau folding, strain propagation, and/or structural stabilization. As a result, the CBD and PSP folds may only partially form in cells. This is potentially reflected in the indistinguishable $P(r, 340\text{-}378)$ data measured from the CBD and PSP lysate-seeded fibrils. Moreover, despite the use of an 18-amino acid flexible linker, the fusion of Tau4RD to a fluorescent protein may induce steric hindrance that inhibits or alters tau aggregation [58]. However, the linker sequence connecting tau and GFP in the studies by Kaniyappan *et al.* (GAPGSAGSAAGSG) is not the same linker sequence used in the cell model used here (EFCSR-RYRGPGIHRSPTA). Others have shown that the presence of a GFP tag on α-synuclein does not interfere with *in vitro* fibrillization [59]. It remains an open question whether the cell-passaged seeds are disrupted by the YFP tag and the protein truncation. The Tau4RD fragment is still important because it is so far the most commonly used construct in cellular seeding assays [11, 25, 60–70]. We, therefore, need to know how to interpret the wealth of critical data previously reported, no matter what the ultimate verdict is (e.g., whether the 4RD seeds do or do not faithfully template disease folds). Furthermore, truncated tau remains biologically relevant because fragments of tau have been identified in AD patient brain samples [69], and in patient cerebrospinal fluid [71–73].

The cell-free seeding conditions used in this study may not favor the propagation of specific disease-relevant strains, resulting in the formation of multiple conformers. For example, no disease-relevant PTMs are present on the recombinant tau used here. Additionally, the exact makeup of the solution conditions, including molecular chaperones, crowding agents, co-factors, and even salt type

and concentration likely play major roles in dictating the tau folding pathway. In fact, a recent study using cryo-EM showed that simply changing salt types and salt concentration in the seeding buffer dramatically alters the shape of the resulting tau fibrils [50]. By systematically and empirically changing the buffer conditions, the authors successfully generated recombinant tau fibrils identical to the conformations reported for AD and CTE tau. Therefore, further optimization of the seeding conditions is needed to study the underlying mechanism(s) for shape selection in tau seeding.

We need to consider two important factors when interpreting cryo-EM structures of tau fibrils. The cryo-EM structures are generated using samples from deceased patients and therefore represent the end stage of the disease. It is possible that the process of tau misfolding in a living patient is more dynamic. Hence, the heterogeneous nature of the tau fibril population reflected in our dataset captured from fibril samples 24 h after seeding may have biological relevance for capturing earlier stages of tau misfolding. Additionally, the class averaging required to resolve these structures likely selects for the most abundant conformations among a population of fibrils. As has been shown for the prion protein, there may be a mixture, or cloud, of conformations present in patients with CBD and PSP. If this is the case, it is less likely that cryo-EM studies will resolve the less abundant tau species, which may be detectable by DEER, demonstrating the need for complementary structural biology tools.

Finally, an important limitation to our current work is the use of one CBD and one PSP patient sample to generate Tau4RD*LM-YFP cells that stably propagate tau aggregates. While this approach was taken to support the initial establishment of the methodology and approach reported here, it is possible that by using a single patient sample, we are not fully capturing the diversity in P(r) that exists across misfolded tau for the two diseases. Future studies will focus on understanding the patient-to-patient variability in tau misfolding that may exist, which will be done using cell lines that avert some of the caveats discussed above.

4 Conclusion

We demonstrate that the recombinant tau fibrils induced by CBD and PSP cell-passaged seeds are structurally distinct from heparin-induced fibrils and that a sub-population of the lysate-seeded fibrils resemble the CBD and PSP fibrils isolated from human patient samples, even when using cellular and in vitro substrates without any PTMs. These findings validate the application of DEER to investigate the self-templating process during tau propagation at high-resolution. Moreover, these data will also be important for guiding the development of strain-specific positron emission tomography (PET) ligands [74–76].

5 Methods

5.1 Human patient samples

The CBD patient sample used was provided by the NIH NeuroBioBank. This sample was from a 65-year-old male patient. The PSP patient sample used was provided by the Massachusetts Alzheimer's Disease Research Center. This sample was from a 70-year-old female patient. Fresh-frozen human tissue was used to create a 10% (wt/vol) homogenate using calcium- and magnesium-free 1X Dulbecco's phosphate-buffered saline (DPBS) using an Omni Tissue Homogenizer (Omni International).

5.2 Cell line development

The human tau DNA sequence encoding residues 244-378 with the P301L and V337M mutations (based on the longest tau isoform, 2N4R) fused to enhanced yellow fluorescent protein by an 18 amino acid flexible linker (EFC-SRRYRGPGIHRSPTA) was synthesized and cloned into the pcDNA3.1(+) expression vector by GenScript. The Tau4RD*LM-YFP sequence was then subcloned into the pIRESpuro3 vector (Takara) using EcoRV (5') and NotI (3'). The same protocol was used to construct a plasmid containing the repeat domain of 3R tau (residues 244-274,306-378) containing the L226V and V337M mutations. Gene sequence and insertion were confirmed by Sanger sequencing before subsequent use.

HEK293T cells (ATCC) were cultured in Dulbecco's modified Eagle's medium (DMEM; Corning) supplemented with 10% fetal bovine serum (FBS), 100 μg/mL penicillin, and 100 μg/mL streptomycin (ThermoFisher), referred to hereafter as complete media. Cultures were maintained in a humidified atmosphere of 5% CO2 at 37 °C. Cells were plated at a density of 5.7 X 10^5 cells per well in a 6-well plate overnight in complete media before adding 1 μg of plasmid DNA incubated with 3.5 μL Lipofectamine 2000 (ThermoFisher) for 20 min. Stable cells were selected in complete media containing 1 μg/mL puromyocin (ThermoFisher) for 48 h before generating monoclonal lines by limiting dilution of polyclonal cells in 384-well plates. The resulting monoclonal lines were frozen in liquid nitrogen. Lysates from the lines were collected in 1X radioimmunoprecipitation assay (RIPA) buffer containing 50 mM Tris-HCl, pH 7.5 (ThermoFisher), 150 mM NaCl (Sigma), 5 mM EDTA (ThermoFisher), 1% nonidet P-40 (ThermoFisher), 0.5% deoxycholate (ThermoFisher), and 0.1% sodium dodecyl sulfate (SDS; ThermoFisher). Cell lysates in RIPA buffer were frozen, thawed, and clarified using two low-speed spins (500 x g for 5 min followed by 1,000 x g for 5 min). Total protein was measured in the supernatants via bicinchoninic acid (BCA) assay (Pierce). To compare the expression of tau-YFP across the clones, a total of 10 μg total protein was prepared in 1X NuPAGE LDS loading buffer and boiled for 10 min. Samples were loaded onto a 10% Bis-Tris gel (ThermoFisher) and SDS-PAGE was performed using the MES buffer system. Protein was transferred

to a polyvinylidene fluoride (PVDF) membrane. The membrane was blocked in blocking buffer [5% (wt/vol) nonfat milk in 1X Tris-buffered saline containing 0.05% (vol/vol) Tween 20 (TBST)] for 30 min at room temperature. Blots were incubated with primary antibody for GFP (1:10,000; Abcam) in block buffer overnight at 4 °C. Membranes were washed three times with 1X TBST before incubating with horseradish peroxidase-conjugated goat antib-rabbit secondary antibody (1:10,000; Abcam) diluted in blocking buffer for 1 h at 4 °C. After washing blots three times in 1X TBST, membranes were developed using the enhanced chemiluminescent detection system (Pierce) for exposure to X-ray film. This same protocol was used to evaluate tau-YFP expression in uninfected vs infected cells.

To generate cells that stably propagate tau prions, fresh frozen human brain tissue samples from one CBD and one PSP patient sample were used. Tau prions were isolated from the samples using sodium phosphotungstic acid (NaPTA; Sigma) as described [11]. Isolated protein pellets were stored at 4 °C. Tau4RD*LM-YFP cells were infected with tau prions isolated from the CBD or PSP patient samples as described below. Monoclonal subclones stably propagating tau prions were generated by isolated Tau4RD*LM-YFP cells by limiting dilution of a polyclonal cell population in 384-well plates. Clones were selected by the presence of YFP-positive aggregates in >95% of cells in the culture. Resulting monoclonal lines were frozen in liquid nitrogen.

5.3 Cellular tau prion bioassay

Tau4RD*LM-YFP cells were plated in 384-well plates with black polystyrene walls (Greiner) with 0.012 μg Hoechst 33342 (ThermoFisher) at a density of 4,000 cells/well. Tau3RD*VM-YFP cells were plated using the same conditions at a density of 2,750 cells/well. Cells were incubated at 37 °C for 2 to 4 h to allow adherence to the plate. To infect Tau4RD*LM-YFP cells, PTA-precipitated patient samples diluted 1:5 in 1X DPBS or 0.05 μM fibrils in 1X DPBS were incubated with Lipofectamine 2000 2% final volume; ThermoFisher) for 1.5 h at room temperature. OptiMEM (78% final volume; ThermoFisher) was added before each sample was plated in six replicate wells. To infect Tau3RD*LM-YFP cells, PTA-precipitated patient samples diluted 1:20 in 1X DPBS or 0.05 μM fibrils in 1X DPBS were incubated with Lipofectamine 2000 1% final volume; ThermoFisher) for 1.5 h at room temperature. OptiMEM (79% final volume; ThermoFisher) was added before each sample was plated in six replicate wells. Plates were then incubated at 37 °C in a humidified atmosphere of 5% (vol/vol) CO_2 for 4 days before imaging on the Molecular Devices XLS.Images of both the DAPI and YFP channels were collected from five different regions in each well. The images were analyzed using an algorithm built in Custom Module Editor (Molecular Devices) to identify intracellular aggregates only in living cells. Data reported as fluorescence/cell X 10^6 arbitrary units (A.U.).

5.4 Tau187 and 0N4R expression and purification

N-terminal truncated, microtubule-binding domain containing Tau187 (residues 255-441 with a His-tag at the N-terminus) and 0N4R were used for *in vitro* studies. Mutated variants of Tau187 were prepared using site-direct mutagenesis: cysteine-less (cysless) Tau187 P301L/Q351C/T373C contains C291S, C322S, Q351C and T373C; cysless Tau187 P301L/G334C/I360C contains C291S, C322S, G334C and I360C; cysless Tau187 P301L/K340/F378C contains C291S, C322S, K340 and F378C. Cysless Tau187 P301L and cysless 0N4R P301L constructs were made with C291S and C322S mutations.

The cloning, expression, and purification of Tau187 have been previously described [77]. Shortly, the gene encoding desired Tau187 tau or 0N4R tau was cloned into a pET-28a vector and was tranfected into E. coli. BL21 (DE3) competent cells (Novagen). E. coli BL21 (DE3) cells were cultured from frozen glycerol stock or from plates overnight in 10 mL luria broth (LB) which was used to inoculate 1 L of fresh LB. Culturing and inoculation were performed at 37 °C with shaking of 200 rpm. At OD 600 of 0.6-0.8, Tau187 variant expression was induced by incubation with 1 mM isopropylß-D-thiogalactoside (Sigma Aldrich) for 2-3 h. Cells were harvested by centrifugation for 10 min at 5000 × g (Beckman J-10; Beckman Instruments,Inc.), and the pellets were stored at -20 °C until further use. Cell pellets were resuspended in lysis buffer (Tris-HCl pH 7.4, 100 mM NaCl, 0.5 mM DTT, 0.1 mM EDTA, 1mM PMSF) with 1 Pierce protease inhibitor tablet (Thermo Fisher). Lysis was initiated by the addition of lysozyme (2 mg/ml), DNase (20 μg/ml), and $MgCl_2$ (10 mM) and incubated for 30 min on ice. Lysate was then heated to 65 °C for 13 min, cooled on ice for 20 min and then centrifuged to remove the precipitant. The supernatant was loaded onto a Ni-NTA agarose column pre-equilibrated with wash buffer A (20 mM sodium phosphate pH 7.0, 500 mM NaCl, 10 mM imidazole, 100 μM EDTA). The column was then washed with 20 ml of buffer A, 15 ml buffer B (20 mM sodium phosphate pH 7.0, 1 M NaCl, 20 mM imidazole, 0.5 mM DTT, 100 μM EDTA). Purified Tau187 was eluted with buffer C (20 mM sodium phosphate pH 7.0, 0.5 mM DTT, 100 mM NaCl) supplemented with varying amounts of imidazole increasing from 100 mM to 300 mM. The protein was then concentrated via Amicon Ultra-15 centrifugal filters (MWCO 10 kDa; Millipore Sigma) and the buffer was exchanged into final buffer (20 mM ammonium acetate buffer at pH 7.0, or 20mM HEPES at pH 7.4) by PD-10 desalting column (GE Healthcare). To further purify the protein, size exclusion chromatography was conducted by injecting 2-5 mL sample onto HiLoad 16/600 Superdex 200 Column (GE Healthcare Life Sciences) connected to a BioRad NGC Quest 10 FPLC system, and 1.5 mL fractions were collected with a BioFrac fraction collector (BioRad). Prior to sample injection, the column was washed by 130 mL degassed Milli-Q water at a flow rate of 0.8 mL/min and equilibrated with 160 mL degassed working buffer (20 mM ammonium acetate buffer at pH 7.0, or 20mM HEPES at pH 7.4) at 0.8 mL/min. Samples were eluted from the column with 150 mL working buffer at 0.6 mL/min after

sample injection. Fractions of each elution peaks were collected right after elution. Elution peak assignment was done by comparing with the elution profile of purified Tau187 monomers. Fractions corresponding to monomer were concentrated using Amicon Ultra-4 centrifugal filters (MWCO 10 kDa; Millipore Sigma). The final protein concentration of Tau187 was determined by UV-Vis absorption at 274 nm using an extinction coefficient of 2.8 $cm^{-1}mM^{-1}$ calculated from absorption of Tyrosine. The final protein concentration of 0N4R was determined by UV-Vis absorption at 274 nm using an extinction coefficient of 7.4 $cm^{-1}mM^{-1}$ calculated from absorption of Tyrosine.

5.5 Protein spin-labeling

Protein was spin-labeled using MTSL ((1-Acetoxy-2,2,5,5-tetramethyl-δ-3-pyrroline-3-methyl) Methanethiosulfonate) purchased from Toronto Research Chemicals. Prior to labeling, samples were treated with 5 mM DTT, which was removed using a PD-10 desalting column. Then, $10\times$ to $15\times$ molar excess MTSL to free cysteine was incubated with the protein at 4 °C overnight. Excess MTSL was removed using a PD-10 desalting column. Labeling efficiency, defined as the molar ratio of tethered spin-labels over the cysteines, was measured to be 50-60% for double-cysteine mutants.

5.6 Heparin-induced tau aggregation

Cell-free tau aggregation was carried out in the working buffer (20mM HEPES at pH 7.4) at 37 °C. 20 μM ThT dye was added into 50 μM Tau187 in Corning™ 384-Well Solid Black Polystyrene Microplates (Thermo Fisher Scientific). Then, heparin was mixed in the samples at the mole ratio of 4:1 (Tau187:heparin) to induce aggregation. ThT fluorescence was monitored by Bio-Tek Synergy 2 microplate reader (excitation 440/30, emission 485/20, number of flash 16, gain 50) over a period of 24h at 37 °C. One data point was taken every 3 minutes. Measurements were done in triplicate. The figures showed the average and standard deviations.

5.7 Cell-free seeding assay

5% to 15% (protein mass percentage) of CBD and PSP cell-passaged seeds were incubated with 50 μM cysless P301 Tau187 in 20 mM HEPES buffer to make seeded fibrils. 20 μM ThT was added to the mixture and the fluorescence was monitored by Bio-Tek Synergy 2 microplate reader using the same setting and as heparin-induced aggregation. One data point was taken every 3 minutes. Measurements were done in triplicate. The figures showed the average and standard deviations.

5.8 TEM

TEM grids were plasma cleaned (for 20s at 60W power) with the shiny side up and used within 1 hour for preparation. 1 drop (5ul) of fibril sample and two

drops (5ul each) of 2 percent UA (Uranyl Acetate) were placed on a parafilm for each grid. The shiny side was then floated on the fibril sample for 1 minute and was blotted once with Wattman filter paper. The grid was then touched with one drop of UA and blotted immediately. It was then placed on the second UA drop for 1 minute and then blotted well. The grids were dried for 2-3 hours, before imaging. Images were collected on a Talos F200X (Thermo Fisher Scientific) operating at 200 kV and equipped with a Ceta II CMOS 4k x 4k camera for high-quality HRTEM imaging.

5.9 Double Electron Electron Resonance (DEER)

Double-cysteine Tau187 was expressed and spin-labeled as doubly-labeled Tau187 that contains two spin-labels in order to probe distances between two target residues (residues 351 and 373, residues 334 and 360, and residues 340 and 378). Cysless Tau187 was expressed and purified to avoid disulfide bonding. doubly-labeled Tau187 and cysless Tau187 were stored in 20 mM HEPES in H_2O. A 1:10 molar ratio doubly-labeled Tau187:cysless Tau187 sample of 57 μM doubly-labeled Tau187 and 570 μM cysless Tu187 was incubated with 157 μM heparin at 37 °C 24 h to prepare heparin-induced fibrils. As for CBD and PSP lysate-seeded fibrils, a 1:10 molar ratio doubly-labeled Tau187:cysless Tau187 sample of 200 μM doubly-labeled Tau187 and 200 μM cysless was incubated with 15% (mass) CBD or PSP cell-passaged seeds at 37 °C for 24 h to prepare seeded fibrils. 50 μL of fibrils were then dialyzed to D_2O-based buffer ((20mM HEPES at pH 7.4) at room temperature for 6 h using Pur-A-Lyzer™ Mini Dialysis Kit (Mini 25000, MWCO 25 kDa). 35 μL samples were then mixed with 15 μL D8-glycerol (30% volume) before transferring to a quartz tube (3 mm o.d., 2 mm i.d.) and frozen using liquid nitrogen.

The DEER experiments were performed with a pulsed Q-band Bruker E580 Elexsys spectrometer, equipped with a Bruker QT-II resonator and a 300 W TWT amplifier with an output power of 20 mW for the recorded data (Applied Systems Engineering, Model 177Ka). The temperature of the cavity was maintained at 65 K using a Bruker/ColdEdge FlexLine Cryostat (Model ER 4118HV-CF100). The bridge is equipped with an Arbitrary Wave Generator to create shaped pulses for increased sensitivity. The following DEER pulse sequence was used: $\pi_{obs}/2 - \tau_1 - \pi_{obs} - (t - \pi_{pump}) - (\tau_2 - t) - \pi_{obs} - \tau_2 - $ echo. Experiment dipolar signal, V(t), was recorded as the integral of the refoucsed echo as a function of time delay, t. Rectangular observe pulses were used with lengths set to $\pi_{obs}/2 = 10$–12 ns and $\pi_{obs} = 20$–24 ns. A chirp π pump pulse was applied with a length of 100 ns and a frequency width of 60 MHz. The observed frequency was 90 MHz higher than the center of the pump frequency range. τ_1 was set to 180 ns for heparin samples and set to the first harmonic in the two-pulse deuterium ESEEM trace (126-128 ns) for cell seeded samples. τ_2 was set according to the SNR profile of the dipolar signal. The data was acquired with Δt of 16 ns, 16-step phase cycling, and signal averaged until desirable SNR was obtained.

5.10 DEER data analysis

All DEER time traces were transformed into distance distributions using Deer-Lab software package for Python. The time traces were phase corrected and truncated by 300 ns to remove possible "2+1"-artifact. One-step analysis was done using the DeerLab fit function with the following models: ex-4deer model with t1, t2, and pulselength set to experiment parameters, bg-strexp model with the stretch parameter feezed to 0.5 (base on previous measurements of dimensions of singularly labeled samples), and dipolarmodel using Tikhonove regularization. The uncertainty analysis was done using bootstrapping method with 100 samples. The time domain fitting results are presented with the fitting to the primary data and the background fit, and the distance distribution fits are presented with 95% confidence intervals.

5.11 DEER Frequency Pattern Recognition

All DEER time traces were decomposed into frequency components using discretized continuous wavelet transform (CWT). The calculated CWT was then converted into spectrograms measuring the range of frequencies present along the DEER trace. These spectrograms were compared using structure similarity index measure (SSIM) analysis by 1) numerical SSIM index, where SSIM index < 1 denotes a meaningful difference in frequency profile and by 2) similarity gradient plots, where the location with frequency differences can be visually identified.

5.12 Statistical analysis

Cell infection data are presented as mean $\pm$ SD. The value represents the averages of five images collected from each well of a 384-well plate. Technical replicates for each sample were averaged across six wells. Statistical comparisons between control and diseased patient samples, and between monomer and fibrils, were performed using a one-way ANOVA with a Dunnett post-hoc analysis. Statistical significance for all tests was determined with a P value $<$ 0.05.

6 Supplementary information

SI is attached as PDF.

7 Acknowledgments

The authors thank the DeerLab developers [53] for making an enhanced DEER analysis software available. Human tissue was obtained from the NIH NeuroBioBank and the Massachusetts Alzheimer's Disease Research Center. We acknowledge support by the Tau Consortium (www.tauconsortium.org) from the Rainwater Charitable Foundation for the study of tau seeding (S.H. and A.L.W.). The W. M. Keck Foundation (www.wmkeck.org) is acknowledged for ongoing method developments for the tau shape propagation study (S.H.). The National Institute of Health (NIH) under grant number R01AG05605 is acknowledged for supporting the aggregation mechanism study of key tau fragments (S.H., K.Z., V.V.). This work was supported by the National Institutes of Health under Grant R35GM136411 for dipolar EPR method developments (K.T.). The NIGMS grants R24GM146107 (M.S.) and R35GM151218 (M.S. and S.P.) supported the development of DEER frequency pattern recognition analysis used in this study. This work was also supported by a Venture Grant (668-2020-06) from the CurePSP Foundation, the University of Massachusetts Amherst (A.L.W.), and Colorado State University.

References

[1] Nelson, P.T., Alafuzoff, I., Bigio, E.H., Bouras, C., Braak, H., Cairns, N.J., Castellani, R.J., Crain, B.J., Davies, P., Del Tredici, K., Duyckaerts, C., Frosch, M.P., Haroutunian, V., Hof, P.R., Hulette, C.M., Hyman, B.T., Iwatsubo, T., Jellinger, K.A., Jicha, G.A., Kövari, E., Kukull, W.A., Leverenz, J.B., Love, S., Mackenzie, I.R., Mann, D.M., Masliah, E., McKee, A.C., Montine, T.J., Morris, J.C., Schneider, J.A., Sonnen, J.A., Thal, D.R., Trojanowski, J.Q., Troncoso, J.C., Wisniewski, T., Woltjer, R.L., Beach, T.G.: Correlation of alzheimer disease neuropathologic changes with cognitive status: a review of the literature. J. Neuropathol. Exp. Neurol. **71**(5), 362–381 (2012)

[2] Arriagada, P.V., Marzloff, K., Hyman, B.T.: Distribution of alzheimer-type pathologic changes in nondemented elderly individuals matches the pattern in alzheimer's disease. Neurology **42**(9), 1681–1688 (1992)

[3] Götz, J., Halliday, G., Nisbet, R.M.: Molecular pathogenesis of the tauopathies. Annu. Rev. Pathol. **14**, 239–261 (2019)

[4] Goedert, M., Eisenberg, D.S., Crowther, R.A.: Propagation of tau aggregates and neurodegeneration. Annual Review of Neuroscience, Vol 40 **40**, 189–210 (2017)

[5] Goedert, M.: Tau filaments in neurodegenerative diseases. FEBS Lett. **592**(14), 2383–2391 (2018)

[6] Walker, L.C., Jucker, M.: Neurodegenerative diseases: Expanding the prion concept. Annual Review of Neuroscience, Vol 38 **38**, 87–103 (2015)

[7] Clavaguera, F., Akatsu, H., Fraser, G., Crowther, R.A., Frank, S., Hench, J., Probst, A., Winkler, D.T., Reichwald, J., Staufenbiel, M., Ghetti, B., Goedert, M., Tolnay, M.: Brain homogenates from human tauopathies induce tau inclusions in mouse brain. Proc. Natl. Acad. Sci. U. S. A. **110**(23), 9535–9540 (2013)

[8] Guo, J.L., Lee, V.M.Y.: Cell-to-cell transmission of pathogenic proteins in neurodegenerative diseases. Nat. Med. **20**(2), 130–138 (2014)

[9] Sanders, D.W., Kaufman, S.K., DeVos, S.L., Sharma, A.M., Mirbaha, H., Li, A., Barker, S.J., Foley, A.C., Thorpe, J.R., Serpell, L.C., Miller, T.M., Grinberg, L.T., Seeley, W.W., Diamond, M.I.: Distinct tau prion strains propagate in cells and mice and define different tauopathies. Neuron **82**(6), 1271–1288 (2014)

[10] Kaufman, S.K., Sanders, D.W., Thomas, T.L., Ruchinskas, A.J., Vaquer-Alicea, J., Sharma, A.M., Miller, T.M., Diamond, M.I.: Tau prion

strains dictate patterns of cell pathology, progression rate, and regional vulnerability in vivo. Neuron **92**(4), 796–812 (2016)

[11] Woerman, A.L., Aoyagi, A., Patel, S., Kazmi, S.A., Lobach, I., Grinberg, L.T., Mckee, A.C., Seeley, W.W., Olson, S.H., Prusiner, S.B.: Tau prions from alzheimer's disease and chronic traumatic encephalopathy patients propagate in cultured cells. Proc. Natl. Acad. Sci. U. S. A. **113**(50), 8187–8196 (2016)

[12] Narasimhan, S., Guo, J.L., Changolkar, L., Stieber, A., McBride, J.D., Silva, L.V., He, Z., Zhang, B., Gathagan, R.J., Trojanowski, J.Q., Lee, V.M.Y.: Pathological tau strains from human brains recapitulate the diversity of tauopathies in nontransgenic mouse brain. J. Neurosci. **37**(47), 11406–11423 (2017)

[13] Nizynski, B., Nieznanska, H., Dec, R., Boyko, S., Dzwolak, W., Nieznanski, K.: Amyloidogenic cross-seeding of tau protein: Transient emergence of structural variants of fibrils. PLoS One **13**(7), 0201182 (2018)

[14] Mirbaha, H., Chen, D., Morazova, O.A., Ruff, K.M., Sharma, A.M., Liu, X., Goodarzi, M., Pappu, R.V., Colby, D.W., Mirzaei, H., Joachimiak, L.A., Diamond, M.I.: Inert and seed-competent tau monomers suggest structural origins of aggregation. Elife **7** (2018)

[15] Skachokova, Z., Martinisi, A., Flach, M., Sprenger, F., Naegelin, Y., Steiner-Monard, V., Sollberger, M., Monsch, A.U., Goedert, M., Tolnay, M., Winkler, D.T.: Cerebrospinal fluid from alzheimer's disease patients promotes tau aggregation in transgenic mice. Acta Neuropathol Commun **7**(1), 72 (2019)

[16] He, Z., McBride, J.D., Xu, H., Changolkar, L., Kim, S.-J., Zhang, B., Narasimhan, S., Gibbons, G.S., Guo, J.L., Kozak, M., Schellenberg, G.D., Trojanowski, J.Q., Lee, V.M.-Y.: Transmission of tauopathy strains is independent of their isoform composition. Nat. Commun. **11**(1), 7 (2020)

[17] Fitzpatrick, A.W.P., Falcon, B., He, S., Murzin, A.G., Murshudov, G., Garringer, H.J., Crowther, R.A., Ghetti, B., Goedert, M., Scheres, S.H.W.: Cryo-EM structures of tau filaments from alzheimer's disease. Nature **547**(7662), 185–190 (2017)

[18] Falcon, B., Zhang, W., Schweighauser, M., Murzin, A.G., Vidal, R., Garringer, H.J., Ghetti, B., Scheres, S.H.W., Goedert, M.: Tau filaments from multiple cases of sporadic and inherited alzheimer's disease adopt a common fold. Acta Neuropathol. **136**(5), 699–708 (2018)

[19] Zhang, W., Tarutani, A., Newell, K.L., Murzin, A.G., Matsubara, T., Falcon, B., Vidal, R., Garringer, H.J., Shi, Y., Ikeuchi, T., Murayama,

S., Ghetti, B., Hasegawa, M., Goedert, M., Scheres, S.H.W.: Novel tau filament fold in corticobasal degeneration. Nature (2020)

[20] Arakhamia, T., Lee, C.E., Carlomagno, Y., Duong, D.M., Kundinger, S.R., Wang, K., Williams, D., DeTure, M., Dickson, D.W., Cook, C.N., Seyfried, N.T., Petrucelli, L., Fitzpatrick, A.W.P.: Posttranslational modifications mediate the structural diversity of tauopathy strains. Cell **180**(4), 633–64412 (2020)

[21] Falcon, B., Zivanov, J., Zhang, W., Murzin, A.G., Garringer, H.J., Vidal, R., Crowther, R.A., Newell, K.L., Ghetti, B., Goedert, M., Scheres, S.H.W.: Novel tau filament fold in chronic traumatic encephalopathy encloses hydrophobic molecules. Nature **568**(7752), 420–423 (2019)

[22] Xiang, X., Arakhamia, T., Carlomagno, Y., Dhingra, S., Thierry, M., DeTure, M., Cook, C.N., Dickson, D.W., Petrucelli, L., Fitzpatrick, A.W.P.: Role of molecular polymorphism in defining tau filament structures in neurodegenerative diseases. bioRxiv, 2021–0524445353 (2021)

[23] Shi, Y., Zhang, W., Yang, Y., Murzin, A.G., Falcon, B., Kotecha, A., van Beers, M., Tarutani, A., Kametani, F., Garringer, H.J., Vidal, R., Hallinan, G.I., Lashley, T., Saito, Y., Murayama, S., Yoshida, M., Tanaka, H., Kakita, A., Ikeuchi, T., Robinson, A.C., Mann, D.M.A., Kovacs, G.G., Revesz, T., Ghetti, B., Hasegawa, M., Goedert, M., Scheres, S.H.W.: Structure-based classification of tauopathies. Nature **598**(7880), 359–363 (2021)

[24] Tarutani, A., Lövestam, S., Zhang, X., Kotecha, A., Robinson, A.C., Mann, D.M.A., Saito, Y., Murayama, S., Tomita, T., Goedert, M., Scheres, S.H.W., Hasegawa, M.: Cryo-EM structures of tau filaments from SH-SY5Y cells seeded with brain extracts from cases of alzheimer's disease and corticobasal degeneration. FEBS Open Bio **13**(8), 1394–1404 (2023)

[25] Mirbaha, H., Holmes, B.B., Sanders, D.W., Bieschke, J., Diamond, M.I.: Tau trimers are the minimal propagation unit spontaneously internalized to seed intracellular aggregation. J. Biol. Chem. **290**(24), 14893–14903 (2015)

[26] Zhang, W., Falcon, B., Murzin, A.G., Fan, J., Crowther, R.A., Goedert, M., Scheres, S.H.: Heparin-induced tau filaments are polymorphic and differ from those in alzheimer's and pick's diseases. Elife **8**, 43584 (2019)

[27] Fichou, Y., Vigers, M., Goring, A.K., Eschmann, N.A., Han, S.I.: Heparin-induced tau filaments are structurally heterogeneous and differ from alzheimer's disease filaments. Chem. Commun. **54**(36), 4573–4576 (2018)

[28] Frost, B., Ollesch, J., Wille, H., Diamond, M.I.: Conformational diversity

of wild-type tau fibrils specified by templated conformation change. J. Biol. Chem. **284**(6), 3546–3551 (2009)

[29] Morozova, O.A., March, Z.M., Robinson, A.S., Colby, D.W.: Conformational features of tau fibrils from alzheimer's disease brain are faithfully propagated by unmodified recombinant protein. Biochemistry **52**(40), 6960–6967 (2013)

[30] Meyer, V., Dinkel, P.D., Rickman Hager, E., Margittai, M.: Amplification of tau fibrils from minute quantities of seeds. Biochemistry **53**(36), 5804–5809 (2014)

[31] Weismiller, H.A., Murphy, R., Wei, G., Ma, B., Nussinov, R., Margittai, M.: Structural disorder in four-repeat tau fibrils reveals a new mechanism for barriers to cross-seeding of tau isoforms. J. Biol. Chem. **293**(45), 17336–17348 (2018)

[32] Ghag, G., Bhatt, N., Cantu, D.V., Guerrero-Munoz, M.J., Ellsworth, A., Sengupta, U., Kayed, R.: Soluble tau aggregates, not large fibrils, are the toxic species that display seeding and cross-seeding behavior: Generation of tau aggregates via sonication. Protein Sci. **27**(11), 1901–1909 (2018)

[33] Kraus, A., Saijo, E., Metrick, M.A. 2nd, Newell, K., Sigurdson, C.J., Zanusso, G., Ghetti, B., Caughey, B.: Seeding selectivity and ultrasensitive detection of tau aggregate conformers of alzheimer disease. Acta Neuropathol. **137**(4), 585–598 (2019)

[34] Xu, H., O'Reilly, M., Gibbons, G.S., Changolkar, L., McBride, J.D., Riddle, D.M., Zhang, B., Stieber, A., Nirschl, J., Kim, S.-J., Hoxha, K.-H., Brunden, K.R., Schellenberg, G.D., Trojanowski, J.Q., Lee, V.M.-Y.: In vitro amplification of pathogenic tau conserves disease-specific bioactive characteristics. Acta Neuropathol. **141**(2), 193–215 (2021)

[35] Lövestam, S., Li, D., Wagstaff, J.L., Kotecha, A., Kimanius, D., McLaughlin, S.H., Murzin, A.G., Freund, S.M.V., Goedert, M., Scheres, S.H.W.: Disease-specific tau filaments assemble via polymorphic intermediates. Nature **625**(7993), 119–125 (2024)

[36] Duan, P., Dregni, A.J., Xu, H., Changolkar, L., Lee, V.M.-Y., Lee, E.B., Hong, M.: Alzheimer's disease seeded tau forms paired helical filaments yet lacks seeding potential. Journal of Biological Chemistry **300**(9), 107730 (2024)

[37] Eschmann, N.A., Georgieva, E.R., Ganguly, P., Borbat, P.P., Rappaport, M.D., Akdogan, Y., Freed, J.H., Shea, J.E., Han, S.: Signature of an aggregation-prone conformation of tau. Sci. Rep. **7** (2017)

[38] Fichou, Y., Lin, Y., Rauch, J.N., Vigers, M., Zeng, Z., Srivastava, M., Keller, T.J., Freed, J.H., Kosik, K.S., Han, S.: Cofactors are essential constituents of stable and seeding-active tau fibrils. Proc. Natl. Acad. Sci. U. S. A. **115**(52), 13234–13239 (2018)

[39] Stelzl, L.S., Fowler, P.W., Sansom, M.S.P., Beckstein, O.: Flexible gates generate occluded intermediates in the transport cycle of LacY. J. Mol. Biol. **426**(3), 735–751 (2014)

[40] Vigers, M.P., Lobo, S., Najafi, S., Dubose, A., Tsay, K., Ganguly, P., Longhini, A.P., Jin, Y., Buratto, S.K., Kosik, K.S., Scott Shell, M., Shea, J.-E., Han, S.: Tau P301L mutation promotes core 4R tauopathy fibril fold through near-surface water structuring and conformational rearrangement. bioRxiv, 2023–1128568818 (2023)

[41] Longhini, A.P., DuBose, A., Lobo, S., Vijayan, V., Bai, Y., Rivera, E.K., Sala-Jarque, J., Nikitina, A., Carrettiero, D.C., Unger, M.T., Sclafani, O.R., Fu, V., Beckett, E.R., Vigers, M., Buée, L., Landrieu, I., Shell, S., Shea, J.E., Han, S., Kosik, K.S.: Precision proteoform design for 4R tau isoform selective templated aggregation. Proc. Natl. Acad. Sci. U. S. A. **121**(15), 2320456121 (2024)

[42] Woerman, A.L., Stöhr, J., Aoyagi, A., Rampersaud, R., Krejciova, Z., Watts, J.C., Ohyama, T., Patel, S., Widjaja, K., Oehler, A., Sanders, D.W., Diamond, M.I., Seeley, W.W., Middleton, L.T., Gentleman, S.M., Mordes, D.A., Südhof, T.C., Giles, K., Prusiner, S.B.: Propagation of prions causing synucleinopathies in cultured cells. Proc. Natl. Acad. Sci. U. S. A. **112**(35), 4949–58 (2015)

[43] Safar, J., Wille, H., Itri, V., Groth, D., Serban, H., Torchia, M., Cohen, F.E., Prusiner, S.B.: Eight prion strains have PrP(sc) molecules with different conformations. Nat. Med. **4**(10), 1157–1165 (1998)

[44] Barghorn, S., Mandelkow, E.: Toward a unified scheme for the aggregation of tau into alzheimer paired helical filaments. Biochemistry **41**(50), 14885–14896 (2002)

[45] Nilsson, M.R.: Techniques to study amyloid fibril formation in vitro. Methods **34**(1), 151–160 (2004)

[46] Ksiezak-Reding, H., Tracz, E., Yang, L.S., Dickson, D.W., Simon, M., Wall, J.S.: Ultrastructural instability of paired helical filaments from corticobasal degeneration as examined by scanning transmission electron microscopy. Am. J. Pathol. **149**(2), 639–651 (1996)

[47] Murray, M.E., Kouri, N., Lin, W.-L., Jack, C.R. Jr, Dickson, D.W., Vemuri, P.: Clinicopathologic assessment and imaging of tauopathies in

neurodegenerative dementias. Alzheimers. Res. Ther. **6**(1), 1 (2014)

[48] Arima, K.: Ultrastructural characteristics of tau filaments in tauopathies: immuno-electron microscopic demonstration of tau filaments in tauopathies. Neuropathology **26**(5), 475–483 (2006)

[49] Falcon, B., Zhang, W., Murzin, A.G., Murshudov, G., Garringer, H.J., Vidal, R., Crowther, R.A., Ghetti, B., Scheres, S.H.W., Goedert, M.: Structures of filaments from pick's disease reveal a novel tau protein fold. Nature **561**(7721), 137–140 (2018)

[50] Lövestam, S., Koh, F.A., van Knippenberg, B., Kotecha, A., Murzin, A.G., Goedert, M., Scheres, S.H.W.: Assembly of recombinant tau into filaments identical to those of alzheimer's disease and chronic traumatic encephalopathy. Elife **11** (2022)

[51] Hubbell, W.L., Altenbach, C.: Investigation of structure and dynamics in membrane proteins using site-directed spin labeling. Curr. Opin. Struct. Biol. **4**(4), 566–573 (1994)

[52] Lerch, M.T., Yang, Z., Brooks, E.K., Hubbell, W.L.: Mapping protein conformational heterogeneity under pressure with site-directed spin labeling and double electron-electron resonance. Proc. Natl. Acad. Sci. U. S. A. **111**(13), 1201–10 (2014)

[53] Fábregas Ibáñez, L., Jeschke, G., Stoll, S.: DeerLab: a comprehensive software package for analyzing dipolar electron paramagnetic resonance spectroscopy data. Magn Reson (Gott) **1**(2), 209–224 (2020)

[54] Sinha Roy, A., Freed, J.H., Srivastava, M.: Differentiating unimodal and multimodal distributions in pulsed dipolar spectroscopy using wavelet transforms. Appl. Magn. Reson. (2023)

[55] Ndajah, P., Kikuchi, H., Yukawa, M., Watanabe, H., Muramatsu, S.: SSIM image quality metric for denoised images (2010)

[56] Schiemann, O., Heubach, C.A., Abdullin, D., Ackermann, K., Azarkh, M., Bagryanskaya, E.G., Drescher, M., Endeward, B., Freed, J.H., Galazzo, L., Goldfarb, D., Hett, T., Esteban Hofer, L., Fábregas Ibáñez, L., Hustedt, E.J., Kucher, S., Kuprov, I., Lovett, J.E., Meyer, A., Ruthstein, S., Saxena, S., Stoll, S., Timmel, C.R., Di Valentin, M., Mchaourab, H.S., Prisner, T.F., Bode, B.E., Bordignon, E., Bennati, M., Jeschke, G.: Benchmark test and guidelines for DEER/PELDOR experiments on nitroxide-labeled biomolecules. J. Am. Chem. Soc. **143**(43), 17875–17890 (2021)

[57] Nam, W.-H., Choi, Y.P.: In vitro generation of tau aggregates conformationally distinct from parent tau seeds of alzheimer's brain. Prion **13**(1), 1–12 (2019)

[58] Kaniyappan, S., Tepper, K., Biernat, J., Chandupatla, R.R., Hübschmann, S., Irsen, S., Bicher, S., Klatt, C., Mandelkow, E.-M., Mandelkow, E.: FRET-based tau seeding assay does not represent prion-like templated assembly of tau filaments. Mol. Neurodegener. **15**(1), 39 (2020)

[59] Trinkaus, V.A., Riera-Tur, I., Martínez-Sánchez, A., Bäuerlein, F.J.B., Guo, Q., Arzberger, T., Baumeister, W., Dudanova, I., Hipp, M.S., Hartl, F.U., Fernández-Busnadiego, R.: In situ architecture of neuronal -synuclein inclusions. Nat. Commun. **12**(1), 2110 (2021)

[60] Furman, J.L., Vaquer-Alicea, J., White, C.L. 3rd, Cairns, N.J., Nelson, P.T., Diamond, M.I.: Widespread tau seeding activity at early braak stages. Acta Neuropathol. **133**(1), 91–100 (2017)

[61] Sanders, D.W., Kaufman, S.K., DeVos, S.L., Sharma, A.M., Mirbaha, H., Li, A., Barker, S.J., Foley, A.C., Thorpe, J.R., Serpell, L.C., Miller, T.M., Grinberg, L.T., Seeley, W.W., Diamond, M.I.: Distinct tau prion strains propagate in cells and mice and define different tauopathies. Neuron **82**(6), 1271–1288 (2014)

[62] Hitt, B.D., Vaquer-Alicea, J., Manon, V.A., Beaver, J.D., Kashmer, O.M., Garcia, J.N., Diamond, M.I.: Ultrasensitive tau biosensor cells detect no seeding in alzheimer's disease CSF. Acta Neuropathol Commun **9**(1), 99 (2021)

[63] Rajasekhar, K., Mirbaha, H., Diamond, M.I., et al.: Site-specific hyperphosphorylation inhibits, rather than promotes, tau fibrillization, seeding capacity and its microtubule binding. Angewandte (2019)

[64] Kfoury, N., Holmes, B.B., Jiang, H., Holtzman, D.M., Diamond, M.I.: Trans-cellular propagation of tau aggregation by fibrillar species. J. Biol. Chem. **287**(23), 19440–19451 (2012)

[65] Frost, B., Jacks, R.L., Diamond, M.I.: Propagation of tau misfolding from the outside to the inside of a cell. J. Biol. Chem. **284**(19), 12845–12852 (2009)

[66] Holmes, B.B., Furman, J.L., Mahan, T.E., Yamasaki, T.R., Mirbaha, H., Eades, W.C., Belaygorod, L., Cairns, N.J., Holtzman, D.M., Diamond, M.I.: Proteopathic tau seeding predicts tauopathy in vivo. Proc. Natl. Acad. Sci. U. S. A. **111**(41), 4376–4385 (2014)

[67] DeVos, S.L., Corjuc, B.T., Oakley, D.H., Nobuhara, C.K., Bannon, R.N.,

Chase, A., Commins, C., Gonzalez, J.A., Dooley, P.M., Frosch, M.P., Hyman, B.T.: Synaptic tau seeding precedes tau pathology in human alzheimer's disease brain. Front. Neurosci. **12**, 267 (2018)

[68] Dujardin, S., Commins, C., Lathuiliere, A., Beerepoot, P., Fernandes, A.R., Kamath, T.V., De Los Santos, M.B., Klickstein, N., Corjuc, D.L., Corjuc, B.T., Dooley, P.M., Viode, A., Oakley, D.H., Moore, B.D., Mullin, K., Jean-Gilles, D., Clark, R., Atchison, K., Moore, R., Chibnik, L.B., Tanzi, R.E., Frosch, M.P., Serrano-Pozo, A., Elwood, F., Steen, J.A., Kennedy, M.E., Hyman, B.T.: Tau molecular diversity contributes to clinical heterogeneity in alzheimer's disease. Nat. Med. **26**(8), 1256–1263 (2020)

[69] Barini, E., Plotzky, G., Mordashova, Y., Hoppe, J., Rodriguez-Correa, E., Julier, S., LePrieult, F., Mairhofer, I., Mezler, M., Biesinger, S., Cik, M., Meinhardt, M.W., Ercan-Herbst, E., Ehrnhoefer, D.E., Striebinger, A., Bodie, K., Klein, C., Gasparini, L., Schlegel, K.: Tau in the brain interstitial fluid is fragmented and seeding-competent. Neurobiol. Aging **109**, 64–77 (2022)

[70] Seidler, P.M., Boyer, D.R., Murray, K.A., Yang, T.P., Bentzel, M., Sawaya, M.R., Rosenberg, G., Cascio, D., Williams, C.K., Newell, K.L., Ghetti, B., DeTure, M.A., Dickson, D.W., Vinters, H.V., Eisenberg, D.S.: Structure-based inhibitors halt prion-like seeding by alzheimer's disease- and tauopathy-derived brain tissue samples. J. Biol. Chem. **294**(44), 16451–16464 (2019)

[71] Blennow, K., Chen, C., Cicognola, C., Wildsmith, K.R., Manser, P.T., Bohorquez, S.M.S., Zhang, Z., Xie, B., Peng, J., Hansson, O., Kvartsberg, H., Portelius, E., Zetterberg, H., Lashley, T., Brinkmalm, G., Kerchner, G.A., Weimer, R.M., Ye, K., Höglund, K.: Cerebrospinal fluid tau fragment correlates with tau PET: a candidate biomarker for tangle pathology. Brain **143**(2), 650–660 (2020)

[72] Barthélemy, N.R., Fenaille, F., Hirtz, C., Sergeant, N., Schraen-Maschke, S., Vialaret, J., Buée, L., Gabelle, A., Junot, C., Lehmann, S., Becher, F.: Tau protein quantification in human cerebrospinal fluid by targeted mass spectrometry at high sequence coverage provides insights into its primary structure heterogeneity. J. Proteome Res. **15**(2), 667–676 (2016)

[73] Horie, K., Barthélemy, N.R., Sato, C., Bateman, R.J.: CSF tau microtubule binding region identifies tau tangle and clinical stages of alzheimer's disease. Brain **144**(2), 515–527 (2021)

[74] Leuzy, A., Chiotis, K., Lemoine, L., Gillberg, P.-G., Almkvist, O., Rodriguez-Vieitez, E., Nordberg, A.: Tau PET imaging in neurodegenerative tauopathies-still a challenge. Mol. Psychiatry **24**(8), 1112–1134

(2019)

[75] Bernard-Gauthier, V., Mossine, A.V., Knight, A., Patnaik, D., Zhao, W.-N., Cheng, C., Krishnan, H.S., Xuan, L.L., Chindavong, P.S., Reis, S.A., Chen, J.M., Shao, X., Stauff, J., Arteaga, J., Sherman, P., Salem, N., Bonsall, D., Amaral, B., Varlow, C., Wells, L., Martarello, L., Patel, S., Liang, S.H., Kurumbail, R.G., Haggarty, S.J., Scott, P.J.H., Vasdev, N.: Structural basis for achieving GSK-3 inhibition with high potency, selectivity, and brain exposure for positron emission tomography imaging and drug discovery. J. Med. Chem. **62**(21), 9600–9617 (2019)

[76] Varlow, C., Vasdev, N.: Evaluation of tau radiotracers in chronic traumatic encephalopathy. J. Nucl. Med. **64**(3), 460–465 (2023)

[77] Pavlova, A., Cheng, C.Y., Kinnebrew, M., Lew, J., Dahlquist, F.W., Han, S.: Protein structural and surface water rearrangement constitute major events in the earliest aggregation stages of tau. Proc. Natl. Acad. Sci. U. S. A. **113**(2), 127–136 (2016)

8 Figures

Fig. 1 (a) Schematic of the cellular amplification of CBD and PSP patient samples followed by cell-free seeding of recombinant Tau187 using cell-passaged seeds. The inset shows the tau constructs used, including full-length 2N4R tau (FL, in green), Tau4RD (4RD, in orange), and Tau187 (in blue). (b) Schematics of the published CBD fold (PDB ID 6TJO) [21] and PSP fold (PDB ID 7P65) [19] with the three selected spin-label pairs: 351 & 373, 334 & 360, and 340 & 378. The mean distances between the spin-label pairs were measured directly in the PDB files, and are labeled within this figure. (c) Theoretical distance distributions simulated from the published cryo-EM structures of the CBD fold [21] and PSP fold [19] by the RotamerConvolveMD method [39] in the 2-8 nm range for the three spin-labeled pairs. The distance distributions across the spin-label pair at residues 351 & 373 are referred to as P(r, 351-373). The same rule applies to P(r, 334-360) and P(r, 340-378)

Fig. 2 (a) Representative images of Tau4RD*LM–YFP cells and Tau3RD*VM-YFP cells infected with CBD and PSP patient samples. YFP is shown in green. (Scale bar, 50 μm). (b) Quantification of cell infection. * = $P < 0.05$. (c) Lysates collected in RIPA buffer from naïve HEK293T cells, Tau3RD*VM-YFP cells, Tau4RD*LM-YFP cells, and Tau4RD*LM-YFP cells stably infected with CBD or PSP tau prions were analyzed for the presence of the tau-YFP fusion protein by Western blot using the GFP primary antibody (top blot). Vinculin (bottom blot) is shown as a loading control.

Fig. 3 (a) ThT fluorescence of tau187 fibrillization induced by heparin (green), Tau4RD cell-passaged seeds (lysates) from CBD (pink) or PSP (blue) patient samples for Tau187 construct, or uninfected Tau4RD cell lysate (negative control, cyan and light green). The CBD and PSP lysate-seeded aggregation generated a relatively low maximum ThT while the control lysate did not generate any ThT signal increase; an enlarged curve is shown in the inset. Tau187 (25 μM) was mixed with stoichiometric amounts of heparin (8.25 μM, 17% by mass) or cell-passaged lysates [either 5% (lighter-color lines) or 15% by mass (darker-colored lines)] and was aggregated in the presence of ThT at 37 °C. (b) Representative negative stain TEM of heparin-induced fibrils and lysate-seeded fibrils (scale bar, 500 nm.), with magnified images shown to the right (scale bar, 100 nm). Full images are shown in Fig. S3. (c) Representative images of Tau4RD*LM–YFP cells infected with monomeric Tau187, heparin-induced fibrils, and lysate-seeded fibrils. While all fibrils induced tau-YFP aggregation, monomeric Tau187 had no effect on the cells. YFP is shown in green. (Scale bar, 50 μm). (d) Quantification of tau-YFP aggregation following infection with Tau187 monomer, heparin-induced fibrils, or lysate-seeded fibrils in Tau3RD*VM-YFP cells (left) and Tau4RD*LM-YFP cells (right). *** = $P < 0.001$

Fig. 4 Intra-molecular DEER distance distribution, P(r), measured for heparin-induced fibrils (green, solid), CBD lysate-seeded fibrils (pink, solid), and PSP lysate-seeded fibrils (blue, solid). Expected distances (dotted curves) were simulated from the published cryo-EM structures of the CBD fold (PDB: 6TJO; pink, dotted) and PSP fold (PDB: 7P65; blue, dotted) [19, 21] by the RotamerConvolveMD method [39] and are illustrated on the schematic view of these folds in Fig. 1. Three spin-label pairs were measured and simulated for (a) residues 351 & 373, (b) residues 334 & 360, and (c) residues 340 & 378. Replicates are shown in SI Figures, Fig. S5 to Fig. S7. The raw DEER signal V(t) are shown with the fit (in black) and the background fit (in black dotted).

Fig. 5 Comparison of the spectrograms of the CWT of raw DEER signal V(t) shown in Fig. 4. Similarity gradient plot of DEER signal of heparin vs CBD-seeded fibrils, heparin vs PSP-seeded fibrils, and PSP vs CBD-seeded fibrils labeled at (a) residues 351 & 373, (b) residues 334 & 360, and (c) residues 340 & 378. The similarity gradient plot is shown as frequency vs time (top to bottom: high to low frequency bands) with the magnitude of similarity displayed in different colors, where red presents the least similarity and pink represents high similarity. Spectrograms are shown in Fig. S8-S10. Spectrograms of replicates are shown in Fig. S11-S14. SSIM of replicates are shown in Table SI.