Migration and standing genetic variation in vaginal and rectal yeast populations in recurrent vulvovaginal candidiasis

Abdul-Rahman Adamu Bukari[a], Rebekah J. Kukurudz-Gorowski[a], Alexia de Graaf[a]*, Devin Habon[a]*, Beamlak Manyaz[a]*, Yana Syvolos[a], Aruni Sumanarathne[a], Vanessa Poliquin[b], Aleeza Gerstein[a,b,#]

[a]Department of Microbiology, Faculty of Science, University of Manitoba, Winnipeg, Manitoba, Canada

[b]Obstetrics, Gynecology and Reproductive Sciences, University of Manitoba, Winnipeg, Manitoba, Canada

[c]Department of Statistics, Faculty of Science, University of Manitoba, Winnipeg, Manitoba, Canada

Running head: Diversity in recurrent vulvovaginal candidiasis

* Alexia de Graaf, Devin Habon, and Beamlak Manyaz contributed equally to this work and are listed alphabetically

Author Correspondence to Aleeza Gerstein, aleeza.gerstein@umanitoba.ca

Abstract

Vulvovaginal candidiasis is one of the most common vaginal and fungal infections. The majority

of infections are successfully treated with antifungal drugs. However, ~8% of cases lead to

chronic recurrent vulvovaginal candidiasis ("RVVC"), and approximately half of RVVC cases are

idiopathic. Previous research has generally found closely-related isolates within vaginal and

rectal populations and between subsequent infections. However, their coarse methods preclude

assessing the fine-scale relationships among closely related isolates and measuring standing

genetic variation, a fundamental property of populations with implications for evolutionary

potential. To address this gap, we isolated 12 vaginal and 12 rectal yeast isolates during

symptomatic relapse from four individuals with a history of RVVC. Three participants had

Candida albicans infections, while the fourth had *Nakaseomyces glabratus.* All isolates were

whole-genome sequenced and phenotyped. The isolates were placed into the global

phylogenies, which included constructing an updated *N. glabratus* tree containing over 500

isolates. Multiple analyses were consistent with frequent migration between sites. Although

there are extremely few comparables, *C. albicans* population nucleotide diversity was similar to

most commensal oral and rectal populations, while *N. glabratus* was similar to some

bloodstream infections, yet higher than others. Diversity was largely driven by single nucleotide

changes; no aneuploidies were found, and although loss-of-heterozygosity tracts were common

in the populations, only a single region on chr1L varied among isolates from one participant.

There was very little phenotypic diversity for drug response or growth and no consistent

difference between isolates from different sites for invasive growth. Combined, this study

provides baseline measurements and describes analysis techniques to quantify within-

population diversity. We highlight a critical need for comparable studies that use the same

sampling effort, sequencing, and analysis methods to understand the interplay between

selection, drift, and migration in shaping fungal microbial communities in this and other

important contexts.

Author Summary

Recurrent vaginal yeast infections are relatively common, and we do not understand why some people experience these chronic infections when many others will have a single infection that is successfully treated and cleared. Many open questions remain about the basic biology of the yeast populations involved. We quantified diversity using modern sequencing technology within vaginal and rectal yeast populations from four individuals with a history of recurrent yeast infections experiencing symptoms. Three participants had a *Candida albicans* infection (the most common causative species), while the fourth had a *Nakaseomyces glabratus* infection (the second most common and increasingly implicated). We found that vaginal and rectal isolates were closely related, indicating the same population is present at the two sites. We found, surprisingly, that diversity was similar to the yeast populations found at other body sites in healthy people. Our study highlights a critical need for additional studies to be done following the same methods in different contexts to understand better the fungal microbial populations that reside in our bodies.

Introduction

Vulvovaginal candidiasis (VVC, colloquially "yeast infection") is a mucosal infection of the lower female reproductive tract. It is very common, affecting approximately 75% of women at least once in their lives [1–3]. The disease burden of VVC results in global annual treatment costs of ~ 1.8 billion USD [4,5], with a loss in productivity in high-income countries of ~ 14 billion USD [4]. Treatment involves topical or oral antifungal medication, which is effective at symptom abatement in most cases. However, ~ 8% of women experience recurrence (RVVC), defined as three or more symptomatic episodes a year [4,5]. Even with weekly drug treatment, most people with RVVC experience symptomatic recurrence within 8-9 months, compared to 2 months in an untreated control group [6]. Many diverse factors are known to trigger recurrence or predispose individuals to RVVC [7,8], and references within], including known human genetic variants, such as mannose-binding lectin deficiency [9,10] and TLR2 Pro631His polymorphism [11]. However, approximately half of all people with RVVC have no identifiable risk factors [12], signifying the need for additional studies on the biological basis of this chronic condition.

Candida albicans is responsible for 50-90% of VVC and RVVC cases (collectively, R/VVC [13–17]). *Nakaseomyces glabratus* (formerly *Candida glabrata*) is the second most prevalent cause of R/VVC, globally attributed to ~ 8% of cases [18–20]. Here, we collectively describe these species using the colloquial term "yeast," which reflects a shared morphology, rather than the historical and still common "*Candida* spp." given our current understanding of their phylogenetic relationships and recent official renaming of *N. glabratus* [21,22]. To be consistent with clinical practice, we will continue to use the R/VVC abbreviations while noting that "candidiasis" does not reflect the updated genera names. Treatment recommendation for R/VVC differs by species, as many isolates from *N. glabratus* (and other non-*albicans* pathogenic yeast species) have intrinsic resistance to the common azole antifungals [23]. Other species-specific differences suggest there may be clinical and phylogenetic reasons not to lump these species together. For example, Pra1, whose expression in *C. albicans* sequesters zinc, promoting neutrophil recruitment and consequently leading to inflammatory vaginitis in an RVVC setting, is absent in *N. glabratus* [24,25].

Understanding the etiology of R/VVC is complicated in part since yeast exists as a common commensal member of the vaginal microbiota without causing symptomatic VVC. Identification of vaginal yeast in the absence of symptoms does not warrant treatment. Approximately 70% of healthy women were found to have vaginal yeast at a single time point through an amplicon-based study [26], which is much higher than the 20-30% asymptomatic commensal colonization estimate frequently cited from culture-based methods (e.g., [27]).

Furthermore, self-diagnosis due to symptoms has also been shown to be problematic, as other

gynecological infections and dermatologic conditions have overlapping symptoms [28]. Genetic

typing and phylogenetic strain analyses in *C. albicans* have repeatedly found no strict

phylogenetic differentiation between commensal and pathogenic strains. However, some

phylogenetic clades are potentially over-represented by commensal strains and those causing

superficial disease [29,30].

Relapse in RVVC, i.e., return of symptoms after treatment, could theoretically be due to

either incomplete eradication of the vaginal yeast population between symptomatic infections or

complete vaginal eradication followed by re-colonization from an endogenous or exogenous

population [31]. There are decades of studies that have sought to understand the etiology of

RVVC, as the answer has potential implications for improving treatment and reducing or

eliminating symptom recurrence. Re-colonization from a sexual partner has been proposed as a

possible mechanism. Yet, recurrence also occurs in non-sexually active individuals [32], and a

review of eight randomized clinical trials found that treating asymptomatic sexual partners did

not influence or cure recurrence rates [33]. The GI tract has also been suggested as a potential

endogenous source population. A study in 1979 that treated RVVC patients with oral nystatin to

reduce the resident GI population found that it did not decrease the time to recurrence [34].

Furthermore, studies examining yeast colonization of the GI tract through feces or rectal swabs

during an active infection find that not all participants are culture-positive [34–40]. However, this

does not necessarily preclude that a small GI population is present in all individuals (below the

culture detection limit in some), which could act as an endogenous reintroduction source under

the right host conditions.

Examining the diversity of strains at different body sites and among recurrent infections

can potentially differentiate between relapse scenarios. If vaginal isolates are closely related to

GI tract/rectal isolates but less diverse, that would be consistent with reintroduction. If vaginal

isolates sampled at different time points consistently have the same genotype that is not present

in the GI tract/rectal isolates, this would be consistent with incomplete eradication. Looking at

the level of diversity within a single time point during an active symptomatic infection ("standing

genetic variation") at different body sites can also help us to understand the adaptive potential

and migration dynamics of the yeast populations within the body.

The earliest studies examined within-population diversity by looking at 'resistogram' or

phenotypic profiles of strains, by typing strains based on their ability to grow in a panel of

diverse chemical challenges [40–42]. This was followed by studies using many different

methods of genomic profiling to assess relatedness among strains, such as rDNA restriction

fragment length [43], DNA/PCR fingerprinting [35,44–46], CHEF gel analysis [47], microsatellite

loci [48–51], and randomly amplified polymorphic DNA (RAPD) [46,52–54]. A higher genomic

resolution and better ability to accurately map phylogenetic relationships among divergent

strains come from recent studies using multilocus sequence typing (MLST) [17,55,56].

Considering this body of existing work, the overall picture does not change drastically with the

method employed. The overarching results are consistent with a role in the maintenance of

genotypes (i.e., the same or very similar genotype is recovered from different body sites and

within the vagina between successive symptomatic episodes), alongside an occasional

opportunity for novel genotypes to arise (i.e. reinfection, sometimes referred to as strain

turnover). However, previous studies do not necessarily rule out a role for standing genetic

variation to have been consistently present yet undetected, i.e., relapse masquerading as

reinfection, as typically only one or a small number of isolates were examined at a given time.

Only a single study sequenced two vaginal isolates from different time points through whole

genome sequencing (WGS) [57] , and no previous studies have used WGS to compare multiple

isolates from the same time point. WGS is required to accurately assess relatedness, diversity,

and the potential source of genetic novelty. WGS removes the need to rely on a single or small

number of markers, which could potentially over or underinflate the true level of diversity. As an

example, Sitterlé et al. showed that while MLST revealed occasional differences among *C.*

albicans oral isolates collected from three healthy individuals, whole-genome sequencing

revealed that the three examined isolates were closely related in each case [58].

Within-species standing genetic variation of yeasts has only been quantified in a handful

of contexts. Two studies in *C. albicans* that whole genome sequenced 3-6 oral and rectal

isolates from six healthy individuals found that although isolates were closely related in most

cases, they differed by numerous single nucleotide polymorphisms, primarily resulting from

short-range loss-of-heterozygosity tracts [30,58]. Two individuals were, however, simultaneously

colonized with oral isolates from different clades [30]. A single *N. glabratus* study sequenced up

to ten isolates from nine patients with bloodstream candidemia [59]; a pairwise SNP analysis

was also consistent with closely related isolates. Variation in RVVC has yet to be quantified

through WGS; hence, it is unknown whether it is similar to commensal populations. It also

needs to be determined how many isolates need to be sequenced to capture population

diversity accurately. It is generally thought a large sample size is required to obtain an accurate

picture of the genetic variation [60]. Unfortunately, larger samples result in extra sequencing and

computational costs. Past studies have shown that population genetics analysis can be

conducted with small sample sizes when using many single nucleotide polymorphisms (e.g.

from WGS or RADseq) [61]. Consequently, simulation and empirical analyses have shown that

the sample size can be reduced to four to six individuals when using a large number of SNPs

[60,62,63]. These studies were conducted in sexually reproducing organisms; it remains

unexplored how many yeast isolates are required to access the population level of diversity.

Here we build on previous work by using WGS paired with high-throughput phenotyping

to quantify standing genetic variation in vaginal and rectal isolates during symptomatic RVVC.

We obtained isolates from four individuals with a history of RVVC in Winnipeg, Canada. Three

participants had *C. albicans* infections, and one had *N. glabratus*. We collectively refer to these

isolates as the "THRIVE-yeast" isolates, following the name of our local umbrella research

program that studies The Host-microbial Relationships and Immune function in different Vaginal

Environments ("THRIVE," http://www.mthrive.ca). From all individuals, we found a complete

phylogenetic overlap of vaginal and rectal isolates consistent with high levels of migration. We

found no evidence diversity in populations from the two sites was different; levels of standing

genetic variation were similar to what has been observed among commensal oral and rectal

populations in *C. albicans* and some bloodstream infections in *N. glabratus*. This suggests that

despite frequent population bottlenecks caused by treatment, diversity can be maintained or

rapidly restored.

Results

THRIVE-yeast isolates in the global species phylogenies

Participants with a history of RVVC were recruited from a monthly clinic specializing in recurrent yeast infections in Winnipeg, Canada. A high number of vaginal and rectal isolates were obtained from swabs taken from four of seventeen participants who were enrolled and screened intermittently between January 2020 and November 2022. Isolates from one participant were *N. glabratus* (YST6), while *C. albicans* was isolated from the other three (TVY4, TVY10, and YST7). We haphazardly isolated 12 vaginal and 12 rectal yeast isolates from each participant. All *C. albicans* isolates were MAT-heterozygous diploids (a/α), while the *N. glabratus* isolates were all MTL1a. We collectively refer to these 93 isolates as the "THRIVE-yeast" isolates.

The phylogenetic relationship of the THRIVE-yeast isolates was evaluated in the context of available whole-genome sequenced (WGS) isolates from each species. We exhaustively searched NCBI for available *N. glabratus* sequences and, in doing so, built the largest *N. glabratus* tree to date. The 526 isolates we found were overwhelmingly blood isolates (n=430); we found only a single vaginal isolate constructed a new global *N. glabratus* phylogeny from 526 previously sequenced isolates. The YST6 isolates are monophyletic and cluster with 68 bloodstream isolates and three isolates of unknown provenance from the United States, Canada, and Australia (Fig. 1A). Unfortunately, over 80% of the sequenced isolates are bloodstream infection isolates, with only one previous isolate annotated as originating from the vagina, rectum, or stool (Table S1), precluding further analysis. The globally published *C. albicans* phylogeny comprises 182 isolates from a wide breadth of geographic and anatomical sites [29]. Isolates from TVY4, TVY10 and YST7 all form monophyletic groups and fall within a subgroup that contains fifteen additional isolates in clade 1, the most common clade (Fig. 1B). TVY4 and TVY10 isolates are beside each other, without other closely related isolates. The YST7 isolates are most closely related to three vaginal isolates (one each from Brazil, Morocco, and China) and one oral isolate from Niger. Five of the remaining twelve isolates in the clade 1 subgroup were also isolated from the vagina. This is a statistical enrichment for vaginal isolates compared to the rest of the isolates in clade 1 (THRIVE-yeast isolates from each participant were counted as a single isolate; Fisher exact test comparing 11 vaginal isolates out of 19 in the subgroup to 5 vaginal isolates out of 24 in the rest of clade 1, p = 0.011). If we discount the 35 predominantly vaginal isolates in clade 13, which is now recognized as likely a separate species (*C. africanus)* [64,65], clade 1 as a whole is also statistically overrepresented for vaginal isolates compared to the *C. albicans* tree in general (16 vaginal isolates in clade 1 out of 42 total isolates, compared to 18 vaginal isolates out of 105 total isolates; Fisher exact test, p = 0.009).

Thus, although the sequenced vaginal isolates are located in six different clades, they are over-
represented in clade 1 relative to a neutral expectation that vaginal isolates are equally likely to
be found anywhere in the existing tree.

The *N. glabratus* phylogeny is composed of long internal branches with only a small
amount of variation within the clades; clades defined through MLST are mainly consistent with
WGS [66,67]. An *in silico* MLST analysis of our WGS data found that the YST6 isolates cluster
with 71 isolates: 67 ST16, two ST60, and two ST187. ST60 is distinguished by a single
polymorphism at position 298 in *LEU2,* while ST60 is distinguished by a single polymorphism at
position 250 in *NMT1*. *C. albicans* diploid sequence types (DST) do not map as well to WGS
clades. Each of the clade 1 subgroup isolates has a different DST [29], and the differences are
mainly registered in 39 positions across the genes.

Vaginal and rectal isolates are closely related and phylogenetically overlapping.

Following the observed monophyly among isolates from the same participant, we next
assessed the relatedness of vaginal and rectal isolates. We first generated phylogenies for each
individual using RAxML [68], as previously done for closely related bacteria strains [69]. The
vaginal and rectal isolates from all four participants are phylogenetically overlapping (Fig. 2A).
Few branches within the four phylogenies had bootstrap support exceeding 80%, yet well-
supported clusters in all participants included isolates from both body sites. For each population,
we also conducted a local principal component analysis (PCA) that examines the regions of
high genomic heterogeneity within populations. From all participants, the regions of high
differences were generally distributed throughout the genome, and the PCA failed to segregate
vaginal and rectal isolates (Fig. S1).

Pairwise differences in single nucleotide polymorphisms among isolates per participant

Although isolates were closely related, WGS data picked up SNP variation among all
pairs of participant isolates. To test whether within-population vaginal diversity was lower than
within-population rectal diversity, we compared pairwise SNP differences among vaginal
isolates to the pairwise SNP differences among rectal isolates. To test whether there was a
signal of divergence between sites, we also compared single-site differences to pairwise
differences between isolates across sites. The average pairwise SNP differences between
vaginal isolates were very similar to average pairwise SNP differences between rectal isolates
and between isolates from different sites (Fig. 2B). The only significant difference was in TVY4,
where the average SNP differences among rectal isolates were significantly lower than the

vaginal isolates (YST7: $F_{2, 250}$ = 2.164, P = 0.117; YST6: $F_{2, 250}$ = 0.785, P =0.457; TVY4: $F_{2, 250}$ =

8.599, p =0.000244; TVY10: $F_{2, 250}$ = 1.66, P = 0.192; Tukey's HSD Test for multiple

comparisons; p = 0.0001). The distributions were fairly normal, as expected for isolates with low

population structure, except that TVY10 isolates showed a bimodal distribution of pairwise SNP

differences in both the vaginal and rectal sites.

Minimum number of isolates for estimating standing genetic variation within participants

A major goal of our work was to compare diversity within RVVC populations to diversity

observed in other contexts to make inferences about the evolutionary process based on the

observed degree of standing genetic variation. However, the small number of comparable

studies all sequenced different numbers of isolates, and nucleotide diversity (π) will decrease

with an increased number of samples taken from a population. To quantify the scale of this

effect of changing the number of isolates, we conducted a bootstrap analysis using our 12

vaginal isolates from each individual. We repeatedly resampled 3-10 isolates and recalculated

diversity. There was subtle variation in nucleotide diversity among the three participants with *C.*

albicans infection. In all cases, the shape of the diversity curve with the number of isolates was

very similar---an elbow was observed around n = 6 (Fig. 3). Nucleotide Diversity in YST6 (*N.*

glabratus) was two orders of magnitude lower, and there was not a consistent change in

diversity with the number of isolates.

THRIVE-yeast isolates share similar diversity as isolates from commensal and other

disease settings

We downloaded the fastq files from two previous *C. albicans* studies on commensal

populations [30,58] and used our pipeline to calculate the average nucleotide diversity for each.

For all populations, including our own, where necessary we downsampled the number of

isolates to three, consistent with the lowest number of isolates sampled from the commensal

populations. The average nucleotide diversity was very similar across most populations (Fig.

4A). The exception was populations from two participants previously shown to have isolates

from different phylogenetic clades. The YST6 vaginal isolate population was compared to

reanalyzed fastq data from nine different BSI populations from Badrane et al., which all had

between 9-10 isolates [59]. The average nucleotide diversity from our population was similar to

the diversity from four participants yet much higher than the average nucleotide diversity from

the other five (Fig. 4B).

Little variation in copy number or loss of heterozygosity within populations

We examined copy number variation (CNV) and loss-of-heterozygosity (LOH) events

among THRIVE-yeast isolates and their closest relatives using Y_{MAP} [70]. No CNVs were

identified in any YST6 isolates (Fig. 5A). A single ~50 kb CNV on the right arm of chr3 was

identified in all YST7 isolates (Fig. 5B). This CNV is also present in the closest relative to the

YST7 isolates, vaginal strain 9518, but is absent in the next two closely related strains that are

also vaginal in origin (B116 and M17). As Y_{MAP} visualizations are based on averages across

5000 bp sliding windows, we examined the region in finer detail. Coverage was measured from

the BAM files to compute the depth at each position in that region. Mapped coverage was

inconsistent with the profile of a typical CNV; the majority of the region had only a slightly

elevated copy number relative to the rest of the genome (Fig. 6C). Two small (< 200 bp) regions

spiked up to ~6-fold and ~14-fold coverage, the first internal to *ALS6* and the second to *ALS7*

(Fig. 6C). A third region comprised of elevated coverage maps to another gene with close

homology to other genes in the genome, *CYP5*, a putative peptidyl-prolyl cis-trans isomerase

[71]. The region identified in Y_{MAP} is thus likely to primarily reflect an error in mapping rather

than a true CNV with potential biological effects. No other CNVs or aneuploidies were identified

in the other THRIVE-yeast isolates.

LOH analysis was consistent with the phylogenetic analysis and diversity metrics.

Generally, all isolates from the same participant shared an LOH profile. Among the THRIVE-

yeast *C. albicans* isolates compared to the SC5314 reference, nearly all chromosomes had at

least one LOH event extending to the telomere (Fig. 6B, Fig. S2). No large interstitial LOH

events were identified. TVY4 isolates share a complicated LOH region on the left arm of chr2

that flips back and forth from the two haplotypes, as well as small LOH regions on both the left

and right arms of chr4. YST7 shares LOH blocks on the right arm of chr2 from the centromere to

the telomere, on the left arm of chr5, and on the left arm of chrR with the vaginal isolates it is

most closely related to (i.e., 9518, B116, M17) but not Niger8, an oral isolate. The exception of

wholly shared LOH blocks among participant isolates was in TVY10, where a single LOH event

on the left arm of chr1 was present in twelve TVY10 isolates that cluster together on the

phylogeny.

In addition to the characteristic LOH regions within populations, some LOH regions were

also shared between isolates from different participants. All isolates have an LOH block on the

left arm of chr3 and the right arm of chr6. TVY10 and TVY4 isolates share a ~300 kb region on

the right arm of chr1. A LOH region on chr7 beside the ancestral chr7 right arm LOH block is

also present in both TVY10 and TVY4, though it is larger in TVY10. TVY10 and YST7 (but not

TVY4) both have LOH blocks on the left arm of chr5, though the allelic profile between them is
different.
298
299   **Phenotypic variation**
300   We quantified within-population phenotypic variation in parallel to genotypic variation.
The average growth rate for YST6 *N. glabratus* isolates was higher than the *C. albicans*
populations in both Roswell Park Memorial Institute (RPMI) medium (Fig. 7A) and vaginal
simulative medium (VSM, Fig. 7B). Growth rates were either the same between vaginal and
rectal isolates or the rectal isolates were higher when grown in either medium (Welch Two
Sample t-test, Table S3, rectal isolates higher in TVY4 grown in RPMI: t = 2.50, df = 16.1, p-
value = 0.023; TVY4 grown in VSM: t = -2.36, df = 18.1, p-value = 0.030; YST7 grown in VSM: t
= -3.63, df = 18.8, p-value = 0.002). Consistent with a visual inspection, single statistical rectal
outliers with increased growth rate relative to other isolates were seen in YST7 and TVY4 grown
in RPMI, and YST6 and TVY10 grown in VSM (Fig. 7, Rosner's test for outliers). No additional
outliers were identified in the other populations.
311   We then compared drug resistance and drug tolerance from vaginal and rectal isolates
for two common drugs prescribed in our local clinic, fluconazole (FLC) and boric acid (BA). In
addition to drug resistance, drug tolerance, the ability of drug-susceptible populations to grow
slowly in the presence of high levels of fungistatic drugs, has also emerged as a trait that varies
among different fungal species and isolates [25,26]. Tolerance may be implicated in the
propensity to cause fungal disease in other infection contexts [27] but has not previously been
examined in the context of R/VVC. Drug response was assessed through disk diffusion assays
done at pH 4.2. The site of isolation was only significant for YST6 BA resistance (vaginal
isolates were slightly more tolerant than rectal isolates; t-test, t = 2.77, df = 18.2, p-value =
0.012, Table S2, Fig. 8). A visual examination of the data also indicated that the majority of
isolates were very similar to each other. Formal outlier statistical tests that grouped vaginal and
rectal isolates were broadly consistent with the qualitative visual assessment, identifying only
two outlier isolates for tolerance (both rectal isolates in BA, one from YST6, one from YST).
324   There was considerable variation among participants and isolates for invasive growth
(Fig. 8, Table S2). However, there was no difference between YST6 or TVY10 vaginal and
rectal isolates. In contrast, YST7 vaginal isolates had higher invasive growth than rectal isolates
(W = 115.5, *P* = 0.0002), and TVY4 rectal isolates exhibited higher invasive growth than vaginal
isolates (W = 372, p-value = 0.032). The overall picture is thus that invasive growth seems to
vary more between participants than between sites of isolation and that statistical differences

between sites are likely due to neutral processes rather than selection for invasive growth in the

vaginal environment.

Discussion

The biological basis of RVVC needs to be better understood. To study the vaginal yeast populations implicated in disease, we quantified the diversity of 12 vaginal and 12 rectal isolates from four people with a history of RVVC who were experiencing symptomatic recurrence at the time of sampling. In each case, the isolates formed monophyletic groups, and the vaginal and rectal isolates were phylogenetically overlapping, consistent with frequent migration between the two sites. The level of diversity within the three *C. albicans* populations was generally similar to the small number of populations from previous studies on the diversity of commensal oral and rectal, while the *N. glabratus* population was similar to or higher than populations from BSI infections. Phenotypically, there was minimal diversity in drug responses or growth ability in clinically relevant medium and no consistent difference between isolates from different isolation sites for invasive growth.

The observed phylogenetic overlap of vaginal and rectal isolates indicates that the yeast populations in these anatomical sites share a high degree of genetic similarity, suggesting a common ancestral source and a high level of ongoing migration (at least during symptomatic recurrence). Our phenotypic analyses are consistent with the genetic results, with generally low variation and non-statistically different phenotypes between the vaginal and rectal isolate populations. We primarily identified species-level differences in growth rate and drug response phenotypes, which was not unexpected, given the well-established intrinsic differences among isolates from *C. albicans* and *N. glabratus* [72–74]. This is also in concordance with previous studies that found high genetic similarity using coarse sequencing methods between vaginal and rectal isolates [52,75,76] as well as similar expression of virulence factors [77]. The proportion of participants we screened with a history of RVVC who were positive for both vaginal and rectal yeast was similar to previous culture-based studies (rectal colonization in tandem with vaginal colonization between 45-86% [35–38,78]).

We found no evidence for selection acting differently at the two sites. This could be due to similar selective pressures in both environments (leading to selection for the same traits) or could indicate that selection cannot overcome migration, potentially coupled with drift due to low effective population size. The isolates come from individuals with a long history of repeatedly taking antifungal drugs, and the yeast population sizes likely undergo many bottleneck cycles over time, functionally reducing the efficacy of selection. Nevertheless, we did observe 100s of SNP differences between all isolate pairs, indicating the presence of what has been termed microvariation, and could be of the magnitude sufficient for adaptation. These results

underscore the need for further investigations to understand the relationship between

populations at different body sites and over time in individuals with RVVC.

The observed average nucleotide diversity for the three RVVC *C. albicans* populations

was similar to what we calculated from previously sequenced commensal oral and rectal

isolates from seven healthy individuals, yet lower than two commensal oral populations that

were previously known to have isolates from different clades [30,58]. The only *N. glabratus*

study we could find that examined intra-population diversity was nine individuals with

bloodstream infections. The RVVC *N. glabratus* population had similar diversity to four BSI

populations yet was much higher than the other five [59]. It was recently shown that *N. glabratus*

isolates causing bloodstream infections were closely related to isolates from commensal sites in

the same individuals [79]. Overall, nucleotide diversity in RVVC populations was higher than we

had anticipated. Of note, we chose isolates for sequencing blind to the phenotypic data to

provide an unbiased estimate of diversity. Although it is tempting in experimental work when

resources are limited to always pick the most diverse isolates for characterization, it makes it

more difficult to compare results among studies [63]. The level of diversity we uncovered

suggests that standing genetic variation may be maintained within the vaginal yeast population,

particularly since the available comparable WGS studies purposefully selected isolates that

maximized phenotypical differences (i.e., [30,59]). Interestingly, the presence of diversity is

consistent with one of the earliest genetic studies, which used DNA fingerprinting to examine

diversity at a single time point in up to 14 vaginal isolates from six RVVC populations [80].

As whole genome sequencing becomes ever more routine, we hope additional studies

will examine intra-population diversity. Our work establishes the importance of sample size in

comparing diversity among studies in *C. albicans*. To our knowledge, our intra-population WGS

characterization of up to 12 isolates at two body sites is the largest number of isolates examined

to date in any context for *C. albicans* or *N. glabratus*. Previous work in other taxonomic groups

suggested that a sample size of four to eight isolates is sufficient to accurately estimate

population genetics parameters such as expected heterozygosity, observed heterozygosity and

pairwise genetic differentiation (i.e., F_{ST}) [60,62,63]. To conduct our analysis in the context of

microbial fungi, we subsampled from the 12 isolates and conducted a bootstrap analysis of

nucleotide diversity. From our results, we suggest that other studies interested in calculating

diversity should sequence six isolates per population, given the tradeoff between precision

(more isolates will always be better) and cost. Our results also highlight that if the goal is to

compare among studies, it is vital to ensure the same number of isolates are used for diversity

calculations. Through the use of pixy to calculate nucleotide diversity, results among studies

should be more comparable than other methods, as pixy has been shown to better deal with

bias that results from the implicit assumption that missing genotypes, including those at sites not

represented in the VCF, are homozygous for the reference allele [81]. We hope that our work

provides a baseline genetic diversity that will be used in the future to contextualize what

happens to the vaginal yeast population between symptomatic episodes and to generally act as

a launching point in studies comparing genetic diversity among different body sites and

research aims.

The use of the WGS enables us to highlight that isolates from the same population

assigned the same MLST can still differ by 100s of SNPs. Furthermore, our work also highlights

that ST groups are not always monophyletic. For example, we found two non-ST16 *N. glabratus*

isolates are nested within the ST16 clade. Hence, conclusions based on MLST should place

their results into a phylogenetic framework to assess relatedness rather than relying on ST

designation alone, and studies relying on MLST should ensure nuance in making statements

about the relatedness of isolates. Although not always feasible, WGS removes the need to rely

on a single or small number of markers, which could over or underinflate the actual level of

diversity, as this demonstrates.

Whole-genome sequencing enabled us to quantify differences among isolates in SNPs

as well as chromosomal aneuploidy, CNVs, and LOH. Non-SNP mutations have a higher

mutation rate than the per-base pair substitution rate [82–90]. Aneuploidy has been commonly

observed in yeast populations evolving to drugs in the lab [90,91] and among clinical isolates

[89,92]. Variation in LOH has similarly been found to be prevalent among closely related

isolates from other contexts in *C. albicans*, e.g., serial isolates from HIV patients with

oropharyngeal candidiasis [93], among oral commensal isolates from healthy individuals [58],

among commensal isolates from the digestive tract of healthy individuals [94], among

populations evolved in the presence of *C. elegans* [95], and among experimentally evolved *S.*

cerevisiae populations in many environments (e.g., rich medium, beer wort, wine must, high

salt, minimal medium, [96,97]). Given that RVVC individuals take drugs repeatedly [98], we

expected instances of aneuploidies, CNVs and LOH among the isolates in this study.

Interestingly, we did not observe any aneuploidy or well-supported CNV within our isolates. We

did, however, see fairly extensive variation in LOH when comparing among populations, even

though they were all from the same subclade, highlighting inter-population diversification despite

being in the same clade. Each of the populations had unique LOH tracks as well as shared LOH

tracks. The only population with variation in LOH was TVY10, where only some of the isolates

carried an LOH on chr1L. Future work will determine whether there is evidence for any unique

LOH signatures common among RVVC vaginal isolates, possibly pinpointing regions under
selection.

All three *C. albicans* RVVC populations were in a subgroup in the global phylogeny in
clade 1. This subgroup is enriched for vaginal isolates compared to the entire tree and even
compared to the other part of clade 1. Compared to different clades, a greater proportion of
clade 1 isolates have previously been noted to be significantly associated with superficial
infections [99], including in the context of VVC [17,55,56,100]. Although vaginal isolates can be
found throughout the phylogenetic tree, there may be something unique to the common
ancestor of the clade 1 subgroup that makes them more amenable to colonizing and invading
epithelial surfaces in general and hence able to cause vaginal diseases [101,102]. Most
phenotypes of potential clinical interest have previously been found to vary among isolates
within the same clade, precluding clear phenotype × clade associations (e.g., [103,104]).
However, it may be that more fine-scale phylogenetic resolution is required to tease apart
relationships; if only a subgroup of clade 1 isolates is enriched for a particular phenotype, that
might not be seen if all clade 1 isolates are grouped together. Sala *et al.* recently found that
VVC isolates induced greater fungal shedding from epithelial cells and differently stimulated
epithelial signaling pathways compared to isolates from healthy women [105]. This is the
clearest *in vitro* assay able to differentiate VVC and healthy isolates phenotypically, and hence,
a strong target for a GWAS analysis to potentially pinpoint the genetic basis of this seemingly
important trait. It will be of great interest in the future to determine whether there is a genotypic
association between common variants in the subgroup of clade 1 isolates (including isolates
from other body sites) and their interaction with vaginal epithelial cells.

Significantly less work has been done to examine *N. glabratus* in the context of R/VVC
compared to *C. albicans*, despite the increasing incidence of *N. glabratus* globally as an
etiological agent of R/VVC [18,106]. We found only a single vaginal isolate out of 526 total
isolates that have been whole genome sequenced. This sharply contrasts with *C. albicans*,
where 35 % of all sequenced isolates in the current phylogeny have been annotated as vaginal
origin [29]. With only two independent vaginal isolates (since the RVVC isolates from our study
are not independent, we only count them as one), this extremely low number precludes making
any conclusions. There are also only two vaginal isolates in the *N. glabratus* MLST database
[107] out of over 2000 isolates. Interestingly, they all belong to ST16, which is also the ST of the
RVVC *N. glabratus* population we analyzed. All three are from North America; ST16 is the most
common group in the US [79] and is seldom found in other areas such as Australia and Asia
[108–111]. Although there is ongoing debate about the utility of renaming *N. glabratus* away

from *C. glabrata* [112], the closest common ancestor is from over 100 million years ago [113].

N. glabratus exhibits different pathogenesis and immunological responses compared to *C.*

albicans [18,114,115] and differs in the approach to inactivating non-beneficial nutrients, such

as excess copper and zinc [116–118]. Given the long evolutionary divergence and biological

differences, it may be that lumping these species together under the colloquial vaginal "yeast

infection" downplays differences that could have disease implications [115].

We have conducted the most extensive study to date that employed whole genome

sequencing, modern methods for calculating and comparing diversity, and high throughput

phenotypic analyses to compare vaginal and rectal isolates from symptomatic RVVC infections.

We find no evidence that rectal isolates are different than vaginal isolates, which is inconsistent

with the hypothesis that the GI tract is a source population for vaginal reinfection. We observed

a near-identical average nucleotide diversity between our populations and some populations

from commensal (*C. albicans*) and BSI (*N. glabratus*) settings. It remains unanswered how the

nucleotide diversity is kept at such elevated levels despite repeated use of antifungals by

individuals with RVVC. Combined, this study provides relevant information on the baseline

diversity of RVVC vaginal and rectal populations. It emphasizes the need to investigate the role

of other body sites in shaping fungal microbial communities across various contexts.

Methods

Clinical isolates

Clinical samples were collected from four non-pregnant female participants between the

ages of 18-50 years with a history of recurrent vulvovaginal candidiasis (RVVC). Seventeen

consenting female participants who attended a clinic for individuals with a history of RVVC in

Winnipeg, Canada, were sampled at the clinic for possible inclusion in this study. Vaginal and

rectal swabs were acquired from all participants. Swabs were kept at -4 ℃ and on ice during

transport and processed within 5 h of acquisition. Swabs were then agitated for ~30 s in 1 mL

PBS, and a dilution series (1, 1:10, 1:100) was conducted. 100 µL from each of the three

dilutions was spread onto SDA and chromogenic *Candida* agar plates. Plates were incubated

for 48 h at 30 °C, at which time 12 vaginal and 12 rectal isolates were haphazardly isolated

from dilution plates that had clear margins between colonies. Colonies were suspended in 1 mL

of 20% glycerol and kept at -70 °C. We collectively refer to all colonies collected in this study as

"THRIVE-yeast" isolates. Of the seventeen participants, vaginal colonies were acquired from

ten. Of those, six were also positive for rectal isolates; we recovered less than ten rectal

colonies from the two participants who were rectal swab positive, leaving us with samples from

four participants for this project. This study has been approved by the University of Manitoba

Biomedical Research Ethics Board (HS24769 (B2021:026)) and Shared Health (SH2021:038).

DNA extraction and sequencing

Genomic DNA was extracted from 24 single colonies (12 each from the vagina and

rectum) from one individual with an *N. glabratus* infection (YST6) and three individuals with *C.*

albicans infections (YST7, TVY4, TVY10). We followed a standard phenol-chloroform protocol

as previously described [91]. DNA quality and concentration were assessed on a Thermo

Scientific™ NanoDrop 2000 and Qubit® 2.0 Fluorometer, respectively. Genomic DNA was sent

to either Microbial Genome Sequencing Center ("MIGS," Pittsburgh, USA; YST6 and YST7) or

SeqCoast Genomics (New Hampshire Ave., USA; TVY4 and TVY10) for sequencing. At MIGS,

sample libraries were prepared using the Illumina DNA Prep kit and IDT 10bp UDI indices and

sequenced on NextSeq 2000 using a 300-cycle flow cell kit, producing 2×151bp reads. The bcl-

convert v3.9.3 software was used to assess read quality, demultiplex and trim adapter

sequences. At SeqCoast Genomics, samples were prepared using an Illumina DNA Prep

tagmentation kit and unique dual indexes. Sequencing was performed on the Illumina

NextSeq2000 platform using a 300-cycle flow cell kit to produce 2×150bp paired reads.

DRAGEN v3.10.11 was used to assess read quality, demultiplex and trim adapter sequences.

The reads have been deposited at the National Center for Biotechnology Information (NCBI)

Sequence Read Archive under BioProject ID PRJNA991137.

In addition to the 93 genomes we sequenced, we downloaded an additional 182 *C.*

albicans FASTQ files from the NCBI Sequence Read Archive database [119] from BioProject

Accession PRJNA432884 [29] and 526 *N. glabratus* FASTQ files from 19 different projects on

the SRA database (assessed on February 05, 2022), including 99 *N. glabratus* FASTQ files

from SRA PRJNA361477 [67] and PRJNA669061 [66] (see Table S1 and Table S3).

Variant calling

The sequence reads were trimmed with Trimmomatic (v0.39) [120] with standard

parameters (LEADING: 10, TRAILING: 3, SLIDINGWINDOW:4:15, MINLEN: 31, TOPHRED33,

following [121]). Quality was assessed with FASTQC

(http://www.bioinformatics.babraham.ac.uk/projects/fastqc/) and MultiQC [122]. *C. albicans*

trimmed paired-end reads were mapped using bwa-mem [123] to the SC5314 haplotype A

reference genome (A22-s07-m01-r160) downloaded from the Candida Genome Database [124].

The resulting SAM file was coordinate-sorted and converted to a Binary Alignment Map (BAM)

file using samtools v1.9 [125]. *N. glabratus* isolates were mapped to the CBS138 reference

genome (GCA000002545v2) downloaded from the Ensembl Genome Database [126].

Alignment quality was assessed with CollectAlignmentSummaryMetrics from Picard v2.26.3

(http://broadinstitute.github.io/picard) and consolidated across all samples with MultiQC [122].

All files had a >95% mapping quality. BAM files were further processed with Picard by adding a

read group annotation so that samples with the same Bioproject ID had the same read group,

removing duplicate PCR amplicons, and fixing mate pairs. Base quality scores for the *C.*

albicans aligned reads were recalibrated with known single-nucleotide polymorphisms obtained

from the Candida Genome Database website

(http://www.candidagenome.org/download/gff/C_albicans_SC5314/Assembly22/A22_Jones_PM

ID_15123810_Polymorphisms.vcf; downloaded on July 29, 2020) [127] using the

BaseRecalibrator and ApplyBQSR from the Genome Analysis Toolkit 4.2.4.0. The average

coverage for each isolate was estimated using samtools v1.9 [125].

The GATK Best Practices were adapted for variant calling. In sequence,

HaplotypeCaller, CombineGVCFs, GenotypeVCFs, VariantFiltration, and SelectVariants [128–

130] were used to identify single nucleotide variants (SNPs) among all sequenced isolates in

diploid and haploid mode for *C. albicans* and *N. glabratus* respectively. The results SNP table

was hard filtered using the suggested parameters and to match [29] (QualByDepth <2.0,

FisherStrand >60.0, root mean square mapping quality <30.0, MappingQualityRankSumTest <

−12.5, ReadPosRankSumTest < −8.0). We excluded variants that were called in known

repetitive regions of the genome, as these are likely to reflect sequencing misalignments rather

than true variants, i.e., the subtelomeric regions (15kb from the start and end of each

chromosome), the centromeres, and the major repeat sequence regions, (Table S4, start and

stop positions from candidagenome.org).

Phylogeny construction

Phylogenetic trees were constructed for *C. albicans* and *N. glabratus* isolates. For each

species, the multi-sample VCF file consisting of only genomic DNA SNPs was converted to a

FASTA alignment using a publicly available Python script that creates an alignment matrix for

phylogenetic analysis (vcf2phylip.py v2.8, downloaded from

https://github.com/edgardomortiz/vcf2phylip) [131]. For heterozygous SNPs in *C. albicans,* the

consensus sequence is preferentially made based on the reference (haplotype A) base.

Ambiguous bases are written following IUPAC nucleotide ambiguity codes in the matrix. Ploidy

is not specified, as the most parsimonious ploidy is detected by the script. The FASTA

alignment was parsed in FastTree (2.1.11) [132] in the double precision mode to construct an

approximate maximum-likelihood phylogenetic tree using the general time reversible model and

the -gamma option to rescale the branch lengths. Compared to the time-consuming ML-based

phylogeny predictors such as RAxML [68], FastTree has been found to produce large dataset

trees with similar accuracy within a significantly shorter amount of time [133]. The phylogeny

was visualized and annotated with the Interactive Tree Of Life (iTOL, v5) [134]. Isolates from *C.*

albicans clade 13, which is believed to be a closely related but different species termed *C.*

africanus [64,65,135] was used to root the phylogeny. The *N. glabratus* phylogeny was rooted at

the midpoint. While constructing the phylogeny with FastTree with this large data set yielded a

phylogeny comparable to *C. albicans* whole-genome phylogenies previously generated by other

groups using RAxML [29,136], an attempt to use FastTree to resolve the relatedness of the

THRIVE-yeast isolates yielded poor resolution, as most of the SNP sites were ignored during

the tree construction. Hence to generate a more-refined phylogeny depicting the relatedness

among the isolates from each participant, we constructed a maximum likelihood phylogeny with

RAxML v8.2.12 [68] following standard practice using the GTR+G model with 20ML inferences

on the alignment, inferring bootstrap replicate trees, applying MRE-based bootstrapping test,

and drawing support values using TBE (Transfer Bootstrap Expectation) on the best-scoring

tree.

Pairwise SNP differences between isolates from the same site in a participant

We also looked at the variation in SNPs among isolates from the same participant within

a site. Using RTG tools with the vcfsplit option [137], the VCF files for each isolate were

extracted from the multi-sample VCF file. Using "bcftools query -f" the VCF files were converted

to bed files. A custom R script was used to do a pairwise comparison of the isolates from a site

to determine differences in SNP positions. An ANOVA test was performed to compare pairwise

SNP differences in the rectal and vaginal sites in each participant.

In silico **MLST typing**

As clades in *N. glabratus* are commonly identified and named through MLST analysis

([66,67]), we conducted *in silico* MLST analyses of the YST6 isolates as well as the cluster in

which the YST6 isolated are associated with using stringMLST [138] which uses a predefined

MLST *N. glabratus* allele library (FKS, LEU2, NMT1, TRP1, UGP1, URA3) in the PubMLST

database [139]. The diploid sequence types (DSTs) of the global *C. albicans* isolates were

determined in a previous study using a PCR-based method [29]. *In silico* identification of DST

from the WGS data was error-prone and unable to replicate known DSTs.

Determining segregating genomic blocks within isolates from the same participant

We conducted a principal component analysis of windows of genomic regions that

differed among isolates from the same participant. Briefly, used the "templated script" from the

R package lostruct (local PCA/population structure, v.0.0.0.9) [140] with run parameters -t

(window type): bp, -s (window size): 5000, -npc (number of pcs): 2 and -m (number of MDS

coordinates): 2. To check for possible segregation of vaginal and rectal isolates, principal

component analysis plots were generated based on the extreme heterogeneous genomic

windows across the entire genome.

Relationship between average nucleotide diversity (π) and number of samples

We assessed the potential limitation of the sample size using average pairwise diversity

differences between all possible isolate pair estimates from Pixy [81] in a randomized

bootstrapping framework. We estimated the π for different numbers of samples (n = 2,3,4,6,10)

from the same population. Each number (n) was randomly selected from the vaginal samples,

and the π was estimated with Pixy [81]. Additionally, n = 12 was generated by randomly

selecting 6 samples each from the rectal and vaginal isolates and combined. All random

selections were bootstrapped 50 times, and the standard deviation was estimated. Finally, the

number of nucleotide diversity of the total number of isolates per participant (i.e. n=23 for

TVY10, TV4, YST7 and n=24 for YST6) was also included. We estimated the nucleotide

diversity within (π) using Pixy (v1.2.6.beta1) [81]. Briefly, variants were called using

GenotypeGVCFs with --all-sites option activated. Vcftools (v0.1.16) [141] was used to filter the

variants (with --max-meanDP 500, --min-meanDP 20, --max-missing 0.8). Indels and

mitochondrial DNA were excluded (--remove-indels, --not-chr). The nucleotide diversity was

then plotted against the number of isolates to assess the value of *n* where near π saturation

was achieved.

Nucleotide diversity from published isolates from other studies

We compared our nucleotide diversities to instances where multiple isolates have been

retrieved in either commensal or disease settings. For *C. albicans*, 3-4 isolates from 6 and 3

individuals respectively from Andersen *et al.* [30] and Sitterlé *et al*. [58] were assessed. All the

isolates are from commensal settings in oral and rectal sites. For *N. grabratus*, nine to ten isolates

from blood cultures in each of the nine patients [59] were compared to the 12 YST6 isolates.

In silico **mating-type locus detection**

To determine mating type-like locus in the *C. albicans* isolates (YST7, TVY4 and

TVY10), the reads were aligned to both haplotypes and consensus sequences of the MAT locus

on chromosome 5 (MATa1 and MATa2 for hapA and MATα1 and MATα2) were extracted. A

BLAST search was then conducted to confirm the locus. Similarly, the mating type-like locus of

N. glabratus (YST6) isolates was determined by determining the consensus sequence for MTL1

(MTLalpha1 and MTLalpha2) and MTL3 on chromosome B, and MTL2 on chromosome E and

confirming the MTL by a BLAST search.

Loss of heterozygosity and copy number analyses

We conducted genome-wide loss of heterozygosity (LOH) and copy number variant

(CNV) analyses using the web-based yeast analysis pipeline (Y_{MAP}) [70]. Mitochondrial DNA is

by default excluded from the pipeline. Paired-end reads of the *N. glabratus* isolates were

uploaded and analyzed to the CBS138 reference genome (CGD: s05-m01-r09). As *N. glabratus*

is haploid, the ploidy was set as 1, which by default excludes LOH analyses. For the *C. albicans*

isolates, paired-end read data for each isolate was uploaded and analyzed against the SC5314

A22-s02-m09-r10 reference genome. Ploidy was left at the default value (two, i.e. diploid), and

correction was enabled for GC-content bias and chromosome-end bias. THRIVE-yeast isolates

were compared against the SC5314 haplotype map to distinguish between ancestral and newly

evolved LOH signatures and determine which allele was retained. CNV profiles were compared

to the reference SC5134 isolate as well as closely related isolates in the phylogeny. The

genomic elements within observed CNV regions were identified using the "gene/sequence

resources" section of the Candida Genome Database (CGD, https://candidagenome.org).

Growth Rate Assay

Two separate growth rate assays were conducted to measure growth in RPMI (10.4 %

w/v RPMI powder, 1.5 % w/v dextrose, 1.73 % w/v 3-(N-morpholino) propanesulfonic acid

(MOPS), adjusted to pH 7 with NaOH tablets) and vaginal simulative medium ("VSM", following

[29]: 1.16 % v/v 5 mM NaCl, 3.6 % v/v 0.5 M KOH, 0.0128 % v/v 99% glycerol, 20 % v/v 0.01 M

$Ca(OH)_2$, 1.34 % v/v 0.5 M Urea, 6.6 % v/v 0.5 M glucose, 0.67% w/v solid YNB, 0.85 % v/v 2 M

acetic acid, 0.192 % v/v lactic acid, adjusted to pH 4.2 with NaOH tablets). For each experiment,

5 µL of frozen glycerol stock from all THRIVE-yeast isolates was inoculated in duplicate into 500

µL of RPMI or VSM and incubated for 48 h at 37 °C with agitation at 250 rpm. Cultures were

then standardized to an optical density (OD) of 0.01 A600 in RPMI or VSM, and 200 µL was

transferred into a 96-well round bottom plate and sealed with a Breathe-Easier sealing

membrane (Electron Microscopy Sciences, PA, United States). OD_{600} readings were taken by

the Epoch plate reader (Biotek) every 15 minutes, with continuous shaking at 37 °C for 48 h.

From each well, the maximal growth rate was calculated as the spline with the highest slope

using a custom R script written by Dr. Richard Fitzjohn (https://github.com/acgerstein/THRIVE-

variation/scripts_real/growthrate_generic.R). The average growth rate between two technical

replicates for each isolate in each growth medium was used for visualization and statistical

analysis. Statistical outliers were determined through Rosner's test of outliers available through

the rosterTest function in the EnvStats R package [142]. For each population, we started with a

k value of one (i.e., testing for a single outlier). If that was significant, we increased k by one

until no additional outliers were identified.

Drug resistance and tolerance

Disk diffusion assays were carried out to measure resistance and tolerance to

fluconazole and boric acid at pH 4.2. Standard CLSI M44 guidelines for antifungal disk diffusion

susceptibility testing [143] were generally followed except for pH adjustment as appropriate. To

initiate each experiment, frozen stock of all THRIVE-yeast was streaked onto Sabouraud

dextrose agar (SDA) plates and incubated for 72 h at room temperature. Isolates were then

subcultured by streaking one colony onto a fresh SDA plate and incubating at 37 ºC for 24 h.

Colonies from each isolate were then suspended in 200 µL of 0.85% saline solution and

standardized to an OD_{600} of 0.01 in 1 mL of saline solution. Within 15 minutes of

standardization, 100 µL of the culture was spread evenly using autoclaved glass beads on

Mueller-Hinton (MH) plates that had been adjusted to a pH of 4.2. Plates were left to dry for 20

minutes before placing a single 5 mg boric acid or 25 mg fluconazole disk in the center of the

plate. Plates were incubated upside down at 37 ºC. After 48 h, photographs of each plate were

taken on a lightbox. Previous work demonstrated consistent drug resistance values at 24 h and

48 h. The entire experiment was conducted twice for each isolate × drug × pH combination, with

two technical replicates for each of the two biological replicates.

The 48 h images were processed in ImageJ as previously described [72]. They were

cropped to a uniform size, the colours were inverted, and brightness and contrast were adjusted

to maximize the contrast between the white disk and the black background. The adjusted

images were run through the diskImageR package [144] for drug resistance (RAD_{20}) and

tolerance (FOG_{20}) quantification. Briefly, diskImageR calculates resistance as RAD_{20}, as the

radius of the zone of inhibition where growth is reduced by 20% relative to growth on the

margins of the plate where there is no drug, and tolerance as FoG_{20}, the fraction of realized

growth between RAD_{20} and the disk.

A Welch two-sample t-test that did not assume equal variance was used for each

participant × drug combination to compare vaginal and rectal isolates. Statistical outliers were

determined through Rosner's test of outliers for growth rates. All statistical analysis was done at

a type I error rate of 0.05.

Invasive growth assay

To examine invasive growth, we revised the methods from [145]. Freezer stock from

THRIVE-yeast isolates were streaked onto 20 mL yeast peptone dextrose (YPD) plates (2% w/v

peptone, 2% w/v yeast extract, 1.8% w/v agar, 1% w/v glucose, 0.00016% w/v adenine

sulphate, 0.00008% w/v uridine, 0.1% v/v of chloramphenicol and ampicillin) and grown for 72 h

at room temperature. A single colony was then randomly chosen from each isolate and

inoculated into 200 μL YPD. If no single colonies were available, a similar amount of culture

from the colony lawn was used. Cultures were standardized to OD_{600} 0.01 in 1 mL of liquid YPD

media, then 2 μL of standardized culture was spotted onto the surface of a 20 mL solid YPD

plate in a hexagonal pattern for a total of 7 spots per plate (i.e., spotted at each vertex and in

the center). Plates were incubated for 96 h at 37 ºC. The surface growth was washed off using

distilled water and a photograph was taken in a dark room on a lightbox. Two biological

replicates were performed for each isolate.

The qualitative amount of invasive growth for each isolate was determined by visual

examination of the post-wash photographs. To develop a five-point scale, two different people

independently went through the post-wash pictures from YST6 and YST7 and selected two to

six representative pictures that fit into five levels of invasive growth (scored as 1-5). The

independent selections were then compared, and one image from each person was chosen as

the most representative for each level of the scale. Using these as a reference, each isolate was

then categorized into the five levels of the scale (0 - no growth/pipette tip indent, 0.25 - pinprick

growth, 0.5 - circular growth evident, 0.75 - circular growth with pinprick, 1 - dense growth

throughout). The maximum score between the two bio-replicates of each isolate was used for

statistical analysis, though the same statistical conclusions were obtained if the mean score was

used instead. For each participant, a Wilcoxon rank sum test was used to compare vaginal and

rectal isolates.

Data availability

FASTQ files generated for this project have been deposited at the National Center for

Biotechnology Information (NCBI) Sequence Read Archive under BioProject ID PRJNA991137.

All phenotypic data and code required to reproduce figures and statistical analyses are available

at https://github.com/acgerstein/THRIVE_yeast-VR. Large files (e.g., BAM, VCF) are available

upon request.

Acknowledgements & Funding

We are extremely thankful to the THRIVE-yeast participants. We are grateful to Dr. Alicia

Berard, Dr. Adam Burgener, Kenzie Birse and other THRIVE investigators for sharing their

knowledge and resources, which were integral in the initiation of this study and thank other

members of the THRIVE-yeast research team including study clinicians, laboratory and

administrative staff for their many contributions. The study was funded through the Manitoba

Medical Service Foundation, a University of Manitoba Faculty of Science Collaborative grant,

the Canadian Institute for Advanced Research (CIFAR), and the National Science and

Engineering Rearch Council of Canada. ACG acknowledges the support of the CIFAR Azrieli

Global Scholars Program and start-up funding from the University of Manitoba. A-RAB and AS

were supported by EvoFunPath (NSERC CREATE) fellowships. DAH was funded by a

University of Manitoba Research Award and the University of Manitoba Department of

Microbiology Fellowship for Education Purposes. AdG was supported by a University of

Manitoba Science Students' Association and the Faculty of Science Undergraduate Student

Research Award (USRA). BM was supported by a Faculty of Science USRA. Some swabs and

Amies buffer tubes were donated by COPAN Italia.

References:

1. Benedict K, Singleton AL, Jackson BR, Molinari NAM. Survey of incidence, lifetime prevalence, and treatment of self-reported vulvovaginal candidiasis, United States, 2020. BMC Womens Health. 2022;22: 147.

2. Yano J, Sobel JD, Nyirjesy P, Sobel R, Williams VL, Yu Q, et al. Current patient perspectives of vulvovaginal candidiasis: incidence, symptoms, management and post-treatment outcomes. BMC Womens Health. 2019;19: 48.

3. Rathod SD, Buffler PA. Highly-cited estimates of the cumulative incidence and recurrence of vulvovaginal candidiasis are inadequately documented. BMC Womens Health. 2014;14: 43.

4. Denning DW, Kneale M, Sobel JD, Rautemaa-Richardson R. Global burden of recurrent vulvovaginal candidiasis: a systematic review. Lancet Infect Dis. 2018;18: e339–e347.

5. Foxman B, Barlow R, D'Arcy H, Gillespie B, Sobel JD. Candida vaginitis: self-reported incidence and associated costs. Sex Transm Dis. 2000;27: 230–235.

6. Donders G, Sziller IO, Paavonen J, Hay P, de Seta F, Bohbot JM, et al. Management of recurrent vulvovaginal candidosis: Narrative review of the literature and European expert panel opinion. Front Cell Infect Microbiol. 2022;12: 934353.

7. Rosati D, Bruno M, Jaeger M, Ten Oever J, Netea MG. Recurrent vulvovaginal candidiasis: An immunological perspective. Microorganisms. 2020;8. doi:10.3390/microorganisms8020144

8. Fidel PL Jr. History and update on host defense against vaginal candidiasis. Am J Reprod Immunol. 2007;57: 2–12.

9. Babula O, Lazdane G, Kroica J, Ledger WJ, Witkin SS. Relation between recurrent vulvovaginal candidiasis, vaginal concentrations of mannose-binding lectin, and a mannose-binding lectin gene polymorphism in Latvian women. Clin Infect Dis. 2003;37: 733–737.

10. Wojitani MDKH, de Aguiar LM, Baracat EC, Linhares IM. Association between mannose-binding lectin and interleukin-1 receptor antagonist gene polymorphisms and recurrent vulvovaginal candidiasis. Arch Gynecol Obstet. 2012;285: 149–153.

11. Rosentul DC, Delsing CE, Jaeger M, Plantinga TS, Oosting M, Costantini I, et al. Gene polymorphisms in pattern recognition receptors and susceptibility to idiopathic recurrent vulvovaginal candidiasis. Front Microbiol. 2014;5: 483.

12. Sobel JD. Management of patients with recurrent vulvovaginal candidiasis. Drugs. 2003;63: 1059–1066.

13. Zhang J-Y, Liu J-H, Liu F-D, Xia Y-H, Wang J, Liu X, et al. Vulvovaginal candidiasis: species distribution, fluconazole resistance and drug efflux pump gene overexpression. Mycoses. 2014;57: 584–591.

14. Shi X-Y, Yang Y-P, Zhang Y, Li W, Wang J-D, Huang W-M, et al. Molecular identification and antifungal susceptibility of 186 Candida isolates from vulvovaginal candidiasis in

southern China. J Med Microbiol. 2015;64: 390–393.

15. Guzel AB, Ilkit M, Akar T, Burgut R, Demir SC. Evaluation of risk factors in patients with
vulvovaginal candidiasis and the value of chromID Candida agar versus CHROMagar
Candida for recovery and presumptive identification of vaginal yeast species. Med Mycol.
2011;49: 16–25.

16. van Schalkwyk J, Yudin MH, INFECTIOUS DISEASE COMMITTEE. Vulvovaginitis:
screening for and management of trichomoniasis, vulvovaginal candidiasis, and bacterial
vaginosis. J Obstet Gynaecol Can. 2015;37: 266–274.

17. Song N, Kan S, Pang Q, Mei H, Zheng H, Li D, et al. A prospective study on vulvovaginal
candidiasis: multicentre molecular epidemiology of pathogenic yeasts in China. J Eur Acad
Dermatol Venereol. 2022;36: 566–572.

18. Kennedy MA, Sobel JD. Vulvovaginal candidiasis caused by non-albicans Candida species:
New insights. Curr Infect Dis Rep. 2010;12: 465–470.

19. Parazzini F, Di Cintio E, Chiantera V, Guaschino S. Determinants of different *Candida*
species infections of the genital tract in women. Eur J Obstet Gynecol Reprod Biol.
2000;93: 141–145.

20. Richter SS, Galask RP, Messer SA, Hollis RJ, Diekema DJ, Pfaller MA. Antifungal
susceptibilities of *Candida* species causing vulvovaginitis and epidemiology of recurrent
cases. J Clin Microbiol. 2005;43: 2155–2162.

21. Borman AM, Johnson EM. Name changes for fungi of medical importance, 2018 to 2019. J
Clin Microbiol. 2021;59. doi:10.1128/JCM.01811-20

22. Kidd SE, Abdolrasouli A, Hagen F. Fungal nomenclature: Managing change is the name of
the game. Open Forum Infect Dis. 2023;10: ofac559.

23. Pfaller MA, Rhomberg PR, Messer SA, Jones RN, Castanheira M. Isavuconazole,
micafungin, and 8 comparator antifungal agents' susceptibility profiles for common and
uncommon opportunistic fungi collected in 2013: temporal analysis of antifungal drug
resistance using CLSI species-specific clinical breakpoints and proposed epidemiological
cutoff values. Diagn Microbiol Infect Dis. 2015;82: 303–313.

24. Citiulo F, Jacobsen ID, Miramón P, Schild L, Brunke S, Zipfel P, et al. *Candida albicans*
scavenges host zinc via Pra1 during endothelial invasion. PLoS Pathog. 2012;8: e1002777.

25. Roselletti E, Pericolini E, Nore A, Takacs P, Kozma B, Sala A, et al. Zinc prevents vaginal
candidiasis by inhibiting expression of an inflammatory fungal protein. Sci Transl Med.
2023;15: eadi3363.

26. Drell T, Lillsaar T, Tummeleht L, Simm J, Aaspõllu A, Väin E, et al. Characterization of the
vaginal micro- and mycobiome in asymptomatic reproductive-age Estonian women. PLoS
One. 2013;8: e54379.

27. Beigi RH, Meyn LA, Moore DM, Krohn MA, Hillier SL. Vaginal yeast colonization in
nonpregnant women: A longitudinal study. Obstetrics & Gynecology. 2004;104: 926.

28. Bullarbo M, Andersch B, Samuelson E, Lindgren A, Kondori N, Baltzer IM. Self-diagnosis of

vulvovaginal candidiasis is poor - A comparison of diagnostic methods introducing β-glucan
as a complement. Bioenergetics. 2017;06. doi:10.4172/2161-038x.1000202

29. Ropars J, Maufrais C, Diogo D, Marcet-Houben M, Perin A, Sertour N, et al. Gene flow
contributes to diversification of the major fungal pathogen *Candida albicans*. Nat Commun.
2018;9: 2253.

30. Anderson FM, Visser ND, Amses KR, Hodgins-Davis A, Weber AM, Metzner KM, et al.
Candida albicans selection for human commensalism results in substantial within-host
diversity without decreasing fitness for invasive disease. PLoS Biol. 2023;21: e3001822.

31. Tasić S, Tasić N, Tasić A, Mitrović S. Recurrent genital candidosis of women; consequence
of reinfection of relapse. Med Biol. 2002;9. Available: http://dx.doi.org/

32. Lanchares JL, Hernández ML. Recurrent vaginal candidiasis changes in etiopathogenical
patterns. Int J Gynaecol Obstet. 2000;71 Suppl 1: S29-35.

33. Giraldo PC, Rodrigues HM, Gosson AM, Amaral R, Eleutério J, Gonçalves AKS.
Vulvovaginitis and the treatment of asymptomatic partners: a systematic review and meta-
analysis. J Bras Doenças Sex Transm. 2014;26: 15–20.

34. Milne JD, Warnock DW. Effect of simultaneous oral and vaginal treatment on the rate of
cure and relapse in vaginal candidosis. Br J Vener Dis. 1979;55: 362–365.

35. El-Din SS, Reynolds MT, Ashbee HR, Barton RC, Evans EG. An investigation into the
pathogenesis of vulvo-vaginal candidosis. Sex Transm Infect. 2001;77: 179–183.

36. Fong IW. The rectal carriage of yeast in patients with vaginal candidiasis. Clin Invest Med.
1994;17: 426–431.

37. Sobel JD. Recurrent vulvovaginal candidiasis. N Engl J Med. 1986;315: 1455–1458.

38. Mårdh P-A, Novikova N, Stukalova E. Colonisation of extragenital sites by Candidain
women with recurrent vulvovaginal candiosis. BJOG. 2003;110: 934–937.

39. Spinillo A, Nicola S, Colonna L, Marangoni E, Cavanna C, Michelone G. Frequency and
significance of drug resistance in vulvovaginal candidiasis. Gynecol Obstet Invest. 1994;38:
130–133.

40. O'Connor MI, Sobel JD. Epidemiology of recurrent vulvovaginal candidiasis: identification
and strain differentiation of *Candida albicans*. J Infect Dis. 1986;154: 358–363.

41. Warnock DW, Speller DC, Day JK, Farrell AJ. Resistogram method for differentiation of
strains of *Candida albicans*. J Appl Bacteriol. 1979;46: 571–578.

42. Odds FC, Abbott AB. A simple system for the presumptive identification of *Candida
albicans* and differentiation of strains within the species. Sabouraudia. 1980;18: 301–317.

43. Stein GE, Sheridan VL, Magee BB, Magee PT. Use of rDNA restriction fragment length
polymorphisms to differentiate strains of *Candida albicans* in women with vulvovaginal
candidiasis. Diagn Microbiol Infect Dis. 1991;14: 459–464.

44. Lockhart SR, Reed BD, Pierson CL, Soll DR. Most frequent scenario for recurrent Candida

vaginitis is strain maintenance with "substrain shuffling": demonstration by sequential DNA
fingerprinting with probes Ca3, C1, and CARE2. J Clin Microbiol. 1996;34: 767–777.

45. Soll DR, Galask R, Isley S, Rao TV, Stone D, Hicks J, et al. Switching of *Candida albicans*
during successive episodes of recurrent vaginitis. J Clin Microbiol. 1989;27: 681–690.

46. Mendling W, Pinto De Andrade M, Gutschmidt J, Gantenberg R, Presber W, Schonian G.
Strain specificity of yeasts isolated from different locations of women suffering from vaginal
candidosis, and their partners Vergleich der Stammspezifitat von Hefepilzen verschiedener
Lokalisation bei Frauen mit Vaginalcandidosen. Mycoses. 2000;43: 387–392.

47. Vazquez JA, Sobel JD, Demitriou R, Vaishampayan J, Lynch M, Zervos MJ. Karyotyping of
Candida albicans isolates obtained longitudinally in women with recurrent vulvovaginal
candidiasis. J Infect Dis. 1994;170: 1566–1569.

48. Amouri I, Sellami H, Abbes S, Hadrich I, Mahfoudh N, Makni H, et al. Microsatellite analysis
of Candida isolates from recurrent vulvovaginal candidiasis. J Med Microbiol. 2012;61:
1091–1096.

49. Sampaio P, Gusmão L, Correia A, Alves C, Rodrigues AG, Pina-Vaz C, et al. New
microsatellite multiplex PCR for *Candida albicans* strain typing reveals microevolutionary
changes. J Clin Microbiol. 2005;43: 3869–3876.

50. Güzel AB, Döğen A, Aydın M, Serin A, Serin MS, Kalkancı A, et al. Genotyping reveals no
link between *Candida albicans* genotype and vaginitis severity in Turkish women.
Mycopathologia. 2013;175: 287–294.

51. Hong N, Lei Y, Chen H, Chen X, Tsui KM, Hu D, et al. Genotyping and drug resistance
profile of clinical isolates of *Candida albicans* from vulvovaginal candidiasis in the Eastern
China. Mycopathologia. 2022;187: 217–224.

52. Araújo Paulo de Medeiros M, Vieira de Melo AP, Gonçalves SS, Milan EP, Chaves GM.
Genetic relatedness among vaginal and anal isolates of *Candida albicans* from women with
vulvovaginal candidiasis in north-east Brazil. J Med Microbiol. 2014;63: 1436–1445.

53. Chong PP, Abdul Hadi SR, Lee YL, Phan CL, Tan BC, Ng KP, et al. Genotyping and drug
resistance profile of *Candida* spp. in recurrent and one-off vaginitis, and high association of
non-albicans species with non-pregnant status. Infect Genet Evol. 2007;7: 449–456.

54. Faria-Gonçalves P, Gaspar C, Oliveira AS, Palmeira-de-Oliveira R, Gonçalves T, Martinez-
de-Oliveira J, et al. Evaluation of overtime phenotypic variation of yeasts in chronic
vulvovaginal candidosis cases. Med Mycol. 2021;59: 1166–1173.

55. Tian J-Y, Yang Y-G, Chen S, Teng Y, Li X-Z. Genetic diversity and molecular epidemiology
of *Candida albicans* from vulvovaginal candidiasis patients. Infect Genet Evol. 2021;92:
104893.

56. Zhu Y, Fang C, Shi Y, Shan Y, Liu X, Liang Y, et al. *Candida albicans* multilocus sequence
typing clade I contributes to the clinical phenotype of vulvovaginal candidiasis patients.
Front Med. 2022;9: 837536.

57. Bradford LL, Chibucos MC, Ma B, Bruno V, Ravel J. Vaginal *Candida* spp. genomes from

women with vulvovaginal candidiasis. Pathog Dis. 2017;75: ftx061

58. Sitterlé E, Maufrais C, Sertour N, Palayret M, d'Enfert C, Bougnoux M-E. Within-host
genomic diversity of *Candida albicans* in healthy carriers. Sci Rep. 2019;9: 2563.

59. Badrane H, Cheng S, Dupont CL, Hao B, Driscoll E, Morder K, et al. Genotypic diversity
and unrecognized antifungal resistance among populations of *Candida glabrata* from
positive blood cultures. Nat Commun. 2023;14: 5918.

60. Willing E-M, Dreyer C, van Oosterhout C. Estimates of genetic differentiation measured by
F(ST) do not necessarily require large sample sizes when using many SNP markers. PLoS
One. 2012;7: e42649.

61. Jeffries DL, Copp GH, Lawson Handley L, Olsén KH, Sayer CD, Hänfling B. Comparing
RADseq and microsatellites to infer complex phylogeographic patterns, an empirical
perspective in the Crucian carp, Carassius carassius, L. Mol Ecol. 2016;25: 2997–3018.

62. Nazareno AG, Bemmels JB, Dick CW, Lohmann LG. Minimum sample sizes for population
genomics: an empirical study from an Amazonian plant species. Mol Ecol Resour. 2017;17:
1136–1147.

63. Qu W-M, Liang N, Wu Z-K, Zhao Y-G, Chu D. Minimum sample sizes for invasion
genomics: Empirical investigation in an invasive whitefly. Ecol Evol. 2020;10: 38–49.

64. Romeo O, Tietz H-J, Criseo G. *Candida africana*: Is it a fungal pathogen? Curr Fungal
Infect Rep. 2013;7: 192–197.

65. Mixão V, Saus E, Boekhout T, Gabaldón T. Extreme diversification driven by parallel events
of massive loss of heterozygosity in the hybrid lineage of *Candida albicans*. Genetics.
2021;217. doi:10.1093/genetics/iyaa004

66. Helmstetter N, Chybowska AD, Delaney C, Da Silva Dantas A, Gifford H, Wacker T, et al.
Population genetics and microevolution of clinical *Candida glabrata* reveals recombinant
sequence types and hyper-variation within mitochondrial genomes, virulence genes, and
drug targets. Genetics. 2022. doi:10.1093/genetics/iyac031

67. Carreté L, Ksiezopolska E, Pegueroles C, Gómez-Molero E, Saus E, Iraola-Guzmán S, et
al. Patterns of genomic variation in the opportunistic pathogen *Candida glabrata* suggest
the existence of mating and a secondary association with humans. Curr Biol. 2018;28: 15-
27.e7.

68. Stamatakis A. RAxML version 8: a tool for phylogenetic analysis and post-analysis of large
phylogenies. Bioinformatics. 2014;30: 1312–1313.

69. Matsutani M, Wakinaka T, Watanabe J, Tokuoka M, Ohnishi A. Comparative genomics of
closely related *Tetragenococcus halophilus* strains elucidate the diversity and
microevolution of CRISPR elements. Front Microbiol. 2021;12: 687985.

70. Abbey DA, Funt J, Lurie-Weinberger MN, Thompson DA, Regev A, Myers CL, et al. YMAP:
a pipeline for visualization of copy number variation and loss of heterozygosity in eukaryotic
pathogens. Genome Med. 2014;6: 100.

71. Pemberton TJ. Identification and comparative analysis of sixteen fungal peptidyl-prolyl

cis/trans isomerase repertoires. BMC Genomics. 2006;7: 244.

72. Salama OE, Gerstein AC. Differential response of species morphologies and isolates to
fluconazole and boric acid. Antimicrob Agents Chemother. 2022;66: e0240621.

73. Bhattacharjee P. Epidemiology and antifungal susceptibility of *Candida* species in a tertiary
care hospital, Kolkata, India. Curr Med Mycol. 2016;2: 20–27.

74. Hashemi SE, Shokohi T, Abastabar M, Aslani N, Ghadamzadeh M, Haghani I. Species
distribution and susceptibility profiles of *Candida* species isolated from vulvovaginal
candidiasis, emergence of C. lusitaniae. Curr Med Mycol. 2019;5: 26–34.

75. Sampaio P, Gusmão L, Alves C, Pina-Vaz C, Amorim A, Pais C. Highly polymorphic
microsatellite for identification of *Candida albicans* strains. J Clin Microbiol. 2003;41: 552–
557.

76. Shi W-M, Mei X-Y, Gao F, Huo K-K, Shen L-L, Qin H-H, et al. Analysis of genital *Candida*
albicans infection by rapid microsatellite markers genotyping. Chin Med J . 2007;120: 975.

77. Araújo Paulo de Medeiros M, Vieira de Melo AP, Maia de Sousa AM, Silva-Rocha WP,
Pipolo Milan E, Maranhão Chaves G. Characterization of virulence factors of vaginal and
anal isolates of *Candida albicans* sequentially obtained from patients with vulvovaginal
candidiasis in north-east Brazil. J Mycol Med. 2017;27: 567–572.

78. Spinillo A, Carratta L, Pizzoli G, Lombardi G, Cavanna C, Michelone G, et al. Recurrent
vaginal candidiasis. Results of a cohort study of sexual transmission and intestinal
reservoir. J Reprod Med. 1992;37: 343–347.

79. Lott TJ, Frade JP, Lyon GM, Iqbal N, Lockhart SR. Bloodstream and non-invasive isolates
of *Candida glabrata* have similar population structures and fluconazole susceptibilities. Med
Mycol. 2012;50: 136–142.

80. Lockhart SR, Fritch JJ, Meier AS, Schröppel K, Srikantha T, Galask R, et al. Colonizing
populations of *Candida albicans* are clonal in origin but undergo microevolution through C1
fragment reorganization as demonstrated by DNA fingerprinting and C1 sequencing. J Clin
Microbiol. 1995;33: 1501–1509.

81. Korunes KL, Samuk K. pixy: Unbiased estimation of nucleotide diversity and divergence in
the presence of missing data. Mol Ecol Resour. 2021;21: 1359–1368.

82. Dutta A, Lin G, Pankajam AV, Chakraborty P, Bhat N, Steinmetz LM, et al. Genome
Dynamics of hybrid *Saccharomyces cerevisiae* during vegetative and meiotic divisions. G3 .
2017;7: 3669–3679.

83. Dutta A, Dutreux F, Schacherer J. Loss of heterozygosity results in rapid but variable
genome homogenization across yeast genetic backgrounds. Elife. 2021;10: e70339

84. Sui Y, Qi L, Wu J-K, Wen X-P, Tang X-X, Ma Z-J, et al. Genome-wide mapping of
spontaneous genetic alterations in diploid yeast cells. Proc Natl Acad Sci U S A. 2020;117:
28191–28200.

85. Sharp NP, Sandell L, James CG, Otto SP. The genome-wide rate and spectrum of
spontaneous mutations differ between haploid and diploid yeast. Proc Natl Acad Sci U S A.

2018;115: E5046–E5055.

86. Zhu YO, Siegal ML, Hall DW, Petrov DA. Precise estimates of mutation rate and spectrum
in yeast. Proc Natl Acad Sci U S A. 2014;111: E2310-8.

87. Smukowski Heil C. Loss of heterozygosity and its importance in evolution. J Mol Evol.
2023;91: 369–377.

88. Lynch M, Sung W, Morris K, Coffey N, Landry CR, Dopman EB, et al. A genome-wide view
of the spectrum of spontaneous mutations in yeast. Proc Natl Acad Sci U S A. 2008;105:
9272–9277.

89. Selmecki A, Gerami-Nejad M, Paulson C, Forche A, Berman J. An isochromosome confers
drug resistance *in vivo* by amplification of two genes, ERG11 and TAC1. Mol Microbiol.
2008;68: 624–641.

90. Todd RT, Selmecki A. Expandable and reversible copy number amplification drives rapid
adaptation to antifungal drugs. Elife. 2020;9: 58349.

91. Kukurudz RJ, Chapel M, Wonitowy Q, Adamu Bukari A-R, Sidney B, Sierhuis R, et al.
Acquisition of cross-azole tolerance and aneuploidy in *Candida albicans* strains evolved to
posaconazole. G3 . 2022;12: jkac156.

92. Yang F, Scopel EFC, Li H, Sun L-L, Kawar N, Cao Y-B, et al. Antifungal Tolerance and
Resistance Emerge at Distinct Drug Concentrations and Rely upon Different Aneuploid
Chromosomes. MBio. 2023;14: e0022723.

93. Ford CB, Funt JM, Abbey D, Issi L, Guiducci C, Martinez DA, et al. The evolution of drug
resistance in clinical isolates of *Candida albicans*. eLife. 2015;4: e00662.

94. Diogo D, Bouchier C, d'Enfert C, Bougnoux M-E. Loss of heterozygosity in commensal
isolates of the asexual diploid yeast *Candida albicans*. Fungal Genet Biol. 2009;46: 159–
168.

95. Smith AC, Morran LT, Hickman MA. Host defense mechanisms induce genome instability
leading to rapid evolution in an opportunistic fungal pathogen. Infect Immun. 2022;90:
e0032821.

96. James TY, Michelotti LA, Glasco AD, Clemons RA, Powers RA, James ES, et al.
Adaptation by loss of heterozygosity in *Saccharomyces cerevisiae* clones under divergent
selection. Genetics. 2019;213: 665–683.

97. Ene IV, Farrer RA, Hirakawa MP, Agwamba K, Cuomo CA, Bennett RJ. Global analysis of
mutations driving microevolution of a heterozygous diploid fungal pathogen. Proc Natl Acad
Sci U S A. 2018;115: E8688–E8697.

98. Pappas PG, Kauffman CA, Andes D, Benjamin DK Jr, Calandra TF, Edwards JE Jr, et al.
Clinical practice guidelines for the management of candidiasis: 2009 update by the
Infectious Diseases Society of America. Clin Infect Dis. 2009;48: 503–535.

99. Odds FC, Bougnoux M-E, Shaw DJ, Bain JM, Davidson AD, Diogo D, et al. Molecular
phylogenetics of *Candida albicans*. Eukaryot Cell. 2007;6: 1041–1052.

100. Ge S-H, Xie J, Xu J, Li J, Li D-M, Zong L-L, et al. Prevalence of specific and
phylogenetically closely related genotypes in the population of *Candida albicans* associated
with genital candidiasis in China. Fungal Genet Biol. 2012;49: 86–93.

101. Schmid J, Herd S, Hunter PR, Cannon RD, Yasin MSM, Samad S, et al. Evidence for a
general-purpose genotype in *Candida albicans*, highly prevalent in multiple geographical
regions, patient types and types of infection. Microbiology. 1999;145 (Pt 9): 2405–2413.

102. Giblin L, Edelmann A, Zhang N, von Maltzahn NB, Cleland SB, Sullivan PA, et al. A DNA
polymorphism specific to *Candida albicans* strains exceptionally successful as human
pathogens. Gene. 2001;272: 157–164.

103. Cravener MV, Do E, May G, Zarnowski R, Andes DR, McManus CJ, et al. Reinforcement
amid genetic diversity in the *Candida albicans* biofilm regulatory network. PLoS Pathog.
2023;19: e1011109.

104. MacCallum DM, Castillo L, Nather K, Munro CA, Brown AJP, Gow NAR, et al. Property
differences among the four major *Candida albicans* strain clades. Eukaryot Cell. 2009;8:
373–387.

105. Sala A, Ardizzoni A, Spaggiari L, Vaidya N, van der Schaaf J, Rizzato C, et al. A new
phenotype in *Candida*-epithelial cell interaction distinguishes colonization- versus
vulvovaginal candidiasis-associated strains. MBio. 2023; e0010723.

106. Makanjuola O, Bongomin F, Fayemiwo SA. An update on the roles of non-albicans
Candida species in vulvovaginitis. J Fungi (Basel). 2018;4: 121.

107. Dodgson AR, Pujol C, Denning DW, Soll DR, Fox AJ. Multilocus sequence typing of
Candida glabrata reveals geographically enriched clades. J Clin Microbiol. 2003;41: 5709–
5717.

108. Chen P-Y, Huang Y-S, Chuang Y-C, Wang J-T, Sheng W-H, Chen Y-C, et al. Implication of
genotypes for prognosis of *Candida glabrata* bloodstream infections. J Antimicrob
Chemother. 2024;22: dkae200.

109. Won EJ, Choi MJ, Kim M-N, Yong D, Lee WG, Uh Y, et al. Fluconazole-resistant *Candida*
glabrata bloodstream isolates, South Korea, 2008-2018. Emerg Infect Dis. 2021;27: 779–
788.

110. Khalifa HO, Arai T, Majima H, Watanabe A, Kamei K. Genetic basis of azole and
echinocandin resistance in clinical *Candida glabrata* in Japan. Antimicrob Agents
Chemother. 2020;64: 10-128.

111. Chew KL, Achik R, Osman NH, Octavia S, Teo JWP. Genomic epidemiology of human
candidaemia isolates in a tertiary hospital. Microb Genom. 2023;9: 001047.

112. de Hoog S, Walsh TJ, Ahmed SA, Alastruey-Izquierdo A, Alexander BD, Arendrup MC, et
al. A conceptual framework for nomenclatural stability and validity of medically important
fungi: a proposed global consensus guideline for fungal name changes supported by ABP,
ASM, CLSI, ECMM, ESCMID-EFISG, EUCAST-AFST, FDLC, IDSA, ISHAM, MMSA, and
MSGERC. J Clin Microbiol. 2023;61: e0087323.

113. Gabaldón T, Carreté L. The birth of a deadly yeast: tracing the evolutionary emergence of
virulence traits in *Candida glabrata*. FEMS Yeast Res. 2016;16: fov110.

114. Galocha M, Pais P, Cavalheiro M, Pereira D, Viana R, Teixeira MC. Divergent Approaches
to Virulence in *C. albicans* and *C. glabrata*: Two Sides of the Same Coin. Int J Mol Sci.
2019;20: 2345.

115. Brunke S, Hube B. Two unlike cousins: *Candida albicans* and *C. glabrata* infection
strategies. Cell Microbiol. 2013;15: 701–708.

116. Thorvaldsen JL, Sewell AK, McCowen CL, Winge DR. Regulation of metallothionein genes
by the ACE1 and AMT1 transcription factors. J Biol Chem. 1993;268: 12512–12518.

117. Weissman Z, Berdicevsky I, Cavari BZ, Kornitzer D. The high copper tolerance of *Candida*
albicans is mediated by a P-type ATPase. Proc Natl Acad Sci U S A. 2000;97: 3520–3525.

118. Riedelberger M, Penninger P, Tscherner M, Hadriga B, Brunnhofer C, Jenull S, et al. Type
I interferons ameliorate zinc intoxication of *Candida glabrata* by macrophages and promote
fungal immune evasion. iScience. 2020;23: 101121.

119. Sayers EW, Bolton EE, Brister JR, Canese K, Chan J, Comeau DC, et al. Database
resources of the national center for biotechnology information. Nucleic Acids Res. 2022;50:
D20–D26.

120. Bolger AM, Lohse M, Usadel B. Trimmomatic: a flexible trimmer for Illumina sequence
data. Bioinformatics. 2014;30: 2114–2120.

121. Todd RT, Wikoff TD, Forche A, Selmecki A. Genome plasticity in *Candida albicans* is
driven by long repeat sequences. Elife. 2019;8: e45954.

122. Ewels P, Magnusson M, Lundin S, Käller M. MultiQC: summarize analysis results for
multiple tools and samples in a single report. Bioinformatics. 2016;32: 3047–3048.

123. Li H. Aligning sequence reads, clone sequences and assembly contigs with BWA-MEM.
arXiv [q-bio.GN]. 2013. Available: http://arxiv.org/abs/1303.3997

124. Skrzypek MS, Binkley J, Binkley G, Miyasato SR, Simison M, Sherlock G. The Candida
Genome Database (CGD): incorporation of Assembly 22, systematic identifiers and
visualization of high throughput sequencing data. Nucleic Acids Res. 2017;45: D592–D596.

125. Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, et al. The sequence
alignment/map format and SAMtools. Bioinformatics. 2009;25: 2078–2079.

126. Yates AD, Allen J, Amode RM, Azov AG, Barba M, Becerra A, et al. Ensembl genomes
2022: an expanding genome resource for non-vertebrates. Nucleic Acids Res. 2022;50:
D996–D1003.

127. Jones T, Federspiel NA, Chibana H, Dungan J, Kalman S, Magee BB, et al. The diploid
genome sequence of *Candida albicans*. Proc Natl Acad Sci U S A. 2004;101: 7329–7334.

128. DePristo MA, Banks E, Poplin R, Garimella KV, Maguire JR, Hartl C, et al. A framework for
variation discovery and genotyping using next-generation DNA sequencing data. Nat
Genet. 2011;43: 491–498.

129. Van der Auwera GA, Carneiro MO, Hartl C, Poplin R, Del Angel G, Levy-Moonshine A, et
al. From FastQ data to high confidence variant calls: the Genome Analysis Toolkit best
practices pipeline. Curr Protoc Bioinformatics. 2013;43: 11.10.1-11.10.33.

130. Poplin R, Ruano-Rubio V, DePristo MA, Fennell TJ, Carneiro MO, Van der Auwera GA, et
al. Scaling accurate genetic variant discovery to tens of thousands of samples. bioRxiv.
2018. p. 201178. doi:10.1101/201178

131. Ortiz EM. vcf2phylip v2.0: convert a VCF matrix into several matrix formats for
phylogenetic analysis. 2019. doi:10.5281/zenodo.2540861

132. Price MN, Dehal PS, Arkin AP. FastTree 2--approximately maximum-likelihood trees for
large alignments. PLoS One. 2010;5: e9490.

133. Liu K, Linder CR, Warnow T. RAxML and FastTree: comparing two methods for large-scale
maximum likelihood phylogeny estimation. PLoS One. 2011;6: e27731.

134. Letunic I, Bork P. Interactive Tree Of Life (iTOL) v5: an online tool for phylogenetic tree
display and annotation. Nucleic Acids Res. 2021;49: W293–W296.

135. Mixão V, Gabaldón T. Genomic evidence for a hybrid origin of the yeast opportunistic
pathogen Candida albicans. BMC Biol. 2020;18: 48.

136. Szarvas J, Rebelo AR, Bortolaia V, Leekitcharoenphon P, Schrøder Hansen D, Nielsen
HL, et al. Danish whole-genome-sequenced *Candida albicans* and *Candida glabrata*
samples fit into globally prevalent clades. J Fungi (Basel). 2021;7. doi:10.3390/jof7110962

137. Cleary JG, Braithwaite R, Gaastra K, Hilbush BS, Inglis S, Irvine SA, et al. Comparing
variant call files for performance benchmarking of next-generation sequencing variant
calling pipelines. bioRxiv. 2015. p. 023754. doi:10.1101/023754

138. Gupta A, Jordan IK, Rishishwar L. stringMLST: a fast k-mer based tool for multilocus
sequence typing. Bioinformatics. 2017;33: 119–121.

139. Jolley KA, Bray JE, Maiden MCJ. Open-access bacterial population genomics: BIGSdb
software, the Pubmlst.org website and their applications. Wellcome Open Res. 2018;3:
124.

140. Li H, Ralph P. Local PCA shows how the effect of population structure differs along the
genome. Genetics. 2019;211: 289–304.

141. Danecek P, Auton A, Abecasis G, Albers CA, Banks E, DePristo MA, et al. The variant call
format and VCFtools. Bioinformatics. 2011;27: 2156–2158.

142. Millard SP. EnvStats: An R Package for environmental statistics. Springer New York; 2013.

143. NCCLS. Reference method for antifungal disk diffusion susceptibility testing of yeasts;
approved guideline. NCCLS document M44-A National Committee for Clinical Laboratory
Standards. Wayne.

144. Gerstein AC, Rosenberg A, Hecht I, Berman J. diskImageR: quantification of resistance
and tolerance to antimicrobial drugs using disk diffusion assays. Microbiology. 2016;162:
1059–1068.

145. Zupan J, Raspor P. Quantitative agar-invasion assay. J Microbiol Methods. 2008;73: 100–
104.

Figures

1148

A *N. glabratus*

Tree scale: 0.1 ├───────┤

Isolation Site
- Rectum / Stool
- Vagina
- Other

B *C. albicans*

Tree scale: 0.1 ├──┤

Fig. 1 Approximate maximum likelihood phylogenies of (A) *N. glabratus* (n = 526 isolates and

eight THRIVE-yeast isolates) and (B) *C. albicans* (n = 182 isolates and 24 THRIVE-yeast

isolates). THRIVE-yeast isolates are indicated with red labels. The YST6 *N. glabratus* isolates

fall in a cluster of predominantly ST16 isolates (ST16 shown by blue branches, black indicates

not ST16). TVY4, TVY10, and YST7 *C. albicans* isolates fall in clade 1 (blue branches). The

physiological sites of isolation are indicated as a colour strip around the phylogeny, indicating

vaginal (purple), rectal (orange), and all other sites (grey). The *N. glabratus* phylogeny was

rooted at the midpoint, and the *C. albicans* tree was rooted by clade 13 (grey labels).

1158

1159

Fig. 2 Within participant phylogenetic and single nucleotide position (SNP) analyses. (A) The

fine-scale phylogenetic structure among THRIVE-yeast isolates. Vaginal (V) and rectal (R)

isolates were acquired during symptomatic infection from four participants with a history of

RVVC (YST6: *N. glabratus*; TVY4, TVY10, YST7: *C. albicans*). Reads were aligned to their

respective reference genomes (SC5314 and CBS138), and variants were called using a custom

variant calling pipeline adopted from the GATK best practices. Mitochondrial variants, as well as

variants in repeat regions, were excluded. Phylogenetic trees were generated with RAxML;

black circles indicate branches with bootstrap support ≥ 0.8. (B) Within-participate pairwise

comparison of single nucleotide positions between isolates from the indicated sites (V-V:

vaginal-vaginal, R-V: rectal-vaginal, R-R: rectal-rectal).

1170

1171

Fig. 3 Relationship between sample size and average nucleotide diversity. π was calculated for different numbers of samples from each individual. For n= 2, 3, 4, 6, or 10, the given number of samples was randomly selected from the vaginal isolate set. For n = 12, the datasets were generated by randomly choosing 6 samples from each of the rectal and vaginal isolates sets. The bootstrap analysis was done 50 times for each sample size, and the mean and standard deviation among data sets were calculated. For n = 23 (*C. albicans*, TVY10, TVY4, YST7) or 24 (*N. glabratus,* YST6), the nucleotide diversity of all samples was calculated.

1182
1183
Fig. 4 Comparison of the average nucleotide diversity between our samples and isolates from

other studies. (A) Comparison of pi in THRIVE *C. albicans* isolates from TVY4, TVY10, and

YST7 to commensal isolates from two previously published studies. For accurate comparison,

three randomly chosen samples were subsampled from each site in the THRIVE-yeast isolates.

(B) Comparison of pi in THRIVE *N. glabratus* (YST6) isolates to bloodstream infection isolates

from 10 individuals in a previous study.

1190

1191

Fig. 5 CNV and LOH profiles of THRIVE-yeast. (A) Representative trace of a YST6 isolate (*N.*

glabratus) and (B) TVY10 isolates, representative traces of TVY4 and YST7 isolates, and the

four isolates that are closely related to YST7 (three vaginal isolates and Niger8, oral isolate).

The relative copy number compared to the reference genomes is shown as the horizontal black

line with the scale on the y-axis. For *C. albicans* isolates, the density of heterozygous SNPs is

shown as the vertical lines spanning the height of each chromosome box, with the intensity

representing the number of SNPs in 5 kb bins. Heterozygous SNPs are gray, homozygous

SNPs are coloured based on the retained SC5314 homolog: cyan for "AA" and magenta for

"BB." White indicates an ancestral LOH in SC5314. For each chromosome, the centromere is

indicated by an indentation in the box. The dots on the bottom line below each box indicate the

positions of major repeat sequences. (C) Fine-scale coverage mapping of the putative CNV on

chr3. Shown is one representative trace from YST7 R2, all isolates have a similar pattern. Gene

positions above the figure are approximated.

1205

1206

1207

1208

1209

1210

1211

1212

1213

1214

1215

1216

1217

1218

1219 **Fig. 6** Growth rate was measured from 12 vaginal and 12 rectal isolates from each population.

1220 Optical density was recorded every 15 minutes in a plate reader with constant shaking and

1221 incubation at 37 ℃. Each point represents the mean of two technical replicates for each of the

1222 two biological isolates, 24 isolates were measured for each group. The growth rate was

1223 calculated as the spline with the highest slope using a custom R script.

1224

Fig. 7 Drug response phenotypes from disk diffusion assays. Drug resistance (top panels) and
drug tolerance (bottom panels) were measured from disk diffusion assays for fluconazole (FLC)
and boric acid (BA). Drug response was measured on pH 4 Mueller-Hinton plates using the R
package *diskImageR* [144], which computationally measures response parameters from images
of disk diffusion assays. Each point represents the mean of four replicates (two technical
replicates for two biological replicates).

1231

1232

1233   **Fig. 8** Invasive growth was qualitatively scored after growth on YPD plates for 96 h. Each point

1234   indicates the maximum score between two bio-replicates for each isolate.

1235