Title

Executioner caspase is proximal to Fasciclin 3 which facilitates non-lethal activation in

Drosophila olfactory receptor neurons

Short title

Fasciclin 3 non-lethally activates caspase in *Drosophila* olfactory neurons

Authors

Masaya Muramoto[a,1], Nozomi Hanawa[a,1], Misako Okumura[b,c], Takahiro Chihara[b,c], Masayuki

Miura[a,2], and Natsuki Shinoda[a,2]

Affiliations & Footnotes

[a]Department of Genetics, Graduate School of Pharmaceutical Sciences, University of Tokyo,

7-3-1 Hongo, Bunkyo-ku, Tokyo 113-0033, Japan

[b]Program of Biomedical Science, Graduate School of Integrated Sciences for Life, Hiroshima

University, Higashi-Hiroshima, Hiroshima 739-8526, Japan

[c]Program of Basic Biology, Graduate School of Integrated Sciences for Life, Hiroshima

University, Higashi-Hiroshima, Hiroshima 739-8526, Japan

[1]Equal contributions

[2]Corresponding authors

Corresponding Authors

Masayuki Miura: Department of Genetics, Graduate School of Pharmaceutical Sciences,

University of Tokyo, 7-3-1 Hongo, Bunkyo-ku, Tokyo 113-0033, Japan. Telephone: +81-3-

5841-4860. E-mail: miura@mol.f.u-tokyo.ac.jp

Natsuki Shinoda: Department of Genetics, Graduate School of Pharmaceutical Sciences,

University of Tokyo 7-3-1 Hongo, Bunkyo-ku, Tokyo 113-0033, Japan. Telephone: +81-3-

5841-4867. E-mail: f-shinoda.natsuki@g.ecc.u-tokyo.ac.jp

Author Contributions

Masaya Muramoto: Formal Analysis, Investigation, Methodology, Resources, Writing –

Review & Editing; Nozomi Hanawa: Formal analysis, Investigation, Methodology, Resources,

Writing – Review & Editing; Misako Okumura: Methodology, Writing – Review & Editing;

Takahiro Chihara: Methodology, Writing – Review & Editing; Masayuki Miura:

Conceptualization, Funding Acquisition, Supervision, Writing – Original Draft Preparation;

Natsuki Shinoda: Conceptualization, Formal Analysis, Funding Acquisition, Investigation,

Project Administration, Supervision, Validation, Visualization, Writing – Original Draft

Preparation

Conflict of Interest

41 The authors declare that they have no conflict of interest.

Abstract

Executioner caspases are activated by initiator caspases, resulting in lethal and non-lethal cellular outcomes. Despite the identification of numerous caspase-dependent non-lethal cellular processes (CDPs), the molecular mechanism that enables precise regulation of the non-lethal functions of executioner caspases remains unclear. Here, we conducted a comprehensive analysis of endogenous proximal proteins of *Drosophila* executioner caspases *in vivo* using TurboID. We discovered that in the adult brain, the executioner caspase Drice is, as an inactive proform, primarily proximal to cell membrane proteins, including the cell adhesion molecule Fasciclin 3 (Fas3). Notably, Drice is proximal to the specific alternative splicing isoforms of Fas3, Fas3G. To investigate whether Fas3G modulates caspase activity, we developed a Gal4-Manipulated Area-Specific CaspaseTracker/CasExpress (MASCaT) system for highly sensitive monitoring of non-lethal caspase activity near cell membrane compartments. Using MASCaT, we demonstrated that *Fas3G* overexpression promotes caspase activation near cell membrane compartments in olfactory receptor neurons without killing them. Mechanistically, *Fas3G* overexpression induces the expression of initiator caspase Dronc, which also comes close to Fas3G, regulating non-lethal caspase activation. Physiologically, *Fas3G* overexpression-facilitated caspase activation regulates innate olfactory attraction behavior. Collectively, our findings suggest that caspase activation is subcellularly restricted, the platform of which is defined by caspase proximal proteins, for non-lethal functions.

Author Summary

Caspases are Cysteine-ASPartic acid proteASEs which are primarily involved in cell death. In addition, caspases play versatile non-lethal (non-cell death) roles in a wide range of cell types including neurons. However, the molecular mechanism that enables the precise regulation of non-lethal activation and functions remains unclear. Here, we focused on the proximal proteins of caspases as potential regulators of non-lethal activation. By comprehensively analyzing the proximal proteins of caspases in the adult *Drosophila* brain using TurboID, we discovered that caspases are primarily proximal to membrane proteins rather than to cytosolic proteins. By developing a genetic tool that enables highly sensitive monitoring of non-lethal caspase activity near membrane compartments, we found that proximal proteins facilitate non-lethal caspase activation in olfactory receptor neurons. Importantly, non-lethal caspase activation in olfactory receptor neurons affects organismal innate attraction behavior. Our research suggests that a caspase activation is compartmentalized, the platform of which is defined by caspase proximal proteins, for non-lethal functions.

Introduction

Caspases are cysteine aspartic acid proteases involved in cell death. Apoptotic stimuli activate initiator caspases that trigger a proteolytic cascade, leading to the activation of executioner caspases and culminating in apoptosis (Fig. 1A) [1]. Caspases exhibit diverse non-lethal cellular functions [2]. In the nervous system, non-lethal caspase activity promotes the maturation of olfactory sensory neurons during development [3]. Furthermore, non-lethal caspase activity induces long-term depression through AKT cleavage [4]. These are termed caspase-dependent non-lethal cellular processes (CDPs) [5]. Despite the discovery of numerous CDPs over the past two decades, the molecular mechanism that underlies specific non-lethal functions without causing cell death remains unclear.

Proteins that interact with initiator caspases are among the key regulators of CDPs. Tango7, the *Drosophila* translation initiation factor eIF3 subunit m, acts as an adapter and regulates the localization of the initiator caspase Dronc. During sperm individualization, Tango7 directs Dronc to the individualization complex [6]. In salivary glands, Tango7 directs Dronc to the plasma membrane [7]. In both cases, caspase activity promotes cellular remodeling without inducing cell death [6,7]. CRINKLED, a *Drosophila* unconventional myosin, functions as an adaptor of Dronc and facilitates the cleavage of Shaggy46 [8]. Shaggy46, activated by caspase-mediated cleavage, suppresses the emergence of ectopic sensory organ precursors [9]. During apoptosis-induced proliferation (AiP), Myo1D, another *Drosophila* unconventional myosin, facilitates Dronc translocation to the basal side of the plasma membrane of imaginal discs to promote AiP [10]. Thus, initiator caspase-interacting proteins regulate caspase activity in CDPs. However, given the importance of executioner caspases, which cleave a broad range of substrates and are essential for apoptosis [11,12], little is known about the subcellular compartmentalization and interacting proteins of those involved in CDPs. This lack of

understanding is partly due to the absence of protein-protein interaction domains (e.g., DED

and CARD) in the N-terminus of executioner caspases [13].

To understand the molecular mechanism regulating caspase activity involved in CDPs,

we focused on the proximal proteins of executioner caspases using TurboID, a proximity

labeling technique that analyzes protein proximity *in vivo* [14,15]. We previously established

C-terminal TurboID knocked-in caspase fly lines for Dronc, Drice, and Dcp-1 [16]. Importantly,

we have demonstrated that one of the preferred substrates of Drice, BubR1, is proximal to Drice

in vivo [17], suggesting that substrate preference is exploited by protein proximity. Therefore,

the proximity of executioner caspases to proteins is emerging as a means of regulating specific

non-lethal functions.

In this study, through a comprehensive analysis of proximal proteins, we discovered

that the cell adhesion molecule Fasciclin 3 (Fas3) is in close proximity to the executioner

caspase Drice, as an inactive proform, in the adult brain. We observed that Drice was

exclusively proximal to specific alternative splicing isoforms of Fas3, Fas3G. To ascertain

whether Fas3G modulates caspase activity, we developed a Gal4-Manipulated Area Specific

CaspaseTracker/CasExpress (MASCaT) system, which permits the monitoring of caspase

activity with high sensitivity and simultaneous genetic manipulation in the cells of interest.

Using MASCaT, we demonstrated that *Fas3G* overexpression enhances caspase activation

without killing cells through the induction of the expression of initiator caspase Dronc, which

comes proximal to Fas3G. Subsequently, we showed that *Fas3G* overexpression-facilitated

activation regulates innate olfactory attraction behavior. Collectively, our findings suggest that

caspase activation is subcellularly restricted, the platform of which is defined by caspase

proximal proteins, for non-lethal functions.

Results

Drice is the major executioner caspase expressed in the adult brains

In *Drosophila*, the caspase-mediated apoptotic signaling pathway is highly conserved [2]. The

activity of executioner caspases, specifically Drice and Dcp-1, is regulated by apoptotic

signaling pathways, including the initiator caspase Dronc (Fig. 1A). To investigate the

expression patterns of *Drosophila* caspases, we employed *Caspase::V5::TurboID* knocked-in

fly lines [16]. We found low levels of expression of the initiator caspase Dronc in adult heads

(Fig. 1B). Conversely, we observed high expression of the executioner caspase Drice, whereas

Dcp-1 was expressed at lower levels in adult heads (Fig. 1B). The proximal proteins of each

caspase were labeled with TurboID in adult heads by administration of 100 µM biotin (Fig. 1B).

Furthermore, we investigated the expression patterns of caspases using histochemical analysis

by examining streptavidin (SA) signal. Consistent with the western blotting findings, Drice

showed higher expression levels than Dronc and Dcp-1, which were expressed at lower levels

(Fig. 1C). We found that Drice was expressed in distinct regions, including the mushroom

bodies (MBs), suboesophageal ganglions (SGs), optic lobes (OLs), and antennal lobes (ALs),

which showed the highest levels of expression (Fig. 1C). The *Drosophila* ALs comprise

synaptic contacts between the axons of olfactory receptor neurons (ORNs), the dendrites of

projection neurons (PNs), and local interneuron (LNs) processes. Neuron-type-specific Gal4s

were used to knockdown *Drice* and the expression patterns were subsequently examined. We

found that Drice was expressed predominantly in ORNs in the AL (Fig. 1D). These results

showcase that Drice is not expressed in the entire brain, but rather in a specific subset of neurons

in the adult brains. Comparisons of the expression patterns of Drice were consistent across both

males and females (Fig. S1A), indicating an absence of sex-based differences, which justifies

the use of males for functional simplicity in this study.

Drice is proximal to cell membrane proteins rather than cytosolic proteins in the adult brains

To comprehensively identify proximal proteins of Drice, we employed a proximity labeling

approach. Specifically, we used *Drice::V5::TurboID* knocked-in flies fed 100 µM biotin. We

confirmed biotinylation of Drice proximal proteins in the adult brains (Fig. 1E). After purifying

the biotinylated proteins using NeutrAvidin magnetic beads, we performed LC-MS/MS

analysis of the samples (Fig. 1F). We identified 643 proteins (Fig. 1G, Table S1), of which 158

were highly specific to *Drice::V5::TurboID* flies compared to wild-type flies

(*Drice::V5::TurboID*/*w^1118* > 10; Fig. 1G, Table S1). Gene ontology analysis revealed that

Drice proximal proteins were enriched in the plasma membrane, synaptic vesicles, postsynaptic

membranes, and adherens junctions (Fig. 1H, Table S2). This finding suggests that Drice is

localized primarily in cell membrane compartments rather than in the cytoplasm of the adult

brain. Additionally, because the SA signal of *Drice::V5::TurboID* is mainly observed in the

AL where the axons of ORNs project (Fig. 1C), the enrichment of cell membrane proteins in

the proximal proteins of Drice is reasonable.

A specific splicing isoform of Fasciclin 3 is in proximity to Drice

During our analysis of proteins proximal to Drice, we discovered that the expression patterns

of Fasciclin 3 (Fas3) in the adult brain closely resembled those of Drice, especially within the

OLs, SGs, and ALs (Fig. 2A, B). Thus, we focused on Fas3 [18], a transmembrane protein

containing immunoglobulin-like domains that regulates synaptic target recognition and axon

fasciculation [19]. *Fas3* is encoded by seven splicing isoforms, resulting in the production of

five protein isoforms that share extracellular domains but differ in intracellular domains with

low-complexity sequences (Fig. 2C, D). Only one specific protein isoform, Fas3 isoform G

(Fas3G), was identified in the LC-MS/MS analysis as a Drice proximal protein in the adult

brain (Fig. 1G, Table S1). To validate this result, we generated an antibody specific to Fas3G

(Fig. 2C, E). We confirmed that only the Fas3G was labeled with Drice::V5::TurboID and

purified using NeutrAvidin magnetic beads as detected using western blotting (Fig. 2F). To

further confirm the isoform specificity to Drice proximity, we overexpressed 3xFLAG-tagged

Fas3 isoforms A, *B/F*, *C*, *D/E*, and *G* in olfactory receptor neurons using *Pebbled-Gal4* in a

Drice::V5::TurboID knocked-in background. After proximity labeling with TurboID, followed

by NeutrAvidin purification, we found that the Fas3G was highly enriched compared to the

other isoforms (Fig. 2G). These findings suggested that Drice is proximal to a specific isoform

of Fas3 in olfactory receptor neurons, regardless of its protein expression level. We also

investigated whether Drice and Fas3G physically interact by co-immunoprecipitation.

Although Fas3G was labelled by Drice::V5::TurboID, indicating its proximity to Drice within

the ORNs of adult brains, Drice did not co-immunoprecipitate with Fas3G (Fig. S2A). This

suggests that while Drice and Fas3G are proximate, they do not physically interact, highlighting

the utility of proximity labeling as a superior technique to conventional co-immunoprecipitation

for identifying proteins that are spatially close.

Fasciclin 3s are not substrates of caspase

It is often observed that proteins in close proximity to caspases are their preferred substrates

[17]. To investigate this, we examined whether any of the Fas3 isoforms are substrates for

caspase by expressing all isoforms in *Drosophila* S2 cells. However, we found that none of the

Fas3 isoforms were cleaved by caspase following the induction of apoptosis (Fig. S2B).

Therefore, Fas3s are not substrates of caspase.

MASCaT: a cell type-specific highly sensitive non-lethal caspase activity reporter

We investigated whether Fas3G regulates caspase activity in lethal or non-lethal processes.

CaspaseTracker/CasExpress is a highly sensitive reporter of non-lethal caspase activity that

uses caspase-activatable Gal4 tethered to the membrane by mCD8 [20,21]. However, reliance

on Gal4 to report their activity presents a challenge in reconciling Gal4/UAS-dependent gene

manipulations. To overcome this limitation, we created a new reporter to capture non-lethal

caspase activation independent of caspase-activatable Gal4. We replaced the caspase-

activatable Gal4 with the caspase-activatable QF2 (*mCD8::DQVD::QF2*), where DQVD is a

caspase cleavage sequence, and first expressed directly downstream of *UAS* sequence (Fig.

S3A). However, the caspase-activatable QF2 induced mNeonGreen expression under *QUAS*

sequence even either with a pan-caspase inhibitor, zVAD-fmk, or an uncleavable DQVA

mutant probe (*mCD8::DQVA::QF2*, insensitive to caspase-mediated cleavage which serves as

a negative control) (Fig. S3B), potentially because the Gal4/UAS binary system expressed

excessive mCD8::DQVD/A::QF2 probes that results in caspase activation-independent

induction of QF2/QUAS system. Thus, to reduce the amount of probes expression, we

generated *Ubi-FRT-STOP-FRT-mCD8::DQVD::QF2*, simultaneously expressing *Flippase*

(*FLP*) under *UAS* to restrict reporter expression to the cells of interest (Fig. 3A). *Drosophila* S2

cells undergo weak caspase activation only by transfection [17]. Taking advantage of that, this

system reports caspase activity through the caspase-activatable QF2/QUAS system, expressing

mNeonGreen, which is not detected by caspase-non-activatable QF2 (Fig. 3B). We named this

novel reporter a Gal4-Manipulated Area Specific CaspaseTracker/CasExpress (MASCaT; Fig.

3A). Utilizing MASCaT, we detected non-lethal caspase activity, specifically in the neurons of

adult brains, using *ELAV-Gal4* (Fig. 3C). Caspase activity was prominent in OLs (Fig. 3C),

which is consistent with the previously reported activation pattern of caspase detected by

CaspaseTracker/CasExpress [20,21]. Previously, we reported that a specific subset of ORNs,

including Or42b, undergoes age-associated caspase activation [22]. Consistent with this finding,

we detected age-dependent caspase activation in the AL (Fig. 3C). Therefore, our newly

established reporter, MASCaT, enables the highly sensitive detection of caspase activity near

the cell membrane in cells of interest reconciling with Gal4/UAS-dependent gene

manipulations.

Fas3G overexpression activates caspase in a non-lethal manner

To investigate the regulatory role of Fas3 in caspase activity *in vivo*, we used Or42b neurons as

a model, which have previously been shown to undergo age-dependent caspase activation [22].

Initially, we evaluated the effects of *Dronc* overexpression on apoptosis by quantifying cell

body numbers in the antenna in young flies. Dronc induced cell loss while catalytically inactive

Dronc (Dronc^{CG}) did not, suggesting that Dronc induce apoptosis in a catalytic-dependent

manner (Fig. 4A, B). We then evaluated the effects of the overexpression of the executioner

caspases *Dcp-1* and *Drice*. Interestingly, although Dcp-1 induced complete cell loss, Drice did

not (Fig. 4A, B), suggesting that Drice requires an initiator caspase for activation, whereas Dcp-

1 does not. These results indicate that Or42b neurons are competent in caspase-mediated cell

loss and thus serve as a model for evaluating apoptosis. However, the expression of *Fas3G*,

even in combination with *Drice*, did not reduce the number of cells, implying that *Fas3G*

overexpression did not induce apoptosis (Fig. 4C, D).

Subsequently, we assessed the regulation of non-lethal caspase activity by Fas3G *in*

vivo. We first confirmed that MASCaT probes were distributed in the axons of ORNs (Fig. 4E).

Using MASCaT, we observed weak non-lethal caspase activation even in the control group at

a young age (Fig. 4F, G). The expression of *Fas3G* significantly promoted caspase activation

(Fig. 4F, G), indicating that Fas3G regulates non-lethal caspase activation. Moreover, the

simultaneous expression of *Drice* and *Fas3G* significantly enhanced caspase activation (Fig.

4F, G), signifying the promoting effect of their proximity. Moreover, the concurrent expression

of Diap-1, an E3 ligase known to inhibit caspase activation [23–25], mitigated the caspase

activation induced by Fas3G overexpression (Fig. 4H, I). This indicates that *Fas3G*

overexpression-facilitated non-lethal caspase activation is dependent on the core apoptotic

machinery. To test whether endogenous Fas3G regulates caspase activation, we generated

Fas3G-specific *shRNA* expressing strain (Fig. S4A, B). Knockdown of *Fas3* or *Fas3G* in Or42b

neurons didn't suppress the inherent, weak caspase activation (Fig. S4C, D). In agreement with

this, Drice's presence in the axon of ORNs with Fas3G was observed, yet its presence was not

strictly contingent on Fas3 or Fas3G (Fig. S4E). Given the numerous other membrane proteins

in proximity to Drice (Fig. 1G, H and Table S1), it is probable that additional proteins may be

responsible for recruiting Drice to the axon. These findings collectively imply that while Fas3G

is not essential for mild endogenous caspase activation, overexpression of *Fas3G* does enhance

non-lethal caspase activation in ORNs, thus providing a valuable model for exploring the

molecular processes that facilitate caspase activation without leading to cell death.

Fas3G is proximal to Dronc which is required for non-lethal caspase activation

Next, we investigated the underlying mechanism that enhances non-lethal activation facilitated

by *Fas3G* overexpression. Executioner caspases, Drice and Dcp-1, are activated by initiator

caspases, Dronc (Fig. 1A). We precisely analyzed Dronc's expression pattern in adult brains

and found that while its expression was low in control, overexpression of *Fas3G* in ORNs

markedly increased Dronc levels in those neurons (Fig. 5A). Moreover, akin to Drice, Dronc

was also found in close proximity to the overexpressed Fas3G, but not to the overexpressed

Fas3C in ORNs (Fig. 5B). Consequently, we examined whether Dronc is required for the non-

lethal caspase activation enhanced by *Fas3G* overexpression. Employing MASCaT, we

discovered that both knockdown of *Dronc* and the expression of the dominant-negative form

of *Dronc* hindered the non-lethal caspase activation facilitated by *Fas3G* overexpression (Fig.

5C, D), indicating a dependence on initiator caspase activity. Importantly1, the genetic

suppression of Dronc activity did not alter the proximity between Drice and Fas3G (Fig. 5E),

suggesting that the inactive Drice proform is naturally near Fas3G. Solo overexpression of

Dronc led to cell death in Or42b neurons (Fig. 4A, B), whereas *Fas3G* overexpression elevated

Dronc expression without causing cell death (Fig. 4C, D, Fig. 5A). Collectively, these findings

imply that the non-lethal caspase activation driven by *Fas3G* overexpression is orchestrated by

an increase in Dronc induced by Fas3G, bringing Dronc in close vicinity to Fas3G and thus

enabling caspase activation near the cell membrane (Fig. 5F).

Non-lethal caspase activation regulates innate attraction behavior

Finally, we examined the function of *Fas3G* overexpression-facilitated non-lethal caspase

activation. Apple cider vinegar (ACV) excites six glomeruli in the AL, including DM1, which

is innervated by the axons of Or42b neurons [26]. We have previously reported that Or42b

neurons undergo aging-dependent caspase activation, which reduces neural activity and

attraction behavior in response to ACV [22]. Inhibition of caspase activity in aged flies restores

attraction behavior but does not affect that of young flies [22]. Using a two-choice preference

assay with ACV (Fig. 6A), we found that the overexpression of *Fas3G*, which activates

caspases, did not impair attraction behavior in young flies (Fig. 6B, C). In contrast,

simultaneous inhibition of *Fas3G* overexpression-facilitated non-lethal caspase activation

promoted attraction to ACV (Fig. 6B, C). Overall, these results suggest that *Fas3G*

overexpression-facilitated non-lethal caspase activation suppresses innate attractive behavior

without killing the cells.

Discussion

Proximal proteins of executioner caspases regulate non-lethal caspase activity

It is important to identify the proximal proteins of executioner caspases, which cleave a wide range of substrates and are essential for apoptosis [11,12], as proximity often dictates substrate specificity [17]. In humans, Caspase-3 and Caspase-7 have been shown to share but also have discrete substrate preferences [27]. Caspase-7 leverages an exosite to facilitate interaction with RNA and enhance the proteolysis of RNA-binding proteins [28,29]. Thus, protein proximity can be used to precisely regulate specific cellular functions by selectively cleaving a limited pool of substrates in non-lethal scenarios compared with lethal cleavages where all the substrates are potentially cleaved during apoptosis. Our current investigation showed that executioner caspases, before activation as proforms, are exclusively sequestered in specific subcellular domains, including the plasma membrane. We found that *Fas3G* overexpression facilitates non-lethal activation, underscoring the importance of protein proximity not only in substrate cleavage, but also in localizing non-lethal activation to the vicinity of plasma membrane (Fig. 5F).

MASCaT as a highly sensitive reporter for identifying non-lethal caspase regulators

Various caspase activity reporters have been developed, including SCAT3, a FRET-based reporter with superior temporal resolution [30], CaspaseTracker/CasExpress, Caspase-activatable Gal4-based reporters with superior sensitivity [20,21]. However, to date, no reporter has been developed that is sensitive enough to identify non-lethal regulators of caspases in cells of interest. In this study, we developed a novel caspase reporter, MASCaT, that can detect non-lethal caspase activation with high sensitivity in a cell type-specific manner following gene manipulation. Using MASCaT, we identified previously overlooked regulators of non-lethal activation of caspases in a cell type-specific manner. Proximal proteins of executioner caspases

are potential non-lethal regulators, and combining TurboID-mediated identification of these

proteins will pave the way for identifying non-lethal components and broaden our

understanding of the molecular mechanism of non-lethal caspase activation.

Non-lethal caspase activation regulates neural activity and behaviors

We previously reported that age-dependent caspase activation reduces neural activity and innate

attraction behavior [22]. Here, we found that, while *Fas3G* overexpression did not impair

attraction behavior by itself, caspase activation upon *Fas3G* overexpression suppressed

attraction behavior toward ACV (Fig. 6A, B). In mice, L1CAM, an immunoglobulin-like cell

adhesion molecule, facilitates neuronal excitability by regulating the voltage-gated Na+

channels [31]. Thus, we hypothesize that non-lethal caspase activation antagonizes the neuronal

excitability promoted by Fas3G. In neurons, caspases have diverse non-lethal functions,

including axon degeneration and macro- and micro-pruning, to regulate synaptic plasticity

[32,33]. In presynaptic terminals, it has been recently reported that Caspase-3 modulates

synaptic vesicle pools via autophagy [34]. Thus, it is possible that non-lethal activation

regulates synaptic pools and neural activity in a reversible manner without killing the cells.

Analyzing the functions mediated by Drice and Fas3G proximity, including the identification

of specific substrates, will pave the way for identifying diverse CDPs and understanding their

molecular mechanisms.

While suppressing innate attraction behavior does not seem beneficial by itself, it should

be a great advantage in decision-making. To survive, flies must constantly make appropriate

behavioral decisions (e.g., feeding versus mating) depending on their physiological conditions.

In males, a subset of neurons regulated by tyramine controls feeding versus mating behaviors

[35]. In regard of that, it is important to adequately promote or suppress neural activity, which

controls behaviors of the organisms, by precisely controlling caspase activation in a

subcellularly restricted manner in neurons.

Proximal proteins of executioner caspases define the site of caspase activation "hotspot"

Although the core apoptotic machinery has been well characterized during the last 30 years [2],

the precise regulation of CDPs, remains incompletely elucidated. In this study, we identified

Fas3G overexpression activates Drice. Mechanistically, we found that *Fas3G* overexpression

induced the expression of initiator caspase Dronc, which also comes close to Fas3G, regulating

non-lethal caspase activation. Direct overexpression of Dronc alone results in cell death within

olfactory receptor neurons (Fig. 4A, B). However, when Dronc expression is upregulated due

to *Fas3G* overexpression, it does not lead to cell death in ORNs (Fig. 4C, D). The exact process

by which Dronc expression is increased in response to *Fas3G* overexpression remains to be

elucidated. It is known that Dronc can be upregulated via the ecdysone signaling pathway [36]

and the Hippo signaling pathway [37]. Therefore, the overexpression of *Fas3G* might

potentially activate Dronc expression through one or both of these signaling pathways. Fas3 is

an immunoglobulin-like domain-containing transmembrane protein that regulates axonal

fasciculation and neuronal projections [18]. NCAM, a mammalian homologue of Fascicilin 2,

is an immunoglobulin-like domain-containing cell adhesion molecule. NCAM can physically

bind to Caspase-8, and depending on NCAM clustering, Caspase-8 and -3 are activated and

contribute to neurite outgrowth by cleaving its substrate spectrin [38]. Thus, it seems likely that

cell adhesion molecule-mediated subcellularly restricted caspase activation is conserved among

species. Indeed, caspase-9/Apaf-1-dependent caspase-3 activation, without resulting in cell

death, has also been observed in the axon of olfactory sensory neurons during mouse

development [3]. A critical insight from our research is that the executioner caspase is pre-

compartmentalized in proximity to a specific set of proteins prior to activation, poised for the

initiator caspase to approach. Consequently, the "hotspot" for non-lethal caspase activity

appears to be determined by these neighboring proteins. While the "hotspot" located in the axon

of the ORN may exert an inhibitory effect on neural activity, it could also play a role in avoiding

cell death by localizing caspase activity, especially during development.

Recently, a protein interactome analysis in humans showed that tissue-preferentially

expressed proteins that interact with known apoptosis regulators participate in apoptosis

sensitization [39]. Given that CDPs are often cell-and tissue type-specific processes, the

identification of modulators of context-dependent apoptotic pathways will aid in understanding

the precise molecular regulation of CDPs. Importantly, alternative splicing can diversify

protein-protein interaction networks. A recent report showed that minor spliceosomes regulate

alternative splicing, including Fas3. A specific isoform of Fas3 has distinct functions and cannot

be substituted by other isoforms during neuronal development [40]. Our results support the idea

that alternative splicing may not only diversify the protein itself, but also its interaction

networks to regulate module-dependent activation. Further interactome analysis using TurboID

will reveal the regulatory mechanisms of the non-lethal activation of caspases.

Materials and Methods

Fly strains and rearing conditions

We raised the flies in standard *Drosophila* medium at 25°C. For in vivo biotin labeling experiments, flies were raised in standard *Drosophila* medium supplemented with 100 μM (+)-biotin (#132-02635, WAKO). The following fly strains were used in this study: w^{1118}, *Drice::V5::TurboID*, *Dcp-1::V5::TurboID*, *Dronc::V5::TurboID* [16], *Pebbled-Gal4* (BL80570), *LN-Gal4* (BL63325), *GH146-Gal4* (BL30026), *UAS-Drice-RNAi* (V28065), *Fas3::EGFP::FlAsH::StrepII::TEV::3xFLAG* (BL59809), *UASz-Fas3A::3xFLAG*, *UASz-Fas3B::3xFLAG*, *UASz-Fas3C::3xFLAG*, *UASz-Fas3D::3xFLAG*, *UASz-Fas3G::3xFLAG* (this study), *Ubi-FRT-STOP-FRT-mCD8::DQVD::QF2*, *Ubi-FRT-STOP-FRT-mCD8::DQVA::QF2* (this study), *UAS-FLP* (K107788), *QUAS-tdTomato::3xHA* (BL30005), *ELAV-Gal4* (BL8765), *Or42b-Gal4* (BL9971), *UAS-RedStinger* (BL8547), *UASz-myc::mNeonGreen*, *UASz-Drice::myc::mNeonGreen* (this study), *UASz-V5::TurboID*, *UASz-Drice::V5::TurboID*, *UASz-Drice^{CG}::V5::TurboID*, *UASz-Dcp-1::V5::TurboID*, *UASz-Dcp-1^{CG}::V5::TurboID*, *UASz-Dronc::V5::TurboID*, *UASz-Dronc^{CG}::V5::TurboID* (this study), *UAS-Diap-1::VENUS* [41], *UAS-Dronc-RNAi* [9], *UAS-Dronc^{DN}* (a gift from H. Richardson), *UAS-LacZ-RNAi*, *UAS-Fas3-RNAi#1* (V940), *UAS-Fas3-RNAi#2* (V3091), *UAS-Fas3G-RNAi* (this study). BL, Bloomington Drosophila Resource Center; K, Kyoto Stock Center; V, Vienna Drosophila Resource Center.

Detailed genotypes

Fig. 1B, C (Left to Right, Top to Bottom)

w^{1118}/Y

w/Y; Dcp-1::V5::TurboID/+

w/Y;; Drice::V5::TurboID/+

w/Y;; Dronc::V5::TurboID/+

Fig. 1D (Left to Right)

w; UAS-Drice-RNAi/+; Drice::V5::TurboID/+

w, Pebbled-Gal4/w; UAS-Drice-RNAi/+; Drice::V5::TurboID/+

w; UAS-Drice-RNAi/GH146-Gal4; Drice::V5::TurboID/+

w; UAS-Drice-RNAi/+; Drice::V5::TurboID/LN-Gal4

Fig. 1E (Left to Right)

w^{1118}/Y

w/Y;; Drice::V5::TurboID/+

Fig. 1G (Left to Right)

w^{1118}/Y

w/Y;; Drice::V5::TurboID/+

Fig. 2A (Left to Right)

w^{1118}/Y

w/Y;; Drice::V5::TurboID/+

Fig. 2B (Top to Bottom, Left to Right)

w^{1118}/Y

w/Y;; Drice::V5::TurboID/+

w/Y; Fas3::EGFP::FlAsH::StrepII::TEV::3xFLAG/+

w/Y; Fas3:: EGFP::FlAsH::StrepII::TEV::3xFLAG/+; Drice::V5::TurboID/+

Fig. 2E (Left to Right)

w, Pebbled-Gal4/w or Y

w, Pebbled-Gal4/w or Y;; UASz-Fas3A::3xFLAG/+

w, Pebbled-Gal4/w or Y;; UASz-Fas3B::3xFLAG/+

w, Pebbled-Gal4/w or Y;; UASz-Fas3C::3xFLAG/+

w, Pebbled-Gal4/w or Y;; UASz-Fas3D::3xFLAG/+

w, Pebbled-Gal4/w or Y;; UASz-Fas3G::3xFLAG/+

Fig. 2F (Left to Right)

w^{1118}/Y

w/Y;; Drice::V5::TurboID/+

Fig. 2G (Left to Right)

w, Pebbled-Gal4/w or Y;; Drice::V5::TurboID/+

w, Pebbled-Gal4/w or Y;; Drice::V5::TurboID/UASz-Fas3A::3xFLAG

w, Pebbled-Gal4/w or Y;; Drice::V5::TurboID/UASz-Fas3B::3xFLAG

w, Pebbled-Gal4/w or Y;; Drice::V5::TurboID/UASz-Fas3C::3xFLAG

w, Pebbled-Gal4/w or Y;; Drice::V5::TurboID/UASz-Fas3D::3xFLAG

w, Pebbled-Gal4/w or Y;; Drice::V5::TurboID/UASz-Fas3G::3xFLAG

Fig. 3C (Top to Bottom)

w/Y; Ubi-FRT-STOP-FRT-mCD8::DQVD::QF2/+; UAS-FLP, QUAS-tdTomato::3xHA/+

w/Y;   Ubi-FRT-STOP-FRT-mCD8::DQVA::QF2/ELAV-Gal4;   UAS-FLP,   QUAS-

tdTomato::3xHA/+

w/Y;   Ubi-FRT-STOP-FRT-mCD8::DQVD::QF2/ELAV-Gal4;   UAS-FLP,   QUAS-

tdTomato::3xHA/+

Fig. 4A, B (Top to Bottom, Left to Right)

w/Y; Or42b-Gal4/+; UAS-RedStinger/UASz-V5::TurboID

w/Y; Or42b-Gal4/+; UAS-RedStinger/UASz-Drice^{CG}::V5::TurboID

w/Y; Or42b-Gal4/+; UAS-RedStinger/UASz-Drice::V5::TurboID

w/Y; Or42b-Gal4/+; UAS-RedStinger/UASz-Dcp-1^{CG}::V5::TurboID

w/Y; Or42b-Gal4/+; UAS-RedStinger/UASz-Dcp-1::V5::TurboID

w/Y; Or42b-Gal4/+; UAS-RedStinger/UASz-Dronc^{CG}::V5::TurboID

w/Y; Or42b-Gal4/+; UAS-RedStinger/UASz-Dronc::V5::TurboID

Fig. 4C, D (Top to Bottom, Left to Right)

w/Y; Or42b-Gal4/UASz-3xFLAG::mNeonGreen; UAS-RedStinger/+

w/Y; Or42b-Gal4/+; UAS-RedStinger/UASz-Fas3G::3xFLAG

w/Y; Or42b-Gal4/UASz-Drice::myc::mNeonGreen; UAS-RedStinger/+

w/Y; Or42b-Gal4/UASz-Drice::myc::mNeonGreen; UAS-RedStinger/UASz-Fas3G::3xFLAG

Fig. 4E (Top to Bottom)

w/Y; Or42b-Gal4/UASz-3xFLAG::mNeonGreen

w/Y; Or42b-Gal4, Ubi-FRT-STOP-FRT-mCD8::DQVD::QF2/UASz-3xFLAG::mNeonGreen;

UAS-FLP, QUAS-tdTomato::3xHA/+

Fig. 4F, G (Top to Bottom, Left to Right)

w/w or Y; Or42b-Gal4, Ubi-FRT-STOP-FRT-mCD8::DQVD::QF2/UASz-

3xFLAG::mNeonGreen; UAS-FLP, QUAS-tdTomato::3xHA/+

w/w or Y; Or42b-Gal4, Ubi-FRT-STOP-FRT-mCD8::DQVD::QF2/+; UAS-FLP, QUAS-

tdTomato::3xHA/UASz-Fas3G::3xFLAG

w/w or Y; Or42b-Gal4, Ubi-FRT-STOP-FRT-mCD8::DQVD::QF2/UASz-

Drice::myc::mNeonGreen; UAS-FLP, QUAS-tdTomato::3xHA/+

w/w or Y; Or42b-Gal4, Ubi-FRT-STOP-FRT-mCD8::DQVD::QF2/UASz-

Drice::myc::mNeonGreen; UAS-FLP, QUAS-tdTomato::3xHA/UASz-Fas3G::3xFLAG

Fig. 4H, I (Top to Bottom, Left to Right)

w/w or Y; Or42b-Gal4, Ubi-FRT-STOP-FRT-mCD8::DQVD::QF2/UASz-

3xFLAG::mNeonGreen; UAS-FLP, QUAS-tdTomato::3xHA/+

w/w or Y; Or42b-Gal4, Ubi-FRT-STOP-FRT-mCD8::DQVD::QF2/+; UAS-FLP, QUAS-

tdTomato::3xHA/UASz-Fas3G::3xFLAG

w/w or Y; Or42b-Gal4, Ubi-FRT-STOP-FRT-mCD8::DQVD::QF2/UAS-Diap-1::VENUS;

UAS-FLP, QUAS-tdTomato::3xHA/UASz-Fas3G::3xFLAG

Fig. 5A (Left to Right)

w, Pebbled-Gal4/Y

w, Pebbled-Gal4/Y;; Dronc::V5::TurboID/+

w, Pebbled-Gal4/Y;; Dronc::V5::TurboID/UASz-Fas3G::3xFLAG

Fig. 5B (Left to Right)

w, Pebbled-Gal4/Y;; Dronc::V5::TurboID/+

w, Pebbled-Gal4/Y;; Dronc::V5::TurboID/UASz-Fas3G::3xFLAG

w, Pebbled-Gal4/Y;; Dronc::V5::TurboID/UASz-Fas3C::3xFLAG

Fig. 5C, D (Top to Bottom, Left to Right)

w/w   or   Y;   Or42b-Gal4,   Ubi-FRT-STOP-FRT-mCD8::DQVD::QF2/UASz-

3xFLAG::mNeonGreen; UAS-FLP, QUAS-tdTomato::3xHA/+

w/w or Y; Or42b-Gal4, Ubi-FRT-STOP-FRT-mCD8::DQVD::QF2/+; UAS-FLP, QUAS-

tdTomato::3xHA/UASz-Fas3G::3xFLAG

w/w or Y; Or42b-Gal4, Ubi-FRT-STOP-FRT-mCD8::DQVD::QF2/UAS-Dronc-RNAi; UAS-

FLP, QUAS-tdTomato::3xHA/UASz-Fas3G::3xFLAG

w/w or Y; Or42b-Gal4, Ubi-FRT-STOP-FRT-mCD8::DQVD::QF2/UAS-Dronc^{DN}; UAS-FLP,

QUAS-tdTomato::3xHA/UASz-Fas3G::3xFLAG

Fig. 5E (Left to Right)

w, Pebbled-Gal4/Y;; Drice::V5::TurboID/+

w, Pebbled-Gal4/Y;; Drice::V5::TurboID/UASz-Fas3G::3xFLAG

w, Pebbled-Gal4/Y; +/UAS-Dronc-RNAi; Drice::V5::TurboID/UASz-Fas3G::3xFLAG

w, Pebbled-Gal4/Y; +/UAS-Dronc^{DN}; Drice::V5::TurboID/UASz-Fas3G::3xFLAG

Fig. 6B, C (Left to Right)

w/Y; Or42b-Gal4/UASz-3xFLAG::mNeonGreen

w/Y; Or42b-Gal4/+; +/UASz-Fas3G::3xFLAG

w/Y; Or42b-Gal4/UAS-Dronc-RNAi; +/UASz-Fas3G::3xFLAG

w/Y; Or42b-Gal4/UAS-Dronc^{DN}; +UASz-Fas3G::3xFLAG

Fig. S1A (Left to Right, Top to Bottom)

w^{1118}/Y

w^{1118}

w/Y;; Drice::V5::TurboID/+

w/w^{1118};; Drice::V5::TurboID/+

Fig. S2A (Left to Right)

w, Pebbled-Gal4/w or Y;; Drice::V5::TurboID/+

w, Pebbled-Gal4/w or Y;; Drice::V5::TurboID/UASz-Fas3C::3xFLAG

w, Pebbled-Gal4/w or Y;; Drice::V5::TurboID/UASz-Fas3G::3xFLAG

Fig. S4C, D (Top to Bottom, Left to Right)

w/w or Y; Or42b-Gal4, Ubi-FRT-STOP-FRT-mCD8::DQVD::QF2/UAS-LacZ-RNAi; UAS-

FLP, QUAS-tdTomato::3xHA/+

w/w or Y; Or42b-Gal4, Ubi-FRT-STOP-FRT-mCD8::DQVD::QF2/+; UAS-FLP, QUAS-

tdTomato::3xHA/UAS-Fas3-RNAi #1

w/w or Y; Or42b-Gal4, Ubi-FRT-STOP-FRT-mCD8::DQVD::QF2/+; UAS-FLP, QUAS-

tdTomato::3xHA/UAS-Fas3-RNAi #2

w/w or Y; Or42b-Gal4, Ubi-FRT-STOP-FRT-mCD8::DQVD::QF2/UAS-Fas3G-RNAi; UAS-

FLP, QUAS-tdTomato::3xHA/+

Fig. S4E (Left to Right, Top to Bottom)

w, Pebbled-Gal4/Y; UASz-Drice::myc::mNeonGreen/+

w, Pebbled-Gal4/Y; UASz-Drice::myc::mNeonGreen/+; UASz-Fas3G::3xFLAG/+

w, Pebbled-Gal4/Y; UASz-Drice::myc::mNeonGreen/+; UAS-Fas3-RNAi #1/+

w, Pebbled-Gal4/Y; UASz-Drice::myc::mNeonGreen/+; UAS-Fas3-RNAi #2/+

w, Pebbled-Gal4/Y; UASz-Drice::myc::mNeonGreen/UAS-Fas3G-RNAi

Protein preparation from Drosophila tissues

Adult heads or brains of the given genotype were dissected and lysed with RIPA buffer (50 mM

Tris-HCl, pH 8.0, 150 mM Sodium Chloride, 0.5 w/v% Sodium Deoxycholate, 0.1% w/v

Sodium Dodecyl Sulfate, and 1.0% w/v NP-40) supplemented with cOmplete ULTRA EDTA-

free protease inhibitor cocktail (#05892953001, Roche). Samples were homogenized and

centrifuged at 20,000 ×g, 4°C for 10 min. The supernatants were collected and snap-frozen in

liquid nitrogen. Protein concentrations were determined using the BCA assay (#297-73101,

Wako) following the manufacturer's protocol. The samples were mixed with 6x Laemmli

buffer, boiled at 95°C for 5 min, and then subjected to purification of biotinylated proteins and

western blot analysis.

Western blotting

Each sample was separated using SDS-PAGE. The proteins were then transferred onto

Immobilon-P PVDF membranes (#IPVH00010; Millipore) for immunoblotting. Membranes

were blocked with 4% skim milk diluted in 1× TBST. Immunoblotting was performed using

the antibodies mentioned below diluted in 4% skim milk. Signals were visualized using

Immobilon Western Chemiluminescent HRP Substrate (#WBKLS0500; Millipore) and

FUSION SOLO. 7S. EDGE imaging station (Vilber-Lourmat). Contrast and brightness

adjustments were applied equally using Fiji (ImageJ) software (NIH Image).

The primary antibodies used included mouse anti-V5 monoclonal antibody (1:5,000,

#46-0705, Invitrogen), mouse anti-FLAG M2 monoclonal antibody (1:5,000, Sigma), mouse

anti-alpha tubulin (DM1A) monoclonal antibody (1:10,000, #T9026, Sigma), mouse anti-Actin

monoclonal antibody (1:5,000; #A4700, Sigma), mouse anti-Fasciclin 3 (7G10) antibody (1:50,

#AB_528238, DSHB), rabbit anti-Fasciclin3 isoform G antibody (1:50, this study), mouse anti-

Drosophila Lamin B monoclonal antibody (1:1000, #ADL67.10, DSHB). The secondary

antibodies used included goat anti-rabbit IgG HRP-conjugated antibody (1:5,000, #7074S,

CST) and goat/rabbit/donkey anti-mouse IgG HRP Conjugate (1:5,000, #W402B, Promega).

The membranes for streptavidin blotting were blocked with 3% BSA. Streptavidin blotting was

performed using streptavidin-horseradish peroxidase (1:10,000, #SA10001; Invitrogen) diluted

in 3% BSA.

Co-immunoprecipitation

Adult heads were lysed on ice using DDM IP lysis buffer (50 mM Tris-HCl (pH 7.5), 150 mM

NaCl, 0.2% N-Dodecyl-β-D-maltoside (DDM) (#341-06161, WAKO)) containing cOmplete

ULTRA EDTA-free protease inhibitor cocktail. The samples were adjusted to 150 μg

protein/150 μL DDM IP lysis buffer. The samples were incubated overnight at 4 °C with 10

μL anti-FLAG M2 Magnetic Beads (#M8823-1ML, Millipore) equilibrated with the DDM IP

lysis buffer. The beads were washed three times in wash buffer (50 mM Tris-HCl (pH 7.5),

150 mM NaCl, 0.1% DDM) and boiled for 5 min at 95 °C with 50 μL 1× Laemmli buffer. The

samples were magnetically separated, and the supernatants were subjected to SDS-PAGE.

Immunohistochemistry

Adult brains were dissected in phosphate-buffered saline (PBS) and fixed in 4%

paraformaldehyde (PFA) at room temperature for 30 min. Adult brains were washed three times

with 0.3% PBST (Triton-X100) for 10 min. After blocking with 5.0% normal donkey serum

(#S30, Millipore)/PBST (PBSTn) for 30 minutes at room temperature, tissues were incubated

with primary antibody/PBSTn overnight at 4°C. The samples were then washed with PBST

three times for 10 min each and incubated with secondary antibody/PBSTn for 2 h at room

temperature. After incubation, the tissues were washed with PBST three times for 10 min.

Streptavidin and Hoechst 33342 were added during secondary antibody incubation. The

samples were mounted on glass slides with SlowFade Gold antifade reagent (#S36939,

Invitrogen). Images were captured using a Leica TCS SP8 micrscope (Leica Microsystems).

Images were analyzed and edited using the Fiji (ImageJ) software. For MASCaT analysis,

newly eclosed adults were raised at 29°C for 1 week. Classification of the MASCaT signal was

performed by optically assessing the tdTomato intensity. For cell number analysis, newly

eclosed adults were raised at 29°C for 1 week. The antennae were dissected as previously

described [42], fixed with 4% PFA at room temperature for 30 min, and washed with 0.4%

PBST (Triton-X100) three times for 10 min. Quantification of cell numbers in antenna was

performed using Fiji (ImageJ) software using "3D object counter" for RedStinger positive cells.

The primary antibodies used were mouse anti-Fasciclin 3 (7G10) antibody (1:20,

#AB_528238, DSHB), rat anti-GFP (GF090R) monoclonal antibody (1:100, #04404-26,

nacalai Tesque), mouse anti-FLAG M2 monoclonal antibody (1:500, F1804, Sigma), and rat

anti-mCD8 monoclonal antibody (1:50, #MCD0800, Invitrogen). The secondary antibodies

used included Alexa Fluor 488-conjugated donkey anti-mouse IgG antibody (1:500, #A-21202,

ThermoFisher Scientific), Alexa Fluor 488-conjugated donkey anti-rat IgG antibody (1:500,

#A-21208, ThermoFisher Scientific), Alexa Fluor 647-conjugated donkey anti-mouse IgG

antibody (1:500, #A-31571, ThermoFisher Scientific), and Alexa Fluor 633-conjugated goat

anti-rat IgG antibody (1:500, #A-21094, ThermoFisher Scientific). The nuclei were stained

with Hoechst 33342 (8 µM, #H3570, Invitrogen). Streptavidin-Cy2 (1:500, #016-220-084,

Jackson ImmunoResearch) and Streptavidin-Cy5 (1:500, #016-170-084, Jackson

ImmunoResearch) were used to stain the biotinylated proteins.

Experimental design for LC-MS/MS analyses

Two biological replicates were analyzed for each experimental condition to determine the

relative abundance of *Drice::V5::TurboID/w^{1118}*.

Purification of biotinylated proteins

Samples were prepared as previously described [17]. Briefly, 100 µg of biotinylated protein-

containing lysate (from 100 brains or 20 heads) was subjected to FG-NeutrAvidin bead

(#TAS8848 N1171, Tamagawa Seiki) purification. The FG-NeutrAvidin beads (25 µL,

approximately 500 µg) were washed three times with RIPA buffer. Benzonase-treated

biotinylated protein samples suspended in 1 mL RIPA buffer were incubated overnight at 4°C.

Conjugated beads were magnetically isolated and washed with 500 µL of an ice-cold RIPA

buffer solution, 1 M KCl solution, 0.1 M Na_2CO_3 solution and 4 M urea solution. For western

blot analyses, the purified samples were mixed with Laemmli buffer, boiled at 95°C for 5 min,

and then subjected to western blot analysis. For LC-MS/MS analysis, the purified samples were

washed with 500 µL ultrapure water (#214-01301, Wako) and 500 µL of 50 mM ammonium

bicarbonate (AMBC). The samples were then mixed with 50 µL of 0.1% Rapigest diluted in 50

mM AMBC, and 5 µL of 50 mM TCEP was subsequently added. Samples were incubated at

60°C for 5 min, and then 2.5 µL 200 mM MMTS was added. One microgram sequence-grade

modified trypsin (#V5111, Promega) was then added for on-bead trypsin digestion at 37°C for

16 h. The beads were magnetically isolated and 60 µL of the supernatant was collected. Then,

3 µL 10% TFA was added to the supernatants, and the samples were incubated at 37°C for 60

min with gentle agitation. The samples were then centrifuged at 20,000 ×g, 4°C for 10 min. The

peptides were desalinated and purified using a GL-tip SDB (#7820-11200, GL Sciences)

following the manufacturer's protocol. The samples were speed-backed at 45°C for 30 min and

dissolved in 25 µL of 0.1% formic acid. The samples were then centrifuged at 20,000 ×g, 4°C

for 10 min, and the supernatants were collected. The peptide concentrations were determined

using the BCA assay. Finally, 500 ng of the purified protein was subjected to LC-MS/MS

analysis.

LC-MS/MS analyses

LC-MS/MS analyses were performed as previously described [17]. Briefly, samples were

loaded onto Acclaim PepMap 100 C18 column (75 μm x 2 cm, 3 μm particle size and 100 Å

pore size; #164946, ThermoFisher Scientific) and separated on nano-capillary C18 column, (75

μm x 12.5 cm, 3 μm particle size, #NTCC-360/75-3-125, Nikkyo Technos) using EASY-nLC

1200 system (ThermoFisher Scientific). The elution conditions are listed in Table S3. The

separated peptides were analyzed using QExactive (ThermoFisher Scientific) in data-dependent

MS/MS mode. The parameters for the MS/MS analysis are listed in Table S4. The collected

data were analyzed using Proteome Discoverer (PD) 2.2 software with the Sequest HT search

engine. The parameters for the PD 2.2 analysis are described in Table S5. Peptides were filtered

at a false discovery rate of 0.01 using the Percolator node. Label-free quantification was

performed based on the intensities of the precursor ions using a precursor-ion quantifier node.

Normalization was performed using the total amount of peptides in all average scaling modes.

Proteins with 10 folds or higher abundance relative to the control (w^{1118}) were considered for

further analysis. MS proteomic data were deposited in the ProteomeXchange Consortium via

the jPOST partner repository with the dataset identifier PXD042922 [43].

Gene Ontology analysis

Gene Ontology (GO) analysis of cellular components was performed using DAVID [44].

Generation of rabbit polyclonal anti-Fasciclin 3 isoform G antibody

The peptide corresponding to the 506–524 AA (N-LKPANEATPATTPAPTTAA-C) of Fas3G

was used by Eurofins Inc. to immunize rabbits to raise a polyclonal antibody. The antibody was

affinity purified using the same peptide provided by Eurofins, Inc.

IDR prediction

Amino acid sequences were obtained from the FlyBase database [https://flybase.org/].

Disordered intracellular regions of Fas3 isoforms were predicted using the PSIPRED protein

sequence analysis workbench in DISOPRED3 [http://bioinf.cs.ucl.ac.uk/psipred/].

Molecular cloning

For all isoforms (A, B/F, C, D/E, G) of *pUASz-Fasciclin 3::3xFLAG*, the coding sequences of

Fasciclin 3s were PCR-amplified using 3xFLAG-tag harboring primers from *Drosophila* cDNA

and were ligated into the *BamH*I/*Kpn*I-digested *pUASz1.0* vector [45] using In-Fusion

(#Z9648N, TaKaRa). For all isoforms of *pAc5-Fasciclin 3::3xFLAG*, the coding sequence of

Fasciclin 3s::3xFLAG were PCR-amplified from corresponding *pUASz-Fasciclin 3::3xFLAG*

vectors and were ligated into the *EcoR*I/*Xho*I-digested *pAc5-STABLE2-neo* vector using

NEBuider HiFi DNA Assembly. using NEBuilder HiFi DNA Assembly (#E2621L, NEB). For

pUASz-3xFLAG::mNeonGreen, the coding sequence of mNeonGreen was PCR-amplified

using 3xFLAG-tag harboring primers from *pAc5-V5::mNeonGreen* [17] and ligated into the

*Xho*I-digested *pUASz1.0* vector [45] using In-Fusion. For *pUASz-Drice::myc::mNeonGreen*,

the coding sequence of mNeonGreen was PCR-amplified from *pAc5-V5::mNeonGreen* [17]

using myc-tag-harboring primers. The coding sequence of Drice was PCR-amplified from

Drosophila cDNA. The two fragments were then ligated into the *Xho*I-digested *pUASz1.0*

vector using In-Fusion. For *pUASz-V5::TurboID* and *pUASz-Caspase::V5::TurboID*, the

coding sequences of *Dcp-1*, *Drice* and *Dronc* were PCR amplified from *Drosophila* cDNA.

The coding sequence of V5::TurboID was PCR-amplified from *V5-TurboID-NES_pcDNA3*

(#107169; Addgene). The two fragments were then ligated into the *Xho*I-digested *pUASz1.0*

vector using In-Fusion. Catalytic dead mutations (Dcp-1: C196G (TGC to GGC), Drice: C211G

(TGC to GGC), and Dronc: C318G (TGC to GGC)) were introduced by PCR-based site-

directed mutagenesis, and the corresponding fragments were ligated into the *Xho*I-digested

pUASz1.0 vector using In-Fusion. For *pJFRC-mCD8::DQVD/A::QF2*, the coding sequence of

mCD8::DQVD/A were PCR-amplified from genomic DNA of *Caspase-Sensitive/Insensitive-*

Gal4 [20]. The coding sequence of *QF2* was PCR-amplified from *pBPQU* vector [46]. These

fragments were ligated into the *Xho*I/*Xba*I-digested *pJFRC-28K* vector [45] using In-Fusion.

For *pUbi-FRT-STOP-FRT-mCD8::DQVD/A::QF2*, the FRT-STOP-FRT cassette sequence

was PCR amplified from *pJFRC201-10xUAS-FRT-STOP-FRT-myr::smGFP-HA* (#63166;

Addgene). The coding sequence of mCD8::DQVD/A was PCR-amplified from the genomic

DNA of *Caspase-Sensitive/Insensitive-Gal4* [20]. The coding sequence of QF2 was PCR

amplified from the *pBPQUw* plasmid [46]. The three fragments were ligated into *EcoR*I-

digested *pUbi83* vector (a gift from D. Umetsu) using In-Fusion. For *pQUAST-mNeonGreen*,

the coding sequence of mNeonGreen was PCR-amplified from *pAc5-V5::mNeonGreen* [17]

and ligated into the *EcoR*I-digested *pQUAST* (#24349, Addgene) vector using In-Fusion. For

pUASz-FLPo, the coding sequences of FLPo were PCR-amplified from *pQUAST-FLPo*

(#24357, Addgene) and ligated into the *Xho*I-digested *pUASz1.0* vector [45] using In-Fusion.

For *pWALIUM20-Fas3G-shRNA*, the top and bottom oligonucleotides including *mRNA*

sequence specific to isoform G (5'-AATGAACCAAAGCAAGACAAA-3') were annealed in

annealing buffer (10 mM Tris-HCl, 0.1 M NaCl, 1 mM EDTA). The annealed fragment was

ligated into *Nhe*I/*Eco*RI-digested *pWALIUM20* (#1472, DGRC) vector using DNA Ligation Kit

<Mighty Mix> (#6023, TaKaRa).

All established plasmids were sequenced by Eurofins, Inc. The detailed plasmid DNA

sequences are available upon request. The established plasmids were injected into y^1, w^{67c23};

P[CaryP]attP40 (*pUASz-3xFLAG::mNeonGreen*, *pUASz-Drice::myc::mNeonGreen* and

pWALIUM20-Fas3G-shRNA) or y^1, w^{67c23}; *P[CaryP]attP2* (*pUASz-Fasciclin 3s::3xFLAG*,

pUbi-FRT-STOP-FRT-mCD8::DQVD/A::QF2, *pUASz-V5::TurboID* and *pUASz-*

Caspase::V5::TurboID,) for PhiC31-mediated transgenesis (BestGene Inc.). Each red-eye-

positive transformant was isogenized. Integration of the attP landing site was confirmed using

genomic PCR.

Cell culture

Drosophila S2 cells were grown at 25°C in Schneider's *Drosophila* medium (#21720001,

GIBCO) supplemented with 10% (v/v) heat-inactivated fetal bovine serum, 100 U/ml penicillin,

and 100 μg/ml streptomycin (#168-23191, WAKO). Cells were seeded in 24-well plates and

transfected with the desired plasmids using Effectene Transfection Reagent (#301427,

QIAGEN) following the manufacturer's protocol. Sixteen hours after transfection, the cells

were reseeded into CELL CULTURE DISH, PS, 35/10 MM (#627870, Grainer BIO-ONE) and

incubated for 24 h. Images were obtained using a Leica AF6000 DMI6000 B microscope (Leica

Microsystems). Images were analyzed and edited using Fiji software (NIH Image).

For the substrate cleavage assay, cells were seeded in 12-well plates and were

transfected with desired *pAc5-Fas3::3xFLAG* plasmids using Effectene Transfection Reagent

following the manufacturer's protocol. After 40 h of incubation, the cells were treated with

10 μg/mL cycloheximide (CHX, #C7698, Sigma) for apoptosis induction. After 6 h of

incubation, cells were collected using RIPA buffer supplemented with cOmplete ULTRA

EDTA-free protease inhibitor cocktail depending on the cell status. For apoptotic cell lysate

collection, cells were resuspended by pipetting and were collected into a 1.5 mL tube. The cells

were then centrifuged at 800 g, 4 °C for 5 min, resuspended in PBS, and centrifuged again using

the same parameters. The cell pellets were then lysed with 100 μL RIPA buffer. For the other

experimental conditions, the cells were washed thrice with PBS and directly lysed on the plate

with 100 μL RIPA buffer. The supernatants were collected and snap-frozen in liquid nitrogen.

Protein concentrations were determined using the BCA assay following the manufacturer's

protocol. The samples were mixed with 6x Laemmli buffer, boiled at 95°C for 5 min, and then

subjected to western blot analysis.

For the validation of Fas3G-shRNA, cells were seeded in 12-well plates and were co-

transfected with *pActin-Gal4,* desired *pUASz-Fas3::3xFLAG* and *pWALIUM20-empty* or

pWALIUM20-Fas3G-shRNA plasmids using Effectene Transfection Reagent following the

manufacturer's protocol. After 40 h of incubation, the cells were washed thrice with PBS and

directly lysed on the plate with 100 μL RIPA buffer. The supernatants were collected and snap-

frozen in liquid nitrogen. Protein concentrations were determined using the BCA assay

following the manufacturer's protocol. The samples were mixed with 6x Laemmli buffer,

boiled at 95°C for 5 min, and then subjected to western blot analysis.

Aging experiment

Adult male flies were raised in vials containing 15 fly/vial, and incubated at 29°C, 65%

humidity with 12h/12h light/dark cycle. The flies were transferred every 2–4 days to vials

containing fresh food.

Two choice preference assay

Newly eclosed adult male flies were raised in vials containing 25 fly/vial, and incubated at 29°C,

65% humidity with 12h/12h light/dark cycle. The flies were transferred every 2–4 days to vials

containing fresh food for 1 week. The flies were starved in MilliQ (MQ) for 4 h prior to the

assay. About 75 flies were put in 1000-mL glass beaker, which contained two standards bottles

filled with 500 µl 1% ACV (Mizkan) and 500 µl MQ. The bottles were then covered with

sponge plugs containing pipette tips. The pipette tip was cut to increase the size of the opening

such that only one fly could climb into the bottle at a time. The glass beaker was covered with

a mesh fabric. The beakers were set in an incubator at 29°C for 2 hr for the assay. The

experiments were evaluated only if at least one of the tested flies entered either of 2 bottles.

The preference index equaled the number of flies in the ACV minus the number of flies in the

MQ, divided by the number of total live flies in both inside and outside of the bottles, and

multiplied by 100. Values greater than zero indicate a preference for 1% ACV and values less

than zero indicate aversion. We repeated the same experiment with different sets of flies and

the number of times is indicated by N.

Statistical analysis

Statistical analyses were performed using the GraphPad Prism 8 software (MDF). Data are

shown as the means ± standard error of the mean. *P*-values were calculated using a one-way

analysis of variance (ANOVA) with Bonferroni correction (Fig. 4B for selected pairs, Fig. 4D

for every pair). *P*-values were calculated using the chi-square test with Bonferroni's correction,

with mNeonGreen as a control (Fig. 4G), LacZ-RNAi as a control (Fig. S4D), mNeonGreen

and Fas3G as controls (Fig. 4I, 5D), and mNeonGreen as a control (Fig. 6B). *P*-values were

calculated using one-way ANOVA with Dunnett's multiple comparison test (Fig. 6C,

mNeonGreen as a control). NS: $P > 0.05$, †: < 0.05.

Acknowledgments

We would like to thank Dr. D. Umetsu for providing plasmids, Dr. Y. Nitta, Dr. A. Sugie, and Dr. T. Ichinose for helpful discussions, and the Bloomington *Drosophila* Stock Center, Vienna Drosophila Resource Center and Kyoto Stock Center for providing the fly strains. We thank Dr. S. Hirayama, Dr. S. Murata, and the one-stop sharing faculty center for future drug discovery at the Graduate School of Pharmaceutical Sciences, University of Tokyo, for the LC-MS/MS analysis. We thank Miura's lab members for their technical assistance and discussions; in particular, K. Takenaga for preparing the fly food, and R. Takamoto for experimental support. The MASCaT was established by MASaya Muramoto with critical advice from members of Shu MASuda at MASayuki Miura Laboratories. We would like to thank *Editage* (www.editage.com) for English language editing of this manuscript.

References

1. Green DR. The Coming Decade of Cell Death Research: Five Riddles. Cell. 2019;177:

1094–1107. doi:10.1016/j.cell.2019.04.024

2. Nakajima YI, Kuranaga E. Caspase-dependent non-apoptotic processes in development.

Cell Death Differ. 2017;24: 1422–1430. doi:10.1038/cdd.2017.36

3. Ohsawa S, Hamada S, Kuida K, Yoshida H, Igaki T, Miura M. Maturation of the

olfactory sensory neurons by Apaf-1/caspase-9–mediated caspase activity. Proc Natl

Acad Sci U S A. 2010;107: 13366–13371. doi:10.1073/pnas.0910488107

4. Li Z, Jo J, Jia JM, Lo SC, Whitcomb DJ, Jiao S, et al. Caspase-3 activation via

mitochondria is required for long-term depression and AMPA receptor internalization.

Cell. 2010;141: 859–871. doi:10.1016/j.cell.2010.03.053

5. Aram L, Yacobi-Sharon K, Arama E. CDPs: Caspase-dependent non-lethal cellular

processes. Cell Death Differ. 2017;24: 1307–1310. doi:10.1038/cdd.2017.111

6. D'Brot A, Chen P, Vaishnav M, Yuan S, Akey CW, Abrams JM. Tango7 directs cellular

remodeling by the Drosophila apoptosome. Genes Dev. 2013;27: 1650–1655.

doi:10.1101/gad.219287.113

7. Kang Y, Neuman SD, Bashirullah A. Tango7 regulates cortical activity of caspases

during reaper-triggered changes in tissue elasticity. Nat Commun. 2017;8: 603.

doi:10.1038/s41467-017-00693-3

8. Orme MH, Liccardi G, Moderau N, Feltham R, Wicky-John S, Tenev T, et al. The

unconventional myosin CRINKLED and its mammalian orthologue MYO7A regulate

caspases in their signalling roles. Nat Commun. 2016;7: 10972.

doi:10.1038/ncomms10972

833 9. Kanuka H, Kuranaga E, Takemoto K, Hiratou T, Okano H, Miura M. Drosophila caspase

834   transduces Shaggy/GSK-3β kinase activity in neural precursor development. EMBO J.

835   2005;24: 3793–3806. doi:10.1038/sj.emboj.7600822

836 10. Amcheslavsky A, Wang S, Fogarty CE, Lindblad JL, Fan Y, Bergmann A. Plasma

837   Membrane Localization of Apoptotic Caspases for Non-apoptotic Functions. Dev Cell.

838   2018;45: 450-464.e3. doi:10.1016/j.devcel.2018.04.020

839 11. Julien O, Wells JA. Caspases and their substrates. Cell Death Differ. 2017;24: 1380–

840   1389. doi:10.1038/cdd.2017.44

841 12. Kumar S. Caspase function in programmed cell death. Cell Death Differ. 2007;14: 32–

842   43. doi:10.1038/sj.cdd.4402060

843 13. Kumar S, Dorstyn L, Lim Y. The role of caspases as executioners of apoptosis. Biochem

844   Soc Trans. 2022;50: 33–45. doi: 10.1042/BST20210751

845 14. Branon TC, Bosch JA, Sanchez AD, Udeshi ND, Svinkina T, Carr SA, et al. Efficient

846   proximity labeling in living cells and organisms with TurboID. Nat Biotechnol. 2018;36:

847   880–887. doi:10.1038/nbt.4201

848 15. Qin W, Cho KF, Cavanagh PE, Ting AY. Deciphering molecular interactions by

849   proximity labeling. Nat Methods. 2021;18: 133–143. doi:10.1038/s41592-020-01010-5

850 16. Shinoda N, Hanawa N, Chihara T, Koto A, Miura M. Dronc-independent basal

851   executioner caspase activity sustains Drosophila imaginal tissue growth. Proc Natl Acad

852   Sci U S A. 2019;116:20539–20544. doi:10.1073/pnas.1904647116

853 17. Shinoda N, Horikoshi M, Taira Y, Muramoto M, Hirayama S, Murata S, et al. Caspase

854   cleaves Drosophila BubR1 to modulate spindle assembly checkpoint function and

855   lifespan of the organism. FEBS J. 2023;290:4147–4371. doi:10.1111/febs.16811

856 18. Chiba A, Snow P, Keshishian H, Hotta Y. Fasciclin III as a synaptic target recognition

857   molecule in Drosophila. Nature. 1995;374: 166–168. doi:10.1038/374166a0

19.   Kose H, Rose D, Zhu X, Chiba A. Homophilic synaptic target recognition mediated by

immunoglobulin-like cell adhesion molecule Fasciclin III. Development. 1997;124:

4143–4152. doi: 10.1242/dev.124.20.4143.

20.   Tang HL, Tang HM, Fung MC, Hardwick JM. In vivo Caspase tracker biosensor system

for detecting anastasis and non-apoptotic caspase activity. Sci Rep. 2015;5: 9015.

doi:10.1038/srep09015

21.   Ding AX, Sun G, Argaw YG, Wong JO, Easwaran S, Montell DJ. CasExpress reveals

widespread and diverse patterns of cell survival of caspase-3 activation during

development in vivo. Elife. 2016;5: e10936. doi:10.7554/eLife.10936

22.   Chihara T, Kitabayashi A, Morimoto M, Takeuchi K, Masuyama K, Tonoki A, et al.

Caspase Inhibition in Select Olfactory Neurons Restores Innate Attraction Behavior in

Aged Drosophila. PLoS Genet. 2014;10: e1004437. doi:10.1371/journal.pgen.1004437

23.   Hay BA, Wassarman DA, Rubin GM. Drosophila homologs of baculovirus inhibitor of

apoptosis proteins function to block cell death. Cell. 1995;83: 1253–1262.

doi:10.1016/0092-8674(95)90150-7

24.   Ditzel M, Broemer M, Tenev T, Bolduc C, Lee T V., Rigbolt KTG, et al. Inactivation of

Effector Caspases through Nondegradative Polyubiquitylation. Mol Cell. 2008;32: 540–

553. doi:10.1016/j.molcel.2008.09.025

25.   Lee T V., Fan Y, Wang S, Srivastava M, Broemer M, Meier P, et al. Drosophila IAP1-

mediated ubiquitylation controls activation of the initiator caspase DRONC independent

of protein degradation. PLoS Genet. 2011;7: e1002261.

doi:10.1371/journal.pgen.1002261

26.   Semmelhack JL, Wang JW. Select Drosophila glomeruli mediate innate olfactory

attraction and aversion. Nature. 2009;459: 218–223. doi:10.1038/nature07983

27.   Walsh JG, Cullen SP, Sheridan C, Lüthi AU, Gerner C, Martin SJ. Executioner caspase-

3 and caspase-7 are functionally distinct proteases. Proc Natl Acad Sci U S A. 2008;105:

12815–12819. doi:10.1073/pnas.0707715105

28.   Boucher D, Blais V, Denault JB. Caspase-7 uses an exosite to promote poly(ADP ribose)

polymerase 1 proteolysis. Proc Natl Acad Sci U S A. 2012;109: 5669–5674.

doi:10.1073/pnas.1200934109

29.   Desroches A, Denault JB. Caspase-7 uses RNA to enhance proteolysis of poly(ADP-

ribose) polymerase 1 and other RNA-binding proteins. Proc Natl Acad Sci U S A.

2019;116: 21521–21528. doi:10.1073/pnas.1909283116

30.   Takemoto K, Nagai T, Miyawaki A, Miura M. Spatio-temporal activation of caspase

revealed by indicator that is insensitive to environmental effects. J Cell Biol. 2003;160:

235–243. doi:10.1083/jcb.200207111

31.   Valente P, Lignani G, Medrihan L, Bosco F, Contestabile A, Lippiello P, et al. Cell

adhesion molecule L1 contributes to neuronal excitability regulating the function of

voltage-gated Na+ channels. J Cell Sci. 2016;129: 1878–1891. doi:10.1242/jcs.182089

32.   Mukherjee A, Williams DW. More alive than dead: Non-apoptotic roles for caspases in

neuronal development, plasticity and disease. Cell Death Differ. 2017;24: 1411–1421.

doi:10.1038/cdd.2017.64

33.   Dehkordi MH, Munn RGK, Fearnhead HO. Non-Canonical Roles of Apoptotic Caspases

in the Nervous System. Front Cell Dev Biol. 2022;10: 840023.

doi:10.3389/fcell.2022.840023

34.   Gu Q, Jiao S, Duan K, Wang YX, Petralia RS, Li Z. The BAD-BAX-Caspase-3 Cascade

Modulates Synaptic Vesicle Pools via Autophagy. J Neurosci. 2021;41: 1174–1190.

doi:10.1523/JNEUROSCI.0969-20.2020

35.   Cheriyamkunnel SJ, Rose S, Jacob PF, Blackburn LA, Glasgow S, Moorse J, et al. A

907   neuronal mechanism controlling the choice between feeding and sexual behaviors in

908   Drosophila. Curr Biol. 2021;31: 4231-4245.e4. doi:10.1016/j.cub.2021.07.029

909   36.   Dorstyn L, Colussi PA, Quinn LM, Richardson H, Kumar S. DRONC, an ecdysone-

910   inducible Drosophila caspase. Proc Natl Acad Sci U S A. 1999;96: 4307–4312.

911   doi:10.1073/pnas.96.8.4307

912   37.   Verghese S, Bedi S, Kango-Singh M. Hippo signalling controls Dronc activity to

913   regulate organ size in Drosophila. Cell Death Differ. 2012;19: 1664–1676.

914   doi:10.1038/cdd.2012.48

915   38.   Westphal D, Sytnyk V, Schachner M, Leshchyns'ka I. Clustering of the neural cell

916   adhesion molecule (NCAM) at the neuronal cell surface induces caspase-8- and -3-

917   dependent changes of the spectrin meshwork required for NCAM-mediated neurite

918   outgrowth. J Biol Chem. 2010;285: 42046–42057. doi:10.1074/jbc.M110.177147

919   39.   Luck K, Kim DK, Lambourne L, Spirohn K, Begg BE, Bian W, et al. A reference map

920   of the human binary protein interactome. Nature. 2020;580: 402–408.

921   doi:10.1038/s41586-020-2188-x

922   40.   Li L, Ding Z, Pang TL, Zhang B, Li CH, Liang AM, et al. Defective minor spliceosomes

923   induce SMA-associated phenotypes through sensitive intron-containing neural genes in

924   Drosophila. Nat Commun. 2020;11: 5608. doi:10.1038/s41467-020-19451-z

925   41.   Koto A, Kuranaga E, Miura M. Temporal regulation of Drosophila IAP1 determines

926   caspase functions in sensory organ development. J Cell Biol. 2009;187: 219–31.

927   doi:10.1083/jcb.200905110

928   42.   Karim MR, Endo K, Moore AW, Taniguchi H. Whole mount immunolabeling of

929   olfactory receptor neurons in the Drosophila antenna. J Vis Exp. 2014;51245.

930   doi:10.3791/51245

931   43.   Okuda S, Watanabe Y, Moriya Y, Kawano S, Yamamoto T, Matsumoto M, et al.

932   JPOSTrepo: An international standard data repository for proteomes. Nucleic Acids Res.

933   2017;45: D1107–D1111. doi:10.1093/nar/gkw1080

934   44.   Huang DW, Sherman BT, Lempicki RA. Bioinformatics enrichment tools: Paths toward

935   the comprehensive functional analysis of large gene lists. Nucleic Acids Res. 2009;37:

936   1–13. doi:10.1093/nar/gkn923

937   45.   Deluca SZ, Spradling AC. Efficient expression of genes in the drosophila germline using

938   a uas promoter free of interference by hsp70 pirnas. Genetics. 2018;209: 381–387.

939   doi:10.1534/genetics.118.300874

940   46.   Kashio S, Obata F, Zhang L, Katsuyama T, Chihara T, Miura M. Tissue nonautonomous

941   effects of fat body methionine metabolism on imaginal disc repair in Drosophila. Proc

942   Natl Acad Sci U S A. 2016;113: 1835–1840. doi:10.1073/pnas.1523681113

Fig 1. Expression patterns and proximal proteins of Drice in the adult brain.

(A) Schematic diagram of *Drosophila* apoptosis signaling. (B) Western blot of expression of

caspases in adult male heads. (C) Representative images of adult male brains. Expression

patterns of each caspase are visualized using Streptavidin (SA, magenta). Nuclei are visualized

by Hoechst33342 (Green). Scale bar: 50 µm. (D) Representative images of antennal lobes (ALs)

in the adult female brains. Expression patterns of Drice are visualized using SA (magenta).

Nuclei are visualized by Hoechst33342 (Green). Scale bar: 50 µm. (E) Western blot of proteins

extracted from adult male brains. (F) Schematic diagram of the TurboID-mediated

identification of proximal proteins. (G) A summary of mass spectrometry analysis of proteins

of the adult male brains. Partial list of Drice proximal proteins and their Fold Change (FC) are

shown. (H) Gene Ontology analysis (Cellular Compartment) of 158 Drice proximal proteins

(*Drice::V5::TurboID/w^{1118}* > 10). Cyan: neural-related fraction, Red: membrane fraction.

Fig 2. A specific isoform of Fasciclin 3 is in proximity to Drice.

(A) Representative expression patterns of Drice (Streptavidin (SA); magenta) and Fasciclin 3 (Fas3, anti-Fasciclin 3 antibody staining; green) in the adult male brains. Scale bar: 50 µm. (B) Representative expression patterns of Drice (SA; magenta) and Fas3 (anti-GFP antibody staining; green) in the antennal lobes (ALs) of the adult male brains. Scale bar: 50 µm. (C) Schematic protein structures of Fas3 isoforms. Intracellular regions differ from each other. A peptide region used to raise anti-Fas3G antibody is shown. (D) IDR predictions of intracellular regions of each protein isoform. (E) Western blot of proteins extracted from adult heads. Anti-Fas3G antibody specifically detects Fas3 isoform G. (F) Western blot of adult male heads. Drice

968   proximal proteins were purified using NeturAvidin. (G) Western blot of adult head proteins.

969   Drice proximal proteins were purified using NeturAvidin.

Fig 3. A Gal4-Manipulated Area Specific CaspaseTracker/CasExpress (MASCaT).

(A) A schematic diagram of MASCaT expressed using the Gal4/UAS system with FLP-mediated recombination. Caspase sensitive-QF2 is activated by caspase-mediated cleavage at the plasma membrane. (B) Representative images of S2 cells expressing caspase sensitive-QF2 probes by the Gal4/UAS system with FLP-mediated recombination. Scale bar: 20 μm. (C) Representative images of adult male brains of young (1 week old) and old (3 weeks old) flies raised at 29°C. Arrowheads indicate optic lobes. Magnified images (white rectangle) show

antennal lobes (ALs; white circles). Arrows indicate age-dependent caspase activation at the

ALs. Scale bar: 100 μm.

Fig 4. *Fasciclin 3 isoform G* overexpression enhances non-lethal caspase activation.

(A) Representative images of the adult male antennae of 1 weeks old flies raised at 29°C. Cell

bodies are visualized using RedStinger (red). Scale bar: 20 μm. (B) Quantification of cell

number of (A). Data are presented as mean ± SEM. *P*-values were calculated using one-way

analysis of variance (ANOVA) with Bonferroni's correction for selected pairs. NS: $P > 0.05$,

†: $P < 0.05$. (C) Representative images of the adult male antennae of 1 week old flies raised at

29°C. Cell bodies are visualized using RedStinger (red). Scale bar: 20 μm. (D) Quantification

of cell number of (C). Data are presented as mean ± SEM. *P*-values were calculated using one-

way ANOVA with Bonferroni's correction for every pair. NS: $P > 0.05$, †: $P < 0.05$. (E)

Representative images of the antennal lobes (ALs) of male brains of 1 week old flies raised at

29°C expressing MASCaT probe (mCD8). Scale bar: 30 μm. (F) Representative images of the

ALs of brains of 1 week old flies raised at 29°C with caspase activation visualized by MASCaT.

Scale bar: 30 μm. (G) Quantifications of caspase activity detected by MASCaT of (F). *P*-values

were calculated using chi-square test with Bonferroni's correction using mNeonGreen as

control. NS: $P > 0.05$, †: $P < 0.05$. Sample sizes are shown in the graph. (H) Representative

images of the ALs of brains of 1 week old flies raised at 29°C with caspase activation visualized

by MASCaT. Scale bar: 30 μm. (I) Quantifications of caspase activity detected by MASCaT of

(H). *P*-values were calculated using chi-square test with Bonferroni's correction using

mNeonGreen and Fas3G as controls, respectively. NS: $P > 0.05$, †: $P < 0.05$. Sample sizes are

shown in the graph.

1001

Fig 5. *Fasciclin 3 isoform G* overexpression enhances non-lethal caspase activation in a

Dronc-dependent manner.

(A) Representative images of adult male brains. Expression patterns of Dronc are visualized using Streptavidin (SA, magenta). Magnified images (white rectangle) show antennal lobes (ALs). Scale bar: 50 µm. (B) Western blot of adult male heads. Dronc proximal proteins were purified using NeturAvidin. (C) Representative images of the AL of brains of 1 week old flies raised at 29°C with caspase activation visualized using MASCaT. Scale bar: 30 µm. (D) Quantifications of caspase activity detected using MASCaT of (C). P-values were calculated using chi-square test with Bonferroni's correction using mNeonGreen and Fas3G as controls, respectively. NS: $P > 0.05$, †: $P < 0.05$. Sample sizes are shown in the graph. (E) Western blot of adult male heads. Drice proximal proteins were purified using NeturAvidin. (F) A schematic model of subcellularly restricted non-lethal caspase activation upon Fas3G overexpression.

Fig 6. *Fasciclin 3 isoform G* overexpression-facilitated caspase activation regulates attraction behavior.

(A) Schematic diagram of two choice preference assay. Flies are starved for 4 hours before the assay. Flies are left for 2 hours in the bottle. Preference index is calculated by counting the number of flies left in MQ, ACV and other (field) compartments. (B) Preference index of 1 week old male flies raised at 29°C in response to 1% apple cider vinegar in two choice preference assay. Data are presented as mean ± SEM. *P*-values were calculated using one-way ANOVA followed by Dunnets' multiple comparison test using mNeonGreen as a control. NS: *P* > 0.05, †: *P* < 0.05. Sample sizes: mNG (N = 19 (1327 flies)), Fas3G (N = 17 (1191 flies)), Fas3G + Dronc-RNAi (N = 19 (1295 flies)), Fas3G + Dronc^{DN} (N = 20 (1320 flies)). (C) The ratio of flies in the given conditions in the two choice preference assay of (A). *P*-values were calculated using chi-square test with Bonferroni's correction using mNeonGreen as control. †: *P* < 0.05. Sample sizes: mNG (N = 1327 flies), Fas3G (N = 1191 flies), Fas3G + Dronc-RNAi (N = 1295 flies), Fas3G + Dronc^{DN} (N = 1320 flies).

1029 **Legends for Figure S1–4**

1030

1031 **Fig S1. Expression patterns of Drice in the adult brain in males and females.**

1032 (A) Representative images of adult male and female brains. Expression patterns of Drice are

1033 visualized using Streptavidin (SA, magenta). Nuclei are visualized by Hoechst33342 (Green).

1034 Scale bar: 50 μm.

1035

Fig S2. Biochemical property of Fas3s with caspases.

(A) Co-immunoprecipitation of 3xFLAG-tagged Fas3s with V5-tagged Drice extracted from
adult male heads. (B) Western blot for 3xFLAG-tagged Fas3s expressed in *Drosophila* S2 cells.
Apoptosis was induced by treating S2 cells with cycloheximide (CHX).

1040

Fig S3. Overexpression of *mCD8::DQVD::QF2* probes in *Drosophila* S2 cells.

(A) A schematic diagram of MASCaT expressed directly using the Gal4/UAS system. Caspase

sensitive-QF2 is activated by caspase-mediated cleavage at the plasma membrane. (B)

Representative images of S2 cells expressing caspase sensitive-QF2 probes directly by the

Gal4/UAS system. A pan-caspase inhibitior, zVAD-fmk, is treated to inhibit caspase activity.

Scale bar: 20 μm.

1047

Fig S4. Loss of function analysis of Fas3G.

(A) Gene structure of Fas3. CDS (orange) and UTR (grey) are shown in boxes. Target sequence of Fas3G-shRNA is shown in red. (B) Western blot for 3xFLAG-tagged Fas3s expressed in S2 cells with Fas3G-shRNA. (C) Representative images of the ALs of brains of 1 week old flies raised at 29°C with caspase activation visualized by MASCaT. Scale bar: 30 μm. (D) Quantifications of caspase activity detected by MASCaT of (C). *P*-values were calculated using chi-square test with Bonferroni's correction using *LacZ-RNAi* as control. NS: *P* > 0.05. Sample sizes are shown in the graph. (E) Representative images of the ALs of brains of 1 week old flies expressing mNeonGreen-tagged Drice (green) and 3xFLAG-tagged Fas3G (magenta). Scale bar: 50 μm.

1058 **Legends for Table S1–5**

1059 **Table S1. Proximal protein list.** A list of the identified proximal proteins (Abundance ratio

1060 (Drice::V5::TurboID/Control) > 10). The UniProt Accessions, Gene Description, and

1061 Abundance Ratio (Drice::V5::TurboID/Control) are listed.

1062

1063 **Table S2. Gene Ontology analysis.** The result of Gene Ontology (GO) analysis of Cellular

1064 Compartment for Drice proximal proteins. GO term, P values, Fold Enrichment, Count and

1065 Genes are listed.

1066

1067 **Table S3. LC settings.** Gradient settings for LC analysis are presented.

1068

1069 **Table S4. MS/MS settings.** Settings used for the MS/MS analysis are presented.

1070

1071 **Table S5. Proteome Discoverer 2.2 settings.** Proteome Discoverer 2.2 settings for proteomics

1072 analysis are presented.