Title

Executioner caspase is proximal to Fasciclin 3 which facilitates non-lethal activation in *Drosophila* olfactory receptor neurons

Short title

Fasciclin 3 non-lethally activates caspase in *Drosophila* olfactory neurons

Authors

Masaya Muramoto[a,1], Nozomi Hanawa[a,1], Misako Okumura[b,c], Takahiro Chihara[b,c], Masayuki Miura[a,2], and Natsuki Shinoda[a,2]

Affiliations & Footnotes

[a]Department of Genetics, Graduate School of Pharmaceutical Sciences, University of Tokyo, 7-3-1 Hongo, Bunkyo-ku, Tokyo 113-0033, Japan

[b]Department of Biological Science, Graduate School of Science, Hiroshima University, Higashi-Hiroshima, Hiroshima 739-8526, Japan

[c]Program of Biomedical Science and Basic Biology, Graduate School of Integrated Sciences for Life, Hiroshima University, Higashi-Hiroshima, Hiroshima 739-8526, Japan

[1]Equal contributions

[2]Corresponding authors

Corresponding Authors

Masayuki Miura: Department of Genetics, Graduate School of Pharmaceutical Sciences, University of Tokyo, 7-3-1 Hongo, Bunkyo-ku, Tokyo 113-0033, Japan. Telephone: +81-3-5841-4860. E-mail: miura@mol.f.u-tokyo.ac.jp

Natsuki Shinoda: Department of Genetics, Graduate School of Pharmaceutical Sciences,

University of Tokyo 7-3-1 Hongo, Bunkyo-ku, Tokyo 113-0033, Japan. Telephone: +81-3-

5841-4867. E-mail: f-shinoda.natsuki@g.ecc.u-tokyo.ac.jp

Author Contributions

Masaya Muramoto: Formal Analysis, Investigation, Methodology, Resources, Writing –

Review & Editing; Nozomi Hanawa: Formal analysis, Investigation, Methodology, Resources,

Writing – Review & Editing; Misako Okumura: Methodology, Writing – Review & Editing;

Takahiro Chihara: Methodology, Writing – Review & Editing; Masayuki Miura:

Conceptualization, Funding Acquisition, Supervision, Writing – Original Draft Preparation;

Natsuki Shinoda: Conceptualization, Formal Analysis, Funding Acquisition, Investigation,

Project Administration, Supervision, Validation, Visualization, Writing – Original Draft

Preparation

Conflict of Interest

The authors declare that they have no conflict of interest.

Abstract

Executioner caspases are activated by initiator caspases, resulting in lethal and non-lethal

cellular outcomes. Despite the identification of numerous caspase-dependent non-lethal cellular

processes (CDPs), the molecular mechanism that enables precise regulation of the non-lethal

functions of executioner caspases remains unclear. Here, we conducted a comprehensive

analysis of endogenous proximal proteins of *Drosophila* executioner caspases *in vivo* using

TurboID. We discovered that in the adult brain, the executioner caspase Drice is primarily

proximal to membrane proteins, including the cell adhesion molecule Fasciclin 3 (Fas3).

Notably, Drice is proximal to the specific alternative splicing isoforms of Fas3G. To investigate

whether Fas3 modulates caspase activity, we developed a Gal4-Manipulated Area-Specific

CaspaseTracker/CasExpress (MASCaT) system for highly sensitive monitoring of non-lethal

caspase activity near membrane compartments. Using MASCaT, we demonstrated that Fas3

promoted caspase activation near membrane compartments in olfactory receptor neurons

without killing them. Activation regulates innate olfactory attraction behavior. Collectively, our

findings suggest that the proximal proteins of executioner caspases provide a

compartmentalized platform for non-lethal caspase activation and function.

Author Summary

Caspases are Cysteine-ASPartic acid proteASEs which are primarily involved in cell death. In

addition, caspases play versatile non-lethal (non-cell death) roles in a wide range of cell types

including neurons. However, the molecular mechanism that enables the precise regulation of

non-lethal activation and functions remains unclear. Here, we focused on the proximal proteins

of caspases as potential regulators of non-lethal activation. By comprehensively analyzing the

proximal proteins of caspases in the adult *Drosophila* brain using TurboID, we discovered that

caspases are primarily proximal to membrane proteins rather than to cytosolic proteins. By

developing a genetic tool that enables highly sensitive monitoring of non-lethal caspase activity

near membrane compartments, we found that proximal proteins facilitate non-lethal caspase

activation in olfactory receptor neurons. Importantly, non-lethal caspase activation in olfactory

receptor neurons affects organismal innate attraction behavior. Our research suggests that a

compartmentalized platform established by proximal proteins enables precise regulation of

non-lethal caspase activation and function.

Introduction

Caspases are cysteine aspartic acid proteases involved in cell death. Apoptotic stimuli activate initiator caspases that trigger a proteolytic cascade, leading to the activation of executioner caspases and culminating in apoptosis (Fig. 1A) [1]. Caspases exhibit diverse non-lethal cellular functions [2]. In the nervous system, non-lethal caspase activity promotes the maturation of olfactory sensory neurons during development [3]. Furthermore, non-lethal caspase activity induces long-term depression through AKT cleavage [4]. These are termed caspase-dependent non-lethal cellular processes (CDPs) [5]. Despite the discovery of numerous CDPs over the past two decades, the molecular mechanism that underlies specific non-lethal functions without causing cell death remains unclear.

Proteins that interact with initiator caspases are among the key regulators of CDPs. Tango7, the *Drosophila* translation initiation factor eIF3 subunit m, acts as an adapter and regulates the localization of the initiator caspase Dronc. During sperm individualization, Tango7 directs Dronc to the individualization complex [6]. In salivary glands, Tango7 directs Dronc to the plasma membrane [7]. In both cases, caspase activity promotes cellular remodeling without inducing cell death [6,7]. CRINKLED, a *Drosophila* unconventional myosin, functions as an adaptor of Dronc and facilitates the cleavage of Shaggy46 [8]. Shaggy46, activated by caspase-mediated cleavage, suppresses the emergence of ectopic sensory organ precursors [9]. During apoptosis-induced proliferation (AiP), Myo1D, another *Drosophila* unconventional myosin, facilitates Dronc translocation to the basal side of the plasma membrane of imaginal discs to promote AiP [10]. Thus, initiator caspase-interacting proteins regulate caspase activity in CDPs. However, given the importance of executioner caspases, which cleave a broad range of substrates and are essential for apoptosis [11,12], little is known about the subcellular compartmentalization and interacting proteins of those involved in CDPs. This lack of

understanding is partly due to the absence of protein-protein interaction domains (e.g., DED

and CARD) in the N-terminus of executioner caspases [13].

To understand the molecular mechanism of function of caspases involved in CDPs, we

focused on the proximal proteins of executioner caspases using TurboID, a proximity labeling

technique that analyzes protein proximity *in vivo* [14,15]. We previously established C-terminal

TurboID knocked-in caspase fly lines for Dronc, Drice, and Dcp-1 [16]. Importantly, we have

demonstrated that one of the preferred substrates of Drice, BubR1, is proximal to Drice *in vivo*

[17], suggesting that substrate preference is exploited by protein proximity. Therefore, the

proximity of executioner caspases to proteins is emerging as a means of regulating specific non-

lethal functions.

In this study, through a comprehensive analysis of proximal proteins, we discovered

that the cell adhesion molecule Fasciclin 3 (Fas3) is in close proximity to the executioner

caspase Drice in the adult brain. We observed that Drice was exclusively proximal to specific

alternative splicing isoforms of Fas3. To ascertain whether Fas3 modulates caspase activity, we

developed a Gal4-Manipulated Area Specific CaspaseTracker/CasExpress (MASCaT) system,

which permits the monitoring of caspase activity with high sensitivity and simultaneous genetic

manipulation in the cells of interest. Using MASCaT, we demonstrated that Fas3 enhances

caspase activation without killing cells. Subsequently, we showed that activation regulates

innate olfactory attraction behavior. Collectively, our findings suggest that the proximal

proteins of executioner caspases provide a compartmentalization platform that enables the

precise regulation of non-lethal caspase activation and function.

Results

Drice is the major executioner caspase expressed in the adult brains

In *Drosophila*, the caspase-mediated apoptotic signaling pathway is highly conserved [2]. The activity of executioner caspases, specifically Drice and Dcp-1, is regulated by apoptotic signaling pathways, including the initiator caspase Dronc (Fig. 1A). To investigate the expression patterns of *Drosophila* caspases, we employed *Caspase::V5::TurboID* knocked-in fly lines [16]. We found low levels of expression of the initiator caspase Dronc in adult heads (Fig. 1B). Conversely, we observed high expression of the executioner caspase Drice, whereas Dcp-1 was expressed at lower levels in adult heads (Fig. 1B). The proximal proteins of each caspase were labeled with TurboID in adult heads by administration of 100 µM biotin (Fig. 1B). Furthermore, we investigated the expression patterns of caspases using histochemical analysis by examining streptavidin (SA) signal. Consistent with the western blotting findings, Drice showed higher expression levels than Dronc and Dcp-1, which were expressed at lower levels (Fig. 1C). We found that Drice was expressed in distinct regions, including the mushroom bodies (MBs), suboesophageal ganglions (SGs), optic lobes (OLs), and antennal lobes (ALs), which showed the highest levels of expression (Fig. 1C). The *Drosophila* AL comprise synaptic contacts between the axons of olfactory receptor neurons (ORNs), the dendrites of projection neurons (PNs), and local interneuron (LNs) processes. Neuron-type-specific Gal4s were used to knockdown *Drice* and the expression patterns were subsequently examined. We found that Drice was expressed predominantly in ORNs in the AL (Fig. 1D). These results suggest that Drice is the major executioner caspase that regulates both lethal and non-lethal functions of adult ORNs.

Drice is proximal to membrane proteins rather than cytosolic proteins in the adult brains

To comprehensively identify proximal proteins of Drice, we employed a proximity labeling

approach. Specifically, we used *Drice::V5::TurboID* flies fed 100 µM biotin. We confirmed

biotinylation of Drice proximal proteins in the adult brain (Fig. 1E). After purifying the

biotinylated proteins using NeutrAvidin magnetic beads, we performed LC-MS/MS analysis of

the samples (Fig. 1F). We identified 643 proteins (Fig. 1G, Table S1), of which 158 were highly

specific to *Drice::V5::TurboID* flies compared to wild-type flies (*Drice::V5::TurboID*/w^{1118} >

10; Fig. 1G, Table S1). Gene ontology analysis revealed that Drice proximal proteins were

enriched in the plasma membrane, synaptic vesicles, postsynaptic membranes, and adherens

junctions (Fig. 1H, Table S2). This finding suggests that Drice is localized primarily in

membrane compartments rather than in the cytoplasm of the adult brain. Additionally, because

the SA signal of *Drice::V5::TurboID* is mainly observed in the AL where the axons of ORNs

project (Fig. 1C), the enrichment of membrane proteins in the proximal proteins of Drice is

reasonable.

A specific splicing isoform of Fasciclin 3 is in proximity to Drice

Among the identified neighboring proteins of Drice, we focused on Fasciclin 3 (Fas3) [18], a

transmembrane protein containing immunoglobulin-like domains that regulates synaptic target

recognition and axon fasciculation [19]. The role of Fas3 in apoptosis remains unclear. The

expression patterns of Fas3 in the adult brain were similar to those of Drice, particularly in the

OLs, SGs, and ALs (Fig. 2A, B). Fas3 is encoded by seven splicing isoforms, resulting in the

production of five protein isoforms that share extracellular domains but differ in intracellular

domains with low-complexity sequences (Fig. 2C-E). Only one specific protein isoform, Fas3

isoform G (Fas3G), was identified in the LC-MS/MS analysis as a Drice proximal protein in

the adult brain (Fig. 1G, Table S1). To validate this result, we generated an antibody specific

to Fas3G (Fig. 2D, F). We confirmed that only the Fas3G was labeled with *Drice::V5::TurboID*

and purified using NeutrAvidin magnetic beads as detected using western blotting (Fig. 2G).

To further confirm the isoform specificity to Drice proximity, we overexpressed 3xFLAG-

tagged Fas3 isoforms A, B/F, C, D/E, and G in olfactory receptor neurons using *Pebbled-Gal4*

in a *Drice::V5::TurboID* background. After proximity labeling with TurboID, followed by

NeutrAvidin purification, we found that the Fas3G was highly enriched compared to the other

isoforms (Fig. 2H). These findings suggested that Drice is proximal to a specific isoform of

Fas3 in olfactory receptor neurons, regardless of its protein expression level.

MASCaT: a cell type-specific highly sensitive non-lethal caspase activity reporter

We investigated whether Fas3 regulates caspase activity in lethal or non-lethal processes.

CaspaseTracker/CasExpress is a highly sensitive reporter of non-lethal caspase activity that

uses caspase-activatable Gal4 tethered to the membrane by mCD8 [20,21]. However, reliance

on Gal4 to report their activity presents a challenge in reconciling Gal4/UAS-dependent gene

manipulations. To overcome this limitation, we developed a novel reporter that captures non-

lethal caspase activation independently of the Gal4/UAS system. Specifically, we replaced

caspase-activatable Gal4 with caspase-activatable QF2 (mCD8::DQVD::QF2), where DQVD

is a caspase cleavage site, and generated *Ubi-FRT-STOP-FRT-mCD8::DQVD::QF2*,

simultaneously expressing *Flippase* (*FLP*) under *UAS* to restrict reporter expression to the cells

of interest (Fig. 3A). As a negative control, we created caspase-non-activatable QF2

(mCD8::DQVA::QF2). *Drosophila* S2 cells undergo weak caspase activation only by

transfection [17]. Taking advantage of that, this system reports caspase activity through the

caspase-activatable QF2/QUAS system, expressing mNeonGreen, which is not detected by

caspase-non-activatable QF2 (Fig. 3B). We named this novel reporter a Gal4-Manipulated Area

Specific CaspaseTracker/CasExpress (MASCaT; Fig. 3A). Utilizing MASCaT, we detected

non-lethal caspase activity, specifically in the neurons of adult brains, using *ELAV-Gal4* (Fig.

3C). Caspase activity was prominent in OLs (Fig. 3C), which is consistent with the previously

reported activation pattern of caspase detected by CaspaseTracker/CasExpress [20,21].

Previously, we reported that a specific subset of ORNs, including Or42b, undergoes age-

associated caspase activation [22]. Consistent with this finding, we detected age-dependent

caspase activation in the AL (Fig. 3C). Therefore, our newly established reporter, MASCaT,

enables the highly sensitive detection of caspase activity near the membrane in cells of interest

in a Gal4/UAS-dependent manner.

Fas3G activates caspase in a non-lethal manner that regulates innate attraction behavior

To investigate the regulatory role of Fas3 in caspase activity *in vivo*, we used Or42b neurons as

a model, which have previously been shown to undergo age-dependent caspase activation [22].

Initially, we evaluated the effects of Dronc overexpression on apoptosis by quantifying cell

body numbers in the antenna in young flies. Dronc induced cell loss while catalytically inactive

Dronc (Dronc^{CG}) did not, suggesting that Dronc induce apoptosis in a catalytic-dependent

manner (Fig. 4A, B). We then evaluated the effects of the overexpression of the executioner

caspases Dcp-1 and Drice. Interestingly, although Dcp-1 induced complete cell loss, Drice did

not (Fig. 4A, B). These results indicate that Drice requires an initiator caspase for activation,

whereas Dcp-1 does not (Fig. 4A, B). Furthermore, the expression of Fas3G, even in

combination with Drice, did not reduce the number of cells, implying that Fas3G did not induce

apoptosis (Fig. 4C, D).

Subsequently, we assessed the regulation of non-lethal caspase activity by Fas3G *in*

vivo. We first confirmed that MASCaT probes were distributed in the axons of olfactory

receptor neurons (Fig. 4E). Using MASCaT, we observed weak non-lethal caspase activation

even in the control group at a young age (Fig. 4F, G). The expression of *Fas3G* significantly

promoted caspase activation (Fig. 4F, G), indicating that Fas3G regulates non-lethal caspase

activation. Moreover, the simultaneous expression of Drice and Fas3G significantly enhanced

caspase activation, signifying the promoting effect of their proximity (Fig. 4F, G). Additionally,

knockdown or expression of the dominant-negative form of Dronc suppressed Fas3G-

facilitated non-lethal caspase activation (Fig. 4H, I), suggesting that this activation is initiator

caspase-dependent. Thus, our findings indicate that the proximal proteins of executioner

caspases facilitate non-lethal caspase activation in close proximity to the membrane (Fig. 4J).

Finally, we examined the function of by Fas3G-facilitated non-lethal caspase activation.

Apple cider vinegar (ACV) excites six glomeruli, including DM1, which are innervated by the

axons of Or42b neurons [23]. We have previously reported that Or42b neurons undergo aging-

dependent caspase activation, which reduces neural activity and attraction behavior in response

to ACV [22]. Inhibition of caspase activity in aged flies restores attraction behavior but does

not affect that of young flies [22]. Using a two-choice preference assay with ACV, we found

that the expression of Fas3G, which activates caspases, did not impair attraction behavior in

young flies (Fig. 4K, L). In contrast, simultaneous inhibition of Fas3G-facilitated non-lethal

caspase activation promoted attraction to ACV (Fig. 4K, L). Overall, these results suggest that

Fas3G-facilitated non-lethal caspase activation suppresses innate attractive behavior without

killing the cells.

Discussion

Proximal proteins of executioner caspases regulate non-lethal caspase activity

It is important to identify the proximal proteins of executioner caspases, which cleave a wide range of substrates and are essential for apoptosis [11,12], as proximity often dictates substrate specificity [17]. In humans, Caspase-3 and Caspase-7 have been shown to share but also have discrete substrate preferences [24]. Caspase-7 leverages an exosite to facilitate interaction with RNA and enhance the proteolysis of RNA-binding proteins [25,26]. Thus, protein proximity can be used to precisely regulate specific cellular functions by selectively cleaving a limited pool of substrates in non-lethal scenarios. Our current investigation showed that executioner caspases are exclusively sequestered in specific subcellular domains, including the plasma membrane. We identified Fas3G as a regulator of non-lethal activation, underscoring the importance of protein proximity not only in substrate cleavage, but also in localizing non-lethal activation to the vicinity of plasma membrane (Fig. 4J).

MASCaT as a highly sensitive reporter for identifying non-lethal caspase regulators

Various caspase activity reporters have been developed, including SCAT3, a FRET-based reporter with superior temporal resolution [27], CaspaseTracker/CasExpress, Caspase-activatable Gal4-based reporters with superior sensitivity [20,21]. However, to date, no reporter has been developed that is sensitive enough to identify non-lethal regulators of caspases in cells of interest. In this study, we developed a novel caspase reporter, MASCaT, that can detect non-lethal caspase activation with high sensitivity in a cell type-specific manner following gene manipulation. Using MASCaT, we identified previously overlooked regulators of non-lethal activation of caspases in a cell type-specific manner. Proximal proteins of executioner caspases are potential non-lethal regulators, and combining TurboID-mediated identification of these

proteins will pave the way for identifying non-lethal components and broaden our

understanding of the molecular mechanism of non-lethal caspase activation.

Non-lethal caspase activation regulates neural activity and behaviors

We previously reported that age-dependent caspase activation reduces neural activity and innate

attraction behavior [22]. Here, we found that, while Fas3G expression did not impair attraction

behavior by itself, caspase activation upon Fas3G overexpression suppressed attraction

behavior toward ACV (Fig. 4K, L). In mice, L1CAM, an immunoglobulin-like cell adhesion

molecule, facilitates neuronal excitability by regulating the voltage-gated Na+ channels [28].

Thus, we hypothesize that non-lethal caspase activation antagonizes the neuronal excitability

promoted by Fas3G. In neurons, caspases have diverse non-lethal functions, including axon

degeneration and macro- and micro-pruning, to regulate synaptic plasticity [29,30]. In

presynaptic terminals, it has been recently reported that Caspase-3 modulates synaptic vesicle

pools via autophagy [31]. Thus, it is possible that non-lethal activation regulates synaptic pools

and neural activity in a reversible manner without killing the cells. Analyzing the functions

mediated by Drice/Fas3G proximity, including the identification of specific substrates, will

pave the way for identifying diverse CDPs and understanding their molecular mechanisms.

While suppressing innate attraction behavior does not seem beneficial by itself, it should

be a great advantage in decision-making. To survive, flies must constantly make appropriate

behavioral decisions (e.g., feeding versus mating) depending on their physiological conditions.

In males, a subset of neurons regulated by tyramine controls feeding versus mating behaviors

[32]. In regard of that, it is important to adequately promote/suppress neural activity, which

controls behaviors of the organisms, by precisely controlling caspase activation in a

compartmentalization-dependent manner in neurons.

How does Fasciclin 3 regulate non-lethal caspase activation?

Although the core apoptotic machinery has been well characterized during the last 30 years [2],

the precise regulation of CDPs, remains incompletely elucidated. In this study, we identified

Fas3 as a context-specific activator of Drice. However, the mechanism by which Fas3G

regulates non-lethal activation is still an unresolved question in our research. Notably, Fas3G

was not a determinant of apoptosis, since overexpression of Fas3G did not reduce the number

of ORNs (Fig. 4C, D). Fas3 is an immunoglobulin-like domain-containing transmembrane

protein that regulates axonal fasciculation and neuronal projections [18]. NCAM, a mammalian

homologue of Fascicilin 2, is an immunoglobulin-like domain-containing cell adhesion

molecule. NCAM can physically bind to Caspase-8, and depending on NCAM clustering,

Caspase-8 and -3 are activated and contribute to neurite outgrowth by cleaving its substrate

spectrin [33]. Thus, it is possible that Fas3G physically binds to the initiator caspase Dronc,

which is required for Fas3G-facilitated non-lethal caspase, to activate Drice.

Recently, a protein interactome analysis in humans showed that tissue-preferentially

expressed proteins that interact with known apoptosis regulators participate in apoptosis

sensitization [34]. Given that CDPs are often cell-and tissue type-specific processes, the

identification of modulators of context-dependent apoptotic pathways will aid in understanding

the precise molecular regulation of CDPs. Importantly, alternative splicing can diversify

protein-protein interaction networks. A recent report showed that minor spliceosomes regulate

alternative splicing, including Fas3. A specific isoform of Fas3 has distinct functions and cannot

be substituted by other isoforms during neuronal development [35]. Our results support the idea

that alternative splicing may not only diversify the protein itself, but also its interaction

networks to regulate module-dependent activation. Further interactome analysis using TurboID

will reveal the regulatory mechanisms of the non-lethal activation of caspases.

Materials and Methods

Fly strains and rearing conditions

We raised the flies in standard *Drosophila* medium at 25°C. For in vivo biotin labeling experiments, flies were raised in standard *Drosophila* medium supplemented with 100 μM (+)-biotin (#132-02635, WAKO). The following fly strains were used in this study: w^{1118}, *Drice::V5::TurboID*, *Dcp-1::V5::TurboID*, *Dronc::V5::TurboID* [16], *Pebbled-Gal4* (BL80570), *LN-Gal4* (BL63325), *GH146-Gal4* (BL30026), *UAS-Drice-RNAi* (V28065), *Fas3::EGFP::FlAsH::StrepII::TEV::3xFLAG* (BL59809), *UASz-Fas3A::3xFLAG*, *UASz-Fas3B::3xFLAG*, *UASz-Fas3C::3xFLAG*, *UASz-Fas3D::3xFLAG*, *UASz-Fas3G::3xFLAG* (this study), *Ubi-FRT-STOP-FRT-mCD8::DQVD::QF2*, *Ubi-FRT-STOP-FRT-mCD8::DQVA::QF2* (this study), *UAS-FLP* (K107788), *QUAS-tdTomato::3xHA* (BL30005), *ELAV-Gal4* (BL8765), *Or42b-Gal4* (BL9971), *UAS-RedStinger* (BL8547), *UASz-myc::mNeonGreen*, *UASz-Drice::myc::mNeonGreen* (this study), *UASz-V5::TurboID*, *UASz-Drice::V5::TurboID*, *UASz-Drice^{CG}::V5::TurboID*, *UASz-Dcp-1::V5::TurboID*, *UASz-Dcp-1^{CG}::V5::TurboID*, *UASz-Dronc::V5::TurboID*, *UASz-Dronc^{CG}::V5::TurboID* (this study), *UAS-Dronc-RNAi* [9], *UAS-Dronc^{DN}* (a gift from H. Richardson). BL, Bloomington Drosophila Resource Center; K, Kyoto Stock Center; V, Vienna Drosophila Resource Center.

Detailed genotypes

Fig. 1B, C (Left to Right, Top to Bottom)

w^{1118}/Y

w/Y; Dcp-1::V5::TurboID/+

w/Y;; Drice::V5::TurboID/+

w/Y;; Dronc::V5::TurboID/+

Fig. 1D (Left to Right)

w; UAS-Drice-RNAi/+; Drice::V5::TurboID/+

w, Pebbled-Gal4/w; UAS-Drice-RNAi/+; Drice::V5::TurboID/+

w; UAS-Drice-RNAi/GH146-Gal4; Drice::V5::TurboID/+

w; UAS-Drice-RNAi/+; Drice::V5::TurboID/LN-Gal4

Fig. 1E (Left to Right)

w^{1118}/Y

w/Y;; Drice::V5::TurboID/+

Fig. 1G (Left to Right)

w^{1118}/Y

w/Y;; Drice::V5::TurboID/+

Fig. 2A (Left to Right)

w^{1118}/Y

w/Y;; Drice::V5::TurboID/+

Fig. 2B (Top to Bottom, Left to Right)

w^{1118}/Y

w/Y;; Drice::V5::TurboID/+

w/Y; Fas3::EGFP::FlAsH::StrepII::TEV::3xFLAG/+

w/Y; Fas3:: EGFP::FlAsH::StrepII::TEV::3xFLAG/+; Drice::V5::TurboID/+

Fig. 2F (Left to Right)

w, Pebbled-Gal4/w or Y

w, Pebbled-Gal4/w or Y;; UASz-Fas3A::3xFLAG/+

w, Pebbled-Gal4/w or Y;; UASz-Fas3B::3xFLAG/+

w, Pebbled-Gal4/w or Y;; UASz-Fas3C::3xFLAG/+

w, Pebbled-Gal4/w or Y;; UASz-Fas3D::3xFLAG/+

w, Pebbled-Gal4/w or Y;; UASz-Fas3G::3xFLAG/+

Fig. 2G (Left to Right)

w^{1118}/Y

w/Y;; Drice::V5::TurboID/+

Fig. 2H (Left to Right)

w, Pebbled-Gal4/w or Y;; Drice::V5::TurboID/+

w, Pebbled-Gal4/w or Y;; Drice::V5::TurboID/UASz-Fas3A::3xFLAG

w, Pebbled-Gal4/w or Y;; Drice::V5::TurboID/UASz-Fas3B::3xFLAG

w, Pebbled-Gal4/w or Y;; Drice::V5::TurboID/UASz-Fas3C::3xFLAG

w, Pebbled-Gal4/w or Y;; Drice::V5::TurboID/UASz-Fas3D::3xFLAG

w, Pebbled-Gal4/w or Y;; Drice::V5::TurboID/UASz-Fas3G::3xFLAG

Fig. 3C (Top to Bottom)

w/Y; Ubi-FRT-STOP-FRT-mCD8::DQVD::QF2/+; UAS-FLP, QUAS-tdTomato::3xHA/+

w/Y;   Ubi-FRT-STOP-FRT-mCD8::DQVA::QF2/ELAV-Gal4;   UAS-FLP,   QUAS-

tdTomato::3xHA/+

w/Y;   Ubi-FRT-STOP-FRT-mCD8::DQVD::QF2/ELAV-Gal4;   UAS-FLP,   QUAS-

tdTomato::3xHA/+

Fig. 4A, B (Top to Bottom, Left to Right)

w/Y; Or42b-Gal4/+; UAS-RedStinger/UASz-V5::TurboID

w/Y; Or42b-Gal4/+; UAS-RedStinger/UASz-Drice^{CG}::V5::TurboID

w/Y; Or42b-Gal4/+; UAS-RedStinger/UASz-Drice::V5::TurboID

w/Y; Or42b-Gal4/+; UAS-RedStinger/UASz-Dcp-1^{CG}::V5::TurboID

w/Y; Or42b-Gal4/+; UAS-RedStinger/UASz-Dcp-1::V5::TurboID

w/Y; Or42b-Gal4/+; UAS-RedStinger/UASz-Dronc^{CG}::V5::TurboID

w/Y; Or42b-Gal4/+; UAS-RedStinger/UASz-Dronc::V5::TurboID

Fig. 4C, D (Top to Bottom, Left to Right)

w/Y; Or42b-Gal4/UASz-3xFLAG::mNeonGreen; UAS-RedStinger/+

w/Y; Or42b-Gal4/+; UAS-RedStinger/UASz-Fas3G::3xFLAG

w/Y; Or42b-Gal4/UASz-Drice::myc::mNeonGreen; UAS-RedStinger/+

w/Y; Or42b-Gal4/UASz-Drice::myc::mNeonGreen; UAS-RedStinger/UASz-Fas3G::3xFLAG

Fig. 4E (Top to Bottom)

w/Y; Or42b-Gal4/UASz-3xFLAG::mNeonGreen

w/Y; Or42b-Gal4, Ubi-FRT-STOP-FRT-mCD8::DQVD::QF2/UASz-3xFLAG::mNeonGreen;

UAS-FLP, QUAS-tdTomato::3xHA/+

Fig. 4F, G (Top to Bottom, Left to Right)

w/w or Y; Or42b-Gal4, Ubi-FRT-STOP-FRT-mCD8::DQVD::QF2/UASz-

3xFLAG::mNeonGreen; UAS-FLP, QUAS-tdTomato::3xHA/+

w/w or Y; Or42b-Gal4, Ubi-FRT-STOP-FRT-mCD8::DQVD::QF2/+; UAS-FLP, QUAS-

tdTomato::3xHA/UASz-Fas3G::3xFLAG

w/w or Y; Or42b-Gal4, Ubi-FRT-STOP-FRT-mCD8::DQVD::QF2/UASz-

Drice::myc::mNeonGreen; UAS-FLP, QUAS-tdTomato::3xHA/+

w/w or Y; Or42b-Gal4, Ubi-FRT-STOP-FRT-mCD8::DQVD::QF2/UASz-

Drice::myc::mNeonGreen; UAS-FLP, QUAS-tdTomato::3xHA/UASz-Fas3G::3xFLAG

Fig. 4H, I (Top to Bottom, Left to Right)

w/w or Y; Or42b-Gal4, Ubi-FRT-STOP-FRT-mCD8::DQVD::QF2/UASz-

3xFLAG::mNeonGreen; UAS-FLP, QUAS-tdTomato::3xHA/+

w/w or Y; Or42b-Gal4, Ubi-FRT-STOP-FRT-mCD8::DQVD::QF2/+; UAS-FLP, QUAS-

tdTomato::3xHA/UASz-Fas3G::3xFLAG

w/w or Y; Or42b-Gal4, Ubi-FRT-STOP-FRT-mCD8::DQVD::QF2/UAS-Dronc-RNAi; UAS-

FLP, QUAS-tdTomato::3xHA/UASz-Fas3G::3xFLAG

w/w or Y; Or42b-Gal4, Ubi-FRT-STOP-FRT-mCD8::DQVD::QF2/UAS-Dronc^{DN}; UAS-FLP,

QUAS-tdTomato::3xHA/UASz-Fas3G::3xFLAG

Fig. 4K, L (Left to Right)

w/Y; Or42b-Gal4/UASz-3xFLAG::mNeonGreen

w/Y; Or42b-Gal4/+; +/UASz-Fas3G::3xFLAG

w/Y; Or42b-Gal4/UAS-Dronc-RNAi; +/UASz-Fas3G::3xFLAG

w/Y; Or42b-Gal4/UAS-Dronc^{DN}; +UASz-Fas3G::3xFLAG

Protein preparation from Drosophila tissues

Adult heads or brains of the given genotype were dissected and lysed with RIPA buffer (50 mM

Tris-HCl, pH 8.0, 150 mM Sodium Chloride, 0.5 w/v% Sodium Deoxycholate, 0.1% w/v

Sodium Dodecyl Sulfate, and 1.0% w/v NP-40) supplemented with cOmplete ULTRA EDTA-

free protease inhibitor cocktail (#05892953001, Roche). Samples were homogenized and

centrifuged at 20,000 ×g, 4°C for 10 min. The supernatants were collected and snap-frozen in

liquid nitrogen. Protein concentrations were determined using the BCA assay (#297-73101,

Wako) following the manufacturer's protocol. The samples were mixed with 6x Laemmli

buffer, boiled at 95°C for 5 min, and then subjected to purification of biotinylated proteins and

western blot analysis.

Western blotting

Each sample was separated using SDS-PAGE. The proteins were then transferred onto

Immobilon-P PVDF membranes (#IPVH00010; Millipore) for immunoblotting. Membranes

were blocked with 4% skim milk diluted in 1× TBST. Immunoblotting was performed using

the antibodies mentioned below diluted in 4% skim milk. Signals were visualized using

Immobilon Western Chemiluminescent HRP Substrate (#WBKLS0500; Millipore) and

FUSION SOLO. 7S. EDGE imaging station (Vilber-Lourmat). Contrast and brightness

adjustments were applied equally using Fiji (ImageJ) software (NIH Image).

The primary antibodies used included mouse anti-V5 monoclonal antibody (1:5,000,

#46-0705, Invitrogen), mouse anti-FLAG M2 monoclonal antibody (1:5,000, Sigma), mouse

anti-alpha tubulin (DM1A) monoclonal antibody (1:10,000, #T9026, Sigma), mouse anti-Actin

monoclonal antibody (1:5,000; #A4700, Sigma), mouse anti-Fasciclin 3 (7G10) antibody (1:50,

#AB_528238, DSHB), rabbit anti-Fasciclin3 isoform G antibody (1:50, this study). The

secondary antibodies used included goat anti-rabbit IgG HRP-conjugated antibody (1:5,000,

#7074S, CST) and goat/rabbit/donkey anti-mouse IgG HRP Conjugate (1:5,000, #W402B,

Promega). The membranes for streptavidin blotting were blocked with 3% BSA. Streptavidin

blotting was performed using streptavidin-horseradish peroxidase (1:10,000, #SA10001;

Invitrogen) diluted in 3% BSA.

Immunohistochemistry

Adult brains were dissected in phosphate-buffered saline (PBS) and fixed in 4%

paraformaldehyde (PFA) at room temperature for 30 min. Adult brains were washed three times

with 0.3% PBST (Triton-X100) for 10 min. After blocking with 5.0% normal donkey serum

(#S30, Millipore)/PBST (PBSTn) for 30 minutes at room temperature, tissues were incubated

with primary antibody/PBSTn overnight at 4°C. The samples were then washed with PBST

three times for 10 min each and incubated with secondary antibody/PBSTn for 2 h at room

temperature. After incubation, the tissues were washed with PBST three times for 10 min.

Streptavidin and Hoechst 33342 were added during secondary antibody incubation. The

samples were mounted on glass slides with SlowFade Gold antifade reagent (#S36939,

Invitrogen). Images were captured using a Leica TCS SP8 micrscope (Leica Microsystems).

Images were analyzed and edited using the Fiji (ImageJ) software. For MASCaT analysis,

newly eclosed adults were raised at 29°C for 1 week. Classification of the MASCaT signal was

performed by optically assessing the tdTomato intensity. For cell number analysis, newly

eclosed adults were raised at 29°C for 1 week. The antennae were dissected as previously

described [36], fixed with 4% PFA at room temperature for 30 min, and washed with 0.4%

PBST (Triton-X100) three times for 10 min. Quantification of cell numbers in antenna was

performed using Fiji (ImageJ) software using "3D object counter" for RedStinger positive cells.

The primary antibodies used were mouse anti-Fasciclin 3 (7G10) antibody (1:20,

#AB_528238, DSHB), rat anti-GFP (GF090R) monoclonal antibody (1:100, #04404-26,

nacalai Tesque), mouse anti-FLAG M2 monoclonal antibody (1:500, F1804, Sigma), and rat

anti-mCD8 monoclonal antibody (1:50, #MCD0800, Invitrogen). The secondary antibodies

used included Alexa Fluor 488-conjugated donkey anti-mouse IgG antibody (1:500, #A-21202,

ThermoFisher Scientific), Alexa Fluor 488-conjugated donkey anti-rat IgG antibody (1:500,

#A-21208, ThermoFisher Scientific), Alexa Fluor 647-conjugated donkey anti-mouse IgG

antibody (1:500, #A-31571, ThermoFisher Scientific), and Alexa Fluor 633-conjugated goat

anti-rat IgG antibody (1:500, #A-21094, ThermoFisher Scientific). The nuclei were stained

with Hoechst 33342 (8 μM, #H3570, Invitrogen). Streptavidin-Cy2 (1:500, #016-220-084,

Jackson ImmunoResearch) and Streptavidin-Cy5 (1:500, #016-170-084, Jackson

ImmunoResearch) were used to stain the biotinylated proteins.

Experimental design for LC-MS/MS analyses

Two biological replicates were analyzed for each experimental condition to determine the

relative abundance of *Drice::V5::TurboID/w^{1118}*.

Purification of biotinylated proteins

Samples were prepared as previously described [17]. Briefly, 100 μg of biotinylated protein-

containing lysate was subjected to FG-NeutrAvidin bead (#TAS8848 N1171, Tamagawa Seiki)

purification. The FG-NeutrAvidin beads (25 μL, approximately 500 μg) were washed three

times with RIPA buffer. Benzonase-treated biotinylated protein samples suspended in 1 mL

RIPA buffer were incubated overnight at 4°C. Conjugated beads were magnetically isolated

and washed with 500 μL of an ice-cold RIPA buffer solution, 1 M KCl solution, 0.1 M Na_2CO_3

solution and 4 M urea solution. For western blot analyses, the purified samples were mixed

with Laemmli buffer, boiled at 95°C for 5 min, and then subjected to western blot analysis. For

LC-MS/MS analysis, the purified samples were washed with 500 μL ultrapure water (#214-

01301, Wako) and 500 μL of 50 mM ammonium bicarbonate (AMBC). The samples were then

mixed with 50 µL of 0.1% Rapigest diluted in 50 mM AMBC, and 5 µL of 50 mM TCEP was

subsequently added. Samples were incubated at 60°C for 5 min, and then 2.5 µL 200 mM

MMTS was added. One microgram sequence-grade modified trypsin (#V5111, Promega) was

then added for on-bead trypsin digestion at 37°C for 16 h. The beads were magnetically isolated

and 60 µL of the supernatant was collected. Then, 3 µL 10% TFA was added to the

supernatants, and the samples were incubated at 37°C for 60 min with gentle agitation. The

samples were then centrifuged at 20,000 ×g, 4°C for 10 min. The peptides were desalinated and

purified using a GL-tip SDB (#7820-11200, GL Sciences) following the manufacturer's

protocol. The samples were speed-backed at 45°C for 30 min and dissolved in 25 µL of 0.1%

formic acid. The samples were then centrifuged at 20,000 ×g, 4°C for 10 min, and the

supernatants were collected. The peptide concentrations were determined using the BCA assay.

Finally, 500 ng of the purified protein was subjected to LC-MS/MS analysis.

LC-MS/MS analyses

LC-MS/MS analyses were performed as previously described [17]. Briefly, samples were

loaded onto Acclaim PepMap 100 C18 column (75 µm x 2 cm, 3 µm particle size and 100 Å

pore size; #164946, ThermoFisher Scientific) and separated on nano-capillary C18 column, (75

µm x 12.5 cm, 3 µm particle size, #NTCC-360/75-3-125, Nikkyo Technos) using EASY-nLC

1200 system (ThermoFisher Scientific). The elution conditions are listed in Table S3. The

separated peptides were analyzed using QExactive (ThermoFisher Scientific) in data-dependent

MS/MS mode. The parameters for the MS/MS analysis are listed in Table S4. The collected

data were analyzed using Proteome Discoverer (PD) 2.2 software with the Sequest HT search

engine. The parameters for the PD 2.2 analysis are described in Table S5. Peptides were filtered

at a false discovery rate of 0.01 using the Percolator node. Label-free quantification was

performed based on the intensities of the precursor ions using a precursor-ion quantifier node.

Normalization was performed using the total amount of peptides in all average scaling modes.

Proteins with 10 folds or higher abundance relative to the control (w^{1118}) were considered for

further analysis. MS proteomic data were deposited in the ProteomeXchange Consortium via

the jPOST partner repository with the dataset identifier PXD042922 [37].

Gene Ontology analysis

Gene Ontology (GO) analysis of cellular components was performed using DAVID [38].

Generation of rabbit polyclonal anti-Fasciclin 3 isoform G antibody

The peptide corresponding to the 506–524 AA (N-LKPANEATPATTPAPTTAA-C) of Fas3G

was used by Eurofins Inc. to immunize rabbits to raise a polyclonal antibody. The antibody was

affinity purified using the same peptide provided by Eurofins, Inc.

IDR prediction

Amino acid sequences were obtained from the FlyBase database [https://flybase.org/].

Disordered intracellular regions of Fas3 isoforms were predicted using the PSIPRED protein

sequence analysis workbench in DISOPRED3 [http://bioinf.cs.ucl.ac.uk/psipred/].

Molecular cloning

For all isoforms (A, B/F, C, D/E, G) of *pUASz-Fasciclin 3::3xFLAG*, the coding sequences of

Fasciclin 3s were PCR-amplified using 3xFLAG-tag harboring primers from *Drosophila* cDNA

and were ligated into the *BamH*I/*Kpn*I-digested *pUASz1.0* vector [39] using In-Fusion

(#Z9648N TaKaRa). For *pUASz-3xFLAG::mNeonGreen*, the coding sequence of mNeonGreen

was PCR-amplified using 3xFLAG-tag harboring primers from *pAc5-V5::mNeonGreen* [17]

and ligated into the *Xho*I-digested *pUASz1.0* vector [39] using In-Fusion. For *pUASz-*

Drice::myc::mNeonGreen, the coding sequence of mNeonGreen was PCR-amplified from

pAc5-V5::mNeonGreen [17] using myc-tag-harboring primers. The coding sequence of Drice

was PCR-amplified from *Drosophila* cDNA. The two fragments were then ligated into the

*Xho*I-digested *pUASz1.0* vector using In-Fusion. For *pUASz-V5::TurboID* and *pUASz-*

Caspase::V5::TurboID, the coding sequences of *Dcp-1*, *Drice* and *Dronc* were PCR amplified

from *Drosophila* cDNA. The coding sequence of V5::TurboID was PCR-amplified from *V5-*

TurboID-NES_pcDNA3 (#107169; Addgene). The two fragments were then ligated into the

*Xho*I-digested *pUASz1.0* vector using In-Fusion. Catalytic dead mutations (Dcp-1: C196G

(TGC to GGC), Drice: C211G (TGC to GGC), and Dronc: C318G (TGC to GGC)) were

introduced by PCR-based site-directed mutagenesis, and the corresponding fragments were

ligated into the *Xho*I-digested *pUASz1.0* vector using In-Fusion. *For pUbi-FRT-STOP-FRT-*

mCD8::DQVD/A::QF2, the FRT-STOP-FRT cassette sequence was PCR amplified from

pJFRC201-10xUAS-FRT-STOP-FRT-myr::smGFP-HA (#63166; Addgene). The coding

sequence of mCD8::DQVD/A was PCR-amplified from the genomic DNA of *Caspase-*

Sensitive/Insensitive-Gal4 [20]. The coding sequence of QF2 was PCR amplified from the

pBPQUw plasmid [40]. The three fragments were ligated into *EcoR*I-digested *pUbi83* vector (a

gift from D. Umetsu) using In-Fusion. For *pQUAST-mNeonGreen*, the coding sequence of

mNeonGreen was PCR-amplified from *pAc5-V5::mNeonGreen* [17] and ligated into the *EcoR*I-

digested *pQUAST* (#24349, Addgene) vector using In-Fusion. For *pUASz-FLPo*, the coding

sequences of FLPo were PCR-amplified from *pQUAST-FLPo* (#24357, Addgene) and ligated

into the *Xho*I-digested *pUASz1.0* vector [39] using In-Fusion.

All established plasmids were sequenced by Eurofins, Inc. The detailed plasmid DNA

sequences are available upon request. The established plasmids were injected into y^1, w^{67c23};

P[CaryP]attP40 (*pUASz-3xFLAG::mNeonGreen* and *pUASz-Drice::myc::mNeonGreen*) or y^1,

w^{67c23}; *P[CaryP]attP2* (*pUASz-Fasciclin 3::3xFLAGs* and *pUbi-FRT-STOP-FRT-*

mCD8::DQVD/A::QF2) for PhiC31-mediated transgenesis (BestGene Inc.). Each red-eye-

positive transformant was isogenized. Integration of the attP landing site was confirmed using

genomic PCR.

Cell culture

Drosophila S2 cells were grown at 25°C in Schneider's *Drosophila* medium (#21720001,

GIBCO) supplemented with 10% (v/v) heat-inactivated fetal bovine serum, 100 U/ml penicillin,

and 100 µg/ml streptomycin (#168-23191, WAKO). Cells were seeded in 24-well plates and

transfected with the desired plasmids using Effectene Transfection Reagent (#301427,

QIAGEN) following the manufacturer's protocol. Sixteen hours after transfection, the cells

were reseeded into CELL CULTURE DISH, PS, 35/10 MM (#627870, Grainer BIO-ONE) and

incubated for 24 h. Images were obtained using a Leica AF6000 DMI6000 B microscope (Leica

Microsystems). Images were analyzed and edited using Fiji software (NIH Image).

Aging experiment

Adult male flies were raised in vials containing 15 fly/vial, and incubated at 29°C, 65%

humidity with 12h/12h light/dark cycle. The flies were transferred every 2–4 days to vials

containing fresh food.

Two choice preference assay

Newly eclosed adult male flies were raised in vials containing 25 fly/vial, and incubated at 29°C,

65% humidity with 12h/12h light/dark cycle. The flies were transferred every 2–4 days to vials

containing fresh food for 1 week. The flies were starved in MilliQ (MQ) for 4 h prior to the

assay. About 75 flies were put in 1000-mL glass beaker, which contained two standards bottles

filled with 500 µl 1% ACV and 500 µl MQ. The bottles were then covered with sponge plugs

containing pipette tips. The pipette tip was cut to increase the size of the opening such that only

one fly could climb into the bottle at a time. The glass beaker was covered with a mesh fabric.

The beakers were set in an incubator at 29°C for 2 hr for the assay. The experiments were

evaluated only if at least one of the tested flies entered either of 2 bottles. The preference index

equaled the number of flies in the ACV minus the number of flies in the MQ, divided by the

number of flies in both bottles and fields, and multiplied by 100. Values greater than zero

indicate a preference for 1% ACV and values less than zero indicate aversion. We repeated the

same experiment with different sets of flies and the number of times is indicated by N.

Statistical analysis

Statistical analyses were performed using the GraphPad Prism 8 software (MDF). Data are

shown as the means ± standard error of the mean. *P*-values were calculated using a one-way

analysis of variance (ANOVA) with Bonferroni correction (Fig. 4B for selected pairs, Fig. 4D

for every pair). *P*-values were calculated using the chi-square test with Bonferroni's correction,

with mNeonGreen as a control (Fig. 4G), mNeonGreen and Fas3G as controls (Fig. 4I), and

mNeonGreen as a control (Fig. 4L). *P*-values were calculated using one-way ANOVA with

Dunnett's multiple comparison test (Fig. 4K, mNeonGreen as a control). NS: $P > 0.05$, †: <

0.05.

Acknowledgments

We would like to thank Dr. D. Umetsu for providing plasmids, Dr. Y. Nitta, Dr. A. Sugie, and Dr. T. Ichinose for helpful discussions, and the Bloomington *Drosophila* Stock Center and Vienna Drosophila Resource Center for providing the fly strains. We thank Dr. S. Hirayama, Dr. S. Murata, and the one-stop sharing faculty center for future drug discovery at the Graduate School of Pharmaceutical Sciences, University of Tokyo, for the LC-MS/MS analysis. We thank Miura's lab members for their technical assistance and discussions; in particular, K. Takenaga for preparing the fly food, and R. Takamoto for experimental support. The MASCaT was established by MASaya Muramoto with critical advice from members of Shu MASuda at MASayuki Miura Laboratories. We would like to thank *Editage* (www.editage.com) for English language editing of this manuscript.

References

634   1.   Green DR. The Coming Decade of Cell Death Research: Five Riddles. Cell. 2019;177:

635   1094–1107. doi:10.1016/j.cell.2019.04.024

636   2.   Nakajima YI, Kuranaga E. Caspase-dependent non-apoptotic processes in development.

637   Cell Death Differ. 2017;24: 1422–1430. doi:10.1038/cdd.2017.36

638   3.   Ohsawa S, Hamada S, Kuida K, Yoshida H, Igaki T, Miura M. Maturation of the

639   olfactory sensory neurons by Apaf-1/caspase-9–mediated caspase activity. Proc Natl

640   Acad Sci U S A. 2010;107: 13366–13371. doi:10.1073/pnas.0910488107

641   4.   Li Z, Jo J, Jia JM, Lo SC, Whitcomb DJ, Jiao S, et al. Caspase-3 activation via

642   mitochondria is required for long-term depression and AMPA receptor internalization.

643   Cell. 2010;141: 859–871. doi:10.1016/j.cell.2010.03.053

644   5.   Aram L, Yacobi-Sharon K, Arama E. CDPs: Caspase-dependent non-lethal cellular

645   processes. Cell Death Differ. 2017;24: 1307–1310. doi:10.1038/cdd.2017.111

646   6.   D'Brot A, Chen P, Vaishnav M, Yuan S, Akey CW, Abrams JM. Tango7 directs cellular

647   remodeling by the Drosophila apoptosome. Genes Dev. 2013;27: 1650–1655.

648   doi:10.1101/gad.219287.113

649   7.   Kang Y, Neuman SD, Bashirullah A. Tango7 regulates cortical activity of caspases

650   during reaper-triggered changes in tissue elasticity. Nat Commun. 2017;8: 603.

651   doi:10.1038/s41467-017-00693-3

652   8.   Orme MH, Liccardi G, Moderau N, Feltham R, Wicky-John S, Tenev T, et al. The

653   unconventional myosin CRINKLED and its mammalian orthologue MYO7A regulate

654   caspases in their signalling roles. Nat Commun. 2016;7: 10972.

655   doi:10.1038/ncomms10972

656   9.   Kanuka H, Kuranaga E, Takemoto K, Hiratou T, Okano H, Miura M. Drosophila caspase

657   transduces Shaggy/GSK-3β kinase activity in neural precursor development. EMBO J.

658   2005;24: 3793–3806. doi:10.1038/sj.emboj.7600822

659   10.   Amcheslavsky A, Wang S, Fogarty CE, Lindblad JL, Fan Y, Bergmann A. Plasma

660   Membrane Localization of Apoptotic Caspases for Non-apoptotic Functions. Dev Cell.

661   2018;45: 450–464.e3. doi:10.1016/j.devcel.2018.04.020

662   11.   Julien O, Wells JA. Caspases and their substrates. Cell Death Differ. 2017;24: 1380–

663   1389. doi:10.1038/cdd.2017.44

664   12.   Kumar S. Caspase function in programmed cell death. Cell Death Differ. 2007;14: 32–

665   43. doi:10.1038/sj.cdd.4402060

666   13.   Kumar S, Dorstyn L, Lim Y. The role of caspases as executioners of apoptosis. Biochem

667   Soc Trans. 2022;50: 33–45. doi: 10.1042/BST20210751

668   14.   Branon TC, Bosch JA, Sanchez AD, Udeshi ND, Svinkina T, Carr SA, et al. Efficient

669   proximity labeling in living cells and organisms with TurboID. Nat Biotechnol. 2018;36:

670   880–887. doi:10.1038/nbt.4201

671   15.   Qin W, Cho KF, Cavanagh PE, Ting AY. Deciphering molecular interactions by

672   proximity labeling. Nat Methods. 2021;18: 133–143. doi:10.1038/s41592-020-01010-5

673   16.   Shinoda N, Hanawa N, Chihara T, Koto A, Miura M. Dronc-independent basal

674   executioner caspase activity sustains Drosophila imaginal tissue growth. Proc Natl Acad

675   Sci U S A. 2019;116: 20539–20544. doi:10.1073/pnas.1904647116

676   17.   Shinoda N, Horikoshi M, Taira Y, Muramoto M, Hirayama S, Murata S, et al. Caspase

677   cleaves Drosophila BubR1 to modulate spindle assembly checkpoint function and

678   lifespan of the organism. FEBS J. 2023; Forthcoming. doi:10.1111/febs.16811

679   18.   Chiba A, Snow P, Keshishian H, Hotta Y. Fasciclin III as a synaptic target recognition

680   molecule in Drosophila. Nature. 1995;374: 166–168. doi:10.1038/374166a0

681   19.   Kose H, Rose D, Zhu X, Chiba A. Homophilic synaptic target recognition mediated by

682   immunoglobulin-like cell adhesion molecule Fasciclin III. Development. 1997;124:

683   4143–4152. doi: 10.1242/dev.124.20.4143.

684   20.   Tang HL, Tang HM, Fung MC, Hardwick JM. In vivo Caspase tracker biosensor system

685   for detecting anastasis and non-apoptotic caspase activity. Sci Rep. 2015;5: 9015.

686   doi:10.1038/srep09015

687   21.   Ding AX, Sun G, Argaw YG, Wong JO, Easwaran S, Montell DJ. CasExpress reveals

688   widespread and diverse patterns of cell survival of caspase-3 activation during

689   development in vivo. Elife. 2016;5: e10936. doi:10.7554/eLife.10936

690   22.   Chihara T, Kitabayashi A, Morimoto M, Takeuchi K, Masuyama K, Tonoki A, et al.

691   Caspase Inhibition in Select Olfactory Neurons Restores Innate Attraction Behavior in

692   Aged Drosophila. PLoS Genet. 2014;10: e1004437. doi:10.1371/journal.pgen.1004437

693   23.   Semmelhack JL, Wang JW. Select Drosophila glomeruli mediate innate olfactory

694   attraction and aversion. Nature. 2009;459: 218–223. doi:10.1038/nature07983

695   24.   Walsh JG, Cullen SP, Sheridan C, Lüthi AU, Gerner C, Martin SJ. Executioner caspase-

696   3 and caspase-7 are functionally distinct proteases. Proc Natl Acad Sci U S A. 2008;105:

697   12815–12819. doi:10.1073/pnas.0707715105

698   25.   Boucher D, Blais V, Denault JB. Caspase-7 uses an exosite to promote poly(ADP ribose)

699   polymerase 1 proteolysis. Proc Natl Acad Sci U S A. 2012;109: 5669–5674.

700   doi:10.1073/pnas.1200934109

701   26.   Desroches A, Denault JB. Caspase-7 uses RNA to enhance proteolysis of poly(ADP-

702   ribose) polymerase 1 and other RNA-binding proteins. Proc Natl Acad Sci U S A.

703   2019;116: 21521–21528. doi:10.1073/pnas.1909283116

704   27.   Takemoto K, Nagai T, Miyawaki A, Miura M. Spatio-temporal activation of caspase

705   revealed by indicator that is insensitive to environmental effects. J Cell Biol. 2003;160:

706   235–243. doi:10.1083/jcb.200207111

707   28.   Valente P, Lignani G, Medrihan L, Bosco F, Contestabile A, Lippiello P, et al. Cell

708   adhesion molecule L1 contributes to neuronal excitability regulating the function of

709   voltage-gated Na+ channels. J Cell Sci. 2016;129: 1878–1891. doi:10.1242/jcs.182089

710   29.   Mukherjee A, Williams DW. More alive than dead: Non-apoptotic roles for caspases in

711   neuronal development, plasticity and disease. Cell Death Differ. 2017;24: 1411–1421.

712   doi:10.1038/cdd.2017.64

713   30.   Dehkordi MH, Munn RGK, Fearnhead HO. Non-Canonical Roles of Apoptotic Caspases

714   in the Nervous System. Front Cell Dev Biol. 2022;10: 840023.

715   doi:10.3389/fcell.2022.840023

716   31.   Gu Q, Jiao S, Duan K, Wang YX, Petralia RS, Li Z. The BAD-BAX-Caspase-3 Cascade

717   Modulates Synaptic Vesicle Pools via Autophagy. J Neurosci. 2021;41: 1174–1190.

718   doi:10.1523/JNEUROSCI.0969-20.2020

719   32.   Cheriyamkunnel SJ, Rose S, Jacob PF, Blackburn LA, Glasgow S, Moorse J, et al. A

720   neuronal mechanism controlling the choice between feeding and sexual behaviors in

721   Drosophila. Curr Biol. 2021;31: 4231-4245.e4. doi:10.1016/j.cub.2021.07.029

722   33.   Westphal D, Sytnyk V, Schachner M, Leshchyns'ka I. Clustering of the neural cell

723   adhesion molecule (NCAM) at the neuronal cell surface induces caspase-8- and -3-

724   dependent changes of the spectrin meshwork required for NCAM-mediated neurite

725   outgrowth. J Biol Chem. 2010;285: 42046–42057. doi:10.1074/jbc.M110.177147

726   34.   Luck K, Kim DK, Lambourne L, Spirohn K, Begg BE, Bian W, et al. A reference map

727   of the human binary protein interactome. Nature. 2020;580: 402–408.

728   doi:10.1038/s41586-020-2188-x

35.   Li L, Ding Z, Pang TL, Zhang B, Li CH, Liang AM, et al. Defective minor spliceosomes

induce SMA-associated phenotypes through sensitive intron-containing neural genes in

Drosophila. Nat Commun. 2020;11: 5608. doi:10.1038/s41467-020-19451-z

36.   Karim MR, Endo K, Moore AW, Taniguchi H. Whole mount immunolabeling of

olfactory receptor neurons in the Drosophila antenna. J Vis Exp. 2014; 51245.

doi:10.3791/51245

37.   Okuda S, Watanabe Y, Moriya Y, Kawano S, Yamamoto T, Matsumoto M, et al.

JPOSTrepo: An international standard data repository for proteomes. Nucleic Acids Res.

2017;45: D1107–D1111. doi:10.1093/nar/gkw1080

38.   Huang DW, Sherman BT, Lempicki RA. Bioinformatics enrichment tools: Paths toward

the comprehensive functional analysis of large gene lists. Nucleic Acids Res. 2009;37:

1–13. doi:10.1093/nar/gkn923

39.   Deluca SZ, Spradling AC. Efficient expression of genes in the drosophila germline using

a uas promoter free of interference by hsp70 pirnas. Genetics. 2018;209: 381–387.

doi:10.1534/genetics.118.300874

40.   Kashio S, Obata F, Zhang L, Katsuyama T, Chihara T, Miura M. Tissue nonautonomous

effects of fat body methionine metabolism on imaginal disc repair in Drosophila. Proc

Natl Acad Sci U S A. 2016;113: 1835–1840. doi:10.1073/pnas.1523681113

Fig 1. Expression patterns and proximal proteins of Drice in the adult brain.

(A) Schematic diagram of *Drosophila* apoptosis signaling. (B) Western blot of expression of

caspases in adult male heads. (C) Representative images of adult male brains. Expression

patterns of each caspase are visualized using Streptavidin (SA, magenta). Scale bar: 50 μm. (D)

Representative images of antennal lobes (ALs) in the adult female brains. Expression patterns

of Drice are visualized using SA (magenta). Scale bar: 50 μm. (E) Western blot of proteins

extracted from adult male brains. (F) Schematic diagram of the TurboID-mediated

identification of proximal proteins. (G) A summary of mass spectrometry analysis of proteins

of the adult male brains. Partial list of Drice proximal proteins and their Fold Change (FC) are

shown. (H) Gene Ontology analysis (Cellular Compartment) of 158 Drice proximal proteins

(*Drice::V5::TurboID*/*w^{1118}* > 10). Cyan: neural-related fraction, Red: membrane fraction.

Fig 2. A specific isoform of Fasciclin 3 is in proximity to Drice.

(A) Representative expression patterns of Drice (Streptavidin (SA); magenta) and Fasciclin 3 (Fas3, anti-Fasciclin 3 antibody staining; green) in the adult male brains. Scale bar: 50 μm. (B) Representative expression patterns of Drice (SA; magenta) and Fas3 (anti-GFP antibody staining; green) in the antennal lobes (ALs) of the adult male brains. Scale bar: 50 μm. (C) Gene structure of Fas3. CDS (orange) and UTR (grey) are shown in boxes. (D) Schematic structures

of Fas3 protein isoforms. Intracellular regions differ from each other. A peptide region used to

raise anti-Fas3G antibody is shown. (E) IDR predictions of intracellular regions of each protein

isoform. (F) Western blot of proteins extracted from adult heads. Anti-Fas3G antibody

specifically detects Fas3 isoform G. (G) Western blot of adult male heads. Drice proximal

proteins were purified using NeturAvidin. (H) Western blot of adult head proteins. Drice

proximal proteins were purified using NeturAvidin.

Fig 3. A Gal4-Manipulated Area Specific CaspaseTracker/CasExpress (MASCaT).

(A) A schematic diagram of MASCaT expressed using the Gal4/UAS system with FLP-mediated recombination. Caspase sensitive-QF2 is activated by caspase-mediated cleavage at the plasma membrane. (B) Representative images of S2 cells expressing MASCaT. Scale bar: 20 µm. (C) Representative images of adult male brains of young (1 week old) and old (3 weeks old) flies raised at 29°C. Arrowheads indicate optic lobes. Magnified images (white rectangle) show antennal lobes (ALs; white circles). Arrows indicate age-dependent caspase activation at the ALs. Scale bar: 100 µm.

Fig 4. Fasciclin 3 isoform G regulates non-lethal caspase activation and functions.

(A) Representative images of the adult male antennas of 1 weeks old flies raised at 29°C. Cell bodies are visualized using RedStinger. Scale bar: 20 μm. (B) Quantification of cell number of (A). Data are presented as mean ± SEM. P-values were calculated using one-way analysis of variance (ANOVA) with Bonferroni's correction for selected pairs. NS: $P > 0.05$, †: $P < 0.05$. (C) Representative images of the adult male antennas of 1 week old flies raised at 29°C. Cell bodies are visualized using RedStinger. Scale bar: 20 μm. (D) Quantification of cell number of (C). Data are presented as mean ± SEM. P-values were calculated using one-way ANOVA with Bonferroni's correction for every pair. NS: $P > 0.05$, †: $P < 0.05$. (E) Representative images of the antennal lobes (ALs) of male brains of 1 week old flies raised at 29°C expressing MASCaT probe (mCD8). Scale bar: 30 μm. (F) Representative images of the ALs of male brains of 1 week old flies raised at 29°C with caspase activation visualized by MASCaT. Scale bar: 30 μm. (G) Quantifications of caspase activity detected by MASCaT of (F). P-values were calculated using chi-square test with Bonferroni's correction using mNeonGreen as control. NS: $P > 0.05$, †: $P < 0.05$. Sample sizes are shown in the graph. (H) Representative images of the AL of male brains of 1 week old flies raised at 29°C with caspase activation visualized using MASCaT. Scale bar: 30 μm. (I) Quantifications of caspase activity detected using MASCaT of (H). P-values were calculated using chi-square test with Bonferroni's correction using mNeonGreen and Fas3G as control, respectively. NS: $P > 0.05$, †: $P < 0.05$. Sample sizes are shown in the graph. (J) A schematic model of compartmentalized non-lethal caspase activation mediated by Fas3G and Dronc. (K) Preference index of 1 week old male flies raised at 29°C in response to 1% apple cider vinegar in two choice preference assay. Data are presented as mean ± SEM. P-values were calculated using one-way ANOVA followed by Dunnets' multiple comparison test using mNeonGreen as a control. NS: $P > 0.05$, †: $P < 0.05$. Sample sizes: mNG (N = 19 (1327 flies)), Fas3G (N = 17 (1191 flies)), Fas3G + Dronc-RNAi (N = 19 (1295 flies)), Fas3G +

Dronc^{DN} (N = 20 (1320 flies)). (L) The ratio of flies in the given conditions in the two choice

preference assay of (K). *P*-values were calculated using chi-square test with Bonferroni's

correction using mNeonGreen as control. †: *P* < 0.05. Sample sizes: mNG (N = 1327 flies),

Fas3G (N = 1191 flies), Fas3G + Dronc-RNAi (N = 1295 flies), Fas3G + Dronc^{DN} (N = 1320

flies).

Legends for Table S1–5

Table S1. Proximal protein list. A list of the identified proximal proteins (Abundance ratio

(Drice-V5-TurboID/Control) > 10). The UniProt Accessions, Gene Description, and

Abundance Ratio (Drice-V5-TurboID/Control) are listed.

Table S2. Gene Ontology analysis. The result of Gene Ontology (GO) analysis of Cellular

Compartment for Drice proximal proteins. GO term, P values, Fold Enrichment, Count and

Genes are listed.

Table S3. LC settings. Gradient settings for LC analysis are presented.

Table S4. MS/MS settings. Settings used for the MS/MS analysis are presented.

Table S5. Proteome Discoverer 2.2 settings. Proteome Discoverer 2.2 settings for proteomics

analysis are presented.