Investigating the effects of age and hearing loss on speech intelligibility and amplitude modulation frequency selectivity

Jonathan Regev,[1,a] Johannes Zaar,[1,2] Helia Relaño-Iborra,[1] and Torsten Dau[1]

[1] *Hearing Systems Section, Department of Health Technology, Technical University of Denmark, Kongens Lyngby,*

2800, Denmark

[2] *Eriksholm Research Centre, Snekkersten, 3070, Denmark*

Abstract

The perception of amplitude modulation (AM), characterized by a frequency-selective process in the modulation domain, is considered critical for speech intelligibility. Previous literature has provided evidence of an age-related decline in AM frequency selectivity. Additionally, a notable sharpening of AM tuning with hearing loss has been observed, which was proposed to be related to a perceptual advantage resulting from peripheral compression loss. This study explores whether such changes in AM tuning with age and hearing loss contribute to the speech intelligibility challenges that older listeners often face in noisy environments. Young (n=10, 22-28 years) and older (n=9, 57–77 years) listeners with normal hearing as well as older listeners with hearing impairment (n=9, 64-77 years) participated in the study. All had previously taken part in studies on AM tuning. Speech-reception thresholds (SRTs) were collected in conditions including stationary, fluctuating, and competing-speech maskers. The results revealed an age-related increase in SRTs, with an additional negative impact of

---

[a] Corresponding author: joreg@dtu.dk

hearing loss. Beyond age and audibility, a measure of AM tuning significantly contributed to explaining the variance in SRTs across listeners for specific maskers. These findings motivate further exploration of the relationship between AM frequency selectivity and speech intelligibility in noise.

I. INTRODUCTION

Temporal envelope fluctuations (also known as "amplitude modulation" or AM) refer to the slow variations in overall amplitude that characterize most sounds in daily life (e.g., Singh and Theunissen, 2003; Elliott and Theunissen, 2009). The auditory system exhibits remarkable sensitivity to AM, as demonstrated by physiological studies (see Joris *et al.*, 2004 for a review) and behavioral investigations of AM detection (e.g., Viemeister, 1979; Kohlrausch *et al.*, 2000). Similar to masking effects in the audio-frequency domain, the detectability of a target AM is disrupted by the presence of distinct masking AM components. AM masking has been extensively studied in behavioral investigations (e.g., Bacon and Grantham, 1989; Houtgast, 1989; Dau *et al.*, 1997; Verhey *et al.*, 2003) as well as through recent neurophysiological research (Viswanathan *et al.*, 2021). Behavioral studies of AM masking have provided evidence of a frequency-selective process in the modulation domain, often conceptualized as a series of bandpass modulation filters known as the modulation filterbank (e.g., Dau *et al.*, 1997, 1999; Ewert and Dau, 2000; Ewert *et al.*, 2002). Computational modelling studies have shown that this concept significantly contributes to quantitatively predicting experimental data from various psychoacoustic tasks (e.g., Dau *et al.*, 1997, 1999; Jepsen *et al.*, 2008; Jepsen and Dau, 2011; McDermott and Simoncelli, 2011; McWalter and Dau, 2017), underscoring the importance of AM in auditory processing and perception.

Beyond its role in 'simple' listening tasks, AM has been argued to be crucial for accurate speech intelligibility (Houtgast and Steeneken, 1985; Drullman *et al.*, 1994; Smith *et al.*, 2002; Elliott and Theunissen, 2009). The significance of AM in speech was highlighted by vocoding studies, which demonstrated that intact envelope information conveyed through a few broad spectral bands was sufficient for precise speech recognition in quiet, even when spectral cues were greatly reduced (e.g., Shannon *et al.*, 1995; Dorman *et al.*, 1997; Xu *et al.*, 2005). Based on this, AM masking has been proposed as the primary mechanism underlying speech intelligibility in noise, whereby speech-

reception thresholds (SRTs) - reflecting the signal-to-noise ratio required for 50% speech recognition

- are related to the modulation-domain signal-to-noise ratio (SNR_{env}) between the speech and masker

(e.g., Dubbelboer and Houtgast, 2008; Jørgensen and Dau, 2011).

The crucial role of AM masking in speech intelligibility in noise was also emphasized by studies

using maskers designed to contain AM components or to have maximally flat envelopes, which

suggested that so-called stationary maskers, usually associated with energetic masking (EM) in the

audio-frequency domain, effectively function as AM maskers due to their inherent envelope

fluctuations (Stone *et al.*, 2012; Stone and Moore, 2014). Computational auditory modeling studies

have successfully incorporated these concepts by employing the SNR_{env} (Jørgensen and Dau, 2011;

Jørgensen *et al.*, 2013; Chabot-Leclerc *et al.*, 2014) or correlation-based measures (Relaño-Iborra *et al.*,

2016, 2019; Steinmetzger *et al.*, 2019) at the output of the modulation filterbank as decision metrics.

Based on these modeling studies, the spectral decomposition of the stimulus envelope carried out by

the modulation filterbank, i.e., AM frequency selectivity, has been suggested as an important factor

for accurately predicting SRTs (see Relaño-Iborra and Dau, 2022 for a review of different speech

intelligibility models using modulation filtering).

Regev *et al.* (2023, 2024a) investigated the effects of age and hearing loss on the selectivity of

hypothetical modulation filters using a perceptual AM masking task. Regev *et al.* (2023) found a general

reduction in AM frequency selectivity associated with age, resulting in broader masking patterns

among older compared to young listeners with normal hearing (referred to as hereafter as NH

listeners). Conversely, Regev *et al.* (2024a) demonstrated a general sharpening of masking patterns

among young and older listeners with hearing impairment (referred to as hereafter as HI listeners)

compared to NH listeners of similar age. This finding suggested that sensorineural hearing loss was

accompanied by a perceptual benefit that partially offset the detrimental effects of age, possibly due

to the HI listeners' ability to exploit an increased internal representation of AM depth resulting from

reduced peripheral compression when "listening in the dips" of the masker. Hence, Regev *et al.* (2024a)

proposed that the sharpening of the masking patterns found in HI listeners was an "apparent" effect,

reflecting the loss of peripheral compression rather than an actual greater selectivity of the hypothetical

modulation filters. Such changes in the sharpness of the hypothetical modulation filters due to age

and hearing loss may affect speech intelligibility in noise by altering the amount of AM masking at the

output of the modulation filters.

The effects of age and hearing loss on speech intelligibility have received significant attention in

the literature. Hearing loss is known to cause a decline in speech intelligibility in both quiet and noisy

environments (e.g., Glasberg and Moore, 1989; Humes and Christopherson, 1991; Frisina and Frisina,

1997; see Humes and Dubno, 2010 for a review). In addition to audibility loss, speech intelligibility

challenges associated with hearing loss have been linked to various supra-threshold deficits, including

reduced frequency selectivity, poorer temporal fine structure processing, reduced recovery from

forward masking, poorer AM depth discrimination, and loudness recruitment (Glasberg and Moore,

1986, 1989; Oxenham and Moore, 1997; Moore and Oxenham, 1998; Moore, 2003; Lorenzi *et al.*,

2006; Strelcyk and Dau, 2009; Schlittenlacher and Moore, 2016). Aided speech intelligibility

performance has also been shown to be strongly correlated with deficits in spectro-temporal

modulation detection in HI listeners (e.g., Bernstein *et al.*, 2016; Zaar *et al.*, 2023, 2024). Recent

neurophysiological studies suggested that poorer speech intelligibility in HI listeners may be related to

increased neural coding of the envelope (Millman *et al.*, 2017; Goossens *et al.*, 2018; Decruy *et al.*, 2020).

This effect may be partly attributed to an increased internal representation of the AM depth resulting

from the loss of cochlear compression commonly associated with sensorineural hearing loss (Moore

et al., 1996; Jennings *et al.*, 2018). In addition to the effects of hearing loss itself, age and reduced

cognitive function have been found to contribute to the decline in speech intelligibility observed in

older HI listeners (e.g., Dubno *et al.*, 1984; Humes and Christopherson, 1991; Frisina and Frisina,

1997; Humes, 2002, 2007; Humes *et al.*, 2006). This suggests that cognitive and age-related perception

deficits may also play a role in speech intelligibility challenges.

To investigate these potential mechanisms, cross-sectional studies focusing on age-related effects

have been conducted, often involving listeners with no or mild hearing loss. These studies have

consistently shown a detrimental effect of age on speech intelligibility in the presence of competing

speech maskers (Rajan and Cainer, 2008; Helfer and Vargo, 2009; Schoof and Rosen, 2014; Füllgrabe

et al., 2015; Gordon-Salant and Cole, 2016; Goossens *et al.*, 2017; Decruy *et al.*, 2019; Sobon *et al.*,

2019). However, the effects of age in stationary and fluctuating noise have been less consistent. Some

studies found differences between young and older listeners (e.g., Goossens *et al.*, 2017; Decruy *et al.*,

2019; Sobon *et al.*, 2019), while others did not (Rajan and Cainer, 2008; Helfer and Vargo, 2009; Schoof

and Rosen, 2014). Overall, the age-related increase in SRTs has been more pronounced for speech

maskers than for stationary or fluctuating noises (e.g., Goossens *et al.*, 2017; Decruy *et al.*, 2019; Sobon

et al., 2019). The observed age-related decline in speech intelligibility has been suggested to be mediated

by deficits in temporal processing (e.g., Gordon-Salant *et al.*, 2006; Anderson *et al.*, 2012; Füllgrabe *et

al.*, 2015; Goossens *et al.*, 2016; Decruy *et al.*, 2019) and by reduced cognitive abilities (Humes *et al.*,

2012; Füllgrabe *et al.*, 2015; Gordon-Salant and Cole, 2016).

Since AM frequency selectivity has been identified as a significant factor contributing to accurate

speech intelligibility, it is reasonable to suggest that the age-related decline in AM selectivity reported

by Regev *et al.* (2023) may contribute to the aforementioned challenges in speech intelligibility. A

reduction in AM tuning would likely result in increased AM masking, thus disrupting the intelligibility

of the target speech. Conversely, while the increased AM frequency selectivity found in older HI

listeners compared to older NH listeners (Regev *et al.*, 2024a) may suggest a decrease in AM masking

with hearing loss, it is hypothesized that this would not directly translate into improved speech

intelligibility. Regev *et al.* (2024a) proposed that the greater AM frequency selectivity in HI listeners,

rather than reflecting a sharpening of the hypothetical modulation filter, is linked to reduced cochlear

compression. Reduced cochlear compression is associated with various supra-threshold deficits that

impair speech intelligibility (e.g., Glasberg and Moore, 1989; Oxenham and Moore, 1997; Moore and

Oxenham, 1998; Moore, 2003). This assumption is consistent with previous studies showing that HI

listeners exhibit higher SRTs than age-matched NH listeners (Goossens *et al.*, 2017).

The present study explored whether the results of speech intelligibility tests align with the

assumption that an age-related decline in AM frequency selectivity would negatively impact speech

intelligibility, while an apparent sharpening of AM tuning with hearing loss would not be beneficial

for speech intelligibility. SRTs were collected for speech in noise, building upon the previous studies

on AM frequency selectivity (Regev *et al.*, 2023, 2024a), using a subset of the participants. Young NH,

older NH, and older HI listeners were tested on sentence recognition in various maskers, including

speech-shaped noise, fluctuating noise, and competing talkers. It was hypothesized that changes in

AM frequency selectivity with age and hearing loss would be linked to increases in SRTs for the older

NH and HI listeners.

II.   METHODS

A. Listeners and previously published data

Ten young NH listeners, aged 22-28 years (mean age = 24.7 years, eight female), nine older NH

listeners, aged 57–77 years (mean age = 66.9 years, six female), and nine older HI listeners, aged 64-

77 years (mean age = 71.2 years, 2 female), participated in the study. All listeners were native Danish

speakers. The experiment was conducted monaurally. Figure 1 shows the audiograms in the

participants' test ears. All young NH listeners had audiometric pure-tone thresholds (PTTs) ≤ 20 dB

HL at frequencies ranging from 125 Hz to 8 kHz. All older NH listeners had PTTs ≤ 20 dB HL at

frequencies ranging from 125 Hz to 4 kHz, PTTs ≤ 25 dB HL at 6 kHz, and PTTs ≤ 45 dB HL at 8

kHz. The older HI listeners exhibited a high-frequency sloping hearing loss, classified as an N2 or N3

standard audiogram (Bisgaard *et al.*, 2010).[1] All HI listeners had a sensorineural hearing loss, with air-

bone gaps $\leq$ 15 dB at 1, 2, 3, and 4 kHz.[2]

Prior to this study, all listeners took part in the AM frequency selectivity studies conducted by Regev

et al. (2023, 2024a). The corresponding datasets are publicly available from Regev *et al.* (2024c, 2024b).

In those studies, masked-threshold patterns (MTPs) were used to assess the masked detection

threshold of a target AM across different masker center frequencies, providing an indication of AM

frequency selectivity. Figure 2 indicates an example of the MTPs for a 4-Hz target modulation

frequency, where the age-related reduction of AM selectivity is evident on both skirts and the

sharpening of the MTP with hearing loss is evident on the low-frequency skirt (see Regev *et al.*, 2023,

and 2024a for a detailed discussion of the possible origins of these effects.). A measure of AM

frequency selectivity at the individual level can be derived from the MTPs by computing the average

range spanned by the MTP (i.e., the MTP's average dynamic range; DR_{MTP}), where a larger dynamic

range is indicative of greater AM selectivity. This is done by identifying the highest masked threshold

as the peak of the MTP, computing the difference between the peak and the lowest masked threshold

on each side of the peak, and averaging the resulting dynamic range across both sides. If the peak of

the MTP is on either extreme, the DR_{MTP} is taken as the dynamic range across the whole MTP. An

illustration of this derivation is provided in supplementary Figure 1. This measure was derived for the

4-Hz MTP of each listener (with the exception of one older HI listener for whom a MTP at 4 Hz

could not be obtained by Regev *et al.*, 2024a), since the 4-Hz modulation frequency has been suggested

to be particularly important for speech perception (e.g., Drullman *et al.*, 1994; Arai *et al.*, 1999;

Greenberg *et al.*, 2003; Elliott and Theunissen, 2009; Varnet *et al.*, 2017) and showed the largest effect

---

[1] The classification of the audiograms was performed using the lowest mean squared error between the standard audiograms and the collected audiometric thresholds at frequencies from 1 to 8 kHz.
[2] Except for one listener who showed a gap of 20 dB at 1 kHz.

sizes across listener groups in Regev *et al.* (2023, 2024a). The resulting individual quantifications of

AM frequency selectivity are shown in Figure 3 for each listener group. As reported by Regev *et al.*

(2023, 2024a), the MTPs and corresponding dynamic ranges indicate a general decline in AM

frequency selectivity with age, which is partially "compensated" by the presence of hearing loss.

Regev *et al.* (2023, 2024a) also evaluated working-memory capacity using a reverse digit span (RDS)

test, which was administered binaurally following the procedure described in Fuglsang *et al.* (2020).

For the HI listeners, a linear gain was applied based on the Cambridge formula (CamEq; Moore and

Glasberg, 1998), with a maximum gain of 30 dB at any frequency. The older NH listeners exhibited

significantly lower average RDS scores (0.40 on a normalized scale) compared to the young NH and

older HI listeners (0.60 and 0.52, respectively).[3]

All participants provided written informed consent and were offered financial compensation for

their participation. The study received ethical approval from the Science Ethics Committee of the

Capital Region of Denmark (reference H-16036391).

Figure 1 (Color online): Individual and average audiograms of the test ear for each listener group. The

thin dashed lines show the individual audiograms, while the dark solid lines represent the average and

---

[3] This was shown by the post-hoc analysis run on a linear model ANOVA using estimated marginal means. The older NH listeners had significantly lower RDS scores than the young NH and older HI listeners [$M = 0.21$, $SE = 0.05$ dB; $M = 0.13$, $SE = 0.05$ dB, respectively].

standard deviation across listeners. Panel A shows the audiograms for the young NH listeners, with

the black horizontal line indicating the 20 dB HL normal-hearing criterion. Panel B displays the

audiograms for the older NH listeners, also with the black horizontal line denoting the 20 dB HL

normal-hearing criterion. Panel C presents the audiograms for the older HI listeners.

Figure 2 (Color online): Average MTPs and AM detection thresholds for a 4-Hz target modulation

frequency. The results for the young NH group are presented as dashed lines and squares, the older

NH group as dashed lines and triangles, and the older HI group as continuous lines and circles. The

bottom open symbols represent the average AM detection threshold for each group. The means and

standard errors are shown. Data adapted from Regev *et al.* (2023, 2024a).

Figure 3 (Color online): Boxplots displaying the average dynamic range of the 4-Hz MTP (DR$_{MTP}$) for

all listeners. The data for the young NH listeners are represented by squares, older NH by triangles,

and older HI by circles. Data derived from Regev *et al.* (2023, 2024a).

B. Apparatus

The listening test was conducted in a single-walled soundproof booth. The experiment was

conducted using MATLAB (The Mathworks Inc., 2015). The stimuli were digitally played using

SoundMexPro (HörTech, Oldenburg, Germany), converted to analog using an RME Fireface

soundcard (Haimhausen, Germany), and played monaurally through Sennheiser HDA200 headphones

(Wedemark, Germany). The output level was calibrated using a GRAS RA0039 ear simulator (Holte,

Denmark) and the frequency response of the headphones was compensated using an inverse digital

filter. The listeners received no feedback during the test.

    C. Procedure and stimuli

    Speech intelligibility in noise was assessed using the Danish Hearing in Noise Test (HINT; Nielsen

    and Dau, 2011). The HINT speech material consists of 10 test lists and 3 training lists, each containing

    20 five-word sentences. The sentences were spoken by a male speaker with an estimated average

    fundamental frequency (F_0) of 123 Hz. The speech signals were combined with five different masking

    noises: i) a stationary speech-shaped noise (SSN); ii) a fluctuating ICRA-5 noise (Dreschler *et al.*, 2001)

    resembling a single male talker in terms of spectral and temporal characteristics; iii) a single male

    competing talker reading a news piece (F_0 = 118 Hz); iv) a single female competing talker reading a

    news piece (F_0 = 173 Hz); and v) a 'cocktail party' condition. The recordings of the competing talkers

    had previously been edited to reduce long pauses to 65 ms. The cocktail-party noise was generated by

    combining a relatively dry recording from a cafeteria with the recordings of the male and female

    competing talkers mentioned earlier. The levels of the competing talkers were each 3 dB higher than

    that of the cafeteria noise. Prior to mixing, each recording was convolved with an impulse response

    from the Aachen Impulse Response database (office setting with dummy head at 1m distance from

    the loudspeaker; Jeub *et al.*, 2009) to introduce reverberation to the original recordings (T_{60} = 0.37 s).

    The spectra of all recordings (prior to mixing the cocktail-party noise) had been pre-processed to

    match the long-term average spectrum of the HINT speech material as closely as possible. The long-

    term average spectra and modulation spectra of the Danish HINT speech material and of all the

    maskers are presented in Figure 4.

    The SRT for each masking noise was determined by conducting two separate measurements using

    different lists. The final SRT was obtained by averaging the SRTs obtained from the two

    measurements. The order of masker presentation and the lists used for each masker were randomized

across the listeners. Both repetitions for each masker were always conducted consecutively.[4] The SRTs

were estimated at the 50%-correct point using an adaptive 1-up 1-down sentence-based scoring rule.

A response was considered correct if all five words of the sentence were repeated accurately. The step

size in SNR was set at 4 dB for the first four sentences and reduced to 2 dB afterwards. To refine the

tracking procedure, a special rule was implemented for the 5[th] sentence: The change in SNR was

computed as the average of the preceding four SNRs and the predicted following SNR, minus the

immediately preceding SNR. The first sentence was repeated until the listener could accurately repeat

the entire sentence. If the sentence had to be repeated more than four times, the run was terminated,

and the starting SNR was increased by 2 dB before starting again. The SRT of a run was calculated as

the average SNR across sentences 5 to 20 plus the SNR required for a fictitious 21[st] sentence,

determined by scoring the 20[th] sentence (Soli and Wong, 2008). The scoring was performed by a

trained native Danish speaker.

The target sentences were played at a fixed level of 60 dB SPL. The masking noise was looped

over the recording, played continuously, and adjusted in level to achieve the desired SNR. For the HI

listeners, linear amplification was added to both the speech and masking noise according to the

Cambridge formula (CamEq; Moore and Glasberg, 1998), with a maximum gain of 30 dB at any

frequency. Each participant underwent training using all three training lists with the SSN. The starting

SNR for each masker and each run was adjusted to be 6 dB lower than the expected SRTs. The

expected SRTs were determined based on a pilot test conducted with young NH listeners and were

defined relative to the SRT measured from the final training run. Therefore, the starting SNR was 6

dB lower than the SRT of the listener's last training list for the SSN, male competing talker, and

---

[4] Due to a mistake in the audibility compensation, one of the older HI listeners was tested twice on the entire HINT corpus. However, more than 7 weeks separated the two testing sessions. Nielsen and Dau (2011) reported an average improvement in SRT of only 0.5 dB across two sessions separated by three weeks. Hence, the current case was likely not affected by memorization of the speech material.

cocktail party conditions; 9 dB lower for the ICRA noise; and 15 dB lower for the female competing

talker condition.

Figure 4 (Color online): Long-term average spectra (top row) and modulation spectra (bottom row)

of the speech material and maskers. Each panel shows the spectra for the HINT speech material (black

line) and one of the maskers (light blue line). Panels A and F: SSN; Panels B and G: ICRA-5; Panels

C and H: Male competing talker; Panels D and I: Female competing talker; Panels E and J: Cocktail-

party condition.

D. Statistical analysis

1. Group-level analysis of SRTs

The degree of variability between the two SRT estimates obtained for each masker was assessed

by calculating the standard deviation across the absolute differences in SRTs, $\sigma_{\Delta|SRT|}$. Notably,

although small, the variability across all maskers was highest for the young NH listeners ($\sigma_{\Delta|SRT|}$ =

2.43 dB) compared to the older NH listeners ($\sigma_{\Delta|SRT|}$ = 1.76 dB) and older HI listeners ($\sigma_{\Delta|SRT|}$ = 1.49

dB). Nevertheless, the degree of variability between SRT estimates was considered sufficiently small

for all statistical analyses to be conducted using the SRTs averaged across both repetitions for each

listener and masker. Analyses of variance (ANOVAs) of linear mixed-effects model fittings were

performed to examine the differences between the listener groups. Two sets of analysis were

conducted: First, the older NH group was compared to the young NH listeners to isolate the effects

of age from hearing loss, and second, the older HI group was compared to the older NH listeners to

isolate the effects of hearing loss from age-related effects. In both analyses, the listeners were treated

as a random effect, while group classification and masker were considered as fixed effects.[5] Post-hoc

analyses were carried out using estimated marginal means, and a Holm-Bonferroni correction criterion

was applied to account for multiple comparisons. The significance level for all analyses was set at 5%.

2. Regression analysis

To assess whether a change in AM frequency selectivity contributes to the variance in SRTs, a

multilinear regression model was built to relate age, audibility, working memory capacity, and AM

frequency to SRTs across listeners. The SRTs from the condition showing the largest effect size across

young and older NH listeners were chosen as the model's outcome variable. The measure of age was

taken as the listeners' chronological age at the time of testing. The measure of audibility was taken as

the 4-frequency pure-tone average threshold (PTA; where the audiometric thresholds at 0.5, 1, 2, and

4 kHz are averaged). The measure of working memory capacity was taken as the listeners' RDS score.

Finally, the measure of AM frequency selectivity was taken as the listeners' DR_{MTP} at a 4-Hz target

modulation frequency, shown in Figure 3. The model included all young and older NH listeners and

8 of the 9 older HI listeners, since a 4-Hz MTP could not be obtained by Regev *et al.* (2024a) for one

of them. The multilinear regression model was built in steps, starting from a model including only age,

---

[5] Levene's test was used to assess the homoskedasticity of the residuals and Shapiro-Wilk's test was used to assess the normality of residual's distribution. Shapiro-Wilk's test was not passed for either model, likely due to the relatively small group sizes and to the presence of outliers in the older HI data. However, no other aspect of the data justified the removal of the outliers, and they were therefore kept in the dataset. Furthermore, the average fittings of the models to the experimental data were found to be accurate, and the models were hence considered as appropriate.

by assessing whether adding the other predictors and their interactions significantly reduced the

model's residual sum of squares, using an F-test. The predictors were added in the above order (PTA,

RDS, and DR_{MTP}) and the addition of their interactions was only assessed if the main effect

significantly improved the model. The significance level was set at 5%.

III.   RESULTS

A. Speech intelligibility

The SRTs for all listeners and groups are shown in Figure 5 for each masker. The young NH

listeners are represented by squares, older NH listeners by triangles, and older HI listeners by circles.

The statistical analysis comparing young and older NH listeners, focusing on the effects of age,

revealed significant main effects of masker [$F_{4,68}$ = 274.2, p < 0.001] and listener group [$F_{1,17}$ = 17.56,

p < 0.001], as well as a significant interaction between both factors [$F_{4,68}$ = 9.28, p < 0.001]. In the

analysis comparing older NH and older HI listeners, examining the effects of hearing loss, significant

main effects of masker [$F_{4,64}$ = 141.6, p < 0.001] and listener group [$F_{1,16}$ = 12.86, p = 0.002], as well

as a significant interaction between both factors [$F_{4,64}$ = 5.32, p < 0.001], were observed. Post-hoc

analyses revealed that the older NH listeners had significantly higher SRTs than the young NH

listeners for the ICRA-5 condition [Effect size (M) = 1.39 dB, standard error (SE) = 0.68 dB], male

competing talker [M = 4.53 dB, SE = 0.68 dB], female competing talker [M = 2.70 dB, SE = 0.68

dB], and cocktail-party condition [M = 1.53 dB, SE = 0.68 dB]. Additionally, the older NH listeners

had significantly lower SRTs than the older HI listeners for the SSN condition [M = 2.36 dB, SE =

1.14 dB], ICRA-5 condition [M = 4.30 dB, SE = 1.14 dB], male competing talker [M = 3.60 dB, SE

= 1.14 dB], and female competing talker [M = 5.52 dB, SE = 1.14 dB]. The increase in SRT for the

older HI listeners in the cocktail-party condition approached, but did not reach, significance [M =

2.24 dB, SE = 1.14 dB, p = 0.06]. Furthermore, the post-hoc analyses revealed that the female

competing talker resulted in significantly lower SRTs than all other maskers for all listener groups.

The male competing talker yielded significantly higher SRTs than all other maskers for the older NH

and older HI listeners (except for the cocktail-party condition in the older NH group), while the

cocktail-party condition yielded the highest SRTs for the young NH listeners. Finally, the ICRA-5

noise resulted in significantly lower SRTs than the SSN for the young and older NH listeners, but not

for the older HI listeners. Additional details and results from all post-hoc analyses can be found in

Appendix A.

Figure 5 (Color online): Boxplots displaying the SRTs for all listeners. Each panel corresponds to a

specific masker: Panel A: Speech-shaped noise (SSN), Panel B: ICRA-5 fluctuating noise, Panel C:

Single male competing talker (same sex), Panel D: Single female competing talker (different sex), Panel

E: Cocktail-party noise. The SRTs for the young NH listeners are represented by squares, older NH

by triangles, and older HI by circles. The stars indicate the level of statistical significance of the

differences between the groups separated by the horizontal black lines (* $p < 0.05$, ** $p < 0.01$, *** p

< 0.001), while "n.s." denotes non-significant differences. Note that the young NH and older HI

groups were not compared in the statistical analysis.

The multilinear regression analysis included the SRTs obtained with the male competing talker as

the outcome variable, as that condition yielded the largest difference in SRTs between young and older

NH listeners, as well as highest SRTs for both the older NH and HI listeners. The results showed that

the best-fitting model included the main effects of age, PTA, and DR_{MTP}, while RDS did not contribute

to explaining the variance in SRTs. Adding DR_{MTP} to age and PTA [$F_{1,23}$ = 9.693, p = 0.005]

significantly increased the amount of variance in SRT explained by the model (adjusted R^2) by 8.4

percentage points to a total of 77%. No interaction term significantly increased the model's R^2. Figure

6 shows the scatter plots of the fitted SRTs as a function of the measured SRTs for each one of the

models iteratively built (i.e., including age, age + PTA, or age + PTA + DR_{MTP} as predictors) in

separate panels. The adjusted R^2 for each model is indicated in each panel. The best-fitting model's

regression coefficient (i.e., β weights) was larger for age (0.773) than for PTA and DR_{MTP} (0.303 and

0.350, respectively). The structure coefficients, i.e., the correlation of each predictor with the outcome

variable estimated from all predictors in the model, showed that both age and PTA were positively

correlated with the fitted SRTs, while DR_{MTP} was negatively correlated (0.896, 0.842, and -0.157,

respectively).

Since all predictors in the best-fitting model (age, PTA, and DR_{MTP}) were likely somewhat collinear,

a commonality analysis (Capraro and Capraro, 2001; Ray-Mukherjee *et al.*, 2014) was implemented to

assess the R^2 either attributable uniquely to each predictor or shared between them. The outcome

showed that age had the greatest unique R^2 (22 %), while PTA and DR_{MTP} had smaller unique R^2 (4.3

and 8.3 %, respectively). Furthermore, the commonality analysis revealed that age and PTA were

highly collinear whereby they shared a large common R^2 (48.8 %), and that age and DR_{MTP} were also

somewhat collinear but shared a negative R^2 (-7.9%). The detailed results of the regression and

commonality analyses can be found in Appendix B.

Figure 6: Scatter plot of the fitted SRTs as a function of the measured SRTs for each one of the

regression models assessed. The predictors included in the model as well as the adjusted R^2 are

indicated in each panel. The diagonal line indicates identity. The models were iteratively built to include

only age (left panel), age and PTA (middle panel), and finally age, PTA, and dynamic range of the 4-

Hz MTP (DR_{MTP}; right panel). The last model provided the best fit of the SRTs.

IV.   DISCUSSION

The aim of the present study was to investigate whether the findings from speech intelligibility

tests would be consistent with the AM frequency selectivity data collected in previous studies by Regev

et al. (2023, 2024a). These previous investigations demonstrated an age-related decline in AM

frequency selectivity and an effective sharpening of AM tuning with hearing loss (see Figure 2 and

Figure 3). A subset of the listener cohort measured by Regev *et al.* (2023, 2024a) was considered here.

It was expected that the age-related reduction in AM frequency selectivity would correspond to poorer

speech intelligibility in older compared to young NH listeners. Conversely, the sharper AM tuning

observed in older HI listeners, assumed to be an "apparent" effect resulting from the loss of cochlear

compression (Regev *et al.*, 2024a), was not anticipated to result in better speech intelligibility compared

to the older NH listeners.

The results revealed an overall decline in speech intelligibility with age, which was further

exacerbated by the presence of hearing loss. These findings are consistent with previous studies that

have shown a deterioration of speech intelligibility with age (e.g., Rajan and Cainer, 2008; Helfer and

Vargo, 2009; Schoof and Rosen, 2014; Füllgrabe *et al.*, 2015; Gordon-Salant and Cole, 2016; Goossens

et al., 2017; Decruy *et al.*, 2019) and hearing loss (e.g., Glasberg and Moore, 1989; Humes and

Christopherson, 1991; Frisina and Frisina, 1997; see Humes and Dubno, 2010 for a review).

Additionally, a multilinear regression model relating the SRTs obtained with a male competing talker

to age, PTA, and DR_{MTP} (a measure of AM frequency selectivity) showed that DR_{MTP} significantly

contributed to explaining the variance in SRTs, increasing the model's R^2 by 8 percentage points.

The observed increase in SRTs with age aligns with the age-related reduction in AM frequency

selectivity reported by Regev *et al.* (2023). Broader modulation filters, indicative of reduced AM

selectivity, may lead to increased AM masking, resulting in a decrease in the SNR_{env} at the output of

these filters and potentially poorer speech intelligibility in noise. However, the lack of a significant

age-related increase in SRT for the SSN condition is inconsistent with this notion. Since the SSN

contains intrinsic envelope fluctuations that act as a relatively broadband modulation masker (as

argued by Stone *et al.*, 2012), one might have expected the broader AM tuning found in older NH

listeners to lead to an increase in AM masking. However, the SRTs obtained with the SSN likely reflect

aspects of EM rather than AM masking. This is consistent with the extensive overlap observed in the

audio spectra of the SSN and HINT speech material (Figure 4, panel A), compared to the highly

positive SNR$_{env}$ observed when comparing their modulation spectra at low modulation frequencies

(Figure 4, panel F). These findings indicate that the SSN provided more EM than AM masking.

Conversely, the differences in modulation spectra between the target speech and masking noise were

much smaller for the other maskers, suggesting that AM masking (and thus potentially AM frequency

selectivity) may have played a more prominent role in those cases (Figure 4, panels G-J). Notably, the

older NH listeners exhibited a reduced release from masking when exposed to the fluctuating ICRA-

5 noise compared to the young NH listeners, consistent with previous studies (Dubno *et al.*, 2002,

2003; Grose *et al.*, 2009; Goossens *et al.*, 2017). In addition to potentially being the result of an age-

related increase in forward masking (e.g., Gifford *et al.*, 2007), the reduced release from masking

observed in the older NH listeners may also be attributed to increased AM masking due to the loss of

AM frequency selectivity, which may have limited the benefit derived from the slow temporal

fluctuations in the ICRA-5 condition.

Regarding the effects of hearing loss, the higher SRTs observed in the older HI listeners compared

to the older NH group are consistent with the expectation that the increase in AM frequency selectivity

observed in HI listeners (as reported in Regev *et al.*, 2024a) would *not* lead to improved speech

intelligibility. Regev *et al.* (2024a) argued that the greater AM frequency selectivity with hearing loss

was an apparent effect, resulting from improved detection of the AM target during temporal dips in

the masker due to the loss of cochlear compression. Since the loss of compression is also associated

with other supra-threshold deficits, such as loudness recruitment, poorer recovery from forward

masking, and reduced frequency selectivity (Glasberg and Moore, 1986, 1989; Oxenham and Moore,

1997; Moore and Oxenham, 1998), it is evident that the process potentially underlying the apparent

improved AM frequency selectivity is also linked to the mechanisms underlying poorer speech

intelligibility in HI listeners. However, while poorer speech intelligibility was expected in the HI

listeners, disentangling the potential impact of the effective sharpening in AM tuning with hearing loss

on speech intelligibility is not straightforward. It remains unclear whether the process underlying the

apparent sharpening of AM tuning with hearing loss a) improves SRTs but is outweighed by co-

occurring supra-threshold deficits typically associated with hearing loss, or b) itself ultimately impairs

speech intelligibility.

The results of the multilinear regression analysis further supported the existence of a link between

alterations in AM frequency selectivity and speech intelligibility challenges in noise for older listeners.

The addition of DR_{MTP} to the model significantly increased the amount of variance in SRTs explained

by the model. Furthermore, the negative structure coefficient for DR_{MTP} indicated that an increase in

the SRTs estimated by the best-fitting model (which included age, PTA, and DR_{MTP}) is linked with a

decrease in DR_{MTP}. This finding aligned with the expectation that poorer AM frequency selectivity

(quantified by a lower DR_{MTP}) would be linked with higher SRTs.[6] The link between poorer AM

frequency selectivity and poorer SRTs across listeners is also broadly consistent with the assumption

that the sharper MTPs observed in older HI listeners reflect a perceptual benefit inherited from the

loss of cochlear compression (represented by PTA in the regression model), rather than a genuine

sharpening of the hypothetical modulation filter (Regev *et al.*, 2024a).

To address the likely collinearity across predictors, the commonality analysis revealed that DR_{MTP}

acts as a suppressor, as shown by the negative R^2 shared between DR_{MTP} and age (Capraro and

Capraro, 2001). As such, the addition of DR_{MTP} to the model helped increase the R^2 uniquely

attributable to age, thus increasing the model's total R^2. Within the model, age can be seen as a "noisy

term" representing the sum of all factors underlying auditory processing deficits in older listeners that

are distinct from the effects of peripheral hearing loss. Such factors, which are likely somewhat

collinear, may include cochlear synaptopathy (Parthasarathy *et al.*, 2019a), reduced neural inhibition

---

[6] Structure coefficients are employed here since they have been suggested to be more robust than regression coefficients in cases involving multicollinearity among predictors (Thompson and Borrello, 1985; Kraha *et al.*, 2012)

(Caspary *et al.*, 2008), or poorer temporal envelope processing (Anderson and Karawani, 2020) such

as a loss of AM frequency selectivity (Regev *et al.*, 2023). Hence, the greater R^2 resulting from the

explicit addition of a measure of AM frequency selectivity to the regression model can reasonably be

interpreted as resulting from a "denoising" of the age term, allowing it to more accurately account for

the variance in SRTs.

However, it should be noted that the significant contribution of DR_{MTP} to the regression model

was not found when the model was applied to SRTs for other maskers than the male competing talker,

or DR_{MTP} derived from MTPs at other target modulation rates than 4 Hz. While the reasons for this

discrepancy remain unclear, the specific choice of the SRTs for the male competing talker and the

DR_{MTP} derived for a 4-Hz modulation rate was soundly motivated by the data. Indeed, the male

competing talker yielded the largest difference in SRTs between young and older NH listeners, as well

as the highest SRTs for both the older NH and HI listeners, compared to all other maskers.

Furthermore, the 4-Hz MTP also showed the largest differences in AM selectivity between young and

older NH listeners (Regev *et al.*, 2023) and was assumed to reflect the prominent role of the 4-Hz

modulation rate in speech perception (e.g., Elliott and Theunissen, 2009; Varnet *et al.*, 2017).

Notably, the present findings indicate that declines in speech intelligibility with age and hearing

loss are most prominent in contexts that involve high cognitive demand and informational masking

(IM), such as in the presence of competing-talker maskers. IM encompasses various factors that

complicate the separation of the target and masker speech signals, such as stream segregation

(Brungart et al., 2001; Darwin et al., 2003; Schneider et al., 2007; Kidd et al., 2016; see Kidd et al.,

2008 for a review). In the groups considered here, the older NH listeners exhibited significantly lower

RDS scores compared to the young NH listeners. Such reduced cognitive abilities have been suggested

to mediate the age-related increase of SRTs under conditions of IM (Rajan and Cainer, 2008; Gordon-

Salant and Cole, 2016; Goossens *et al.*, 2017). Therefore, the poorer speech intelligibility observed in

the older NH listeners in the present study may be partly attributed to a reduction in working-memory

capacity. However, the older NH listeners also had higher SRTs than the young NH listeners in the

ICRA-5 condition, which does not exert large amounts of IM or high cognitive demand. Furthermore,

the multilinear regression analysis showed that the RDS scores did not contribute to explaining the

variance in SRTs for the male competing talker across all listeners. Theses finding are broadly

consistent with studies reporting no significant correlation between reduced cognitive abilities and

increased SRTs in babble for older listeners (e.g., Schoof and Rosen, 2014), suggesting that other

factors may also contribute to the age-related decline in speech intelligibility. These factors may be

associated with age-related temporal processing deficits, as suggested by perceptual (Pichora-Fuller

and Souza, 2003; Pichora-Fuller and MacDonald, 2008; Helfer and Vargo, 2009; Füllgrabe et al., 2015;

see Fitzgibbons and Gordon-Salant, 2010 for a review) and neurophysiological research (e.g.,

Anderson et al., 2012; Decruy et al., 2019; Parthasarathy et al., 2019b; Roque et al., 2019; for reviews

see Parthasarathy et al., 2019a and Anderson and Karawani, 2020). The hypothesis that the age-related

reduction in AM frequency selectivity (as reported by Regev *et al.*, 2023) may contribute to increased

SRTs aligns with the notion that temporal processing deficits underlie the challenges in speech

intelligibility associated with age.

In the case of hearing loss, the differences in SRTs between older NH and older HI listeners were

also largest for the fluctuating and competing-talker maskers. This observation is consistent with the

reduced release from masking commonly found in HI listeners (e.g., Bacon *et al.*, 1998; Bernstein and

Grant, 2009; Strelcyk and Dau, 2009; Goossens *et al.*, 2017), as well as with the suggestion that the

effects of hearing loss on speech intelligibility are most pronounced in cognitively demanding settings

(e.g., Schneider *et al.*, 2010; Goossens *et al.*, 2017). Although the older HI listeners showed higher RDS

scores than their older NH counterparts, they did exhibit a marginally reduced working-memory

capacity compared to the young NH listeners. Hence, the large effect of hearing loss in conditions

involving IM may be partly connected to the age-related cognitive decline discussed earlier.

Conversely, the older HI listeners also had higher SRTs than their older NH counterparts in the SSN

and ICRA-5 conditions, which are commonly assumed not to involve high cognitive demand.

Notably, the multilinear regression analysis showed that the RDS scores, when added to age and

PTA, did not contribute to explaining the variance in SRTs for the male competing talker across

listeners. This suggests that working-memory capacity did not influence the speech intelligibility results

obtained in the present study. However, working-memory capacity is only one of several measures of

cognitive ability that may be reduced in older listeners, thus contributing to speech intelligibility

challenges in noise (e.g., Dryden *et al.*, 2017). Overall, while reduced cognitive ability in older listeners

may have partly mediated the observed effects of age and hearing loss on speech intelligibility, the

results also indicate that supra-threshold deficits influenced the SRTs, likely related to reduced

cochlear compression and temporal envelope processing deficits.

Although the data from the present study revealed significant group-level and regression results

that support a relationship between changes in AM tuning and speech intelligibility in noise for older

listeners, it is important to consider several co-occurring factors that may have influenced the findings.

Even though all the older NH listeners exhibited clinically normal hearing up to 4 kHz, their

audiometric thresholds were on average 8 dB higher than those of the young NH listeners between

125 Hz and 8 kHz, with the greatest differences occurring at 8 kHz. Although these differences in

audiometric thresholds are small, previous studies have suggested that they can have a considerable

effect on speech intelligibility (Dubno and Ahlstrom, 1997; Dubno *et al.*, 2000), possibly due to related

supra-threshold processing deficits (e.g., Léger *et al.*, 2012). Moreover, the older NH listeners included

in the present study may have experienced reduced audibility at extended high frequencies (> 8 kHz),

which have also been suggested to play a role in speech intelligibility (e.g., Hunter *et al.*, 2020; Polspoel

et al., 2022). These aspects may have contributed to the observed age-related increase in SRTs.

Concerning the experimental approach, since the tests provided listeners with cues beyond the

temporal envelope, factors such as IM, EM, and stream segregation likely interacted with age- and

hearing loss-related perception deficits, as well as the age-related reduction in working-memory

capacity discussed above. Therefore, the connection between SRTs and AM frequency selectivity

suggested by the present results should be investigated further. An experimental paradigm limiting the

perceptual cues available in the speech intelligibility test to the temporal envelope may help establish

such a link with more certainty.

V.   SUMMARY AND CONCLUSION

Speech intelligibility in noise was examined in three groups: young listeners with normal hearing

(aged 22-28 years), older listeners with normal hearing (aged 57-77 years), and older listeners with

hearing impairment (aged 64 – 77) years. All participants had previously taken part in AM masking

studies investigating AM frequency selectivity, which revealed an age-related reduction in AM tuning

and an effective sharpening in AM tuning with hearing loss (Regev *et al.*, 2023, 2024a). This study

aimed to determine if the effects of age and hearing loss on speech intelligibility in noise support the

hypothesis that the age-related reduction in AM tuning negatively impacts speech intelligibility, while

the improvement in AM tuning with hearing loss does not provide a benefit.

Consistent with the hypothesis, the results showed an age-related decline and an additional

negative effect of hearing loss on speech intelligibility, especially for fluctuating maskers. Furthermore,

a multilinear regression analysis showed that adding a measure of AM frequency selectivity to a model

using age and audibility as predictors significantly increased the amount of variance in SRTs explained.

This finding supported the assumption that an age-related loss of AM frequency selectivity is linked

to speech intelligibility difficulties in noise, once audibility is accounted for. Moreover, the results

broadly aligned with the proposition made by Regev *et al.* (2024a) that the improved AM tuning in

listeners with hearing impairment is an "apparent" effect resulting from reduced peripheral

compression, which is known to contribute to other supra-threshold deficits that impair speech intelligibility.

Overall, this study's results support the notion that an age-related reduction in AM frequency selectivity may contribute to speech intelligibility challenges in noise and warrant further investigation into this potential mechanism. To more directly assess the effects of AM tuning, and a loss thereof, on speech intelligibility, future studies may focus on speech-intelligibility testing paradigms that specifically examine envelope-based cues.

ACKNOWLEDGMENTS

The authors thank Borgný Súsonnudóttir Hansen for her contribution to the data collection. The authors also thank Christian Stender Simonsen for kindly sharing the experimental framework used to run the listening test, as well as several of the signal recordings, and Jens Hjortkjær and Jonatan Märcher-Rørsted for kindly providing their implementations of the reverse digit span test and of the Cambridge formula (CamEq; Moore and Glasberg, 1998). Finally, the authors thank Raul Sanchez-Lopez for helpful insights and discussions. This study was supported by the Oticon Centre of Excellence for Hearing and Speech Sciences (CHeSS).

AUTHOR DECLARATIONS

Conflict of Interest

The authors have no conflict of interest to report.

Ethics Approval

All participants provided written informed consent and were offered financial compensation for their participation. Ethical approval for the study was provided by the Science Ethics Committee of the Capital Region in Denmark (reference H-16036391).

DATA AVAILABILITY

This manuscript has been submitted to the Journal of the Acoustical Society of America. Once it

has been accepted, the data that support the findings of this study will be openly available from Regev,

J., Zaar, J., Relaño-Iborra, H., and Dau, T. (2024). "Dataset for: 'Investigating the effects of age and

hearing loss on speech intelligibility and amplitude modulation frequency selectivity'", Technical

University of Denmark. Dataset. https://doi.org/10.11583/DTU.25771884.

537
APPENDIX

A. Post-hoc analyses

Table 1 summarizes the results of the post-hoc analyses that compared the SRTs obtained with

young and older NH listeners for each masker.

Table 1: Post-hoc analysis of the differences in SRT between young and older NH listeners for each

masker. The effect sizes (M), standard errors (SE, and corrected p-values are reported. The significant

p-values are highlighted in bold.

Masker	M [dB]	SE [dB]	Corrected p-value
SSN	0.75		0.279
ICRA-5	1.39		**0.047**
Male competing	4.53	0.68	**< 0.001**
Female competing	2.70		**< 0.001**
Cocktail party	1.53		**0.030**

548   Table 2 summarizes the results of the post-hoc analyses that compared the SRTs obtained with

549   older NH and older HI listeners for each masker.

551   Table 2: Post-hoc analysis of the differences in SRT between older NH and older HI listeners for each

552   masker. The effect sizes (M), standard errors (SE, and corrected p-values are reported. The significant

553   p-values are highlighted in bold.

Masker	M [dB]	SE [dB]	Corrected p-value
SSN	2.36		**0.048**
ICRA-5	4.30		**< 0.001**
Male competing	3.60	1.14	**0.004**
Female competing	5.52		**< 0.001**
Cocktail party	2.24		0.060

Table 3 summarizes the results of the post-hoc analyses that compared the SRTs obtained with

the different maskers for each of the three listener groups (young NH, older NH, and older HI). Since

this post-hoc analysis was run twice for the older NH listeners (once for each statistical model applied),

the values reported reflect the most conservative p-values and standard errors, obtained from the

model fitted to the older NH and older HI listeners. However, the effect sizes and overall statistical

outcomes did not differ between the analyses run in the two models.

565   Table 3: Post-hoc analysis of the differences in SRT between older the different maskers for each

566   group. The effect sizes (M), standard errors (SE, and corrected p-values are reported. The significant

567   p-values are highlighted in bold.

Listener group	Maskers compared	M [dB]	SE [dB]	Corrected p-value
Young NH	SSN - ICRA-5	2.35	0.48	**< 0.001**
	SSN – Male competing	0.83		0.086
	SSN – Female competing	9.23		**< 0.001**
	SSN – Cocktail party	-1.42		**0.008**
	ICRA-5 – Male competing	-1.52		**0.006**
	ICRA-5 – Female competing	6.88		**< 0.001**
	ICRA-5 – Cocktail party	-3.77		**< 0.001**
	Male competing – Female competing	8.40		**< 0.001**
	Male competing – Cocktail party	-2.25		**< 0.001**
	Female competing – Cocktail party	-10.65		**< 0.001**
Older NH	SSN - ICRA-5	1.71	0.60	**0.011**
	SSN – Male competing	-2.95		**< 0.001**
	SSN – Female competing	7.28		**< 0.001**
	SSN – Cocktail party	-2.20		**0.001**
	ICRA-5 – Male competing	-4.66		**< 0.001**
	ICRA-5 – Female competing	5.57		**< 0.001**
	ICRA-5 – Cocktail party	-3.91		**< 0.001**
	Male competing – Female competing	10.23		**< 0.001**
	Male competing – Cocktail party	0.75		0.212

	Female competing – Cocktail party	-9.48		**< 0.001**
Older HI	SSN - ICRA-5	-0.23	0.60	0.696
	SSN – Male competing	-4.20		**< 0.001**
	SSN – Female competing	4.12		**< 0.001**
	SSN – Cocktail party	-2.09		**0.003**
	ICRA-5 – Male competing	-3.96		**< 0.001**
	ICRA-5 – Female competing	4.35		**< 0.001**
	ICRA-5 – Cocktail party	-1.85		**0.006**
	Male competing – Female competing	8.32		**< 0.001**
	Male competing – Cocktail party	2.11		**0.003**
	Female competing – Cocktail party	-6.20		**< 0.001**

 B. Regression analysis

    Table 4 summarizes the details of the best-fitting model used in the multilinear regression analysis.

 Table 4: Details of the multilinear regression analysis relating the SRTs obtained for the male

 competing talker to age, PTA, and DR_{MTP}. For each predictor, the table reports the standardized

 regression coefficients (β), the 95%-confidence interval for β (lower | upper), as well as the structure

 coefficients (r_s).

Predictor	Standardized β	95% CI β (lower | upper)	r_s
Age	0.7726	0.4892 | 1.056	0.8964
PTA	0.3025	0.0452 | 0.5598	0.8422
DR_{MTP}	0.3504	0.111 | 0.5898	-0.1569

    Table 5 summarizes the details of the commonality analysis conducted on the best-fitting model

 (relating SRTs to age, PTA, and DR_{MTP}), which identifies the R^2 either uniquely attributable to each

 predictor or shared among them.

 Table 5: Outcome of the commonality analysis applied on the best-fitting model, that related SRTs

 obtained for a male competing talker to age, PTA, and the 4-Hz MTP's average dynamic range

 (DR_{MTP}). The table reports the amount of variance in SRTs explained (R^2) either uniquely explained

 by each predictor (single variables) or commonly shared between predictors (two or more variables).

Predictor	Age	PTA	DR_{MTP}	Age and PTA	Age and DR_{MTP}	PTA and DR_{MTP}	Age, PTA, and DR_{MTP}	Total
R^2 [%]	21.96	4.33	8.30	48.78	-7.89	2.49	-1.00	76.97

REFERENCES

Anderson, S., and Karawani, H. (**2020**). "Objective evidence of temporal processing deficits in older adults," Hear Res, **397**, 108053. doi:10.1016/j.heares.2020.108053

Anderson, S., Parbery-Clark, A., White-Schwoch, T., and Kraus, N. (**2012**). "Aging Affects Neural Precision of Speech Encoding," J Neurosci, **32**, 14156–14164. doi:10.1523/JNEUROSCI.2176-12.2012

Arai, T., Pavel, M., Hermansky, H., and Avendano, C. (**1999**). "Syllable intelligibility for temporally filtered LPC cepstral trajectories," J Acoust Soc Am, **105**, 2783–2791. doi:10.1121/1.426895

Bacon, S. P., and Grantham, D. W. (**1989**). "Modulation masking: Effects of modulation frequency, depth, and phase," J Acoust Soc Am, **85**, 2575–2580. doi:10.1121/1.397751

Bacon, S. P., Opie, J. M., and Montoya, D. Y. (**1998**). "The Effects of Hearing Loss and Noise Masking on the Masking Release for Speech in Temporally Complex Backgrounds," J Speech Lang Hear Res, **41**, 549–563. doi:10.1044/jslhr.4103.549

Bernstein, J. G. W., Danielsson, H., Hällgren, M., Stenfelt, S., Rönnberg, J., and Lunner, T. (**2016**). "Spectrotemporal Modulation Sensitivity as a Predictor of Speech-Reception Performance in Noise With Hearing Aids," Trends Hear, **20**, 233121651667038. doi:10.1177/2331216516670387

Bernstein, J. G. W., and Grant, K. W. (**2009**). "Auditory and auditory-visual intelligibility of speech in fluctuating maskers for normal-hearing and hearing-impaired listeners," J Acoust Soc Am, **125**, 3358. doi:10.1121/1.3110132

Bisgaard, N., Vlaming, M. S. M. G., and Dahlquist, M. (**2010**). "Standard Audiograms for the IEC 60118-15 Measurement Procedure," Trends Amplif, **14**, 113–120. doi:10.1177/1084713810379609

Brungart, D. S., Simpson, B. D., Ericson, M. A., and Scott, K. R. (**2001**). "Informational and

energetic masking effects in the perception of multiple simultaneous talkers," J Acoust

Soc Am, **110**, 2527–2538. doi:10.1121/1.1408946

Capraro, R. M., and Capraro, M. M. (**2001**). "Commonality Analysis: Understanding Variance

Contributions to Overall Canonical Correlation Effects of Attitude Toward Mathematics

on Geometry Achievement," Multiple Linear Regression Viewpoints, **27**, 16–23.

Caspary, D. M., Ling, L., Turner, J. G., and Hughes, L. F. (**2008**). "Inhibitory neurotransmission,

plasticity and aging in the mammalian central auditory system," J Exp Biol, **211**, 1781–

1791. doi:10.1242/jeb.013581

Chabot-Leclerc, A., Jørgensen, S., and Dau, T. (**2014**). "The role of auditory spectro-temporal

modulation filtering and the decision metric for speech intelligibility prediction," J

Acoust Soc Am, **135**, 3502–3512. doi:10.1121/1.4873517

Darwin, C. J., Brungart, D. S., and Simpson, B. D. (**2003**). "Effects of fundamental frequency

and vocal-tract length changes on attention to one of two simultaneous talkers," J Acoust

Soc Am, **114**, 2913. doi:10.1121/1.1616924

Dau, T., Kollmeier, B., and Kohlrausch, A. (**1997**). "Modeling auditory processing of amplitude

modulation. I. Detection and masking with narrow-band carriers," J Acoust Soc Am, **102**,

2892–2905. doi:10.1121/1.420344

Dau, T., Verhey, J., and Kohlrausch, A. (**1999**). "Intrinsic envelope fluctuations and modulation-

detection thresholds for narrow-band noise carriers," J Acoust Soc Am, **106**, 2752–2760.

doi:10.1121/1.428103

Decruy, L., Vanthornhout, J., and Francart, T. (**2019**). "Evidence for enhanced neural tracking

of the speech envelope underlying age-related speech-in-noise difficulties," J

Neurophysiol, **122**, 601–615. doi:10.1152/jn.00687.2018

Decruy, L., Vanthornhout, J., and Francart, T. (**2020**). "Hearing impairment is associated with

enhanced neural tracking of the speech envelope," Hear Res, **393**, 107961.

doi:10.1016/j.heares.2020.107961

Dorman, M. F., Loizou, P. C., and Rainey, D. (**1997**). "Speech intelligibility as a function of the

number of channels of stimulation for signal processors using sine-wave and noise-band

outputs," J Acoust Soc Am, **102**, 2403–2411. doi:10.1121/1.419603

Dreschler, W. A., Verschuure, H., Ludvigsen, C., and Westermann, S. (**2001**). "ICRA Noises:

Artificial Noise Signals with Speech-like Spectral and Temporal Properties for Hearing

Instrument Assessment," Int J Audiol, **40**, 148–157. doi:10.3109/00206090109073110

Drullman, R., Festen, J. M., and Plomp, R. (**1994**). "Effect of temporal envelope smearing on

speech reception," J Acoust Soc Am, **95**, 1053–1064. doi:10.1121/1.408467

Dryden, A., Allen, H. A., Henshaw, H., and Heinrich, A. (**2017**). "The Association Between

Cognitive Performance and Speech-in-Noise Perception for Adult Listeners: A

Systematic Literature Review and Meta-Analysis," Trends Hear, **21**, 233121651774467.

doi:10.1177/2331216517744675

Dubbelboer, F., and Houtgast, T. (**2008**). "The concept of signal-to-noise ratio in the modulation

domain and speech intelligibility," J Acoust Soc Am, **124**, 3937–3946.

doi:10.1121/1.3001713

Dubno, J. R., and Ahlstrom, J. B. (**1997**). "Additivity of Multiple Maskers of Speech," In W.

Jesteadt (Ed.), Modeling Sensorineural Hearing Loss, Routledge, New York, pp. 253–

272. doi:10.4324/9781315789392-21

Dubno, J. R., Ahlstrom, J. B., and Horwitz, A. R. (**2000**). "Use of context by young and aged

adults with normal hearing," J Acoust Soc Am, **107**, 538–546. doi:10.1121/1.428322

Dubno, J. R., Dirks, D. D., and Morgan, D. E. (**1984**). "Effects of age and mild hearing loss on
speech recognition in noise," J Acoust Soc Am, **76**, 87–96. doi:10.1121/1.391011

Dubno, J. R., Horwitz, A. R., and Ahlstrom, J. B. (**2002**). "Benefit of modulated maskers for
speech recognition by younger and older adults with normal hearing," J Acoust Soc Am,
111, 2897–2907. doi:10.1121/1.1480421

Dubno, J. R., Horwitz, A. R., and Ahlstrom, J. B. (**2003**). "Recovery from prior stimulation:
Masking of speech by interrupted noise for younger and older adults with normal
hearing," J Acoust Soc Am, **113**, 2084–2094. doi:10.1121/1.1555611

Elliott, T. M., and Theunissen, F. E. (**2009**). "The Modulation Transfer Function for Speech
Intelligibility," PLoS Comput Biol, **5**, e1000302. doi:10.1371/journal.pcbi.1000302

Ewert, S. D., and Dau, T. (**2000**). "Characterizing frequency selectivity for envelope
fluctuations," J Acoust Soc Am, **108**, 1181–1196. doi:10.1121/1.1288665

Ewert, S. D., Verhey, J. L., and Dau, T. (**2002**). "Spectro-temporal processing in the envelope-
frequency domain," J Acoust Soc Am, **112**, 2921–2931. doi:10.1121/1.1515735

Fitzgibbons, P. J., and Gordon-Salant, S. (**2010**). "Behavioral Studies With Aging Humans:
Hearing Sensitivity and Psychoacoustics," In S. Gordon-Salant, R. Frisina, A. Popper,
and R. Fay (Eds.), The Aging Auditory System. Springer Handbook of Auditory
Research, Springer, New York, Ny, Vol. 34, pp. 111–134. doi:10.1007/978-1-4419-0993-
0_5

Frisina, D. R., and Frisina, R. D. (**1997**). "Speech recognition in noise and presbycusis: relations
to possible neural mechanisms," Hear Res, **106**, 95–104. doi:10.1016/S0378-
5955(97)00006-3

Fuglsang, S. A., Märcher-Rørsted, J., Dau, T., and Hjortkjær, J. (**2020**). "Effects of Sensorineural

Hearing Loss on Cortical Synchronization to Competing Speech during Selective

Attention," J Neurosci, **40**, 2562–2572. doi:10.1523/JNEUROSCI.1936-19.2020

Füllgrabe, C., Moore, B. C. J., and Stone, M. A. (**2015**). "Age-group differences in speech

identification despite matched audiometrically normal hearing: contributions from

auditory temporal processing and cognition," Front Aging Neurosci, **6**, 1–25.

doi:10.3389/fnagi.2014.00347

Gifford, R. H., Bacon, S. P., and Williams, E. J. (**2007**). "An Examination of Speech Recognition

in a Modulated Background and of Forward Masking in Younger and Older Listeners,"

J Speech Lang Hear Res, **50**, 857–864. doi:10.1044/1092-4388(2007/060)

Glasberg, B. R., and Moore, B. C. J. (**1986**). "Auditory filter shapes in subjects with unilateral and

bilateral cochlear impairments," J Acoust Soc Am, **79**, 1020–1033. doi:10.1121/1.393374

Glasberg, B. R., and Moore, B. C. J. (**1989**). "Psychoacoustic abilities of subjects with unilateral

and bilateral cochlear hearing impairments and their relationship to the ability to

understand speech," Scand Audiol Suppl, **32**, 1–25.

Goossens, T., Vercammen, C., Wouters, J., and Wieringen, A. van (**2016**). "Aging Affects Neural

Synchronization to Speech-Related Acoustic Modulations," Front Aging Neurosci, **8**, 1–

16. doi:10.3389/fnagi.2016.00133

Goossens, T., Vercammen, C., Wouters, J., and van Wieringen, A. (**2017**). "Masked speech

perception across the adult lifespan: Impact of age and hearing impairment," Hear Res,

344, 109–124. doi:10.1016/j.heares.2016.11.004

Goossens, T., Vercammen, C., Wouters, J., and van Wieringen, A. (**2018**). "Neural envelope

encoding predicts speech perception performance for normal-hearing and hearing-

impaired adults," Hear Res, **370**, 189–200. doi:10.1016/j.heares.2018.07.012

Gordon-Salant, S., and Cole, S. S. (**2016**). "Effects of Age and Working Memory Capacity on

Speech Recognition Performance in Noise Among Listeners With Normal Hearing," Ear

Hear, **37**, 593–602. doi:10.1097/AUD.0000000000000316

Gordon-Salant, S., Yeni-Komshian, G. H., Fitzgibbons, P. J., and Barrett, J. (**2006**). "Age-related

differences in identification and discrimination of temporal cues in speech segments," J

Acoust Soc Am, **119**, 2455–2466. doi:10.1121/1.2171527

Greenberg, S., Carvey, H., Hitchcock, L., and Chang, S. (**2003**). "Temporal properties of

spontaneous speech—a syllable-centric perspective," J Phon, **31**, 465–485.

doi:10.1016/j.wocn.2003.09.005

Grose, J. H., Mamo, S. K., and Hall, J. W. (**2009**). "Age Effects in Temporal Envelope Processing:

Speech Unmasking and Auditory Steady State Responses," Ear Hear, **30**, 568–575.

doi:10.1097/AUD.0b013e3181ac128f

Helfer, K. S., and Vargo, M. (**2009**). "Speech recognition and temporal processing in middle-

aged women," J Am Acad Audiol, **20**, 264–271. doi:10.3766/jaaa.20.4.6

Houtgast, T. (**1989**). "Frequency selectivity in amplitude-modulation detection," J Acoust Soc

Am, **85**, 1676–1680. doi:10.1121/1.397956

Houtgast, T., and Steeneken, H. J. M. (**1985**). "A review of the MTF concept in room acoustics

and its use for estimating speech intelligibility in auditoria," J Acoust Soc Am, **77**, 1069–

1077. doi:10.1121/1.392224

Humes, L. E. (**2002**). "Factors underlying the speech-recognition performance of elderly hearing-

aid wearers," J Acoust Soc Am, **112**, 1112–1132. doi:10.1121/1.1499132

Humes, L. E. (**2007**). "The Contributions of Audibility and Cognitive Factors to the Benefit

Provided by Amplified Speech to Older Adults," J Am Acad Audiol, **18**, 590–603.

doi:10.3766/jaaa.18.7.6

Humes, L. E., and Christopherson, L. (**1991**). "Speech Identification Difficulties of Hearing-

Impaired Elderly Persons," J Speech Lang Hear Res, **34**, 686–693.

doi:10.1044/jshr.3403.686

Humes, L. E., and Dubno, J. R. (**2010**). "Factors Affecting Speech Understanding in Older

Adults," In S. Gordon-Salant, R. Frisina, and A. F. R. Popper (Eds.), The Aging Auditory

System. Springer Handbook of Auditory Research, Springer, New York, Vol. 34, pp.

211–257. doi:10.1007/978-1-4419-0993-0_8

Humes, L. E., Dubno, J. R., Gordon-Salant, S., Lister, J. J., Cacace, A. T., Cruickshanks, K. J.,

Gates, G. A., et al. (**2012**). "Central Presbycusis: A Review and Evaluation of the

Evidence," J Am Acad Audiol, **23**, 635–666. doi:10.3766/jaaa.23.8.5

Humes, L. E., Lee, J. H., and Coughlin, M. P. (**2006**). "Auditory measures of selective and divided

attention in young and older adults using single-talker competition.," J Acoust Soc Am,

120, 2926–37. doi:10.1121/1.2354070

Hunter, L. L., Monson, B. B., Moore, D. R., Dhar, S., Wright, B. A., Munro, K. J., Zadeh, L. M.,

et al. (**2020**). "Extended high frequency hearing and speech perception implications in

adults and children," Hear Res, **397**, 107922. doi:10.1016/j.heares.2020.107922

Jennings, S. G., Chen, J., Fultz, S. E., Ahlstrom, J. B., and Dubno, J. R. (**2018**). "Amplitude

modulation detection with a short-duration carrier: Effects of a precursor and hearing

loss," J Acoust Soc Am, **143**, 2232–2243. doi:10.1121/1.5031122

Jepsen, M. L., and Dau, T. (**2011**). "Characterizing auditory processing and perception in

individual listeners with sensorineural hearing loss," J Acoust Soc Am, **129**, 262–281.

doi:10.1121/1.3518768

Jepsen, M. L., Ewert, S. D., and Dau, T. (**2008**). "A computational model of human auditory

signal processing and perception," J Acoust Soc Am, **124**, 422–438.

Jeub, M., Schafer, M., and Vary, P. (**2009**). "A binaural room impulse response database for the

753   evaluation of dereverberation algorithms," 2009 16th International Conference on

754   Digital Signal Processing, IEEE, 1–5. doi:10.1109/ICDSP.2009.5201259

755   Jørgensen, S., and Dau, T. (**2011**). "Predicting speech intelligibility based on the signal-to-noise

756   envelope power ratio after modulation-frequency selective processing," J Acoust Soc

757   Am, **130**, 1475–1487. doi:10.1121/1.3621502

758   Jørgensen, S., Ewert, S. D., and Dau, T. (**2013**). "A multi-resolution envelope-power based model

759   for speech intelligibility," J Acoust Soc Am, **134**, 436–446. doi:10.1121/1.4807563

760   Joris, P. X., Schreiner, C. E., and Rees, A. (**2004**). "Neural Processing of Amplitude-Modulated

761   Sounds," Physiol Rev, **84**, 541–577. doi:10.1152/physrev.00029.2003

762   Kidd, G., Mason, C. R., Richards, V. M., Gallun, F. J., and Durlach, N. I. (**2008**). "Informational

763   Masking," In W. A. Yost, A. N. Popper, and R. R. Fay (Eds.), Auditory Perception of

764   Sound Sources. Springer Handbook of Auditory Research, Springer, Boston, MA, Vol.

765   29, pp. 143–189. doi:10.1007/978-0-387-71305-2_6

766   Kidd, G., Mason, C. R., Swaminathan, J., Roverud, E., Clayton, K. K., and Best, V. (**2016**).

767   "Determining the energetic and informational components of speech-on-speech

768   masking," J Acoust Soc Am, **140**, 132–144. doi:10.1121/1.4954748

769   Kohlrausch, A., Fassel, R., and Dau, T. (**2000**). "The influence of carrier level and frequency on

770   modulation and beat-detection thresholds for sinusoidal carriers," J Acoust Soc Am, **108**,

771   723–734. doi:10.1121/1.429605

772   Kraha, A., Turner, H., Nimon, K., Zientek, L. R., and Henson, R. K. (**2012**). "Tools to support

773   interpreting multiple regression in the face of multicollinearity," Front Psychol, **3**, 1–16.

774   doi:10.3389/fpsyg.2012.00044

Léger, A. C., Moore, B. C. J., and Lorenzi, C. (**2012**). "Abnormal speech processing in frequency regions where absolute thresholds are normal for listeners with high-frequency hearing loss," Hear Res, **294**, 95–103. doi:10.1016/j.heares.2012.10.002

Lorenzi, C., Gilbert, G., Carn, H., Garnier, S., and Moore, B. C. J. (**2006**). "Speech perception problems of the hearing impaired reflect inability to use temporal fine structure," Proc Natl Acad Sci, **103**, 18866–18869. doi:10.1073/pnas.0607364103

McDermott, J. H., and Simoncelli, E. P. (**2011**). "Sound texture perception via statistics of the auditory periphery: Evidence from sound synthesis," Neuron, **71**, 926–940. doi:10.1016/j.neuron.2011.06.032

McWalter, R., and Dau, T. (**2017**). "Cascaded Amplitude Modulations in Sound Texture Perception," Front Neurosci, **11**, 1–12. doi:10.3389/fnins.2017.00485

Millman, R. E., Mattys, S. L., Gouws, A. D., and Prendergast, G. (**2017**). "Magnified neural envelope coding predicts deficits in speech perception in noise," J Neurosci, **37**, 7727–7736. doi:10.1523/JNEUROSCI.2722-16.2017

Moore, B. C. J. (**2003**). "Speech processing for the hearing-impaired: successes, failures, and implications for speech mechanisms," Speech Commun, **41**, 81–91. doi:10.1016/S0167-6393(02)00095-X

Moore, B. C. J., and Glasberg, B. R. (**1998**). "Use of a loudness model for hearing-aid fitting. I. Linear hearing aids," Br J Audiol, **32**, 317–335. doi:10.3109/03005364000000083

Moore, B. C. J., and Oxenham, A. J. (**1998**). "Psychoacoustic consequences of compression in the peripheral auditory system.," Psychol Rev, **105**, 108–124. doi:10.1037/0033-295X.105.1.108

Moore, B. C. J., Wojtczak, M., and Vickers, D. A. (**1996**). "Effect of loudness recruitment on the

perception of amplitude modulation," J Acoust Soc Am, **100**, 481–489.

doi:10.1121/1.415861

Nielsen, J. B., and Dau, T. (**2011**). "The Danish hearing in noise test," Int J Audiol, **50**, 202–208.

doi:10.3109/14992027.2010.524254

Oxenham, A. J., and Moore, B. C. J. (**1997**). "Modeling the Effects of Peripheral Nonlinearity in

Listeners With Normal and Impaired Hearing," in Jesteadt W. (Ed), Modeling

Sensorineural Hearing Loss, Routledge, New York, pp. 273–288.

doi:10.4324/9781315789392-22

Parthasarathy, A., Bartlett, E. L., and Kujawa, S. G. (**2019a**). "Age-related Changes in Neural

Coding of Envelope Cues: Peripheral Declines and Central Compensation," Neurosci,

407, 21–31. doi:10.1016/j.neuroscience.2018.12.007

Parthasarathy, A., Herrmann, B., and Bartlett, E. L. (**2019b**). "Aging alters envelope

representations of speech-like sounds in the inferior colliculus," Neurobiol Aging, **73**,

30–40. doi:10.1016/j.neurobiolaging.2018.08.023

Pichora-Fuller, M. K., and MacDonald, E. (**2008**). "Auditory temporal processing deficits in older

listeners: From a review to a future view of presbycusis," Proceedings of the International

Symposium on Auditory and Audiological Research, Nyborg, Denmark, 291–300.

Pichora-Fuller, M. K., and Souza, P. E. (**2003**). "Effects of aging on auditory processing of

speech," Int J Audiol, **42**, 11–16. doi:10.3109/14992020309074638

Polspoel, S., Kramer, S. E., van Dijk, B., and Smits, C. (**2022**). "The Importance of Extended

High-Frequency Speech Information in the Recognition of Digits, Words, and Sentences

in Quiet and Noise," Ear Hear, **43**, 913–920. doi:10.1097/AUD.0000000000001142

Rajan, R., and Cainer, K. E. (**2008**). "Ageing without hearing loss or cognitive impairment causes

a decrease in speech intelligibility only in informational maskers," Neurosci, **154**, 784–

795. doi:10.1016/j.neuroscience.2008.03.067

Ray-Mukherjee, J., Nimon, K., Mukherjee, S., Morris, D. W., Slotow, R., and Hamer, M. (**2014**).

"Using commonality analysis in multiple regressions: A tool to decompose regression

effects in the face of multicollinearity," Methods Ecol Evol, **5**, 320–328.

doi:10.1111/2041-210X.12166

Regev, J., Relaño-Iborra, H., Zaar, J., and Dau, T. (**2024a**). "Disentangling the effects of hearing

loss and age on amplitude modulation frequency selectivity," J Acoust Soc Am, **155**,

2589–2602. doi:10.1121/10.0025541

Regev, J., Relaño-Iborra, H., Zaar, J., and Dau, T. (**2024b**). "Dataset for: 'Disentangling the

effects of hearing loss and age on amplitude modulation frequency selectivity,'" Technical

University of Denmark. doi:10.11583/DTU.25134611

Regev, J., Zaar, J., Relaño-Iborra, H., and Dau, T. (**2023**). "Age-related reduction of amplitude

modulation frequency selectivity," J Acoust Soc Am, **153**, 2298. doi:10.1121/10.0017835

Regev, J., Zaar, J., Relaño-Iborra, H., and Dau, T. (**2024c**). "Dataset for: 'Age-related reduction

of amplitude modulation frequency selectivity,'" Technical University of Denmark.

doi:10.11583/DTU.25134527

Relaño-Iborra, H., and Dau, T. (**2022**). "Speech intelligibility prediction based on modulation

frequency-selective processing," Hear Res, **426**, 108610.

doi:10.1016/j.heares.2022.108610

Relaño-Iborra, H., May, T., Zaar, J., Scheidiger, C., and Dau, T. (**2016**). "Predicting speech

intelligibility based on a correlation metric in the envelope power spectrum domain," J

Acoust Soc Am, **140**, 2670–2679. doi:10.1121/1.4964505

Relaño-Iborra, H., Zaar, J., and Dau, T. (**2019**). "A speech-based computational auditory signal
processing and perception model," J Acoust Soc Am, **146**, 3306–3317.
doi:10.1121/1.5129114

Roque, L., Karawani, H., Gordon-Salant, S., and Anderson, S. (**2019**). "Effects of age, cognition,
and neural encoding on the perception of temporal speech cues," Front Neurosci, **13**, 1–
15. doi:10.3389/fnins.2019.00749

Schlittenlacher, J., and Moore, B. C. J. (**2016**). "Discrimination of amplitude-modulation depth
by subjects with normal and impaired hearing," J Acoust Soc Am, **140**, 3487–3495.
doi:10.1121/1.4966117

Schneider, B. A., Li, L., and Daneman, M. (**2007**). "How Competing Speech Interferes with
Speech Comprehension in Everyday Listening Situations," J Am Acad Audiol, **18**, 559–
572. doi:10.3766/jaaa.18.7.4

Schneider, B. A., Pichora-Fuller, K., and Daneman, M. (**2010**). "Effects of Senescent Changes in
Audition and Cognition on Spoken Language Comprehension," In S. Gordon-Salant, R.
Frisina, A. Popper, and R. Fay (Eds.), The Aging Auditory System. Springer Handbook
of Auditory Research, Springer, New York, NY, Vol. 34, pp. 167–210. doi:10.1007/978-
1-4419-0993-0_7

Schoof, T., and Rosen, S. (**2014**). "The role of auditory and cognitive factors in understanding
speech in noise by normal-hearing older listeners," Front Aging Neurosci, **6**, 1–14.
doi:10.3389/fnagi.2014.00307

Shannon, R. V., Zeng, F.-G., Kamath, V., Wygonski, J., and Ekelid, M. (**1995**). "Speech
Recognition with Primarily Temporal Cues," Science (1979), **270**, 303–304.
doi:10.1126/science.270.5234.303

Singh, N. C., and Theunissen, F. E. (**2003**). "Modulation spectra of natural sounds and
ethological theories of auditory processing," J Acoust Soc Am, **114**, 3394–3411.
doi:10.1121/1.1624067

Smith, Z. M., Delgutte, B., and Oxenham, A. J. (**2002**). "Chimaeric sounds reveal dichotomies in
auditory perception," Nature, **416**, 87–90. doi:10.1038/416087a

Sobon, K. A., Taleb, N. M., Buss, E., Grose, J. H., and Calandruccio, L. (**2019**). "Psychometric
function slope for speech-in-noise and speech-in-speech: Effects of development and
aging," J Acoust Soc Am, **145**, EL284–EL290. doi:10.1121/1.5097377

Soli, S. D., and Wong, L. L. N. (**2008**). "Assessment of speech intelligibility in noise with the
hearing in noise test," Int J Audiol, **47**, 356–361. doi:10.1080/14992020801895136

Steinmetzger, K., Zaar, J., Relaño-Iborra, H., Rosen, S., and Dau, T. (**2019**). "Predicting the
effects of periodicity on the intelligibility of masked speech: An evaluation of different
modelling approaches and their limitations," J Acoust Soc Am, **146**, 2562–2576.
doi:10.1121/1.5129050

Stone, M. A., Füllgrabe, C., and Moore, B. C. J. (**2012**). "Notionally steady background noise acts
primarily as a modulation masker of speech," J Acoust Soc Am, **132**, 317–326.
doi:10.1121/1.4725766

Stone, M. A., and Moore, B. C. J. (**2014**). "On the near non-existence of 'pure' energetic masking
release for speech," J Acoust Soc Am, **135**, 1967–1977. doi:10.1121/1.4868392

Strelcyk, O., and Dau, T. (**2009**). "Relations between frequency selectivity, temporal fine-
structure processing, and speech reception in impaired hearing," J Acoust Soc Am, **125**,
3328. doi:10.1121/1.3097469

The Mathworks Inc. (**2015**). "MATLAB 2015b - MathWorks,"
www.mathworks.com/products/matlab. Last accessed January 1st, 2016.

Thompson, B., and Borrello, G. M. (**1985**). "The Importance of Structure Coefficients in

Regression Research," Educ Psychol Meas, **45**, 203–209.

doi:10.1177/001316448504500202

Varnet, L., Ortiz-Barajas, M. C., Erra, R. G., Gervain, J., and Lorenzi, C. (**2017**). "A cross-

linguistic study of speech modulation spectra," J Acoust Soc Am, **142**, 1976–1989.

doi:10.1121/1.5006179

Verhey, J. L., Ewert, S. D., and Dau, T. (**2003**). "Modulation masking produced by complex tone

modulators," J Acoust Soc Am, **114**, 2135–2146. doi:10.1121/1.1612489

Viemeister, N. F. (**1979**). "Temporal modulation transfer functions based upon modulation

thresholds," J Acoust Soc Am, **66**, 1364–1380. doi:10.1121/1.383531

Viswanathan, V., Bharadwaj, H. M., Shinn-Cunningham, B. G., and Heinz, M. G. (**2021**).

"Modulation masking and fine structure shape neural envelope coding to predict speech

intelligibility across diverse listening conditions," J Acoust Soc Am, **150**, 2230–2244.

doi:10.1121/10.0006385

Xu, L., Thompson, C. S., and Pfingst, B. E. (**2005**). "Relative contributions of spectral and

temporal cues for phoneme recognition," J Acoust Soc Am, **117**, 3255–3267.

doi:10.1121/1.1886405

Zaar, J., Simonsen, L. B., Dau, T., and Laugesen, S. (**2023**). "Toward a clinically viable spectro-

temporal modulation test for predicting supra-threshold speech reception in hearing-

impaired listeners," Hear Res, **427**, 108650. doi:10.1016/j.heares.2022.108650

Zaar, J., Simonsen, L. B., and Laugesen, S. (**2024**). "A spectro-temporal modulation test for

predicting speech reception in hearing-impaired listeners with hearing aids," Hear Res,

443, 108949. doi:10.1016/j.heares.2024.108949