Investigating the effects of age and hearing loss on speech intelligibility and amplitude modulation frequency selectivity

Jonathan Regev,[1,a] Johannes Zaar,[1,2] Helia Relaño-Iborra,[1] and Torsten Dau[1]

[1] Hearing Systems Section, Department of Health Technology, Technical University of Denmark, Kongens Lyngby,

2800, Denmark

[2] Eriksholm Research Centre, Snekkersten, 3070, Denmark

Abstract

The perception of amplitude modulation (AM), characterized by a frequency-selective process in the modulation domain, is considered critical for speech intelligibility. Previous literature has provided evidence of an age-related decline in AM frequency selectivity. Additionally, a notable sharpening of AM tuning with hearing loss has been observed, which was proposed to be related to a perceptual advantage resulting from peripheral compression loss. This study explores whether such changes in AM tuning with age and hearing loss contribute to the speech intelligibility challenges that older listeners often face in noisy environments. Young (n=10, 22-28 years) and older (n=9, 57-77 years) listeners with normal hearing as well as older listeners with hearing impairment (n=9, 64-77 years) participated in the study. All had previously taken part in studies on AM tuning. Speech-reception thresholds (SRTs) were collected in conditions including stationary, fluctuating, and competing-speech maskers. The results revealed an age-related increase in SRTs, with an additional negative impact of

[a] joreg@dtu.dk

hearing loss. Beyond age and audibility, a measure of AM tuning significantly contributed to explaining the variance in SRTs across listeners for specific maskers. These findings motivate further exploration of the relationship between AM frequency selectivity and speech intelligibility in noise.

I. INTRODUCTION

Temporal envelope fluctuations (also known as "amplitude modulation" or AM) refer to the slow variations in overall amplitude that characterize most sounds in daily life (e.g., Singh and Theunissen, 2003; Elliott and Theunissen, 2009). The auditory system exhibits remarkable sensitivity to AM, as demonstrated by physiological studies (see Joris *et al.*, 2004 for a review) and behavioral investigations of AM detection (e.g., Viemeister, 1979; Kohlrausch *et al.*, 2000). Similar to masking effects in the audio-frequency domain, the detectability of a target AM is disrupted by the presence of distinct masking AM components. AM masking has been extensively studied in behavioral investigations (e.g., Bacon and Grantham, 1989; Houtgast, 1989; Dau *et al.*, 1997; Verhey *et al.*, 2003) as well as through recent neurophysiological research (Viswanathan *et al.*, 2021). Behavioral studies of AM masking have provided evidence of a frequency-selective process in the modulation domain, often conceptualized as a series of bandpass modulation filters known as the modulation filterbank (e.g., Dau *et al.*, 1997, 1999; Ewert and Dau, 2000; Ewert *et al.*, 2002). Computational modelling studies have shown that this concept significantly contributes to quantitatively predicting experimental data from various psychoacoustic tasks (e.g., Dau *et al.*, 1997, 1999; Jepsen *et al.*, 2008; Jepsen and Dau, 2011; McDermott and Simoncelli, 2011; McWalter and Dau, 2017), underscoring the importance of AM in auditory processing and perception.

Beyond its role in 'simple' listening tasks, AM has been argued to be crucial for accurate speech intelligibility (Houtgast and Steeneken, 1985; Drullman *et al.*, 1994; Smith *et al.*, 2002; Elliott and Theunissen, 2009). The significance of AM in speech was highlighted by vocoding studies, which demonstrated that intact envelope information conveyed through a few broad spectral bands was sufficient for precise speech recognition in quiet, even when spectral cues were greatly reduced (e.g., Shannon *et al.*, 1995; Dorman *et al.*, 1997; Xu *et al.*, 2005). Based on this, AM masking has been proposed as the primary mechanism underlying speech intelligibility in noise, whereby speech-

reception thresholds (SRTs) - reflecting the signal-to-noise ratio required for 50% speech recognition

- are related to the modulation-domain signal-to-noise ratio (SNR_{env}) between the speech and masker

(e.g., Dubbelboer and Houtgast, 2008; Jørgensen and Dau, 2011).

The crucial role of AM masking in speech intelligibility in noise was also emphasized by studies

using maskers designed to contain AM components or to have maximally flat envelopes, which

suggested that so-called stationary maskers, usually associated with energetic masking (EM) in the

audio-frequency domain, effectively function as AM maskers due to their inherent envelope

fluctuations (Stone *et al.*, 2012; Stone and Moore, 2014). Computational auditory modeling studies

have successfully incorporated these concepts by employing the SNR_{env} (Jørgensen and Dau, 2011;

Jørgensen *et al.*, 2013; Chabot-Leclerc *et al.*, 2014) or correlation-based measures (Relaño-Iborra *et al.*,

2016, 2019; Steinmetzger *et al.*, 2019) at the output of the modulation filterbank as decision metrics.

Based on these modeling studies, the spectral decomposition of the stimulus envelope carried out by

the modulation filterbank, i.e., AM frequency selectivity, has been suggested as an important factor

for accurately predicting SRTs (see Relaño-Iborra and Dau, 2022 for a review of different speech

intelligibility models using modulation filtering).

Regev *et al.* (2023, 2024a) investigated the effects of age and hearing loss on the selectivity of

hypothetical modulation filters using a perceptual AM masking task. Regev *et al.* (2023) found a general

reduction in AM frequency selectivity associated with age, resulting in broader masking patterns

among older compared to young listeners with normal hearing (referred to as hereafter as NH

listeners). Conversely, Regev *et al.* (2024a) demonstrated a general sharpening of masking patterns

among young and older listeners with hearing impairment (referred to hereafter as HI listeners)

compared to NH listeners of similar age. This finding suggested that sensorineural hearing loss was

accompanied by a perceptual benefit that partially offset the detrimental effects of age, possibly due

to the HI listeners' ability to exploit an increased internal representation of AM depth resulting from

reduced peripheral compression when "listening in the dips" of the masker. Hence, Regev *et al.* (2024a)

proposed that the sharpening of the masking patterns found in HI listeners was an "apparent" effect,

reflecting the loss of peripheral compression rather than an actual greater selectivity of the hypothetical

modulation filters. In other words, Regev *et al.* (2024a) proposed that older HI listeners also experience

an age-related reduction in AM frequency selectivity, which is concealed by a perceptual benefit

stemming from the loss of peripheral compression. Such age-related changes in the sharpness of the

hypothetical modulation filters may affect speech intelligibility in noise by altering the amount of AM

masking at the output of the modulation filters.

The effects of age and hearing loss on speech intelligibility have received significant attention in

the literature. Hearing loss is known to cause a decline in speech intelligibility in both quiet and noisy

environments (e.g., Glasberg and Moore, 1989; Humes and Christopherson, 1991; Frisina and Frisina,

1997; see Humes and Dubno, 2010 for a review). In addition to audibility loss, speech intelligibility

challenges associated with hearing loss have been linked to various supra-threshold deficits, including

reduced frequency selectivity, poorer temporal fine structure processing, reduced recovery from

forward masking, poorer AM depth discrimination, and loudness recruitment (Glasberg and Moore,

1986, 1989; Oxenham and Moore, 1997; Moore and Oxenham, 1998; Moore, 2003; Lorenzi *et al.*,

2006; Strelcyk and Dau, 2009; Schlittenlacher and Moore, 2016; Gallun *et al.*, 2022). Aided speech

intelligibility performance has also been shown to be strongly correlated with deficits in spectro-

temporal modulation detection in HI listeners (e.g., Bernstein *et al.*, 2016; Zaar *et al.*, 2023, 2024).

Recent neurophysiological studies suggested that poorer speech intelligibility in HI listeners may be

related to increased neural coding of the envelope (Millman *et al.*, 2017; Goossens *et al.*, 2018; Decruy

et al., 2020). This effect may be partly attributed to an increased internal representation of the AM

depth resulting from the loss of cochlear compression commonly associated with sensorineural

hearing loss (Moore *et al.*, 1996; Jennings *et al.*, 2018). In addition to the effects of hearing loss itself,

age and reduced cognitive function have been found to contribute to the decline in speech intelligibility

observed in older HI listeners (e.g., Dubno *et al.*, 1984; Humes and Christopherson, 1991; Frisina and

Frisina, 1997; Humes, 2002, 2007; Humes *et al.*, 2006). This suggests that cognitive and age-related

perception deficits may also play a role in speech intelligibility challenges.

To investigate these potential mechanisms, cross-sectional studies focusing on age-related effects

have been conducted, often involving listeners with no or mild hearing loss. These studies have

consistently shown a detrimental effect of age on speech intelligibility in the presence of competing

speech maskers (Rajan and Cainer, 2008; Helfer and Vargo, 2009; Schoof and Rosen, 2014; Füllgrabe

et al., 2015; Gordon-Salant and Cole, 2016; Goossens *et al.*, 2017; Decruy *et al.*, 2019; Sobon *et al.*,

2019). However, the effects of age in stationary and fluctuating noise have been less consistent. Some

studies found differences between young and older listeners (e.g., Goossens *et al.*, 2017; Decruy *et al.*,

2019; Sobon *et al.*, 2019), while others did not (Rajan and Cainer, 2008; Helfer and Vargo, 2009; Schoof

and Rosen, 2014). Overall, the age-related increase in SRTs has been more pronounced for speech

maskers than for stationary or fluctuating noises (e.g., Goossens *et al.*, 2017; Decruy *et al.*, 2019; Sobon

et al., 2019). The observed age-related decline in speech intelligibility has been suggested to be mediated

by deficits in temporal processing (e.g., Gordon-Salant *et al.*, 2006; Anderson *et al.*, 2012; Füllgrabe *et

al.*, 2015; Goossens *et al.*, 2016; Decruy *et al.*, 2019; McFarlane and Sanchez, 2024) and by reduced

cognitive abilities (Humes *et al.*, 2012; Füllgrabe *et al.*, 2015; Gordon-Salant and Cole, 2016).

Since AM frequency selectivity has been identified as a significant factor contributing to accurate

speech intelligibility, it is reasonable to suggest that the age-related decline in NH listeners' AM

selectivity reported by Regev *et al.* (2023) may contribute to the aforementioned challenges in speech

intelligibility. A reduction in AM tuning would likely result in increased AM masking, thus disrupting

the intelligibility of the target speech. Similarly, it is hypothesized that older HI listeners also experience

an age-related reduction in AM selectivity that contributes to a decline in speech intelligibility.

Although the increased AM frequency selectivity observed in older HI listeners compared to older

NH listeners (Regev *et al.*, 2024a) might suggest reduced AM masking with hearing loss, it was

proposed that this greater AM selectivity in HI listeners is an apparent effect caused by reduced

cochlear compression, rather than a true sharpening of the hypothetical modulation filter.

Consequently, the contribution of an age-related reduction in AM selectivity to speech intelligibility

difficulties in older HI listeners should become evident once hearing loss is accounted for.

Furthermore, reduced cochlear compression is associated with supra-threshold deficits that impair

speech intelligibility (e.g., Glasberg and Moore, 1989; Oxenham and Moore, 1997; Moore and

Oxenham, 1998; Moore, 2003). This aligns with studies showing that HI listeners exhibit higher SRTs

than age-matched NH listeners even when audibility is restored (Goossens *et al.*, 2017).

The present study investigated whether the results of speech intelligibility tests support the

assumption that an age-related decline in AM frequency selectivity negatively impacts speech

intelligibility in both older NH and older HI listeners, once the effects of cochlear compression loss

in HI listeners are accounted for. SRTs were collected for speech in noise, building upon the previous

studies on AM frequency selectivity (Regev *et al.*, 2023, 2024a), using a subset of the participants.

Young NH, older NH, and older HI listeners were tested on sentence recognition in various maskers,

including speech-shaped noise, fluctuating noise, and competing talkers. It was hypothesized that age-

related changes in AM frequency selectivity would be linked to increases in SRTs for the older NH

and HI listeners.

II.   METHODS

A. Listeners and previously published data

All young NH and older NH listeners tested by Regev *et al.* (2023) as well as all older HI listeners

tested by Regev *et al.* (2024a) were invited to participate in the present study. Ten young NH listeners,

aged 22-28 years (mean age = 24.7 years, eight female), nine older NH listeners, aged 57–77 years

(mean age = 66.9 years, six female), and nine older HI listeners, aged 64-77 years (mean age = 71.2

years, 2 female), accepted this invitation. All listeners were native Danish speakers. The experiment

was conducted monaurally in the same test ear as used by Regev *et al.* (2023, 2024a). Figure 1 shows

the audiograms in the participants' test ears. All young NH listeners had audiometric pure-tone

thresholds (PTTs) ≤ 20 dB HL at frequencies ranging from 125 Hz to 8 kHz. All older NH listeners

had PTTs ≤ 20 dB HL at frequencies ranging from 125 Hz to 4 kHz, PTTs ≤ 25 dB HL at 6 kHz,

and PTTs ≤ 45 dB HL at 8 kHz. The older HI listeners exhibited a high-frequency sloping hearing

loss, classified as an N2 or N3 standard audiogram (Bisgaard *et al.*, 2010),[1] and audiometric thresholds

in the range of 50-60 dB HL at 3 kHz as required by Regev *et al.* (2024a). All HI listeners had

sensorineural hearing loss, with air-bone gaps ≤ 15 dB at 1, 2, 3, and 4 kHz.[2]

Prior to this study, all listeners took part in the AM frequency selectivity studies conducted by Regev

et al. (2023, 2024a). The corresponding datasets are publicly available from Regev *et al.* (2024c, 2024b).

In those studies, masked-threshold patterns (MTPs) were used to assess the masked detection

threshold of a target AM across different masker center frequencies, providing an indication of AM

frequency selectivity. Figure 2 indicates an example of the MTPs for a 4-Hz target modulation

frequency, where the age-related reduction in AM selectivity is evident on both skirts and the

sharpening of the MTP with hearing loss is evident on the low-frequency skirt (see Regev *et al.*, 2023,

and 2024a for a detailed discussion of the possible origins of these effects). A measure of AM

frequency selectivity at the individual level can be derived from the MTPs by computing the average

range spanned by the MTP (i.e., the MTP's average dynamic range; DR_{MTP}), where a larger dynamic

range is indicative of greater AM selectivity. This is done by identifying the highest masked threshold

as the peak of the MTP, computing the difference between the peak and the lowest masked threshold

on each side of the peak, and averaging the resulting dynamic range across both sides. If the peak of

the MTP is on either extreme, the DR_{MTP} is taken as the dynamic range across the whole MTP. An

illustration of this derivation is provided in supplementary Figure 1. This measure was derived for the

4-Hz MTP of each listener (with the exception of one older HI listener for whom a MTP at 4 Hz

could not be obtained by Regev *et al.*, 2024a), since the 4-Hz modulation frequency has been suggested

to be particularly important for speech perception (e.g., Drullman *et al.*, 1994; Arai *et al.*, 1999;

Greenberg *et al.*, 2003; Elliott and Theunissen, 2009; Varnet *et al.*, 2017) and showed the largest effect

sizes across listener groups in Regev *et al.* (2023, 2024a). The resulting individual quantifications of

AM frequency selectivity are shown in Figure 3 for each listener group. As reported by Regev *et al.*

(2023, 2024a), the MTPs and corresponding dynamic ranges indicate a general decline in AM

frequency selectivity with age, which is partially "compensated" by the presence of hearing loss.

Regev *et al.* (2023, 2024a) also evaluated working-memory capacity using a reverse digit span (RDS)

test, which was administered binaurally following the procedure described in Fuglsang *et al.* (2020).

For the HI listeners, a linear gain was applied based on the Cambridge formula (CamEq; Moore and

Glasberg, 1998), with a maximum gain of 30 dB at any frequency. The older NH listeners exhibited

significantly lower average RDS scores (0.40 on a normalized scale) compared to the young NH and

older HI listeners (0.60 and 0.52, respectively).[3]

All participants provided written informed consent and were offered financial compensation for

their participation. The study received ethical approval from the Science Ethics Committee of the

Capital Region of Denmark (reference H-16036391).

Figure 1 (Color online): Individual and average audiograms of the test ear for each listener group. The

thin dashed lines show the individual audiograms, while the dark solid lines represent the average and

standard deviation across listeners. Panel A shows the audiograms for the young NH listeners, with

the black horizontal line indicating the 20 dB HL normal-hearing criterion. Panel B displays the

audiograms for the older NH listeners, also with the black horizontal line denoting the 20 dB HL

normal-hearing criterion. Panel C presents the audiograms for the older HI listeners.

Figure 2 (Color online): Average MTPs and AM detection thresholds for a 4-Hz target modulation

frequency. The results for the young NH group are presented as dashed lines and squares, the older

NH group as dashed lines and triangles, and the older HI group as continuous lines and circles. The

bottom open symbols represent the average AM detection threshold for each group. The means and

standard errors are shown. Data adapted from Regev *et al.* (2023, 2024a).

Figure 3 (Color online): Boxplots displaying the average dynamic range of the 4-Hz MTP (DR_{MTP}) for

all listeners. The data for the young NH listeners are represented by squares, older NH by triangles,

and older HI by circles. Data derived from Regev *et al.* (2023, 2024a).

B. Apparatus

The listening test was conducted in a single-walled soundproof booth. The experiment was

conducted using MATLAB (The Mathworks Inc., 2015). The stimuli were digitally played using

SoundMexPro (HörTech, Oldenburg, Germany), converted to analog using an RME Fireface

soundcard (Haimhausen, Germany), and played monaurally through Sennheiser HDA200 headphones

(Wedemark, Germany). The output level was calibrated using a GRAS RA0039 ear simulator (Holte,

Denmark) and the frequency response of the headphones was compensated using an inverse digital

filter to achieve a flat response at the eardrum. The listeners received no feedback during the test.

C. Procedure and stimuli

Speech intelligibility in noise was assessed using the Danish Hearing in Noise Test (HINT; Nielsen

and Dau, 2011). The HINT speech material consists of 10 test lists and 3 training lists, each containing

20 five-word sentences. The sentences were spoken by a male speaker with an estimated average

fundamental frequency (F_0) of 123 Hz. The speech signals were combined with five different masking

noises: i) a stationary speech-shaped noise (SSN); ii) a fluctuating ICRA-5 noise (Dreschler *et al.*, 2001)

resembling a single male talker in terms of spectral and temporal characteristics; iii) a single male

competing talker reading a news piece ($F_0 = 118$ Hz); iv) a single female competing talker reading a

news piece ($F_0 = 173$ Hz); and v) a 'cocktail party' condition. The recordings of the competing talkers

had previously been edited to reduce long pauses to 65 ms. The cocktail-party noise was generated by

combining a relatively dry recording from a cafeteria with the recordings of the male and female

competing talkers mentioned earlier. The levels of the competing talkers were each 3 dB higher than

that of the cafeteria noise. Prior to mixing, both competing talker recordings and the cafeteria noise

were convolved with an impulse response from the Aachen Impulse Response database (office setting

with dummy head at 1 m distance from the loudspeaker; Jeub *et al.*, 2009) to introduce reverberation

to the original recordings ($T_{60} = 0.37$ s). The spectra of the SSN and ICRA-5 noises, as well as that

of the cafeteria noise used to create the cocktail-party noise, had been pre-processed to match the

long-term average spectrum of the HINT speech material as closely as possible. The recording for the

male and female competing talkers had not been spectrally shaped, as their spectra were deemed

sufficiently close to the long-term average spectrum of the HINT material. The long-term average

spectra and modulation spectra of the Danish HINT speech material and all maskers are presented in

Figure 4.

Each participant underwent training using all three training lists mixed with SSN. The rule defining

the starting SNR (SNR_{start}) for each masker and testing run was based on pilot data from two young

NH listeners, whose SRTs were measured for all maskers. The pilot data indicated expected SRT

offsets (SRT_{offset}) relative to the SSN condition as follows: 0 dB for the male competing talker and

cocktail-party conditions, -3 dB for the ICRA noise, and -9 dB for the female competing talker. Based

on these offsets, SNR_{start} for each masker was adjusted to be 6 dB below the expected SRTs,

calculated relative to the SRT measured in the final training run with SSN ($SRT_{final\ train}$) and rounded

to the nearest integer (see Equation 1). Therefore, SNR_{start} was 6 dB lower than $SRT_{final\ train}$ for

the male competing talker, SSN, and cocktail-party conditions; 9 dB lower for the ICRA noise; and

15 dB lower for the female competing talker condition.

$$SNR_{start} = SRT_{final\ train} + SRT_{offset} - 6 \qquad (1)$$

The SRT for each masking noise was determined by averaging two separate measurements using

different lists, with masker presentation and lists order randomized across listeners. Both

measurements for each masker were conducted consecutively.[4] The SRTs were estimated using an

adaptive 1-up 1-down procedure targeting 50%-correct performance with sentence-based scoring. A

response was correct if all five words of the sentence were accurately repeated. The initial step size in

SNR was 4 dB for the first four sentences, then reduced to 2 dB. A refinement rule was applied to the

5[th] sentence: the change in SNR was computed as the average of the preceding four SNRs and the

predicted next SNR, minus the immediately preceding SNR. The first sentence was repeated until the

listener repeated it correctly. If the sentence required more than four repetitions, the run was

terminated, and the starting SNR was increased by 2 dB before restarting. The SRT for a run was

calculated as the average SNR from sentences 5 to 20, plus the SNR required for a fictitious 21[st]

sentence, determined by scoring the 20[th] sentence (Soli and Wong, 2008). The scoring was performed

by a trained native Danish speaker.

Target sentences were presented at a fixed level of 60 dB SPL, with masking noise looped

continuously and adjusted in level to achieve the desired SNR. For the HI listeners, linear amplification

was applied to both speech and masking noise using the Cambridge formula (CamEq; Moore and

Glasberg, 1998), with a maximum gain of 30 dB at any frequency.[5]

Figure 4 (Color online): Long-term average spectra (top row) and modulation spectra (bottom row)

of the speech material and maskers. Each panel shows the spectra for the HINT speech material (black

line) and one of the maskers (light blue line). Panels A and F: SSN; Panels B and G: ICRA-5; Panels

C and H: Male competing talker; Panels D and I: Female competing talker; Panels E and J: Cocktail-

party condition.

D. Statistical analysis

1. Group-level analysis of SRTs

The degree of variability between the two SRT estimates obtained for each masker was assessed

by calculating the standard deviation across the absolute differences in SRTs, $\sigma_{\Delta|SRT|}$. Notably,

although small, the variability across all maskers was highest for the young NH listeners ($\sigma_{\Delta|SRT|} =$

2.43 dB) compared to the older NH listeners ($\sigma_{\Delta|SRT|} = 1.76$ dB) and older HI listeners ($\sigma_{\Delta|SRT|} = 1.49$

dB). Nevertheless, the degree of variability between SRT estimates was considered sufficiently small

for all statistical analyses to be conducted using the SRTs averaged across both repetitions for each

listener and masker. Analyses of variance (ANOVAs) of linear mixed-effects model fittings were

performed to examine the differences between the listener groups. Two sets of analysis were

conducted: First, the older NH group was compared to the young NH listeners to isolate the effects

of age from hearing loss, and second, the older HI group was compared to the older NH listeners to

isolate the effects of hearing loss from age-related effects. In both analyses, the listeners were treated

as a random effect, while group classification and masker were considered as fixed effects.[6] Post-hoc

analyses were carried out using estimated marginal means, and a Holm-Bonferroni correction criterion

was applied to account for multiple comparisons. The significance level for all analyses was set at 5%.

2. Regression analysis

To assess whether a change in AM frequency selectivity contributes to the variance in SRTs, a

multilinear regression model was built to relate age, audibility, working memory capacity, and AM

frequency to SRTs across listeners. The SRTs from the condition showing the largest effect size across

young and older NH listeners were chosen as the model's outcome variable. The measure of age was

taken as the listeners' chronological age at the time of testing. The measure of audibility was taken as

the 4-frequency pure-tone average threshold (PTA; where the audiometric thresholds at 0.5, 1, 2, and

4 kHz are averaged). The measure of working memory capacity was taken as the listeners' RDS score.

Finally, the measure of AM frequency selectivity was taken as the listeners' DR_{MTP} at a 4-Hz target

modulation frequency, shown in Figure 3. The model included all young and older NH listeners and

8 of the 9 older HI listeners, since a 4-Hz MTP could not be obtained by Regev *et al.* (2024a) for one

of them. The multilinear regression model was built in steps, starting from a model including only age,

by assessing whether adding the other predictors and their interactions significantly reduced the

model's residual sum of squares, using an F-test. The predictors were added in the above order (PTA,

RDS, and DR_{MTP}) and the addition of their interactions was only assessed if the main effect

significantly improved the model. The significance level was set at 5%.

III.   RESULTS

A. Speech intelligibility

The SRTs for all listeners and groups are shown in Figure 5 for each masker. The young NH

listeners are represented by squares, older NH listeners by triangles, and older HI listeners by circles.

The statistical analysis comparing young and older NH listeners, focusing on the effects of age,

revealed significant main effects of masker [$F_{4,68}$ = 274.2, p < 0.001] and listener group [$F_{1,17}$ = 17.56,

p < 0.001], as well as a significant interaction between both factors [$F_{4,68}$ = 9.28, p < 0.001]. In the

analysis comparing older NH and older HI listeners, examining the effects of hearing loss, significant

main effects of masker [$F_{4,64}$ = 141.6, p < 0.001] and listener group [$F_{1,16}$ = 12.86, p = 0.002], as well

as a significant interaction between both factors [$F_{4,64}$ = 5.32, p < 0.001], were observed. Post-hoc

analyses revealed that the older NH listeners had significantly higher SRTs than the young NH

listeners for the ICRA-5 condition [Effect size (*M*) = 1.39 dB, standard error (*SE*) = 0.68 dB], male

competing talker [*M* = 4.53 dB, *SE* = 0.68 dB], female competing talker [*M* = 2.70 dB, *SE* = 0.68

dB], and cocktail-party condition [*M* = 1.53 dB, *SE* = 0.68 dB]. Additionally, the older NH listeners

had significantly lower SRTs than the older HI listeners for the SSN condition [*M* = 2.36 dB, *SE* =

1.14 dB], ICRA-5 condition [M = 4.30 dB, SE = 1.14 dB], male competing talker [M = 3.60 dB, SE = 1.14 dB], and female competing talker [M = 5.52 dB, SE = 1.14 dB]. The increase in SRT for the older HI listeners in the cocktail-party condition approached, but did not reach, significance [M = 2.24 dB, SE = 1.14 dB, p = 0.06]. Furthermore, the post-hoc analyses revealed that the female competing talker resulted in significantly lower SRTs than all other maskers for all listener groups. The cocktail-party condition yielded the highest SRTs for the young NH listeners, while the male competing talker resulted in significantly higher SRTs than all other maskers for the older HI listeners. For the older NH listeners, both the male competing talker and cocktail-party conditions produced similar SRTs, which were significantly higher than those for all other maskers. Finally, the ICRA-5 noise resulted in significantly lower SRTs than the SSN for the young and older NH listeners, but not for the older HI listeners. Additional details and results from all post-hoc analyses can be found in Appendix A.

Figure 5 (Color online): Boxplots displaying the SRTs for all listeners. Each panel corresponds to a

specific masker: Panel A: Speech-shaped noise (SSN), Panel B: ICRA-5 fluctuating noise, Panel C:

Single male competing talker (same sex), Panel D: Single female competing talker (different sex), Panel

E: Cocktail-party noise. The SRTs for the young NH listeners are represented by squares, older NH

by triangles, and older HI by circles. The stars indicate the level of statistical significance of the

differences between the groups separated by the horizontal black lines (* $p < 0.05$, ** $p < 0.01$, *** p

< 0.001), while "n.s." denotes non-significant differences. Note that the young NH and older HI

groups were not compared in the statistical analysis.

The multilinear regression analysis included the SRTs obtained with the male competing talker as

the outcome variable, as that condition yielded the largest difference in SRTs between young and older

NH listeners, as well as highest SRTs for both the older NH and HI listeners. The results showed that

the best-fitting model included the main effects of age, PTA, and DR_{MTP}, while RDS did not contribute

to explaining the variance in SRTs. Adding DR_{MTP} to age and PTA [$F_{1,23} = 9.693$, $p = 0.005$]

significantly increased the amount of variance in SRT explained by the model (adjusted R^2) by 8.4

percentage points to a total of 77%. No interaction term significantly increased the model's R^2. Figure

6 shows the scatter plots of the fitted SRTs as a function of the measured SRTs for each one of the

models iteratively built (i.e., including age, age + PTA, or age + PTA + DR_{MTP} as predictors) in

separate panels. The adjusted R^2 for each model is indicated in each panel. The best-fitting model's

regression coefficient (i.e., β weights) was larger for age (0.773) than for PTA and DR_{MTP} (0.303 and

0.350, respectively). The structure coefficients, i.e., the correlation of each predictor with the outcome

variable estimated from all predictors in the model, showed that both age and PTA were positively

correlated with the fitted SRTs, while DR$_{MTP}$ was negatively correlated (0.896, 0.842, and -0.157,

respectively).

Since all predictors in the best-fitting model (age, PTA, and DR$_{MTP}$) were likely somewhat collinear,

a commonality analysis (Capraro and Capraro, 2001; Ray-Mukherjee *et al.*, 2014) was implemented to

assess the R^2 either attributable uniquely to each predictor or shared between them. The outcome

showed that age had the greatest unique R^2 (22 %), while PTA and DR$_{MTP}$ had smaller unique R^2 (4.3

and 8.3 %, respectively). Furthermore, the commonality analysis revealed that age and PTA were

highly collinear whereby they shared a large common R^2 (48.8 %), and that age and DR$_{MTP}$ were also

somewhat collinear but shared a negative R^2 (-7.9%). The detailed results of the regression and

commonality analyses can be found in Appendix B.

Figure 6: Scatter plot of the fitted SRTs as a function of the measured SRTs for each one of the

regression models assessed. The predictors included in the model as well as the adjusted R^2 are

indicated in each panel. The SRTs for the young NH listeners are represented by squares, older NH

by triangles, and older HI by circles. The diagonal line indicates identity. The models were iteratively

built to include only age (left panel), age and PTA (middle panel), and finally age, PTA, and dynamic

range of the 4-Hz MTP (DR_{MTP}; right panel). The last model provided the best fit of the SRTs.

IV. DISCUSSION

The aim of the present study was to investigate whether the findings from speech intelligibility

tests would be consistent with the AM frequency selectivity data collected in previous studies by Regev

et al. (2023, 2024a). These previous investigations demonstrated an age-related decline in AM

frequency selectivity and an effective sharpening of AM tuning with hearing loss (see Figure 2 and

Figure 3). However, the latter was proposed to be an "apparent" result of reduced cochlear

compression (Regev *et al.*, 2024a), rather than a true sharpening of the theoretical modulation filters.

Thus, older HI listeners are also hypothesized to experience an age-related decline in AM selectivity.

A subset of the listener cohort measured by Regev *et al.* (2023, 2024a) was considered here. It was

expected that the age-related reduction in AM frequency selectivity would contribute to poorer speech

intelligibility in older NH and HI listeners compared to young NH listeners.

The results revealed an overall decline in speech intelligibility with age, which was further

exacerbated by the presence of hearing loss. These findings are consistent with previous studies that

have shown a deterioration of speech intelligibility with age (e.g., Rajan and Cainer, 2008; Helfer and

Vargo, 2009; Schoof and Rosen, 2014; Füllgrabe *et al.*, 2015; Gordon-Salant and Cole, 2016; Goossens

et al., 2017; Decruy *et al.*, 2019) and hearing loss (e.g., Glasberg and Moore, 1989; Humes and

Christopherson, 1991; Frisina and Frisina, 1997; see Humes and Dubno, 2010 for a review).

Additionally, a multilinear regression model relating the SRTs obtained with a male competing talker

to age, PTA, and DR_{MTP} (a measure of AM frequency selectivity) showed that DR_{MTP} significantly

contributed to explaining the variance in SRTs across all listeners, increasing the model's R^2 by 8

percentage points.

The observed increase in SRTs with age aligns with the age-related reduction in AM frequency

selectivity reported by Regev *et al.* (2023). Broader modulation filters, indicative of reduced AM

selectivity, may lead to increased AM masking, resulting in a decrease in the SNR_{env} at the output of

these filters and potentially poorer speech intelligibility in noise. However, the lack of a significant

age-related increase in SRT for the SSN condition is inconsistent with this notion. Since the SSN

contains intrinsic envelope fluctuations that act as a relatively broadband modulation masker (as

argued by Stone *et al.*, 2012), one might have expected the broader AM tuning found in older NH

listeners to lead to an increase in AM masking. However, the SRTs obtained with the SSN likely reflect

aspects of EM rather than AM masking. This is consistent with the extensive overlap observed in the

audio spectra of the SSN and HINT speech material (Figure 4, panel A), compared to the highly

positive SNR_{env} observed when comparing their modulation spectra at low modulation frequencies

(Figure 4, panel F). These findings indicate that the SSN provided more EM than AM masking.

Conversely, the differences in modulation spectra between the target speech and masking noise were

much smaller for the other maskers, suggesting that AM masking (and thus potentially AM frequency

selectivity) may have played a more prominent role in those cases (Figure 4, panels G-J). Notably, the

older NH listeners exhibited a reduced release from masking when exposed to the fluctuating ICRA-

5 noise compared to the young NH listeners, consistent with previous studies (Dubno *et al.*, 2002,

2003; Grose *et al.*, 2009; Goossens *et al.*, 2017). In addition to potentially being the result of an age-

related increase in forward masking (e.g., Gifford *et al.*, 2007), the reduced release from masking

observed in the older NH listeners may also be attributed to increased AM masking due to the loss of

AM frequency selectivity, which may have limited the benefit derived from the slow temporal

fluctuations in the ICRA-5 condition.

Regarding the effects of hearing loss, the higher SRTs observed in the older HI listeners compared

to the older NH group are consistent with the expectation that the increase in AM frequency selectivity

observed in HI listeners (as reported in Regev *et al.*, 2024a) would *not* lead to improved speech

intelligibility. Instead, it was hypothesized that older HI listeners also experience an age-related loss of

AM selectivity that contributes to poorer speech intelligibility. This was based on the proposition by

Regev *et al.* (2024a) that the greater AM frequency selectivity observed with hearing loss is an apparent

effect resulting from improved detection of the AM targets during temporal dips in the masker, due

to reduced cochlear compression. Since the loss of compression is also associated with other supra-

threshold deficits, such as loudness recruitment, poorer recovery from forward masking, and reduced

frequency selectivity (Glasberg and Moore, 1986, 1989; Oxenham and Moore, 1997; Moore and

Oxenham, 1998), it is evident that the process potentially underlying the apparent improved AM

frequency selectivity is also linked to the mechanisms underlying poorer speech intelligibility in HI

listeners. However, while poorer speech intelligibility was expected in the HI listeners, disentangling

the potential impact of the effective sharpening in AM tuning with hearing loss on speech intelligibility

is not straightforward. It remains unclear whether the process underlying the apparent sharpening of

AM tuning with hearing loss a) improves SRTs but is outweighed by co-occurring supra-threshold

deficits typically associated with hearing loss, or b) itself ultimately impairs speech intelligibility. The

group-level results obtained for the older HI listeners do not clarify whether an age-related reduction

in AM selectivity contributed to the increase in SRTs, beyond the effects of hearing loss.

However, the results of the multilinear regression analysis allowed for a separate investigation of

these effects, further supporting the existence of a link between alterations in AM frequency selectivity

and speech intelligibility challenges in noise for older listeners. The addition of DR_{MTP} to the model

significantly increased the amount of variance in SRTs explained by the model. Furthermore, the

negative structure coefficient for DR_{MTP} indicated that an increase in the SRTs estimated by the best-

fitting model (which included age, PTA, and DR_{MTP}) is linked with a decrease in DR_{MTP}. This finding

aligned with the expectation that poorer AM frequency selectivity (quantified by a lower DR_{MTP}) would

be linked with higher SRTs.[7] This link, identified in a model including all listeners, is also broadly

consistent with the proposition by Regev *et al.* (2024a) that older HI listeners experience an age-related

reduction in AM frequency selectivity, which is concealed in the MTPs by a perceptual benefit

stemming from the loss of cochlear compression (represented by PTA in the regression model).

Employed to address the likely collinearity across predictors, the commonality analysis revealed

that DR_{MTP} acts as a suppressor, as shown by the negative R^2 shared between DR_{MTP} and age (Capraro

and Capraro, 2001). As such, the addition of DR_{MTP} to the model helped increase the R^2 uniquely

attributable to age, thus increasing the model's total R^2. Within the model, age can be seen as a "noisy

term" representing the sum of all factors underlying auditory and cognitive processing deficits in older

listeners that are distinct from the effects of peripheral hearing loss and reduced working-memory

capacity. Such factors, which are likely somewhat collinear, may include cochlear synaptopathy

(Parthasarathy *et al.*, 2019a), reduced neural inhibition (Caspary *et al.*, 2008), or poorer temporal

envelope processing (Anderson and Karawani, 2020) such as a loss of AM frequency selectivity (Regev

et al., 2023). Hence, the greater R^2 resulting from the explicit addition of a measure of AM frequency

selectivity to the regression model can reasonably be interpreted as resulting from a "denoising" of

the age term, allowing it to more accurately account for the variance in SRTs.

However, it should be noted that the significant contribution of DR_{MTP} to the regression model

was not observed when the model was applied to SRTs for maskers other than the male competing

talker or DR_{MTP} derived from MTPs at modulation rates other than 4 Hz. For the SSN and cocktail-

party conditions, this may partly result from the limited variance in SRTs across listeners (see Figure

5 A and E). For the ICRA-5 and female competing talker conditions, only the main effect of PTA

significantly contributed to explaining the variance in SRTs. This dominance of the PTA effect and

the unexpected lack of an age effect may arise from the overlap in SRTs between young and older NH

listeners (though significantly different) for these conditions, while the older HI listeners exhibited

substantially higher SRTs compared to the NH groups (see Figure 5 B and D). Hence, the age effect

in these conditions may not have been strong enough for an age-related loss in AM frequency

selectivity to influence intelligibility. Overall, while the reasons for the inconsistent effects of age and

DR_{MTP} remain unclear, the specific selection of the SRTs for the male competing talker and the DR_{MTP}

derived from a 4-Hz modulation rate was intentional in order to maximize the variability of dependent

and independent variables in the regression model. The male competing talker yielded the largest SRT

differences between young and older NH listeners and the highest SRTs for the older NH and HI

listeners compared to other maskers. Additionally, the 4-Hz MTP showed the largest differences in

AM selectivity between young and older NH listeners (Regev *et al.*, 2023). Finally, the 4-Hz MTP likely

reflects the prominent role of this modulation rate in speech perception (e.g., Elliott and Theunissen,

2009; Varnet *et al.*, 2017), making it more perceptually relevant than MTPs obtained at other rates.

Notably, the present findings indicate that declines in speech intelligibility with age and hearing

loss are most pronounced in contexts involving high cognitive demand and informational masking

(IM), such as in the presence of competing-talker maskers. IM includes factors that complicate the

separation target and masker speech signals, such as stream segregation (Brungart et al., 2001; Darwin

et al., 2003; Schneider et al., 2007; Kidd et al., 2016; see Kidd et al., 2008 for a review)). In this study,

both older NH and older HI listeners exhibited lower RDS scores compared to the young NH listeners

(significantly and marginally so, respectively). Such reduced cognitive abilities have been implicated in

age-related increases of SRTs under conditions of IM (Rajan and Cainer, 2008; Gordon-Salant and

Cole, 2016; Goossens *et al.*, 2017) and in the pronounced effects of hearing loss in cognitively

demanding settings (e.g., Schneider *et al.*, 2010; Goossens *et al.*, 2017). Thus, the poorer speech

intelligibility observed in the older NH and HI listeners in the present study may partly reflect reduced

working-memory capacity. However, significant effects of hearing loss and age were also found for

the ICRA-5 condition, which does not impose high IM or cognitive demand. Furthermore, the

multilinear regression analysis showed that RDS scores did not contribute to explaining the variance

in SRTs for the male competing talker across all listeners, suggesting that working-memory capacity

did not significantly influence the observed speech intelligibility results. It is important to note that

working-memory capacity is only one of several cognitive abilities that may be reduced in older

listeners, potentially contributing to challenges in noisy environments (e.g., Dryden *et al.*, 2017).

Overall, while reduced cognitive ability may have partly mediated the effects of age and hearing loss

on speech intelligibility, the results also point to supra-threshold deficits. These may involve age-

related temporal processing deficits, as suggested by perceptual (Pichora-Fuller and Souza, 2003;

Pichora-Fuller and MacDonald, 2008; Helfer and Vargo, 2009; Füllgrabe et al., 2015; see Fitzgibbons

and Gordon-Salant, 2010 for a review) and neurophysiological research (e.g., Anderson et al., 2012;

Decruy et al., 2019; Parthasarathy et al., 2019b; Roque et al., 2019; McFarlane and Sanchez, 2024; for

reviews see Parthasarathy et al., 2019a and Anderson and Karawani, 2020), or reduced cochlear

compression (e.g., Glasberg and Moore, 1989; Oxenham and Moore, 1997; Moore and Oxenham,

1998; Moore, 2003). The hypothesis that an age-related reduction in AM frequency selectivity

contributes to increased SRTs aligns with these findings from previous studies.

Although the data from the present study revealed significant group-level and regression results

supporting a relationship between changes in AM tuning and speech intelligibility in noise for older

NH and HI listeners, it is important to consider several co-occurring factors that may have influenced

the findings. Even though all the older NH listeners exhibited clinically normal hearing up to 4 kHz,

their audiometric thresholds were on average 8 dB higher than those of the young NH listeners

between 125 Hz and 8 kHz, with the greatest differences occurring at 8 kHz. Although these

differences in audiometric thresholds are small, previous studies have suggested that they can have a

considerable effect on speech intelligibility (Dubno and Ahlstrom, 1997; Dubno *et al.*, 2000), possibly

due to related supra-threshold processing deficits (e.g., Léger *et al.*, 2012). Moreover, the older NH

listeners included in the present study may have experienced reduced audibility at extended high

frequencies (> 8 kHz), which have also been suggested to play a role in speech intelligibility (e.g.,

Hunter *et al.*, 2020; Polspoel *et al.*, 2022). These aspects may have contributed to the observed age-

related increase in SRTs and are consistent with the significant effect of PTA found in the regression

analysis of the SRTs.

Concerning the experimental approach, since the tests provided listeners with cues beyond the

temporal envelope, factors such as IM, EM, and stream segregation likely interacted with age- and

hearing loss-related perception deficits, as well as the age-related reduction in working-memory

capacity discussed above. Therefore, the connection between SRTs and AM frequency selectivity

suggested by the present results warrants further investigation. Notably, a regression analysis using the

SRTs from the ICRA-5 condition, which is not affected by IM, revealed that only PTA significantly

contributed to explaining the variance in SRTs. This suggests that the effect of age on SRTs in this

condition was not sufficiently strong for an age-related loss of AM selectivity to play a role. Overall,

an experimental paradigm limiting the perceptual cues available in the speech intelligibility test to the

temporal envelope could help establish a more definitive link between declines in AM selectivity and

speech intelligibility.

Finally, the outcomes of the multilinear regression model are limited by the bimodal distributions

of age and PTA across listeners, which may have contributed to the strong effect of age found in the

model. Since the studies by Regev *et al.* (2023, 2024a) focused on groups of young NH, older NH, and

older HI listeners, both the age and PTA dimensions are constrained to low or high values. A study

recruiting a larger sample with more continuous distributions of age and PTA would enable a stronger

regression model and more conclusive outcomes. Moreover, recruiting NH listeners across a

continuum of ages would allow for a regression analysis focusing exclusively on the effects of AM

selectivity on SRTs in isolation from hearing loss, potentially providing clearer evidence of a link

between these measures.

V.   SUMMARY AND CONCLUSION

Speech intelligibility in noise was examined in three groups: young listeners with normal hearing

(aged 22-28 years), older listeners with normal hearing (aged 57-77 years), and older listeners with

hearing impairment (aged 64 – 77) years. All participants had previously taken part in AM masking

studies investigating AM frequency selectivity. These studies revealed an age-related reduction in AM

tuning (Regev *et al.*, 2023), and an apparent sharpening in AM tuning with hearing loss (Regev *et al.*,

2024a), which was attributed to reduced cochlear compression rather than a true sharpening of the

theoretical modulation filters. This study aimed to determine whether the effects of age and hearing

loss on speech intelligibility in noise support the hypothesis that an age-related reduction in AM tuning

in older NH and HI listeners negatively impacts speech intelligibility.

Consistent with the hypothesis, the results showed an age-related decline and an additional

negative effect of hearing loss on speech intelligibility, especially for fluctuating maskers. Furthermore,

a multilinear regression analysis showed that adding a measure of AM frequency selectivity to a model

using age and audibility as predictors significantly increased the amount of variance in SRTs explained

across all listeners. This finding supported the assumption that an age-related loss of AM frequency

selectivity is linked to speech intelligibility difficulties in noise in older NH and HI listeners, once

audibility is accounted for. Moreover, the results broadly aligned with the proposition made by Regev

et al. (2024a) that the improved AM tuning in listeners with hearing impairment is an "apparent" effect

resulting from reduced peripheral compression, which is known to contribute to other supra-threshold

deficits that impair speech intelligibility.

Overall, this study's results support the notion that an age-related reduction in AM frequency

selectivity may contribute to speech intelligibility challenges in noise and warrant further investigation

into this potential mechanism. To more directly assess the effects of AM tuning, and a loss thereof, on speech intelligibility, future studies may focus on speech-intelligibility testing paradigms that specifically examine envelope-based cues.

ACKNOWLEDGMENTS

The authors thank Borgný Súsonnudóttir Hansen for her contribution to the data collection. The authors also thank Christian Stender Simonsen for kindly sharing the experimental framework used to run the listening test, as well as several of the signal recordings, and Jens Hjortkjær and Jonatan Märcher-Rørsted for kindly providing their implementations of the reverse digit span test and of the Cambridge formula (CamEq; Moore and Glasberg, 1998). We thank Dr. Christopher Brown and two anonymous reviewers for their insightful comments on earlier versions of this manuscript. Finally, the authors thank Raul Sanchez-Lopez for helpful insights and discussions. This study was supported by the Oticon Centre of Excellence for Hearing and Speech Sciences (CHeSS).

AUTHOR DECLARATIONS

Conflict of Interest

The authors have no conflict of interest to report.

Ethics Approval

All participants provided written informed consent and were offered financial compensation for their participation. Ethical approval for the study was provided by the Science Ethics Committee of the Capital Region in Denmark (reference H-16036391).

 DATA AVAILABILITY

This manuscript has been submitted to the Journal of the Acoustical Society of America. Once it

has been accepted, the data that support the findings of this study will be openly available from: Regev,

J., Zaar, J., Relaño-Iborra, H., and Dau, T. (2024). "Dataset for: 'Investigating the effects of age and

hearing loss on speech intelligibility and amplitude modulation frequency selectivity'", Technical

University of Denmark. Dataset. https://doi.org/10.11583/DTU.25771884.

APPENDIX

A. Post-hoc analyses

Table 1 summarizes the results of the post-hoc analyses that compared the SRTs obtained with

young and older NH listeners for each masker.

Table 1: Post-hoc analysis of the differences in SRT between young and older NH listeners for each

masker. The effect sizes (M), standard errors (SE, and corrected p-values are reported. The significant

p-values are highlighted in bold.

Masker	M [dB]	SE [dB]	Corrected p-value
SSN	0.75		0.279
ICRA-5	1.39		**0.047**
Male competing	4.53	0.68	**< 0.001**
Female competing	2.70		**< 0.001**
Cocktail party	1.53		**0.030**

577   Table 2 summarizes the results of the post-hoc analyses that compared the SRTs obtained with

578   older NH and older HI listeners for each masker.

580   Table 2: Post-hoc analysis of the differences in SRT between older NH and older HI listeners for each

581   masker. The effect sizes (M), standard errors (SE, and corrected p-values are reported. The significant

582   p-values are highlighted in bold.

Masker	M [dB]	SE [dB]	Corrected p-value
SSN	2.36		**0.048**
ICRA-5	4.30		**< 0.001**
Male competing	3.60	1.14	**0.004**
Female competing	5.52		**< 0.001**
Cocktail party	2.24		0.060

Table 3 summarizes the results of the post-hoc analyses that compared the SRTs obtained with

the different maskers for each of the three listener groups (young NH, older NH, and older HI). Since

this post-hoc analysis was run twice for the older NH listeners (once for each statistical model applied),

the values reported reflect the most conservative p-values and standard errors, obtained from the

model fitted to the older NH and older HI listeners. However, the effect sizes and overall statistical

outcomes did not differ between the analyses run in the two models.

Table 3: Post-hoc analysis of the differences in SRT between older the different maskers for each group. The effect sizes (M), standard errors (SE, and corrected p-values are reported. The significant p-values are highlighted in bold.

Listener group	Maskers compared	M [dB]	SE [dB]	Corrected p-value
Young NH	SSN - ICRA-5	2.35	0.48	**< 0.001**
	SSN – Male competing	0.83		0.086
	SSN – Female competing	9.23		**< 0.001**
	SSN – Cocktail party	-1.42		**0.008**
	ICRA-5 – Male competing	-1.52		**0.006**
	ICRA-5 – Female competing	6.88		**< 0.001**
	ICRA-5 – Cocktail party	-3.77		**< 0.001**
	Male competing – Female competing	8.40		**< 0.001**
	Male competing – Cocktail party	-2.25		**< 0.001**
	Female competing – Cocktail party	-10.65		**< 0.001**
Older NH	SSN - ICRA-5	1.71	0.60	**0.011**
	SSN – Male competing	-2.95		**< 0.001**
	SSN – Female competing	7.28		**< 0.001**
	SSN – Cocktail party	-2.20		**0.001**
	ICRA-5 – Male competing	-4.66		**< 0.001**

	ICRA-5 – Female competing	5.57		**< 0.001**
	ICRA-5 – Cocktail party	-3.91		**< 0.001**
	Male competing – Female competing	10.23		**< 0.001**
	Male competing – Cocktail party	0.75		0.212
	Female competing – Cocktail party	-9.48		**< 0.001**
Older HI	SSN - ICRA-5	-0.23	0.60	0.696
	SSN – Male competing	-4.20		**< 0.001**
	SSN – Female competing	4.12		**< 0.001**
	SSN – Cocktail party	-2.09		**0.003**
	ICRA-5 – Male competing	-3.96		**< 0.001**
	ICRA-5 – Female competing	4.35		**< 0.001**
	ICRA-5 – Cocktail party	-1.85		**0.006**
	Male competing – Female competing	8.32		**< 0.001**
	Male competing – Cocktail party	2.11		**0.003**
	Female competing – Cocktail party	-6.20		**< 0.001**

B. Regression analysis

Table 4 summarizes the details of the best-fitting model used in the multilinear regression analysis.

Table 4: Details of the multilinear regression analysis relating the SRTs obtained for the male

competing talker to age, PTA, and DR_{MTP}. For each predictor, the table reports the standardized

regression coefficients (β), the 95%-confidence interval for β (lower | upper), as well as the structure

coefficients (r_s).

Predictor	Standardized β	95% CI β (lower | upper)	r_s
Age	0.7726	0.4892 | 1.056	0.8964
PTA	0.3025	0.0452 | 0.5598	0.8422
DR_{MTP}	0.3504	0.111 | 0.5898	-0.1569

605   Table 5 summarizes the details of the commonality analysis conducted on the best-fitting model

606   (relating SRTs to age, PTA, and DR_{MTP}), which identifies the R^2 either uniquely attributable to each

607   predictor or shared among them.

612 Table 5: Outcome of the commonality analysis applied on the best-fitting model, that related SRTs

613 obtained for a male competing talker to age, PTA, and the 4-Hz MTP's average dynamic range

614 (DR$_{MTP}$). The table reports the amount of variance in SRTs explained (R^2) either uniquely explained

615 by each predictor (single variables) or commonly shared between predictors (two or more variables).

Predictor	Age	PTA	DR$_{MTP}$	Age and PTA	Age and DR$_{MTP}$	PTA and DR$_{MTP}$	Age, PTA, and DR$_{MTP}$	Total
R^2 [%]	21.96	4.33	8.30	48.78	-7.89	2.49	-1.00	76.97

---

[1] The classification of the audiograms was performed using the lowest mean squared error between the standard audiograms and the collected audiometric thresholds at frequencies from 1 to 8 kHz.

[2] Except for one listener who showed a gap of 20 dB at 1 kHz.

[3] This was shown by the post-hoc analysis run on a linear model ANOVA using estimated marginal means. The older NH listeners had significantly lower RDS scores than the young NH and older HI listeners [$M = 0.21$, $SE = 0.05$ dB; $M = 0.13$, $SE = 0.05$ dB, respectively].

[4] Due to a mistake in the audibility compensation, one of the older HI listeners was tested twice on the entire HINT corpus. However, more than 7 weeks separated the two testing sessions. Nielsen and Dau (2011) reported an average improvement in SRT of only 0.5 dB across two sessions separated by three weeks. Hence, the current case was likely not affected by memorization of the speech material.

[5] The amplification produced aided thresholds of at most 35 dB until 4 kHz, 45 dB at 6 kHz, and 55 dB at 8 kHz. Hence, while reasonable audibility was achieved at low frequencies, audibility was not fully restored to levels comparable to those of the older NH listeners. However, a maximum gain of 30 dB was applied to ensure that no listener was exposed to uncomfortable or potentially hazardous sound levels during the speech intelligibility test, where the unaided masker levels could reach up to 80 dB SPL.

[6] Levene's test was used to assess the homoskedasticity of the residuals and Shapiro-Wilk's test was used to assess the normality of residual's distribution. Shapiro-Wilk's test was not passed for either model, likely due to the relatively small group sizes and to the presence of outliers in the older HI data. However, no other aspect of the data justified the removal of the outliers, and they were therefore kept in the dataset. Furthermore, the average fittings of the models to the experimental data were found to be accurate, and the models were hence considered as appropriate.

[7] Structure coefficients are employed here since they have been suggested to be more robust than regression coefficients in cases involving multicollinearity among predictors (Thompson and Borrello, 1985; Kraha *et al.*, 2012)

REFERENCES

Anderson, S., and Karawani, H. (**2020**). "Objective evidence of temporal processing deficits in older adults," Hear Res, **397**, 108053. doi:10.1016/j.heares.2020.108053

Anderson, S., Parbery-Clark, A., White-Schwoch, T., and Kraus, N. (**2012**). "Aging affects neural precision of speech encoding," J Neurosci, **32**, 14156–14164. doi:10.1523/JNEUROSCI.2176-12.2012

Arai, T., Pavel, M., Hermansky, H., and Avendano, C. (**1999**). "Syllable intelligibility for temporally filtered LPC cepstral trajectories," J Acoust Soc Am, **105**, 2783–2791. doi:10.1121/1.426895

Bacon, S. P., and Grantham, D. W. (**1989**). "Modulation masking: Effects of modulation frequency, depth, and phase," J Acoust Soc Am, **85**, 2575–2580. doi:10.1121/1.397751

Bernstein, J. G. W., Danielsson, H., Hällgren, M., Stenfelt, S., Rönnberg, J., and Lunner, T. (**2016**). "Spectrotemporal modulation sensitivity as a predictor of speech-reception performance in noise with hearing aids," Trends Hear, **20**, 233121651667038. doi:10.1177/2331216516670387

Bisgaard, N., Vlaming, M. S. M. G., and Dahlquist, M. (**2010**). "Standard audiograms for the IEC 60118-15 measurement procedure," Trends Amplif, **14**, 113–120. doi:10.1177/1084713810379609

Brungart, D. S., Simpson, B. D., Ericson, M. A., and Scott, K. R. (**2001**). "Informational and energetic masking effects in the perception of multiple simultaneous talkers," J Acoust Soc Am, **110**, 2527–2538. doi:10.1121/1.1408946

Capraro, R. M., and Capraro, M. M. (**2001**). "Commonality analysis: understanding variance contributions to overall canonical correlation effects of attitude toward mathematics on geometry achievement," Multiple Linear Regression Viewpoints, **27**, 16–23.

Caspary, D. M., Ling, L., Turner, J. G., and Hughes, L. F. (**2008**). "Inhibitory neurotransmission, plasticity and aging in the mammalian central auditory system," J Exp Biol, **211**, 1781–1791. doi:10.1242/jeb.013581

Chabot-Leclerc, A., Jørgensen, S., and Dau, T. (**2014**). "The role of auditory spectro-temporal modulation filtering and the decision metric for speech intelligibility prediction," J Acoust Soc Am, **135**, 3502–3512. doi:10.1121/1.4873517

Darwin, C. J., Brungart, D. S., and Simpson, B. D. (**2003**). "Effects of fundamental frequency and vocal-tract length changes on attention to one of two simultaneous talkers," J Acoust Soc Am, **114**, 2913. doi:10.1121/1.1616924

Dau, T., Kollmeier, B., and Kohlrausch, A. (**1997**). "Modeling auditory processing of amplitude modulation. I. Detection and masking with narrow-band carriers," J Acoust Soc Am, **102**, 2892–2905. doi:10.1121/1.420344

Dau, T., Verhey, J., and Kohlrausch, A. (**1999**). "Intrinsic envelope fluctuations and modulation-detection thresholds for narrow-band noise carriers," J Acoust Soc Am, **106**, 2752–2760. doi:10.1121/1.428103

Decruy, L., Vanthornhout, J., and Francart, T. (**2019**). "Evidence for enhanced neural tracking of the speech envelope underlying age-related speech-in-noise difficulties," J Neurophysiol, **122**, 601–615. doi:10.1152/jn.00687.2018

Decruy, L., Vanthornhout, J., and Francart, T. (**2020**). "Hearing impairment is associated with enhanced neural tracking of the speech envelope," Hear Res, **393**, 107961. doi:10.1016/j.heares.2020.107961

Dorman, M. F., Loizou, P. C., and Rainey, D. (**1997**). "Speech intelligibility as a function of the number of channels of stimulation for signal processors using sine-wave and noise-band outputs," J Acoust Soc Am, **102**, 2403–2411. doi:10.1121/1.419603

Dreschler, W. A., Verschuure, H., Ludvigsen, C., and Westermann, S. (**2001**). "ICRA noises: artificial noise signals with speech-like spectral and temporal properties for hearing instrument assessment," Int J Audiol, **40**, 148–157. doi:10.3109/00206090109073110

Drullman, R., Festen, J. M., and Plomp, R. (**1994**). "Effect of temporal envelope smearing on speech reception," J Acoust Soc Am, **95**, 1053–1064. doi:10.1121/1.408467

Dryden, A., Allen, H. A., Henshaw, H., and Heinrich, A. (**2017**). "The association between cognitive performance and speech-in-noise perception for adult listeners: a systematic literature review and meta-analysis," Trends Hear, **21**, 233121651774467. doi:10.1177/2331216517744675

Dubbelboer, F., and Houtgast, T. (**2008**). "The concept of signal-to-noise ratio in the modulation domain and speech intelligibility," J Acoust Soc Am, **124**, 3937–3946. doi:10.1121/1.3001713

Dubno, J. R., and Ahlstrom, J. B. (**1997**). "Additivity of multiple maskers of speech," In W. Jesteadt (Ed.), Modeling Sensorineural Hearing Loss, Routledge, New York, pp. 253–272. doi:10.4324/9781315789392-21

Dubno, J. R., Ahlstrom, J. B., and Horwitz, A. R. (**2000**). "Use of context by young and aged adults with normal hearing," J Acoust Soc Am, **107**, 538–546. doi:10.1121/1.428322

Dubno, J. R., Dirks, D. D., and Morgan, D. E. (**1984**). "Effects of age and mild hearing loss on speech recognition in noise," J Acoust Soc Am, **76**, 87–96. doi:10.1121/1.391011

Dubno, J. R., Horwitz, A. R., and Ahlstrom, J. B. (**2002**). "Benefit of modulated maskers for speech recognition by younger and older adults with normal hearing," J Acoust Soc Am, **111**, 2897–2907. doi:10.1121/1.1480421

Dubno, J. R., Horwitz, A. R., and Ahlstrom, J. B. (**2003**). "Recovery from prior stimulation:

Masking of speech by interrupted noise for younger and older adults with normal

hearing," J Acoust Soc Am, **113**, 2084–2094. doi:10.1121/1.1555611

Elliott, T. M., and Theunissen, F. E. (**2009**). "The modulation transfer function for speech

intelligibility," PLoS Comput Biol, **5**, e1000302. doi:10.1371/journal.pcbi.1000302

Ewert, S. D., and Dau, T. (**2000**). "Characterizing frequency selectivity for envelope

fluctuations," J Acoust Soc Am, **108**, 1181–1196. doi:10.1121/1.1288665

Ewert, S. D., Verhey, J. L., and Dau, T. (**2002**). "Spectro-temporal processing in the envelope-

frequency domain," J Acoust Soc Am, **112**, 2921–2931. doi:10.1121/1.1515735

Fitzgibbons, P. J., and Gordon-Salant, S. (**2010**). "Behavioral studies with aging humans: hearing

sensitivity and psychoacoustics," In S. Gordon-Salant, R. Frisina, A. Popper, and R. Fay

(Eds.), The Aging Auditory System. Springer Handbook of Auditory Research, Springer,

New York, Ny, Vol. 34, pp. 111–134. doi:10.1007/978-1-4419-0993-0_5

Frisina, D. R., and Frisina, R. D. (**1997**). "Speech recognition in noise and presbycusis: relations

to possible neural mechanisms," Hear Res, **106**, 95–104. doi:10.1016/S0378-

5955(97)00006-3

Fuglsang, S. A., Märcher-Rørsted, J., Dau, T., and Hjortkjær, J. (**2020**). "Effects of sensorineural

hearing loss on cortical synchronization to competing speech during selective attention,"

J Neurosci, **40**, 2562–2572. doi:10.1523/JNEUROSCI.1936-19.2020

Füllgrabe, C., Moore, B. C. J., and Stone, M. A. (**2015**). "Age-group differences in speech

identification despite matched audiometrically normal hearing: contributions from

auditory temporal processing and cognition," Front Aging Neurosci, **6**, 1–25.

doi:10.3389/fnagi.2014.00347

Gallun, F. J., Coco, L., Koerner, T. K., Larrea-Mancera, E. S. L. de, Molis, M. R., Eddins, D. A.,

and Seitz, A. R. (**2022**). "Relating suprathreshold auditory processing abilities to speech

understanding in competition," Brain Sci, **12**, 695. doi:10.3390/brainsci12060695

Gifford, R. H., Bacon, S. P., and Williams, E. J. (**2007**). "An examination of speech recognition

in a modulated background and of forward masking in younger and older listeners," J

Speech Lang Hear Res, **50**, 857–864. doi:10.1044/1092-4388(2007/060)

Glasberg, B. R., and Moore, B. C. J. (**1986**). "Auditory filter shapes in subjects with unilateral and

bilateral cochlear impairments," J Acoust Soc Am, **79**, 1020–1033. doi:10.1121/1.393374

Glasberg, B. R., and Moore, B. C. J. (**1989**). "Psychoacoustic abilities of subjects with unilateral

and bilateral cochlear hearing impairments and their relationship to the ability to

understand speech," Scand Audiol Suppl, **32**, 1–25.

Goossens, T., Vercammen, C., Wouters, J., and Wieringen, A. van (**2016**). "Aging affects neural

synchronization to speech-related acoustic modulations," Front Aging Neurosci, **8**, 1–

16. doi:10.3389/fnagi.2016.00133

Goossens, T., Vercammen, C., Wouters, J., and van Wieringen, A. (**2017**). "Masked speech

perception across the adult lifespan: Impact of age and hearing impairment," Hear Res,

344, 109–124. doi:10.1016/j.heares.2016.11.004

Goossens, T., Vercammen, C., Wouters, J., and van Wieringen, A. (**2018**). "Neural envelope

encoding predicts speech perception performance for normal-hearing and hearing-

impaired adults," Hear Res, **370**, 189–200. doi:10.1016/j.heares.2018.07.012

Gordon-Salant, S., and Cole, S. S. (**2016**). "Effects of age and working memory capacity on

speech recognition performance in noise among listeners with normal hearing," Ear

Hear, **37**, 593–602. doi:10.1097/AUD.0000000000000316

Gordon-Salant, S., Yeni-Komshian, G. H., Fitzgibbons, P. J., and Barrett, J. (**2006**). "Age-related

differences in identification and discrimination of temporal cues in speech segments," J

Acoust Soc Am, **119**, 2455–2466. doi:10.1121/1.2171527

Greenberg, S., Carvey, H., Hitchcock, L., and Chang, S. (**2003**). "Temporal properties of

spontaneous speech—a syllable-centric perspective," J Phon, **31**, 465–485.

doi:10.1016/j.wocn.2003.09.005

Grose, J. H., Mamo, S. K., and Hall, J. W. (**2009**). "Age effects in temporal envelope processing:

speech unmasking and auditory steady state responses," Ear Hear, **30**, 568–575.

doi:10.1097/AUD.0b013e3181ac128f

Helfer, K. S., and Vargo, M. (**2009**). "Speech recognition and temporal processing in middle-

aged women," J Am Acad Audiol, **20**, 264–271. doi:10.3766/jaaa.20.4.6

Houtgast, T. (**1989**). "Frequency selectivity in amplitude-modulation detection," J Acoust Soc

Am, **85**, 1676–1680. doi:10.1121/1.397956

Houtgast, T., and Steeneken, H. J. M. (**1985**). "A review of the MTF concept in room acoustics

and its use for estimating speech intelligibility in auditoria," J Acoust Soc Am, **77**, 1069–

1077. doi:10.1121/1.392224

Humes, L. E. (**2002**). "Factors underlying the speech-recognition performance of elderly hearing-

aid wearers," J Acoust Soc Am, **112**, 1112–1132. doi:10.1121/1.1499132

Humes, L. E. (**2007**). "The contributions of audibility and cognitive factors to the benefit

provided by amplified speech to older adults," J Am Acad Audiol, **18**, 590–603.

doi:10.3766/jaaa.18.7.6

Humes, L. E., and Christopherson, L. (**1991**). "Speech identification difficulties of hearing-

impaired elderly persons," J Speech Lang Hear Res, **34**, 686–693.

doi:10.1044/jshr.3403.686

Humes, L. E., and Dubno, J. R. (**2010**). "Factors affecting speech understanding in older adults,"

In S. Gordon-Salant, R. Frisina, and A. F. R. Popper (Eds.), The Aging Auditory System.

Springer Handbook of Auditory Research, Springer, New York, Vol. 34, pp. 211–257.

doi:10.1007/978-1-4419-0993-0_8

Humes, L. E., Dubno, J. R., Gordon-Salant, S., Lister, J. J., Cacace, A. T., Cruickshanks, K. J.,

Gates, G. A., et al. (**2012**). "Central presbycusis: A review and evaluation of the evidence,"

J Am Acad Audiol, **23**, 635–666. doi:10.3766/jaaa.23.8.5

Humes, L. E., Lee, J. H., and Coughlin, M. P. (**2006**). "Auditory measures of selective and divided

attention in young and older adults using single-talker competition.," J Acoust Soc Am,

120, 2926–37. doi:10.1121/1.2354070

Hunter, L. L., Monson, B. B., Moore, D. R., Dhar, S., Wright, B. A., Munro, K. J., Zadeh, L. M.,

et al. (**2020**). "Extended high frequency hearing and speech perception implications in

adults and children," Hear Res, **397**, 107922. doi:10.1016/j.heares.2020.107922

Jennings, S. G., Chen, J., Fultz, S. E., Ahlstrom, J. B., and Dubno, J. R. (**2018**). "Amplitude

modulation detection with a short-duration carrier: Effects of a precursor and hearing

loss," J Acoust Soc Am, **143**, 2232–2243. doi:10.1121/1.5031122

Jepsen, M. L., and Dau, T. (**2011**). "Characterizing auditory processing and perception in

individual listeners with sensorineural hearing loss," J Acoust Soc Am, **129**, 262–281.

doi:10.1121/1.3518768

Jepsen, M. L., Ewert, S. D., and Dau, T. (**2008**). "A computational model of human auditory

signal processing and perception," J Acoust Soc Am, **124**, 422–438.

Jeub, M., Schafer, M., and Vary, P. (**2009**). "A binaural room impulse response database for the

evaluation of dereverberation algorithms," 2009 16th International Conference on

Digital Signal Processing, IEEE, 1–5. doi:10.1109/ICDSP.2009.5201259

Jørgensen, S., and Dau, T. (**2011**). "Predicting speech intelligibility based on the signal-to-noise

envelope power ratio after modulation-frequency selective processing," J Acoust Soc

Am, **130**, 1475–1487. doi:10.1121/1.3621502

Jørgensen, S., Ewert, S. D., and Dau, T. (**2013**). "A multi-resolution envelope-power based model

for speech intelligibility," J Acoust Soc Am, **134**, 436–446. doi:10.1121/1.4807563

Joris, P. X., Schreiner, C. E., and Rees, A. (**2004**). "Neural processing of amplitude-modulated

sounds," Physiol Rev, **84**, 541–577. doi:10.1152/physrev.00029.2003

Kidd, G., Mason, C. R., Richards, V. M., Gallun, F. J., and Durlach, N. I. (**2008**). "Informational

masking," In W. A. Yost, A. N. Popper, and R. R. Fay (Eds.), Auditory Perception of

Sound Sources. Springer Handbook of Auditory Research, Springer, Boston, MA, Vol.

29, pp. 143–189. doi:10.1007/978-0-387-71305-2_6

Kidd, G., Mason, C. R., Swaminathan, J., Roverud, E., Clayton, K. K., and Best, V. (**2016**).

"Determining the energetic and informational components of speech-on-speech

masking," J Acoust Soc Am, **140**, 132–144. doi:10.1121/1.4954748

Kohlrausch, A., Fassel, R., and Dau, T. (**2000**). "The influence of carrier level and frequency on

modulation and beat-detection thresholds for sinusoidal carriers," J Acoust Soc Am, **108**,

723–734. doi:10.1121/1.429605

Kraha, A., Turner, H., Nimon, K., Zientek, L. R., and Henson, R. K. (**2012**). "Tools to support

interpreting multiple regression in the face of multicollinearity," Front Psychol, **3**, 1–16.

doi:10.3389/fpsyg.2012.00044

Léger, A. C., Moore, B. C. J., and Lorenzi, C. (**2012**). "Abnormal speech processing in frequency

regions where absolute thresholds are normal for listeners with high-frequency hearing

loss," Hear Res, **294**, 95–103. doi:10.1016/j.heares.2012.10.002

Lorenzi, C., Gilbert, G., Carn, H., Garnier, S., and Moore, B. C. J. (**2006**). "Speech perception problems of the hearing impaired reflect inability to use temporal fine structure," Proc Natl Acad Sci, **103**, 18866–18869. doi:10.1073/pnas.0607364103

McDermott, J. H., and Simoncelli, E. P. (**2011**). "Sound texture perception via statistics of the auditory periphery: Evidence from sound synthesis," Neuron, **71**, 926–940. doi:10.1016/j.neuron.2011.06.032

McFarlane, K. A., and Sanchez, J. T. (**2024**). "Effects of temporal processing on speech-in-noise perception in middle-aged adults," Biology (Basel), **13**, 371. doi:10.3390/biology13060371

McWalter, R., and Dau, T. (**2017**). "Cascaded amplitude modulations in sound texture perception," Front Neurosci, **11**, 1–12. doi:10.3389/fnins.2017.00485

Millman, R. E., Mattys, S. L., Gouws, A. D., and Prendergast, G. (**2017**). "Magnified neural envelope coding predicts deficits in speech perception in noise," J Neurosci, **37**, 7727–7736. doi:10.1523/JNEUROSCI.2722-16.2017

Moore, B. C. J. (**2003**). "Speech processing for the hearing-impaired: successes, failures, and implications for speech mechanisms," Speech Commun, **41**, 81–91. doi:10.1016/S0167-6393(02)00095-X

Moore, B. C. J., and Glasberg, B. R. (**1998**). "Use of a loudness model for hearing-aid fitting. I. Linear hearing aids," Br J Audiol, **32**, 317–335. doi:10.3109/03005364000000083

Moore, B. C. J., and Oxenham, A. J. (**1998**). "Psychoacoustic consequences of compression in the peripheral auditory system.," Psychol Rev, **105**, 108–124. doi:10.1037/0033-295X.105.1.108

Moore, B. C. J., Wojtczak, M., and Vickers, D. A. (**1996**). "Effect of loudness recruitment on the

perception of amplitude modulation," J Acoust Soc Am, **100**, 481–489.

doi:10.1121/1.415861

Nielsen, J. B., and Dau, T. (**2011**). "The Danish hearing in noise test," Int J Audiol, **50**, 202–208.

doi:10.3109/14992027.2010.524254

Oxenham, A. J., and Moore, B. C. J. (**1997**). "Modeling the effects of peripheral nonlinearity in

listeners with normal and impaired hearing," in W. Jesteadt (Ed), Modeling Sensorineural

Hearing Loss, Routledge, New York, pp. 273–288. doi:10.4324/9781315789392-22

Parthasarathy, A., Bartlett, E. L., and Kujawa, S. G. (**2019a**). "Age-related changes in neural

coding of envelope cues: Peripheral declines and central compensation," Neurosci, **407**,

21–31. doi:10.1016/j.neuroscience.2018.12.007

Parthasarathy, A., Herrmann, B., and Bartlett, E. L. (**2019b**). "Aging alters envelope

representations of speech-like sounds in the inferior colliculus," Neurobiol Aging, **73**,

30–40. doi:10.1016/j.neurobiolaging.2018.08.023

Pichora-Fuller, M. K., and MacDonald, E. (**2008**). "Auditory temporal processing deficits in older

listeners: From a review to a future view of presbycusis," Proceedings of the International

Symposium on Auditory and Audiological Research, Nyborg, Denmark, 291–300.

Pichora-Fuller, M. K., and Souza, P. E. (**2003**). "Effects of aging on auditory processing of

speech," Int J Audiol, **42**, 11–16. doi:10.3109/14992020309074638

Polspoel, S., Kramer, S. E., van Dijk, B., and Smits, C. (**2022**). "The importance of extended

high-frequency speech information in the recognition of digits, words, and sentences in

quiet and noise," Ear Hear, **43**, 913–920. doi:10.1097/AUD.0000000000001142

Rajan, R., and Cainer, K. E. (**2008**). "Ageing without hearing loss or cognitive impairment causes
a decrease in speech intelligibility only in informational maskers," Neurosci, **154**, 784–
795. doi:10.1016/j.neuroscience.2008.03.067

Ray-Mukherjee, J., Nimon, K., Mukherjee, S., Morris, D. W., Slotow, R., and Hamer, M. (**2014**).
"Using commonality analysis in multiple regressions: A tool to decompose regression
effects in the face of multicollinearity," Methods Ecol Evol, **5**, 320–328.
doi:10.1111/2041-210X.12166

Regev, J., Relaño-Iborra, H., Zaar, J., and Dau, T. (**2024a**). "Disentangling the effects of hearing
loss and age on amplitude modulation frequency selectivity," J Acoust Soc Am, **155**,
2589–2602. doi:10.1121/10.0025541

Regev, J., Relaño-Iborra, H., Zaar, J., and Dau, T. (**2024b**). "Dataset for: 'Disentangling the
effects of hearing loss and age on amplitude modulation frequency selectivity,'" Technical
University of Denmark,. doi:10.11583/DTU.25134611

Regev, J., Zaar, J., Relaño-Iborra, H., and Dau, T. (**2023**). "Age-related reduction of amplitude
modulation frequency selectivity," J Acoust Soc Am, **153**, 2298. doi:10.1121/10.0017835

Regev, J., Zaar, J., Relaño-Iborra, H., and Dau, T. (**2024c**). "Dataset for: 'Age-related reduction
of amplitude modulation frequency selectivity,'" Technical University of Denmark,.
doi:10.11583/DTU.25134527

Relaño-Iborra, H., and Dau, T. (**2022**). "Speech intelligibility prediction based on modulation
frequency-selective processing," Hear Res, **426**, 108610.
doi:10.1016/j.heares.2022.108610

Relaño-Iborra, H., May, T., Zaar, J., Scheidiger, C., and Dau, T. (**2016**). "Predicting speech
intelligibility based on a correlation metric in the envelope power spectrum domain," J
Acoust Soc Am, **140**, 2670–2679. doi:10.1121/1.4964505

Relaño-Iborra, H., Zaar, J., and Dau, T. (**2019**). "A speech-based computational auditory signal

processing and perception model," J Acoust Soc Am, **146**, 3306–3317.

doi:10.1121/1.5129114

Roque, L., Karawani, H., Gordon-Salant, S., and Anderson, S. (**2019**). "Effects of age, cognition,

and neural encoding on the perception of temporal speech cues," Front Neurosci, **13**, 1–

15. doi:10.3389/fnins.2019.00749

Schlittenlacher, J., and Moore, B. C. J. (**2016**). "Discrimination of amplitude-modulation depth

by subjects with normal and impaired hearing," J Acoust Soc Am, **140**, 3487–3495.

doi:10.1121/1.4966117

Schneider, B. A., Li, L., and Daneman, M. (**2007**). "How competing speech interferes with speech

comprehension in everyday listening situations," J Am Acad Audiol, **18**, 559–572.

doi:10.3766/jaaa.18.7.4

Schneider, B. A., Pichora-Fuller, K., and Daneman, M. (**2010**). "Effects of senescent changes in

audition and cognition on spoken language comprehension," In S. Gordon-Salant, R.

Frisina, A. Popper, and R. Fay (Eds.), The Aging Auditory System. Springer Handbook

of Auditory Research, Springer, New York, NY, Vol. 34, pp. 167–210. doi:10.1007/978-

1-4419-0993-0_7

Schoof, T., and Rosen, S. (**2014**). "The role of auditory and cognitive factors in understanding

speech in noise by normal-hearing older listeners," Front Aging Neurosci, **6**, 1–14.

doi:10.3389/fnagi.2014.00307

Shannon, R. V., Zeng, F.-G., Kamath, V., Wygonski, J., and Ekelid, M. (**1995**). "Speech

recognition with primarily temporal cues," Science (1979), **270**, 303–304.

doi:10.1126/science.270.5234.303

Singh, N. C., and Theunissen, F. E. (**2003**). "Modulation spectra of natural sounds and ethological theories of auditory processing," J Acoust Soc Am, **114**, 3394–3411. doi:10.1121/1.1624067

Smith, Z. M., Delgutte, B., and Oxenham, A. J. (**2002**). "Chimaeric sounds reveal dichotomies in auditory perception," Nature, **416**, 87–90. doi:10.1038/416087a

Sobon, K. A., Taleb, N. M., Buss, E., Grose, J. H., and Calandruccio, L. (**2019**). "Psychometric function slope for speech-in-noise and speech-in-speech: Effects of development and aging," J Acoust Soc Am, **145**, EL284–EL290. doi:10.1121/1.5097377

Soli, S. D., and Wong, L. L. N. (**2008**). "Assessment of speech intelligibility in noise with the hearing in noise test," Int J Audiol, **47**, 356–361. doi:10.1080/14992020801895136

Steinmetzger, K., Zaar, J., Relaño-Iborra, H., Rosen, S., and Dau, T. (**2019**). "Predicting the effects of periodicity on the intelligibility of masked speech: An evaluation of different modelling approaches and their limitations," J Acoust Soc Am, **146**, 2562–2576. doi:10.1121/1.5129050

Stone, M. A., Füllgrabe, C., and Moore, B. C. J. (**2012**). "Notionally steady background noise acts primarily as a modulation masker of speech," J Acoust Soc Am, **132**, 317–326. doi:10.1121/1.4725766

Stone, M. A., and Moore, B. C. J. (**2014**). "On the near non-existence of 'pure' energetic masking release for speech," J Acoust Soc Am, **135**, 1967–1977. doi:10.1121/1.4868392

Strelcyk, O., and Dau, T. (**2009**). "Relations between frequency selectivity, temporal fine-structure processing, and speech reception in impaired hearing," J Acoust Soc Am, **125**, 3328. doi:10.1121/1.3097469

The Mathworks Inc. (**2015**). "MATLAB 2015b – MathWorks." Retrieved from www.mathworks.com/products/matlab (Last viewed January 1st, 2016)

Thompson, B., and Borrello, G. M. (**1985**). "The importance of structure coefficients in

regression research," Educ Psychol Meas, **45**, 203–209.

doi:10.1177/001316448504500202

Varnet, L., Ortiz-Barajas, M. C., Erra, R. G., Gervain, J., and Lorenzi, C. (**2017**). "A cross-

linguistic study of speech modulation spectra," J Acoust Soc Am, **142**, 1976–1989.

doi:10.1121/1.5006179

Verhey, J. L., Ewert, S. D., and Dau, T. (**2003**). "Modulation masking produced by complex tone

modulators," J Acoust Soc Am, **114**, 2135–2146. doi:10.1121/1.1612489

Viemeister, N. F. (**1979**). "Temporal modulation transfer functions based upon modulation

thresholds," J Acoust Soc Am, **66**, 1364–1380. doi:10.1121/1.383531

Viswanathan, V., Bharadwaj, H. M., Shinn-Cunningham, B. G., and Heinz, M. G. (**2021**).

"Modulation masking and fine structure shape neural envelope coding to predict speech

intelligibility across diverse listening conditions," J Acoust Soc Am, **150**, 2230–2244.

doi:10.1121/10.0006385

Xu, L., Thompson, C. S., and Pfingst, B. E. (**2005**). "Relative contributions of spectral and

temporal cues for phoneme recognition," J Acoust Soc Am, **117**, 3255–3267.

doi:10.1121/1.1886405

Zaar, J., Simonsen, L. B., Dau, T., and Laugesen, S. (**2023**). "Toward a clinically viable spectro-

temporal modulation test for predicting supra-threshold speech reception in hearing-

impaired listeners," Hear Res, **427**, 108650. doi:10.1016/j.heares.2022.108650

Zaar, J., Simonsen, L. B., and Laugesen, S. (**2024**). "A spectro-temporal modulation test for

predicting speech reception in hearing-impaired listeners with hearing aids," Hear Res,

443, 108949. doi:10.1016/j.heares.2024.108949