Impact of age versus hearing loss on amplitude modulation frequency selectivity and speech intelligibility

Jonathan Regev,[1,a] Johannes Zaar,[1,2] Helia Relaño-Iborra,[1] and Torsten Dau[1]

[1] *Hearing Systems Section, Department of Health Technology, Technical University of Denmark, Kongens Lyngby, 2800, Denmark*

[2] *Eriksholm Research Centre, Snekkersten, 3070, Denmark*

Abstract

The perception of amplitude modulations (AMs), which is characterized by a frequency-selective process in the modulation domain, is considered critical for speech intelligibility. Previous studies have provided evidence of an age-related decline in AM frequency selectivity, as well as a notable sharpening of AM tuning associated with hearing loss, possibly due to a perceptual advantage resulting from peripheral compression loss. This study aimed to examine whether speech intelligibility in noisy environments would support the following ideas: i) age-related declines in AM tuning might lead to poorer speech intelligibility, and ii) sharper AM tuning associated with hearing loss would not result in improved speech intelligibility. Young (n=10, 22-28 years) and older listeners with normal hearing (n=9, 57–77 years) as well as older listeners with hearing impairment (n=9, 64-77 years) were included in the investigation. All participants had previously taken part in studies on AM frequency selectivity. Speech intelligibility was tested in various listening conditions, including stationary, fluctuating, and

---
[a] Corresponding author: joreg@dtu.dk

competing-speech maskers. Consistent with the hypothesis, the results revealed an age-related increase in speech reception thresholds, with an additional negative impact of hearing loss. These findings motivate further exploration of the relationship between AM frequency selectivity and speech intelligibility in noisy environments.

I. INTRODUCTION

Temporal envelope fluctuations (also referred to as "amplitude modulations" or AMs) are commonly used to describe sounds that exhibit dynamic variations over time (e.g., Singh and Theunissen, 2003; Elliott and Theunissen, 2009). The auditory system displays a remarkable sensitivity to AMs, as evidenced by physiological studies (see Joris *et al.*, 2004 for a review) and behavioral investigations of AM detection (e.g., Viemeister, 1979; Kohlrausch *et al.*, 2000). Analogous to masking effects in the audio-frequency domain, the detectability of target AMs is disrupted in the presence of separate masking AM components. AM masking has been extensively studied by behavioral investigations (e.g., Bacon and Grantham, 1989; Houtgast, 1989; Dau *et al.*, 1997; Verhey *et al.*, 2003) as well as through recent neurophysiological research (Viswanathan *et al.*, 2021). Behavioral studies of AM masking have provided evidence of a frequency-selective process in the modulation domain, often conceptualized as a series of bandpass modulation filters known as the modulation filterbank (e.g., Dau *et al.*, 1997, 1999; Ewert and Dau, 2000; Ewert *et al.*, 2002). Computational modelling studies have shown that this concept significantly contributes to quantitatively predicting experimental data from various psychoacoustic tasks (e.g., Dau *et al.*, 1997, 1999; Jepsen *et al.*, 2008; Jepsen and Dau, 2011; McDermott and Simoncelli, 2011; McWalter and Dau, 2017), underscoring the importance of AMs in auditory processing and perception.

Beyond their role in 'simple' listening tasks, AMs have been argued to be crucial for accurate speech intelligibility (Houtgast and Steeneken, 1985; Drullman *et al.*, 1994; Smith *et al.*, 2002; Elliott and Theunissen, 2009). The significance of AMs in speech was highlighted by vocoding studies, which demonstrated that intact envelope information conveyed through a few broad spectral bands was sufficient for precise speech recognition in quiet, even when spectral cues were greatly reduced (e.g., Shannon *et al.*, 1995; Dorman *et al.*, 1997; Xu *et al.*, 2005). Based on this, AM masking has been proposed as the primary mechanism underlying speech intelligibility in noise, whereby speech

reception thresholds (SRTs) - reflecting the signal-to-noise ratio required for 50% speech recognition

- are proportional to the modulation-domain signal-to-noise ratio (SNR_{env}) between speech and the

masker (e.g., Dubbelboer and Houtgast, 2008; Jørgensen and Dau, 2011). The central role of AM

masking in speech intelligibility in noise was also emphasized by studies using maskers designed to

contain AMs or to have maximally flat envelopes, which suggested that steady maskers, usually

associated with energetic masking (EM) in the audio-frequency domain, effectively function as AM

maskers due to their inherent envelope fluctuations (Stone *et al.*, 2012; Stone and Moore, 2014).

Computational auditory modeling studies have successfully incorporated these concepts by employing

the SNR_{env} (Jørgensen and Dau, 2011; Jørgensen *et al.*, 2013; Chabot-Leclerc *et al.*, 2014) or correlation-

based measures (Relaño-Iborra *et al.*, 2016, 2019; Steinmetzger *et al.*, 2019) at the output of the

modulation filterbank as decision metrics. Based on these modeling studies, the spectral

decomposition of the stimulus envelope carried out by the modulation filterbank, i.e., AM frequency

selectivity, has been suggested as an important factor for accurately predicting SRTs (see Relaño-

Iborra and Dau, 2022, for a review of different speech intelligibility models using modulation filtering).

Regev *et al.* (2023a, 2023b) investigated the effects of age and hearing loss on the selectivity of

hypothetical modulation filters using a perceptual AM masking task. Regev *et al.* (2023a) found a

general reduction in AM frequency selectivity associated with age, resulting in broader masking

patterns among older compared to young listeners with normal hearing (NH listeners). Conversely,

Regev *et al.* (2023b) demonstrated a general sharpening of masking patterns among young and older

listeners with hearing impairment (HI listeners) compared to similarly aged NH listeners. This finding

suggested that sensorineural hearing loss was accompanied by a perceptual benefit that partially offset

the detrimental effects of age, possibly due to the HI listeners' ability to exploit an increased internal

representation of AM depth resulting from reduced peripheral compression when "listening in the

dips" of the masker. Such alterations in the selectivity of the hypothetical modulation filters due to

age and hearing loss may affect speech intelligibility in noise by either increasing or reducing the

amount of AM masking at the output of the modulation filters.

The effects of age and hearing loss on speech intelligibility have received significant attention in

the literature. Hearing loss is known to cause a decline in speech intelligibility in both quiet and noisy

environments (e.g., Glasberg and Moore, 1989; Humes and Christopherson, 1991; Frisina and Frisina,

1997; see Humes and Dubno, 2010 for a review). In addition to audibility loss, speech intelligibility

challenges associated with hearing loss have been linked to various supra-threshold deficits, including

reduced frequency selectivity, poorer temporal fine structure processing, reduced recovery from

forward masking, poorer AM depth discrimination, and loudness recruitment (Glasberg and Moore,

1986, 1989; Oxenham and Moore, 1997; Moore and Oxenham, 1998; Moore, 2003; Lorenzi *et al.*,

2006; Strelcyk and Dau, 2009; Schlittenlacher and Moore, 2016). Aided speech intelligibility

performance has also been shown to be strongly correlated with deficits in spectro-temporal

modulation detection in HI listeners (e.g., Bernstein *et al.*, 2016; Zaar *et al.*, 2023). Recent

neurophysiological studies have suggested that poorer speech intelligibility in HI listeners may be

related to increased neural coding of the envelope (Millman *et al.*, 2017; Goossens *et al.*, 2018; Decruy

et al., 2020). This effect may be partly attributed to an increased internal representation of the AM

depth resulting from the loss of cochlear compression commonly associated with sensorineural

hearing loss (Moore *et al.*, 1996; Jennings *et al.*, 2018). In addition to the effects of hearing loss itself,

age and reduced cognitive function have been found to contribute to the decline in speech intelligibility

observed in older HI listeners (e.g., Dubno *et al.*, 1984; Humes and Christopherson, 1991; Frisina and

Frisina, 1997; Humes, 2002, 2007; Humes *et al.*, 2006), suggesting that cognitive and age-related

perception deficits may also play a role in speech intelligibility challenges.

To investigate these potential mechanisms, cross-sectional studies focusing on age-related effects

have been conducted, often in listeners without or with mild hearing loss. While these studies have

consistently shown a detrimental effect of age on speech intelligibility in the presence of competing

speech maskers (Rajan and Cainer, 2008; Helfer and Vargo, 2009; Schoof and Rosen, 2014; Füllgrabe

et al., 2015; Gordon-Salant and Cole, 2016; Goossens *et al.*, 2017; Decruy *et al.*, 2019; Sobon *et al.*,

2019), the effects of age in stationary and fluctuating noise have been less consistent. Some studies

have found differences between young and older listeners (e.g., Goossens *et al.*, 2017; Decruy *et al.*,

2019; Sobon *et al.*, 2019), while others have not (Rajan and Cainer, 2008; Helfer and Vargo, 2009;

Schoof and Rosen, 2014). Overall, the age-related increase in SRTs has been found to be more

pronounced for speech maskers than for stationary or fluctuating noises (e.g., Goossens *et al.*, 2017;

Decruy *et al.*, 2019; Sobon *et al.*, 2019). The observed age-related decline in speech intelligibility has

been suggested to be mediated by deficits in temporal processing (e.g., Gordon-Salant *et al.*, 2006;

Anderson *et al.*, 2012; Füllgrabe *et al.*, 2015; Goossens *et al.*, 2016; Decruy *et al.*, 2019), as well as by

reduced cognitive abilities (Humes *et al.*, 2012; Füllgrabe *et al.*, 2015; Gordon-Salant and Cole, 2016).

Since AM frequency selectivity has been identified as a significant factor contributing to accurate

speech intelligibility, it is reasonable to suggest that the age-related decline in AM selectivity reported

by Regev *et al.* (2023a) may contribute to the aforementioned challenges in speech intelligibility. A

reduction in AM tuning would likely result in increased AM masking, which could disrupt the

intelligibility of the target speech. Additionally, the observed beneficial effects of hearing loss on AM

frequency selectivity, as reported by Regev *et al.* (2023b), may also impact speech intelligibility in noisy

environments. However, although the increased AM frequency selectivity found in older HI listeners

compared to older NH listeners (Regev *et al.*, 2023a, 2023b) may suggest a decrease in AM masking

with hearing loss, it is hypothesized that this would not directly translate into improved speech

intelligibility. Indeed, the greater AM frequency selectivity in HI listeners is believed, as proposed by

Regev *et al.* (2023b), to be linked to reduced cochlear compression, which is known to be connected

to various supra-threshold deficits that impair speech intelligibility (e.g., Glasberg and Moore, 1989;

Oxenham and Moore, 1997; Moore and Oxenham, 1998; Moore, 2003). Furthermore, the assumption

that a sharpening in AM tuning with hearing loss would not lead to an improvement in speech

intelligibility is consistent with previous studies showing that HI listeners exhibit higher SRTs than

age-matched NH listeners (Goossens *et al.*, 2017).

The present study explored whether the results of speech intelligibility tests align with the

assumption that an age-related decline in AM frequency selectivity may negatively impact speech

intelligibility, while a sharpening of AM tuning with hearing loss would not be beneficial for speech

intelligibility. SRTs were collected for speech in noise, building upon the previous studies on AM

frequency selectivity (Regev *et al.*, 2023a, 2023b), using a subset of the participants. Young NH, older

NH, and older HI listeners were tested on sentence recognition in various maskers, including speech-

shaped noise, fluctuating noise, and competing talkers. It was hypothesized that the older NH listeners

would exhibit an age-related increase in SRTs, and that the older HI listeners would demonstrate a

further increase in SRTs associated with their hearing loss.

II.   METHODS

A. Listeners and previously published data

Ten young NH listeners, aged 22-28 years (mean age = 24.7 years, eight female), nine older NH

listeners, aged 57–77 years (mean age = 66.9 years, six female), and nine older HI listeners, aged 64-

77 years (mean age = 71.2 years, 2 female), participated in the study. All listeners were native Danish

speakers. The experiment was conducted monaurally. Figure 1 shows the audiograms in the

participants' test ears. Among the young NH listeners, all had audiometric pure-tone thresholds

(PTTs) ≤ 20 dB HL at frequencies ranging from 125 Hz to 8 kHz. In the older NH listeners, all had

PTTs ≤ 20 dB HL at frequencies ranging from 125 Hz to 4 kHz, PTTs ≤ 25 dB HL at 6 kHz, and

PTTs ≤ 45 dB HL at 8 kHz. The older HI listeners exhibited a high-frequency sloping hearing loss,

classified as an N2 or N3 standard audiogram (Bisgaard *et al.*, 2010).[1] All HI listeners had a

sensorineural hearing loss, with air-bone gaps ≤ 15 dB at 1, 2, 3, and 4 kHz.[2]

Prior to this study, all listeners took part in the AM frequency selectivity studies conducted by

Regev *et al.* (2023a, 2023b). In those studies, masked-threshold patterns (MTPs) were used to assess

the detection threshold of a target AM across different masker center frequencies, providing an

indication of AM frequency selectivity. Figure 2 indicates an example of the MTPs for a 4-Hz target

modulation frequency, while Table 1 presents the slopes obtained from linear fits applied to the group-

level MTPs for target modulation frequencies ranging from 4 to 128 Hz, following the method

described in Regev *et al.* (2023b). As reported by Regev *et al.* (2023a, 2023b), the MTPs and

corresponding fitted slopes indicate a general decline in AM frequency selectivity with age, which is

partially compensated by the presence of hearing loss, as reflected by the shallower slopes observed

in the older NH listeners compared to the young NH and older HI listeners.[3]

Regev *et al.* (2023a, 2023b) also evaluated working-memory capacity using a reverse digit span

(RDS) test, which was administered binaurally following the procedure described in Fuglsang *et al.*

(2020). For the HI listeners, a linear gain was applied based on the Cambridge formula (CamEq; Moore

and Glasberg, 1998), with a maximum gain of 30 dB at any frequency. The older NH listeners

exhibited significantly lower average RDS scores (0.40 on a normalized scale) compared to the young

NH and older HI listeners (0.60 and 0.52, respectively).[4]

---

[1] The classification of the audiograms was performed using the lowest mean squared error between the standard audiograms and the collected audiometric thresholds at frequencies from 1 to 8 kHz.

[2] Except for one listener who showed a gap of 20 dB at 1 kHz.

[3] The similar fitted slopes obtained with the older NH and older HI listeners for a 128-Hz target modulation frequency reflect the overall lower masked thresholds obtained from the older HI listeners (see Regev et al., 2023b for further details).

[4] This was shown by the post-hoc analysis run on a linear model ANOVA using estimated marginal means. The older NH listeners had significantly lower RDS scores than the young NH and older HI listeners [$M = 0.21$, $SE = 0.05$ dB; $M = 0.13$, $SE = 0.05$ dB, respectively].

All participants provided written informed consent and were offered financial compensation for

their participation. The study received ethical approval from the Science Ethics Committee of the

Capital Region of Denmark (reference H-16036391).

Figure 1: Individual and average audiograms of the test ear for each listener group. The light lines

show the individual audiograms, while the dark lines represent the average and standard deviation

across listeners. Panel A shows the audiograms for the young NH listeners, with the black horizontal

line indicating the 20 dB HL normal-hearing criterion. Panel B displays the audiograms for the older

NH listeners, also with the black horizontal line denoting the 20 dB HL normal-hearing criterion.

Panel C presents the audiograms for the older HI listeners.

Figure 2: Average MTPs and AM detection thresholds for a 4-Hz target modulation frequency. The
results for the young NH group are presented as dashed lines and squares, the older NH group as
dashed lines and triangles, and the older HI group as continuous lines and circles. The bottom open
symbols represent the average AM detection threshold for each group. The means and standard errors
are shown. Data adapted from Regev *et al.* (2023a) and (2023b).

Table 1: Slopes obtained from the asymmetric linear fitting approach applied to the low- and high-
frequency skirts of the group-level MTPs, following the method described in Regev *et al.* (2023b). The
slopes are presented for the young NH (YNH), older NH (ONH), and older HI (OHI) groups at
target modulation frequencies of 4, 16, 64, and 128 Hz. The slope values are reported in dB/octave.

Target modulation frequency [Hz]	Fitted slope [dB/octave]					
	Low-frequency skirt			*High-frequency skirt*		
	YNH	*ONH*	*OHI*	*YNH*	*ONH*	*OHI*
4	3.7	1.5	3.2	-3.4	-1.0	-0.8
16	3.8	3.3	6.0	-4.3	-3.2	-4.8
64	4.4	2.5	3.7	-6.2	-6.3	-9.1
128	4.2	1.7	1.7	-	-	-

B. Apparatus

The listening test was conducted in a single-walled soundproof booth. The experiment was

conducted using MATLAB (The Mathworks Inc., 2015). The stimuli were digitally played using

SoundMexPro (HörTech, Oldenburg, Germany), converted to analog using an RME Fireface

soundcard (Haimhausen, Germany), and played monaurally through Sennheiser HDA200 headphones

(Wedemark, Germany). The output level was calibrated using a GRAS RA0039 ear simulator (Holte,

Denmark) and the frequency response of the headphones was compensated using an inverse digital

filter. The listeners received no feedback during the test.

C. Procedure and stimuli

Speech intelligibility in noise was assessed using the Danish Hearing in Noise Test (HINT; Nielsen

and Dau, 2011). The HINT speech material consists of 10 test lists and 3 training lists, each containing

20 five-word sentences. The sentences were spoken by a male speaker with an estimated average

fundamental frequency (F_0) of 123 Hz. The speech signals were combined with five different masking

noises: i) a stationary speech-shaped noise (SSN); ii) a fluctuating ICRA-5 noise (Dreschler *et al.*, 2001)

resembling a single male talker in terms of spectral and temporal characteristics; iii) a single male

competing talker reading a news piece (F_0 = 118 Hz); iv) a single female competing talker reading a

news piece (F_0 = 173 Hz); and v) a 'cocktail party' condition. The recordings of the competing talkers

had previously been edited to reduce long pauses to 65 ms. The cocktail-party noise was generated by

combining a relatively dry recording from a cafeteria with the recordings of the male and female

competing talkers mentioned earlier. The levels of the competing talkers were each 3 dB higher than

that of the cafeteria noise. Prior to mixing, each recording was convolved with an impulse response

from the Aachen Impulse Response database (office setting with a dummy head at a 1m distance from

the loudspeaker; Jeub et al., 2009) to introduce reverberation to the original recordings (T_{60} = 0.37 s).

The spectra of all recordings (prior to mixing the cocktail-party noise) had been pre-processed to

match the long-term average spectrum of the HINT speech material as closely as possible. The long-

term average spectra and modulation spectra of the Danish HINT speech material and of all the

maskers are presented in Figure 3.

The SRT for each masking noise was determined by conducting two separate measurements using

different lists. The final SRT was obtained by averaging the SRTs obtained from the two

measurements. The order of masker presentation and the lists used for each masker were randomized

across the listeners. Both repetitions for each masker were always conducted consecutively.[5] The SRTs

were estimated at the 50%-correct point using an adaptive 1-up 1-down sentence-based scoring rule.

A response was considered correct if all five words of the sentence were repeated accurately. The step

size in SNR was set at 4 dB for the first four sentences and reduced to 2 dB afterwards. To refine the

tracking procedure, a special rule was implemented for the 5[th] sentence: The change in SNR was

computed as the average of the preceding four SNRs and the predicted following SNR, minus the

immediately preceding SNR. The first sentence was repeated until the listener could accurately repeat

the entire sentence. If the sentence had to be repeated more than four times, the run was terminated,

and the starting SNR was increased by 2 dB before starting again. The SRT of a run was calculated as

the average SNR across sentences 5 to 20 plus the SNR required for a fictitious 21[st] sentence,

determined by scoring the 20[th] sentence (Soli and Wong, 2008). The scoring of the repeated words

was performed by a trained native Danish speaker.

The speech was played at a fixed level of 60 dB SPL. The masking noise was looped over the

recording, played continuously, and adjusted in level to achieve the desired SNR. For the HI listeners,

linear amplification was added to both the speech and masking noise according to the Cambridge

---

[5] Due to a mistake in the audibility compensation, one of the older HI listeners was tested twice on the entire HINT
corpus. However, more than 7 weeks separated the two testing sessions. Nielsen and Dau (2011) reported an average
improvement in SRT of only 0.5 dB across two sessions separated by three weeks. Hence, the current case was likely not
affected by memorization of the speech material.

formula (CamEq; Moore and Glasberg, 1998), with a maximum gain of 30 dB at any frequency. Each

participant underwent training using all three training lists with the SSN. The starting SNR for each

masker and each run was adjusted to be 6 dB lower than the expected SRTs. The expected SRTs were

determined based on a pilot test conducted with young NH listeners and were defined relative to the

SRT measured from the final training run. Therefore, the starting SNR was 6 dB lower than the SRT

of the listener's last training list for the SSN, male competing talker, and cocktail party conditions; 9

dB lower for the ICRA noise; and 15 dB lower for the female competing talker condition.

Figure 3: Long-term average spectra (top row) and modulation spectra (bottom row) of the speech

material and maskers. Each panel shows the spectra for the HINT speech material (black line) and

one of the maskers (gray line). Panels A and F: SSN; Panels B and G: ICRA-5; Panels C and H: Male

competing talker; Panels D and I: Female competing talker; Panels E and J: Cocktail-party condition.

D. Statistical analysis

All statistical analyses were conducted using the SRTs averaged across both repetitions for each

listener and masker. This was assumed to represent performance, given that the degree of variability

between the two SRT estimates across maskers (assessed by calculating the standard deviation across

the absolute differences in SRTs, $\sigma_{\Delta|SRT|}$) was considered sufficiently small for all listener groups.

Notably, although small, the variability across all maskers was highest for the young NH listeners

($\sigma_{\Delta|SRT|}$ = 2.43 dB) compared to the older NH listeners ($\sigma_{\Delta|SRT|}$= 1.76 dB) and older HI listeners

($\sigma_{\Delta|SRT|}$= 1.49 dB). Analyses of variances (ANOVAs) of linear mixed-effects model fittings were

performed to examine the differences between the listener groups. Two sets of analysis were

conducted: First, the older NH group was compared to the young NH listeners to isolate the effects

of age from hearing loss, and second, the older HI group was compared to the older NH listeners to

isolate the effects of hearing loss from age-related effects. In both analyses, the listeners were treated

as a random effect, while group classification and masker were considered as fixed effects.[6] Post-hoc

analyses were carried out using estimated marginal means, and a Holm-Bonferroni correction criterion

was applied to account for multiple comparisons. The significance level for all analyses was set at 5%.

---

[6] Levene's test was used to assess the homoskedasticity of the residuals and Shapiro-Wilk's test was used to assess the normality of residual's distribution. Shapiro-Wilk's test was not passed for either model, likely due to the relatively small group sizes and to the presence of outliers in the older HI data. However, no other aspect of the data justified the removal of the outliers, and they were therefore kept in the dataset. Furthermore, the average fittings of the models to the experimental data were found to be accurate, and the models were hence considered as appropriate.

III.   RESULTS

A. Speech intelligibility

Figure 4: Boxplots displaying the SRTs for all listeners. Each panel corresponds to a specific masker:
Panel A: Speech-shaped noise (SSN), Panel B: ICRA-5 fluctuating noise, Panel C: Single male
competing talker (same sex), Panel D: Single female competing talker (different sex), Panel E: Cocktail-
party noise. The SRTs for the young NH listeners are represented by squares, older NH by triangles,
and older HI by circles. The stars indicate the level of statistical significance of the differences between
the groups separated by the horizontal black lines (*$p < 0.05$, **$p < 0.01$, ***$p < 0.001$), while "n.s."
denotes non-significant differences. Note that the young NH and older HI groups were not compared
in the statistical analysis.

The SRTs for all listeners and groups are shown in Figure 4 for each masker. The young NH
listeners are represented by squares, older NH listeners by triangles, and older HI listeners by circles.

The statistical analysis comparing young and older NH listeners, focusing on the effects of age,

revealed significant main effects of masker [$F_{4,68} = 274.2, p < 0.001$] and listener group [$F_{1,17} = 17.56$,

$p < 0.001$], as well as a significant interaction between both factors [$F_{4,68} = 9.284, p < 0.001$]. In the

analysis comparing older NH and older HI listeners, examining the effects of hearing loss, significant

main effects of masker [$F_{4,64} = 141.6, p < 0.001$] and listener group [$F_{1,16} = 12.86, p = 0.002$], as well

as a significant interaction between both factors [$F_{4,64} = 5.321, p < 0.001$], were observed. Post-hoc

analyses revealed that the older NH listeners had significantly higher SRTs than the young NH

listeners for the ICRA-5 condition [Effect size (M) = 1.39 dB, standard error (SE) = 0.68 dB], male

competing talker [$M = 4.53$ dB, $SE = 0.68$ dB], female competing talker [$M = 2.70$ dB, $SE = 0.68$

dB], and cocktail-party condition [$M = 1.53$ dB, $SE = 0.68$ dB]. Additionally, the older NH listeners

had significantly lower SRTs than the older HI listeners for the SSN condition [$M = 2.36$ dB, $SE =$

1.14 dB], ICRA-5 condition [$M = 4.30$ dB, $SE = 1.14$ dB], male competing talker [$M = 3.60$ dB, SE

$= 1.14$ dB], and female competing talker [$M = 5.52$ dB, $SE = 1.14$ dB]. The increase in SRT for the

older HI listeners in the cocktail-party condition approached, but did not reach, significance [$M =$

2.24 dB, $SE = 1.14$ dB, $p = 0.06$]. Furthermore, the post-hoc analyses revealed that the female

competing talker resulted in significantly lower SRTs than all other maskers for all listener groups.

The male competing talker yielded significantly higher SRTs than all other maskers for the older NH

and older HI listeners (except for the cocktail-party condition in the older NH group), while the

cocktail-party condition yielded the highest SRTs for the young NH listeners. Finally, the ICRA-5

noise resulted in significantly lower SRTs than the SSN for the young and older NH listeners, but not

for the older HI listeners. Additional details and results from all post-hoc analyses can be found in

Appendix A.

IV. DISCUSSION

The aim of the present study was to investigate whether the findings from speech intelligibility tests would be consistent with the AM frequency selectivity data collected in previous studies by Regev *et al.* (2023a, 2023b). These previous investigations demonstrated an age-related decline in AM frequency selectivity and a sharpening of AM tuning with hearing loss (see Figure 2 and Table 1). A subset of the listener cohort measured by Regev *et al.* (2023a, 2023b) was considered here. It was expected that the age-related reduction in AM frequency selectivity would correspond to poorer speech intelligibility in older compared to young NH listeners. Conversely, the sharper AM tuning observed in older HI listeners was not anticipated to result in better speech intelligibility compared to the older NH listeners. The results revealed an overall decline in speech intelligibility with age, which was further exacerbated by the presence of hearing loss. These findings are consistent with previous studies that have shown a deterioration of speech intelligibility with age (e.g., Rajan and Cainer, 2008; Helfer and Vargo, 2009; Schoof and Rosen, 2014; Füllgrabe *et al.*, 2015; Gordon-Salant and Cole, 2016; Goossens *et al.*, 2017; Decruy *et al.*, 2019) and hearing loss (e.g., Glasberg and Moore, 1989; Humes and Christopherson, 1991; Frisina and Frisina, 1997; see Humes and Dubno, 2010 for a review).

The observed increase in SRTs with age aligns with the age-related reduction in AM frequency selectivity reported by Regev *et al.* (2023a). Broader modulation filters, indicative of reduced AM selectivity, may lead to increased AM masking, resulting in a decrease in the SNR_{env} at the output of these filters and potentially poorer speech intelligibility in noise. However, the lack of a significant age-related increase in SRT for the SSN condition is not consistent with this notion, contrary to expectations. Since the SSN contains intrinsic envelope fluctuations that act as a relatively broadband modulation masker (as argued by Stone *et al.*, 2012), one might have expected the broader AM tuning found in older NH listeners to lead to an increase in AM masking. However, the SRTs obtained with

the SSN likely reflect aspects of EM rather than AM masking. This is suggested by the extensive

overlap observed in the audio spectra of the SSN and HINT speech material (Figure 3, panel A),

compared to the highly positive SNR_{env} observed when comparing their modulation spectra at low

modulation frequencies (Figure 3, panel F), indicating that the SSN provided more EM than AM

masking. Conversely, the differences in modulation spectra between the target speech and masking

noise were much smaller for the other maskers, suggesting that AM masking (and thus potentially AM

frequency selectivity) may have played a more prominent role in those cases (Figure 3, panels G-J).

Notably, the older NH listeners exhibited a reduced release from masking when exposed to the

fluctuating ICRA-5 noise compared to the young NH listeners, consistent with previous studies

(Dubno *et al.*, 2002, 2003; Grose *et al.*, 2009; Goossens *et al.*, 2017). In addition to potentially being the

result of an age-related increase in forward masking (e.g., Gifford *et al.*, 2007), the reduced release from

masking observed in the older NH listeners may also be attributed to increased AM masking due to

the loss of AM frequency selectivity, which may have limited the benefit derived from the slow

temporal fluctuations in the ICRA-5 condition.

Regarding the effects of hearing loss, the higher SRTs observed in the older HI listeners compared

to the older NH group are consistent with the expectation that the increase in AM frequency selectivity

observed in HI listeners (as reported in Regev *et al.*, 2023b) would *not* lead to improved speech

intelligibility. Regev *et al.* (2023b) argued that the greater AM frequency selectivity with hearing loss

was an "apparent" effect, resulting from improved detection of the AM target during temporal dips

in the masker due to the loss of cochlear compression. Since the loss of compression is also associated

with other supra-threshold deficits, such as loudness recruitment, poorer recovery from forward

masking, and reduced frequency selectivity (Glasberg and Moore, 1986, 1989; Oxenham and Moore,

1997; Moore and Oxenham, 1998), it is evident that the process potentially underlying the improved

AM frequency selectivity is also linked to the mechanisms underlying poorer speech intelligibility in

HI listeners. However, while poorer speech intelligibility was expected in the HI listeners,
disentangling the potential impact of the effective sharpening in AM tuning with hearing loss on
speech intelligibility is not straightforward. It remains unclear whether a) the sharper AM tuning with
hearing loss improves SRTs but is outweighed by co-occurring supra-threshold deficits typically
associated with hearing loss, or b) the process underlying the sharpening of AM tuning with hearing
loss itself ultimately impairs speech intelligibility.

The present findings indicate that declines in speech intelligibility with age and hearing loss are
most prominent in contexts that involve high cognitive demand and informational masking (IM), such
as in the presence of competing-talker maskers. IM encompasses various factors that complicate the
separation of the target and masker speech signals, such as stream segregation (Brungart et al., 2001;
Darwin et al., 2003; Schneider et al., 2007; Kidd et al., 2016; see Kidd et al., 2008 for a review). In the
groups considered here, the older NH listeners exhibited significantly lower RDS scores compared to
the young NH listeners. Such reduced cognitive abilities have been suggested to mediate the age-
related increase of SRTs under conditions of IM (Rajan and Cainer, 2008; Gordon-Salant and Cole,
2016; Goossens *et al.*, 2017). Therefore, the poorer speech intelligibility observed in the older NH
listeners in the present study may be partly attributed to a reduction in working-memory capacity.
However, the older NH listeners were also found to have higher SRTs than the young NH listeners
in the ICRA-5 condition, which does not involve IM or high cognitive demand. This finding is broadly
consistent with studies that reported no significant correlation between reduced cognitive abilities and
increased SRTs in older listeners (e.g., Schoof and Rosen, 2014), suggesting that other factors may
also contribute to the age-related decline in speech intelligibility. These factors may be associated with
age-related temporal processing deficits, as suggested by perceptual (Pichora-Fuller and Souza, 2003;
Pichora-Fuller and MacDonald, 2008; Helfer and Vargo, 2009; Füllgrabe et al., 2015; see Fitzgibbons
and Gordon-Salant, 2010 for a review) and neurophysiological research (e.g., Anderson et al., 2012;

Decruy et al., 2019; Parthasarathy et al., 2019b; Roque et al., 2019; for reviews see Parthasarathy et al.,

2019a and Anderson and Karawani, 2020). The hypothesis that the age-related reduction in AM

frequency selectivity (as reported by Regev *et al.*, 2023a) may contribute to increased SRTs aligns with

the notion that temporal processing deficits underlie the challenges in speech intelligibility associated

with age. In the case of hearing loss, the differences in SRTs between older NH and older HI listeners

were also largest for the fluctuating and competing-talker maskers. This observation is consistent with

the reduced release from masking commonly found in HI listeners (e.g., Bacon *et al.*, 1998; Bernstein

and Grant, 2009; Strelcyk and Dau, 2009; Goossens *et al.*, 2017), as well as with the suggestion that

the effects of hearing loss on speech intelligibility are most pronounced in cognitively demanding

settings (e.g., Schneider *et al.*, 2010; Goossens *et al.*, 2017). Although the older HI listeners showed

higher RDS scores than their older NH counterparts, they did exhibit a marginally reduced working-

memory capacity compared to the young NH listeners. Hence, the large effect of hearing loss in

conditions involving IM may be partly connected to the age-related cognitive decline discussed earlier.

Conversely, the older HI listeners were also found to have higher SRTs than their older NH

counterparts in the SSN and ICRA-5 conditions, which are commonly assumed not to involve high

cognitive demand. Overall, while reduced cognitive ability in older listeners may have partly mediated

the observed effects of age and hearing loss on speech intelligibility, the results also indicate that supra-

threshold deficits influenced the SRTs, likely related to reduced cochlear compression and temporal

envelope processing deficits.

Although the data from the present study revealed significant group-level results that support a

relationship between AM tuning and speech intelligibility in noise with age, as well as the absence of

a benefit in speech intelligibility from the sharper AM tuning with hearing loss, it is important to

consider several co-occurring factors that may have influenced the results. Even though all the older

NH listeners exhibited clinically normal hearing up to 4 kHz, their audiometric thresholds were on

average 8 dB higher than those of the young NH listeners between 125 Hz and 8 kHz, with the greatest

differences occurring at 8 kHz. Although these differences in audiometric thresholds are small,

previous studies have suggested that they can have a considerable effect on speech intelligibility

(Dubno and Ahlstrom, 1997; Dubno *et al.*, 2000). Moreover, the older NH listeners included in the

present study may have experienced reduced audibility at extended high frequencies (> 8 kHz), which

have also been suggested to play a role in speech intelligibility (e.g., Hunter *et al.*, 2020; Polspoel *et al.*,

2022). These aspects may have contributed to the observed age-related increase in SRTs. Concerning

the experimental approach, since the tests provided listeners with cues beyond the temporal envelope,

factors such as IM, EM, and stream segregation likely interacted with age- and hearing loss-related

perception deficits. Therefore, establishing an unambiguous connection between SRTs and AM

frequency selectivity based solely on the present data is not possible.

V.   SUMMARY AND CONCLUSION

Speech intelligibility in noise was examined in three groups: young listeners with normal hearing,

aged 22-28 years, older listeners with normal hearing, aged 57-77 years, and older listeners with hearing

impairment, aged 64 – 77 years. All participants had previously taken part in AM masking studies

investigating AM frequency selectivity, which revealed an age-related reduction in AM tuning and a

sharpening in AM tuning with hearing loss (Regev *et al.*, 2023a, 2023b). The aim of this study was to

determine if the effects of age and hearing loss on speech intelligibility in noise support the hypothesis

that the age-related reduction in AM tuning negatively impacts speech intelligibility, while the

improvement in AM tuning with hearing loss does not provide a benefit. Consistent with the

hypotheses, the results showed an age-related decline and an additional negative effect of hearing loss

on speech intelligibility. The detrimental effect of hearing loss on speech intelligibility suggests that

the underlying process responsible for the greater AM frequency selectivity observed in listeners with

hearing impairment (Regev *et al.*, 2023b) is connected to the mechanisms that ultimately lead to poorer

speech intelligibility in noise. This finding aligns with the proposition made by Regev *et al.* (2023b)

that the improved AM tuning in listeners with hearing impairment is an "apparent" effect resulting

from reduced peripheral compression, which is known to contribute to other supra-threshold deficits

that impair speech intelligibility. However, it remains unclear to what extent the apparent greater AM

frequency selectivity itself may compensate for, or contribute to, deficits in speech intelligibility. With

regards to the effects of age, the present results provided clear evidence of an age-related decline in

speech intelligibility, which warrants further investigation into its potential association with a decrease

in AM frequency selectivity. Although the age-related decline in AM frequency selectivity is consistent

with the observed detrimental effects of age on speech intelligibility, this study does not establish a

definitive link between these two aspects, as the employed speech intelligibility test is influenced by

various other factors such as informational masking, energetic masking, and stream segregation.

Therefore, a more controlled experimental paradigm that specifically examines envelope-based cues

may be necessary to directly assess the effects of AM tuning on speech intelligibility.

ACKNOWLEDGMENTS

The authors thank Borgný Súsonnudóttir Hansen for her contribution to the data collection. The

authors also thank Christian Stender Simonsen for kindly sharing the experimental framework used

to run the listening test, as well as several of the signal recordings, and Jens Hjortkjær and Jonatan

Märcher-Rørsted for kindly providing their implementations of the reverse digit span test and of the

Cambridge formula (CamEq; Moore and Glasberg, 1998). Finally, the authors thank Raul Sanchez-

Lopez for helpful insights and discussions. This study was supported by the Oticon Centre of

Excellence for Hearing and Speech Sciences (CHeSS).

AUTHOR DECLARATIONS

Conflict of Interest

The authors have no conflict of interest to report.

Ethics Approval

All participants provided written informed consent and were offered financial compensation for

their participation. Ethical approval for the study was provided by the Science Ethics Committee of

the Capital Region in Denmark (reference H-16036391).

DATA AVAILABILITY

The data that support the findings of this study are available from the corresponding author upon

request.

APPENDIX

A. Post-hoc analyses

Table 2 summarizes the results of the post-hoc analyses that compared the SRTs obtained with

young and older NH listeners for each masker.

Table 2: Post-hoc analysis of the differences in SRT between young and older NH listeners for each

masker. The effect sizes (M), standard errors (SE, and corrected p-values are reported. The significant

p-values are highlighted in bold.

Masker	*M [dB]*	*SE [dB]*	*Corrected p-value*
SSN	0.75		0.279
ICRA-5	1.39		**0.047**
Male competing	4.53	0.68	**< 0.001**
Female competing	2.70		**< 0.001**
Cocktail party	1.53		**0.030**

437   Table 3 summarizes the results of the post-hoc analyses that compared the SRTs obtained with

438   older NH and older HI listeners for each masker.

440   Table 3: Post-hoc analysis of the differences in SRT between older NH and older HI listeners for each

441   masker. The effect sizes (M), standard errors (SE, and corrected p-values are reported. The significant

442   p-values are highlighted in bold.

Masker	M [dB]	SE [dB]	Corrected p-value
SSN	2.36		**0.048**
ICRA-5	4.30		**< 0.001**
Male competing	3.60	1.14	**0.004**
Female competing	5.52		**< 0.001**
Cocktail party	2.24		0.060

Table 4 summarizes the results of the post-hoc analyses that compared the SRTs obtained with

the different maskers for each of the three listener groups (young NH, older NH, and older HI). Since

this post-hoc analysis was run twice for the older NH listeners (once for each statistical model applied),

the values reported reflect the most conservative p-values and standard errors, obtained from the

model fitted to the older NH and older HI listeners. However, the effect sizes and overall statistical

outcomes did not differ between the analyses run in the two models.

Table 4: Post-hoc analysis of the differences in SRT between older the different maskers for each
group. The effect sizes (M), standard errors (SE, and corrected p-values are reported. The significant
p-values are highlighted in bold.

Listener group	Maskers compared	M [dB]	SE [dB]	Corrected p-value
Young NH	SSN - ICRA-5	2.35		**< 0.001**
	SSN – Male competing	0.83		0.086
	SSN – Female competing	9.23		**< 0.001**
	SSN – Cocktail party	-1.42		**0.008**
	ICRA-5 – Male competing	-1.52		**0.006**
	ICRA-5 – Female competing	6.88		**< 0.001**
	ICRA-5 – Cocktail party	-3.77	0.48	**< 0.001**
	Male competing – Female competing	8.40		**< 0.001**
	Male competing – Cocktail party	-2.25		**< 0.001**
	Female competing – Cocktail party	-10.65		**< 0.001**
Older NH	SSN - ICRA-5	1.71		**0.011**
	SSN – Male competing	-2.95		**< 0.001**
	SSN – Female competing	7.28		**< 0.001**
	SSN – Cocktail party	-2.20	0.60	**0.001**
	ICRA-5 – Male competing	-4.66		**< 0.001**
	ICRA-5 – Female competing	5.57		**< 0.001**
	ICRA-5 – Cocktail party	-3.91		**< 0.001**

	Male competing – Female competing	10.23		**< 0.001**
	Male competing – Cocktail party	0.75		0.212
	Female competing – Cocktail party	-9.48		**< 0.001**
Older HI	SSN - ICRA-5	-0.23		0.696
	SSN – Male competing	-4.20		**< 0.001**
	SSN – Female competing	4.12		**< 0.001**
	SSN – Cocktail party	-2.09		**0.003**
	ICRA-5 – Male competing	-3.96		**< 0.001**
	ICRA-5 – Female competing	4.35		**< 0.001**
	ICRA-5 – Cocktail party	-1.85	0.60	**0.006**
	Male competing – Female competing	8.32		**< 0.001**
	Male competing – Cocktail party	2.11		**0.003**
	Female competing – Cocktail party	-6.20		**< 0.001**

REFERENCES

Anderson, S., and Karawani, H. (**2020**). "Objective evidence of temporal processing deficits in

older adults," Hear Res, **397**, 108053. doi:10.1016/j.heares.2020.108053

Anderson, S., Parbery-Clark, A., White-Schwoch, T., and Kraus, N. (**2012**). "Aging Affects

Neural Precision of Speech Encoding," Journal of Neuroscience, **32**, 14156–14164.

doi:10.1523/JNEUROSCI.2176-12.2012

Bacon, S. P., and Grantham, D. W. (**1989**). "Modulation masking: Effects of modulation

frequency, depth, and phase," J Acoust Soc Am, **85**, 2575–2580. doi:10.1121/1.397751

Bacon, S. P., Opie, J. M., and Montoya, D. Y. (**1998**). "The Effects of Hearing Loss and Noise

Masking on the Masking Release for Speech in Temporally Complex Backgrounds,"

Journal of Speech, Language, and Hearing Research, **41**, 549–563.

doi:10.1044/jslhr.4103.549

Bernstein, J. G. W., Danielsson, H., Hällgren, M., Stenfelt, S., Rönnberg, J., and Lunner, T. (**2016**).

"Spectrotemporal Modulation Sensitivity as a Predictor of Speech-Reception

Performance in Noise With Hearing Aids," Trends Hear, **20**, 233121651667038.

doi:10.1177/2331216516670387

Bernstein, J. G. W., and Grant, K. W. (**2009**). "Auditory and auditory-visual intelligibility of

speech in fluctuating maskers for normal-hearing and hearing-impaired listeners," J

Acoust Soc Am, **125**, 3358. doi:10.1121/1.3110132

Bisgaard, N., Vlaming, M. S. M. G., and Dahlquist, M. (**2010**). "Standard Audiograms for the

IEC   60118-15   Measurement   Procedure,"   Trends   Amplif,   **14**,   113–120.

doi:10.1177/1084713810379609

Brungart, D. S., Simpson, B. D., Ericson, M. A., and Scott, K. R. (**2001**). "Informational and

energetic masking effects in the perception of multiple simultaneous talkers," J Acoust

Soc Am, **110**, 2527–2538. doi:10.1121/1.1408946

Chabot-Leclerc, A., Jørgensen, S., and Dau, T. (**2014**). "The role of auditory spectro-temporal

modulation filtering and the decision metric for speech intelligibility prediction," J

Acoust Soc Am, **135**, 3502–3512. doi:10.1121/1.4873517

Darwin, C. J., Brungart, D. S., and Simpson, B. D. (**2003**). "Effects of fundamental frequency

and vocal-tract length changes on attention to one of two simultaneous talkers," J Acoust

Soc Am, **114**, 2913. doi:10.1121/1.1616924

Dau, T., Kollmeier, B., and Kohlrausch, A. (**1997**). "Modeling auditory processing of amplitude

modulation. I. Detection and masking with narrow-band carriers," J Acoust Soc Am, **102**,

2892–2905. doi:10.1121/1.420344

Dau, T., Verhey, J., and Kohlrausch, A. (**1999**). "Intrinsic envelope fluctuations and modulation-

detection thresholds for narrow-band noise carriers," J Acoust Soc Am, **106**, 2752–2760.

doi:10.1121/1.428103

Decruy, L., Vanthornhout, J., and Francart, T. (**2019**). "Evidence for enhanced neural tracking

of the speech envelope underlying age-related speech-in-noise difficulties," J

Neurophysiol, **122**, 601–615. doi:10.1152/jn.00687.2018

Decruy, L., Vanthornhout, J., and Francart, T. (**2020**). "Hearing impairment is associated with

enhanced neural tracking of the speech envelope," Hear Res, **393**, 107961.

doi:10.1016/j.heares.2020.107961

Dorman, M. F., Loizou, P. C., and Rainey, D. (**1997**). "Speech intelligibility as a function of the

number of channels of stimulation for signal processors using sine-wave and noise-band

outputs," J Acoust Soc Am, **102**, 2403–2411. doi:10.1121/1.419603

Dreschler, W. A., Verschuure, H., Ludvigsen, C., and Westermann, S. (**2001**). "ICRA Noises:

Artificial Noise Signals with Speech-like Spectral and Temporal Properties for Hearing

Instrument Assessment," Int J Audiol, **40**, 148–157. doi:10.3109/00206090109073110

Drullman, R., Festen, J. M., and Plomp, R. (**1994**). "Effect of temporal envelope smearing on

speech reception," J Acoust Soc Am, **95**, 1053–1064. doi:10.1121/1.408467

Dubbelboer, F., and Houtgast, T. (**2008**). "The concept of signal-to-noise ratio in the modulation

domain and speech intelligibility," J Acoust Soc Am, **124**, 3937–3946.

doi:10.1121/1.3001713

Dubno, J. R., and Ahlstrom, J. B. (**1997**). "Additivity of Multiple Maskers of Speech," Modeling

Sensorineural Hearing Loss, Routledge, pp. 253–272. doi:10.4324/9781315789392-21

Dubno, J. R., Ahlstrom, J. B., and Horwitz, A. R. (**2000**). "Use of context by young and aged

adults with normal hearing," J Acoust Soc Am, **107**, 538–546. doi:10.1121/1.428322

Dubno, J. R., Dirks, D. D., and Morgan, D. E. (**1984**). "Effects of age and mild hearing loss on

speech recognition in noise," J Acoust Soc Am, **76**, 87–96. doi:10.1121/1.391011

Dubno, J. R., Horwitz, A. R., and Ahlstrom, J. B. (**2002**). "Benefit of modulated maskers for

speech recognition by younger and older adults with normal hearing," J Acoust Soc Am,

111, 2897–2907. doi:10.1121/1.1480421

Dubno, J. R., Horwitz, A. R., and Ahlstrom, J. B. (**2003**). "Recovery from prior stimulation:

Masking of speech by interrupted noise for younger and older adults with normal

hearing," J Acoust Soc Am, **113**, 2084–2094. doi:10.1121/1.1555611

Elliott, T. M., and Theunissen, F. E. (**2009**). "The Modulation Transfer Function for Speech

Intelligibility," (K. J. Friston, Ed.) PLoS Comput Biol, **5**, e1000302.

doi:10.1371/journal.pcbi.1000302

Ewert, S. D., and Dau, T. (**2000**). "Characterizing frequency selectivity for envelope

526   fluctuations," J Acoust Soc Am, **108**, 1181–1196. doi:10.1121/1.1288665

527   Ewert, S. D., Verhey, J. L., and Dau, T. (**2002**). "Spectro-temporal processing in the envelope-

528   frequency domain," J Acoust Soc Am, **112**, 2921–2931. doi:10.1121/1.1515735

529   Fitzgibbons, P. J., and Gordon-Salant, S. (**2010**). "Behavioral Studies With Aging Humans:

530   Hearing Sensitivity and Psychoacoustics," Springer, New York, NY, pp. 111–134.

531   doi:10.1007/978-1-4419-0993-0_5

532   Frisina, D. R., and Frisina, R. D. (**1997**). "Speech recognition in noise and presbycusis: relations

533   to possible neural mechanisms," Hear Res, **106**, 95–104. doi:10.1016/S0378-

534   5955(97)00006-3

535   Füllgrabe, C., Moore, B. C. J., and Stone, M. A. (**2015**). "Age-group differences in speech

536   identification despite matched audiometrically normal hearing: contributions from

537   auditory temporal processing and cognition," Front Aging Neurosci, **6**, 1–25.

538   doi:10.3389/fnagi.2014.00347

539   Gifford, R. H., Bacon, S. P., and Williams, E. J. (**2007**). "An Examination of Speech Recognition

540   in a Modulated Background and of Forward Masking in Younger and Older Listeners,"

541   Journal of Speech, Language, and Hearing Research, **50**, 857–864. doi:10.1044/1092-

542   4388(2007/060)

543   Glasberg, B. R., and Moore, B. C. J. (**1986**). "Auditory filter shapes in subjects with unilateral and

544   bilateral cochlear impairments," J Acoust Soc Am, **79**, 1020–1033. doi:10.1121/1.393374

545   Glasberg, B. R., and Moore, B. C. J. (**1989**). "Psychoacoustic abilities of subjects with unilateral

546   and bilateral cochlear hearing impairments and their relationship to the ability to

547   understand speech," Scand Audiol Suppl, **32**, 1–25.

Goossens, T., Vercammen, C., Wouters, J., and Wieringen, A. van (**2016**). "Aging Affects Neural
Synchronization to Speech-Related Acoustic Modulations," Front Aging Neurosci, **8**, 1–
16. doi:10.3389/fnagi.2016.00133

Goossens, T., Vercammen, C., Wouters, J., and van Wieringen, A. (**2017**). "Masked speech
perception across the adult lifespan: Impact of age and hearing impairment," Hear Res,
344, 109–124. doi:10.1016/j.heares.2016.11.004

Goossens, T., Vercammen, C., Wouters, J., and van Wieringen, A. (**2018**). "Neural envelope
encoding predicts speech perception performance for normal-hearing and hearing-
impaired adults," Hear Res, **370**, 189–200. doi:10.1016/j.heares.2018.07.012

Gordon-Salant, S., and Cole, S. S. (**2016**). "Effects of Age and Working Memory Capacity on
Speech Recognition Performance in Noise Among Listeners With Normal Hearing," Ear
Hear, **37**, 593–602. doi:10.1097/AUD.0000000000000316

Gordon-Salant, S., Yeni-Komshian, G. H., Fitzgibbons, P. J., and Barrett, J. (**2006**). "Age-related
differences in identification and discrimination of temporal cues in speech segments," J
Acoust Soc Am, **119**, 2455–2466. doi:10.1121/1.2171527

Grose, J. H., Mamo, S. K., and Hall, J. W. (**2009**). "Age Effects in Temporal Envelope Processing:
Speech Unmasking and Auditory Steady State Responses," Ear Hear, **30**, 568–575.
doi:10.1097/AUD.0b013e3181ac128f

Helfer, K. S., and Vargo, M. (**2009**). "Speech recognition and temporal processing in middle-
aged women," J Am Acad Audiol, **20**, 264–271. doi:10.3766/jaaa.20.4.6

Houtgast, T. (**1989**). "Frequency Selectivity In Amplitude-Modulation Detection," Journal of the
Acoustical Society of America, **85**, 1676–1680. doi:10.1121/1.397956

Houtgast, T., and Steeneken, H. J. M. (**1985**). "A review of the MTF concept in room acoustics

and its use for estimating speech intelligibility in auditoria," J Acoust Soc Am, **77**, 1069–

1077. doi:10.1121/1.392224

Humes, L. E. (**2002**). "Factors underlying the speech-recognition performance of elderly hearing-

aid wearers," J Acoust Soc Am, **112**, 1112–1132. doi:10.1121/1.1499132

Humes, L. E. (**2007**). "The Contributions of Audibility and Cognitive Factors to the Benefit

Provided by Amplified Speech to Older Adults," J Am Acad Audiol, **18**, 590–603.

doi:10.3766/jaaa.18.7.6

Humes, L. E., and Christopherson, L. (**1991**). "Speech Identification Difficulties of Hearing-

Impaired Elderly Persons," Journal of Speech, Language, and Hearing Research, **34**, 686–

693. doi:10.1044/jshr.3403.686

Humes, L. E., and Dubno, J. R. (**2010**). "Factors Affecting Speech Understanding in Older

Adults," In S. Gordon-Salant, R. Frisina, and A. F. R. Popper (Eds.), The Aging Auditory

System. Springer Handbook of Auditory Research, Springer, New York, Vol. 34, pp.

211–257. doi:10.1007/978-1-4419-0993-0_8

Humes, L. E., Dubno, J. R., Gordon-Salant, S., Lister, J. J., Cacace, A. T., Cruickshanks, K. J.,

Gates, G. A., et al. (**2012**). "Central Presbycusis: A Review and Evaluation of the

Evidence," J Am Acad Audiol, **23**, 635–666. doi:10.3766/jaaa.23.8.5

Humes, L. E., Lee, J. H., and Coughlin, M. P. (**2006**). "Auditory measures of selective and divided

attention in young and older adults using single-talker competition.," J Acoust Soc Am,

120, 2926–37. doi:10.1121/1.2354070

Hunter, L. L., Monson, B. B., Moore, D. R., Dhar, S., Wright, B. A., Munro, K. J., Zadeh, L. M.,

et al. (**2020**). "Extended high frequency hearing and speech perception implications in

adults and children," Hear Res, **397**, 107922. doi:10.1016/j.heares.2020.107922

Jennings, S. G., Chen, J., Fultz, S. E., Ahlstrom, J. B., and Dubno, J. R. (**2018**). "Amplitude

595   modulation detection with a short-duration carrier: Effects of a precursor and hearing

596   loss," J Acoust Soc Am, **143**, 2232–2243. doi:10.1121/1.5031122

597   Jepsen, M. L., and Dau, T. (**2011**). "Characterizing auditory processing and perception in

598   individual listeners with sensorineural hearing loss," J Acoust Soc Am, **129**, 262–281.

599   doi:10.1121/1.3518768

600   Jepsen, M. L., Ewert, S. D., and Dau, T. (**2008**). "A computational model of human auditory

601   signal processing and perception," J Acoust Soc Am, **124**, 422–438.

602   Jeub, M., Schafer, M., and Vary, P. (**2009**). "A binaural room impulse response database for the

603   evaluation of dereverberation algorithms," 2009 16th International Conference on

604   Digital Signal Processing, IEEE, 1–5. doi:10.1109/ICDSP.2009.5201259

605   Jørgensen, S., and Dau, T. (**2011**). "Predicting speech intelligibility based on the signal-to-noise

606   envelope power ratio after modulation-frequency selective processing," J Acoust Soc

607   Am, **130**, 1475–1487. doi:10.1121/1.3621502

608   Jørgensen, S., Ewert, S. D., and Dau, T. (**2013**). "A multi-resolution envelope-power based model

609   for speech intelligibility," J Acoust Soc Am, **134**, 436–446. doi:10.1121/1.4807563

610   Joris, P. X., Schreiner, C. E., and Rees, A. (**2004**). "Neural Processing of Amplitude-Modulated

611   Sounds," Physiol Rev, **84**, 541–577. doi:10.1152/physrev.00029.2003

612   Kidd, G., Mason, C. R., Richards, V. M., Gallun, F. J., and Durlach, N. I. (**2008**). "Informational

613   Masking," pp. 143–189. doi:10.1007/978-0-387-71305-2_6

614   Kidd, G., Mason, C. R., Swaminathan, J., Roverud, E., Clayton, K. K., and Best, V. (**2016**).

615   "Determining the energetic and informational components of speech-on-speech

616   masking," J Acoust Soc Am, **140**, 132–144. doi:10.1121/1.4954748

Kohlrausch, A., Fassel, R., and Dau, T. (**2000**). "The influence of carrier level and frequency on modulation and beat-detection thresholds for sinusoidal carriers," J Acoust Soc Am, **108**, 723–734. doi:10.1121/1.429605

Lorenzi, C., Gilbert, G., Carn, H., Garnier, S., and Moore, B. C. J. (**2006**). "Speech perception problems of the hearing impaired reflect inability to use temporal fine structure," Proceedings of the National Academy of Sciences, **103**, 18866–18869. doi:10.1073/pnas.0607364103

McDermott, J. H., and Simoncelli, E. P. (**2011**). "Sound texture perception via statistics of the auditory periphery: Evidence from sound synthesis," Neuron, **71**, 926–940. doi:10.1016/j.neuron.2011.06.032

McWalter, R., and Dau, T. (**2017**). "Cascaded amplitude modulations in sound texture perception," Front Neurosci, **11**, 1–12. doi:10.3389/fnins.2017.00485

Millman, R. E., Mattys, S. L., Gouws, A. D., and Prendergast, G. (**2017**). "Magnified neural envelope coding predicts deficits in speech perception in noise," Journal of Neuroscience, **37**, 7727–7736. doi:10.1523/JNEUROSCI.2722-16.2017

Moore, B. C. J. (**2003**). "Speech processing for the hearing-impaired: successes, failures, and implications for speech mechanisms," Speech Commun, **41**, 81–91. doi:10.1016/S0167-6393(02)00095-X

Moore, B. C. J., and Glasberg, B. R. (**1998**). "Use of a loudness model for hearing-aid fitting. I. Linear hearing aids," Br J Audiol, **32**, 317–335. doi:10.3109/03005364000000083

Moore, B. C. J., and Oxenham, A. J. (**1998**). "Psychoacoustic consequences of compression in the peripheral auditory system.," Psychol Rev, **105**, 108–124. doi:10.1037/0033-295X.105.1.108

Moore, B. C. J., Wojtczak, M., and Vickers, D. A. (**1996**). "Effect of loudness recruitment on the

perception of amplitude modulation," J Acoust Soc Am, **100**, 481–489.

doi:10.1121/1.415861

Nielsen, J. B., and Dau, T. (**2011**). "The Danish hearing in noise test," Int J Audiol, **50**, 202–208.

doi:10.3109/14992027.2010.524254

Oxenham, A. J., and Moore, B. C. J. (**1997**). "Modeling the Effects of Peripheral Nonlinearity in

Listeners With Normal and Impaired Hearing," Modeling Sensorineural Hearing Loss,

Routledge, New York, Jesteadt, W., pp. 273–288. doi:10.4324/9781315789392-22

Parthasarathy, A., Bartlett, E. L., and Kujawa, S. G. (**2019a**). "Age-related Changes in Neural

Coding of Envelope Cues: Peripheral Declines and Central Compensation,"

Neuroscience, **407**, 21–31. doi:10.1016/j.neuroscience.2018.12.007

Parthasarathy, A., Herrmann, B., and Bartlett, E. L. (**2019b**). "Aging alters envelope

representations of speech-like sounds in the inferior colliculus," Neurobiol Aging, **73**,

30–40. doi:10.1016/j.neurobiolaging.2018.08.023

Pichora-Fuller, M. K., and MacDonald, E. (**2008**). "Auditory temporal processing deficits in older

listeners: From a review to a future view of presbycusis," Proceedings of the International

Symposium on Auditory and Audiological Research, **1**, 291–300.

Pichora-Fuller, M. K., and Souza, P. E. (**2003**). "Effects of aging on auditory processing of

speech," Int J Audiol, **42**, 11–16. doi:10.3109/14992020309074638

Polspoel, S., Kramer, S. E., van Dijk, B., and Smits, C. (**2022**). "The Importance of Extended

High-Frequency Speech Information in the Recognition of Digits, Words, and Sentences

in Quiet and Noise," Ear Hear, **43**, 913–920. doi:10.1097/AUD.0000000000001142

Rajan, R., and Cainer, K. E. (**2008**). "Ageing without hearing loss or cognitive impairment causes

a decrease in speech intelligibility only in informational maskers," Neuroscience, **154**,

784–795. doi:10.1016/j.neuroscience.2008.03.067

Regev, J., Zaar, J., Relaño-Iborra, H., and Dau, T. (**2023a**). "Age-related reduction of amplitude

modulation frequency selectivity," J Acoust Soc Am, **153**, 2298–2298.

doi:10.1121/10.0017835

Regev, J., Relaño-Iborra, H., Zaar, J., and Dau, T. (**2023b**). "Disentangling the effects of hearing

loss and age on amplitude modulation frequency selectivity," BioRxiv.

doi:10.1101/2023.07.15.549131

Relaño-Iborra, H., and Dau, T. (**2022**). "Speech intelligibility prediction based on modulation

frequency-selective processing," Hear Res, **426**, 108610.

doi:10.1016/j.heares.2022.108610

Relaño-Iborra, H., May, T., Zaar, J., Scheidiger, C., and Dau, T. (**2016**). "Predicting speech

intelligibility based on a correlation metric in the envelope power spectrum domain," J

Acoust Soc Am, **140**, 2670–2679. doi:10.1121/1.4964505

Relaño-Iborra, H., Zaar, J., and Dau, T. (**2019**). "A speech-based computational auditory signal

processing and perception model," J Acoust Soc Am, **146**, 3306–3317.

doi:10.1121/1.5129114

Roque, L., Karawani, H., Gordon-Salant, S., and Anderson, S. (**2019**). "Effects of age, cognition,

and neural encoding on the perception of temporal speech cues," Front Neurosci, **13**, 1–

15. doi:10.3389/fnins.2019.00749

Schlittenlacher, J., and Moore, B. C. J. (**2016**). "Discrimination of amplitude-modulation depth

by subjects with normal and impaired hearing," J Acoust Soc Am, **140**, 3487–3495.

doi:10.1121/1.4966117

Schneider, B. A., Li, L., and Daneman, M. (**2007**). "How Competing Speech Interferes with

Speech Comprehension in Everyday Listening Situations," J Am Acad Audiol, **18**, 559–

572. doi:10.3766/jaaa.18.7.4

Schneider, B. A., Pichora-Fuller, K., and Daneman, M. (**2010**). "Effects of Senescent Changes in

Audition and Cognition on Spoken Language Comprehension," Springer, New York,

NY, pp. 167–210. doi:10.1007/978-1-4419-0993-0_7

Schoof, T., and Rosen, S. (**2014**). "The role of auditory and cognitive factors in understanding

speech in noise by normal-hearing older listeners," Front Aging Neurosci, **6**, 1–14.

doi:10.3389/fnagi.2014.00307

Shannon, R. V., Zeng, F.-G., Kamath, V., Wygonski, J., and Ekelid, M. (**1995**). "Speech

Recognition with Primarily Temporal Cues," Science (1979), **270**, 303–304.

doi:10.1126/science.270.5234.303

Singh, N. C., and Theunissen, F. E. (**2003**). "Modulation spectra of natural sounds and

ethological theories of auditory processing," J Acoust Soc Am, **114**, 3394–3411.

doi:10.1121/1.1624067

Smith, Z. M., Delgutte, B., and Oxenham, A. J. (**2002**). "Chimaeric sounds reveal dichotomies in

auditory perception," Nature, **416**, 87–90. doi:10.1038/416087a

Sobon, K. A., Taleb, N. M., Buss, E., Grose, J. H., and Calandruccio, L. (**2019**). "Psychometric

function slope for speech-in-noise and speech-in-speech: Effects of development and

aging," J Acoust Soc Am, **145**, EL284–EL290. doi:10.1121/1.5097377

Soli, S. D., and Wong, L. L. N. (**2008**). "Assessment of speech intelligibility in noise with the

hearing in noise test," Int J Audiol, **47**, 356–361. doi:10.1080/14992020801895136

Steinmetzger, K., Zaar, J., Relaño-Iborra, H., Rosen, S., and Dau, T. (**2019**). "Predicting the

effects of periodicity on the intelligibility of masked speech: An evaluation of different

modelling approaches and their limitations," J Acoust Soc Am, **146**, 2562–2576.

711   doi:10.1121/1.5129050

712 Stone, M. A., Füllgrabe, C., and Moore, B. C. J. (**2012**). "Notionally steady background noise acts

713   primarily as a modulation masker of speech," J Acoust Soc Am, **132**, 317–326.

714   doi:10.1121/1.4725766

715 Stone, M. A., and Moore, B. C. J. (**2014**). "On the near non-existence of 'pure' energetic masking

716   release for speech," J Acoust Soc Am, **135**, 1967–1977. doi:10.1121/1.4868392

717 Strelcyk, O., and Dau, T. (**2009**). "Relations between frequency selectivity, temporal fine-

718   structure processing, and speech reception in impaired hearing," J Acoust Soc Am, **125**,

719   3328. doi:10.1121/1.3097469

720 The Mathworks Inc. (**2015**). "MATLAB 2015b - MathWorks,"

721   www.mathworks.com/products/matlab,.

722 Verhey, J. L., Ewert, S. D., and Dau, T. (**2003**). "Modulation masking produced by complex tone

723   modulators," J Acoust Soc Am, **114**, 2135–2146. doi:10.1121/1.1612489

724 Viemeister, N. F. (**1979**). "Temporal modulation transfer functions based upon modulation

725   thresholds," J Acoust Soc Am, **66**, 1364–1380. doi:10.1121/1.383531

726 Viswanathan, V., Bharadwaj, H. M., Shinn-Cunningham, B. G., and Heinz, M. G. (**2021**).

727   "Modulation masking and fine structure shape neural envelope coding to predict speech

728   intelligibility across diverse listening conditions," J Acoust Soc Am, **150**, 2230–2244.

729   doi:10.1121/10.0006385

730 Xu, L., Thompson, C. S., and Pfingst, B. E. (**2005**). "Relative contributions of spectral and

731   temporal cues for phoneme recognition," J Acoust Soc Am, **117**, 3255–3267.

732   doi:10.1121/1.1886405

733   Zaar, J., Simonsen, L. B., Dau, T., and Laugesen, S. (**2023**). "Toward a clinically viable spectro-

734   temporal modulation test for predicting supra-threshold speech reception in hearing-

735   impaired listeners," Hear Res, **427**, 108650. doi:10.1016/j.heares.2022.108650