Title: No evidence for the association between COVID-19 and neuroinflammation: A diffusion basis spectrum imaging study.

Wei Zhang [a,#], **Aaron J Gorelik** [b], **Qing Wang** [a], **Sara A Norton** [b], **Tamara Hershey** [a,b,c,d], **Arpana Agrawal** [c], **Janine D Bijsterbosch** [a,#,*], **Ryan Bogdan** [b,*]

[a] Mallinckrodt Institute of Radiology, Washington University School of Medicine, St. Louis, MO, United States
[b] Department of Psychological & Brain Sciences, Washington University, St. Louis, MO, United States
[c] Department of Psychiatry, Washington University School of Medicine, St. Louis, MO, United States
[d] Department of Neurology, Washington University School of Medicine, St. Louis, MO, United States

[#] Corresponding authors
[*] Shared senior authorship

Keywords: Neuroinflammation, COVID, Long COVID, UK Biobank, neuroimaging, Diffusion Basis Spectrum Imaging, DBSI.

Abstract

COVID continues to be a major international public health concern, the underlying mechanisms of which are not fully understood. Recent studies suggest that COVID may cause prolonged inflammation within the central nervous system. However, the evidence so far has been limited to few small-scale case studies. To address this, this study leveraged a longitudinal dataset from the UK Biobank that included neuroimaging data prior to and following COVID testing (analytic N=416 including n=224 COVID-positive cases) and applied a novel and non-invasive Diffusion Basis Spectrum Imaging (DBSI) technique to derive putative indices of neuroinflammation (i.e., restricted fraction; DBSI-RF) for gray matter structures and white matter tracts in the brain. We hypothesized that SARS-CoV-2 infection would be associated with elevated DBSI markers of putative neuroinflammation and conducted linear regression analyses with adjustment for age, sex, race, body mass index, smoking frequency, and data acquisition interval. After multiple testing correction using false discovery rate, we found no evidence that COVID is associated with variability in neuroinflammation. Several brain regions showed nominally significant differences in DBSI-RF between COVID cases and controls including psychopathology-related regions linked that are either part of (i.e., orbitofrontal cortex) or functionally connected to the olfactory network (e.g., amygdala, caudate). It remains possible that there are acute and transitory neuroinflammatory effects associated with COVID that were not observed in our study due to potential resolution of COVID prior to the scan. Future research is warranted to examine whether neuroinflammation is associated with SARS-CoV-2 infection in a time- and/or symptom-dependent manner.

Introduction

The ongoing COVID19 ("COVID") pandemic remains a threat to global health and economies. In America alone, at least 42% of adults have been diagnosed with COVID during their lifetime (1), and approximately 20% of them experience the longer-term consequences of COVID (also known as long-COVID; (2), including loss of smell, psychopathology (e.g., sleep disorders, depression, anxiety), tiredness, cognitive impairment (e.g., "brain fog"), and increased mortality (3–6). It is now widely recognized that SARS-CoV-2 infection leads to a wide range of medical conditions from respiratory symptoms to the development of neurological symptomatology (2). COVID also continues to pose a significant global economic burden (7), with the total estimated economic burden of 3.7 trillion USD in 2022 (8). These ongoing challenges necessitate a comprehensive understanding of the underlying mechanisms, through which COVID impacts health and behavior.

Accumulating research suggests that COVID may impact brain structure (6,9–11). For instance, widespread changes in brain structure have been observed within individuals who were scanned prior to and following a COVID diagnosis, with the most profound differences observed in the orbitofrontal cortex and related regions (e.g., piriform cortex) involved in the olfactory network (16). Evidence showing the association between COVID-related changes in brain structure and impaired cognitive performance further suggests that some COVID-related symptomatology may be driven by changes in brain structure (17). Interestingly, these brain-behavior associations may not resolve with time as COVID-related differences in brain structure have been observed one- and two-years following hospitalization (18,19).

COVID may impact brain structure by affecting inflammatory processes in the central nervous system (20,21). SARS-CoV-2 infection leads to activation of inflammatory signaling cascades, which can become dysregulated resulting in excessively high inflammation, known as cytokine storms (22–24). This hyperactivity of the immune system can disrupt the blood brain barrier (25), enabling peripheral inflammatory markers to gain access into the CNS, which may then impact brain structure (26). As such, neuroinflammation (e.g., encephalitis) may occur in COVID patients as early as two weeks after the infection (27,28) and persist in long-COVID patients with neuropsychiatric symptoms (29). Brain olfactory regions may be particularly vulnerable to these hyperactive responses. For example, a study of golden hamsters found that COVID infection may increase microglial and infiltrating macrophage activation in olfactory tissues, which was associated with behavioral alterations in scent-based food finding (30). Further evidence comes from post-mortem data in human COVID patients and a rhesus macaque model of COVID showing inflammation in brain barrier (i.e., choroid plexus; (31)), T-cell infiltration, and microglia activation (32). While there are intriguing *in vivo* data from position emission tomography (PET) case studies linking COVID to elevated neuroinflammation (29,33,34), there have yet to be any large-scale investigations of neuroinflammation in the context of COVID.

Here, we tested whether COVID is associated with changes in putative neuroinflammation (analytic n=416; cases n=244, controls n=192). To this end, we used Diffusion Basis Spectrum Imaging (DBSI) to estimate neuroinflammation within the "COVID-19 repeat imaging" sub-study of the UK Biobank (35). This sub-study collected neuroimaging data from individuals prior to and following a COVID positive or negative test (e.g., based on diagnoses comprising PCR test, hospital impatient admission or GP records, as well as home-based test; see details in source of positive test result below in Methods). Neuroinflammation was indexed by DBSI-Restricted Fraction (DBSI-RF), which provides a putative measure of inflammation-related cellularity (36,37). We hypothesized that COVID would be associated with increases in DBSI-RF across the brain with the most profound differences in brain regions that showed the strongest COVID-related structural changes (e.g., orbitofrontal cortex, piriform

cortex) in humans (16)) or changes in neuroinflammation (e.g., microglia activation in olfactory bulb) in post-mortem or animal studies (30,32,38).

Materials and Methods
Sample.
The UK Biobank (UKB) is a large-scale study (N > 500,000 participants) designed to examine the genetic, environmental, biological, and behavioral correlates of broad-spectrum health outcomes and related phenotypes (39). In February 2021, a UKB sub-study, the 'COVID-19 repeat imaging study,' was launched to collect neuroimaging data at a second timepoint, following either a positive (cases) or negative (controls) COVID test, among individuals who completed a neuroimaging session prior to the COVID-19 pandemic to study longitudinal neuroimaging correlates of SARS-CoV-2 infection. COVID positivity/negativity prior to the second neuroimaging session was determined from 3 sources: 1) hospital records contained in the Hospital Episode Statistics (a database containing admissions, outpatient appointments, and attendances and emergency admissions to English National Health Service Hospitals), or 2) primary care (GP) data, and 3) a record of a positive COVID antibody test obtained from a home-based lateral flow kit sent to participants. For individuals who completed home testing and were vaccinated, a second testing kit was collected to ensure that any antibodies detected were from infection as opposed to recent vaccination (16). Participants were identified as COVID-positive cases if they had a positive test record on any of these data sources. COVID-negative participants (i.e., controls) were then selected in this sub-study from the remaining participants that were previously imaged before the pandemic to achieve 1:1 matching to the COVID-positive cases on sex, race (white/non-white), age (date of birth), location and date of the first imaging assessment. Details of inclusion criteria and case-control matching are provided in online   documentation (https://biobank.ndph.ox.ac.uk/showcase/showcase/docs/casecontrol_covidimaging.pdf).

Data from the 'COVID-19 repeat imaging study' has been released on a rolling basis. As of March 24, 2023, we identified n = 416 (including n = 224 COVID-positive) participants from the matched case-control list (variable ID 41000 in the UKB Data Showcase), who met the following inclusion criteria of the current study: 1) no mismatch between self-reported and genetic sexes, and 2) no missing data in any of the measures used in the current study. Demographic characteristics of the present study sample are summarized in **Table 1** and consistent with study design, were comparable between COVID case and control participants. We also identified COVID testing date for n=219 of the COVID cases (UKB variable ID 40100). For this group of participants, the COVID testing preceded the second imaging session by 128.3±70.8 days on average (range 37-372 days). Unfortunately, COVID-related symptomatology data and vaccination status were unavailable for the current study.

Table 1. Demographics of the Study Sample.

	Group		Between-group Difference [a]
	Covid-positive (n=224 case)	Covid-negative (n=192 control)	
Genetic Sex (Female)	122	112	$X^2 = 0.63$ (p=0.43)
Race (White)	219	188	$X^2 = 0.01$ (p=0.92)
Age [b] (M/SD Year)	59.1/7.03	58/7.13	t = 1.0 (p=0.32)
Δ Age [c] (M/SD Year)	3.03/1.59	3.11/1.62	t = -0.52 (p=0.60)
Δ BMI [c] (M/SD)	-0.08/2.12	0.17/1.96	t = -1.2 (p=0.23)
Δ Smoking Status (M/SD Frequency)	-0.05/0.24	-0.02/0.18	t = -1.07 (p0.29)
Longitudinal Assessment Interval [d] (M/SD Days)	1114/570	1157/577	t = -0.76 (p=0.45)
SARS-CoV-2 infection **Time** [d] (M/SD Days)	128.3/70.2	NA	NA

[a] Chi-squared and two-sample t-tests were run to test between-group differences in categorical and numeric measures, respectively.
[b] Age at the first imaging scan prior to COVID
[c] Δ, Delta = differential measures between pre- and post-Pandemic assessments.
[d] Longitudinal assessment interval refers to the time difference between the acquisition of two imaging scans, whereas SARS-CoV-2 infection time refers to the time difference between the Covid testing date and the second scan for Covid-positive participants.

Imaging Acquisition and Processing.
Diffusion Weighted Imaging (DWI) and T1-weighted structural MRI scans were acquired in the present study and processed using a pipeline developed and executed by the UKB Brain Imaging group (50). Briefly, DWI data were acquired using a multi-shell approach (b_1 = 1000 and b_2 = 2000 s/mm2) with 50 distinct diffusion-encoding directions and then preprocessed to correct for eddy currents, head motion, outlier slice, and gradient distortion. The preprocessed data are available for download from the UKB database. T1 structural MRI data were acquired using an in-plane acceleration sequence and preprocessed to remove the skull and corrected for gradient distortion. Further processing on the T1 images was then carried out using FreeSurfer software, which produced images, surface files, and summary outputs, all available for download from the UKB database (https://biobank.ndph.ox.ac.uk/showcase/). More information about the acquisition protocols, image processing pipeline, and derived imaging measures can be found in the UK Biobank Imaging Documentation (https://biobank.ctsu.ox.ac.uk/crystal/crystal/docs/brain_mri.pdf) and studies by Alfaro-Almagro et al. (48) and Miller et al. (35) .

Diffusion Basis Spectrum Imaging and A Putative Marker of Neuroinflammation.
Diffusion basis spectrum imaging (DBSI) is an extension of standard Diffusion Tensor Imaging (DTI). In contrast to DTI that only estimates direction-dependent parameters of water movement (i.e., using anisotropic tensors), DBSI describes the diffusion-weighted imaging signal with multiple anisotropic and isotropic tensors, which improves specificity and sensitivity of diffusion properties (36,41,49). More importantly, the DBSI approach provides a spectrum of isotropic diffusion metrics, including restricted fraction (DBSI-RF) that has been shown to indicate inflammation-related cellularity (i.e., either from cell proliferation or infiltration), with higher values indicating higher levels of inflammatory cell fraction. DBSI-RF as a putative neuroinflammation marker has been validated in a series of experiments showing that DBSI-RF is associated with inflammation-related cellularity change as examined by immunohistochemistry in an experimental mouse model of induced autoimmune encephalomyelitis (41), and with stain-quantitated nuclei and microglia density (i.e., cellularity) from post-mortem human brain tissues (40). Higher levels of DBSI-RF have also been linked to inflammation-related conditions, including multiple sclerosis (37,49–51), obesity (45,46), HIV (52), Alzheimer's disease (40,43,45), and depression (49), as welle as markers of disease progression (40,53).

To indicate neuroinflammation levels for specific brain structures, we applied the DBSI analysis package that was developed in house using the MATLAB (36) to the DWI data and used pre-defined brain structures (i.e., regions of interest, ROIs) as masks to extract region-specific neuroinflammation indices. Specifically, we used gray matter parcellations generated by the Automatic subcortical segmentation (54) and Desikan-Killiany cortical atlases (55). The resulting gray matter ROIs consisted of 14 bilateral subcortical and 66 cortical parcellations, representing n = 7 and n = 33 subcortical and cortical structures, respectively (see full list in Table S1). Additionally, n = 20 white matter tracts from both left and right hemispheres were extracted from a probabilistic tractography atlas (JHU-ICBM-tracts) with the highest probability of 25% at a given brain voxel (56). We used this probability threshold to ensure that each individual white matter tract could be identified in the subject-specific diffusion images (i.e., increasing probability would result in uneven tract numbers identified at the individual level).

To extract one neuroinflammation index per individual for each of the gray and white matter ROIs, we first created individual-specific ROI masks by registering T1 structural (i.e., FreeSurfer outputs) and MNI standard diffusion images (i.e., white matter tracts) with native diffusion images. The mean DBSI-RF values across all voxels within each ROI were then calculated for each individual participant. As more than half of the entire gray matter structures (n = 21) showed a correlation less than 0.6 between the left and right hemispheric homologues for the baseline imaging scan (i.e., pre-COVID), we considered each parcellation as an independent ROI for statistical analysis (i.e., not combined across hemispheres). Similarly, all 20 white matter tracts were considered as separate ROI as the left-right correlations for all tracts from the first imaging scan were below 0.6.

Covariates
Following a prior study (16), we included genetic sex (UKB variable ID 22001), ethnicity (UKB variable ID 21000), as well as differences between pre- and post-COVID assessments in age (UKB variable ID 21003) as covariates in this study. We further included changes in body mass index (BMI; UKB variable ID 21001), in smoking status (UKB variable ID 20116), and in date (i.e., number of days; UKB variable ID 53) between two assessments to adjust for potential confounds that may contribute to the changes in neuroinflammation from pre- to post-COVID. As in this prior study (16), we also used white versus non-white for ethnicity in all models.

Statistical Analysis
We tested whether COVID cases differed from controls on neuroinflammation, as indexed by DBSI-RF. Mirroring the analytic strategy of a prior study linking COVID case status to brain structural changes (16), we conducted a series of linear regressions in which COVID case/control status was modeled as a predictor of post-COVID regional neuroinflammation, while accounting for pre-COVID neuroinflammation and the covariates described above. Separate models were conducted for each individual ROI and false discovery rate (FDR) was applied to adjust for multiple testing within gray and white matter ROIs, separately (i.e., 40 tests for gray matter ROIs, and 20 tests for white matter ROIs).

As it is plausible that neuroinflammation may reflect a transient phenomenon resolving over time, we further conducted post-hoc analyses on data of participants whose positive COVID test occurred ≤ 60 (n = 75) and ≤90 (n = 23) days prior to the neuroimaging session and conducted group comparisons of each of these two subgroups with controls separately.

As changes in putative neuroinflammation within each ROI may not occur uniformly, we further explored the association between COVID and whole-brain voxel-wise neuroinflammation, as indexed by DBSI-RF. To this end, we first registered all participants' DBSI-RF maps (i.e., both pre- and post-COVID maps) with the standard MNI brain template and obtained a differential map between two scans per individual (i.e., scan 2 minus scan 1; delta DBSI-map in MNI space). Using these delta maps as input, we then conducted permutation tests (permuted n = 5000) with FSL *randomise (57)* and a Threshold-Free Cluster Enhancement (TFCE) method (58) to identify delta DBSI-RF "clusters" in the brain that differ between COVID-negative and positive individuals (i.e., case-minus-control contrast and control-minus-case contrast), while accounting for covariates (i.e., sex, ethnicity, data acquisition interval, and changes in age, BMI, and smoking status). Categorical variables (e.g., sex, group) were dummy coded before permutation testing.

Results

   After multiple testing correction, COVID infection was not associated with any change in DBSI-RF values across all regional gray matter regions and white matter tracts **(Table S1)**. There were only a handful of nominally significant associations (i.e., uncorrected p<0.05; **Figure 1; Table 2**). Gray matter regions included the left caudal middle frontal gyrus (β = -0.14, $p_{uncorrected}$ = 0.048), superior parietal lobule (β = -0.14, $p_{uncorrected}$ = 0.047), and postcentral gyrus (β = -0.14, $p_{uncorrected}$ = 0.041), as well as the right caudate (β = 0.18, $p_{uncorrected}$ = 0.039), amygdala (β = 0.22, $p_{uncorrected}$ = 0.013), caudal anterior cingulate cortex (β=0.25, $p_{uncorrected}$ = 0.003), rostral anterior cingulate cortex (β = 0.27, $p_{uncorrected}$ = 0.003), lateral orbitofrontal cortex (β=0.19, $p_{uncorrected}$ = 0.037), and fusiform gyrus (β = 0.19, $p_{uncorrected}$ = 0.029). Only one region was nominally significant in white matter: the temporal proportion of the left superior longitudinal fasciculus (β = 0.25, $p_{uncorrected}$ = 0.008).

Figure 1. Gray Matter Regions with Nominal Increase in Neuroinflammation.

Table 2. Nominally significant between-group differences in neuroinflammation.

Brain Structure	ROI (L/R) [a]	Statistics [b]			
		B [95% CI] [c]	*t*	Uncorrected P-value	FDR corrected P-value [d]
Subcortical					
	Caudate (R)	0.18 [0.01, 0.35]	2.08	0.039	0.424
	Amygdala (R)	0.22 [0.04, 0.41]	2.49	0.013	0.356
Cortical					
	Caudal Middle Frontal (L)	-0.14 [-0.28, 0.001]	-1.99	0.048	0.424
	Postcentral (L)	-0.14 [-0.27, 0.003]	-2.05	0.041	0.424
	Superior Parietal (L)	-0.13 [-.027, 0.001]	-1.99	0.047	0.424
	Caudal Anterior Cingulate (R)	0.25	2.54	0.011	0.356

		[0.05, 0.44]			
	Fusiform (R)	0.19 [0.02, 0.36]	2.20	0.029	0.424
	Lateral Orbitofrontal (R)	0.19 [0.01, 0.38]	2.09	0.037	0.424
	Rostral Anterior Cingulate (R)	0.27 [0.09, 0.46]	2.94	0.003	0.279
White Matter Tracts					
	Superior Longitudinal Fasciculus (L; Temporal Part)	0.25 [0.06, 0.44]	2.66	0.008	0.165

[a] ROI = Regions of Interest; L/R = Left / Right hemisphere
[b] Group difference between the COVID-positive and negative participants after adjusting for the covariates.
[c] CI = Confidence Interval
[d] FDR = False Discovery Rate

Proximity of Scan to COVID Diagnosis.
Participants who had COVID positive tests that were closer in time to the neuroimaging session (i.e., ≤ 60 or ≤ 90 days following COVID test) did not differ in DBSI-RF in any regions or tracts under investigation after accounting for multiple testing. Yet, nominally higher DBSI-RF values were observed in several brain structures and tracts, including rostral anterior cingulate cortex in the right hemisphere (β's > 0.3, uncorrected p's < 0.05), which were also observed in the full sample (**Table 3**).

Whole-brain voxel-wise differences in putative neuroinflammation.
After applying threshold-free cluster enhancement (TFCE), we did not observe any clusters that reached a statistical significance (all p's>0.05).

Table 3. Nominally significant between-group differences in neuroinflammation in subsets.

Brain Structure	ROI (L/R) [a]	Statistics							
		≤ 60 Days [b]				≤ 90 Days [b]			
		β [95% CI] [c]	t	Uncorrected P-value	FDR corrected P-value	β [95% CI] [c]	t	Uncorrected P-value	FDR corrected P-value [d]
Subcortical									
	Hippocampus (R)					0.27 [0.03, 0.52]	2.22	0.027	0.463
Cortical									
	Pars Triangularis (L)	-0.39 [-0.75, -0.02]	-2.11	0.036	0.597	-0.24 [-0.46, -0.02]	-2.22	0.027	0.463
	Postcentral (L)					-0.21 [-0.41, -0.02]	-2.19	0.029	0.463
	Cuneus (R)	0.37 [0.02, 0.72]	2.13	0.035	0.597	0.23 [0.01, 0.45]	2.12	0.035	0.463
	Fusiform (R)					0.28 [0.04, 0.53]	2.33	0.020	0.463
	Pericalcarine Cortex (R)	0.41 [0.02, 0.80]	2.10	0.037	0.597				
	Rostral Anterior Cingulate (R)	0.52 [0.09, 0.95]	2.40	0.011	0.597	0.32 [0.06, 0.58]	2.45	0.015	0.463
	Rostral Middle Frontal (R)	-0.41 [-0.76, 0.07]	-2.39	0.029	0.597				
White Matter Tracts									
	Superior Longitudinal Fasciculus (L; Temporal Part)					0.34 [0.07, 0.60]	2.53	0.012	0.243

[a] ROI = Regions of Interest; L/R = Left / Right hemisphere
[b] Group comparisons of COVID-negative individuals with COVID-positive individuals whose COVID positive testing occurred within 60 or 90 days of their second imaging scan.
[c] CI = Confidence Interval
[d] FDR = False Discovery Rate

Discussion

In this study we examined whether COVID is associated with changes in neuroinflammation, as indexed by DBSI-RF, using a unique UKB prospective case-control cohort (n=224 cases; n=129 controls) that was scanned prior to and following COVID testing. In contrast to our hypotheses and prior studies of post-mortem brain tissue (31,38) and *in vivo* imaging of long-COVID cases (29), we found that COVID was not associated with changes in DBSI-RF following multiple testing correction. Several brain regions (e.g., orbitofrontal cortex, anterior cingulate cortex, ventral striatum) that have previously been shown to have the largest structural changes following COVID (16) had increased in DBSI-RF at nominal levels of significance among COVID cases (**Table 2**). However, given that these findings did not survive multiple testing correction and the directionality of nominally significant findings was not consistent across regions (e.g., reduced DBSI-RF among COVID cases in the caudal middle frontal gyrus), these nominal findings should be interpreted with caution. While our null findings were recapitulated when restricting analyses to cases that were scanned closer to their COVID test date (≤ 60 or 90 days), it is possible that COVID-related neuroinflammation may be temporally constrained and may have resolved prior to the post-COVID scan (e.g., 60-90 days post COVID following functional recovery of glia cells; (59–61)), while structural differences are observed in this sample (16). Alternatively, neuroinflammation may only be present during active symptomatology.

Infection-induced Neuroinflammation Progression and Clinical Heterogeneity

In contrast to our null findings, other studies suggest that the acute infection of COVID induces neuroinflammation and may have long-term consequences including neurological and psychiatric syndromes (63). Specifically, encephalitis may occur as early as 2 weeks after the onset of SARS-CoV-2 infection (27,28) and last up to two years after the infection with widespread influences in the brain in individuals suffering from long COVID (29). In additional to acute production of proinflammatory microglia (i.e., the most dominant immune cells in the brain) and a chronic loss of microglia (64), COVID has also been linked to damages in astrocytes and oligodendrocytes, which may gain a functional recovery 60-90 days after the infection onset and subsequently lead to reductions in neuroinflammation (59–61), but may also ultimately lead to neuronal cell death in the cerebral cortex (59,62). These findings suggest that COVID-related inflammatory processes may diverge depending on the assessment timing (65). As DBSI-RF is an indirect proxy of inflammation-related cellularity from either immune cell proliferation or infiltration (36,41,43,49), it remains possible that these inflammatory processes may have subsided by the time of the second imaging scan. Although DBSI-RF holds potential to capture a wide range of inflammatory processes, it may also require future investigation to determine its sensitivity in detecting inflammatory changes induced by COVID infection in the central nervous system. Further, due to the lack of information regarding the disease recovery and the real-time symptoms for COVID-positive cases, our null findings may have also been influenced by the clinical heterogeneity of COVID manifestations with some case participants possibly having fully recovered from the infection while some others still experiencing some degrees of symptoms by the time of the second imaging assessment.

It should also be noted that the present null findings were obtained after correcting for potential confounding effects of age, sex, ethnicity, BMI, smoking frequency, and data acquisition intervals. Additional factors may contribute to variations in clinical presentation and related neuroinflammation. For instance, a "two-hit" hypothesis on the link between microglial activation and COVID severity suggests that predisposed conditions such as exposures to childhood trauma (i.e., first hit) may sensitize individuals' microglia responses when facing the second immune challenge, such as COVID (66). In line with a survey study reporting increased risks of post-COVID conditions for individuals who had indicated prior-COVID distress (67),

future research may consider including stress-related factors to better understand the COVID-related neuroinflammation and its implications.

Nominal Associations of Neuroinflammation with COVID
A few nominally significant results (i.e., before FDR correction) are worth mentioning in the context of prior research findings. First, the lateral orbitofrontal cortex exhibited a nominally higher level of post-COVID neuroinflammation in individuals who were tested COVID-positive before the second imaging assessment. This effect is consistent with previous findings that the olfactory system, including orbitofrontal areas are particularly vulnerable to SARS-CoV-2 virus invasion (68–72). These vulnerabilities may manifest at the structural level of the brain such that the infected individuals exhibited greater reduction in grey matter thickness and tissue contrast in the orbitofrontal cortex (16). Additionally, our results indicated that regions that are functionally connected to the piriform cortex (i.e., the primary component of the olfactory network) including the amygdala, caudate, and the anterior cingulate cortex also showed nominally higher neuroinflammation levels in COVID-positive individuals (**Table 2**). Interestingly, these regions were previously reported with longitudinal anatomical changes in the same cohort (UK Biobank) of COVID-positive individuals as in the current study (16). In addition to these gray matter structures, the COVID-positive individuals had higher neuroinflammation in the temporal part of the superior longitudinal fasciculus (SLF), a white matter tract mainly connects the frontal and parietal cortices and plays a crucial role in language, attention, memory, and emotion (73,74). Altered diffusion properties in SLF have been reported in post-COVID individuals who had mild to moderate acute COVID infection (75,76), and these alterations appear to be persistent long after recovery of COVID (19,75). Thus, the nominally elevated neuroinflammation in SLF might also be indicative of this pathological process induced by COVID. However, as these findings did not survive FDR correction, they should be interpreted with caution.

Limitations
It is important to consider study limitations when interpreting these data. First, COVID-positive cases in this study were defined solely by SARS-Cov-2 testing while symptom and severity assessments, as well as vaccination status were unavailable (77). It is possible that assessment timing and clinical heterogeneity might have contributed to our null findings in COVID-induced neuroinflammation. Second, while the UK Biobank is population-based cohort, it is also self-selective (78) and predominantly White (**Table 1**), which may limit the generalizability of these findings. Third, information of COVID testing date for control participants were unavailable, making it impossible to model within-subject variability in neuroinflammation changes due to differences in the time interval between COVID testing and the second imaging assessment for the full study sample.

Conclusion
In contrast to our hypotheses, we found no evidence that COVID is associated with changes in neuroinflammation, at least as measured using DBSI-RF in a relatively large case (n = 224) - control (n = 192) sample with MRI scans prior to and following COVID testing results. Given previous findings in altered brain structure and function in the same cohort (n = 785 including n = 401 COVID cases; (16), our data suggest that DBSI-derived neuroinflammation index may not contribute to COVID-related changes in brain structure and function observed in this sample. Alternatively, neuroinflammation may occur in a more transient manner, contributing to structural changes in the brain that remain like a scar. As elevated neuroinflammation has been reported in postmortem brain tissues from severe COVID patients (31,62) and in long-COVID patients with persistent neurological or psychiatric symptoms (29,34), and different timing of

analysis may capture divergent inflammatory responses to COVID (65), it remains possible that neuroinflammation is time- and symptom-dependent. Due to the lack of relevant information in the current study, our null findings should be interpreted with caution.

Statement of Ethics

All participants in UK Biobank study have provided their written informed consent and that the current study has been granted an exemption from requiring ethics approval by the IRB at Washington University in St. Louis. This study was conducted under the UK Biobank Application ID 47267.

Conflict of Interest Statement

The authors have no conflicts of interest to declare.

Funding Sources

WZ was supported by McDonnell Center for Systems Neuroscience at Washington University in St. Louis. AG was supported by NSF DGE-213989. TH receives funding from NIH (R01 DK126826, NS109487 and HD070855). RB receives funding from NIH (R01 AG061162, R21 AA027827, R01 DA054750, U01 DA055367). JB receives funding from the NIH (NIMH R01 MH128286) and from the McDonnell Center for Systems Neuroscience.

Author Contributions

AA and RB conceived the idea. QW and TH contributed to DBSI processing. WZ analyzed the data and wrote the first draft of the article together with AG. JB and RB contributed to supervision. All authors contributed to the manuscript preparation.

Data Availability Statement

The UK Biobank data used in this study can be accessed by researchers upon application (https://www.ukbiobank.ac.uk/register-apply).

Reference

1. Akinbami LJ. SARS-CoV-2 Serology and Self-Reported Infection Among Adults — National Health and Nutrition Examination Survey, United States, August 2021–May 2022. MMWR Morb Mortal Wkly Rep [Internet]. 2022 [cited 2023 Jun 20];71. Available from: https://www.cdc.gov/mmwr/volumes/71/wr/mm7148a4.htm

2. Baj J, Karakuła-Juchnowicz H, Teresiński G, Buszewicz G, Ciesielka M, Sitarz R, et al. COVID-19: Specific and Non-Specific Clinical Manifestations and Symptoms: The Current State of Knowledge. J Clin Med. 2020 Jun 5;9(6):1753.

3. Ma Y, Deng J, Liu Q, Du M, Liu M, Liu J. Long-Term Consequences of COVID-19 at 6 Months and Above: A Systematic Review and Meta-Analysis. Int J Environ Res Public Health. 2022 Jun 3;19(11):6865.

4. Mehandru S, Merad M. Pathological sequelae of long-haul COVID. Nat Immunol. 2022 Feb;23(2):194–202.

5. Chasco EE, Dukes K, Jones D, Comellas AP, Hoffman RM, Garg A. Brain Fog and Fatigue following COVID-19 Infection: An Exploratory Study of Patient Experiences of Long COVID. Int J Environ Res Public Health. 2022 Nov 23;19(23):15499.

6. Premraj L, Kannapadi NV, Briggs J, Seal SM, Battaglini D, Fanning J, et al. Mid and long-term neurological and neuropsychiatric manifestations of post-COVID-19 syndrome: A meta-analysis. J Neurol Sci. 2022 Mar 15;434:120162.

7. Naseer S, Khalid S, Parveen S, Abbass K, Song H, Achim MV. COVID-19 outbreak: Impact on global economy. Front Public Health. 2022;10:1009393.

8. Cutler DM. The Costs of Long COVID. JAMA Health Forum. 2022 May 12;3(5):e221809.

9. Parsons N, Outsikas A, Parish A, Clohesy R, D'Aprano F, Toomey F, et al. Modelling the Anatomic Distribution of Neurologic Events in Patients with COVID-19: A Systematic Review of MRI Findings. AJNR Am J Neuroradiol. 2021 Jul;42(7):1190–5.

10. Lu Y, Li X, Geng D, Mei N, Wu PY, Huang CC, et al. Cerebral Micro-Structural Changes in COVID-19 Patients - An MRI-based 3-month Follow-up Study. EClinicalMedicine. 2020 Aug;25:100484.

11. Kumar PR, Shilpa B, Jha RK. Brain disorders: Impact of mild SARS-CoV-2 may shrink several parts of the brain. Neurosci Biobehav Rev. 2023 Jun;149:105150.

12. Du YY, Zhao W, Zhou XL, Zeng M, Yang DH, Xie XZ, et al. Survivors of COVID-19 exhibit altered amplitudes of low frequency fluctuation in the brain: a resting-state functional magnetic resonance imaging study at 1-year follow-up. Neural Regeneration Research. 2022 Jul;17(7):1576.

13. Hosp JA, Dressing A, Blazhenets G, Bormann T, Rau A, Schwabenland M, et al. Cognitive impairment and altered cerebral glucose metabolism in the subacute stage of COVID-19. Brain. 2021 May 7;144(4):1263–76.

14. Guedj E, Million M, Dudouet P, Tissot-Dupont H, Bregeon F, Cammilleri S, et al. 18F-FDG brain PET hypometabolism in post-SARS-CoV-2 infection: substrate for persistent/delayed disorders? Eur J Nucl Med Mol Imaging. 2021 Feb;48(2):592–5.

15. Churchill NW, Roudaia E, Chen JJ, Gilboa A, Sekuler A, Ji X, et al. Effects of post-acute COVID-19 syndrome on the functional brain networks of non-hospitalized individuals. Front Neurol. 2023;14:1136408.

16. Douaud G, Lee S, Alfaro-Almagro F, Arthofer C, Wang C, McCarthy P, et al. SARS-CoV-2 is associated with changes in brain structure in UK Biobank. Nature. 2022;604(7907):697–707.

17. Rau A, Schroeter N, Blazhenets G, Dressing A, Walter LI, Kellner E, et al. Widespread white matter oedema in subacute COVID-19 patients with neurological symptoms. Brain. 2022 Sep 14;145(9):3203–13.

18. Huang S, Zhou Z, Yang D, Zhao W, Zeng M, Xie X, et al. Persistent white matter changes in recovered COVID-19 patients at the 1-year follow-up. Brain. 2021 Dec 16;145(5):1830–8.

19. Huang S, Zhou X, Zhao W, Du Y, Yang D, Huang Y, et al. Dynamic white matter changes in recovered COVID-19 patients: a two-year follow-up study. Theranostics. 2023 Jan 1;13(2):724–35.

20. Merad M, Martin JC. Pathological inflammation in patients with COVID-19: a key role for monocytes and macrophages. Nat Rev Immunol. 2020 Jun;20(6):355–62.

21. Darif D, Hammi I, Kihel A, El Idrissi Saik I, Guessous F, Akarid K. The pro-inflammatory cytokines in COVID-19 pathogenesis: What goes wrong? Microb Pathog. 2021 Apr;153:104799.

22. Rabaan AA, Al-Ahmed SH, Muhammad J, Khan A, Sule AA, Tirupathi R, et al. Role of Inflammatory Cytokines in COVID-19 Patients: A Review on Molecular Mechanisms, Immune Functions, Immunopathology and Immunomodulatory Drugs to Counter Cytokine Storm. Vaccines (Basel). 2021 Apr 29;9(5):436.

23. Channappanavar R, Perlman S. Pathogenic human coronavirus infections: causes and consequences of cytokine storm and immunopathology. Semin Immunopathol. 2017;39(5):529–39.

24. Ragab D, Salah Eldin H, Taeimah M, Khattab R, Salem R. The COVID-19 Cytokine Storm; What We Know So Far. Frontiers in Immunology [Internet]. 2020 [cited 2023 Jun 24];11. Available from: https://www.frontiersin.org/articles/10.3389/fimmu.2020.01446

25. Meinhardt J, Radke J, Dittmayer C, Franz J, Thomas C, Mothes R, et al. Olfactory transmucosal SARS-CoV-2 invasion as a port of central nervous system entry in individuals with COVID-19. Nat Neurosci. 2021 Feb;24(2):168–75.

26. Krasemann S, Haferkamp U, Pfefferle S, Woo MS, Heinrich F, Schweizer M, et al. The blood-brain barrier is dysregulated in COVID-19 and serves as a CNS entry route for SARS-CoV-2. Stem Cell Reports. 2022 Feb 8;17(2):307–20.

27. Siow I, Lee KS, Zhang JJY, Saffari SE, Ng A. Encephalitis as a neurological complication of COVID-19: A systematic review and meta-analysis of incidence, outcomes, and predictors. Eur J Neurol. 2021 Oct;28(10):3491–502.

28. Ross Russell AL, Hardwick M, Jeyanantham A, White LM, Deb S, Burnside G, et al. Spectrum, risk factors and outcomes of neurological and psychiatric complications of COVID-19: a UK-wide cross-sectional surveillance study. Brain Commun. 2021 Jul 22;3(3):fcab168.

29. Visser D, Golla SSV, Verfaillie SCJ, Coomans EM, Rikken RM, Giessen EM van de, et al. Long COVID is associated with extensive in-vivo neuroinflammation on [18F]DPA-714 PET [Internet]. medRxiv; 2022 [cited 2023 Jun 20]. p. 2022.06.02.22275916. Available from: https://www.medrxiv.org/content/10.1101/2022.06.02.22275916v1

30. Frere JJ, Serafini RA, Pryce KD, Zazhytska M, Oishi K, Golynker I, et al. SARS-CoV-2 infection in hamsters and humans results in lasting and unique systemic perturbations after recovery. Science Translational Medicine. 2022 Jun 7;14(664):eabq3059.

31. Yang AC, Kern F, Losada PM, Agam MR, Maat CA, Schmartz GP, et al. Dysregulation of brain and choroid plexus cell types in severe COVID-19. Nature. 2021 Jul;595(7868):565–71.

32. Philippens IHCHM, Böszörményi KP, Wubben JA, Fagrouch ZC, Driel N van, Mayenburg AQ, et al. SARS-CoV-2 causes brain inflammation and induces Lewy body formation in macaques [Internet]. bioRxiv; 2021 [cited 2023 Jun 20]. p. 2021.02.23.432474. Available from: https://www.biorxiv.org/content/10.1101/2021.02.23.432474v2

33. Wischmann J, Bartos LM, Brendel M, Albert NL, Forbrig R, Straube A, et al. Translocator protein (TSPO)-PET as diagnostic and monitoring tool in COVID-19 related MRI-negative brainstem encephalitis: a case report. J Neurol. 2023;270(6):2853–6.

34. Braga J, Lepra M, Kish SJ, Rusjan PabloM, Nasser Z, Verhoeff N, et al. Neuroinflammation After COVID-19 With Persistent Depressive and Cognitive Symptoms. JAMA Psychiatry [Internet]. 2023 May 31 [cited 2023 Jul 5]; Available from: https://doi.org/10.1001/jamapsychiatry.2023.1321

35. Miller KL, Alfaro-Almagro F, Bangerter NK, Thomas DL, Yacoub E, Xu J, et al. Multimodal population brain imaging in the UK Biobank prospective epidemiological study. Nat Neurosci. 2016 Nov;19(11):1523–36.

36. Wang Y, Wang Q, Haldar JP, Yeh FC, Xie M, Sun P, et al. Quantification of increased cellularity during inflammatory demyelination. Brain. 2011 Dec 1;134(12):3590–601.

37. Cross AH, Song SK. "A new imaging modality to non-invasively assess multiple sclerosis pathology." J Neuroimmunol. 2017 Mar 15;304:81–5.

38. Schwabenland M, Salié H, Tanevski J, Killmer S, Lago MS, Schlaak AE, et al. Deep spatial profiling of human COVID-19 brains reveals neuroinflammation with distinct microanatomical microglia-T-cell interactions. Immunity. 2021 Jul 13;54(7):1594-1610.e11.

39. Sudlow C, Gallacher J, Allen N, Beral V, Burton P, Danesh J, et al. UK Biobank: An Open Access Resource for Identifying the Causes of a Wide Range of Complex Diseases of Middle and Old Age. PLOS Medicine. 2015 Mar 31;12(3):e1001779.

40. Wu W, Wang Q, Sun Z, Flores S, Hobbs DA, Franklin EE, et al. Initial Correlation Analysis of Diffusion Basis Spectrum Imaging of Alzheimer's Brain and Quantitative Histology. Alzheimer's & Dementia. 2022;18(S6):e064265.

41. Wang X, Cusick MF, Wang Y, Sun P, Libbey JE, Trinkaus K, et al. Diffusion basis spectrum imaging detects and distinguishes coexisting subclinical inflammation, demyelination and axonal injury in experimental autoimmune encephalomyelitis mice. NMR Biomed. 2014 Jul;27(7):843–52.

42. Zhan J, Lin TH, Libbey JE, Sun P, Ye Z, Song C, et al. Diffusion Basis Spectrum and Diffusion Tensor Imaging Detect Hippocampal Inflammation and Dendritic Injury in a Virus-Induced Mouse Model of Epilepsy. Front Neurosci. 2018;12:77.

43. Wang Q, Wang Y, Liu J, Sutphen CL, Cruchaga C, Blazey T, et al. Quantification of white matter cellularity and damage in preclinical and early symptomatic Alzheimer's disease. Neuroimage Clin. 2019;22:101767.

44. Shirani A, Sun P, Trinkaus K, Perantie DC, George A, Naismith RT, et al. Diffusion basis spectrum imaging for identifying pathologies in MS subtypes. Ann Clin Transl Neurol. 2019 Nov;6(11):2323–7.

45. Ly M, Raji CA, Yu GZ, Wang Q, Wang Y, Schindler SE, et al. Obesity and White Matter Neuroinflammation Related Edema in Alzheimer's Disease Dementia Biomarker Negative Cognitively Normal Individuals. J Alzheimers Dis. 2021;79(4):1801–11.

46. Samara A, Murphy T, Strain J, Rutlin J, Sun P, Neyman O, et al. Neuroinflammation and White Matter Alterations in Obesity Assessed by Diffusion Basis Spectrum Imaging. Frontiers in Human Neuroscience [Internet]. 2020 [cited 2022 Nov 1];13. Available from: https://www.frontiersin.org/articles/10.3389/fnhum.2019.00464

47. Zhang W, Rutlin J, Eisenstein SA, Wang Y, Barch D, Hershey T, et al. Neuroinflammation in the amygdala is associated with recent depressive symptoms. Biological Psychiatry:

Cognitive Neuroscience and Neuroimaging [Internet]. 2023 May 9 [cited 2023 Jun 20]; Available from: https://www.sciencedirect.com/science/article/pii/S2451902223001192

48. Alfaro-Almagro F, Jenkinson M, Bangerter NK, Andersson JLR, Griffanti L, Douaud G, et al. Image processing and Quality Control for the first 10,000 brain imaging datasets from UK Biobank. NeuroImage. 2018 Feb 1;166:400–24.

49. Wang Y, Sun P, Wang Q, Trinkaus K, Schmidt RE, Naismith RT, et al. Differentiation and quantification of inflammation, demyelination and axon injury or loss in multiple sclerosis. Brain. 2015 May;138(Pt 5):1223–38.

50. Shirani A, Sun P, Schmidt RE, Trinkaus K, Naismith RT, Song SK, et al. Histopathological correlation of diffusion basis spectrum imaging metrics of a biopsy-proven inflammatory demyelinating brain lesion: A brief report. Mult Scler. 2019 Dec;25(14):1937–41.

51. Ye Z, George A, Wu AT, Niu X, Lin J, Adusumilli G, et al. Deep learning with diffusion basis spectrum imaging for classification of multiple sclerosis lesions. Ann Clin Transl Neurol. 2020 May;7(5):695–706.

52. Strain JF, Burdo TH, Song SK, Sun P, El-Ghazzawy O, Nelson B, et al. Diffusion Basis Spectrum Imaging Detects Ongoing Brain Inflammation in Virologically Well Controlled HIV+ Patients. J Acquir Immune Defic Syndr. 2017 Dec 1;76(4):423–30.

53. Vavasour I, Sun P, Graf C, Yik J, Kolind S, Li D, et al. Characterisation of multiple sclerosis neuroinflammation and neurodegeneration with relaxation and diffusion basis spectrum imaging. Mult Scler. 2022 Mar;28(3):418–28.

54. Fischl B, Salat DH, Busa E, Albert M, Dieterich M, Haselgrove C, et al. Whole Brain Segmentation: Automated Labeling of Neuroanatomical Structures in the Human Brain. Neuron. 2002 Jan 31;33(3):341–55.

55. Desikan RS, Ségonne F, Fischl B, Quinn BT, Dickerson BC, Blacker D, et al. An automated labeling system for subdividing the human cerebral cortex on MRI scans into gyral based regions of interest. Neuroimage. 2006 Jul 1;31(3):968–80.

56. Mazziotta J, Toga A, Evans A, Fox P, Lancaster J, Zilles K, et al. A probabilistic atlas and reference system for the human brain: International Consortium for Brain Mapping (ICBM). Philos Trans R Soc Lond B Biol Sci. 2001 Aug 29;356(1412):1293–322.

57. Winkler AM, Ridgway GR, Webster MA, Smith SM, Nichols TE. Permutation inference for the general linear model. Neuroimage. 2014 May 15;92(100):381–97.

58. Smith SM, Nichols TE. Threshold-free cluster enhancement: addressing problems of smoothing, threshold dependence and localisation in cluster inference. Neuroimage. 2009 Jan 1;44(1):83–98.

59. Plantone D, Locci S, Bergantini L, Manco C, Cortese R, Meocci M, et al. Brain neuronal and glial damage during acute COVID-19 infection in absence of clinical neurological manifestations. J Neurol Neurosurg Psychiatry. 2022 Dec 1;93(12):1343–8.

60. Steardo L, Steardo L, Scuderi C. Astrocytes and the Psychiatric Sequelae of COVID-19: What We Learned from the Pandemic. Neurochem Res. 2023 Apr;48(4):1015–25.

61. Monje M, Iwasaki A. The neurobiology of long COVID. Neuron. 2022 Nov 2;110(21):3484–96.

62. Boroujeni ME, Simani L, Bluyssen HAR, Samadikhah HR, Zamanlui Benisi S, Hassani S, et al. Inflammatory Response Leads to Neuronal Death in Human Post-Mortem Cerebral Cortex in Patients with COVID-19. ACS Chem Neurosci. 2021 Jun 16;12(12):2143–50.

63. Spudich S, Nath A. Nervous system consequences of COVID-19. Science. 2022 Jan 21;375(6578):267–9.

64. Jeong GU, Lyu J, Kim KD, Chung YC, Yoon GY, Lee S, et al. SARS-CoV-2 Infection of Microglia Elicits Proinflammatory Activation and Apoptotic Cell Death. Microbiol Spectr. 2022 Jun 29;10(3):e0109122.

65. Liu C, Martins AJ, Lau WW, Rachmaninoff N, Chen J, Imberti L, et al. Time-resolved systems immunology reveals a late juncture linked to fatal COVID-19. Cell. 2021 Apr 1;184(7):1836-1857.e22.

66. Bouayed J, Bohn T. The link between microglia and the severity of COVID-19: The "two-hit" hypothesis. J Med Virol. 2021 Jul;93(7):4111–3.

67. Wang S, Quan L, Chavarro JE, Slopen N, Kubzansky LD, Koenen KC, et al. Associations of Depression, Anxiety, Worry, Perceived Stress, and Loneliness Prior to Infection With Risk of Post–COVID-19 Conditions. JAMA Psychiatry. 2022 Nov 1;79(11):1081–91.

68. Butowt R, Bilinska K. SARS-CoV-2: Olfaction, Brain Infection, and the Urgent Need for Clinical Samples Allowing Earlier Virus Detection. ACS Chem Neurosci. 2020 May 6;11(9):1200–3.

69. Netland J, Meyerholz DK, Moore S, Cassell M, Perlman S. Severe acute respiratory syndrome coronavirus infection causes neuronal death in the absence of encephalitis in mice transgenic for human ACE2. J Virol. 2008 Aug;82(15):7264–75.

70. Brann DH, Tsukahara T, Weinreb C, Lipovsek M, Van den Berge K, Gong B, et al. Non-neuronal expression of SARS-CoV-2 entry genes in the olfactory system suggests mechanisms underlying COVID-19-associated anosmia. Sci Adv. 2020 Jul 31;6(31):eabc5801.

71. Carmichael ST, Clugnet MC, Price JL. Central olfactory connections in the macaque monkey. J Comp Neurol. 1994 Aug 15;346(3):403–34.

72. Palouzier-Paulignan B, Lacroix MC, Aimé P, Baly C, Caillol M, Congar P, et al. Olfaction Under Metabolic Influences. Chem Senses. 2012 Nov;37(9):769–97.

73. Janelle F, Iorio-Morin C, D'amour S, Fortin D. Superior Longitudinal Fasciculus: A Review of the Anatomical Descriptions With Functional Correlates. Frontiers in Neurology [Internet]. 2022 [cited 2023 Jun 26];13. Available from: https://www.frontiersin.org/articles/10.3389/fneur.2022.794618

74. Wang X, Pathak S, Stefaneanu L, Yeh FC, Li S, Fernandez-Miranda JC. Subcomponents and connectivity of the superior longitudinal fasciculus in the human brain. Brain Struct Funct. 2016 May 1;221(4):2075–92.

75. Petersen M, Nägele FL, Mayer C, Schell M, Petersen E, Kühn S, et al. Brain imaging and neuropsychological assessment of individuals recovered from a mild to moderate SARS-CoV-2 infection. Proc Natl Acad Sci U S A. 2023 May 30;120(22):e2217232120.

76. Bispo DD de C, Brandão PR de P, Pereira DA, Maluf FB, Dias BA, Paranhos HR, et al. Brain microstructural changes and fatigue after COVID-19. Front Neurol. 2022 Nov 10;13:1029302.

77. Ssentongo P, Ssentongo AE, Voleti N, Groff D, Sun A, Ba DM, et al. SARS-CoV-2 vaccine effectiveness against infection, symptomatic and severe COVID-19: a systematic review and meta-analysis. BMC Infectious Diseases. 2022 May 7;22(1):439.

78. Fry A, Littlejohns TJ, Sudlow C, Doherty N, Adamska L, Sprosen T, et al. Comparison of Sociodemographic and Health-Related Characteristics of UK Biobank Participants With Those of the General Population. Am J Epidemiol. 2017 Nov 1;186(9):1026–34.

Table S1. Associations between neuroinflammation in ROIs and COVID.

Brain Structure	Index	ROI Name (L/R) [a]	Statistics	
			DBSI-RF L-R Correlation (Pearson's r)	Between-group Difference (*t*) [b]
Subcortical				
	1	Accumbens Area	0.34	1.12 / 1.30
	2	Amygdala	0.55	0.82 / 2.49[*]
	3	Caudate	0.28	-0.05 / 2.08[*]
	4	Hippocampus	0.76	0.21 / 0.99
	5	Pallidum	0.73	-1.37 / 0.63
	6	Putamen	0.62	-0.59 / -0.34
	7	Thalamus	0.72	0.40 / 0.14
Cortical				
	1	Caudal Anterior Cingulate	0.50	0.18 / 2.54[*]
	2	Caudal Middle Frontal	0.64	-1.99[*] / -1.61
	3	Cuneus	0.73	0.02 / 1.39
	4	Entorhinal	0.57	0.57 / 0.85
	5	Frontal Pole	0.59	-1.00 / -1.31
	6	Fusiform	0.66	1.24 / 2.20[*]
	7	Inferior Parietal	0.54	-0.81 / 0.30
	8	Inferior Temporal	0.45	0.54 / 0.21
	9	Insula	0.55	1.69 / 0.56
	10	Isthmus Cingulate	0.69	1.09 / 1.55
	11	Lateral Occipital	0.75	0.05 / -0.43

	12	Lateral Orbitofrontal	0.64	1.31 / 2.09[*]
	13	Lingual	0.75	0.23 / 1.56
	14	Medial Orbitofrontal	0.64	-0.36 / 1.18
	15	Middle Temporal	0.32	0.62 / 0.60
	16	Paracentral	0.53	-1.00 / -0.56
	17	Parahippocampal	0.57	0.50 / 0.62
	18	Pars Opercularis	0.46	-0.30 / -0.03
	19	Pars Orbitalis	0.46	1.02 / 0.59
	20	Pars Triangularis	0.23	-1.45 / 0.44
	21	Pericalcarine	0.70	-0.20 / 0.90
	22	Postcentral	0.63	-2.05[*] / -0.94
	23	Posterior Cingulate	0.70	-1.07 / 0.24
	24	Precentral	0.59	-1.33 / -0.61
	25	Precuneus	0.80	-0.66 / 0.29
	26	Rostral Anterior Cingulate	0.46	-0.62 / 2.94[*]
	27	Rostral Middle Frontal	0.71	-0.66 / -1.10
	28	Superior Frontal	0.86	-1.13 / -0.70
	29	Superior Parietal	0.85	-1.99[*] / -1.49
	30	Superior Temporal	0.51	1.41 / -0.63
	31	Supramarginal	0.56	-0.34 / -0.39
	32	Temporal Pole	0.38	-0.06 / 1.04

	33	Transverse Temporal	0.55	-0.59 / 0.08
White Matter Tracts				
	1	Anterior Thalamic Radiation	0.59	-0.60 / -0.23
	2	Cingulum (Cingulate Gyrus)	0.42	-1.24 / -0.80
	3	Cingulum (Hippocampus)	0.34	0.51 / -0.48
	4	Corticospinal Tract	0.46	-1.41 / -0.77
	5	Forceps (Major/Minor)	0.23	-0.26 / -0.40
	6	Inferior Fronto-occipital Fasciculus	0.41	-1.27 / -0.17
	7	Inferior Longitudinal Fasciculus	0.60	0.85 / 1.06
	8	Superior Longitudinal Fasciculus	0.45	-0.39 / -1.16
	9	Superior Longitudinal Fasciculus	0.13	2.66[*] / 0.24
	10	Uncinate Fasciculus	0.32	-1.37 / -0.59

[a] ROI = Region of Interest; L/R = Left / Right hemisphere
[b] Group difference between the COVID-positive and negative participants after adjusting for the covariates.
[*] p < 0.05 before FDR correction.