Title: Evolutionary trade-offs in dormancy phenology

Théo Constant [1] , F. Stephen Dobson[1,2] , Caroline Habold[1] * & Sylvain Giroud[3,4] *

[1] UMR 7178, Centre National de la Recherche Scientifique, Institut Pluridisciplinaire Hubert
CURIEN, Université de Strasbourg, Strasbourg, France

[2] Department of Biological Sciences, Auburn University, Auburn, AL, United States

[3] Department of Interdisciplinary Life Sciences, Research Institute of Wildlife Ecology,
University of Veterinary Medicine Vienna, Vienna, Austria

[4] Energetic Lab, Department of Biology, Northern Michigan University, Marquette, MI, USA

* Co-senior authorship

Corresponding authors: Sylvain Giroud; sgiroud@nmu.edu ; Energetic Lab, Department of
Biology, Northern Michigan University, Marquette, MI, USA & Caroline Habold ;
caroline.habold@iphc.cnrs.fr ; Institut Pluridisciplinaire Hubert Curien CNRS, 23 rue du Lœss,
Strasbourg 67037, France

Data accessibility statement: the data and computer code supporting the results are available at

https://github.com/Theo-Constant/Evolutionary-trade-offs-in-dormancy-phenology.git

Acknowledgments: the authors are grateful to the researchers for providing specific information

on previous studies including published data, in particular to Carina Siutz, Danielle Levesque,

Andrey Tchabovsky, Eric Rickart, Philip Leitner and John Hoogland for sharing data, some of

which have been used in the meta-analysis of this article. The authors are also grateful to John

Drake and two referees who provided excellent suggestions for manuscript improvements. SG was

financially supported by the Austrian Science Fund (FWF, Grant No. P31577-B25) and the

Austrian Agency for International Cooperation in Education and Research (OeAD – Scientific and

Technological Cooperation, Grant No. FR 09/2020).

Abstract

Seasonal animal dormancy, hibernation or diapause, is widely interpreted as a physiological response for surviving energetic challenges during the harshest times of the year (the physiological constraint hypothesis). However, there are other mutually non-exclusive hypotheses to explain the timing of animal dormancy, that is, entry into and emergence from hibernation (i.e. dormancy phenology). Other survival advantages of dormancy that have been proposed are reduced risks of predation and competition (the "life-history" hypothesis), but comparative tests across animal species are not yet available. Under this hypothesis, dormancy phenology is influenced by a trade-off between reproductive advantages of being active and survival benefits of dormancy. Within a species, males and females differ in the amount of time and energy they invest in reproduction. Thus, the trade-off between reproduction and survival may be reflected by within-species sex differences in the phenology of dormancy. To examine this hypothesis, we used two complementary approaches: (i) a set of phylogenetic comparative analyses on mammals (mainly holarctic rodents), and (ii) a comparison between endotherm and ectotherm dormancy, via analyses of endotherms (including mainly holoarctic rodents) and the existing literature on ectotherms.

Using the phylogenetic comparative method applied to more than 20 hibernating mammalian species, we found support for both hypotheses as explanations for the phenology of dormancy. In accordance with the life history hypotheses, sex differences in emergence and immergence were favored by the sex difference in reproductive effort. In addition, physiological constraint may influence the trade-off between survival and reproduction such that, low temperature and precipitation as well as smaller body mass influence sex differences in phenology. We also compiled initial evidence that ectotherm dormancy (invertebrates and reptiles) may be 1) less temperature dependent than previously thought and 2) associated with trade-offs consistent

with the life history hypothesis. Dormancy in some endotherms and ectotherms show staggered

phenology with respect to the growing season (earlier emergence and immergence than expected)

which illustrates the selection pressure exerted by the trade-off between reproduction (earlier

emergence than expected) and adult survival (earlier immergence than expected). Thus, dormancy

during non-life-threatening periods that are unfavorable for reproduction may be more widespread

than previously appreciated.

Keywords: annual cycle, dormancy, energy balance, hibernation, metabolic suppression,

overwintering, protandry, reproductive investment, risk spreading.

1) Introduction

A large number of species across the tree of life enter prolonged dormancy each year (Wilsterman et al. 2021). From a physiological point of view, dormancy occurs under a combination of high energy reserves and a significant reduction in energy demand, thus allowing prolonged inactivity for several months to several years (Hoehler and Jørgensen 2013, Staples 2016). From an evolutionary perspective, dormancy improves survival during stressful environmental periods, until subsequent conditions that favor reproduction (Watts and Tenhumberg 2021). The evolutionary tactic of dormancy has been studied independently among different major phylogenetic groups (Wilsterman et al. 2021), and these separate considerations have limited the development of a global evolutionary framework to explain dormancy.

The dormancy of plants, micro-organisms and some invertebrates has been extensively studied from an evolutionary point of view. In these species and in addition to energetic benefits, dormancy occurs in a large number of situations that reduce competition and predation (Danks 1992, Satterthwaite 2010, Blath and Tóbiás 2020). The evolution of dormancy is explained as based, for example, on "evolutionarily stable strategies" (Hairston Jr and Munns Jr 1984, Kortessis and Chesson 2019), "life-history theory" (Ji 2011, Watts and Tenhumberg 2021), or as a "bet-hedging strategy" (Hopper 1999, Joschinski and Bonte 2021).

In most animals, however, the topic of prolonged dormancy has primarily focused on ecophysiology rather than evolutionary biology. Dormancy duration increases with physiological constraints in the environment (e.g. energy shortage, thermic, or water constraints), demonstrating its adaptive role in response to short growing season (Pianka 1970, Turbill and Prior 2016, Wilsterman et al. 2021). However, some animals immerge into dormancy while environnemental conditions would allow (from a physiological point of view) a continuation of activity, suggesting

other survival benefits than coping with a short growing season (Jameson and Allison 1976,

Wiklund et al. 1996, Humphries et al. 2002). A reduction in the risk of predation or competition

during animal dormancy has been suggested based on increased survival during hibernation,

compared to the active season (Turbill et al. 2011, Ruf et al. 2012, Constant et al. 2020), in

particular from studies of the hibernating edible dormouse (*Glis glis*) (Bieber and Ruf 2009, Bieber

et al. 2014, Hoelzl et al. 2015, Ruf and Bieber 2022). To date, however, the generality of an

influence of these factors on the evolution of prolonged dormancy lacks attention. This raises the

question of whether the timing of animal dormancy, that is dormancy phenology, might be

exclusively explained by physiological constraint or whether other ecological factors may be

involved.

Among dormant animals, a general distinction is made between heterothermic endotherms

(some mammals and birds) that are able to actively influence their metabolic rate via oxidative

metabolism and ectotherms (invertebrates, fish, amphibians, and reptiles) whose metabolic rates

are more subject to microclimatic fluctuations (Staples 2016). In both cases of heterothermy,

although dormancy drastically reduces energy expenditure, some energy is nonetheless lost in the

absence of an external source. As a consequence, if dormancy phenology is explained solely by

physiological issues (the physiological constraint hypothesis), selection should favor remaining

active until a positive energy balance (in endotherms) or thermal window favorable for activity (in

ectotherms ; see Gunderson and Leal 2016) is no longer possible (Fig. 1a). If, however, there are

other benefits to dormancy such as improved survival due to a reduction of predation risk, these

survival benefits may produce a trade-off between being active and investing in reproduction

versus being dormant for a time to increase survival (the life-history hypothesis). Nevertheless,

these two hypotheses are not mutually exclusive, since harsh climatic conditions and low food

availability may reduce the benefits of being active for reproductive success and thus influence

this trade-off. Within species, this trade-off may be reflected by sex differences in the phenology

of dormancy, since males and females exhibit differences in reproductive timing and investment

(Emlen and Oring 1977).

In a recent study, predation avoidance and sexual selection received support for explaining

intraspecific variation in hibernation phenology in the northern Idaho ground squirrel (*Urocitellus*

brunneus, Allison et al. n.d.) Males often emerged from dormancy and arrived at mating sites some

days or weeks before females (termed "protandry"), and mating occurred shortly after female

emergenced from dormancy. Sexual selection may favor a life history in which relatively early-

emerging males benefit from greater reproductive success (the "mating opportunity hypothesis,"

Morbey and Ydenberg 2001). Males that are physiologically prepared to mate (Breedveld and

Fitze 2016) and have established intrasexual dominance or territories (Manno and Dobson 2008,

Hibbitts et al. 2012) prior to mating are likely to have greater reproductive success (Michener

1983, 1984). Thus, greater protandry is assumed to have evolved with intraspecific competition

and longer periods of mating preparation. For females, emergence phenology may promote

breeding and/or care of offspring during the most favorable annual period (e.g., a match of the

peak in lactational energy demand and maximum food availability, Fig. 1) or beginning early to

afford long active seasons for offspring while not compromising the survival of parents. Although

males are active above ground before females, the latter sex may not emerge until later to limit

mortality risks (see the "waiting cost hypothesis," Morbey and Ydenberg 2001). Later in the year,

both sexes are expected to enter dormancy when they are no longer investing in or recovering from

reproduction, and after acquiring adequate energy stores for overwintering.

In the present study, we investigated the "physiological constraints" and "life-history"

hypotheses, two non-mutually exclusive evolutionary explanations for the phenology of

dormancy, especially in regard to sex differences within species. To examine these hypotheses, we

used two complementary approaches: (i) a set of phylogenetic comparative analyses, and (ii) a

comparison between endotherm and ectotherm dormancy via the conducted analyses (from

endotherms, including mainly holoarctic rodents) and the existing literature (from ectotherms).

First, using phylogenetic comparative analyses, we compared traits associated with climat,

reproduction, body mass, and hibernation to explain sex differences in phenology in more than 20

hibernating mammals (Fig. 2). To begin with, we predicted from the life history hypothesis that

(1) males of species with high energetic cost of reproduction would exhibit greater protandry due

to longer mating preparation. We tested whether body mass lost during mating was associated

with greater protandry. For immergence, males were expected to be constrained by the long-term

negative effects of reproductive stress (Millesi et al. 1998), whereas female immergence was

contrained by maternal effort (Levesque et al. 2013). Thus, we expected that (2) the species with

higher sex differences in energetic cost of reproduction would have a greater sex difference in the

timing of immergence, with the sex that invested more energy or time in reproduction also being

the one that immerged last. If sex differences in hibernation phenology were solely explain by the

physiological constraint hypothesis, then we would predict that species with high sexual

dimophism (proxy of sex difference in energetic demand) in body mass would exhibit greater sex

differences in hibernation phenology. On the contrary, if physiological constraints influenced the

trade-off between survival and reproduction, then we would predict that that species living in short

growing seasons (low temperature and precipitation) and smaller species would exhibit lower sex

differences in hibernation phenology due to higher cost of thermoregulation during activity.

Secondly, there is a need to unify the study of dormancy across ectotherms and endotherms for

answering fundamental evolutionary questions (Wilsterman et al. 2021). While there are

insufficient data on dormancy phenology and reproductive investment in ectotherms to allow a

statistical comparative analysis, we examined, as a second step, the relationships between

reproductive investment, energy balance, thermal contraints, and dormancy phenology of

ectotherms that are already available in the literature. We compared these studies for ectotherms

to our results for endotherms. Finally, we highlighted evidence from the literature for the

independence of dormancy phenology from energy balance.

2) Materials and methods

a) Review Criteria

Our literature review was based on 152 hibernating species of mammals (see

supplementary material 1 in Constant et al., 2020). In this study, we addressed what can be called

ecological hibernation, i.e. the seasonal duration that an animal remains sequestered in its burrow

or den, which is assumed to be directly linked to the reduced risk of predation. In contrast, we did

not consider heterothermic hibernation, which corresponds to the time between the beginning and

end of the use of torpor. So when we mention hibernation, emergence or immergence, the specific

reference is to ecological hibernation. We excluded non-seasonal hibernating species that do not

have a consistent seasonal hibernation phenology (elephant shrew and marsupial species except

Burramys parvus (the mountain pygmy possum)). We did not include species from the order

Carnivora and *Chiroptera* because of a difference in reproductive phenology compared to the

majority of other hibernators, especially due to delayed embryo implantation (Sandell 1990). This

implies different trade-offs between hibernation and reproduction that require separate analyses.

In addition, there were few data available on both reproduction and hibernation for hibernating bat

species (see below for traits needed for inclusion; also the Discussion for hypotheses applied to

these groups). Only one bird species is considered to be a hibernator, and no information is

available on sex differences in hibernation phenology (Woods and Brigham 2004, Woods et al.

2019).

Each of the following literature reviews was conducted using the search engine Google

Scholar with specific keywords and considered articles up to and including January 2021. In total,

our literature search allowed inclusion of 29 hibernating mammals in the analyses for which we

have both reproduction and hibernation data including mainly rodents, a monotreme, a primate

and an *Eulipotyphla* species.

b) Sex difference in hibernation phenology

We searched for hibernation phenology for each sex based on average date of emergence

and immergence in the same population. When these types of data were not available, we accepted

the date at which first/last individuals of each sex were observed or the approximate sex difference

available in the text (see supplementary material S1 for search criteria).

From the remaining data, we calculated protandry and the sex difference in immergence

into hibernation as female Julian date – male Julian date.

We have excluded immergence data for *Tamias sibericus* because this species does not

follow the general phenology of other species (see "Study limitations" section for details).

c)   Sex differences in reproductive investment and sexual dimorphism

For males, relative body mass changes during the mating period (hereafter referred to as

"Δ body mass during mating") was calculated as follows: (body mass at the end of mating - body

mass at the beginning of mating) / body mass at the beginning of mating. Male immergence is

expected to be constrained by the long-term negative effects of reproductive stress (Millesi et al.

1998). In addition to energetic costs of mating, some recently emerged males lost body mass before

females emerged from hibernation, which may have resulted in physiological stress. Thus, we

calculated relative body mass change between emergence and the end of mating, hereafter referred

to as "Δ body mass through the end of mating" as follows: (body mass at the end of mating - body

mass at emergence) / body mass at emergence.

Maternal effort may have contrained female immergence in two ways: the duration and

energy cost of maternal care. Maternal effort duration was calculated as the sum of the gestation

and lactation periods. Energy cost of reproductive effort also known as "specific reproductive

effort" is calculated as the mean annual litter size times juvenile body mass at weaning divided by

the body mass of adult female before reproduction (Armitage 1981).

For the analyses, we only accepted reproductive and body mass data from the same or very

close study sites as the hibernation phenology data. We extracted body mass at immergence and

emergence for both sexes and calculated sexual dimorphism as female body mass divided by male

body mass. We made a few exceptions for juvenile body mass at weaning, for which some data

were obtained in Hayssen (2008). As maternal effort duration and lactation data were not available

in the literature for *Urocitellus brunneus* and *Ictidomys parvidens* respectively, we used averages

for the clade *Marmotini* obtained from Hayssen (2008).

d) Climate data

Species living in harsh conditions may be constrained by a shorter active season that might

influence sex differences in hibernation phenology. To take this into account in the models (see

section "Statistics"), the location (latitude and longitude) of hibernation study sites were recorded,

and when not provided we determined their location using Google Earth and the location

description. Then the location data were used to extract values of the mean temperature of the

coldest month (known as "BIO6" in the database and hereafter referred to as minimum

temperature) and annual precipitation (known as "BIO12" in the database) from an interpolated

climate surface (BIOCLIM) with 1 km^2 resolution (30 sec) based on data for the period 1970-2000

(Hijmans et al. 2005).

e) Statistics

We used phylogenetic generalized least squares (PGLS) models (see supplementary

materials S1 for details and mammalian phylogeny) to account for the non-independence of

phylogeny-related species with the "ape 5.0," "apTreeshape 1.5," and "caper 1.0" packages in R

v. 3.6.2 (Orme et al., 2013; Paradis, 2011; Paradis and Schliep, 2019; R Core Team, 2019). Each

PGLS model produces a λ parameter representing the effect of phylogeny ranging between 0 (no

phylogeny effect) and 1 (covariance entirely explained by co-ancestry).

The PGLS models used only one value per species for each factor. For hibernation

phenology, body mass, reproductive effort and climatic variable, we first averaged by study when

data were available over several years, and then we averaged the data for the species. This produces

equal weighting between studies on the same species.

To test the life history and physiological constraint hypotheses on emergence, protandry

was the dependent variable while , Δ body mass during mating, male body mass at emergence,

sexual dimorphism at emergence, minimum temperature, annual precipitation, foodstoring and late

mating were independent variables. The "late mating" factor aims to test for lower protandry when

mating was more greatly delayed after the onset of the annual activity period, as has been shown

for reptiles (Olsson et al. 1999), hereafter referred to « delay in mating ». The "food-storing" factor

tested for lower protandry for species that store food in a burrow and consume it after the last

torpor bouts, which may allow them to prepare for reproduction without emerging above ground

(Williams et al. 2014). Food-storing species have been identified in several studies (Kenagy et al.

1989, Vander Wall 1990, Michener 1992, Bieber and Ruf 2004). To test the life history and

physiological constraint hypotheses on immergence, sex difference in immergence was the

dependent variable while Δ body mass through the end of mating, duration of maternal effort,

specific reproductive effort of female, sexual dimorphism in impmergence, minimum temperature

and annual precipitation were independent variables. In both models, we tested for a "two factor

interaction" between either minimum temperature, annual precipitation male and female

reproductive effort.   were independent variables.

The two models are described hereafter (see supplementary materials S2 for datasets) :

Protandry ~ late_mating + foodstoring + male body mass + dimorphisme body mass at

emergence + Δ body mass during mating * Minimum temperature + Δ body mass during mating

* annual precipitation

Sex difference in immergence ~ log(specific reproductive effort) * Minimum temperature +

log(specific reproductive effort) * precipitation + Δ body mass through the end of mating *

Minimum temperature + Δ body mass through the end of mating * Precipitation + Maternal

effort duration* Precipitation + Maternal effort duration * Minimum temperature + Body mass

immergence + Dimorphisme at immergence

For both models, we used the dredge function of the MuMIn package (version 1.43.17;

Barto 2020) to select the best model based on the corrected Akaike information criterion (AICc).

Normality and homoscedasticity were checked by graphical observation. We tested for

multicollinearity using variance inflation factors (we required VIF < 3) on linear models including

the factors of the best models. PGLS models do not include calculations of VIFs (Wartel et al.

2019, Ancona et al. 2020). Female-specific reproductive effort was log-transformed to improve

the fit to normality of the residuals. All independent variables were standardized (using *z-scores*)

in multi-factor models, so that their coefficients are directly comparable as estimates of effect sizes

(Abdi 2007).

Results

a) Emergence

Protandry increased significantly with male body mass loss during mating and increased

ambient temperature (model 1 in Table 1 and Fig. 3). At the same time, protandry decreased in

species for which males are smaller. The effect of ambient temperature on protandry is almost 2

times greater than male body mass and body mass loss during mating. There is also a trend

towards lower protandry with increasing time between female emergence and the start of mating

but this relationship is not significant ("late mating" in model 1, Table 1). This model showed no

influence of phylogeny.

285   b) Immergence

The sex difference in immergence date was associated with maternal effort duration, specific

reproductive effort of female, Δ body mass through the end of mating and precipitation (model 2,

Table 1). High reproductive effort for both males and females delayed their immergence (Fig. 4

and 5). At the same time, males immerged before females for species living in an environment

with high annual precipitation. The effect of maternal effort duration was influenced by an outlier,

Tachyglossus aculeatus, a species for which gestation and lactation for females was an extremely

long period (*i.e.* 167 day). Maternal effort duration was not conserved in the best model when the

outlier was removed (model 3, Table 1). There is also a trend towards early male immergence in

species with male biaised body mass dimorphism at immergence but this relationship is not

significant (model 3, Table 1). Finally, phylogeny had no influence on this model.

296
297

298   **4) Discussion**

299   a) Sex difference in dormancy phenology

In this study, we investigated two mutually non-exclusive hypotheses about dormancy phenology : the physiological contraints and life history hypotheses. The sex difference in hibernation phenology is a good opportunity to confront these hypotheses, because the sexes are faced with somewhat different life-history challenges. The life-history hypothesis predicts a trade-off between the survival benefit of hibernation and the reproductive benefit of activity, such that the sex with lesser investment of time and energy in reproduction should spend more time in hibernation. If the physiological constraints hypothesis was the sole explanation for hibernation phenology, then species with high sexual dimorphism (proxy for energy demand) should show a large sex difference in hibernation phenology. As well, if physiological constraints influence the trade-off between survival and reproduction, then small species and species living in short growing season should show lower sex differences due to the higher costs of being active. Based on phylogenetic comparative analyses, we found supports for both hypotheses. In accordance with the life history hypothesis, sex difference in emergence and immergence seem to reflect sex differences in reproductive effort. The comparative method, however, did not allow assignment of causation of one variable on the other; that is, of the causal direction of selection pressures between reproductive investments and hibernation phenology. In addition, physiological constraint may influence the trade off between survival and reproduction such that, low temperature and precipitation as well as smaller body mass influence sex differences in phenology.

Males with higher body mass loss during mating showed higher protandry. The males of some species increased locomotion and are subject to strong competition during mating, which greatly reduces their foraging time (Millesi et al. 1998). Thus, an early emergence of males may have evolved in response to sexual selection to accumulate energy reserves in anticipation of reproductive efforts. Females, on the contrary, are not subject to intraspecific competition for

reproduction and may have sufficient time before (generally one week after emergence) and during

the breeding period to improve their body condition. Protandry may also allow the maturation of

the reproductive system or monopolization of territories (Barnes et al. 1988, Millesi et al. 2008b,

Thompson et al. 2023). In addition, protandry decreases in species living in cold environments

and in species with lower male body mass in accordance with the physiological contraints

hypothesis. Despite the reproductive benefits of early emergence, the cost of activity may be too

high for these species, probably due to excessive heat loss in a cold environment.

In some species, males may have adaptations that improve their physical condition before

reproduction without early emergence in cold climate. For example, males of some fat-storing

species (e.g. *Urocitellus Parryii)* hoard food in their burrows. This energetic supply would support

a return to euthermia of up to a few weeks prior to behavioral emergence and allow for testes

maturation and fat accumulation (Michener 1992, Williams et al. 2014), while remaining sheltered

in the burrow. Thus, males might gain energy benefits without paying survival costs of above-

ground activity (Turbill et al. 2011, Constant et al. 2020). Although food storage in the burrow

may have evolved to overcome short growing season or predation, model selection did not retain

the food-storing factor. Thus, the ability to accumulate food in the burrow was not by itself likely

to keep males of some species from emerging earlier (e.g. *Cricetus cricetus*, protandry : 20 day,

Siutz et al., 2016). Early emerging males may benefit from consuming higher quality food or in

competition with other males (e.g., dominance assertion or territory establishment, Manno and

Dobson 2008, Thompson et al. 2023).

There is a trend for decrease protandry in species for which reproduction occurs several

weeks after female emergence, as demonstrated in lizards and snakes (Graves and Duvall 1990,

Olsson et al. 1999). According to the life history hypothesis, the benefits for males to emerge

before females decreases with the delay in the mating period (relative to female emergence),

because they are less constrained by time for mating preparation. This phenomenon concerns only

a few species in our comparative analysis, which may explain its non-significant effect.

In ectotherms, testes maturation has been proposed as a major influence on protandry. In

Zootoca vivipara (a viviparous lizard), male lizards generally emerged earlier than females

(Breedveld and Fitze 2016). The sex difference did not seem to be explained by a difference in the

maturation duration of the reproductive organs. In this species, females did not have developed

follicles at emergence and ovulation occured several weeks after mating (Bauwens and Verheyen

1985). In addition, by experimentally manipulating the emergence from dormancy of males but

not females, it was shown that the degree of protandry affected the order of sperm presence in

males, but not the probability of copulation. Thus, protandry may have increased the chances of

fertilizing eggs for males and decreased the probability of copulating with an infertile male for

females (Breedveld and Fitze 2016). In *Gonepteryx rhamni* (the common brimstone butterfly),

males emerged from dormancy 3 weeks before females. Males were quickly ready to reproduce,

but this delay would have allowed them to increase their amount of sperm before mating and thus

reproduce more successfully (Wiklund et al. 1996).

Unlike emergence, the sex that immerges into hibernation first varies among species. For

males, high body mass loss before and during mating was associated with a delays in male

immergence (for the same date of female immergence). In *Spermophilus citellus* (the European

ground squirrel), the most actively mating males delay the onset of post-mating accumulation of

body mass and also delay hibernation, presumably due to the long-term negative effects of

reproductive stress (Millesi et al. 1998). For females, the higher the specific reproductive effort,

the later the females immerged for the same date of male immergence. As for males, high

reproductive investment by females may induce stress that delays immergence. The duration of

maternal effort may also delay entry into hibernation, but its influence seems to result from an

extreme value (*Tachyglossus aculeatus,* maternal effort duration :167 day). Therefore, it was the

sex difference in the time and energy invested in reproduction which was associated with the order

of immergence. As for emergence, physiological constraints due to short growing season may

influence immergence. High precipitation seem to favor early immerge of males compared to

females. High precipitation may be associated with high water and food availability enable a fast

recovery from reproductive effort. However, female are constrained by maternal effort duration

(gestation and lactation) which may result in early immergence of males compared to females. On

the contrary, low precipitation may constrain recovery, and thus delay male immergence to a

greater extent than females in comparison with an environment with high precipitation (in both

environments, female are constrained by maternal care). In ectotherms, very few data are available

on sex differences in immergence date and therefore do not allow evaluation of this hypothesis.

Although the sex difference in dormancy phenology seems to be widely supported for a

trade-off between survival and reproduction, evidence exists at other scales of life. In mammals,

males and females that invest little or not at all in reproduction exhibit advances in energy reserve

accumulation and earlier immergence for up to several weeks, while reproductive congeners

continue activity (Neuhaus 2000, Millesi et al. 2008a). Another surprising example of this trade-

off is *Glis glis* (the edible dormouse), for which emergence became earlier with age. The authors

posited that, as younger individuals had a greater chance of reproducing in subsequent years, they

delayed their emergence for survival benefit at the expense of their immediate reproductive success

(Bieber et al. 2018). In ectotherms, several studies suggest that the benefits for reproduction

(Diamond et al. 2011, Navarro-Cano et al. 2015) and the benefits for survival such as avoiding

predators (Slusarczyk 1995, Kroon et al. 2008, Ji 2011) or intra- (Tougeron et al. 2018) and

interspecific competition (Dyugmedzhiev et al. 2019), influence dormancy phenology at the

species level. Thus, the life history hypothesis finds important support to explain the phenology of

dormancy at different scales (e.g. the individual scale, differences between sexes). Although the

energy limitation hypothesis seems to explain part of dormancy duration (Pianka 1970, Turbill and

Prior 2016, Wilsterman et al. 2021), it cannot, by itself, explain the sex differences in dormancy

phenology. In the next section, we examine how physiological constraints may be integrated into

the life history hypothesis framework, that is, the trade-off between reproduction and survival, to

explain the phenology of dormancy in endotherms and ectotherms.

404   b) Dormancy and physiological constraints

From an energetic point of view, it is favorable to remain active as long as the energy

balance is stable or positive, or conversely to remain in hibernation as long as the environment

does not allow for a positive energy balance. Any deviation from this principle may highlight a

balance rather in favor of reproduction (by sexual or non-sexual forms of natural selection) or

survival (Fig. 1) as expected by the life history hypothesis. Presumably, reducing the risk of

extrinsic mortality favors dormancy while the environment allows a positive energy balance. In

contrast, preparation for reproduction promotes activity while the environment does not allow a

positive energy balance (Snyder et al. 1961). Thus, a dormancy phenology staggered with respect

to the growing season (earlier emergence and immergence than expected) illustrates the selection

pressure exerted by the trade-off between reproduction (earlier emergence than expected) and adult

survival (earlier immergence than expected).

In our study, several elements might suggest that hibernation occurs even when

environmental conditions allow for a positive energy balance. Gains in body mass observed for

some individuals, even in species not known to hoard food, may indicate that the environment

allows a positive energy balance for other individuals with comparable energy demands. In several

species, females stay in hibernation (up to almost 2 months more) while males gain body mass of

up to 9% after emergence, or one sex immerges while the second continues to accumulate energy

reserves (Table 2). Sexual dimorphism may not be responsible for this sex differences as these

observations concern species with a sexual size dimorphism biased towards both males or females

(Table 4). To our knowledge, the only study that measured energy balance at the time of

immergence showed that *Tamias striatus* (the eastern chipmunk) immerged while a positive

energy balance could be maintained (Humphries et al. 2002). Observations of other species suggest

immergence when little food is available, but supposedly enough to support activity (Dobson et

al. 1992, Grigg and Beard 2000, Munro et al. 2008, Hoelzl et al. 2015). On the contrary, years of

low productivity can lead to later immergence (Alcorn 1940, O'Farrell et al. 1975), probably due

to a delay in the accumulation of reserves. This contradicts the view that hibernation duration

should necessarily increase with energetic constraints.

In ectotherms as well, some observations confirmed a dormancy phenology staggered with

respect to the growing season. In some reptiles and insects, individuals enter dormancy while

ambient temperature and food were still high enough to promote activity (Jameson Jr 1974,

Jameson and Allison 1976, Etheridge et al. 1983, Wiklund et al. 1996). In *Elaphe obsoleta* (the

black rat snake), part of the variation in emergence date was explained by the fact that smaller and

younger individuals emerged later than others (Blouin-Demers et al. 2000). This result is the

opposite of what is expected from a thermoregulation perspective, since small individuals should

reach their preferred temperature for activity more quickly (due to their low inertia) and should be

the first to emerge (Stevenson 1985). The authors proposed, on the contrary, that small individuals,

subjected to a higher predation rate in spring, benefited from increased survival while remaining

inactive. Males of *Vipera berus* (the common European adder) emerge before females in thermally

unfavorable periods, leading to significant body mass loss because of the reproductive benefits of

early emergence (Herczeg et al. 2007).

In the same way, the majority of insects enter dormancy long before environmental

conditions deteriorate, and remain dormant sometimes long after favorable conditions return

(Tauber and Tauber 1976, Koštál 2006). This strategy has been called "temporal conservative bet-

hedging" (Hopper 1999). Temporal bet-hedging strategies reduce fitness variation across the years

in a temporally fluctuating environment and supposedly result in higher average long-term fitness.

In this case, all individuals in a population (conservative because of low phenotypic variability)

reproduce only during the period that is always favorable through the years and avoid the period

with adverse conditions in some years at the expense of possible reproductive benefits in years

with favorable conditions. Temporal diversified bet-hedging exists in species for which the

duration of dormancy varies within a single cohort (diversified because of high phenotypic

variability) from one to several years (i.e. prolonged diapause), regardless of external conditions.

Thus, whatever the environmental conditions, a small proportion of the progeny will experience

optimal conditions to reproduce (Hopper 1999).

The independence of ectotherm dormancy towards harsh environmental conditions is in

contradiction with the vision of a passive inactivity induced by suboptimal temperature. Several

physiological and behavioral thermoregulation mechanisms may facilitate entering into dormancy

when ambient temperature above ground is still high. Indeed, some ectotherms enter dormancy in

summer (i.e., estivation or summer dormancy) and use deep burrows or crevices where the ambient

temperature is much colder. Thus, by exploiting their habitat, some ectotherms are able to reduce

their energy consumption (Pinder et al. 1992). On the other hand, some species are capable of an

active reduction in metabolism below that required under the simple passive effect of ambient

temperature on metabolism (Q10 effect) (Staples 2016). Ectotherm dormancy could therefore be

less temperature dependent than previously thought and would allow survival under a wider

spectrum of biotic and abiotic pressures.

c) Study limits

Our statistical analyses were based on a limited number of hibernators, and we have

accepted approximations concerning the phenology data for some species (e.g., date at which

first/last individuals of each sex were observed) due to the limited data in the literature. The

analysis included mainly Holarctic rodents except for *Erinaceus europaeus, Microcebus*

murinus and *Tachyglossus aculeatus.* Our analyses included two non-Holarctic species

(*Microcebus murinus* and *Tachyglossus aculeatus)*. Although they represent a very small minority

of hibernating species, the results obtained seem to be consistent with Holarctic species.

Hibernation in non-Holarctic species is supposed to have evolved in response to other

environmental factors than food shortage, such as water shortage (Bintz 1984; Nowack et al.,

2020). However, similar selection pressures may therefore exist and should encourage further

comparative research on hibernation in non-Holarctic and Holarctic species.

Bats were not included in this meta-analysis but represent an interesting model for

hibernation biology, because the sex difference in reproduction phenology is very different from

most hibernators. Thus, bats introduce varied and unique patterns to an understanding of

hibernation phenology (Willis 2017). In temperate bats, mating takes place just before hibernation

during "fall swarming" (Thomas et al. 1979). Females store sperm during winter and ovulation

takes place shortly after emergence (Buchanan 1987). In *Myotis lucifugus* (the little brown bat),

males immerge after females, likely increasing their mating opportunities and subsequently

recovering from body mass lost during mating (Norquay and Willis 2014). Female bats likely

emerged first because early parturition increased juvenile survival. The patterns observed were

consistent with the life-history hypothesis. Although few data are currently available, future

comparative studies between bat species should enhance our understanding of the life-history

hypothesis. *Tamias sibericus* was also excluded from the analyse of immergence. Chipmunks are

food storing hibernators and wait until autumn to fill the hibernaculum with acorns (from masting

in autumn) for winter feeding. Thus, reproductive effort may no longer have an impact on

immergence at this time, but rather males may wait until all females hibernate in order to confirm

the location of female burrows and increase mating success in the spring (Kawamichi 1996). Thus,

chipmunks may delay their immergence for reproductive benefits, in line with the life-history

hypothesis. Although the analyses performed in this study only included some of hibernators, the

available information suggested that the life history hypothesis may apply to the phenology of

other hibernators.

3) Conclusion

The sex difference in dormancy phenology observed in endotherms and ectotherms may be a widespread consequence of the trade-off between the benefits of being active for reproduction and the benefits of dormancy for survival (viz., the life-history hypothesis). Other non-exclusive and context-specific hypotheses have also been proposed (Morbey and Ydenberg 2001) and further studies are needed to test them. Energy constraints explain a part of dormancy phenology in both endotherms and ectotherms (Wilsterman et al. 2021), but evidence from our study shows some independence of energy balance at the specific times of emergence and immergence into hibernation. This study shows that physiological constraints must be integrated into the survival-reproduction trade-off of hibernation phenology to be meaningful. Thus, we expect that dormancy phenology will exhibit multiple evolutionary causes, especially when many species are studied and compared. The occurance of dormancy at high altitude and latitude where few or no energy resources are available over part of the year appears to be supported for the physiological constraints hypothesis (Ruf et al. 2012), although this hypothesis appears to be of limited or partial importance in explaining the phenology of the transition from dormancy to activity and *vice-versa*. Dormancy in energetically benign periods, but unfavorable for reproduction, may be more widespread than previously thought. Such research highlights the opportunities of studying dormancy across a broad spectrum of species (Wilsterman et al. 2021).

4) References

Abdi, H. 2007. Z-scores. Encyclopedia of measurement and statistics 3:1055–1058.

Alcorn, J. R. 1940. Life History Notes on the Piute Ground Squirrel. Journal of Mammalogy 21:160–170.

Allison, A. Z., C. J. Conway, and A. E. Morris. (n.d.). Why hibernate? Tests of four hypotheses to explain intraspecific variation in hibernation phenology. Functional Ecology.

Ancona, S., A. Liker, M. C. Carmona-Isunza, and T. Székely. 2020. Sex differences in age-to-maturation
relate to sexual selection and adult sex ratios in birds. Evolution letters 4:44–53.

Armitage, K. B. 1981. Sociality as a Life-History Tactic of Ground Squirrels. Oecologia 48:36–49.

Aydin, O., and M. Y. Yassikaya. 2021. Validity and reliability analysis of the PlotDigitizer software
program for data extraction from single-case graphs. Perspectives on Behavior Science:1–19.

Barnes, B. M., M. Kretzmann, I. Zucker, and P. Licht. 1988. Plasma androgen and gonadotropin levels
during hibernation and testicular maturation in golden-mantled ground squirrels. Biology of
reproduction 38:616–622.

Barto, K. 2020. MuMIn: multi-model inference, R package version 1.43. 17.

Bauwens, D., and R. F. Verheyen. 1985. The timing of reproduction in the lizard Lacerata vivipara:
differences between individual females. Journal of Herpetology:353–364.

Bieber, C., K. Lebl, G. Stalder, F. Geiser, and T. Ruf. 2014. Body mass dependent use of hibernation: why
not prolong the active season, if they can? Functional Ecology:167–177.

Bieber, C., and T. Ruf. 2004. Seasonal timing of reproduction and hibernation in the edible dormouse
(Glis glis). Life in the cold: Evolution, mechanism, adaptation, and application:113–125.

Bieber, C., and T. Ruf. 2009. Summer dormancy in edible dormice (Glis glis) without energetic
constraints. Naturwissenschaften 96:165–171.

Bieber, C., C. Turbill, and T. Ruf. 2018. Effects of aging on timing of hibernation and reproduction.
Scientific Reports 8:1–11.

Blath, J., and A. Tóbiás. 2020. Invasion and fixation of microbial dormancy traits under competitive
pressure. Stochastic Processes and their Applications 130:7363–7395.

Blouin-Demers, G., K. A. Prior, and P. J. Weatherhead. 2000. Patterns of variation in spring emergence by
black rat snakes (Elaphe obsoleta obsoleta). Herpetologica:175–188.

Breedveld, M. C., and P. S. Fitze. 2016. Experimental evidence that sperm maturation drives protandry in
an ectotherm. Oecologia 182:129–137.

Buchanan, G. D. 1987. Timing of ovulation and early embryonic development in Myotis lucifugus
(Chiroptera: Vespertilionidae) from northern central Ontario. American Journal of Anatomy 178:335–
340.

Constant, T., S. Giroud, V. A. Viblanc, M. L. Tissier, P. Bergeron, F. S. Dobson, and C. Habold. 2020.
Integrating Mortality Risk and the Adaptiveness of Hibernation. Frontiers in Physiology 11.

Danks, H. V. 1992. LONG LIFE CYCLES IN INSECTS. The Canadian Entomologist 124:167–187.

Diamond, S. E., A. M. Frame, R. A. Martin, and L. B. Buckley. 2011. Species' traits predict phenological
responses to climate change in butterflies. Ecology 92:1005–1012.

Dobson, F. S., M. J. Badry, and C. Geddes. 1992. Seasonal activity and body mass of Columbian ground
squirrels. Canadian Journal of Zoology 70:1364–1368.

Dobson, F. S., and D. E. Davis. 1986. Hibernation and Sociality in the California Ground Squirrel. Journal
of Mammalogy 67:416–421.

Dunford, C. 1974. Annual cycle of cliff chipmunks in the Santa Catalina Mountains, Arizona. Journal of
mammalogy 55:401–416.

Dyugmedzhiev, A., M. Slavchev, and B. Naumov. 2019. Emergence and dispersal of snakes after syntopic
hibernation. Herpetozoa 32:149.

Emlen, S. T., and L. W. Oring. 1977. Ecology, Sexual Selection, and the Evolution of Mating Systems.
Science 197:215–223.

Etheridge, K., L. C. Wit, and J. C. Sellers. 1983. Hibernation in the lizard Cnemidophorus sexlineatus
(Lacertilia: Teiidae). Copeia:206–214.

Friebe, A., A. L. Evans, J. M. Arnemo, S. Blanc, S. Brunberg, G. Fleissner, J. E. Swenson, and A. Zedrosser.
2014. Factors Affecting Date of Implantation, Parturition, and Den Entry Estimated from Activity and
Body Temperature in Free-Ranging Brown Bears. PLOS ONE 9:e101410.

Gashwiler, J. S. 1976. Biology of Townsend's chipmunks in western Oregon. The Murrelet 57:26–31.

Grigg, G., and L. Beard. 2000. Hibernation by Echidnas in Mild Climates: Hints about the Evolution of
Endothermy? Pages 5–19 in G. Heldmaier and M. Klingenspor, editors. Life in the Cold. Springer Berlin
Heidelberg.

Gunderson, A. R., and M. Leal. 2016. A conceptual framework for understanding thermal constraints on
ectotherm activity with implications for predicting responses to global change. Ecology Letters 19:111–
120.

Hairston Jr, N. G., and W. R. Munns Jr. 1984. The timing of copepod diapause as an evolutionarily stable
strategy. The American Naturalist 123:733–751.

Hayssen, V. 2008. Reproductive Effort in Squirrels: Ecological, Phylogenetic, Allometric, and Latitudinal
Patterns. Journal of Mammalogy 89:582–606.

Herczeg, G., J. SAARIKIVI, A. GONDA, J. PERÄLÄ, A. TUOMOLA, and J. MERILÄ. 2007. Suboptimal
thermoregulation in male adders (Vipera berus) after hibernation imposed by spermiogenesis. Biological
Journal of the Linnean Society 92:19–27.

Hibbitts, T. J., W. E. Cooper, and M. J. Whiting. 2012. Spatial distribution and activity patterns in African
barking geckos: implications for mating system and reproduction. Journal of Herpetology 46:456–460.

Hijmans, R. J., S. E. Cameron, J. L. Parra, P. G. Jones, and A. Jarvis. 2005. Very high resolution
interpolated climate surfaces for global land areas. International Journal of Climatology: A Journal of the
Royal Meteorological Society 25:1965–1978.

Hoehler, T. M., and B. B. Jørgensen. 2013. Microbial life under extreme energy limitation. Nature
Reviews Microbiology 11:83–94.

Hoelzl, F., C. Bieber, J. S. Cornils, H. Gerritsmann, G. L. Stalder, C. Walzer, and T. Ruf. 2015. How to spend
the summer? Free-living dormice (Glis glis) can hibernate for 11 months in non-reproductive years.
Journal of Comparative Physiology B 185:931–939.

Holekamp, K. E., S. Nunes, and F. Talamantes. 1988. Circulating prolactin in free-living California ground
squirrels (Spermophilus beecheyi). General and comparative endocrinology 71:484–492.

Hopper, K. R. 1999. Risk-spreading and bet-hedging in insect population biology. Annual review of
entomology 44:535–560.

Humphries, M. M., D. W. Thomas, C. L. Hall, J. R. Speakman, and D. L. Kramer. 2002. The energetics of
autumn mast hoarding in eastern chipmunks. Oecologia 133:30–37.

Huwaldt, J. A., and S. Steinhorst. 2015. Plot Digitizer, version 2.6. 8. Software.

Jameson, E. W., and A. Allison. 1976. Fat and Breeding Cycles in Two Montane Populations of Sceloporus
occidentalis (Reptilia, Lacertilia, Iguanidae). Journal of Herpetology 10:211.

Jameson Jr, E. W. 1974. Fat and breeding cycles in a montane population of Sceloporus graciosus.
Journal of Herpetology:311–322.

Ji, R. 2011. Calanus finmarchicus diapause initiation: new view from traditional life history-based model.
Marine Ecology Progress Series 440:105–114.

Jones, K. E., J. Bielby, M. Cardillo, S. A. Fritz, J. O'Dell, C. D. L. Orme, K. Safi, W. Sechrest, E. H. Boakes, C.
Carbone, C. Connolly, M. J. Cutts, J. K. Foster, R. Grenyer, M. Habib, C. A. Plaster, S. A. Price, E. A. Rigby,
J. Rist, A. Teacher, O. R. P. Bininda-Emonds, J. L. Gittleman, G. M. Mace, and A. Purvis. 2009. PanTHERIA:
a species-level database of life history, ecology, and geography of extant and recently extinct mammals.
Ecology 90:2648–2648.

Joschinski, J., and D. Bonte. 2021. Diapause and bet-hedging strategies in insects: a meta-analysis of
reaction norm shapes. Oikos.

Kawamichi, M. 1996. Ecological Factors Affecting Annual Variation in Commencement of Hibernation in
Wild Chipmunks (Tamias sibiricus). Journal of Mammalogy 77:731–744.

Kenagy, G. J., S. M. Sharbaugh, and K. A. Nagy. 1989. Annual cycle of energy and time expenditure in a
golden-mantled ground squirrel population. Oecologia 78:269–282.

Kortessis, N., and P. Chesson. 2019. Germination variation facilitates the evolution of seed dormancy
when coupled with seedling competition. Theoretical Population Biology 130:60–73.

Koštál, V. 2006. Eco-physiological phases of insect diapause. Journal of insect physiology 52:113–127.

Kroon, A., R. L. Veenendaal, J. Bruin, M. Egas, and M. W. Sabelis. 2008. "Sleeping with the enemy"—
predator-induced diapause in a mite. Naturwissenschaften 95:1195–1198.

Levesque, D. L., O. M. A. Lovasoa, S. N. Rakotoharimalala, and B. G. Lovegrove. 2013. High mortality and
annual fecundity in a free-ranging basal placental mammal, Setifer setosus (Tenrecidae: Afrosoricida).
Journal of Zoology 291:205–212.

Magalhães, J. P. D., and J. Costa. 2009. A database of vertebrate longevity records and their relation to
other life-history traits. Journal of Evolutionary Biology 22:1770–1774.

Manno, T. G., and F. S. Dobson. 2008. Why are male Columbian ground squirrels territorial? Ethology
114:1049–1060.

Michener, G. R. 1983. Spring emergence schedules and vernal behavior of Richardson's ground squirrels:
why do males emerge from hibernation before females? Behavioral Ecology and Sociobiology 14:29–38.

Michener, G. R. 1984. Age, sex, and species differences in the annual cycles of ground-dwelling sciurids:
implications for sociality.

Michener, G. R. 1992. Sexual differences in over-winter torpor patterns of Richardson's ground squirrels
in natural hibernacula. Oecologia 89:397–406.

Millesi, E., A. Divjak, and A. Strauss. 2008a. Seasonal timing and pre-hibernation fattening in breeding
and non-breeding European Ground Squirrels. In: Lovegrove BG, McKechnie AE (eds) Hypometabo-lism
in animals: hibernation, torpor and cryobiology. University of KwaZulu-Natal, Pietermaritzburg,:p 289-
294.

Millesi, E., S. Huber, J. Dittami, I. Hoffmann, and S. Daan. 1998. Parameters of Mating Effort and Success
in Male European Ground Squirrels, Spermophilus citellus. Ethology 104:298–313.

Millesi, E., A. Strauss, T. Burger, I. E. Hoffmann, and M. Walzl. 2008b. Follicular development in European
ground squirrels (Spermophilus citellus) in different phases of the annual cycle. Reproduction 136:205–
210.

Morbey, Y. E., and R. C. Ydenberg. 2001. Protandrous arrival timing to breeding areas: a review. Ecology
Letters 4:663–673.

Mouhoub sayah, C., J.-P. Robin, P. Pévet, S. Monecke, S. Doumandji, and M. Saboureau. 2009. Road
mortality of the Algerian hedgehog (Atelerix algirus) in the Soummam Valley (Algeria). Revue d'écologie.

Munro, D., D. W. Thomas, and M. M. Humphries. 2008. Extreme suppression of aboveground activity by
a food-storing hibernator, the eastern chipmunk (Tamias striatus). Canadian Journal of Zoology 86:364–
370.

Munroe, K. E. 2011. The Socioecology, Mating System and Behavior of Round-Tailed Ground Squirrels
(Xerospermophilus tereticaudus). Ph.D. dissertation, University of Arizona, Tucson, AZ.

Navarro-Cano, J. A., B. Karlsson, D. Posledovich, T. Toftegaard, C. Wiklund, J. Ehrlén, and K. Gotthard.
2015. Climate change, phenology, and butterfly host plant utilization. AMBIO 44:78–88.

Neuhaus, P. 2000. Timing of Hibernation and Molt in Female Columbian Ground Squirrels. Journal of
Mammalogy 81:571–577.

Norquay, K. J. O., and C. K. R. Willis. 2014. Hibernation phenology of M yotis lucifugus. Journal of
Zoology 294:85–92.

O'Farrell, T. P., R. J. Olson, R. O. Gilbert, and J. D. Hedlund. 1975. A Population of Great Basin Pocket
Mice, Perognathus parvus, in the Shrub-Steppe of South-Central Washington. Ecological Monographs
45:1–28.

Olsson, M., T. Birkead, and R. Shine. 1999. Can relaxed time constraints on sperm production eliminate
protandry in an ectotherm? Biological Journal of the Linnean Society 66:159–170.

Orme, D., R. Freckleton, G. Thomas, T. Petzoldt, and S. Fritz. 2013. The caper package: comparative
analysis of phylogenetics and evolution in R. R package version 5:1–36.

Paradis, E. 2011. Analysis of Phylogenetics and Evolution with R. Springer Science & Business Media.

Paradis, E., and K. Schliep. 2019. ape 5.0: an environment for modern phylogenetics and evolutionary
analyses in R. Bioinformatics 35:526–528.

Pianka, E. R. 1970. Comparative Autecology of the Lizard Cnemidophorus Tigris in Different Parts of Its
Georgraphic Range. Ecology 51:703–720.

Pinder, A. W., K. B. Storey, and G. R. Ultsch. 1992. Estivation and Hibernation. Environmental physiology
of the amphibians:250.

Randrianambinina, B., D. Rakotondravony, U. Radespiel, and E. Zimmermann. 2003. Seasonal changes in
general activity, body mass and reproduction of two small nocturnal primates: a comparison of the
golden brown mouse lemur (Microcebus ravelobensis) in Northwestern Madagascar and the brown
mouse lemur (Microcebus rufus) in Eastern Madagascar. Primates 44:321–331.

Ruf, T., and C. Bieber. 2022. Why hibernate? Predator avoidance in the edible dormouse. Mammal
Research:1–11.

Ruf, T., C. Bieber, and C. Turbill. 2012. Survival, Aging, and Life-History Tactics in Mammalian
Hibernators. Pages 123–132 in T. Ruf, C. Bieber, W. Arnold, and E. Millesi, editors. Living in a Seasonal
World: Thermoregulatory and Metabolic Adaptations. Springer Berlin Heidelberg, Berlin, Heidelberg.

Sandell, M. 1990. The Evolution of Seasonal Delayed Implantation. The Quarterly Review of Biology
65:23–42.

Satterthwaite, W. H. 2010. Competition for space can drive the evolution of dormancy in a temporally
invariant environment. Plant Ecology 208:167–185.

Siutz, C., C. Franceschini, and E. Millesi. 2016. Sex and age differences in hibernation patterns of
common hamsters: adult females hibernate for shorter periods than males. Journal of Comparative
Physiology B 186:801–811.

Slusarczyk, M. 1995. Predator-induced diapause in Daphnia. Ecology 76:1008–1013.

Snyder, R. L., D. E. Davis, and J. J. Christian. 1961. Seasonal Changes in the Weights of Woodchucks.
Journal of Mammalogy 42:297–312.

Staples, J. F. 2016. Metabolic Flexibility: Hibernation, Torpor, and Estivation. Pages 737–771
Comprehensive Physiology. American Cancer Society.

Stevenson, R. D. 1985. Body Size and Limits to the Daily Range of Body Temperature in Terrestrial
Ectotherms. The American Naturalist 125:102–117.

Tauber, M. J., and C. A. Tauber. 1976. Insect seasonality: diapause maintenance, termination, and
postdiapause development. Annual review of entomology 21:81–107.

Thomas, D. W., M. B. Fenton, and R. M. Barclay. 1979. Social behavior of the little brown bat, Myotis
lucifugus: I. Mating behavior. Behavioral Ecology and Sociobiology:129–136.

Thompson, M. J., F. S. Dobson, D. W. Coltman, J. O. Murie, S. Raveh, and J. E. Lane. 2023. Sexes in sync:
phenotypic plasticity, sexual selection and phenological synchrony between the sexes in a wild
hibernator. Proceedings of the Royal Society B: Biological Sciences 290:20231113.

Tougeron, K., G. Hraoui, C. Le Lann, J. van Baaren, and J. Brodeur. 2018. Intraspecific maternal
competition induces summer diapause in insect parasitoids. Insect science 25:1080–1088.

Turbill, C., C. Bieber, and T. Ruf. 2011. Hibernation is associated with increased survival and the
evolution of slow life histories among mammals. Proceedings of the Royal Society B: Biological Sciences
278:3355–3363.

Turbill, C., and S. Prior. 2016. Thermal climate-linked variation in annual survival rate of hibernating
rodents: shorter winter dormancy and lower survival in warmer climates. Functional Ecology 30:1366–
1372.

Upham, N. S., J. A. Esselstyn, and W. Jetz. 2019. Inferring the mammal tree: Species-level sets of
phylogenies for questions in ecology, evolution, and conservation. PLoS Biology 17.

Vander Wall, S. B. 1990. Food hoarding in animals. University of Chicago Press, Chicago, Illinois, USA.

Wartel, A., P. Lindenfors, and J. Lind. 2019. Whatever you want: Inconsistent results are the rule, not the
exception, in the study of primate brain evolution. PloS one 14:e0218655.

Watts, J. C., and B. Tenhumberg. 2021. Optimal resource allocation and prolonged dormancy strategies
in herbaceous plants. Journal of Ecology 109:218–233.

Wiklund, C., V. Lindfors, and J. Forsberg. 1996. Early male emergence and reproductive phenology of the
adult overwintering butterfly Gonepteryx rhamni in Sweden. Oikos:227–240.

Williams, C. T., B. M. Barnes, G. J. Kenagy, and C. L. Buck. 2014. Phenology of hibernation and
reproduction in ground squirrels: integration of environmental cues with endogenous programming.
Journal of Zoology 292:112–124.

Willis, C. K. R. 2017. Trade-offs Influencing the Physiological Ecology of Hibernation in Temperate-Zone
Bats. Integrative and Comparative Biology 57:1214–1224.

Wilsterman, K., M. A. Ballinger, and C. M. Williams. 2021. A unifying, eco-physiological framework for
animal dormancy. Functional Ecology 35:11–31.

Woods, C. P., and R. M. Brigham. 2004. The avian enigma:'"hibernation"'by common poorwills
(Phalaenoptilus nuttallii). Pages 129–138 Life in the cold: evolution, mechanisms, adaptation, and
application. Twelfth International Hibernation Symposium. Institute of Arctic Biology, University of
Alaska Fairbanks, USA.

Woods, C. P., Z. J. Czenze, and R. M. Brigham. 2019. The avian "hibernation" enigma: thermoregulatory
patterns and roost choice of the common poorwill. Oecologia 189:47–53.

Figure 1 Schematic representation of "the physiological constraint hypothesis" and "the life-

history hypothesis" and their predictions. Physiological constraints in the environment refers to

variation over time of the energetic balance (mainly for heterothermic endotherms) and the

thermal window favorable to activity (mainly for ectotherms). Hypothesis H1 assumes that

dormancy phenology occurs at the time of transition between favorable and unfavorable

energetic or thermal conditions or *vice versa*. It predicts that the sex difference in dormancy

phenology is explained by sex differences in physiological constraints, and that reproductive

investment should be independent of this sex difference in phenology. In contrast, hypothesis

H2 predicts that a phenology that would occur before or after the energetic transition may be

associated with benefits to survival or reproduction. It predicts that the sex difference in

dormancy phenology is associated with a sex difference in reproductive investment. This pattern

is expected for species without paternal effort. But the concept can be applied to other types of

mating strategies. The hibernation phenology presented for prediction (H1) are those expected

for hibernating mammals. Note that the magnitude and order of sex differences in phenology is

not an expected general trend, because it is assumed to vary between species according to

beneficial envioronmental conditions for energy demand (prediction H1) and reproductive

investment (prediction H2). Nevertheless, the sex difference is assumed to be smaller with the

H1 prediction because there is less sex difference in energy demand than in reproductive

investment. Black, grey and dark blue horizontal arrows represent respectively time over the

year, reproductive investment in males and reproductive investment in females. Black, grey and

dark blue horizontal arrows represent respectively time over the year, reproductive investment

in males and reproductive investment in females. Red and orange triangles represent

respectively immergence and emergence timing and black cross represent mating timing.

782   Figure 2 World map showing the study sites of the hibernating species studied. Green dots indicate

783   the location of study sites. A red gradient represents the variation in minimum ambient

784   temperature, a parameter used in this study.

Figure 3 Effects of body mass variation during mating on protandry. The minimum temperatures

of the study sites are indicated by a color gradient with the warmest temperatures in red. The

regression line in black indicates the effect of body mass variation during mating on protandry.

Body mass variation during mating was represented as a percentage of body mass before mating.

Figure 4 Effects of body mass variation through the end of mating on sex differences in immergence. Annual precipitation at the study sites are indicated by a color gradient with the highest precipitation in dark blue. The regression line in black indicates the effect of body mass variation through the end of mating on sex difference in immergence. Body mass variation through the end of mating was represented as a percentage of body mass at emergence. Males immerged earlier in species living in an environment with high precipitation (dark blue dots).

Figure 5 Effects of female specific reproductive effort on sex difference in immergence. The annual precipitation of the study sites are indicated by a color gradient with the highest precipitation in dark blue. The regression line in black indicates the effect of female specific reproductive effort on sex difference in immergence. Males immerged earlier in species living in an environment with high precipitation (dark blue dots).

Table 1 Regression results for the best models explaining variation in protandry and sex difference in immergence. The Z standardized model estimates and the phylogenetic effect are reciprocally estimated by β and γ_M. The abbreviations "diff" and "rel" "Min" stand respectively for "difference", "relative" and "Minimum". A negative value for the sex difference in immergence indicates that males immerge before females and a positive value indicates that females immerge before males. Relative testes mass, Δ body mass before mating, Δ body mass during mating, Δ body mass through the end of mating was represented respectively as a percentage of body mass, body mass at emergence, body mass before mating and body mass at emergence.

	R^2	γ_{ML}	Dependent variable	Independent variable	β ± SE	t-value	p- value
				γ_{ML}			
Model 1 (28 species)	0.59	0.00 (NA, 0.69)	Protandry	Intercept	16.31 ± 1.27	12.76	< 0.001***
				z- Male body mass at emergence	3.25 ± 1.37	2.37	0.02*
				z-Δ body mass during mating	-3.32 ± 1.39	-2.38	0.02*
				z-Min temperature	5.94 ± 1.36	4.34	< 0.001***
				z- late mating	-2.58 ± 1.41	-1.83	0.08 .
Model 2 (21 species)	0.50	0.00 (NA, 0.38)	Sex diff in immergence	Intercept	-5.18 ± 2.18	-2.36	0.03*
				z- log specific reproductive effort	-13.77 ± 3.88	-3.54	0.002**
				z-Δ body mass through the end of mating	-9.07 ± 2.37	-3.81	0.001**
				z- precipitation	-7.67 ± 2.59	-2.95	0.009**
				z- Maternal effort duration	-12.66 ± 3.58	-3.52	0.002**
Model 3 (20 species)	0.63	0.00 (NA, 0.73)	Sex diff in immergence	Intercept	-4.24 ± 1.89	-2.23	0.04*
				z- log specific reproductive effort	-10.20 ± 2.34	-4.35	< 0.001***
				z-Δ body mass through the end of mating	-7.05 ± 2.05	-3.42	0.003**
				z- precipitation	-12.48 ± 2.70	-4.61	< 0.001***
				z-dimorphism at immergence	-4.75 ± 2.42	-1.96	0.06 .

Table 2 Species with dimorphisms biased in favor of males or females and their body mass gain during the year. Body size dimorphism is calculated as male body size divided by female body size. See section "Sex differences in reproductive investment" for the determination of the body mass gain before mating (See supplementary materials S2 for references).

Species	Body size dimorphism	Male body mass gain before mating (% of emergence body mass)	The end of reserve accumulation before hibernation for females
Cricetus cricetus	1.14	9.35	27 days after male
Cynomys leucurus	1.04	4.89	11 days after male
Glis glis	0.97	6.63	14 days after male
Microcebus murinus	0.96	9.01	Same time as male
Urocitellus parryii	0.97	0.49	35 days before male

Supplementary materials S1

Materials and methods

  a) Sex difference in hibernation phenology

The search criteria were based on combining the following terms: (scientific OR common names of species) AND (phenology OR annual cycle OR hibernation). Because of their imprecision, we excluded the studies for which hibernation season phenology was deduced from the presence of active individuals on a monthly basis. Because of their imprecision, we excluded the studies for which hibernation season phenology was deduced from the presence of active individuals on a monthly basis. This excluded four studies [3–6]. As the data were averaged for each species (see section "Statistics") we did not use data with exceptional variation between years within the same study site. This excluded data from [7] on the sex difference in immergence date (55 days difference between the two years) for *Xerospermophilus tereticaudus* (the round-tailed ground squirrel). *Otospermophilus beecheyii* (the California ground squirrel) appeared to be a species with great variation in hibernation phenology and whether males and females hibernated [8,9]. These data were therefore not included in this study.

  b) Sex differences in reproductive investment

For all data on changes in body mass, the search was conducted by combining the following terms: (scientific OR common names of species) AND (body mass change OR annual body mass). To be as accurate as possible, we have obtained data only when measured at the same or nearby the study site that was used for hibernation data. In cases where information

were not directly available in the text or table, we used the software Plot Digitizer [10] to extract the data from graphs. This software has recently been validated for this use [11]. The start and end dates of mating were estimated from information available in the text or from other studies at the same study site. When the mating period could not be clearly determined, studies were omitted from analyses.

For the analyses, we give priority to reproductive and body mass data from the same or very close study sites as the hibernation phenology data. We made a few exceptions for juvenile body mass at weaning, for which some data were obtained in Hayssen (2008), male reproductive effort for *Spermophilus brunneus* and *Zapus princeps* and the length of gestation and lactation. Maternal effort duration is calculated as the sum of the gestation and lactation periods. We obtained data on the length of gestation and lactation from the AnAge database (The Animal Aging and Longevity Database; Magalhães and Costa, 2009), and complemented these data with a specific search combining the following terms: (scientific OR common names of species) AND (lactation duration OR gestation duration). As maternal effort duration and lactation data were not available in the literature for *Urocitellus brunneus* and *Ictidomys parvidens* respectively, we used averages for the clade *Marmotini* obtained from Hayssen (2008). The body mass at weaning for *Urocitellus brunneus* correpsonds to the body mass at weaning of *Urocitellus townsendii* (phylogenetically close species) corrected for the difference in body mass of the females from Hayssen (2008) as body mass accounts for 76–84% of the variation in weaning mass in the clade *Marmotini* (Hayssen, 2008)

c) Statistics

For each PGLS model tested we downloaded 100 phylogenetic mammalian trees (see Upham et al., 2019). Then, strict consensus trees for which the included clades were those present in all the 100 phylogenetic mammalian trees were constructed [15]. For each consensus trees (see Figure S1), branch lengths were calculated with the "compute.brlen" function from

the "ape" package based on Grafen's (1989) method, and were used to compute PGLS models with the "caper" package in R.

Figure S1. Consensus phylogenetic trees for the species under study: (a) model 1 (b) model 2 and (c) model 3. Each consensus tree was built from 100 trees obtained from http://vertlife.org/phylosubsets/. Branch lengths were calculated using Grafen's computations with the 'ape' package in R (see Materials and Methods).

References

1.   Sandell, M. (1990). The Evolution of Seasonal Delayed Implantation. Q. Rev. Biol. *65*, 23–42.
10.1086/416583.

2.   Friebe, A., Evans, A.L., Arnemo, J.M., Blanc, S., Brunberg, S., Fleissner, G., Swenson, J.E., and Zedrosser, A.
(2014). Factors Affecting Date of Implantation, Parturition, and Den Entry Estimated from Activity and Body
Temperature in Free-Ranging Brown Bears. PLOS ONE *9*, e101410. 10.1371/journal.pone.0101410.

3.   Dunford, C. (1974). Annual cycle of cliff chipmunks in the Santa Catalina Mountains, Arizona. J. Mammal.
55, 401–416.

4.   Gashwiler, J.S. (1976). Biology of Townsend's chipmunks in western Oregon. The Murrelet *57*, 26–31.

5.   Mouhoub sayah, C., Robin, J.-P., Pévet, P., Monecke, S., Doumandji, S., and Saboureau, M. (2009). Road
mortality of the Algerian hedgehog (Atelerix algirus) in the Soummam Valley (Algeria). Rev. Décologie.

6.   Randrianambinina, B., Rakotondravony, D., Radespiel, U., and Zimmermann, E. (2003). Seasonal changes
in general activity, body mass and reproduction of two small nocturnal primates: a comparison of the golden
brown mouse lemur (Microcebus ravelobensis) in Northwestern Madagascar and the brown mouse lemur
(Microcebus rufus) in Eastern Madagascar. Primates *44*, 321–331.

7.   Munroe, K.E. (2011). The Socioecology, Mating System and Behavior of Round-Tailed Ground Squirrels
(Xerospermophilus tereticaudus).

8.   Dobson, F.S., and Davis, D.E. (1986). Hibernation and Sociality in the California Ground Squirrel. J.
Mammal. *67*, 416–421. 10.2307/1380902.

9.   Holekamp, K.E., Nunes, S., and Talamantes, F. (1988). Circulating prolactin in free-living California ground
squirrels (Spermophilus beecheyi). Gen. Comp. Endocrinol. *71*, 484–492.

10.   Huwaldt, J.A., and Steinhorst, S. (2015). Plot Digitizer, version 2.6. 8 (Software).

11.   Aydin, O., and Yassikaya, M.Y. (2021). Validity and reliability analysis of the PlotDigitizer software program
for data extraction from single-case graphs. Perspect. Behav. Sci., 1–19.

12.   Magalhães, J.P.D., and Costa, J. (2009). A database of vertebrate longevity records and their relation to
other life-history traits. J. Evol. Biol. *22*, 1770–1774. 10.1111/j.1420-9101.2009.01783.x.

13.   Jones, K.E., Bielby, J., Cardillo, M., Fritz, S.A., O'Dell, J., Orme, C.D.L., Safi, K., Sechrest, W., Boakes, E.H.,
Carbone, C., et al. (2009). PanTHERIA: a species-level database of life history, ecology, and geography of extant and
recently extinct mammals. Ecology *90*, 2648–2648. 10.1890/08-1494.1.

14.   Upham, N.S., Esselstyn, J.A., and Jetz, W. (2019). Inferring the mammal tree: Species-level sets of
phylogenies for questions in ecology, evolution, and conservation. PLoS Biol. *17*. 10.1371/journal.pbio.3000494.

15.   Paradis, E. (2011). Analysis of Phylogenetics and Evolution with R (Springer Science & Business Media).

1035

Supplementary materials S2

1037

1038

1039   Table S1 Data on dependent and independent factors used in models 1. Protandry was used as the dependent factor and Body mass

1040   change during mating, late mating, body mass male, strategy fatstoring or foodstoring, minimum temperature, precipitation and

1041   dimorphism of body mass at emergence were considered as independent factors. Protandry was calculated as follows: Female Julian

1042   date – male Julian date. The exact hibernation phenology data for *Cricetus cricetus* have been confirmed by the authors. See materials

1043   and methods for the acquisition of minimum temperature data.

1044

1045

1046

1047

1048

1049

1050

1051

1052

1053

species	Late mating	Food/Fat	protandry	Body mass change during mating	Body mass male	Dimorphism body mass emergence	precipitation	Min temperature
Callospermophilus lateralis	1[1]	Fat	21[1]	-11,126[2]	173[2]	1,26[2]	411	-7
Callospermophilus saturatus	1[3]	Food	4[3]	-6,38[3]	248[3]	1,45[3]	863	-7,3
Chaetodipus formosus	4[4]	Food	14[4]	0,8[4]	18,6[4]	1,12[4]	863	-7,3
Cricetus cricetus	2,5[5,6]	Food	20[7]	-5,6[8]	400,9[8]	1,58[6,8]	595	-2,7
Cynomys leucurus	1[9]	Fat	21[9,10]	-21,61[9]	829,4[9]	1,69[9]	337	-11,5
Erinaceus europaeus	1[11,12]	Fat	17,6[12,13]	-12,5[12,13]	1028,67[12,13]	1,32[12,13]	803	-5,3
Glis glis	1[14]	Fat	30[15]	-8,6[14]	105,5[14]	1,1[14]	758	-2,5
Ictidomys parvidens	6[16]	Fat	7,5[16]	-10,5[16]	192[16]	1,27[16]	347	-2,4
Marmota monax	1[17]	Fat	22[18]	-6,2[17]	2701[17]	1,03[17]	1156	-11,5
Microcebus murinus	1[19]	Fat	48[19]	-20,7[20]	62[20]	1,02[20]	922	14,6
Perognathus longimembris	4[4]	Food	14[21]	3,2[21]	7[21]	1,06[21]	863	-7,3
Poliocitellus franklinii	1[22,23]	Fat	11[22,23]	-2,62[22,23]	417,1[22,23]	1,27[22,23]	517	-21,1
Spermophilus citellus	1[24]	Fat	23[25]	-9,8[25]	316,26[25]	1,68[25]	556	-2,5
Spermophilus xanthoprymnus	1[24]	Fat	7,5[24]	-11,62[24]	292[24]	1,69[24]	407	-5,2
Tachyglossus aculeatus	1[26]	Fat	45[26]	-12,7[27]	3950[27]	1,06[27]	625	3,6
Tamias amoenus	1[21]	Food	10,5[21]	-5,4[21]	42,3[21]	0,91[21]	846	-7,3
Tamias sibiricus	1[28]	Food	20[28]	1,5[29]	90,6[29]	0,96[29]	942	-9,4
Urocitellus armatus	1[24]	Fat	3,66[30]	6,82[30]	333,7[30]	1,25[30]	402	-13
Urocitellus beldingi	1[24]	Fat	9[31]	-0,4[31]	240,1[31]	1,14[31]	568	-12,4
Urocitellus brunneus	1[32]	Fat	9[32]	-6,57[33]	175[33]	1,44[33]	610	-11
Urocitellus columbianus	1[24]	Fat	7[34]	-0,4[34]	430[34]	1,18[34]	346	-17,5
Urocitellus elegans	1[35]	Fat	17,3[35]	-3,8[35]	249,3[35]	1,37[35]	281	-16,6
Urocitellus mollis	1[36]	Fat	15[36]	4,17[37]	181[37]	1,49[37]	248	-5,4
Urocitellus parryii	1[24]	Food	14[38]	-21,6[39]	941[39]	1,58[39]	237	-30,2
Urocitellus richardsonii	1[24]	Food	16[40–43]	-10[41,43]	392,5[41,43]	1,7[41,43]	371,5	-14,4
Xerospermophilus tereticaudus	4[44]	Fat	5,3[44]	4,2[44]	142[44]	1,27[44]	250	2,4
Zapus hudsonius	1[45]	Fat	14[38]	-2,67[45]	17[45]	1,02[45]	837	-8,5

1054

1055

1056

| Zapus princeps | 1[46] | Fat | 10,33[46] | -7,91[47] | 22,48[47] | 0,95[47] | 570 | -14,1 |

1057

1058

1. Bronson, M. T. Altitudinal variation in emergence time of golden-mantled ground squirrels (Spermophilus lateralis). *Journal of Mammalogy* **61**, 124–126 (1980).

2. McKeever, S. The biology of the golden-mantled ground squirrel, Citellus lateralis. *Ecological Monographs* **34**, 383–401 (1964).

3. Kenagy, G. J., Sharbaugh, S. M. & Nagy, K. A. Annual cycle of energy and time expenditure in a golden-mantled ground squirrel population. *Oecologia* **78**, 269–282 (1989).

4. Kenagy, G. J. & Bartholomew, G. A. Seasonal Reproductive Patterns in Five Coexisting California Desert Rodent Species: Ecological Archives M055-002. *Ecological Monographs* **55**, 371–397 (1985).

5. Siutz, C., Valent, M., Ammann, V., Niebauer, A. & Millesi, E. Sex-specific effects of food supplementation on hibernation performance and reproductive timing in free-ranging common hamsters. *Sci Rep* **8**, 1–10 (2018).

6. Hufnagl, S., Franceschini-Zink, C. & Millesi, E. Seasonal constraints and reproductive performance in female Common hamsters (Cricetus cricetus). *Mammalian Biology* **76**, 124–128 (2011).

7. Siutz, C., Franceschini, C. & Millesi, E. Sex and age differences in hibernation patterns of common hamsters: adult females hibernate for shorter periods than males. *J Comp Physiol B* **186**, 801–811 (2016).

8. Lebl, K. & Millesi, E. Yearling male Common hamsters and the trade-off between growth and reproduction. *Biosystematics and Ecology Series* **25**, 115–126 (2008).

9. Bakko, E. B. & Brown, L. N. Breeding Biology of The White-Tailed Prairie Dog, Cynomys Leucurus, in Wyoming. *Journal of Mammalogy* **48**, 100–112 (1967).

10. Clark, T. W. Ecology and ethology of the white-tailed prairie dog (Cynomys leucurus). *Milwaukee Public Museum Publications in Biology and Geology* **3**, 1–96 (1977).

11. Rautio, A., Valtonen, A. & Kunnasranta, M. The effects of sex and season on home range in European hedgehogs at the northern edge of the species range. in *Annales Zoologici Fennici* vol. 50 107–123 (BioOne, 2013).

12. Haigh, A., O'Riordan, R. M. & Butler, F. Nesting behaviour and seasonal body mass changes in a rural Irish population of the Western hedgehog
(Erinaceus europaeus). *Acta Theriologica* **57**, 321–331 (2012).

13. Rautio, A., Valtonen, A., Auttila, M. & Kunnasranta, M. Nesting patterns of European hedgehogs (Erinaceus europaeus) under northern conditions.
Acta theriologica **59**, 173–181 (2014).

14. Bieber, C. Population dynamics, sexual activity, and reproduction failure in the fat dormouse (Myoxus glis). *Journal of Zoology* **244**, 223–229 (1998).

15. Bieber, C. & Ruf, T. Seasonal timing of reproduction and hibernation in the edible dormouse (Glis glis). *Life in the cold: Evolution, mechanism,*
adaptation, and application 113–125 (2004).

16. Schwanz, L. E. Annual cycle of activity, reproduction, and body mass in Mexican ground squirrels (Spermophilus mexicanus). *Journal of Mammalogy*
87, 1086–1095 (2006).

17. Maher, C. R. & Duron, M. Mating system and paternity in woodchucks (Marmota monax). *Journal of Mammalogy* **91**, 628–635 (2010).

18. Maher, C. R. Social organization in woodchucks (Marmota monax) and its relationship to growing season. *Ethology* **112**, 313–324 (2006).

19. Schmid, J. Sex-specific differences in activity patterns and fattening in the gray mouse lemur (Microcebus murinus) in Madagascar. *Journal of*
Mammalogy **80**, 749–757 (1999).

20. Schmid, J. & Kappeler, P. M. Fluctuating sexual dimorphism and differential hibernation by sex in a primate, the gray mouse lemur (Microcebus
murinus). *Behavioral Ecology and Sociobiology* **43**, 125–132 (1998).

21. Kenagy, G. J. & Barnes, B. M. Seasonal Reproductive Patterns in Four Coexisting Rodent Species from the Cascade Mountains, Washington. *J Mammal*
69, 274–292 (1988).

22. Iverson, S. L. & Turner, B. N. Natural history of a Manitoba population of Franklin's ground squirrels. *Canadian Field-Naturalist* **86**, 145–149 (1972).

23. Choromanski-Norris, J., Fritzell, E. K. & Sargeant, A. B. Seasonal Activity Cycle and Weight Changes of the Franklin's Ground Squirrel. *American Midland*
Naturalist **116**, 101 (1986).

24. Gür, H. & Gür, M. K. Annual cycle of activity, reproduction, and body mass of Anatolian ground squirrels (Spermophilus xanthoprymnus) in Turkey.
Journal of Mammalogy **86**, 7–14 (2005).

25. Millesi, E., Strijkstra, A. M., Hoffmann, I. E., Dittami, J. P. & Daan, S. Sex and Age Differences in Mass, Morphology, and Annual Cycle in European
Ground Squirrels, Spermophilus citellus. *J Mammal* **80**, 218–231 (1999).

26.   Nicol, S. C., Morrow, G. E. & Harris, R. L. Energetics meets sexual conflict: The phenology of hibernation in Tasmanian echidnas. *Functional Ecology* **33**,
2150–2160 (2019).

27.   Nicol, S. C., Andersen, N. A., Morrow, G. E. & Harris, R. L. Spurs, sexual dimorphism and reproductive maturity in Tasmanian echidnas (Tachyglossus
aculeatus setosus). *Australian Mammalogy* **41**, 161–169 (2018).

28.   Kawamichi, M. Nest Structure Dynamics and Seasonal Use of Nests by Siberian Chipmunks (Eutamias sibiricus). *J Mammal* **70**, 44–57 (1989).

29.   Kawamichi, M. Ecological Factors Affecting Annual Variation in Commencement of Hibernation in Wild Chipmunks (Tamias sibiricus). *J Mammal* **77**,
731–744 (1996).

30.   Knopf, F. L. & Balph, D. F. Annual Periodicity of Uinta Ground Squirrels. *The Southwestern Naturalist* **22**, 213–224 (1977).

31.   Morton, M. L. & Sherman, P. W. Effects of a spring snowstorm on behavior, reproduction, and survival of Belding's ground squirrels. *Canadian Journal*
of Zoology **56**, 2578–2590 (1978).

32.   Goldberg, A. R. *Diet, disease, and hibernation behavior of northern Idaho ground squirrels*. (University of Idaho, 2018).

33.   Barrett, J. S. Population viability of the southern Idaho ground squirrel (Spermophilus brunneus endemicus): effects of an altered landscape. (2005).

34.   Raveh, S. *et al.* Mating order and reproductive success in male Columbian ground squirrels (Urocitellus columbianus). *Behavioral Ecology* **21**, 537–547
(2010).

35.   Fagerstone, K. A. The annual cycle of Wyoming ground squirrels in Colorado. *Journal of Mammalogy* **69**, 678–687 (1988).

36.   Van Horne, B., Schooley, R. L., Olson, G. S. & Burnham, K. P. Patterns of density, reproduction, and survival in Townsend's ground squirrels. *Snake River*
Birds of Prey Area, research and monitoring, annual report. Edited by K. Steenhof. US Department of the Interior, Bureau of Land Management, Boise, Idaho 158
(1993).

37.   Van Horne, B., Olson, G. S., Schooley, R. L., Corn, J. G. & Burnham, K. P. EFFECTS OF DROUGHT AND PROLONGED WINTER ON TOWNSEND'S GROUND
SQUIRREL DEMOGRAPHY IN SHRUBSTEPPE HABITATS. *Ecological Monographs* **67**, 295–315 (1997).

38.   Sheriff, M. J. *et al.* Phenological variation in annual timing of hibernation and breeding in nearby populations of Arctic ground squirrels. *Proceedings of*
the Royal Society B: Biological Sciences **278**, 2369–2375 (2011).

39.   Buck, C. L. & Barnes, B. M. Annual cycle of body composition and hibernation in free-living arctic ground squirrels. *Journal of Mammalogy* **80**, 430–442
(1999).

40.   Michener, G. R. & Locklear, L. Over-winter weight loss by Richardson's ground squirrels in relation to sexual differences in mating effort. *Journal of*
Mammalogy **71**, 489–499 (1990).

41.   Michener, G. R. Sexual differences in reproductive effort of Richardson's ground squirrels. *Journal of Mammalogy* **79**, 1–19 (1998).

42.   Michener, G. R. Sexual differences in over-winter torpor patterns of Richardson's ground squirrels in natural hibernacula. *Oecologia* **89**, 397–406
(1992).

43.   Michener, G. R. & Locklear, L. Differential costs of reproductive effort for male and female Richardson's ground squirrels. *Ecology* **71**, 855–868 (1990).

44.   Munroe, K. E. & Koprowski, J. L. Annual cycles in the desert: body mass, activity and reproduction in round-tailed ground squirrels (Xerospermophilus
tereticaudus). *THE SOCIOECOLOGY, MATING SYSTEM AND BEHAVIOR OF ROUND-TAILED GROUND SQUIRRELS (XEROSPERMOPHILUS TERETICAUDUS)* 29 (2011).

45.   Hoyle, J. & Boonstra, R. Life history traits of the meadow jumping mouse, Zapus hudsonius, in southern Ontario. (1986).

46.   Brown, L. N. Seasonal activity patterns and breeding of the western jumping mouse (Zapus princeps) in Wyoming. *American Midland Naturalist* 460–
470 (1967).

47.   Cranford, J. A. Ecological strategies of a small hibernator, the western jumping mouse *Zapus princeps*. *Can. J. Zool.* **61**, 232–240 (1983).

1140

1141

1142

1143

1144

1145

1146

1147

1148

1149

Table S2 Data on dependent and independent factors used in models 2 and 3 (same data base without *Tachyglossus aculeatus*). Sex

difference in immergence was used as the dependent factor whereas strategy fatstoring or foodstoring, body mass change through

the end of mating, body mass immergence, dimorphism at immergence maternal effort, maternal effort duration, female specific

reproductive effort, minimum temperature and precipitation were considered as independent factors. The exact hibernation

phenology data for *Cricetus cricetus* have been confirmed by the authors. Weaning body mass for *Zapus princeps* and *Urocitellus*

mollis was calculated from body mass data at 52-60 days and 29 days, respectively, and corrected at 48 and 34 days, corresponding

to weaning dates. For maternal effort duration of *Urocitellus brunneus* and *Ictidomys parvidens*, we used respectively the averages

for the clade *Marmotini* obtained from Hayssen (2008). The body mass at weaning for *Urocitellus brunneus* correpsonds to the body

mass at weaning of *Urocitellus townsendii* (phylogenetically close species) corrected for the difference in body mass of the females

from Hayssen (2008) as body mass accounts for 76–84% of the variation in weaning mass in the clade *Marmotini* (Hayssen, 2008)

Sex difference in immergence was calculated as follows: female Julian date – male Julian date.

1161

1162

1163

1164

1165

Species	Food/Fat-	Sex diff immerg	Minimum temper	Body mass change end of mating	Precipitation	Body mass immerg	Maternal effort duration	Littersize	Weaning mass	Body mass femelle	Dimorphism immergence	Specific reprod effort
Callospermophilus saturatus	Food	3[2]	-7,3	-13,1[2]	863	273[2]	70[1,3]	2,7[2]	76[2]	185[2]	0,97[2]	110,7
Cricetus cricetus	Food	27[4]	-2,7	3,2[8]	595	344[5,6]	43[3]	7,31[5]	105[5]	245[6]	1,44[5,6]	313,2
Cynomys leucurus	Fat	11[10]	-11,5	-17,7[7]	337	1288[7]	67[1]	5,64[7]	198,5[7]	491[7]	1,55[7]	227,6
Erinaceus europaeus	Fat	15,7[8,9]	-5,2	-10,1[8,9]	803,5	1152[8,9]	72[2]	5[10]	235[11]	606[8,9]	1,16[8,9]	193,8
Glis glis	Fat	14[12]	-2,5	-2,5[13]	758	122,26[13]	58[3]	4,9[14]	39,2[15]	99,32[13]	1,21[13]	193,3
Ictidomys parvidens	Fat	15[16]	-2,4	24,4[16]	347	282,5[16]	68[1]	6,4[16]	43,8[16]	187[16]	1,24[16]	149,9
Marmota monax	Fat	-2[17]	-11,5	-6,2[18]	1156	3798,7[18]	76,05[1]	3,5[19]	490[17]	2610[18]	1,08[18]	65,7
Microcebus murinus	Fat	0[20]	14,6	-13,5[21]	922	74,5[21]	98[3]	2[22]	36,6[23]	60[21]	0,71[21]	121,6
Poliocitellus franklinii	Fat	15,75[24,25]	-21,1	-4,5[24,25]	517	577,88[24,25]	58[1]	6,75[25]	101,33[1]	322[24,25]	1,26[24,25]	211,7
Spermophilus citellus	Fat	-21,25[26]	-2,5	-13,3[26]	556	352,47[26]	64,5[1]	3,9[26]	51,5[27]	188[26]	1,31[26]	106,8
Tachyglossus aculeatus	Fat	24[28]	3,5	-6,6[29]	625	4180[29]	167,5[30]	1[31]	1650[31]	3806[29]	1,13[29]	43,3
Urocitellus armatus	Fat	-8,6[32]	-13	2,2[32]	402	521,83[32]	45[1]	6[33]	60,3[34]	262[32]	1,32[32]	137,7
Urocitellus beldingi	Fat	20[35]	-12,3	-4,5[35]	568	381,8[35]	51,53[1]	6,31[36]	69[37]	217[35]	1,21[35]	200,6
Urocitellus brunneus	Fat	-2[38]	-11	-6,5[39]	610	269,5[39]	66,5[1]	6,2[40]	36,55[1]	121[39]	1,35[39]	186,7
Urocitellus columbianus	Fat	11,7[41]	-16,1	8,4[41]	693	579[41]	53,2[1]	3,6[42]	112,6[43]	380[41]	1,32[41]	106,6

1166

1167

1168

Urocitellus elegans	Fat	-9[44]	-16,6	-3,8[44]	281	363,68[44]	55,58[1]	5,88[45]	80,5[46]	203[44]	1,25[44]	233,1
Urocitellus mollis	Fat	-3,66[47]	-5,4	-3,4[48]	248	227,83[48]	58[1]	8,1[47]	32,8[47]	122,5[48]	1,38[48]	216,8
Urocitellus parryii	Food	-28[49]	-30,2	-21,2[50]	237	1034,74[50]	53	5,8	234,5	615[50]	1,05[50]	221,1
Urocitellus richardsonii	Food	12,16[51–54]	-14,42	-4,4[52,54]	370	524,52[52,54]	51,79	7,1	83,75	229[52,54]	1,30[52,54]	259,6
Zapus hudsonius	Fat	14[49]	-8,5	-9,5[55]	837	22,78[55]	47	5,5	7,8	16,7[55]	0,92[55]	256,8
Zapus princeps	Fat	0[56]	-14,1	-7,9[57]	570	35,24[57]	48	5,4	12,9	23[57]	1[57]	296,9

References

1.   Hayssen, V. Reproductive Effort in Squirrels: Ecological, Phylogenetic, Allometric, and Latitudinal Patterns. *J Mammal* **89**, 582–606 (2008).

2.   Kenagy, G. J., Sharbaugh, S. M. & Nagy, K. A. Annual cycle of energy and time expenditure in a golden-mantled ground squirrel population. *Oecologia* **78**, 269–282 (1989).

3.   Magalhães, J. P. D. & Costa, J. A database of vertebrate longevity records and their relation to other life-history traits. *Journal of Evolutionary Biology* **22**, 1770–1774 (2009).

4.   Siutz, C., Franceschini, C. & Millesi, E. Sex and age differences in hibernation patterns of common hamsters: adult females hibernate for shorter periods than males. *J Comp Physiol B* **186**, 801–811 (2016).

5.   Hufnagl, S., Franceschini-Zink, C. & Millesi, E. Seasonal constraints and reproductive performance in female Common hamsters (Cricetus cricetus). *Mammalian Biology* **76**, 124–128 (2011).

6.   Lebl, K. & Millesi, E. Yearling male Common hamsters and the trade-off between growth and reproduction. *Biosystematics and Ecology*
Series **25**, 115–126 (2008).

7.   Bakko, E. B. & Brown, L. N. Breeding Biology of The White-Tailed Prairie Dog, Cynomys Leucurus, in Wyoming. *Journal of Mammalogy* **48**,
100–112 (1967).

8.   Rautio, A., Valtonen, A. & Kunnasranta, M. The effects of sex and season on home range in European hedgehogs at the northern edge of
the species range. in *Annales Zoologici Fennici* vol. 50 107–123 (BioOne, 2013).

9.   Haigh, A., O'Riordan, R. M. & Butler, F. Nesting behaviour and seasonal body mass changes in a rural Irish population of the Western
hedgehog (Erinaceus europaeus). *Acta Theriologica* **57**, 321–331 (2012).

10.   Kristiansson, H. Young production of European hedgehog in Sweden and Britain. *Acta Theriologica* **26**, 504–507 (1981).

11.   Haigh, A. The ecology of the European hedgehog (Erinaceus europaeus) in rural Ireland. (University College Cork, 2011).

12.   Bieber, C. & Ruf, T. Seasonal timing of reproduction and hibernation in the edible dormouse (Glis glis). *Life in the cold: Evolution,*
mechanism, adaptation, and application 113–125 (2004).

13.   Bieber, C. Population dynamics, sexual activity, and reproduction failure in the fat dormouse (Myoxus glis). *Journal of Zoology* **244**, 223–
229 (1998).

14.   Vekhnik, V. A., Ruf, T. & Bieber, C. A Review on the Edible dormouse reproduction (Glis glis Linnaeus, 1766). *Journal of Wildlife and*
Biodiversity **6**, 24–45 (2022).

15.   Kryštufek, B. Glis glis (Rodentia: Gliridae). *Mammalian Species* **42**, 195–206 (2010).

16.   Schwanz, L. E. Annual cycle of activity, reproduction, and body mass in Mexican ground squirrels (Spermophilus mexicanus). *Journal of*
Mammalogy **87**, 1086–1095 (2006).

17.   Maher, C. R. Social organization in woodchucks (Marmota monax) and its relationship to growing season. *Ethology* **112**, 313–324 (2006).

18.   Maher, C. R. & Duron, M. Mating system and paternity in woodchucks (Marmota monax). *Journal of Mammalogy* **91**, 628–635 (2010).

19.   Maher, C. R. Effects of relatedness on social interaction rates in a solitary marmot. *Animal Behaviour* **78**, 925–933 (2009).

20.   Schmid, J. Sex-specific differences in activity patterns and fattening in the gray mouse lemur (Microcebus murinus) in Madagascar.
Journal of Mammalogy **80**, 749–757 (1999).

21.   Schmid, J. & Kappeler, P. M. Fluctuating sexual dimorphism and differential hibernation by sex in a primate, the gray mouse lemur
(Microcebus murinus). *Behavioral Ecology and Sociobiology* **43**, 125–132 (1998).

22.   Eberle, M. & Kappeler, P. M. Sex in the dark: determinants and consequences of mixed male mating tactics in Microcebus murinus, a
small solitary nocturnal primate. *Behav Ecol Sociobiol* **57**, 77–90 (2004).

23.   Zimmermann, E., Radespiel, U., Mestre-Francés, N. & Verdier, J.-M. 8 Life history variation in mouse lemurs (Microcebus murinus, M.
lehilahytsara): the effect of environmental and phylogenetic determinants. *The Dwarf and Mouse Lemurs of Madagascar: Biology, Behavior and*
Conservation Biogeography of the Cheirogaleidae **73**, 174 (2016).

24.   Choromanski-Norris, J., Fritzell, E. K. & Sargeant, A. B. Seasonal Activity Cycle and Weight Changes of the Franklin's Ground Squirrel.
American Midland Naturalist **116**, 101 (1986).

25.   Iverson, S. L. & Turner, B. N. Natural history of a Manitoba population of Franklin's ground squirrels. *Canadian Field-Naturalist* **86**, 145–
149 (1972).

26.   Millesi, E., Strijkstra, A. M., Hoffmann, I. E., Dittami, J. P. & Daan, S. Sex and Age Differences in Mass, Morphology, and Annual Cycle in
European Ground Squirrels, Spermophilus citellus. *J Mammal* **80**, 218–231 (1999).

27.   Huber, S., Hoffmann, I. E., Millesi, E., Dittami, J. & Arnold, W. Explaining the seasonal decline in litter size in European ground squirrels.
Ecography **24**, 205–211 (2001).

28.   Nicol, S. C., Morrow, G. E. & Harris, R. L. Energetics meets sexual conflict: The phenology of hibernation in Tasmanian echidnas.
Functional Ecology **33**, 2150–2160 (2019).

29.   Nicol, S. C., Andersen, N. A., Morrow, G. E. & Harris, R. L. Spurs, sexual dimorphism and reproductive maturity in Tasmanian echidnas
(Tachyglossus aculeatus setosus). *Australian Mammalogy* **41**, 161–169 (2018).

30.   Morrow, G., Andersen, N. A. & Nicol, S. C. Reproductive strategies of the short-beaked echidna–a review with new data from a long-term
study on the Tasmanian subspecies (Tachyglossus aculeatus setosus). *Australian Journal of Zoology* **57**, 275–282 (2009).

31.   Nicol, S. & Andersen, N. A. The life history of an egg-laying mammal, the echidna (Tachyglossus aculeatus). *Ecoscience* **14**, 275 (2007).

32.   Knopf, F. L. & Balph, D. F. Annual Periodicity of Uinta Ground Squirrels. *The Southwestern Naturalist* **22**, 213–224 (1977).

33.   Slade, N. A. & Balph, D. F. Population Ecology of Uinta Ground Squirrels. *Ecology* **55**, 989–1003 (1974).

34.   Eshelman, B. D. & Sonnemann, C. S. Spermophilus armatus. *Mamm Species* 1–6 (2000) doi:10.2307/0.637.1.

35.   Morton, M. L. & Sherman, P. W. Effects of a spring snowstorm on behavior, reproduction, and survival of Belding's ground squirrels.
Canadian Journal of Zoology **56**, 2578–2590 (1978).

36.   Morton, M. L. & Gallup, J. S. Reproductive cycle of the Belding ground squirrel (Spermophilus beldingi beldingi): seasonal and age
differences. *The Great Basin Naturalist* 427–433 (1975).

37.   Morton, M. L. & Tung, H. L. Growth and development in the Belding ground squirrel (Spermophilus beldingi beldingi). *Journal of
Mammalogy* **52**, 611–616 (1971).

38.   Goldberg, A. R. *Diet, disease, and hibernation behavior of northern Idaho ground squirrels*. (University of Idaho, 2018).

39.   Barrett, J. S. Population viability of the southern Idaho ground squirrel (Spermophilus brunneus endemicus): effects of an altered
landscape. (2005).

40.   Yensen, E. & Sherman, P. W. Spermophilus brunneus. *Mamm Species* 1–5 (1997) doi:10.2307/3504405.

41.   Raveh, S. *et al.* Mating order and reproductive success in male Columbian ground squirrels (Urocitellus columbianus). *Behavioral Ecology*
21, 537–547 (2010).

42.   Neuhaus, P. Parasite removal and its impact on litter size and body condition in Columbian ground squirrels (Spermophilus columbianus).
Proc. R. Soc. Lond. B **270**, (2003).

43.   Skibiel, A. L., Dobson, F. S. & Murie, J. O. Maternal influences on reproduction in two populations of Columbian ground squirrels.
Ecological Monographs **79**, 325–341 (2009).

44.   Fagerstone, K. A. The annual cycle of Wyoming ground squirrels in Colorado. *Journal of Mammalogy* **69**, 678–687 (1988).

45.   Clark, T. W. Richardson's ground squirrel (Spermophilus richardsonii) in the Laramie Basin, Wyoming. *The Great Basin Naturalist* 55–70
(1970).

46.   Clark, T. W. Early growth, development, and behavior of the Richardson ground squirrel (Spermophilus richardsoni elegans). *American*
Midland Naturalist 197–205 (1970).

47.   Van Horne, B., Schooley, R. L., Olson, G. S. & Burnham, K. P. Patterns of density, reproduction, and survival in Townsend's ground
squirrels. *Snake River Birds of Prey Area, research and monitoring, annual report. Edited by K. Steenhof. US Department of the Interior, Bureau of*
Land Management, Boise, Idaho 158 (1993).

48.   Van Horne, B., Olson, G. S., Schooley, R. L., Corn, J. G. & Burnham, K. P. EFFECTS OF DROUGHT AND PROLONGED WINTER ON
TOWNSEND'S GROUND SQUIRREL DEMOGRAPHY IN SHRUBSTEPPE HABITATS. *Ecological Monographs* **67**, 295–315 (1997).

49.   Sheriff, M. J. *et al.* Phenological variation in annual timing of hibernation and breeding in nearby populations of Arctic ground squirrels.
Proceedings of the Royal Society B: Biological Sciences **278**, 2369–2375 (2011).

50.   Buck, C. L. & Barnes, B. M. Annual cycle of body composition and hibernation in free-living arctic ground squirrels. *Journal of*
Mammalogy **80**, 430–442 (1999).

51.   Michener, G. R. & Locklear, L. Over-winter weight loss by Richardson's ground squirrels in relation to sexual differences in mating effort.
Journal of Mammalogy **71**, 489–499 (1990).

52.   Michener, G. R. Sexual differences in reproductive effort of Richardson's ground squirrels. *Journal of Mammalogy* **79**, 1–19 (1998).

53.   Michener, G. R. Sexual differences in over-winter torpor patterns of Richardson's ground squirrels in natural hibernacula. *Oecologia* **89**,
397–406 (1992).

54.   Michener, G. R. & Locklear, L. Differential costs of reproductive effort for male and female Richardson's ground squirrels. *Ecology* **71**,
855–868 (1990).

55.   Hoyle, J. & Boonstra, R. Life history traits of the meadow jumping mouse, Zapus hudsonius, in southern Ontario. (1986).

56.   Brown, L. N. Seasonal activity patterns and breeding of the western jumping mouse (Zapus princeps) in Wyoming. *American Midland*
Naturalist 460–470 (1967).

57.   Cranford, J. A. Ecological strategies of a small hibernator, the western jumping mouse *Zapus princeps*. *Can. J. Zool.* **61**, 232–240 (1983).

1275

1276

1277

1278

1279

1280

1281

1282

1283

1284

1285

1286

1287

1288

1289

1290