Visuomotor prediction during action planning in the human frontoparietal cortex and cerebellum

Abbreviated title: Visuomotor prediction during action planning

Authors: Felix Quirmbach[1,2] and Jakub Limanowski[2,3]

[1]Faculty of Psychology, Technical University of Dresden, 01069 Dresden, Germany

[2]Center for Tactile Internet with Human-in-the-Loop, Technical University of Dresden, 01187 Dresden, Germany

[3]Institute of Psychology, University of Greifswald, 17486 Greifswald, Germany

Corresponding author email address: jakub.limanowski@uni-greifswald.de

Abstract

The concept of forward models in the brain, classically applied to describing on-line motor control, can in principle be extended to action planning; i.e., assuming forward sensory predictions are issued during the mere preparation of movements. To test this idea, we combined a delayed movement task with a virtual reality based manipulation of visuomotor congruence during functional magnetic resonance imaging (fMRI). Participants executed simple hand movements after a delay. During the delay, two aspects of the upcoming movement could be cued: the movement type and the visuomotor mapping (i.e., (in)congruence of executed hand movements and visual movement feedback by a glove-controlled virtual hand). Frontoparietal areas showed increased delay period activity when preparing pre-specified movements (cued > uncued). The cerebellum showed increased activity during the preparation for incongruent > congruent visuomotor mappings. The left anterior intraparietal sulcus (aIPS) showed an interaction effect, responding most strongly when a pre-specified (cued) movement was prepared under expected visuomotor incongruence. These results suggest that motor planning entails a forward prediction of visual body movement feedback, which can be adjusted in anticipation of nonstandard visuomotor mappings, and which is likely computed by the cerebellum and integrated with state estimates for (planned) control in the aIPS.

Introduction

Our current understanding of adaptive motor control largely builds upon the concept of internal

forward models that predict the sensory consequences of one's movement based on motor signals

(Miall & Wolpert, 1996; Wolpert & Kawato, 1998; Shadmehr, & Krakauer, 2008). This forward

prediction can be used to calculate a dynamic estimate of the effector's state, enabling 'on-line' control

despite sensory delays (Todorov & Jordan, 2002; Wolpert & Ghahramani 2000). Moreover, forward

predictions can be updated by sensory prediction errors, enabling, for instance, the adaptation of

control to new sensorimotor mappings (e.g., altered visual movement feedback, Tseng et al., 2007;

Grafton et al., 2008; Shadmehr et al., 2010). The respective computations have been linked to the

posterior parietal cortex (PPC) and the cerebellum (Miall et al. 1993; Wolpert et al. 1998; Blakemore

et al., 1998; Kilteni et al., 2023), and their interaction with (pre)motor areas and the basal ganglia

implementing the task and control policies (Shadmehr & Krakauer, 2008; Scott, 2012; Parr et al., 2021).

Thus, activity of the PPC and cerebellum has been associated with the detection of 'nonstandard'

visuomotor mappings (Leube et al., 2003; Nahab et al., 2011; Limanowski et al., 2017; Arikan et al.,

2019), and with the learning process required to adapt motor control to those mappings (Ogawa et al.,

2006; Izawa et al., 2012; Van Kemenade et al., 2017, 2019; Ohata et al., 2020; Kufer et al., 2024). In

line with this, brain stimulation and clinical case studies have linked the efficacy of sensory forward

predictions directly to intact PPC and cerebellar function (Desmurget et al., 1999; Synofzik et al., 2008;

Parr et al., 2021; Schmitter & Straube, 2024).

Besides on-line control and learning, however, the internal forward model framework can, in principle,

also be applied to action *planning* (Stanley & Miall, 2009; Scott, 2012); in line with the proposed

functional equivalence of planning and overt action control in the brain (Jeannerod & Decety, 1995;

Lotze, 1999). A derivative hypothesis of this extension is that the motor system should also issue

sensory predictions when merely planning actions; i.e., in the absence of overt action and associated

sensory movement feedback (Kuang et al., 2015; Pilacinski et al., 2018; Kilteni et al., 2018).

We tested this idea by combining a delayed movement paradigm with a virtual reality based

manipulation of visuomotor mapping inside an MR scanner. In delayed movement tasks, movements

are pre-cued (Rosenbaum, 1980); neural activity during the delay period is thought to encode specific

aspects of motor planning, such as the selection of effectors or goals, and the preparation of the

appropriate control policy and motor commands (Crammond & Kalaska, 1989; Hoshi & Tanji, 2002;

Fernandez-Ruiz et al., 2007; Bernier et al., 2012; Ariani et al., 2022). These planning processes have

been linked to activity in the dorsal premotor cortex (PMd) and the PPC (Churchland et al., 2010; Cisek

& Kalaska, 2005; Ferraina & Bianchi, 1994; Kaufman et al., 2014; Kuang et al., 2015; Tanji & Evarts,

1976; Schaffelhofer et al., 2015; Gertz & Fiehler, 2015; Lindner et al., 2010; Verhagen et al., 2012).

In our experiment, participants had to execute one of two possible hand movements after a variable

delay (open or close, see Fig. 1). During the delay, two aspects of the upcoming movement could be

independently cued (or not): the movement type and the visuomotor mapping. We assumed that

cueing (pre-specifying) movements would establish a specific motor plan during the delay; whereas in

the uncued case, a motor plan would be initiated only 'ad-hoc' upon presentation of the execution

prompt (Rosenbaum, 1980; cf. Ames et al., 2014; Pilacinski et al., 2018). Consequently, we expected

relatively increased frontoparietal blood oxygenation level dependent signal (BOLD) during the delay

period when participants were preparing for cued > uncued movements. If, alternatively, both

movements were prepared in parallel during the delay, frontoparietal BOLD should show the opposite

effect (uncued > cued).

By cueing incongruent (i.e., inverted) visual hand movements, we aimed to establish an anticipation of

an unnatural visuomotor mapping. In other words, we assumed that participants would naturally

expect visuomotor congruence, based on a life-long, learned association between executed

movements and their visual feedback; and that incongruent movements would violate this expectation

(Yon et al., 2018). When these 'nonstandard' visuomotor mappings were cued before the actual

movement execution, we expected an anticipatory adjustment of the forward model's predictions,

detectable as increased delay period BOLD. Importantly, as our study design entailed no spatial targets

during planning and execution, these predictions should specifically relate to visual body movement

feedback, i.e.: *What my hand movement will look like*. Moreover, we expected motor planning to

require more neuronal resources when integrating the (updated) predictions of nonstandard visual

feedback in the state estimate—with corresponding interaction effects predominantly in the PPC and

PMd.

Figure 1. Experimental task to manipulate motor planning and visuomotor prediction. The participants' real hand movements (left, hidden from view) were measured via a data glove and fed to a virtual hand (VH) model presented on screen (right). In each trial, participants had to execute one of two possible movements (open or close their hand) after a variable delay, starting from a neutral position (hand half-way closed). During the delay (preparation period), we presented a visual cue that could provide two kinds of predictive information: Firstly, the cue's *shape* could specify the movement type to be executed after the delay (up- or downward triangle = open or close their hand, 25% of all trials each, i.e., together 50%) or leave it ambiguous (square = upcoming movement unknown, 50% of all trials). Secondly, the cue's *color* indicated the to-be-expected visuomotor congruence; i.e., whether the virtual hand would next reflect the real hand movement (here: cyan cue) or the respective opposing movement (i.e., inverting the real hand movement, here: pink cue). Thus, we orthogonally manipulated processes related to motor planning and visuomotor prediction (the predicted mapping of visual movement feedback onto motor commands; i.e., what the hand movement will look like), respectively; in the absence of spatial movement targets during planning or execution. After a 'Go signal' and appearance of the virtual hand on screen, the participants executed the instructed hand movement while maintaining central fixation. In this particular example, the participant was cued that she would need to open her hand (triangle pointing upwards) and that the virtual hand would move incongruently during execution (pink cue), i.e., close. To ensure task engagement and attention to the cues, participants had to monitor and, in occasional catch trials, report whether the virtual hand movement matched the color-cued predictions (in 25% of trials, the virtual hand mapping was invalidly cued). Trials were separated by a brief (2s) interval. In this figure, the visual cues have been enlarged for display purposes.

Materials and Methods

Participants

30 healthy, right-handed volunteers (18 females, mean age = 25.4 years, range 19-35) participated in

the experiment after signing informed consent. The experiment was approved by the ethics committee

of the Technische Universität Dresden and conducted in accordance with this approval.

Experimental design and procedure

Participants lay comfortably inside an MR scanner. On their left hand, they wore an MR-compatible

data glove (5DT Data Glove 14 Ultra MRI, 60 Hz sampling rate, full-speed USB 1.1 connection), which

measured each finger's flexion via sewn-in optical sensors. The glove data were fed to a photo-realistic

virtual hand model, which was thus controllable by the participants' hand movements. We could not

measure the exact intrinsic delay of the setup, but even including conservative estimates for USB

transmission and projector latencies, the intrinsic delay should be less than visuomotor delays typically

detected above chance level in such setups (e.g., 167 ms in our previous study using the same glove

and hand model, Limanowski et al., 2017). The virtual hand received an average of the four sensor

values to ensure smooth and coherent visual motion (cf. Limanowski & Friston, 2020). The participants'

real hand was placed outside of their view; i.e., during the entire experiment, participants only saw

the virtual hand, but not their real hand. Prior to the experiment, the glove was calibrated carefully;

i.e., the virtual hand model was shown and the participant moved her fingers to establish the possible

range of flexion (minimum and maximum sensor values), and the virtual hand model's finger flexion

was adjusted to closely match the real hand posture and movement range according to the

participant's perception. If necessary, the calibration was repeated during the pauses between

individual scanning runs. Through a mirror on the head coil, participants saw a screen, onto which a

projector (Casio XJ-M156, Laser/LED hybrid, 1024 x 768 pixels resolution, 60 Hz refresh rate) displayed

the virtual reality environment; i.e., the instructions, cues, and the virtual hand model. The virtual

environment was created with the Blender 3D graphics software package (https://www.blender.org/,

Version 2.79) and its Python programming interface. The experimenter observed the virtual display

(via a mirrored screen) at all times, thus ensuring task compliance.

The participants' task was to perform, in each trial, one of two types of cued hand movements (open

or close, from a neutral starting position) after an instructed delay. During movement, the virtual hand

model moved congruently or incongruently to the participants' actual hand movements*.* Each trial

consisted of two periods – a *preparation period,* in which participants could be cued about various

aspects of the upcoming movement (i.e., inducing motor planning and visuomotor expectations, see

below); followed by an *execution period*, where participants executed the hand movements, while

receiving (condition-specific, see below) visual movement feedback from the virtual hand model. See

Figure 1 for an example trial.

At the beginning of each trial, participants had to bring their real, unseen hand to the neutral starting

position (i.e., fingers half-way closed). For this, a small white bar was presented on screen, whose

vertical position was controlled by the participants' hand posture via the data glove (i.e., the degree

to which the hand was opened or closed). Participants had to align this bar with a central target

(another bar, see Fig. 1); which, on average, took 1.96 s (SD = 0.48 s). Upon alignment, the bar was

then replaced by a central white fixation dot for 1 s. Participants were instructed to keep their gaze on

this fixation dot throughout the entire trial; i.e., until the end of the execution period.

Then, a cue was presented and remained visible throughout the delay period (3 to 5 s jittered, pseudo-

randomized, evenly distributed across conditions). This cue could provide participants with two kinds

of predictive information: The *type of the upcoming movement* (i.e., hand opening or closing) and the

congruence of the virtual and real hand movements during this movement (see below). These two

predictions were coded by cue shape and color, respectively. Thus, in half of all trials, the cue *shape*

specified the movement that the participant would need to execute next. Specifically, in 25% of trials

an upward-pointing triangle informed participants that they would have to *open* their hand, while in

another 25% a downward-pointing triangle informed them they would have to *close* their hand. In the

remaining half of all trials, the cue was square-shaped and, thus, did not reveal the to-be-executed

movement type in advance. In other words, only in half of the trials participants were able to generate

a specific motor prediction about the movement they would have to execute after the delay period.

Furthermore, the *color* of the cue was predictive of whether, during the upcoming movement, the

virtual hand would move congruently (i.e., reflecting the participants' real hand movement) or

incongruently (performing the respective opposite, i.e., inverted, movement type) with the

participant's real hand movements. In the incongruent conditions, the values fed to the hand model

from the glove's sensors were inverted so that an opening movement would close the virtual hand,

and vice versa. Thus, the movements of the virtual hand were controlled by the participant in both

cases, ensuring matching virtual and real hand kinematics during congruent and incongruent

conditions. This was important because our 'incongruent' virtual hand movements were, therefore,

also predictable in principle from the real hand movements; i.e., from motor signals.

The two movement types were designed to produce, from the neutral starting position, a comparable

range of visual motion; i.e., cover the same Euclidean distance on screen: At the beginning of the trial,

the virtual hand model was displayed on the screen in the neutral starting position. In this position,

the tip of the middle finger was positioned exactly in the middle of the screen (i.e., in the same position

as the fixation dot); and the Euclidean distance between this fixation dot and the fingertip for either

the fully opened or closed hand position was the same. While opening vs closing the hand were not

diametrical opposites in terms of actual joint rotations (cf. Fig. 2A), all conditions contained the same

number of open and close movements (see above); and all fMRI contrasts were calculated on

conditional contrast estimates pooled over both movement types. Thus, kinetic or kinematic

differences between opening vs closing movements were balanced across conditions; i.e., any

potentially related effects were factored out in our fMRI analysis.

The cue colors (pink and cyan) were counter-balanced across participants. Following the preparatory

period, a 'Go' signal (white triangle, centrally presented) appeared. Again, the pointing direction of this

triangle instructed participants to open (triangle pointing up) or close (triangle pointing down) their

hand. Note that, if the prior cue had not yet specified the movement type (i.e., square-shaped cue),

the participants would only now know which movement they should execute.

Participants were instructed to execute the movement as soon as the 'Go' signal disappeared (after

500 ms) and, instead, the virtual hand model was presented on the screen. This was done to ensure

that participants did not start to move reflexively before the virtual hand was visible. Participants then

had to execute the hand movement fluently, for which they had up to 2 s (if no movement was made

during that period, the trial ended). Participants were instructed and trained to make similar

movements in all conditions and, after movement, to keep the hand in the final position (completely

opened or closed) until the end of the trial. Then, the virtual hand disappeared again and a blank screen

was shown for 2 s. As in the preparation period, participants still had to maintain fixation on the central

dot during and post execution. In a prior training session, participants were able to practice the

movements and familiarize themselves with all conditions. In sum, this experimental setup resulted in

a factorial design with the factors *Movement type* (Cued, Uncued) and *Visuomotor mapping*

(Congruent, Incongruent); and, correspondingly, four conditions: CC (cued movement with expected

congruent mapping), CI (cued movement with expected incongruent mapping), UC (uncued movement

with expected congruent mapping) and UI (uncued movement with expected incongruent mapping).

In other words, movement planning was orthogonal to expectations about visuomotor congruence.

One important characteristic of our experimental design was the absence of spatial movement targets

or 'goals' during planning or execution. The presence of such targets, during planning or executing

movements towards (or away from) those targets, can make it difficult to distinguish activity related

to the processing or encoding of the properties of visual movement feedback *per se* from, for instance,

spatial eye-hand coordination or the visual-spatial representation of the movement goal (see Pilacinski

et al., 2018; Lindner et al., 2010, for discussions). Therefore, in our design, we ensured that the

visuomotor predictions pertained exclusively to the mapping of executed and observed hand

movements, in other words: a prediction of what the planned hand movement would look like.

One implication of this design was that participants could have abstained from movement planning

and simply waited for the 'Go' signal, presented in the form of a triangle. To ensure that participants

nevertheless paid attention to the cues and, consequently, generated the respective visuomotor

expectations or predictions, we included occasional trials in which the virtual hand moved *contrary* to

the color-cued expectation (25% of all trials, balanced across conditions). For instance, the virtual hand

moved congruently although the cue had predicted it to move incongruently (but note that the cues

indicating the to be executed 'own' movement type were always valid). Following these invalidly cued

trials, and following the same number of validly cued trials in each condition, participants had to report

whether the virtual hand behavior matched the predictions of the cue color (cf. Yon et al., 2018). For

this, a probe question ("Prediction correct?") was presented centrally on the screen alongside two

response options ("Yes"/"No"). Participants had to choose between those by pressing one of the

response device buttons twice (2-Button Fiber Optic Response pad, Current Designs, placed in the

participant's right hand, on their lap). If they did so within 1.5 s, feedback (i.e. "right" or "wrong") was

shown for 0.75 s, otherwise they were visually instructed to respond faster the next time. A blank

screen was shown for 2 s before the next trial. To increase motivation, participants received an

additional monetary reward after the experiment, depending on the proportion of correct answers to

the probe questions in the catch trials (1 € if at least 75% were answered correctly, 3 € for ≥85% and

5 € for ≥95%). We chose a relatively large number of catch trials (half of all trials in total) to maximize

the task relevance of the cues during the preparation period; i.e., participants had to assume that the

cue color could be probed by chance. Thus, the design for the execution period entailed *Visuomotor*

cue validity (Valid, Invalid), as another nested factor (see below). This manipulation also allowed us to

examine neuronal "surprise" responses to expectation violation; i.e., by comparing invalidly with

validly cued catch trials.

Each trial lasted between 8.5 and 15 s. For each participant, we recorded 8 runs of 32 trials each (6.5-

8.5 mins run length), resulting in 256 trials (64 per condition and preparation/execution period,

respectively) and around 60 mins scanning time in total (depending on hand positioning and response

times). Participants were free to rest as much as they wanted in between the individual scanning runs.

Behavioral data analysis

We tested for systematic differences in movement onset, duration, or amplitude between conditions;

using the participants' hand movement data from the glove sensors (averaged over all fingers but the

thumb; the movements had a similar kinematic profile, comparable across conditions, without

averaging the contribution of the individual fingers, see Results). We segmented the continuous glove

data to select the execution period time window of each trial, smoothed (Gaussian-weighted moving

average filter, time window 10 frames i.e. 167 ms), and re-adjusted the values to a baseline defined by

the mean hand position during the 500 ms before appearance of the virtual hand (i.e. the instructed

start of execution). Movement onset was defined, for each trial, as the respectively last time point

before movement with zero velocity. Note that we could only approximate the actual movement onset

from the recorded glove data; intrinsic delays due to the design of the glove (see above) should be

taken into account when interpreting the onset values, but these delays were constant and hence

irrelevant to our factorial analyses. The end of the movement was defined as the time point where

hand velocity reached zero again. Movement duration and amplitude, respectively, were defined as

the time difference and amplitude difference between movement onset and -end. Mean values for

movement onset, duration and amplitude per condition were calculated for each subject. Based on

these, a four-way ANOVA with the factors *Movement type* (Open, Close) *Movement type cueing* (Cued,

Uncued), *Visuomotor mapping* (Congruent, Incongruent) and *Visuomotor cue validity* (Valid, Invalid)

was used to test for differences between conditions. Reaction time in catch trials was defined as the

difference between presentation of the question on screen and the time point at which participants

confirmed their answer. All analyses were performed using MATLAB (MathWorks).

FMRI data preprocessing and analysis

The fMRI data were recorded with a 3T scanner (MAGNETOM Prisma; Siemens) equipped with a 32-

channel head coil. T2*-weighted images were acquired using an echo-planar imaging sequence

(GRAPPA acceleration factor = 3, multi-band factor = 6, TR = 869 ms, TE = 38 ms, flip angle = 58°, voxel

size = 2.4 mm^3, matrix size = 88 x 88, 60 interleaved slices). Depending on the length of the individual

session (see above), 3000-4000 functional image volumes were recorded per participant. Additionally,

for each participant we recorded a T1-weighted anatomical image (3D, voxel size = 0.9 x 0.9 x 0.9 mm^3,

FOV = 272 x 272 mm, 240 slices, TR = 2.4 s, TE = 2.2 ms, flip angle = 8°) and two GRE field maps (64

layers, voxel size 2 x 2 x 2 mm^3; matrix size = 88 x 88; TR = 698 ms, TE_1 = 5.19 ms, TE_2 = 7.65 ms, flip

angle 54°), after the second and sixth scanning run, respectively. All fMRI preprocessing steps and

analyses were performed using MATLAB (MathWorks) and SPM12 (Wellcome Trust Centre for

Neuroimaging, University College London; https://www.fil.ion.ucl.ac.uk/spm). The functional images

were realigned and unwarped using voxel displacement maps calculated from the two field maps (the

first map was used for runs 1 to 4, the second for runs 5 to 8). These images were then co-registered

with the individual T1 image, spatially normalized to MNI space via previously estimated deformation

fields, resliced to 2 mm^3 voxels, and smoothed with an 8 mm full width at half maximum Gaussian

kernel. The T1 images were also normalized (but not smoothed) and averaged to create a group

averaged structural image.

In our fMRI analysis, we adopted a mass univariate approach; i.e., we tested for voxel-wise differences

in the amplitude of the BOLD signal between our experimental conditions. Implicit in this is the

assumption that a relatively higher or lower BOLD signal in one condition compared to another (i.e., a

differential contrast) indicates a regionally in- or decreased metabolic demand in this condition,

respectively. We interpret such differences as evincing an increased engagement of brain regions in

experimentally manipulated processes related to motor planning and visuomotor prediction, as

identified per our factorial between-condition comparisons (see Fernandez-Ruiz et al., 2007; Lindner

et al., 2010; Bernier et al., 2012; Gertz & Fiehler, 2015; or Pilacinski et al., 2018, for similar approaches).

For each participant, we fitted a general linear model (GLM, 128 s high-pass filter, including temporal

and dispersion derivatives for each regressor) to the preprocessed data. We included stick (0 s

duration) regressors modeling each combination of the two factors *Movement type* (Cued, Uncued)

and *Visuomotor mapping* (Congruent, Incongruent) in the preparation and movement period,

respectively. The resulting four regressors were thereby aligned with the onsets of the predictive cues;

or with the onsets of the movement execution period including the virtual hand presentation. To

model any potential 'surprise' effects, the regressors of the execution period were each supplemented

by a mean-centered parametric modulator encoding trials in which the virtual hand moved unlike cued

with +1, and trials in which it moved like cued with -1. Thereby, we applied the modulation to the catch

trials only; i.e., ensuring the surprise effect was calculated on the same amount of trials (all with

subsequent button presses) in invalidly and validly cued movements.

We added the six realignment parameters as noise regressors to each run, alongside a session

constant. Furthermore, we added three regressors of no interest modelling the differences in

movement amplitude, movement duration, and reaction time (i.e., the time it took participants to

initiate a movement after onset of the execution period) during the execution period. Each kinematic

regressor modelled trial-to-trial variations (in amplitude, duration, and reaction time) across the entire

scanning session; i.e., it also modelled differences between conditions. Two additional regressors of

no interest were included: one modeling the initial hand positioning (a 0s duration stick regressor with

an additional orthogonalized parametric regressor encoding the duration from trial start until the

correct hand position was reached and the fixation dot presented), and one modeling the response

periods after the occasional catch trials (a 0s duration stick regressor with an additional orthogonalized

parametric regressor encoding the respective variations in reaction time until the confirmatory key

press). On the first (single participant) level, T-contrast images were created for each regressor vs

baseline, across all 8 runs per participant (i.e., over all 64 trials per preparation or execution period per

condition. These first-level contrast images were entered into group-level GLMs using a flexible

factorial design, in which the group (second-level) T- and F-contrasts were calculated. We thereby used

2x2 factorial designs with the aforementioned factors to evaluate the main effects of cued vs uncued

movements and congruent vs incongruent visuomotor mapping, as well as their potential interaction,

during the preparation and execution periods, respectively.

We furthermore tested for the *consistency* of main effects; i.e., whether a given main effect of one

experimental factor was significant at each level of the respective other factor. To do this, we

calculated 'null' conjunction contrasts; a statistical test whether a BOLD signal difference in a given

voxel is significant in each of multiple tested contrasts. Thus, we calculated a conjunction across the

contrasts CC>UC and CI>UI (testing for consistent significance of the main effect movement type cued

> uncued across both levels of the factor visuomotor mapping); and another one across the contrasts

CI>CC and UI>UC (testing for consistent significance of the main effect incongruent > congruent

visuomotor mapping across both levels of the factor movement type cueing). These conjunction

contrasts were evaluated for statistical significance using a voxel-wise threshold of p_{FWE}<0.05,

corrected within the restricted search region defined by the significant voxels obtained from the

respective main effect.

Furthermore, we tested for any influence of the (cued) preparatory processes on activity in the

respectively identified areas during subsequent movement execution. As we did not have any specific

prediction about the directionality of response differences during movement execution, we used non-

directional F-contrasts, with post-hoc t-tests to examine directionality, in this case. To assess potential

surprise effects during execution associated with invalidly > validly cued movements, we calculated an

analogous 2x2 factorial group level design on the parametric 'surprise' modulators of each condition

(i.e., on T contrast images weighting invalidly cued catch trials in each condition with +1 and validly

cued ones with -1, see above). Finally, supplementing our main analysis (which focused on differences

in metabolic demand via estimating the amplitude of the hemodynamic response), we looked for

analogous conditional differences in the loadings on the temporal and dispersion derivatives of our

model's regressors; using an analogous second-level factorial design as in the main (amplitude)

analysis on the first-level contrast images of the respective derivative. These estimates can provide

additional information about the shape of the BOLD signal; i.e., a relatively larger estimate of the

temporal derivative may indicate a relatively earlier hemodynamic response, and a relatively larger

estimate of the dispersion derivative may indicate a relatively narrower response (Friston et al., 1998;

Henson et al., 2002). This can supplement the interpretation of the hemodynamic (amplitude)

response in terms of underlying neuronal, e.g. predictive, processes (cf. Kavroulakis et al., 2022).

Activations in the whole brain obtained from the group-level contrasts were assessed for statistical

significance using a voxel-wise threshold of $p<0.05$, family-wise error corrected ($p_{FWE}<0.05$). In some

cases, we had specific regions of interest defined a priori by orthogonal contrasts, and restricted the

search space accordingly, thus applying a small volume correction under the same voxel-wise threshold

of $p_{FWE}<0.05$ for those results (similar to the conjunction contrasts described above): Firstly, we looked

for an interaction of motor and visuomotor planning in brain regions showing a response to both

individual effects. Therefore, we defined the search space for these interaction effects based on all

(whole-brain) significant voxels obtained from the null conjunction of the respective main effects (Fig.

5). For completeness, we also tested for whole-brain results. Based on the previously reported overlap

between action planning and execution (Lotze et al., 1999; Ames et al., 2014), we expected that brain

regions involved in motor or visuomotor planning would also show condition specific effects during

movement execution. Therefore, for the analysis of responses during the execution period, we

restricted the search space for effects to all significant voxels obtained from each respective analogous

contrast during the preparation period. Thirdly, we hypothesized that surprise effects might emerge

in brain regions implementing the forward model, i.e., issuing the respective forward predictions. In

other words, in those regions, we expected a relative increase in BOLD in response to a (sensory)

prediction error elicited by the unpredicted visuomotor mapping. As the generation of those

predictions should be highlighted by the main effect of preparation for incongruent > congruent

movements (see Introduction), we used those significant activations as a mask image to look for

corresponding surprise effects; but also calculated an additional whole-brain analysis.

The significant activation differences resulting from the group level contrasts are presented as renders

on SPM12's template brains, or superimposed onto the group averaged normalized structural image.

The displayed significant activations contain only voxels surviving the chosen threshold of p_{FWE}<0.05.

The contrast estimate plots were created by extracting and averaging the first-level contrast estimates

for each condition from the peak voxel of each anatomically distinct cluster, using the rfxplot toolbox

(Gläscher, 2009). For anatomical reference, we used the SPM Anatomy toolbox (Eickhoff et al., 2005).

Eye tracking control experiment

Our non-spatial (i.e., without spatial movement targets) movement task was intended to render eye

movements useless and, thus, unlikely. As we did not measure gaze on-line inside the scanner, we

conducted an eye-tracking control experiment post-hoc, with the same design. The aim of this

experiment was to determine whether participants could, in principle in this task, maintain fixation

onto the central cues/dot; and whether gaze behavior was comparable across experimental

conditions. A sub-sample of our fMRI study participants (*N*=6) completed the same task as in the

scanner (with 4 instead of 8 runs in total), with stimuli presented at a size corresponding to those seen

in the scanner environment. Participants were seated comfortably in a constantly lit room in front of

a screen (at a distance of 93 cm) with their head stabilized via a chin rest, while we measured gaze

position with an Eyetracker (EyeLink 1000 Plus, Desktop Mount, SR Research). We segmented the

resulting eye position data to select 2500 ms time windows for both the preparation and execution

periods of each trial, with phases of data loss (e.g. due to eye blinks) corrected via linear interpolation.

Gaze position was defined relative to the position of the central fixation dot; i.e., the measured fixation

position in the horizontal and vertical planes was combined to a measure of Euclidean distance from

the fixation point, and converted to degrees of visual angle. To evaluate the quality of fixation, we

tested for differences in mean gaze position and in the variability of fixation (i.e., the standard

deviation); and we tested for this separately for the two possible movements. Thus, we performed a

four-way repeated-measures ANOVA with the factors *Movement* (Open, Close), *Movement type* (Cued,

Uncued), *Visuomotor mapping* (Congruent, Incongruent), and *Visuomotor cue validity* (Valid, Invalid)

on the participants' condition averages of both fixation position and variability, respectively.

FMRI right-hand control experiment

To follow up on the finding that the main effects of cueing (and to a lesser extent also execution) were

more pronounced in the left than in the right PPC (i.e., ipsilateral to the moving hand, see Figs. 3-6),

we scanned 6 of our original participants again in a separate session; using an identical experimental

setup and design (with 5 instead of 8 runs in total) but with right-hand instead of left-hand movements

(and a right instead of left virtual hand model). These data were preprocessed and analyzed as

described for the main experiment. As the small sample size limited the power of the random-effects

group analysis, we evaluated and displayed the resulting statistical parametric maps at a threshold of

$p < 0.001$, uncorrected (see supplement, Fig. S6).

Results

All participants showed very good task performance; i.e., all performing >97% correct movements (on

average, only one incorrect movement per recording session) and correctly identifying >93% of the

catch trials (on average, >98% of responses were correct). The average reaction time in catch trials was

0.920±0.246 s. There were no significant effects of visuomotor cue validity, visuomotor mapping, or

motor planning on reaction times. The participants' hand movements were consistent across

conditions (Fig. 2). On average, movement execution was detectable from the glove data 121±119 ms

after the onset of the execution period, with a mean movement duration of 1181±316 ms. Movement

amplitude, in terms of average flexion values relative to the starting position, averaged to -0.26±0.08

(hand opening) and 0.48±0.09 (hand closing), respectively. Open vs close hand movements differed in

terms of movement amplitude ($F_{(2,29)}$ = 73.880, p<0.001), but not in terms of onset or duration of the

movements. Crucially, there was no significant effect of visuomotor mapping (congruent vs

incongruent) on movement onset, duration, or amplitude of the hand movements. Likewise, motor

planning (cued vs uncued) had no significant effect on movement amplitude or duration. However,

movements in cued trials were on average initiated 51 ms earlier than those in uncued trials ($F_{(2,29)}$ =

7.34, p=0.011). This very likely indicates a reaction time benefit of prior motor planning in cued trials

(which speeded up subsequent execution upon prompting), in line with previous findings (Rosenbaum,

1980; cf. Ames et al., 2014). Finally, there was a significant three-way-interaction (cueing x mapping x

cue validity) on movement duration ($F_{(2,29)}$ = 6.82, p=0.014), which was, however, difficult to interpret.

Interestingly, movements in invalidly cued visuomotor mapping trials were, on average, initiated 72

ms later ($F_{(2,29)}$ = 42.09, p<0.001) and 155 ms shorter ($F_{(2,29)}$ = 56.67, p<0.001) than movements in validly

cued trials; but movement amplitude was unaffected by cue validity ($F_{(2,29)}$ = 0.38, n.s). This could

indicate very early surprise responses to the expectation violation, leading to a slower onset (which

was defined as the glove sensor flexion data crossing a certain threshold, see Materials and Methods).

Importantly, however, this effect was comparable across experimental factors (i.e., no significant

interaction with either visuomotor mapping or motor planning), hence our fMRI contrasts of interest

– between conditions – were unlikely to have been affected by this. Furthermore, kinematic

differences between trials and conditions were included as covariates of no interest in the fMRI

analysis (see Materials and Methods). All of the above significant conditional effects were replicated

when analyzing the data of the four fingers individually (significant for the index, middle, and ring

finger, Fs > 5.9, all ps < 0.05; non-significant for the pinky, Fs > 1.9, ps < 0.17); there were no additional

significant main or interaction effects when looking at each finger individually. A supplementary eye-

tracking experiment (*N*=6) showed that participants were, in principle, able to maintain very stable

fixation onto the central cues independently of condition throughout the preparation period, and

overall also during movement execution (see Fig. S1 for details).

In sum, the participants' behavior suggested a high task engagement and comparable hand

movements (and fixation) across conditions, while showing a likely reaction time benefit of pre-cueing

movements. Moreover, the almost perfect responses to catch trials, in which the validity of the color-

cued and the actual visuomotor mapping was probed, indicated that participants paid attention to the

visuomotor cues. Thus, we were confident that participants generated the desired (cued) motor plans

and visuomotor predictions during scanning.

Figure 2. Average hand movements during the execution period. **(A)** The curves show the group-averaged unsmoothed flexion values with associated standard deviations as shaded areas (plotted for each condition, CC = cued/congruent, CI = cued/incongruent, UC = uncued/congruent, UI = uncued/incongruent, and separately for validly vs invalidly cued trials pooled over conditions). Positive flexion values indicating closing and negative values indicating opening, relative to the neutral starting position; time = 0 marks the instructed start and appearance of the virtual hand model. **(B)** The bar plots show the average movement onsets with associated standard errors of the mean (for each pair of bars, the first bar corresponds to the "open" hand movement, the second bar to the "close" movement). Movements were initiated significantly faster in cued > uncued trials, and in valid > invalid trials(*$p<0.05$, ***$p<0.001$). See Results for details.

In our fMRI analysis, our primary aim was to identify brain areas involved in motor planning and

visuomotor prediction; i.e., we looked for corresponding BOLD signal differences during the delay

period. First, we contrasted the delay period of trials in which a specific movement type could be

prepared – based on the cue shape – with trials in which this was not possible (i.e., main effect of

movement type cued > uncued). As expected, this contrast revealed significantly ($p_{FWE}<0.05$) stronger

BOLD signal predominantly in the bilateral frontoparietal cortices; including the bilateral aIPS (and, in

the left hemisphere, larger portions of the IPS), the bilateral SPL, the SMA, medial parts of the bilateral

PMd, the left ventral premotor cortex (PMv); and the bilateral lateral occipital cortices (LOC, including

area V5). See Table 1 and Figure 3A for details. A conjunction analysis revealed that, in almost all of

these regions, the main effect of cued > uncued movements was consistent across congruent and

incongruent visuomotor mappings (i.e., levels of the second experimental factor; p_{FWE}<0.05, see Table

1). In other words, this suggested a consistently elevated BOLD signal by motor planning during

preparation for congruent and incongruent visuomotor mappings alike. None of these regions showed

significant differences in an analysis of the analogous contrasts calculated on the temporal and

dispersion derivatives (Fig. S2). The converse contrast (i.e., delay period of uncued > cued movement

types) revealed smaller clusters of significant voxels mainly in the visual and somatosensory cortices

(Table 1 and Fig. 3B).

Then, we tested for increased responses during the delay period of trials in which an incongruent >

congruent mapping of the virtual hand movements relative to the real hand movements was cued;

assuming that here, predictions of the forward model would need updating. There was a significant

main effect of cues predicting an incongruent > congruent mapping in the cerebellum (centered on

Lobule VI in each hemisphere), premotor (the bilateral PMd, and the left preSMA and PMv) and

posterior parietal (the left aIPS and the bilateral SPL) areas (see Figure 4 and Table 1 for all effects).

However, a conjunction analysis revealed that this effect was consistent across cued and uncued

movement types in only some of those regions (p_{FWE}<0.05, see Fig. 4): Most notably, the strongest

consistent effects across the CI > CC and the UI > UC contrasts were located in the bilateral Lobules VI

of the cerebellum (with the strongest effect located in the right cerebellum, see slice overlays in Fig.

4). Only few other voxels in the PMd, the SMA, and in the left precuneus showed consistent effects.

No significant activation differences were observed for the converse contrast, cued congruent >

incongruent visuomotor mapping. Furthermore, an analysis of the analogous contrasts calculated on

the temporal and dispersion derivatives revealed significantly earlier and narrower hemodynamic

responses related to the preparation of congruent > incongruent mappings in the same areas of the

cerebellum (Fig. 4; further temporal but not dispersion effects were observed in the left IPS, PMd, and

PMv, see Fig. S2). In other words, the (right) cerebellum's hemodynamic response peaked relatively

later and was wider during the preparation for visuomotor incongruence, compared with

congruence—suggesting, together with its higher amplitude, more sustained neural activity (cf. Friston

et al., 1998; Henson et al., 2002).

Figure 3. Brain activations related to motor planning during the delay period. **(A)** The renders show significant (p_{FWE}<0.05) voxels obtained from contrasting the delay period of trials in which the to-be-executed movement type was cued, with trials in which this was left ambiguous until execution (cued > uncued). Inlay: Significant (p_{FWE}<0.05) voxels obtained from a conjunction testing for consistency of the effect cued > uncued across congruent and incongruent mappings. **(B)** The converse contrast (uncued > cued) yielded significant in visual and somatosensory cortices. CC = cued/congruent, CI = cued/incongruent, UC = uncued/congruent, UI = uncued/incongruent. See Table 1 for details.

Figure 4. Brain activations related to visuomotor prediction during the delay period. The renders show significant (p_{FWE}<0.05) voxels obtained from contrasting the delay period of trials in which the cue color predicted incongruent, compared with congruent visuomotor mapping (incongruent > congruent). The converse contrast yielded no significant effects. Inlay, left: Significant (p_{FWE}<0.05) voxels obtained from a conjunction testing for consistency of the effect incongruent > congruent visuomotor mapping across cued and uncued movements. CC = cued/congruent, CI = cued/incongruent, UC = uncued/congruent, UI = uncued/incongruent. CR = cerebellum. See Table 1 for details. Inlay, right: An analysis of the loadings on the temporal derivative (TD, top) and dispersion derivative (DD, bottom) revealed voxels in corresponding areas of the cerebellum showing significantly later and wider hemodynamic responses when incongruent movements were prepared.

Table 1. Brain areas showing significant main effects during the delay.

Movement type cued > uncued (CC+CI)-(UC+UI)		MNI			Peak	
Anatomical location	*Voxels*	*x*	*y*	*z*	*T*	*p_{FWE}*
L. Lateral occipital cortex (V5)*	390	-40	-82	-4	8.72	<0.001
R. Lateral occipital cortex (V5)*	183	46	-78	-8	7.85	<0.001
L. Superior frontal gyrus (SMA & PMd)*	475	-10	-4	64	7.06	<0.001
L. Intraparietal sulcus and superior parietal lobe*	695	-38	-38	46	6.68	<0.001
L. Precentral/inferior frontal gyrus (PMv)*	60	-60	10	26	6.26	<0.001
R. Superior parietal lobe (BA 7A)*	68	20	-68	58	6.08	0.001
R. Fusiform gyrus*	16	42	-58	12	5.85	0.002
R. Anterior intraparietal sulcus	46	34	-36	42	5.72	0.004

Movement type uncued > cued (UC+UI)-(CC+CI)		MNI			Peak	
Anatomical location	Voxels	*x*	*y*	*z*	*T*	*pFWE*
L. Pre-/postcentral gyrus (S1/M1)*	115	-28	-24	48	6.18	<0.001
L. Parietal operculum (OP)	19	-38	-24	22	5.78	0.003
L. Paracentral lobule*	17	-8	-20	50	5.69	0.004
R. Parietal operculum (OP)	16	46	-30	20	5.55	0.008
L. Medial temporal lobe*	8	-54	-10	-22	5.49	0.009
L. Occipital pole (V3)*	7	-18	-100	0	5.34	0.016
L. Parahippocampal gyrus*	1	-32	-40	-8	5.15	0.032
R. Anterior cingulate gyrus*	1	6	32	-8	5.07	0.041
R. Postcentral gyrus (S1)	1	22	-36	64	5.04	0.047
R. Postcentral gyrus (S1)	1	20	-38	62	5.03	0.048
Mapping incongruent > congruent (CI+UI)-(CC+UC)		MNI			Peak	
Anatomical location	Voxels	*x*	*y*	*z*	*T*	*pFWE*
R. Middle frontal gyrus (PMd)*	688	32	-2	56	7.94	<0.001
R. Cerebellum (Lobule VI/VII) *	781	38	-64	-32	7.86	<0.001
L. Superior/middle frontal gyrus (PMd/preSMA)*	1852	-10	12	54	7.22	<0.001
L. Cerebellum (Lobule VI)*	263	-26	-58	-34	6.64	<0.001
L. Superior parietal lobe (BA 7A)*	200	-8	-60	48	6.52	<0.001
L. Anterior intraparietal sulcus	486	-46	-42	52	6.07	0.001
R. Anterior intraparietal sulcus	243	36	-50	52	5.96	0.002
R. Cerebellum (Lobule I)	56	2	-56	-12	5.89	0.002
R. Cerebellum (Lobule VII/VIII)*	74	32	-64	-48	5.86	0.002
L. Superior temporal sulcus	13	-46	-50	8	5.76	0.003
R. Paracingulate gyrus	34	8	22	36	5.70	0.004
Cerebellum (Vermis)	96	0	-72	-18	5.57	0.007
L. Anterior insular cortex	51	-28	26	2	5.54	0.008
R. Superior parietal lobe (BA 7A)	62	-16	-66	54	5.47	0.010
R. Caudate	23	16	2	16	5.44	0.011
L. Middle frontal gyrus	5	-38	30	28	5.28	0.020
R. Anterior insular cortex	5	34	20	10	5.26	0.020
R. Intraparietal sulcus	4	30	-62	38	5.21	0.025
L. Supramarginal gyrus	2	-56	-44	26	5.18	0.029
L. Paracingulate gyrus	3	-8	20	36	5.17	0.029
R. Superior frontal gyrus	2	16	8	56	5.07	0.041
R. Intraparietal sulcus	1	28	-62	42	5.07	0.042
L. Cerebellum (Crus I)	1	-46	-58	-34	5.03	0.047
L. Middle frontal gyrus	2	-44	24	38	5.02	0.048

CC = cued/congruent, CI = cued/incongruent, UC = uncued/congruent, UI = uncued/incongruent.
* clusters with significant voxels in a conjunction contrast across both levels of the respective other factor.

Next, we specifically tested for brain areas that would jointly encode both preparatory processes in

our design; i.e., those related to motor planning and visuomotor mappings. A null conjunction across

both main effects (cued > uncued movement type and incongruent > congruent visuomotor mapping,

Fig. 3A and Fig. 4) revealed consistently significant voxels within the bilateral aIPS and SPL, and the left

medial PMd/SMA (p_{FWE}<0.05, Table 2 and Fig. S3). Then, we evaluated our hypothesis that motor

planning and visuomotor prediction would interact; i.e., we tested for relatively stronger movement

planning effects (cued > uncued movements) under expected visuomotor incongruence, compared

with congruence; i.e., via the contrast (CI-UI)-(CC-UC). Only the left aIPS/IPL showed a significant

corresponding interaction effect: a relatively stronger increase in fMRI activity for delay periods in

which the cue predicted a specific movement type with incongruent > congruent visuomotor mappings

(CI>CC), compared with when the movement type was left ambiguous (UI>UC; see Figure 5 and Table

2). In other words, in the left aIPS, while the main effect of cueing itself was consistent, the main effect

of preparing for incongruent visuomotor mappings was significantly stronger in the cued movement

conditions. No other significant effects were found; a similar interaction effect in the left PMd did not

reach significance (Fig. S3).

Table 2. Significant activation differences obtained from the conjunction of main effects and the interaction effect.

Null conjunction (CC+CI)-(UC+UI) ∩ (CI+UI)-(CC+UC)		MNI			Peak	
Anatomical location	*Voxels*	*x*	*y*	*z*	*T*	*p_{FWE}*
L. Superior frontal gyrus (PMd/SMA)	82	-12	0	64	6.17	0.001
L. Anterior intraparietal sulcus	197	-46	-42	52	6.01	0.001
L. Superior parietal lobule	29	-8	-60	54	5.86	0.002
R. Superior parietal lobule	11	18	-66	56	5.43	0.012
L. Superior frontal gyrus	1	-6	4	50	5.33	0.034
R. Anterior intraparietal sulcus	3	40	-36	44	5.13	0.034
R. Anterior intraparietal sulcus	1	36	-40	42	5.09	0.039
L. Superior frontal gyrus	2	-4	2	54	5.08	0.041
Interaction (IC-IU)-(CC-CU)		MNI			Peak	
Anatomical location	*Voxels*	*x*	*y*	*z*	*T*	*p_{FWE}*
L. Inferior parietal lobule / intraparietal sulcus	8	-46	-46	-56	3.70	0.015
L. Anterior intraparietal sulcus	5	-36	-42	40	3.47	0.040

CC = cued/congruent, CI = cued/incongruent, UC = uncued/congruent, UI = uncued/incongruent.

Figure 5. During the delay period, voxels in the left IPS/IPL showed a significant ($p_{FWE}<0.05$) interaction effect, within the areas showing both main effects as identified by a null conjunction contrast (see main text): the highest responses were observed during the preparation of specified (cued) movements when an incongruent visuomotor mapping was expected. The bar plots show the contrast estimates with associated standard errors of the mean. CC = cued/congruent, CI = cued/incongruent, UC = uncued/congruent, UI = uncued/incongruent.

Subsequently, we looked at condition-specific activation differences during the execution period

depending on the cues presented during the prior delay period. The left aIPS and the left PMd showed

significant BOLD signal differences depending on cueing (Fig. 6A and Table 3; cf. Fig. S4 for uncorrected

renders). A post-hoc t-test in these areas revealed significantly increased activation in trials where the

movement type was left uncued until the 'Go' signal, compared with cued trials ($p_{FWE}<0.05$). A whole-

brain analysis revealed one further significant voxel in the ventromedial prefrontal cortex. Exploratory

correlation analyses revealed that, in the key frontoparietal areas, stronger preparatory activity during

cued>uncued trials was associated with lower execution related activity (i.e., cued<uncued) and with

faster movement onsets across participants (Fig. S6, but not all of these correlations were significant).

Furthermore, an analysis of the loadings on the temporal derivative suggested significantly earlier

hemodynamic responses in bilateral frontoparietal areas, but not M1, for uncued > cued movement

execution (Fig. S2). Movement execution also depended on cued visuomotor mapping (congruent vs

incongruent virtual hand movement) in the left PMd, the left aIPS, and the left anterior insula (Fig. 6B

and Table 3; cf. Fig. S4 for uncorrected renders). A post-hoc t-test in these areas revealed significantly

increased activation during movements following cued incongruent > congruent visuomotor mapping

($p_{FWE}<0.05$). The main effects overlapped in the left aIPS (conjunction of main effects: x = -38, y = -50,

z = 50, peak F = 14.57, Fig. S4), but this effect did not reach corrected significance; there were no

significant interaction effects. Most of the key brain regions identified in the planning period contrasts

(Figs. 3-5) showed substantially stronger BOLD responses during movement compared with planning

(Fig. 6C)—with the notable exception of preparation for cued movements under expected

incongruence in the left aIPS (CI, cf. Fig. 5), and preparation for incongruent mappings in the right

cerebellum (CI and UI, cf. Fig. 4).

Figure 6. Brain activation differences during movement execution, related to prior movement planning and visuomotor predictions. **(A)** Voxels showing significant BOLD signal differences during movement execution in trials where the movement type had been cued in advance versus when it was left ambiguous (F-test, cued vs uncued, p_{FWE}<0.05; a post-hoc t-test revealed significantly stronger responses for uncued > cued, see Results). **(B)** Voxels showing significant BOLD signal differences during movement execution in trials where congruent vs incongruent visuomotor mapping had been cued (F-test, congruent vs incongruent, p_{FWE}<0.05; a post-hoc t-test revealed significantly stronger responses for incongruent > congruent, see Results). **(C)** Estimated hemodynamic responses, throughout the different trial phases of the experiment, in the main frontoparietal and cerebellar areas identified in our analysis (the renders show the peak coordinates marked with red dots). The plots show contrast estimates with associated standard errors of the mean: The first bar shows the estimate for the pre-

positioning period prior to each trial (cf. Fig. 1); the next bars show the contrast estimates for each condition, first for the delay (preparation period), then for the execution period. CC = cued/congruent, CI = cued/incongruent, UC = uncued/congruent, UI = uncued/incongruent. **(D)** Significantly (p_{FWE}<0.05) stronger activations by unexpected visuomotor mappings ('surprise' effect) within brain areas showing effects of visuomotor prediction (cf. Fig. 3B and Fig. S5). CC = cued/congruent, CI = cued/incongruent, UC = uncued/congruent, UI = uncued/incongruent. See Table 3 for details and supplementary Figures S4-S5 for (uncorrected) whole-brain results.

Table 3. Brain areas showing significant activation differences during movement execution related to prior (cued) movement planning and visuomotor mapping predictions.

Execution after Movement type cued vs uncued		MNI			Peak	
Anatomical location	*Voxels*	*x*	*y*	*z*	*F*	*p_{FWE}*
L. Anterior intraparietal sulcus	12	-42	-44	50	18.88	0.020
L. Superior frontal gyrus (PMd)	7	-20	4	70	18.69	0.021
Ventromedial prefrontal cortex	3	0	28	-6	27.74	0.047*
Execution after Mapping cued congruent vs incongruent		MNI			Peak	
Anatomical location	*Voxels*	*x*	*y*	*z*	*F*	*p_{FWE}*
L. Precentral / middle frontal gyrus (PMd)	14	-26	-6	50	21.59	0.017
L. Anterior intraparietal sulcus	7	-36	-50	42	20.85	0.022
L. Anterior insular cortex	5	-30	24	0	20.51	0.025

* whole-brain analysis.

We also tested for potential 'expectation violation' effects observable during movement execution in

trials where the virtual hand behaved contrary to the cued expectations (i.e., the cued visuomotor

mapping was invalidated). We tested for such responses in brain regions showing increased activity

during the preparation for incongruent > congruent mappings, as we assumed this contrast identified

brain areas generating visuomotor predictions. Indeed, we found significant (p_{FWE}<0.05) surprise

effects in the (pre)SMA, the left PMd and PMv, the bilateral IPL/IPS, the left anterior insula, and the

right anterior cerebellum (Fig. 6C). An additional whole-brain analysis revealed further surprise effects

in the dorsomedial prefrontal cortex, the bilateral inferior frontal gyri, the bilateral angular gyri, the

right anterior insula, the right middle temporal and fusiform gyri, the putamen and thalamus, and in

the left primary sensory and motor cortex (Fig. S5). There were no significant differences in surprise-

related activations between conditions (i.e., no voxels showed a significant main or interaction effect).

The converse contrast (expected > unexpected visuomotor mapping) yielded no significant activations

either.

Finally, following the observation that the movement cueing effects in our left-hand experiment were

strongly left-lateralized in the IPS, six of our original study's participants were invited to a separate

scanning session, in which they repeated a right-hand version of the task. These results are reported

in Figure S7. Most importantly, as in the main experiment, the preparation of cued > uncued

movements was associated with stronger BOLD signal in bilateral frontoparietal areas and the V5.

Within the left, compared with the right IPS, the effects were more extensive (spanning to anterior

portions of the IPS only in the left hemisphere) and stronger (all peak T-values were higher in the left

> right PPC). Again, the preparation of movements with an expected incongruent > congruent

visuomotor mapping was associated with stronger BOLD signal in the cerebellum and in bilateral

frontoparietal areas. Contrasting the execution of uncued > cued movements likewise revealed

activation differences corresponding to those in the main experiment; and again stronger left-

hemispheric effects in the PPC. In sum, the pattern of results obtained during *right-hand* planning and

movement replicated our main results; together, our results suggest that motor planning, in our

design, predominantly engaged the left IPS for either hand side.

Discussion

In our non-spatial delayed movement task with altered visuomotor mapping, delay period activity in

the premotor cortex (PMd, PMv, and the SMA) and the PPC (IPS and SPL) was increased if one of two

alternative movements was cued ahead of execution. This supports the general association of those

regions with motor planning (see Introduction) and aligns with frontoparietal BOLD signal increases

observed during cued reach planning (Fernandez-Ruiz et al., 2007; Lindner et al., 2010; Gertz & Fiehler,

2015; Pilacinski et al., 2018). The PMd and aIPS showed the inverse pattern during execution (relatively

higher BOLD when executing uncued movements), as similarly observed in previous uncued vs cued

effector selection tasks (Bernier et al., 2012). We propose this indicates the fast establishment of an

ad-hoc motor plan, which our participants did not already prepare during the delay (cf. Ames et al.,

2014; Pilacinski et al., 2018). In other words, our participants seemed to simply wait for the execution

prompt in the uncued trials, instead of preparing both possible movements during the delay. Whether

(or perhaps rather: when) the motor system plans actions serially, one at a time, or multiple potential

plans simultaneously, is a key question of the current motor control literature—with evidence for both

cases (e.g., Dekleva et al., 2018; Cisek & Kalaska, 2002, 2005; Cui & Andersen, 2011). The results of

such studies depend, among other things, on the kind of decision, the number of alternatives, and the

kind of movement used in each case. For instance, "nonspatial motor decisions" – like in our task –

may be processed more serially than others (Cui & Andersen, 2011); intransitive actions may also be

processed differently than goal-directed actions (see below). In our case, the frontoparietal BOLD

pattern and the behavioral reaction time differences clearly indicated the establishment of one specific

movement plan during the delay of cued trials, rather than parallel planning of both possible

movements (and subsequent selection of the prompted movement) during the delay of uncued trials.

Our second key finding was the relatively higher cerebellar activity during the preparation for

incongruent > congruent visuomotor mappings. In addition to its higher amplitude, the (right)

cerebellar hemodynamic response was relatively delayed (i.e., it peaked later) and wider in those

cases. This suggests that, when the associated visual movement feedback was expected to be

incongruent (i.e., inverted) while otherwise identical muscle (hand) movements were planned,

cerebellar neural activity was elevated and more sustained. Thereby, in the right cerebellum, the

amplitude of the BOLD response during preparation for incongruence even exceeded that observed

during actual movement execution. Note that the incongruent virtual hand movements were

predictable in principle from (planned) motor signals, just like the congruent ones, because they

reflected the (inverted) sensor values from the data glove; i.e., their kinematics were also controlled

by the participant's hand movements. Taken together, we propose that the associated stronger

cerebellar delay activity likely indicated an anticipatory update of forward predictions of visual

movement feedback—of "What my hand movement will look like"—specifically in those conditions

where the expected visual hand movement feedback violated the participants' life-long learned

visuomotor associations (i.e., of "congruence" between executed hand movements and their visual

feedback). Conversely, the 'natural' predictions of visuomotor congruence could have been more

easily and more quickly generated, requiring less computational resources (cf. Kavroulakis et al., 2022).

That the preparation for incongruent mappings increased cerebellar BOLD regardless of whether a

specific movement was planned or not (as indicated by the significant conjunction contrast of

preparation for incongruence > congruence across cued and uncued movements) suggests a general

update of the effector's forward model to a nonstandard visuomotor mapping; i.e., not only of those

predictions associated with the specific movement selected for execution.

It has long been proposed that the cerebellum implements forward models (Miall et al., 1993; Wolpert

et al., 1998). Lobule VI, specifically, responds to unpredicted sensory movement feedback (Blakemore

et al., 1998, 2001; Kilteni et al., 2023), including visuomotor incongruencies (Leube et al., 2003;

Limanowski et al., 2017; Van Kemenade et al., 2019). During sensorimotor adaptation, cerebellar

activity may indicate trial-to-trial updates of forward models by sensory prediction errors (Shadmehr

& Krakauer 2008; Izawa et al., 2012; see Tseng et al. 2007; Synofzik et al., 2008; Donchin et al. 2012;

Tzvi et al. 2022 for visuomotor adaptation, specifically). Our results suggest that cerebellar predictions

—likely informing state estimates about visuomotor mapping—can be updated ahead of, i.e., without

movement execution, and thus without any associated sensory prediction errors. This supports the

hypothesized generation of forward sensory predictions during action planning (see Introduction). The

other main regions showing increased activity during the preparation for incongruence were the PMd,

which has been shown to implement adjustments to "nonstandard sensorimotor mapping" (Wise et

al., 1996); and the SMA. The SMA (the dorsomedial frontal cortex in general) is involved in processing

conflicts during action selection (Taylor et al., 2007; Duque et al., 2013), error learning (Ulllsperger et

al., 2014), and visuomotor adaptation (Grafton et al., 2008; Nahab et al., 2011; Limanowski et al., 2017;

Ohata et al., 2020). Furthermore, electrophysiological recordings suggested that the "readiness

potential", which likely originates from the SMA, may also represent a prediction of (voluntary) action

feedback prior to execution—much like proposed by the efference copy concept (Reznik et al., 2018;

Pinhero et al., 2020; Ody et al., 2023). Both premotor areas could therefore have contributed to

forward modeling (although the clearest predictive effects were observed in the cerebellum), or more

generally anticipated the upcoming conflict (here: a nonstandard visuomotor mapping).

Thirdly, and crucially, the left aIPS showed a significant interaction effect, characterized by relatively

increased BOLD during the planning of a pre-specified (cued > uncued) movement under an expected

nonstandard (incongruent > congruent) visuomotor mapping. While several fronto-parietal areas

showed overlapping activations by both main effects (in line with suggestions that these regions may

be involved in more than one type of computation for motor control, cf. Scott, 2012), only the left

aIPS/IPL showed a significant interaction effect. Thus, movement planning in the aIPS was more

metabolically demanding when nonstandard>standard visuomotor mappings were expected. We

propose that, while cerebellar activity could be linked to forward model updates, the interaction effect

in the aIPS indicated a different computation; namely, the *integration* of the respective updated

(cerebellar) forward predictions of nonstandard visual body movement feedback with the state

estimate used for movement planning. The fact the interaction was confined to the anterior part of

the IPS fits with this region's proposed role of a specifically "body-related" state estimator (Medendorp

& Heed, 2019; Kuang et al., 2015). It also aligns with previous demonstrations of the aIPS's key role in

adaptive visuomotor control and upper limb state estimation (Desmurget et al., 1999; Brass et al.,

2001; Ehrsson et al., 2004; Grefkes & Fink, 2005; de Lange et al., 2006; Zimmermann et al., 2012;

Limanowski & Blankenburg, 2016, 2017; Limanowski, 2022). Generally, it supports the notion of the

PPC as a high-level motor control region (Andersen and Buneo 2002; Blakemore & Sirigu, 2003).

Our results highlighted specifically the *left* aIPS, irrespective of the hand side used (i.e., analogous

effects and topography in our main, left-hand experiment and a right-hand replication, see

Supplementary Material). This speaks to previous findings: The PPC represents actions from contra-

and ipsilateral hands (Gallivan et al., 2013; Turella et al., 2020); and the left (but not right) PPC has

been found activated during the selection, preparation, and execution of hand actions irrespective of

hand laterality (Schluter et al., 2001; Rushworth et al., 2001; Grafton et al., 2002; Hardwick et al., 2013).

The left PPC has also been identified as playing a key role in motor imagery, visuomotor adaptation,

and the visuoproprioceptive integration it entails, for the left and right hand (Bonda et al., 1995; Servos

et a., 2002; de Lange et al., 2006; Mutha et al., 2011; Hétu et al., 2013; Limanowski & Blankenburg,

2017). Notably, left PPC lesions have been associated with ideamotor apraxia (Haaland et al., 2000);

fMRI results correspondingly suggest a "left-lateralized praxis network", centered on the PPC, which

plans actions with either hand (Bohlhalter et al., 2009). Specifically, intransitive actions (i.e., without

object interaction) may be more strongly left-lateralized than transitive ones (Króliczak & Frey, 2009).

Our results, obtained with such intransitive actions, fit very well into this view; moreover, they show

that the left PPC is essential for integrating visual movement predictions with the respective motor

plans.

Two regions showing a main effect of planning but not visuomotor incongruence were the PMv and

the V5. The PMv is clearly linked to upper limb motor control (Graziano, 1999); the V5 activation could

indicate the generation of visual postural or motion templates (Zimmermann et al., 2012, 2016; Yon et

al., 2018), perhaps as 'simulated' sensory input for state estimation. Finally, the cerebellum, the SMA,

and several other regions that increased activity during the preparation for incongruent visuomotor

mappings showed surprise related responses. Thus, brain regions updating predictions also responded

to the respective prediction errors. That surprise responses did not differ between congruent and

incongruent mappings suggests that the forward model's predictions were, generally, successfully

updated in response to the cues. Further surprise responses were located in medial and inferior frontal

areas associated with general error detection (Ullsperger et al., 2014; cf. Quirmbach & Limanowski,

2022) and in temporoparietal areas implied in visuomotor mismatch detection (Leube et al., 2003;

Balslev et al., 2005; Farrer et al., 2008; van Kemenade et al., 2017, 2019; Limanowski et al., 2017). That

not all of the 'predictive' regions showed surprise responses, and vice versa, can potentially help refine

neuroanatomical models of visuomotor adaptation (Synofzik et al., 2006; Grafton et al., 2008).

Together, our results support the idea of forward sensory prediction during action planning, and

associate the underlying computations mainly with the cerebellum and the PPC. This opens up

interesting avenues for future work, for instance, using brain stimulation. Thus, previous work has

shown that transcranial magnetic or direct current stimulation of the aIPS or the cerebellum directly

influenced adaptive motor control (Desmurget et al., 1999; Yavari et al., 2016; Cao et al., 2017; cf.

Weightman et al., 2023). Stimulation of frontoparietal and cerebellar areas improved sensorimotor

delay detection and recalibration in schizophrenic patients, which presumably suffer from deficits in

forward modeling (Straube et al., 2020; Schmitter & Straube, 2024; cf. Voss et al., 2010). One question

following from our work is whether such cerebellar or posterior parietal manipulations would similarly

affect the updating and integration of sensory predictions during action *planning*. Future studies could

also use multivariate approaches to reveal the encoding of more abstract action planning related

information such as, e.g., different sensory content representations (cf. Ariani et al., 2015, 2018;

Gallivan et al., 2013; Yon et al., 2018; Turella et al., 2020). Our study raises further questions: One

limitation of our design were the relatively frequent catch trials that more likely followed invalid

(surprising) than valid movements. Participants could thus have associated surprising movements with

subsequent button presses, introducing a bias by movement preparation, initiation, or reward

expectations. Thus, some surprise effects, e.g. in motor areas, could partly reflect anticipatory or more

extensive movements of the responding (right) hand; which needs to be clarified by future work.

However, this did not affect the between-condition contrasts of the movement (or preparation)

periods, as surprise and catch trials were balanced across conditions. Furthermore, we cannot exclude

potential eye movement influences on our results (cf. Gorbet & Sergio, 2016). However, our non-

spatial movement task and the instructed central fixation rendered systematic eye movements

useless; and our control experiment suggested only minimal differences in gaze behavior between

conditions.

To conclude, our results suggest visual feedback prediction is an inherent component of motor

planning; and a key role of the cerebellum and the left aIPS in the underlying forward predictions and

integrative processes, respectively.

Conflict of interest statement: The authors declare no competing financial interests.

Acknowledgments: This work was funded by the German Research Foundation (DFG, Deutsche Forschungsgemeinschaft) as part of Germany's Excellence Strategy – EXC 2050/1 – Project ID 390696704 – Cluster of Excellence "Centre for Tactile Internet with Human-in-the-Loop" (CeTI) of Technische Universität Dresden. JL was supported by a Freigeist Fellowship of the VolkswagenStiftung (AZ 97-932). We thank Gesche Vigh for help with implementing the virtual reality.

References

Andersen, R. A., & Buneo, C. A. (2002). Intentional maps in posterior parietal cortex. Annual review of neuroscience, 25(1), 189-220.

Ames, K. C., Ryu, S. I., & Shenoy, K. V. (2014). Neural Dynamics of Reaching following Incorrect or Absent Motor Preparation. *Neuron*, *81*(2), 438–451. https://doi.org/10.1016/j.neuron.2013.11.003

Ariani, G., Wurm, M. F., & Lingnau, A. (2015). Decoding internally and externally driven movement plans. *Journal of Neuroscience, 35*(42), 14160-14171.

Ariani, G., Oosterhof, N. N., & Lingnau, A. (2018). Time-resolved decoding of planned delayed and immediate prehension movements. *Cortex*, *99*, 330–345. https://doi.org/10.1016/j.cortex.2017.12.007

Ariani, G., Pruszynski, J. A., & Diedrichsen, J. (2022). Motor planning brings human primary somatosensory cortex into action-specific preparatory states. *elife, 11*, e69517.

Arikan, B. E., van Kemenade, B. M., Podranski, K., Steinsträter, O., Straube, B., & Kircher, T. (2019). Perceiving your hand moving: BOLD suppression in sensory cortices and the role of the cerebellum in the detection of feedback delays. Journal of Vision, 19(14), 4-4.

Balslev, D., Nielsen, F. Å., Paulson, O. B., & Law, I. (2005). Right Temporoparietal Cortex Activation during Visuo-proprioceptive Conflict. *Cerebral Cortex*, *15*(2), 166–169. https://doi.org/10.1093/cercor/bhh119

Bernier, P.-M., Cieslak, M., & Grafton, S. T. (2012). Effector selection precedes reach planning in the dorsal parietofrontal cortex. *Journal of Neurophysiology*, *108*(1), 57–68. https://doi.org/10.1152/jn.00011.2012

Blakemore, S. J., & Sirigu, A. (2003). Action prediction in the cerebellum and in the parietal lobe. Experimental Brain Research, 153, 239-245.

Blakemore, S.-J., Wolpert, D. M., & Frith, C. D. (1998). Central cancellation of self-produced tickle sensation. *Nature Neuroscience*, *1*(7), 635–640. https://doi.org/10.1038/2870

Bohlhalter, S., Hattori, N., Wheaton, L., Fridman, E., Shamim, E. A., Garraux, G., & Hallett, M. (2009). Gesture subtype–dependent left lateralization of praxis planning: An event-related fMRI study. *Cerebral cortex, 19*(6), 1256-1262.

Bonda, E., Petrides, M., Frey, S., & Evans, A. (1995). Neural correlates of mental transformations of the body-in-space. *Proceedings of the National Academy of Sciences*, *92*(24), 11180–11184. https://doi.org/10.1073/pnas.92.24.11180

Brass, M., Zysset, S., & von Cramon, D. Y. (2001). The Inhibition of Imitative Response Tendencies. *NeuroImage*, *14*(6), 1416–1423. https://doi.org/10.1006/nimg.2001.0944

Cao, L., Veniero, D., Thut, G., & Gross, J. (2017). Role of the cerebellum in adaptation to delayed action effects. Current Biology, 27(16), 2442-2451.

Churchland, M. M., Cunningham, J. P., Kaufman, M. T., Ryu, S. I., & Shenoy, K. V. (2010). Cortical Preparatory Activity: Representation of Movement or First Cog in a Dynamical Machine? *Neuron*, *68*(3), 387–400. https://doi.org/10.1016/j.neuron.2010.09.015

Cisek, P., & Kalaska, J. F. (2002). Simultaneous Encoding of Multiple Potential Reach Directions in Dorsal Premotor Cortex. *Journal of Neurophysiology*, *87*(2), 1149–1154. https://doi.org/10.1152/jn.00443.2001

Cisek, P., & Kalaska, J. F. (2005). Neural Correlates of Reaching Decisions in Dorsal Premotor Cortex: Specification of Multiple Direction Choices and Final Selection of Action. *Neuron*, *45*(5), 801–814. https://doi.org/10.1016/j.neuron.2005.01.027

Crammond, D. J., & Kalaska, J. F. (1989). Neuronal activity in primate parietal cortex area 5 varies with intended movement direction during an instructed-delay period. *Experimental Brain Research*, *76*(2). https://doi.org/10.1007/BF00247902

Cui, H., & Andersen, R. A. (2011). Different representations of potential and selected motor plans by distinct parietal areas. Journal of Neuroscience, 31(49), 18130-18136.

Cunnington, R., Bradshaw, J. L., & Iansek, R. (1996). The role of the supplementary motor area in the control of voluntary movement. *Human Movement Science*, *15*(5), 627–647. https://doi.org/10.1016/0167-9457(96)00018-8

de Lange, F. P., Helmich, R. C., & Toni, I. (2006). Posture influences motor imagery: An fMRI study. *NeuroImage*, *33*(2), 609–617. https://doi.org/10.1016/j.neuroimage.2006.07.017

Dekleva, B. M., Kording, K. P., & Miller, L. E. (2018). Single reach plans in dorsal premotor cortex during a two-target task. *Nature Communications*, *9*(1), 3556. https://doi.org/10.1038/s41467-018-05959-y

Desmurget, M., Epstein, C. M., Turner, R. S., Prablanc, C., Alexander, G. E., & Grafton, S. T. (1999). Role of the posterior parietal cortex in updating reaching movements to a visual target. *Nature Neuroscience*, *2*(6), 563–567. https://doi.org/10.1038/9219

Donchin, O., Rabe, K., Diedrichsen, J., Lally, N., Schoch, B., Gizewski, E. R., & Timmann, D. (2012). Cerebellar regions involved in adaptation to force field and visuomotor perturbation. Journal of neurophysiology, 107(1), 134-147.

Duque, J., Olivier, E., & Rushworth, M. (2013). Top–down inhibitory control exerted by the medial frontal cortex during action selection under conflict. Journal of cognitive neuroscience, 25(10), 1634-1648.

Ehrsson, H. H., Spence, C., & Passingham, R. E. (2004). That's My Hand! Activity in Premotor Cortex Reflects Feeling of Ownership of a Limb. *Science*, *305*(5685), 875–877. https://doi.org/10.1126/science.1097011

Eickhoff, S. B., Stephan, K. E., Mohlberg, H., Grefkes, C., Fink, G. R., Amunts, K., & Zilles, K. (2005). A new SPM toolbox for combining probabilistic cytoarchitectonic maps and functional imaging data. *NeuroImage*, *25*(4), 1325–1335. https://doi.org/10.1016/j.neuroimage.2004.12.034

Fabbri, S., Strnad, L., Caramazza, A., & Lingnau, A. (2014). Overlapping representations for grip type and reach direction. Neuroimage, 94, 138-146.

Farrer, C., Frey, S. H., Van Horn, J. D., Tunik, E., Turk, D., Inati, S., & Grafton, S. T. (2008). The angular gyrus computes action awareness representations. *Cerebral cortex*, *18*(2), 254-261.

Fernandez-Ruiz, J., Goltz, H. C., DeSouza, J. F. X., Vilis, T., & Crawford, J. D. (2007). Human Parietal "Reach Region" Primarily Encodes Intrinsic Visual Direction, Not Extrinsic Movement Direction, in a Visual–Motor Dissociation Task. *Cerebral Cortex*, *17*(10), 2283–2292. https://doi.org/10.1093/cercor/bhl137

Ferraina, S., & Bianchi, L. (1994). Posterior parietal cortex: Functional properties of neurons in area 5 during an instructed-delay reaching task within different parts of space. *Experimental Brain Research*, *99*(1). https://doi.org/10.1007/BF00241423

Friston, K., Fletcher, P., Josephs, O., Holmes, A., Rugg, M. D., & Turner, R. (1998). Event-related fMRI: Characterizing differential responses. NeuroImage, 7, 30–40.

Gallivan, J. P., McLean, D. A., Flanagan, J. R., & Culham, J. C. (2013). Where One Hand Meets the Other: Limb-Specific and Action-Dependent Movement Plans Decoded from Preparatory Signals in Single Human Frontoparietal Brain Areas. *The Journal of Neuroscience*, *33*(5), 1991–2008. https://doi.org/10.1523/JNEUROSCI.0541-12.2013

Gertz, H., & Fiehler, K. (2015). Human posterior parietal cortex encodes the movement goal in a pro-/anti-reach task. *Journal of Neurophysiology*, *114*(1), 170–183. https://doi.org/10.1152/jn.01039.2014

Gläscher, J. (2009). Visualization of group inference data in functional neuroimaging. *Neuroinformatics*, 7, 73-82.

Gorbet, D. J., & Sergio, L. E. (2016). Don't watch where you're going: The neural correlates of decoupling eye and arm movements. *Behavioural Brain Research*, *298*, 229–240. https://doi.org/10.1016/j.bbr.2015.11.012

Grafton, S. T., Hazeltine, E., & Ivry, R. B. (2002). Motor sequence learning with the nondominant left hand: A PET functional imaging study. Experimental brain research, 146, 369-378.

Grafton, S. T., Schmitt, P., Van Horn, J., & Diedrichsen, J. (2008). Neural substrates of visuomotor learning based on improved feedback control and prediction. Neuroimage, 39(3), 1383-1395.

Graziano, M. S. A. (1999). Where is my arm? The relative role of vision and proprioception in the neuronal representation of limb position. *Proceedings of the National Academy of Sciences*, *96*(18), 10418–10421. https://doi.org/10.1073/pnas.96.18.10418

Grefkes, C., & Fink, G. R. (2005). REVIEW: The functional organization of the intraparietal sulcus in humans and monkeys: Intraparietal sulcus in monkeys and humans, C. Grefkes and G. R. Fink. *Journal of Anatomy*, *207*(1), 3–17. https://doi.org/10.1111/j.1469-7580.2005.00426.x

Haaland, K. Y., Harrington, D. L., & Knight, R. T. (2000). Neural representations of skilled movement. Brain, 123(11), 2306-2313.

Hardwick, R. M., Rottschy, C., Miall, R. C., & Eickhoff, S. B. (2013). A quantitative meta-analysis and review of motor learning in the human brain. *NeuroImage*, *67*, 283–297. https://doi.org/10.1016/j.neuroimage.2012.11.020

Henson, R. N., Price, C. J., Rugg, M. D., Turner, R., & Friston, K. J. (2002). Detecting latency differences in event-related BOLD responses: application to words versus nonwords and initial versus repeated face presentations. Neuroimage, 15(1), 83-97.

Hétu, S., Grégoire, M., Saimpont, A., Coll, M. P., Eugène, F., Michon, P. E., & Jackson, P. L. (2013). The neural network of motor imagery: an ALE meta-analysis. Neuroscience & Biobehavioral Reviews, 37(5), 930-949.

Hoshi, E., & Tanji, J. (2002). Contrasting Neuronal Activity in the Dorsal and Ventral Premotor Areas During Preparation to Reach. *Journal of Neurophysiology*, *87*(2), 1123–1128. https://doi.org/10.1152/jn.00496.2001

Izawa, J., Criscimagna-Hemminger, S. E., & Shadmehr, R. (2012). Cerebellar contributions to reach adaptation and learning sensory consequences of action. Journal of Neuroscience, 32(12), 4230-4239.

Jeannerod, M. & Decety, J. Mental motor imagery: a window into the representational stages of action. Curr. Opin. Neurobiol. 5, 727–732 (1995).

Kaufman, M. T., Churchland, M. M., Ryu, S. I., & Shenoy, K. V. (2014). Cortical activity in the null space: Permitting preparation without movement. *Nature Neuroscience*, *17*(3), 440–448. https://doi.org/10.1038/nn.3643

Kavroulakis, E., van Kemenade, B. M., Arikan, B. E., Kircher, T., & Straube, B. (2022). The effect of self-generated versus externally generated actions on timing, duration, and amplitude of blood oxygen level dependent response for visual feedback processing. Human Brain Mapping, 43(16), 4954-4969.

Kilteni, K., Andersson, B. J., Houborg, C., & Ehrsson, H. H. (2018). Motor imagery involves predicting the sensory consequences of the imagined movement. Nature communications, 9(1), 1617.

Kilteni, K., Houborg, C., & Ehrsson, H. H. (2023). Brief temporal perturbations in somatosensory reafference disrupt perceptual and neural attenuation and increase supplementary motor area–cerebellar connectivity. Journal of Neuroscience.

Klaes, C., Westendorff, S., Chakrabarti, S., & Gail, A. (2011). Choosing Goals, Not Rules: Deciding among Rule-Based Action Plans. *Neuron*, *70*(3), 536–548. https://doi.org/10.1016/j.neuron.2011.02.053

Króliczak, G., & Frey, S. H. (2009). A common network in the left cerebral hemisphere represents planning of tool use pantomimes and familiar intransitive gestures at the hand-independent level. *Cerebral Cortex, 19*(10), 2396-2410.

Kuang, S., Morel, P., & Gail, A. (2015). Planning Movements in Visual and Physical Space in Monkey Posterior Parietal Cortex. *Cerebral Cortex*, bhu312. https://doi.org/10.1093/cercor/bhu312

Kufer, K., Schmitter, C. V., Kircher, T., & Straube, B. (2024). Temporal recalibration in response to delayed visual feedback of active versus passive actions: an fMRI study. Scientific Reports, 14(1), 4632.

Leube, D., Knoblich, G., Erb, M., Grodd, W., Bartels, M., & Kircher, T. T. (2003). The neural correlates of perceiving one's own movements. *NeuroImage*, *20*(4), 2084–2090. https://doi.org/10.1016/j.neuroimage.2003.07.033

Limanowski, J., & Blankenburg, F. (2016). Integration of Visual and Proprioceptive Limb Position Information in Human Posterior Parietal, Premotor, and Extrastriate Cortex. *The Journal of Neuroscience*, *36*(9), 2582–2589. https://doi.org/10.1523/JNEUROSCI.3987-15.2016

Limanowski, J., & Blankenburg, F. (2017). Posterior parietal cortex evaluates visuoproprioceptive congruence based on brief visual information. *Scientific Reports*, *7*(1), 16659. https://doi.org/10.1038/s41598-017-16848-7

Limanowski, J., & Friston, K. (2020). Attentional Modulation of Vision Versus Proprioception During Action. *Cerebral Cortex*, *30*(3), 1637–1648. https://doi.org/10.1093/cercor/bhz192

Limanowski, J., Kirilina, E., & Blankenburg, F. (2017). Neuronal correlates of continuous manual tracking under varying visual movement feedback in a virtual reality environment. *NeuroImage*, *146*, 81–89. https://doi.org/10.1016/j.neuroimage.2016.11.009

Lindner, A., Iyer, A., Kagan, I., & Andersen, R. A. (2010). Human Posterior Parietal Cortex Plans Where to Reach and What to Avoid. *The Journal of Neuroscience*, *30*(35), 11715–11725. https://doi.org/10.1523/JNEUROSCI.2849-09.2010

Lotze, M., Montoya, P., Erb, M., Hülsmann, E., Flor, H., Klose, U., Birbaumer, N., & Grodd, W. (1999). Activation of Cortical and Cerebellar Motor Areas during Executed and Imagined Hand Movements: An fMRI Study. *Journal of Cognitive Neuroscience*, *11*(5), 491–501. https://doi.org/10.1162/089892999563553

Makoshi, Z., Kroliczak, G., & Van Donkelaar, P. (2011). Human Supplementary Motor Area Contribution to Predictive Motor Planning. *Journal of Motor Behavior*, *43*(4), 303–309. https://doi.org/10.1080/00222895.2011.584085

Medendorp, W. P., & Heed, T. (2019). State estimation in posterior parietal cortex: Distinct poles of environmental and bodily states. *Progress in neurobiology, 183*, 101691.

Miall, R. C., Weir, D. J., Wolpert, D. M., & Stein, J. F. (1993). Is the Cerebellum a Smith Predictor? *Journal of Motor Behavior*, *25*(3), 203–216. https://doi.org/10.1080/00222895.1993.9942050

Miall, R. C., & Wolpert, D. M. (1996). Forward models for physiological motor control. Neural networks, 9(8), 1265-1279.

Michaels, J. A., Dann, B., Intveld, R. W., & Scherberger, H. (2018). Neural Dynamics of Variable Grasp-Movement Preparation in the Macaque Frontoparietal Network. *The Journal of Neuroscience*, *38*(25), 5759–5773. https://doi.org/10.1523/JNEUROSCI.2557-17.2018

Mutha, P. K., Sainburg, R. L., & Haaland, K. Y. (2011). Left parietal regions are critical for adaptive visuomotor control. Journal of Neuroscience, 31(19), 6972-6981.

Nahab, F. B., Kundu, P., Gallea, C., Kakareka, J., Pursley, R., Pohida, T., Miletta, N., Friedman, J., & Hallett, M. (2011). The Neural Processes Underlying Self-Agency. *Cerebral Cortex*, *21*(1), 48–55. https://doi.org/10.1093/cercor/bhq059

Ody, E., Kircher, T., Straube, B., & He, Y. (2023). Pre-movement event-related potentials and multivariate pattern of EEG encode action outcome prediction. Human Brain Mapping, 44(17), 6198-6213.

Ogawa, K., Inui, T., & Sugio, T. (2006). Separating brain regions involved in internally guided and visual feedback control of moving effectors: An event-related fMRI study. *NeuroImage*, *32*(4), 1760–1770. https://doi.org/10.1016/j.neuroimage.2006.05.012

Ohata, R., Asai, T., Kadota, H., Shigemasu, H., Ogawa, K., & Imamizu, H. (2020). Sense of Agency Beyond Sensorimotor Process: Decoding Self-Other Action Attribution in the Human Brain. *Cerebral Cortex*, *30*(7), 4076–4091. https://doi.org/10.1093/cercor/bhaa028

Parr, T., Limanowski, J., Rawji, V., & Friston, K. (2021). The computational neurology of movement under active inference. Brain, 144(6), 1799-1818.

Pilacinski, A., Wallscheid, M., & Lindner, A. (2018). Human posterior parietal and dorsal premotor cortex encode the visual properties of an upcoming action. *PLOS ONE*, *13*(10), e0198051. https://doi.org/10.1371/journal.pone.0198051

Pinheiro, A. P., Schwartze, M., Gutierrez-Dominguez, F., & Kotz, S. A. (2020). Real and imagined sensory feedback have comparable effects on action anticipation. Cortex, 130, 290-301.

Quirmbach, F., & Limanowski, J. (2022). A Crucial Role of the Frontal Operculum in Task-Set Dependent Visuomotor Performance Monitoring. *Eneuro*, *9*(2), ENEURO.0524-21.2021. https://doi.org/10.1523/ENEURO.0524-21.2021

Reznik, D., Simon, S., & Mukamel, R. (2018). Predicted sensory consequences of voluntary actions modulate amplitude of preceding readiness potentials. Neuropsychologia, 119, 302–307.

Rosenbaum, D. A. (1980). Human movement initiation: specification of arm, direction, and extent. Journal of Experimental Psychology: General, 109(4), 444.

Rushworth, M. F., Krams, M., & Passingham, R. E. (2001). The attentional role of the left parietal cortex: the distinct lateralization and localization of motor attention in the human brain. Journal of cognitive neuroscience, 13(5), 698-710.

Sakai, K., Hikosaka, O., Miyauchi, S., Sasaki, Y., Fujimaki, N., & Pütz, B. (1999). Presupplementary Motor Area Activation during Sequence Learning Reflects Visuo-Motor Association. *The Journal of Neuroscience*, *19*(10), RC1–RC1. https://doi.org/10.1523/JNEUROSCI.19-10-j0002.1999

Schaffelhofer, S., Agudelo-Toro, A., & Scherberger, H. (2015). Decoding a Wide Range of Hand Configurations from Macaque Motor, Premotor, and Parietal Cortices. *The Journal of Neuroscience*, *35*(3), 1068–1081. https://doi.org/10.1523/JNEUROSCI.3594-14.2015

Schluter, N. D., Krams, M., Rushworth, M. F. S., & Passingham, R. E. (2001). Cerebral dominance for action in the human brain: the selection of actions. Neuropsychologia, 39(2), 105-113.

Schmitter, C. V., & Straube, B. (2024). Facilitation of sensorimotor temporal recalibration mechanisms by cerebellar tDCS in patients with schizophrenia spectrum disorders and healthy individuals. Scientific Reports, 14(1), 2627.

Scott, S. H. (2012). The computational and neural basis of voluntary motor control and planning. Trends in cognitive sciences, 16(11), 541-549.

Servos, P., Osu, R., Santi, A., & Kawato, M. (2002). The neural substrates of biological motion perception: an fMRI study. *Cerebral Cortex, 12*(7), 772-782.

Shadmehr, R., & Krakauer, J. W. (2008). A computational neuroanatomy for motor control. Experimental brain research, 185, 359-381.

Shadmehr, R., Smith, M. A., & Krakauer, J. W. (2010). Error correction, sensory prediction, and adaptation in motor control. Annual review of neuroscience, 33, 89-108.

Stanley, J., & Miall, R. C. (2007). Functional activation in parieto-premotor and visual areas dependent on congruency between hand movement and visual stimuli during motor-visual priming. Neuroimage, 34(1), 290-299.

Stanley, J., & Miall, R. C. (2009). Using predictive motor control processes in a cognitive task: behavioral and neuroanatomical perspectives. In: D. Sternad (ed.), Progress in motor control: A multidisciplinary perspective, Springer Science & Business Media, LLC 2009, 337-354.

Straube, B., van Kemenade, B. M., Kircher, T., & Schülke, R. (2020). Transcranial direct current stimulation improves action-outcome monitoring in schizophrenia spectrum disorder. Brain communications, 2(2), fcaa151.

Synofzik, M., Thier, P., & Lindner, A. (2006). Internalizing agency of self-action: perception of one's own hand movements depends on an adaptable prediction about the sensory action outcome. Journal of neurophysiology, 96(3), 1592-1601.

Synofzik, M., Lindner, A., & Thier, P. (2008). The cerebellum updates predictions about the visual consequences of one's behavior. *Current Biology, 18*(11), 814-818.

Tanji, J., & Evarts, E. V. (1976). Anticipatory activity of motor cortex neurons in relation to direction of an intended movement. *Journal of Neurophysiology*, *39*(5), 1062–1068. https://doi.org/10.1152/jn.1976.39.5.1062

Taylor, P. C., Nobre, A. C., & Rushworth, M. F. (2007). Subsecond changes in top–down control exerted by human medial frontal cortex during conflict and action selection: a combined transcranial magnetic stimulation–electroencephalography study. Journal of Neuroscience, 27(42), 11343-11353.

Todorov, E., & Jordan, M. I. (2002). Optimal feedback control as a theory of motor coordination. Nature neuroscience, 5(11), 1226-1235.

Tseng YW, Diedrichsen J, Krakauer JW, Shadmehr R, Bastian AJ (2007) Sensory prediction errors drive cerebellum-dependent adaptation of reaching. J Neurophysiol 98:54–62https://doi.org/10.1523/JNEUROSCI.2451-11.2011

Turella, L., Rumiati, R., & Lingnau, A. (2020). Hierarchical action encoding within the human brain. Cerebral cortex, 30(5), 2924-2938.

Tzvi, E., Loens, S., & Donchin, O. (2022). Mini-review: the role of the cerebellum in visuomotor adaptation. The Cerebellum, 21(2), 306-313.

Ullsperger, M., Danielmeier, C., & Jocham, G. (2014). Neurophysiology of Performance Monitoring and Adaptive Behavior. *Physiological Reviews*, *94*(1), 35–79. https://doi.org/10.1152/physrev.00041.2012

Van Kemenade, B. M., Arikan, B. E., Kircher, T., & Straube, B. (2017). The angular gyrus is a supramodal comparator area in action–outcome monitoring. *Brain Structure and Function*, *222*(8), 3691–3703. https://doi.org/10.1007/s00429-017-1428-9

Van Kemenade, B. M., Arikan, B. E., Podranski, K., Steinsträter, O., Kircher, T., & Straube, B. (2019). Distinct Roles for the Cerebellum, Angular Gyrus, and Middle Temporal Gyrus in Action–Feedback Monitoring. *Cerebral Cortex*, *29*(4), 1520–1531. https://doi.org/10.1093/cercor/bhy048

Verhagen, L., Dijkerman, H. C., Medendorp, W. P., & Toni, I. (2012). Cortical Dynamics of Sensorimotor Integration during Grasp Planning. *The Journal of Neuroscience*, *32*(13), 4508–4519. https://doi.org/10.1523/JNEUROSCI.5451-11.2012

Voss, M., Moore, J., Hauser, M., Gallinat, J., Heinz, A., & Haggard, P. (2010). Altered awareness of action in schizophrenia: a specific deficit in predicting action consequences. Brain, 133(10), 3104-3112.

Weightman, M., Lalji, N., Lin, C. H. S., Galea, J. M., Jenkinson, N., & Miall, R. C. (2023). Short duration event related cerebellar TDCS enhances visuomotor adaptation. Brain stimulation, 16(2), 431-441.

Wise, S. P., di Pellegrino, G., & Boussaoud, D. (1996). The premotor corte× and nonstandard sensorimotor mapping. *Canadian Journal of Physiology and Pharmacology*, *74*(4), 469–482. https://doi.org/10.1139/cjpp-74-4-469

Wolpert, D. M., & Kawato, M. (1998). Multiple paired forward and inverse models for motor control. *Neural networks, 11*(7-8), 1317-1329.

Wolpert, D. M., Miall, R. C., & Kawato, M. (1998). Internal models in the cerebellum. Trends in cognitive sciences, 2(9), 338-347.

Wolpert, D. M., & Ghahramani, Z. (2000). Computational principles of movement neuroscience. Nature neuroscience, 3(11), 1212-1217.

Yavari, F., Mahdavi, S., Towhidkhah, F., Ahmadi-Pajouh, M. A., Ekhtiari, H., & Darainy, M. (2016). Cerebellum as a forward but not inverse model in visuomotor adaptation task: a tDCS-based and modeling study. Experimental brain research, 234, 997-1012.

Yomogida, Y., Sugiura, M., Sassa, Y., Wakusawa, K., Sekiguchi, A., Fukushima, A., Takeuchi, H., Horie, K., Sato, S., & Kawashima, R. (2010). The neural basis of agency: An fMRI study. *NeuroImage*, *50*(1), 198–207. https://doi.org/10.1016/j.neuroimage.2009.12.054

Yon, D., Gilbert, S. J., de Lange, F. P., & Press, C. (2018). Action sharpens sensory representations of expected outcomes. *Nature Communications*, *9*(1), 4288. https://doi.org/10.1038/s41467-018-06752-7

Zimmermann, M., Verhagen, L., de Lange, F. P., & Toni, I. (2016). The extrastriate body area computes desired goal states during action planning. Eneuro, 3(2).

Zimmermann, M., Meulenbroek, R. G. J., & de Lange, F. P. (2012). Motor Planning Is Facilitated by

Adopting an Action's Goal Posture: An fMRI Study. *Cerebral Cortex*, *22*(1), 122–131.

https://doi.org/10.1093/cercor/bhr098