Visuomotor prediction during action planning in the human frontoparietal cortex and cerebellum

Abbreviated title: Action planning and visuomotor prediction

Authors: Felix Quirmbach[1,2] and Jakub Limanowski[2,3]

[1]Faculty of Psychology, Technical University of Dresden, 01069 Dresden, Germany

[2]Center for Tactile Internet with Human-in-the-Loop, Technical University of Dresden, 01187 Dresden, Germany

[3]Institute of Psychology, University of Greifswald**,** 17486 Greifswald, Germany

Corresponding author email address: jakub.limanowski@uni-greifswald.de

Conflict of interest statement: The authors declare no competing financial interests.

Acknowledgments: This work was funded by the German Research Foundation (DFG, Deutsche Forschungsgemeinschaft) as part of Germany's Excellence Strategy – EXC 2050/1 – Project ID 390696704 – Cluster of Excellence "Centre for Tactile Internet with Human-in-the-Loop" (CeTI) of Technische Universität Dresden. JL was supported by a Freigeist Fellowship of the VolkswagenStiftung (AZ 97-932). We thank Gesche Vigh for help with implementing the virtual reality.

Abstract

A prominent concept in the domain of motor control is that of forward models in the brain, which predict the sensory consequences of one's actions for improved state estimation and on-line control. This idea can be extended to action planning; i.e., assuming that sensory forward predictions are also issued during the preparation of movements, in the absence of overt movement or related sensory feedback. To test this idea, we combined a delayed movement task with a virtual reality based manipulation of visuomotor congruence during functional magnetic resonance imaging (fMRI). Participants (18 female, 22 male) executed hand movements after a delay. During the delay, two aspects of the upcoming movement could be pre-cued: the movement type (opening or closing) and the visuomotor mapping (i.e., of executed hand movements and virtual hand movement feedback). Frontoparietal areas showed increased delay period activity when a specific movement was cued (vs uncued) and when an incongruent (inverted vs congruent) visuomotor mapping was cued. The cerebellum showed increased responses during the preparation for incongruent mappings independently of planning (i.e., cued and uncued movements). The left anterior intraparietal sulcus (aIPS) showed an interaction effect; i.e., a relatively stronger activity increase when a specific movement was prepared under expected visuomotor incongruence. These results suggest that motor planning entails a forward prediction of visual body movement consequences, likely implemented by the cerebellum; which can be adjusted in anticipation of novel visuomotor mappings, and integrated with state estimates for (planned) control in frontoparietal areas, most notably, in the aIPS.

Introduction

Our current understanding of motor control largely builds upon the concept of internal forward models that predict the sensory consequences of one's movement, based on motor signals (Miall & Wolpert, 1996; Wolpert & Kawato, 1998; Shadmehr, & Krakauer, 2008). This prediction can be used to calculate a dynamic estimate of the effector's state, enabling 'on-line' control despite sensory delays (Todorov & Jordan, 2002; Wolpert & Ghahramani 2000). The respective computations have been linked to the posterior parietal cortex (PPC) and the cerebellum (Miall et al. 1993; Wolpert et al. 1998; Desmurget et al., 1999; Blakemore et al., 1998; Synofzik et al., 2008; Kilteni et al., 2023), and their interaction with (pre)motor areas and the basal ganglia implementing the task and control policies (Shadmehr & Krakauer, 2008; Scott, 2012; Parr et al., 2021). The forward predictions can be updated by sensory prediction errors to adapt to new sensorimotor mappings (e.g., altered visual movement feedback, Tseng et al., 2007; Grafton et al., 2008; Shadmehr et al., 2010).

The internal forward model framework can, in principle, also be applied to action *planning* (Stanley & Miall, 2009; Scott, 2012); in line with the proposed functional equivalence of planning and overt action control in the brain (Jeannerod & Decety, 1995; Lotze, 1999). A derivative hypothesis is that the motor system should also issue sensory predictions when merely planning actions; i.e., in the absence of overt action and sensory movement feedback (Kuang et al., 2015; Pilacinski et al., 2018; Kilteni et al., 2018).

We tested this idea by combining a delayed movement paradigm with a virtual reality based manipulation of visuomotor mapping inside an MR scanner. In delayed movement tasks, movements are pre-cued (Rosenbaum, 1980); neural activity during the delay period is thought to encode specific aspects of motor planning, such as the selection of effectors or goals, and the preparation of the appropriate control policy and motor commands (Crammond & Kalaska, 1989; Hoshi & Tanji, 2002; Fernandez-Ruiz et al., 2007; Bernier et al., 2012; Ariani et al., 2022). Predominantly, these planning processes have been linked to activity in the dorsal premotor cortex (PMd) and the PPC (Churchland et al., 2010; Cisek & Kalaska, 2005; Ferraina & Bianchi, 1994; Kaufman et al., 2014; Kuang et al., 2015;

Tanji & Evarts, 1976; Schaffelhofer et al., 2015; Gertz & Fiehler, 2015; Lindner et al., 2010; Verhagen et al., 2012).

Here, participants had to execute one of two possible hand movements after a variable delay (Fig. 1). During the delay, two aspects of the upcoming movement could be independently cued (or not): the movement type and the visuomotor mapping. We assumed that cueing movements would establish a specific motor plan during the delay; whereas in the uncued case, a fast planning process and movement execution would be initiated only upon the execution prompt (Rosenbaum, 1980; cf. Ames et al., 2014; Pilacinski et al., 2018). This should yield higher frontoparietal blood oxygenation level dependent signal (BOLD) when preparing for cued > uncued movements. If, alternatively, both movements were prepared in parallel, BOLD should show the opposite effect (uncued > cued).

By cueing incongruent (inverted) visual hand movements, we aimed to establish an anticipation of an unnatural visuomotor mapping. We assumed that participants would naturally expect visuomotor congruence, based on a life-long learned association between executed movements and their visual feedback; and that incongruent movements would violate this expectation (Yon et al., 2018). When these 'nonstandard' visuomotor mappings were cued, we expected an anticipatory adjustment of the forward model's predictions, detectable as increased delay period BOLD. As there were no spatial targets during planning and execution, these predictions should specifically relate to visual body movement feedback, i.e.: *What my hand movement will look like*. Moreover, we expected motor planning to require more neuronal resources when integrating the (updated) predictions of nonstandard visual feedback in the state estimate. We anticipated these (interaction) effects predominantly in the PPC and PMd.

Figure 1. Experimental task to manipulate motor planning and visuomotor prediction. The participants' real hand movements (left, hidden from view) were measured via a data glove and fed to a virtual hand (VH) model presented on screen (right). In each trial, participants had to execute one of two possible movements (open or close their hand) after a variable delay, starting from a neutral position (hand half-way closed). During the delay (preparation period), we presented a visual cue that could provide two kinds of predictive information: Firstly, the cue's *shape* could specify the movement type to be executed after the delay (up- or downward triangle = open or close their hand, 25% of all trials each, i.e., together 50%) or leave it ambiguous (square = upcoming movement unknown, 50% of all trials). Secondly, the cue's *color* indicated the to-be-expected visuomotor congruence; i.e., whether the virtual hand would next reflect the real hand movement (here: cyan cue) or the respective opposing movement (i.e., inverting the real hand movement, here: pink cue). Thus, we orthogonally manipulated processes related to motor planning and visuomotor prediction (the predicted mapping of visual movement feedback onto motor commands; i.e., what the hand movement will look like), respectively; in the absence of spatial movement targets during planning or execution. After a 'Go signal' and appearance of the virtual hand on screen, the participants executed the instructed hand movement while maintaining central fixation. In this particular example, the participant was cued that she would need to open her hand (triangle pointing upwards) and that the virtual hand would move incongruently during execution (pink cue), i.e., close. To ensure task engagement and attention to the cues, participants had to monitor and, in occasional catch trials, report whether the virtual hand movement matched the color-cued predictions (in 25% of trials, the virtual hand mapping was invalidly cued). Trials were separated by a brief (2s) interval. In this figure, the visual cues have been enlarged for display purposes.

Materials and Methods

Participants

30 healthy, right-handed volunteers (18 female, mean age = 25.4 years, range 19-35) participated in the experiment after signing informed consent. The experiment was approved by the ethics committee of the Technische Universität Dresden and conducted in accordance with this approval.

Experimental design and procedure

Participants lay comfortably inside an MR scanner. On their left hand, they wore an MR-compatible data glove (5DT Data Glove 14 Ultra MRI, 60 Hz sampling rate, full-speed USB 1.1 connection), which measured each finger's flexion via sewn-in optical sensors. The glove data were fed to a photo-realistic virtual hand model, which was thus controllable in real-time by the participants' hand movements. The virtual hand received an average of the fingers' (index, middle, ring, and pinky) sensor values to ensure smooth and coherent visual motion (cf. Limanowski & Friston, 2020). The participants' real hand was placed outside of their view; i.e., during the entire experiment, participants only saw the virtual hand, but not their real hand. Prior to the experiment, the glove was calibrated carefully; i.e., the virtual hand model was shown and the participant moved her fingers to establish the possible range of flexion (minimum and maximum sensor values), and the virtual hand model's finger flexion was adjusted to closely match the real hand posture and movement range according to the participant's perception. If necessary, the calibration was repeated during the pauses between individual scanning runs. Through a mirror on the head coil, participants saw a screen, onto which a projector (Casio XJ-M156, Laser/LED hybrid, 1024 x 768 pixels resolution, 60 Hz refresh rate) displayed the virtual reality environment; i.e., the instructions, cues, and the virtual hand model. The virtual environment was created with the Blender 3D graphics software package (https://www.blender.org/, Version 2.79) and its Python programming interface. The experimenter observed the virtual display (via a mirrored screen) at all times, thus ensuring task compliance.

The participants' task was to perform, in each trial, one of two types of cued hand movements (open or close, from a neutral starting position) after an instructed delay. During movement, the virtual hand model moved congruently or incongruently to the participants' actual hand movements. Each trial consisted of two periods – a *preparation period,* in which participants could be cued about various aspects of the upcoming movement (i.e., inducing motor planning and visuomotor expectations, see below); followed by an *execution period*, where participants executed the hand movements, while receiving (condition-specific, see below) visual movement feedback from the virtual hand model. See Figure 1 for an example trial.

At the beginning of each trial, participants had to bring their real, unseen hand to the neutral starting position (i.e., fingers half-way closed). For this, a small white bar was presented on screen, whose vertical position was controlled by the participants' hand posture via the data glove (i.e., the degree to which the hand was opened or closed). Participants had to align this bar with a central target (another bar, see Fig.1), which, on average, took 1.96 s (SD = 0.48 s). Upon alignment, the bar was then replaced by a central white fixation dot for 1 s. Participants were instructed to keep their gaze on this fixation dot throughout the entire trial; i.e., until the end of the execution period.

Then, a cue was presented and remained visible throughout the delay period (3 to 5 s jittered, pseudo-randomized, evenly distributed across conditions). This cue could provide participants with two kinds of predictive information: The *type of the upcoming movement* (i.e., hand opening or closing) and the *congruence of the virtual and real hand movements* during this movement (see below). These two predictions were coded by cue shape and color, respectively. Thus, in half of all trials, the cue *shape* specified the movement that the participant would need to execute next. Specifically, in 25% of trials an upward-pointing triangle informed participants that they would have to *open* their hand, while in another 25% a downward-pointing triangle informed them they would have to *close* their hand. In the remaining half of all trials, the cue was square-shaped and, thus, did not reveal the to-be-executed movement type in advance. In other words, only in half of the trials participants were able to generate a specific motor prediction about the movement they would have to execute after the delay period.

Furthermore, the *color* of the cue was predictive of whether, during the upcoming movement, the virtual hand would move congruently (i.e., reflecting the participants' real hand movement) or incongruently (performing the respective opposite, i.e., inverted, movement type) with the participant's real hand movements. In the incongruent conditions, the values fed to the hand model from the glove's sensors were inverted so that an opening movement would close the virtual hand, and vice versa. Thus, the movements of the virtual hand were controlled in real time by the participant in both cases, ensuring matching virtual and real hand kinematics during congruent and incongruent conditions. This was important because our 'incongruent' virtual hand movements were, therefore, also predictable in principle from the real hand movements; i.e., from motor signals.

The two movement types were designed to produce, from the neutral starting position, a comparable range of visual motion; i.e., cover the same Euclidean distance on screen: At the beginning of the trial, the virtual hand model was displayed on the screen in the neutral starting position. In this position, the tip of the middle finger was positioned exactly in the middle of the screen (i.e., in the same position as the fixation dot); and the Euclidean distance between this fixation dot and the finger tip for either the fully opened or closed hand position was the same. While opening vs closing the hand were not diametrical opposites in terms of actual joint rotations (cf. Fig. 2A), all conditions contained the same number of open and close movements (see above); and all fMRI contrasts were calculated on conditional contrast estimates pooled over both movement types. Thus, kinetic or kinematic differences between opening vs closing movements were balanced across conditions; i.e., any potentially related effects were factored out in our fMRI analysis.

The cue colors (pink and cyan) were counter-balanced across participants. Following the preparatory period, a 'Go' signal (white triangle, centrally presented) appeared. Again, the pointing direction of this triangle instructed participants to open (triangle pointing up) or close (triangle pointing down) their hand. Note that, if the prior cue had not yet specified the movement type (i.e., square-shaped cue), the participants would only now know which movement they should execute.

Participants were instructed to execute the movement as soon as the 'Go' signal disappeared (after 500 ms) and, instead, the virtual hand model was presented on the screen. This was done to ensure that participants did not start to move reflexively before the virtual hand was visible. Participants then had to execute the hand movement fluently, for which they had up to 2 s. Participants were instructed and trained to make similar movements in all conditions and, after completing the movement, to keep the hand in the final position (completely opened or closed) until the end of the trial. Then, the virtual hand disappeared again and a blank screen was shown for 2 s. As in the preparation period, participants still had to maintain fixation on the central dot during and post execution. In a prior training session, participants were able to practice the movements and familiarize themselves with all conditions. In sum, this experimental setup resulted in a factorial design with the factors *Movement type cueing* (Cued, Uncued) and *Visuomotor mapping* (Congruent, Incongruent); and, correspondingly, four conditions: CC (cued movement with expected congruent mapping), CI (cued movement with expected incongruent mapping), UC (uncued movement with expected congruent mapping) and UI (uncued movement with expected incongruent mapping). In other words, movement planning was orthogonal to expectations about visuomotor congruence.

One novelty of our experimental design was that there were no spatial movement targets during planning or execution. The presence of such targets, during planning or executing movements towards (or away from) those targets, can make it difficult to distinguish activity related to the processing or encoding of the properties of visual movement feedback *per se* from, for instance, spatial eye-hand coordination or the visual-spatial representation of the movement goal (see Pilacinski et al., 2018; Lindner et al., 2010, for discussions). Therefore, in our design, we ensured that the visuomotor predictions pertained exclusively to the mapping of executed and observed hand movements, in other words: a prediction of what the planned hand movement would look like.

One implication of this design was that participants could have abstained from movement planning and simply waited for the 'Go' signal, presented in the form of a triangle. To ensure that participants nevertheless paid attention to the cues and, consequently, generated the respective visuomotor

expectations or predictions, we included occasional trials in which the virtual hand moved *contrary* to the color-cued expectation (25% of all trials, balanced across conditions). For instance, the virtual hand moved congruently although the cue had predicted it to move incongruently (but note that the cues indicating the to be executed 'own' movement type were always valid). Following these invalidly cued trials, and following the same number of validly cued trials in each condition, participants had to report whether the virtual hand behavior matched the predictions of the cue color (cf. Yon et al., 2018). For this, a probe question ("Prediction correct?") was presented centrally on the screen alongside two response options ("Yes"/"No"). Participants had to choose between those by pressing one of the response device buttons twice (2-Button Fiber Optic Response pad, Current Designs, placed in the participant's right hand, on their lap). If they did so within 1.5 s, feedback (i.e. "right" or "wrong") was shown for 0.75 s, otherwise they were visually instructed to respond faster the next time. A blank screen was shown for 2 s before the next trial. To increase motivation, participants received an additional monetary reward after the experiment, depending on the proportion of correct answers to the probe questions in the catch trials (1 € if at least 75% were answered correctly, 3 € for ≥85% and 5 € for ≥95%). We chose a relatively large number of catch trials (half of all trials in total) to maximize the task relevance of the cues during the preparation period; i.e., participants had to assume that the cue color could be probed by chance. Thus, the design for the execution period entailed *Visuomotor cue validity* (Valid, Invalid), as another nested factor (see below). This manipulation also allowed us to examine neuronal "surprise" responses to expectation violation; i.e., by comparing invalidly with validly cued catch trials.

Each trial lasted between 8.5 and 15 s. For each participant, we recorded 8 runs of 32 trials each (6.5-8.5 mins run length), resulting in 256 trials (64 per condition and preparation/execution period, respectively) and around 60 mins scanning time in total (depending on hand positioning and response times). Participants were free to rest as much as they wanted in between the individual scanning runs.

Behavioral data analysis

We tested for systematic differences in movement onset, duration, or amplitude between conditions; using the participants' hand movement data from the glove sensors (averaged over all fingers but the thumb; the movements had a similar kinematic profile, comparable across conditions, without averaging the contribution of the individual fingers, see Results). We segmented the continuous glove data to select the execution period time window of each trial, smoothed (Gaussian-weighted moving average filter, time window 10 frames i.e. 167 ms), and re-adjusted the values to a baseline defined by the mean hand position during the 500 ms before appearance of the virtual hand (i.e. the instructed start of execution). Movement onset was defined, for each trial, as the respectively last prior time point with zero velocity. Note that we could only approximate the actual movement onset from the recorded glove data; intrinsic delays due to the design of the glove (see above) should be taken into account when interpreting the onset values, but these delays were constant and hence irrelevant to our factorial analyses. The end of the movement was defined as the time point where hand velocity reached zero again. Movement duration and amplitude, respectively, were defined as the time difference and amplitude difference between movement onset and -end. Mean values for movement onset, duration and amplitude per condition were calculated for each subject. Based on these, a four-way repeated-measures ANOVA with the factors *Movement type* (Open, Close) *Movement type cueing* (Cued, Uncued), *Visuomotor mapping* (Congruent, Incongruent) and *Visuomotor cue validity* (Valid, Invalid) was used to test for differences between conditions. Reaction time in catch trials was defined as the difference between presentation of the question on screen and the time point at which participants confirmed their answer. All analyses were performed using MATLAB (MathWorks).

fMRI data preprocessing and analysis

The fMRI data were recorded with a 3T scanner (MAGNETOM Prisma; Siemens) equipped with a 32-channel head coil. T2*-weighted images were acquired using an echo-planar imaging sequence (GRAPPA acceleration factor = 3, multi-band factor = 6, TR = 869 ms, TE = 38 ms, flip angle = 58°, voxel size = 2.4 mm^3, matrix size = 88 x 88, 60 interleaved slices). Depending on the length of the individual session (see above), 3000-4000 functional image volumes were recorded per participant. Additionally,

for each participant we recorded a T1-weighted anatomical image (3D, voxel size = 0.9 x 0.9 x 0.9 mm^3, FOV = 272 x 272 mm, 240 slices, TR = 2.4 s, TE = 2.2 ms, flip angle = 8°) and two GRE field maps (64 layers, voxel size 2 x 2 x 2 mm^3; matrix size = 88 x 88; TR = 698 ms, TE$_1$ = 5.19 ms, TE$_2$ = 7.65 ms, flip angle 54°), after the second and sixth scanning run, respectively. All fMRI preprocessing steps and analyses were performed using MATLAB (MathWorks) and SPM12 (Wellcome Trust Centre for Neuroimaging, University College London; https://www.fil.ion.ucl.ac.uk/spm). The functional images were realigned and unwarped using voxel displacement maps calculated from the two field maps (the first map was used for runs 1 to 4, the second for runs 5 to 8). These images were then co-registered with the individual T1 image, spatially normalized to MNI space via previously estimated deformation fields, resliced to 2 mm^3 voxels, and smoothed with an 8 mm full with at half maximum Gaussian kernel. The T1 images were also normalized (but not smoothed) and averaged to create a group averaged structural image.

In our fMRI analysis, we adopted a mass univariate approach; i.e., we tested for voxel-wise differences in the BOLD signal between our experimental conditions. Implicit in this is the assumption that a relatively higher or lower BOLD signal in one condition compared to another (i.e., a differential contrast) indicates a regionally in- or decreased metabolic demand in this condition, respectively. We interpret such differences as evincing an increased engagement of brain regions in experimentally manipulated processes related to motor planning and visuomotor prediction, as identified per our factorial between-condition comparisons (see Fernandez-Ruiz et al., 2007; Lindner et al., 2010; Bernier et al., 2012; Gertz & Fiehler, 2015; or Pilacinski et al., 2018, for similar approaches). For each participant, a general linear model (GLM, 128 s high-pass filter) was fitted to the preprocessed data with stick (0 s duration) regressors modeling each combination of the two factors *Movement type cueing* (Cued, Uncued) and *Visuomotor mapping* (Congruent, Incongruent) in the preparation and movement period, respectively. The resulting four regressors were thereby aligned with the onsets of the predictive cues; or with the onsets of the movement execution period including the virtual hand presentation. To model any potential 'surprise' effects, the regressors of the execution period were

each supplemented by a mean-centered parametric modulator encoding trials in which the virtual hand moved unlike cued with +1, and trials in which it moved like cued with -1. Thereby, we applied the modulation to the catch trials only; i.e., ensuring the surprise effect was calculated on the same amount of trials (all with subsequent button presses) in invalidly and validly cued movements.

The six realignment parameters were added as noise regressors to each run, alongside a session constant. Furthermore, we added three regressors of no interest modelling the differences in movement amplitude, duration, and onset (i.e., the reaction time it took participants to initiate a movement after onset of the execution period) during the execution period. Each kinematic regressor modelled trial-to-trial variations (in amplitude, duration, and onset) across the entire scanning session; i.e., it also modelled differences between conditions. Two additional regressors of no interest were included: one modeling the initial hand positioning (a 0s duration stick regressor with an additional orthogonalized parametric regressor encoding the duration from trial start until the correct hand position was reached and the fixation dot presented), and one modeling the response periods after the occasional catch trials (a 0s duration stick regressor with an additional orthogonalized parametric regressor encoding the respective variations in reaction time until the confirmatory key press). On the first (single participant) level, T-contrast images were created for each regressor vs baseline, across all 8 runs per participant (i.e., over all 64 trials per preparation or execution period per condition. These first-level contrast images were entered into group-level GLMs using a flexible factorial design, in which the group (second-level) T- and F-contrasts were calculated. We thereby used 2x2 factorial designs with the aforementioned factors to evaluate the main effects of cued vs uncued movements and congruent vs incongruent visuomotor mapping, as well as their potential interaction, during the preparation and execution periods, respectively.

We furthermore tested for the *consistency* of the cueing main effects; i.e., whether a given main effect of one experimental factor (movement type cueing or visuomotor mapping) was significant at each level of the respective other factor. To do this, we calculated 'null' conjunction contrasts; a statistical test whether a BOLD signal difference in a given voxel is significant in each of multiple tested contrasts.

Thus, we calculated a conjunction across the contrasts CC>UC and CI>UI (testing for consistent significance of the main effect movement type cued > uncued across both levels of the factor visuomotor mapping); and another one across the contrasts CI>CC and UI>UC (testing for consistent significance of the main effect incongruent > congruent visuomotor mapping across both levels of the factor movement type cueing). These conjunction contrasts were evaluated for statistical significance using a voxel-wise threshold of p_{FWE}<0.05, within the search region defined by the significant voxels obtained from the respective main effect.

Furthermore, we tested for any influence of the (cued) preparatory processes on activity in the respectively identified areas during subsequent movement execution. As we did not have any specific prediction about the directionality of response differences during movement execution, we used non-directional F-contrasts in this case. To assess potential surprise effects during execution associated with invalidly > validly cued movements, we calculated an analogous 2x2 factorial group level design on the parametric 'surprise' modulators of each condition (i.e., on T contrast images weighting invalidly cued catch trials in each condition with +1 and validly cued ones with -1, see above).

Activations in the whole brain obtained from the group-level contrasts were assessed for statistical significance using a voxel-wise threshold of p<0.05, family-wise error corrected (p_{FWE}<0.05). In three cases, we had specific regions of interest defined a priori by orthogonal contrasts, and restricted the search space accordingly, thus applying a small volume correction under the same voxel-wise threshold of p_{FWE}<0.05 for those results: Firstly, we looked for an interaction of motor and visuomotor planning in brain regions showing a response to both individual effects. Therefore, we defined the search space for these interaction effects based on all (whole-brain) significant voxels obtained from the null conjunction of the respective main effects (Fig. 5). For completeness, we also tested for, and show the corresponding whole-brain results. Secondly, based on the previously reported overlap between action planning and execution (Lotze et al., 1999; Ames et al., 2014), we expected that brain regions involved in motor or visuomotor planning would also show condition specific effects during movement execution. Therefore, for the analysis of responses during the execution period, we restricted the

search space for effects to all significant voxels obtained from each respective analogous contrast during the preparation period. Thirdly, we hypothesized that surprise effects might emerge in brain regions implementing the forward model, i.e., issuing the respective forward predictions. In other words, in those regions, we expected a relative increase in BOLD in response to a (sensory) prediction error elicited by the unpredicted visuomotor mapping. As the generation of those predictions should be highlighted by the main effect of preparation for incongruent > congruent movements (see Introduction), we used those significant activations as a mask image to look for corresponding surprise effects; but also calculated an additional whole-brain analysis.

The significant activation differences resulting from the group level contrasts are presented as renders on SPM12's template brains, or superimposed onto the group averaged normalized structural image. The displayed significant activations contain only voxels surviving the chosen threshold of $p_{FWE}<0.05$; any statistical maps displayed at $p<0.001$, uncorrected, were included solely to demonstrate that the obtained significant activations were very focal; i.e., did not spread into other, nonrelated areas.

The contrast estimate plots were created by extracting and averaging the first-level contrast estimates for each condition from the peak voxel of each anatomically distinct cluster, using the rfxplot toolbox (Gläscher, 2009). Note that our trial structure was designed to maximize the number of repetitions per condition, for optimal sensitivity to the factorial (between-condition) effects during the planning or execution period, respectively; i.e., relative hemodynamic activity differences. Thus, there were only very brief periods of no stimulation or movement within and between trials throughout the entire scanning session (trials were separated by 2 s, see above), which in other designs may serve as a BOLD signal 'baseline'. As most (high-level) motor areas could be expected to be involved in planning and execution (more or less continually), the *absolute* values of the contrast estimates should be interpreted with some caution; i.e., positive or negative values need not necessarily indicate (de)activations vs a rest period baseline. Thus, our contrast estimates should be interpreted in terms of a relative contribution of each regressor to the continuous hemodynamic activity in the respective brain region; the corresponding bar plots should be taken mainly as a visualization of the *relative,*

between-condition BOLD signal differences in regions identified by our group-level contrasts. For anatomical reference, we used the SPM Anatomy toolbox (Eickhoff et al., 2005).

Eye tracking control experiment

Our non-spatial (i.e., without spatial movement targets) movement task was intended to render eye movements useless and, thus, unlikely. As we did not measure gaze on-line inside the scanner, we conducted an eye-tracking control experiment post-hoc, with the same design. The aim of this experiment was to determine whether participants could, in principle in this task, maintain fixation onto the central cues/dot; and whether gaze behavior was comparable across experimental conditions. A sub-sample of our fMRI study participants (*N*=6) completed the same task as in the scanner (with 4 instead of 8 runs in total), with stimuli presented at a size corresponding to those seen in the scanner environment. Participants were seated comfortably in a constantly lit room in front of a screen (at a distance of 93 cm) with their head stabilized via a chin rest, while we measured gaze position with an Eyetracker (EyeLink 1000 Plus, Desktop Mount, SR Research). We segmented the resulting eye position data to select 2500 ms time windows for both the preparation and execution periods of each trial, with phases of data loss (e.g. due to eye blinks) corrected via linear interpolation. Gaze position was defined relative to the position of the central fixation dot; i.e., the measured fixation position in the horizontal and vertical planes was combined to a measure of Euclidean distance from the fixation point, and converted to degrees of visual angle. To evaluate the quality of fixation, we tested for differences in mean gaze position and in the variability of fixation (i.e., the standard deviation); and we tested for this separately for the two possible movements. Thus, we performed a four-way repeated-measures ANOVA with the factors *Movement type* (Open, Close), *Movement type cueing* (Cued, Uncued), *Visuomotor mapping* (Congruent, Incongruent), and *Visuomotor cue validity* (Valid, Invalid) on the participants' condition averages of both fixation position and variability, respectively.

FMRI control experiment

To follow up on the finding that the main effects of cueing (and to a lesser extent also execution) were more pronounced in the left than in the right PPC (i.e., ipsilateral to the moving hand, see Figs. 3-6), we scanned 6 of our original participants again in a separate session; using an identical experimental setup and design (with 5 instead of 8 runs in total) but with right-hand instead of left-hand movements (and a right instead of left virtual hand model). These data were preprocessed and analyzed as described for the main experiment. As the small sample size limited the power of the random-effects group analysis, we evaluated and displayed the resulting statistical parametric maps at a threshold of $p < 0.001$, uncorrected.

Results

All participants showed very good task performance; i.e., all performing >97% correct movements (on average, only one incorrect movement per recording session) and correctly identifying >93% of the catch trials (on average, >98% of responses were correct). The participants' hand movements were consistent across conditions (Fig. 2A). On average, movement execution was detectable from the glove data 143±149 ms after the onset of the execution period, with a mean movement duration of 1145±352 ms. Movement amplitude, in terms of average flexion values relative to the starting position, averaged to -0.26±0.08 (hand opening) and 0.48±0.09 (hand closing), respectively. There was no significant effect of visuomotor mapping (congruent vs incongruent) on movement onset, duration, or amplitude of the hand movements. Likewise, motor planning (cued vs uncued) had no significant effect on movement amplitude or duration. However, movements in cued trials were on average initiated 49 ms earlier than those in uncued trials ($F_{(2,29)}$=7.34, p=0.011, η_p^2=0.20). Movements under an invalidly cued visuomotor mapping trials were, on average, initiated 72 ms later ($F_{(2,29)}$=42.09, p=0.0000004, η_p^2=0.59) and 149 ms shorter ($F_{(2,29)}$=56.67, p=0.00000003, η_p^2=0.66) than movements in validly cued trials; movement amplitude was not significantly affected by cue validity. The above significant differences were replicated when analyzing the data of the four fingers individually (all Fs>6.5, all ps<0.05, all η_p^2>0.18). These differences could indicate early surprise responses to the expectation violation, leading to a delay in movement initiation and, possibly, shorter movements due to

perceptual processing of the surprising mapping. There were no significant interaction effects, except a three-way interaction on movement duration (visuomotor mapping x motor planning x cue validity, $F_{(2,29)}$=6.82, *p*=0.014, η_p^2=0.19), which was difficult to interpret (see Fig. 2A). Importantly, kinematic differences between trials and conditions – including those above – were modeled as covariates of no interest in the fMRI first-level GLMs (see Materials and Methods). The average reaction time in catch trials was 0.920±0.246 s. There were no significant effects of visuomotor cue validity, visuomotor mapping, or motor planning on reaction times.

A supplementary eye-tracking experiment (*N*=6) showed that participants were, in principle, able to maintain very stable fixation onto the central cues throughout the preparation and execution periods (see Fig. 2B). Moreover, none of our experimental factors (visuomotor mapping, motor planning, or cue validity) significantly affected average fixation or fixation variability during the preparation period. Likewise, visuomotor mapping and motor planning had no significant effect on average fixations and fixation variability during movement execution. The execution of movements with incongruent > congruent visuomotor mappings was associated with larger fixation variability ($F_{(1,5)}$=7.25, *p*=0.043, η_p^2=0.59), but did not significantly affect average fixation. However, fixation variability was in itself very small: the average standard deviation of any condition was <0.17 degrees visual angle, whereas the hand movements spanned 10.6 degrees. The planned or executed hand movement type (open vs close) had no significant effects on average gaze position or variability during preparation and execution, respectively. There were no significant interaction effects between any two factors either (there were, for average fixation but not variability, a significant four-way interaction during planning, $F_{(1,5)}$=11.89, *p*=0.018, η_p^2=0.70 and a significant three-way interaction, mapping x cue validity x movement type, during execution, $F_{(1,5)}$=6.82, *p*=0.048, η_p^2=0.58 see Fig. 2B). In sum, the eye tracking results suggested that the results obtained from our fMRI main effect and interaction contrasts (Figs. 3-7) were likely not biased by systematic differences in gaze behavior.

Figure 2. Comparability of hand movements and fixation across conditions. (A) Average hand movements during the execution period (Time = 0 marks the instructed start and appearance of the virtual hand model). The curves show the group-averaged raw (unsmoothed) flexion values, with associated standard deviation indicated by shaded areas; positive values indicating closing and negative values indicating opening, relative to the neutral starting position (0 on the y-axis). We show the closing (positive flexion values) and opening (negative flexion values) movement separately per condition and, furthermore, for movements in which the cued visuomotor mapping was valid vs invalid. All fMRI contrasts were pooled over both movement types per condition, balancing out any kinetic or kinematic differences. CC = cued/congruent, CI = cued/incongruent, UC = uncued/congruent, UI = uncued/incongruent. **(B)** A supplementary eye tracking experiment (*N*=6 of the original study participants, without scanning) showed overall very stable gaze behavior during the preparation and execution periods. The right plots show, for open (top rows) and close (bottom rows) separately, a small section of the screen (1° x 1° visual angle, marked by the white squares in the screenshots) centered on the fixation dot (here in grey). The crosses mark the respective average fixation, with horizontal and vertical standard deviations, during the preparation period (PREP, dotted bars) and the execution period (MOVE, solid bars).

In sum, the participants' behavior suggested high task engagement (as indicated by almost perfect execution of cued movements) and comparable hand movements (and fixation) across conditions. The fact that movements in cued trials were initiated faster than in uncued trials likely indicates a reaction time benefit of prior motor planning, in line with previous findings (Rosenbaum, 1980; cf. Ames et al., 2014). Moreover, the almost perfect responses to catch trials, in which the validity of the color-cued and the actual visuomotor mapping was probed, indicated that participants paid attention to the

visuomotor cues. Thus, we were confident that participants generated the desired (cued) motor plans and visuomotor predictions during scanning.

In our fMRI analysis, our primary aim was to identify brain areas involved in both motor planning and visuomotor prediction; i.e., we looked for corresponding BOLD signal differences during the delay period. First, we contrasted the delay period of trials in which a specific movement type could be prepared – based on the cue shape – with trials in which this was not possible (i.e., main effect of movement type cued > uncued). As expected, this contrast revealed significantly (p_{FWE}<0.05) stronger BOLD predominantly in the bilateral frontoparietal cortices; including the bilateral aIPS (and, in the left hemisphere, larger portions of the IPS), the bilateral SPL, the SMA, medial parts of the bilateral PMd, the left ventral premotor cortex (PMv); and the bilateral lateral occipital cortices (LOC, including area V5). See Table 1 and Figure 3A for details. A conjunction analysis revealed that, in almost all of these regions, the main effect of cued > uncued movements was consistent across congruent and incongruent visuomotor mappings (i.e., levels of the second experimental factor; p_{FWE}<0.05, see Table 1). In other words, this suggested a consistently elevated BOLD signal by motor planning during preparation for congruent and incongruent visuomotor mappings. The converse contrast (i.e., delay period of uncued > cued movement types) revealed smaller clusters of significant voxels mainly in the visual and somatosensory cortices (see Table 1 and Fig. 3B).

Then, we tested for increased responses during the delay period of trials in which an incongruent > congruent mapping of the virtual hand movements relative to the real hand movements was cued; assuming that here, predictions of the forward model would need updating. There was a significant main effect of cues predicting an incongruent > congruent mapping in the cerebellum (centered on Lobule VI in each hemisphere), premotor (the bilateral PMd, and the left preSMA and PMv) and posterior parietal (the left aIPS and the bilateral SPL) areas (see Figure 4 and Table 1 for further effects). However, a conjunction analysis revealed that this effect was consistent across cued and uncued movement types in only some of those regions (p_{FWE}<0.05, see Fig. 4): Most notably, the strongest consistent effects across the CI > CC and the UI > UC contrasts were located in the bilateral Lobules VI

of the cerebellum (see slice overlays in Fig. 4). Only few other voxels in the PMd, the SMA, and the left precuneus showed consistent effects. No significant activation differences were observed for the converse contrast, cued congruent > incongruent visuomotor mapping.

Figure 3. Brain activation differences related to motor planning during the delay period. (A) The renders show significant (p_{FWE}<0.05) voxels obtained from contrasting the delay period of trials in which the to-be-executed movement type was cued, with trials in which this was left ambiguous (cued > uncued). Inlay: Significant (p_{FWE}<0.05) voxels obtained from a conjunction testing for consistency of the effect cued > uncued across congruent and incongruent mappings. **(B)** The converse contrast (uncued > cued) yielded significant activation differences in visual and somatosensory cortices. The bar plots show the contrast estimates at the peak voxel of each respective cluster, with associated standard errors of the mean. These estimates indicate each regressor's relative contribution to the overall BOLD signal, as our design did not include a proper rest period baseline, see Materials and Methods. CC = cued/congruent, CI = cued/incongruent, UC = uncued/congruent, UI = uncued/incongruent. See Table 1 for details.

Figure 4. Brain activation differences related to visuomotor prediction during the delay period. The renders show significant (p_{FWE}<0.05) voxels obtained from contrasting the delay period of trials in which the cue color predicted incongruent, compared with congruent visuomotor mapping (incongruent > congruent). Inlay: Significant (p_{FWE}<0.05) voxels obtained from a conjunction testing for consistency of the effect incongruent > congruent visuomotor mapping across cued and uncued movements. The bar plots show the contrast estimates at the peak voxel of each respective cluster, with associated standard errors of the mean. These estimates indicate each regressor's relative contribution to the overall BOLD signal, as our design did not include a proper rest period baseline, see Materials and Methods. CC = cued/congruent, CI = cued/incongruent, UC = uncued/congruent, UI = uncued/incongruent. CR = cerebellum. See Table 1 for details.

Table 1. Brain areas showing significant main effects during the delay.

Movement type cued > uncued (CC+CI)-(UC+UI)		MNI			Peak	
Anatomical location	*Voxels*	*x*	*y*	*z*	*T*	*p_{FWE}*
L. Lateral occipital cortex (V5)*	390	-40	-82	-4	8.72	<0.001
R. Lateral occipital cortex (V5)*	183	46	-78	-8	7.85	<0.001
L. Superior frontal gyrus (SMA & PMd)*	475	-10	-4	64	7.06	<0.001
L. Intraparietal sulcus and superior parietal lobe*	695	-38	-38	46	6.68	<0.001
L. Precentral/inferior frontal gyrus (PMv)*	60	-60	10	26	6.26	<0.001
L. Superior parietal lobe (BA 7A)*	68	20	-68	58	6.08	0.001
R. Fusiform gyrus*	16	42	-58	12	5.85	0.002
R. Anterior intraparietal sulcus	46	34	-36	42	5.72	0.004
Movement type uncued > cued (UC+UI)-(CC+CI)		**MNI**			**Peak**	
Anatomical location	*Voxels*	*x*	*y*	*z*	*T*	*p_{FWE}*
L. Pre-/postcentral gyrus (S1/M1)*	115	-28	-24	48	6.18	<0.001

	Voxels	x	y	z	T	p
L. Parietal operculum (OP)	19	-38	-24	22	5.78	0.003
L. Paracentral lobule*	17	-8	-20	50	5.69	0.004
R. Parietal operculum (OP)	16	46	-30	20	5.55	0.008
L. Medial temporal lobe*	8	-54	-10	-22	5.49	0.009
L. Occipital pole (V3)*	7	-18	-100	0	5.34	0.016
L. Parahippocampal gyrus*	1	-32	-40	-8	5.15	0.032
R. Anterior cingulate gyrus*	1	6	32	-8	5.07	0.041
R. Postcentral gyrus (S1)	1	22	-36	64	5.04	0.047
R. Postcentral gyrus (S1)	1	20	-38	62	5.03	0.048

Mapping incongruent > congruent (CI+UI)-(CC+UC)		MNI			Peak	
Anatomical location	Voxels	x	y	z	T	p_{FWE}
R. Middle frontal gyrus (PMd)*	688	32	-2	56	7.94	<0.001
R. Cerebellum (Lobule VI/VII) *	781	38	-64	-32	7.86	<0.001
L. Superior/middle frontal gyrus (PMd/preSMA)*	1852	-10	12	54	7.22	<0.001
L. Cerebellum (Lobule VI)*	263	-26	-58	-34	6.64	<0.001
L. Superior parietal lobe (BA 7A)*	200	-8	-60	48	6.52	<0.001
L. Anterior intraparietal sulcus	486	-46	-42	52	6.07	0.001
R. Anterior intraparietal sulcus	243	36	-50	52	5.96	0.002
R. Cerebellum (Lobule I)	56	2	-56	-12	5.89	0.002
R. Cerebellum (Lobule VII/VIII)*	74	32	-64	-48	5.86	0.002
L. Superior temporal sulcus	13	-46	-50	8	5.76	0.003
R. Paracingulate gyrus	34	8	22	36	5.70	0.004
Cerebellum (Vermis)	96	0	-72	-18	5.57	0.007
L. Anterior insular cortex	51	-28	26	2	5.54	0.008
R. Superior parietal lobe (BA 7A)	62	-16	-66	54	5.47	0.010
R. Caudate	23	16	2	16	5.44	0.011
L. Middle frontal gyrus	5	-38	30	28	5.28	0.020
R. Anterior insular cortex	5	34	20	10	5.26	0.020
R. Intraparietal sulcus	4	30	-62	38	5.21	0.025
L. Supramarginal gyrus	2	-56	-44	26	5.18	0.029
L. Paracingulate gyrus	3	-8	20	36	5.17	0.029
R. Superior frontal gyrus	2	16	8	56	5.07	0.041
R. Intraparietal sulcus	1	28	-62	42	5.07	0.042
L. Cerebellum (Crus I)	1	-46	-58	-34	5.03	0.047
L. Middle frontal gyrus	2	-44	24	38	5.02	0.048

CC = cued/congruent, CI = cued/incongruent, UC = uncued/congruent, UI = uncued/incongruent.
* marks clusters with significant voxels in a conjunction contrast across both levels of the respective other factor.

Next, we specifically tested for brain areas that would jointly encode both preparatory processes in

our design; i.e., those related to motor planning and visuomotor mappings. A null conjunction contrast

across the respective main effects of (cued > uncued movement type and incongruent > congruent visuomotor mapping, Fig. 3A and Fig. 4) revealed that identical voxels within the bilateral aIPS and SPL, and the left medial PMd/SMA were significantly activated when participants anticipated a specific movement type or an incongruent visuomotor mapping, see Figure 5A and Table 2. Then, we evaluated our hypothesis that motor planning and visuomotor prediction would interact; i.e., we tested for relatively stronger movement planning effects (cued > uncued movements) under expected visuomotor incongruence, compared with congruence. Only the left aIPS showed a significant corresponding interaction effect: a relatively stronger increase in fMRI activity for delay periods in which the cue predicted a specific movement type with incongruent > congruent visuomotor mappings (CI>CC), compared with when the movement type was left ambiguous (UI>UC; see Figure 5B and Table 2). In other words, in the left aIPS, while the main effect of cueing itself was consistent, the main effect of preparing for incongruent visuomotor mappings was significantly stronger in the cued movement conditions. No other significant effects were found in the whole brain; a similar interaction effect in the left PMd did not reach significance (x = -26, y = -4, z = 58, T = 3.71, *p*<0.001, uncorrected).

Table 2. Significant activation differences obtained from the null conjunction of main effects and the interaction effect.

Null conjunction (CC+CI)-(UC+UI) ∩ (CI+UI)-(CC+UC)		*MNI*			*Peak*	
Anatomical location	*Voxels*	*x*	*y*	*z*	*T*	*pFWE*
L. Superior frontal gyrus (PMd/SMA)	82	-12	0	64	6.17	0.001
L. Anterior intraparietal sulcus	197	-46	-42	52	6.01	0.001
L. Superior parietal lobule	29	-8	-60	54	5.86	0.002
R. Superior parietal lobule	11	18	-66	56	5.43	0.012
L. Superior frontal gyrus	1	-6	4	50	5.33	0.034
R. Anterior intraparietal sulcus	3	40	-36	44	5.13	0.034
R. Anterior intraparietal sulcus	1	36	-40	42	5.09	0.039
L. Superior frontal gyrus	2	-4	2	54	5.08	0.041
Interaction (CI-CC)-(UI-UC)		*MNI*			*Peak*	
Anatomical location	*Voxels*	*x*	*y*	*z*	*T*	*pFWE*
L. Inferior parietal lobule / intraparietal sulcus	8	-46	-46	-56	3.70	0.015
L. Anterior intraparietal sulcus	5	-36	-42	40	3.47	0.040

CC = cued/congruent, CI = cued/incongruent, UC = uncued/congruent, UI = uncued/incongruent.

Figure 5. Overlap and interaction of motor planning and visuomotor preparation. (A) Renders showing voxels with significantly (p_{FWE}<0.05) stronger BOLD during the preparation of cued > uncued movements *and* during the preparation for incongruent > congruent visuomotor mappings; i.e., a null conjunction of the main effects shown in Figs. 3A and 4. **(B)** Voxels in the left IPS/IPL showed a significant (p_{FWE}<0.05) interaction between those effects, within the areas identified by the null conjunction. The bar plots show the contrast estimates with associated standard errors of the mean. These estimates indicate each regressor's relative contribution to the overall BOLD signal, as our design did not include a proper rest period baseline, see Materials and Methods. The inlay shows the corresponding whole-brain render, thresholded at *p*<0.001, uncorrected. CC = cued/congruent, CI = cued/incongruent, UC = uncued/congruent, UI = uncued/incongruent. See Table 2 for details.

Subsequently, we looked at condition-specific activation differences during the movement execution period; which would indicate specific effects of the cues presented during the prior delay period. The left aIPS and the left PMd showed significant BOLD signal differences depending on cueing (Fig. 6A and Table 3). Specifically, they showed increased activation levels in trials where the movement type was left uncued until the 'Go' signal, compared with trials where it had been cued in advance. A whole-brain analysis revealed one further significant voxel in the ventromedial prefrontal cortex. Movement execution depended on cued visuomotor mapping (congruent vs incongruent virtual hand movement)

in the left PMd, the left aIPS, and the left anterior insula (Table 3). Activations of those regions were relatively increased during movements following cued incongruent visuomotor mapping (Fig. 6B). In a whole-brain analysis at more liberal thresholds ($p<0.001$, uncorrected, see Fig. 6B inlay), we found further activation differences in the right PMd, anterior insuls, and IPS, among other areas. The main effects overlapped in the left aIPS (conjunction of main effects: x = -38, y = -50, z = 50, peak F = 14.57, Fig. 6C), but this effect did not reach corrected significance; and there were no significant interaction effects.

Table 3. Brain areas showing significant activation differences during movement execution related to prior (cued) movement planning and visuomotor mapping predictions.

Execution after Movement type cued vs uncued		MNI			Peak	
Anatomical location	*Voxels*	*x*	*y*	*z*	*F*	*p$_{FWE}$*
L. Anterior intraparietal sulcus	12	-42	-44	50	18.88	0.020
L. Superior frontal gyrus (PMd)	7	-20	4	70	18.69	0.021
Ventromedial prefrontal cortex	3	0	28	-6	27.74	0.047*
Execution after Mapping cued congruent vs incongruent		MNI			Peak	
Anatomical location	*Voxels*	*x*	*y*	*z*	*F*	*p$_{FWE}$*
L. Precentral / middle frontal gyrus (PMd)	14	-26	-6	50	21.59	0.017
L. Anterior intraparietal sulcus	7	-36	-50	42	20.85	0.022
L. Anterior insular cortex	5	-30	24	0	20.51	0.025

* whole-brain analysis.

Figure 6. Brain activation differences during movement execution related to prior movement planning and visuomotor predictions. (A) Renders of voxels showing significant ($p_{FWE}<0.05$) BOLD signal differences during movement execution in trials where the movement type had been cued in advance versus when it was left ambiguous (uncued). **(B)** Renders of voxels showing significant ($p_{FWE}<0.05$) BOLD signal differences during movement execution in trials where congruent vs incongruent visuomotor mapping had been cued. **(C)** Voxels showing an overlap of both main effects (conjunction contrast, $p<0.001$, uncorrected). The bar plots show the contrast estimates indicating each condition's relative effect (see Materials and Methods) at the peak voxel of each respective cluster, with associated standard errors of the mean. CC = cued/congruent, CI = cued/incongruent, UC = uncued/congruent, UI = uncued/incongruent. aIC = anterior insular cortex. See Table 3 for details.

We also tested for potential 'expectation violation' effects observable during movement execution in trials where the virtual hand behaved contrary to the cued expectations (i.e., the cued visuomotor mapping was invalidated). We tested for such responses in brain regions showing increased activity during the preparation for incongruent > congruent mappings, as we assumed this contrast identified brain areas generating visuomotor predictions (see Introduction). Indeed, we found significant

(p_{FWE}<0.05) surprise effects in the (pre)SMA, the left PMd and PMv, the bilateral IPL/IPS, the left

anterior insula, and the right anterior cerebellum (see Fig. 7A). An additional whole-brain analysis

revealed further surprise effects in the dorsomedial prefrontal cortex, the bilateral inferior frontal gyri,

the bilateral angular gyri, the right anterior insula, the right middle temporal and fusiform gyri, the

putamen and thalamus, and in the left primary sensory and motor cortex (p_{FWE}<0.05, Fig. 7B). There

were no significant differences in surprise-related activations between conditions (i.e., no voxels

showed a significant main or interaction effect). The converse contrast (expected > unexpected

visuomotor mapping) yielded no significant activations either.

Figure 7. Effects of unexpected > expected visuomotor mappings ('surprise'). A: Significant (p_{FWE}<0.05) surprise effects within brain areas showing effects of visuomotor prediction (cf. Fig. 4). **B:** Render of significant (p_{FWE}<0.05) activations in a whole-brain analysis.

Finally, following the observation that the movement cueing effects in our left-hand experiment were

strongly left-lateralized in the IPS, six of our original study's participants were invited to a separate

scanning session, in which they repeated a right-hand version of the task. As in the main experiment,

the preparation of cued > uncued movements was associated with stronger BOLD signal in bilateral

frontoparietal areas (including PMd, PMv, SMA, SPL and IPS) and the V5 (Fig. 8A; cf. Fig. 3A). Again,

within the left compared with the right IPS, the effects were more extensive (spanning to anterior

portions of the IPS only in the left hemisphere) and stronger (all peak T-values were higher in the left

> right PPC). The corresponding overlap of main effect also revealed a left-hemispheric dominance (see

conjunction contrast in Fig. 8A). The interaction effect did not reach significance but was present,

among other regions, in the left but not right IPL (x = -48, y = -58, z = 50, T = 3.13, *p*<0.005). As in the main experiment, the preparation of movements with an expected incongruent > congruent visuomotor mapping was associated with stronger BOLD signal in the bilateral PMd and SPL, the SMA, and in the cerebellum (Fig. 8A; cf. Fig. 4). Contrasting the execution of uncued > cued movements likewise revealed activation differences corresponding to those in the main experiment; and again stronger left-hemispheric effects in the PPC (Fig. 8B; cf. Fig. 6). Surprise responses were also observable in similar brain regions, including notable effects in the inferior and dorsomedial frontal cortex, temporoparietal areas, and in the cerebellum (Fig. 8C; cf. Fig. 7). In sum, the pattern of results obtained during right-hand planning and movement replicated our main results; together, our results suggest that cueing effects, particularly planning, predominantly engaged the left IPS for either hand side.

Figure 8. Supplementary right-hand fMRI experiment. N=6 of the original study participants repeated the main experiment with right hand movements. The renders show group-level activation differences (*p*<0.001, uncorrected) obtained from the respective contrasts. The conjunction contrast was significant at *p*<0.001, but is displayed at *p*<0.005 to better show the spatial extent of left > right-hemispheric PPC effects. **(A)** Preparation period (cf. Figs. 3A and 4); **(B)** execution period (cf. Fig. 6); **(C)** surprise effect (cf. Fig. 7).

Discussion

In our non-spatial delayed movement task with altered visuomotor mapping, delay period activity in the premotor cortex (PMd, PMv, and the SMA) and the PPC (IPS and SPL) was increased if one of two alternative movements was cued ahead of execution. This supports the general association of those regions with motor planning (see Introduction) and aligns with frontoparietal BOLD increases observed during cued reach planning (Fernandez-Ruiz et al., 2007; Lindner et al., 2010; Gertz & Fiehler, 2015; Pilacinski et al., 2018). The PMd and aIPS showed the inverse pattern during execution (relatively higher BOLD when executing uncued movements), as similarly observed in uncued vs cued effector selection tasks (Bernier et al., 2012). We propose this indicates a fast planning process that still needed to be established (cf. Ames et al., 2014; Pilacinski et al., 2018). Together, we interpret the frontoparietal BOLD pattern as indicating the establishment of a movement plan during the delay of cued trials, rather than parallel planning of both possible movements during the delay of uncued trials. Some recent studies suggest the motor system plans one action at a time (Dekleva et al., 2018), whereas other results suggest parallel planning (e.g. Cisek & Kalaska, 2002, 2005).Task design, including the kind of decision and movement used, may play a role in these differences; for instance, "nonspatial motor decisions" – like in our task – may be processed more serially than others (Cui & Andersen, 2011).

Our second key finding was the relatively higher cerebellar activity during the preparation for incongruent > congruent visuomotor mappings. Note that the incongruent virtual hand movements were predictable in principle from (planned) motor signals, because they reflected inverted sensor values from the data glove and were therefore also controlled on-line during execution. Thus, cerebellar delay activity likely indicated an anticipatory update of forward predictions of visual movement feedback. That this effect was significant regardless of whether a specific movement was planned or not suggests a general update of the effector's forward model to a nonstandard (inverted) visuomotor mapping, not only of the specific predictions associated with the movement selected for execution.

It has long been proposed that the cerebellum implements forward models (Miall et al., 1993; Wolpert et al., 1998). Lobule VI, specifically, responds to unpredicted sensory movement feedback (Blakemore et al., 1998, 2001; Kilteni et al., 2023), including visuomotor incongruencies (Leube et al., 2003; Limanowski et al., 2017; Van Kemenade et al., 2019). During sensorimotor adaptation, cerebellar activity may indicate trial-to-trial updates of forward models by sensory prediction errors (Shadmehr & Krakauer 2008; Izawa et al., 2012; see Tseng et al. 2007; Synofzik et al., 2008; Donchin et al. 2012; Tzvi et al. 2022 for visuomotor adaptation). Our results suggest that cerebellar predictions – informing state estimates about visuomotor mapping – can be updated ahead of, i.e., without movement execution. This supports the hypothesized generation of forward sensory predictions during action planning (see Introduction). The other main regions showing increased activity during the preparation for incongruence were the PMd and SMA. The PMd implements adjustments to "nonstandard sensorimotor mapping" (Wise et al., 1996); the dorsomedial frontal cortex is involved in processing conflicts during action selection (Taylor et al., 2007; Duque et al., 2013), error learning (Ulllsperger et al., 2014), and visuomotor adaptation (Grafton et al., 2008; Nahab et al., 2011; Limanowski et al., 2017; Ohata et al., 2020). Both premotor areas could therefore have contributed to forward modeling, or more generally anticipated the upcoming conflict (here: a nonstandard visuomotor mapping).

Thirdly, the bilateral PPC and the left PMd showed overlapping main effects of motor planning and visuomotor prediction, suggesting they contributed to more than one kind of predictive computation (cf. Scott, 2012). Crucially, the left aIPS showed a significant interaction, characterized by relatively increased BOLD during the planning of a specific movement under an expected nonstandard visuomotor mapping. Following our interpretation of the main effects, we propose this indicates the integration of the updated forward predictions of nonstandard visual body movement feedback with the state estimate used for movement planning, when a specific movement was pre-cued. The fact that this interaction was confined to the anterior part of the IPS fits with this region's proposed role of a specifically "body-related" state estimator (Medendorp & Heed, 2019; Kuang et al., 2015). It also aligns with previous demonstrations of the aIPS's key role in visuomotor control and upper limb state

estimation (Desmurget et al., 1999; Brass et al., 2001; Ehrsson et al., 2004; Grefkes & Fink, 2005; de Lange et al., 2006; Zimmermann et al., 2012; Limanowski & Blankenburg, 2016, 2017; Limanowski, 2022). It also supports the notion of the PPC as a high-level motor control region (Andersen and Buneo 2002; Blakemore & Sirigu, 2003).

Our results highlighted specifically the left aIPS (Figs. 3 and 7). This speaks to previous findings: The PPC represents actions from contra- and ipsilateral hands (Gallivan et al., 2013; Turella et al., 2020); and the left (but not right) PPC has been found activated during the selection, preparation, and execution of hand actions irrespective of hand laterality (Schluter et al., 2001; Rushworth et al., 2001; Grafton et al., 2002; Hardwick et al., 2013). The left PPC has also been identified as playing a key role in motor imagery, visuomotor adaptation, and the visuoproprioceptive integration it entails, for the left and right hand (Bonda et al., 1995; Servos et a., 2002; de Lange et al., 2006; Mutha et al., 2011; Hétu et al., 2013; Limanowski & Blankenburg, 2017). Notably, left PPC lesions have been associated with ideamotor apraxia (Haaland et al., 2000); fMRI results correspondingly suggest a "left-lateralized praxis network", centered on the PPC, which plans actions with either hand (Bohlhalter et al., 2009). Specifically, intransitive actions (i.e., without object interaction) may be more strongly left-lateralized than transitive ones (Króliczak & Frey, 2009). Our results, obtained with such actions, fit very well into this view; moreover, they show that the left PPC is essential for integrating visual movement predictions with these motor plans.

Two regions showing a main effect of planning but not visuomotor incongruence were the PMv and the V5. The PMv is clearly linked to upper limb motor control (Graziano, 1999); the V5 activation could indicate the generation of visual postural or motion templates (Zimmermann et al., 2012, 2016; Yon et al., 2018; Muckli et al., 2015), perhaps as 'simulated' sensory input for state estimation.

Finally, the cerebellum, the SMA, and several other regions that increased activity during the preparation for incongruent visuomotor mappings showed surprise related responses. Thus, brain regions updating predictions also responded to the respective prediction errors. Further surprise responses were located in medial and inferior frontal areas associated with general error detection

(Ullsperger et al., 2014; cf. Quirmbach & Limanowski, 2022) and in temporoparietal areas implied in visuomotor mismatch detection (Leube et al., 2003; Balslev et al., 2005; Farrer et al., 2008; van Kemenade et al., 2017, 2019; Limanowski et al., 2017). That not all of the 'predictive' regions showed surprise responses, and vice versa, can inform neuroanatomical models of visuomotor adaptation (Synofzik et al., 2006; Grafton et al., 2008). That surprise responses did not differ between congruent and incongruent mappings suggests that the forward model's predictions were, generally, successfully updated in response to the cues.

A limitation of our study were the relatively frequent catch trials that more likely followed invalid (surprising) than valid movements. Participants could thus have associated surprising movements with subsequent button presses, introducing a bias by movement preparation, initiation, or reward expectations. Thus, some surprise effects, e.g. in motor areas, could partly reflect anticipatory or more extensive movements of the responding (right) hand; which needs to be clarified by future work. However, this did not affect the between-condition contrasts of the movement (or preparation) periods, as surprise and catch trials were balanced across conditions. Furthermore, we cannot exclude potential eye movement influences on our results (cf. Gorbet & Sergio, 2016). However, our non-spatial movement task and the instructed central fixation rendered systematic eye movements useless; and our control experiment suggested that gaze behavior did not significantly differ between conditions. While our study design yielded strong experimental effects between conditions, the absence of a proper BOLD baseline limits the interpretation of the individual regressors' contrast estimates as "(de-)activations" from rest. Thus, planning related changes in brain metabolism relative to a resting state should be addressed with alternative designs. Our univariate results highlight regional metabolic differences depending on planning and predictive processes – these insights could be complemented by multivariate approaches, which may reveal the encoding of more abstract action planning related information such as different sensory content representations (cf. Ariani et al., 2015, 2018; Gallivan et al., 2013; Yon et al., 2018; Townsend et al., 2011; Turella et al., 2020).

To conclude, our results suggest visual feedback prediction is an inherent component of motor planning; and a key role of the cerebellum and the left aIPS in the underlying forward predictions and integrative processes, respectively.

References

Andersen, R. A., & Buneo, C. A. (2002). Intentional maps in posterior parietal cortex. Annual review of neuroscience, 25(1), 189-220.

Ames, K. C., Ryu, S. I., & Shenoy, K. V. (2014). Neural Dynamics of Reaching following Incorrect or Absent Motor Preparation. *Neuron*, *81*(2), 438–451. https://doi.org/10.1016/j.neuron.2013.11.003

Ariani, G., Wurm, M. F., & Lingnau, A. (2015). Decoding internally and externally driven movement plans. *Journal of Neuroscience, 35*(42), 14160-14171.

Ariani, G., Oosterhof, N. N., & Lingnau, A. (2018). Time-resolved decoding of planned delayed and immediate prehension movements. *Cortex*, *99*, 330–345. https://doi.org/10.1016/j.cortex.2017.12.007

Ariani, G., Pruszynski, J. A., & Diedrichsen, J. (2022). Motor planning brings human primary somatosensory cortex into action-specific preparatory states. *elife, 11*, e69517.

Balslev, D., Nielsen, F. Å., Paulson, O. B., & Law, I. (2005). Right Temporoparietal Cortex Activation during Visuo-proprioceptive Conflict. *Cerebral Cortex*, *15*(2), 166–169. https://doi.org/10.1093/cercor/bhh119

Bernier, P.-M., Cieslak, M., & Grafton, S. T. (2012). Effector selection precedes reach planning in the dorsal parietofrontal cortex. *Journal of Neurophysiology*, *108*(1), 57–68. https://doi.org/10.1152/jn.00011.2012

Blakemore, S. J., & Sirigu, A. (2003). Action prediction in the cerebellum and in the parietal lobe. Experimental Brain Research, 153, 239-245.

Blakemore, S.-J., Wolpert, D. M., & Frith, C. D. (1998). Central cancellation of self-produced tickle sensation. *Nature Neuroscience*, *1*(7), 635–640. https://doi.org/10.1038/2870

Bohlhalter, S., Hattori, N., Wheaton, L., Fridman, E., Shamim, E. A., Garraux, G., & Hallett, M. (2009). Gesture subtype–dependent left lateralization of praxis planning: An event-related fMRI study. *Cerebral cortex, 19*(6), 1256-1262.

Bonda, E., Petrides, M., Frey, S., & Evans, A. (1995). Neural correlates of mental transformations of the body-in-space. *Proceedings of the National Academy of Sciences*, *92*(24), 11180–11184. https://doi.org/10.1073/pnas.92.24.11180

Brass, M., Zysset, S., & von Cramon, D. Y. (2001). The Inhibition of Imitative Response Tendencies. *NeuroImage*, *14*(6), 1416–1423. https://doi.org/10.1006/nimg.2001.0944

Churchland, M. M., Cunningham, J. P., Kaufman, M. T., Ryu, S. I., & Shenoy, K. V. (2010). Cortical Preparatory Activity: Representation of Movement or First Cog in a Dynamical Machine? *Neuron*, *68*(3), 387–400. https://doi.org/10.1016/j.neuron.2010.09.015

Cisek, P., & Kalaska, J. F. (2002). Simultaneous Encoding of Multiple Potential Reach Directions in Dorsal Premotor Cortex. *Journal of Neurophysiology*, *87*(2), 1149–1154. https://doi.org/10.1152/jn.00443.2001

Cisek, P., & Kalaska, J. F. (2005). Neural Correlates of Reaching Decisions in Dorsal Premotor Cortex: Specification of Multiple Direction Choices and Final Selection of Action. *Neuron*, *45*(5), 801–814. https://doi.org/10.1016/j.neuron.2005.01.027

Crammond, D. J., & Kalaska, J. F. (1989). Neuronal activity in primate parietal cortex area 5 varies with intended movement direction during an instructed-delay period. *Experimental Brain Research*, *76*(2). https://doi.org/10.1007/BF00247902

Cui, H., & Andersen, R. A. (2011). Different representations of potential and selected motor plans by distinct parietal areas. Journal of Neuroscience, 31(49), 18130-18136.

Cunnington, R., Bradshaw, J. L., & Iansek, R. (1996). The role of the supplementary motor area in the control of voluntary movement. *Human Movement Science*, *15*(5), 627–647. https://doi.org/10.1016/0167-9457(96)00018-8

de Lange, F. P., Helmich, R. C., & Toni, I. (2006). Posture influences motor imagery: An fMRI study. *NeuroImage*, *33*(2), 609–617. https://doi.org/10.1016/j.neuroimage.2006.07.017

Dekleva, B. M., Kording, K. P., & Miller, L. E. (2018). Single reach plans in dorsal premotor cortex during a two-target task. *Nature Communications*, *9*(1), 3556. https://doi.org/10.1038/s41467-018-05959-y

Desmurget, M., Epstein, C. M., Turner, R. S., Prablanc, C., Alexander, G. E., & Grafton, S. T. (1999). Role of the posterior parietal cortex in updating reaching movements to a visual target. *Nature Neuroscience*, *2*(6), 563–567. https://doi.org/10.1038/9219

Donchin, O., Rabe, K., Diedrichsen, J., Lally, N., Schoch, B., Gizewski, E. R., & Timmann, D. (2012). Cerebellar regions involved in adaptation to force field and visuomotor perturbation. Journal of neurophysiology, 107(1), 134-147.

Duque, J., Olivier, E., & Rushworth, M. (2013). Top–down inhibitory control exerted by the medial frontal cortex during action selection under conflict. Journal of cognitive neuroscience, 25(10), 1634-1648.

Ehrsson, H. H., Spence, C., & Passingham, R. E. (2004). That's My Hand! Activity in Premotor Cortex Reflects Feeling of Ownership of a Limb. *Science*, *305*(5685), 875–877. https://doi.org/10.1126/science.1097011

Eickhoff, S. B., Stephan, K. E., Mohlberg, H., Grefkes, C., Fink, G. R., Amunts, K., & Zilles, K. (2005). A new SPM toolbox for combining probabilistic cytoarchitectonic maps and functional imaging data. *NeuroImage*, *25*(4), 1325–1335. https://doi.org/10.1016/j.neuroimage.2004.12.034

Fabbri, S., Strnad, L., Caramazza, A., & Lingnau, A. (2014). Overlapping representations for grip type and reach direction. Neuroimage, 94, 138-146.

Farrer, C., Frey, S. H., Van Horn, J. D., Tunik, E., Turk, D., Inati, S., & Grafton, S. T. (2008). The angular gyrus computes action awareness representations. *Cerebral cortex*, *18*(2), 254-261.

Fernandez-Ruiz, J., Goltz, H. C., DeSouza, J. F. X., Vilis, T., & Crawford, J. D. (2007). Human Parietal "Reach Region" Primarily Encodes Intrinsic Visual Direction, Not Extrinsic Movement Direction, in a Visual–Motor Dissociation Task. *Cerebral Cortex*, *17*(10), 2283–2292. https://doi.org/10.1093/cercor/bhl137

Ferraina, S., & Bianchi, L. (1994). Posterior parietal cortex: Functional properties of neurons in area 5 during an instructed-delay reaching task within different parts of space. *Experimental Brain Research*, *99*(1). https://doi.org/10.1007/BF00241423

Gallivan, J. P., McLean, D. A., Flanagan, J. R., & Culham, J. C. (2013). Where One Hand Meets the Other: Limb-Specific and Action-Dependent Movement Plans Decoded from Preparatory Signals in Single Human Frontoparietal Brain Areas. *The Journal of Neuroscience*, *33*(5), 1991–2008. https://doi.org/10.1523/JNEUROSCI.0541-12.2013

Gertz, H., & Fiehler, K. (2015). Human posterior parietal cortex encodes the movement goal in a pro-/anti-reach task. *Journal of Neurophysiology*, *114*(1), 170–183. https://doi.org/10.1152/jn.01039.2014

Gläscher, J. (2009). Visualization of group inference data in functional neuroimaging. *Neuroinformatics*, 7, 73-82.

Gorbet, D. J., & Sergio, L. E. (2016). Don't watch where you're going: The neural correlates of decoupling eye and arm movements. *Behavioural Brain Research*, *298*, 229–240. https://doi.org/10.1016/j.bbr.2015.11.012

Grafton, S. T., Hazeltine, E., & Ivry, R. B. (2002). Motor sequence learning with the nondominant left hand: A PET functional imaging study. Experimental brain research, 146, 369-378.

Grafton, S. T., Schmitt, P., Van Horn, J., & Diedrichsen, J. (2008). Neural substrates of visuomotor learning based on improved feedback control and prediction. Neuroimage, 39(3), 1383-1395.

Graziano, M. S. A. (1999). Where is my arm? The relative role of vision and proprioception in the neuronal representation of limb position. *Proceedings of the National Academy of Sciences*, *96*(18), 10418–10421. https://doi.org/10.1073/pnas.96.18.10418

Grefkes, C., & Fink, G. R. (2005). REVIEW: The functional organization of the intraparietal sulcus in humans and monkeys: Intraparietal sulcus in monkeys and humans, C. Grefkes and G. R. Fink. *Journal of Anatomy*, *207*(1), 3–17. https://doi.org/10.1111/j.1469-7580.2005.00426.x

Haaland, K. Y., Harrington, D. L., & Knight, R. T. (2000). Neural representations of skilled movement. Brain, 123(11), 2306-2313.

Hardwick, R. M., Rottschy, C., Miall, R. C., & Eickhoff, S. B. (2013). A quantitative meta-analysis and review of motor learning in the human brain. *NeuroImage*, *67*, 283–297. https://doi.org/10.1016/j.neuroimage.2012.11.020

Hétu, S., Grégoire, M., Saimpont, A., Coll, M. P., Eugène, F., Michon, P. E., & Jackson, P. L. (2013). The neural network of motor imagery: an ALE meta-analysis. Neuroscience & Biobehavioral Reviews, 37(5), 930-949.

Hoshi, E., & Tanji, J. (2002). Contrasting Neuronal Activity in the Dorsal and Ventral Premotor Areas During Preparation to Reach. *Journal of Neurophysiology*, *87*(2), 1123–1128. https://doi.org/10.1152/jn.00496.2001

Izawa, J., Criscimagna-Hemminger, S. E., & Shadmehr, R. (2012). Cerebellar contributions to reach adaptation and learning sensory consequences of action. Journal of Neuroscience, 32(12), 4230-4239.

Jeannerod, M. & Decety, J. Mental motor imagery: a window into the representational stages of action. Curr. Opin. Neurobiol. 5, 727–732 (1995).

Kaufman, M. T., Churchland, M. M., Ryu, S. I., & Shenoy, K. V. (2014). Cortical activity in the null space: Permitting preparation without movement. *Nature Neuroscience*, *17*(3), 440–448. https://doi.org/10.1038/nn.3643

Kilteni, K., Andersson, B. J., Houborg, C., & Ehrsson, H. H. (2018). Motor imagery involves predicting the sensory consequences of the imagined movement. Nature communications, 9(1), 1617.

Kilteni, K., Houborg, C., & Ehrsson, H. H. (2023). Brief temporal perturbations in somatosensory reafference disrupt perceptual and neural attenuation and increase supplementary motor area–cerebellar connectivity. Journal of Neuroscience.

Klaes, C., Westendorff, S., Chakrabarti, S., & Gail, A. (2011). Choosing Goals, Not Rules: Deciding among Rule-Based Action Plans. *Neuron*, *70*(3), 536–548. https://doi.org/10.1016/j.neuron.2011.02.053

Króliczak, G., & Frey, S. H. (2009). A common network in the left cerebral hemisphere represents planning of tool use pantomimes and familiar intransitive gestures at the hand-independent level. *Cerebral Cortex, 19*(10), 2396-2410.

Kuang, S., Morel, P., & Gail, A. (2015). Planning Movements in Visual and Physical Space in Monkey Posterior Parietal Cortex. *Cerebral Cortex*, bhu312. https://doi.org/10.1093/cercor/bhu312

Leube, D., Knoblich, G., Erb, M., Grodd, W., Bartels, M., & Kircher, T. T. (2003). The neural correlates of perceiving one's own movements. *NeuroImage*, *20*(4), 2084–2090. https://doi.org/10.1016/j.neuroimage.2003.07.033

Limanowski, J., & Blankenburg, F. (2016). Integration of Visual and Proprioceptive Limb Position Information in Human Posterior Parietal, Premotor, and Extrastriate Cortex. *The Journal of Neuroscience*, *36*(9), 2582–2589. https://doi.org/10.1523/JNEUROSCI.3987-15.2016

Limanowski, J., & Blankenburg, F. (2017). Posterior parietal cortex evaluates visuoproprioceptive congruence based on brief visual information. *Scientific Reports*, *7*(1), 16659. https://doi.org/10.1038/s41598-017-16848-7

Limanowski, J., & Friston, K. (2020). Attentional Modulation of Vision Versus Proprioception During Action. *Cerebral Cortex*, *30*(3), 1637–1648. https://doi.org/10.1093/cercor/bhz192

Limanowski, J., Kirilina, E., & Blankenburg, F. (2017). Neuronal correlates of continuous manual tracking under varying visual movement feedback in a virtual reality environment. *NeuroImage*, *146*, 81–89. https://doi.org/10.1016/j.neuroimage.2016.11.009

Lindner, A., Iyer, A., Kagan, I., & Andersen, R. A. (2010). Human Posterior Parietal Cortex Plans Where to Reach and What to Avoid. *The Journal of Neuroscience*, *30*(35), 11715–11725. https://doi.org/10.1523/JNEUROSCI.2849-09.2010

Lotze, M., Montoya, P., Erb, M., Hülsmann, E., Flor, H., Klose, U., Birbaumer, N., & Grodd, W. (1999). Activation of Cortical and Cerebellar Motor Areas during Executed and Imagined Hand Movements: An fMRI Study. *Journal of Cognitive Neuroscience*, *11*(5), 491–501. https://doi.org/10.1162/089892999563553

Makoshi, Z., Kroliczak, G., & Van Donkelaar, P. (2011). Human Supplementary Motor Area Contribution to Predictive Motor Planning. *Journal of Motor Behavior*, *43*(4), 303–309. https://doi.org/10.1080/00222895.2011.584085

Medendorp, W. P., & Heed, T. (2019). State estimation in posterior parietal cortex: Distinct poles of environmental and bodily states. *Progress in neurobiology, 183*, 101691.

Miall, R. C., Weir, D. J., Wolpert, D. M., & Stein, J. F. (1993). Is the Cerebellum a Smith Predictor? *Journal of Motor Behavior*, *25*(3), 203–216. https://doi.org/10.1080/00222895.1993.9942050

Miall, R. C., & Wolpert, D. M. (1996). Forward models for physiological motor control. Neural networks, 9(8), 1265-1279.

Michaels, J. A., Dann, B., Intveld, R. W., & Scherberger, H. (2018). Neural Dynamics of Variable Grasp-Movement Preparation in the Macaque Frontoparietal Network. *The Journal of Neuroscience*, *38*(25), 5759–5773. https://doi.org/10.1523/JNEUROSCI.2557-17.2018

Muckli, L., De Martino, F., Vizioli, L., Petro, L. S., Smith, F. W., Ugurbil, K., ... & Yacoub, E. (2015). Contextual feedback to superficial layers of V1. *Current Biology, 25*(20), 2690-2695.

Mutha, P. K., Sainburg, R. L., & Haaland, K. Y. (2011). Left parietal regions are critical for adaptive visuomotor control. Journal of Neuroscience, 31(19), 6972-6981.

Nahab, F. B., Kundu, P., Gallea, C., Kakareka, J., Pursley, R., Pohida, T., Miletta, N., Friedman, J., & Hallett, M. (2011). The Neural Processes Underlying Self-Agency. *Cerebral Cortex*, *21*(1), 48–55. https://doi.org/10.1093/cercor/bhq059

Ogawa, K., Inui, T., & Sugio, T. (2006). Separating brain regions involved in internally guided and visual feedback control of moving effectors: An event-related fMRI study. *NeuroImage*, *32*(4), 1760–1770. https://doi.org/10.1016/j.neuroimage.2006.05.012

Ohata, R., Asai, T., Kadota, H., Shigemasu, H., Ogawa, K., & Imamizu, H. (2020). Sense of Agency Beyond Sensorimotor Process: Decoding Self-Other Action Attribution in the Human Brain. *Cerebral Cortex*, *30*(7), 4076–4091. https://doi.org/10.1093/cercor/bhaa028

Parr, T., Limanowski, J., Rawji, V., & Friston, K. (2021). The computational neurology of movement under active inference. Brain, 144(6), 1799-1818.

Pilacinski, A., Wallscheid, M., & Lindner, A. (2018). Human posterior parietal and dorsal premotor cortex encode the visual properties of an upcoming action. *PLOS ONE*, *13*(10), e0198051. https://doi.org/10.1371/journal.pone.0198051

Quirmbach, F., & Limanowski, J. (2022). A Crucial Role of the Frontal Operculum in Task-Set Dependent Visuomotor Performance Monitoring. *Eneuro*, *9*(2), ENEURO.0524-21.2021. https://doi.org/10.1523/ENEURO.0524-21.2021

Rosenbaum, D. A. (1980). Human movement initiation: specification of arm, direction, and extent. Journal of Experimental Psychology: General, 109(4), 444.

Rushworth, M. F., Krams, M., & Passingham, R. E. (2001). The attentional role of the left parietal cortex: the distinct lateralization and localization of motor attention in the human brain. Journal of cognitive neuroscience, 13(5), 698-710.

Sakai, K., Hikosaka, O., Miyauchi, S., Sasaki, Y., Fujimaki, N., & Pütz, B. (1999). Presupplementary Motor Area Activation during Sequence Learning Reflects Visuo-Motor Association. *The Journal of Neuroscience*, *19*(10), RC1–RC1. https://doi.org/10.1523/JNEUROSCI.19-10-j0002.1999

Schaffelhofer, S., Agudelo-Toro, A., & Scherberger, H. (2015). Decoding a Wide Range of Hand Configurations from Macaque Motor, Premotor, and Parietal Cortices. *The Journal of Neuroscience*, *35*(3), 1068–1081. https://doi.org/10.1523/JNEUROSCI.3594-14.2015

Schluter, N. D., Krams, M., Rushworth, M. F. S., & Passingham, R. E. (2001). Cerebral dominance for action in the human brain: the selection of actions. Neuropsychologia, 39(2), 105-113.

Scott, S. H. (2012). The computational and neural basis of voluntary motor control and planning. Trends in cognitive sciences, 16(11), 541-549.

Servos, P., Osu, R., Santi, A., & Kawato, M. (2002). The neural substrates of biological motion perception: an fMRI study. *Cerebral Cortex, 12*(7), 772-782.

Shadmehr, R., & Krakauer, J. W. (2008). A computational neuroanatomy for motor control. Experimental brain research, 185, 359-381.

Shadmehr, R., Smith, M. A., & Krakauer, J. W. (2010). Error correction, sensory prediction, and adaptation in motor control. Annual review of neuroscience, 33, 89-108.

Stanley, J., & Miall, R. C. (2007). Functional activation in parieto-premotor and visual areas dependent on congruency between hand movement and visual stimuli during motor-visual priming. Neuroimage, 34(1), 290-299.

Stanley, J., & Miall, R. C. (2009). Using predictive motor control processes in a cognitive task: behavioral and neuroanatomical perspectives. In: D. Sternad (ed.), Progress in motor control: A multidisciplinary perspective, Springer Science & Business Media, LLC 2009, 337-354.

Synofzik, M., Thier, P., & Lindner, A. (2006). Internalizing agency of self-action: perception of one's own hand movements depends on an adaptable prediction about the sensory action outcome. Journal of neurophysiology, 96(3), 1592-1601.

Synofzik, M., Lindner, A., & Thier, P. (2008). The cerebellum updates predictions about the visual consequences of one's behavior. *Current Biology, 18*(11), 814-818.

Tanji, J., & Evarts, E. V. (1976). Anticipatory activity of motor cortex neurons in relation to direction of an intended movement. *Journal of Neurophysiology*, *39*(5), 1062–1068. https://doi.org/10.1152/jn.1976.39.5.1062

Taylor, P. C., Nobre, A. C., & Rushworth, M. F. (2007). Subsecond changes in top–down control exerted by human medial frontal cortex during conflict and action selection: a combined transcranial magnetic stimulation–electroencephalography study. Journal of Neuroscience, 27(42), 11343-11353.

Todorov, E., & Jordan, M. I. (2002). Optimal feedback control as a theory of motor coordination. Nature neuroscience, 5(11), 1226-1235.

Townsend, B. R., Subasi, E., & Scherberger, H. (2011). Grasp Movement Decoding from Premotor and Parietal Cortex. *The Journal of Neuroscience*, *31*(40), 14386–14398.

Tseng YW, Diedrichsen J, Krakauer JW, Shadmehr R, Bastian AJ (2007) Sensory prediction errors drive cerebellum-dependent adaptation of reaching. J Neurophysiol 98:54–62https://doi.org/10.1523/JNEUROSCI.2451-11.2011

Turella, L., Rumiati, R., & Lingnau, A. (2020). Hierarchical action encoding within the human brain. Cerebral cortex, 30(5), 2924-2938.

Tzvi, E., Loens, S., & Donchin, O. (2022). Mini-review: the role of the cerebellum in visuomotor adaptation. The Cerebellum, 21(2), 306-313.

Ullsperger, M., Danielmeier, C., & Jocham, G. (2014). Neurophysiology of Performance Monitoring and Adaptive Behavior. *Physiological Reviews*, *94*(1), 35–79. https://doi.org/10.1152/physrev.00041.2012

Van Kemenade, B. M., Arikan, B. E., Kircher, T., & Straube, B. (2017). The angular gyrus is a supramodal comparator area in action–outcome monitoring. *Brain Structure and Function*, *222*(8), 3691–3703. https://doi.org/10.1007/s00429-017-1428-9

Van Kemenade, B. M., Arikan, B. E., Podranski, K., Steinsträter, O., Kircher, T., & Straube, B. (2019). Distinct Roles for the Cerebellum, Angular Gyrus, and Middle Temporal Gyrus in Action–Feedback Monitoring. *Cerebral Cortex*, *29*(4), 1520–1531. https://doi.org/10.1093/cercor/bhy048

Verhagen, L., Dijkerman, H. C., Medendorp, W. P., & Toni, I. (2012). Cortical Dynamics of Sensorimotor Integration during Grasp Planning. *The Journal of Neuroscience*, *32*(13), 4508–4519. https://doi.org/10.1523/JNEUROSCI.5451-11.2012

Wise, S. P., di Pellegrino, G., & Boussaoud, D. (1996). The premotor cortex and nonstandard sensorimotor mapping. *Canadian Journal of Physiology and Pharmacology*, *74*(4), 469–482. https://doi.org/10.1139/cjpp-74-4-469

Wolpert, D. M., & Kawato, M. (1998). Multiple paired forward and inverse models for motor control. *Neural networks, 11*(7-8), 1317-1329.

Wolpert, D. M., Miall, R. C., & Kawato, M. (1998). Internal models in the cerebellum. Trends in cognitive sciences, 2(9), 338-347.

Wolpert, D. M., & Ghahramani, Z. (2000). Computational principles of movement neuroscience. Nature neuroscience, 3(11), 1212-1217.

Yomogida, Y., Sugiura, M., Sassa, Y., Wakusawa, K., Sekiguchi, A., Fukushima, A., Takeuchi, H., Horie, K., Sato, S., & Kawashima, R. (2010). The neural basis of agency: An fMRI study. *NeuroImage*, *50*(1), 198–207. https://doi.org/10.1016/j.neuroimage.2009.12.054

Yon, D., Gilbert, S. J., de Lange, F. P., & Press, C. (2018). Action sharpens sensory representations of expected outcomes. *Nature Communications*, *9*(1), 4288. https://doi.org/10.1038/s41467-018-06752-7

Zimmermann, M., Verhagen, L., de Lange, F. P., & Toni, I. (2016). The extrastriate body area computes desired goal states during action planning. Eneuro, 3(2).

Zimmermann, M., Meulenbroek, R. G. J., & de Lange, F. P. (2012). Motor Planning Is Facilitated by Adopting an Action's Goal Posture: An fMRI Study. *Cerebral Cortex*, *22*(1), 122–131. https://doi.org/10.1093/cercor/bhr098